1

Promiscuous esterases counterintuitively are less flexible
than specific ones

Christina Nutschel[1], Cristina Coscolín[2], Daniel Mulnaes[3], Benoit David[3], Manuel Ferrer[2],

Karl-Erich Jaeger[4,5], Holger Gohlke[1,3,*]

[1] John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC) and

Institute of Biological Information Processing (IBI-7: Structural Biochemistry),

Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

[2] Institute of Catalysis, Consejo Superior de Investigaciones Científicas, 28049 Madrid, Spain

[3] Institute for Pharmaceutical and Medicinal Chemistry, Heinrich Heine University Düsseldorf,

40225 Düsseldorf, Germany

[4] Institute of Molecular Enzyme Technology, Heinrich Heine University Düsseldorf, 52425

Jülich, Germany

[5] Institute of Bio- and Geosciences IBG-1: Biotechnology, Forschungszentrum Jülich GmbH,

52425 Jülich, Germany

Running title: Flexibility and promiscuity of esterases

Keywords: Esterase promiscuity, structural flexibility, thermostability, Constraint Network

Analysis, TopModel, TopScore, conformational proofreading

*Corresponding author:

John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC), and

Institute of Biological Information Processing (IBI-7: Structural Biochemistry)

Forschungszentrum Jülich GmbH

Wilhelm-Johnen-Straße

52425 Jülich

Germany

Email: gohlke@uni-duesseldorf.de or h.gohlke@fz-juelich.de

Abstract

Understanding mechanisms of promiscuity is increasingly important from a fundamental and application point of view. As to enzyme structural dynamics, more promiscuous enzymes generally have been recognized to also be more flexible. However, examples for the opposite received much less attention. Here, we exploit comprehensive experimental information on the substrate promiscuity of 147 esterases tested against 96 esters together with computationally efficient rigidity analyses to understand the molecular origin of the observed promiscuity range. Unexpectedly, our data reveal that promiscuous esterases are significantly less flexible than specific ones, are significantly more thermostable, and have a significantly increased specific activity. These results may be reconciled with a model according to which structural flexibility in the case of specific esterases serves for conformational proofreading. Our results signify that esterase sequence space can be screened by rigidity analyses for promiscuous esterases as starting points for further exploration in biotechnology and synthetic chemistry.

1. Introduction

Enzymes involved in primary metabolism typically exquisitely discriminate against other metabolites. Yet, evolution of specificity is only pushed by nature to the point at which 'unauthorized' reactions do not impair the fitness of the organism (1). As a result, the universe of promiscuous activities available in nature has been suggested to be enormous (2, 3). Understanding mechanisms of promiscuity has thus become increasingly important for the fundamental understanding of molecular recognition and how enzyme function has evolved over time(4) but also to optimize enzyme engineering applications (5). A particular challenge in the latter case is the ability to discover a suitable enzyme with 'sufficient' promiscuous activity to serve as a starting point for further exploration (1).

Enzyme structural dynamics, besides its role in catalysis (6, 7) and allosteric regulation (8-11), has been recognized as likely the single most important mechanism by which promiscuity can be achieved (5). Prominent examples are human cytochrome P450 (CYP) enzymes, for which crystallographic studies and molecular simulations demonstrated that more promiscuous CYPs show larger structural plasticity and mobility (12-14), or TEM-1 β-lactamase and a resurrected progenitor, for which molecular simulations show that the pocket of the ancestral, and more promiscuous, enzyme fluctuates to a greater extent (15). However, examples for the opposite, i.e., conformational changes selected in evolution such that they enhance specificity in molecular recognition (16), have received much less attention in the context of enzyme promiscuity.

A clear limitation for scrutinizing the link between enzyme structural dynamics and substrate promiscuity is the general lack of large-scale data on one enzyme (super)family tested against a multitude of ligands (17) (cf. ref. (1) for notable exceptions). Likewise, acquiring information on enzyme dynamics at the atomistic level by experimental techniques or classical molecular dynamics (MD) simulations is burdensome. Here, we exploit comprehensive experimental information on the substrate promiscuity (18) of esterases (abbreviated EHs, for "Ester Hydrolases") (19) together with computationally efficient rigidity analyses (20-23) of comparative models of EHs to understand the molecular origin of the observed promiscuity range. Enzyme rigidity, or its opposite flexibility, are static properties that denote the *impossibility*, or *possibility*, *of motions* in an enzyme under force, without giving information about directions and magnitudes of movements (23). Enzyme flexibility, thus, should not be confused with enzyme mobility, which describes *actual motions* in an enzyme. Rigidity analysis results do not rely on the correct description of the time-dependency of processes (23), which

makes them valuable in cases where timescales over multiple orders of magnitude may govern

such processes, like in enzyme dynamics (6, 7).

In recent years, EHs have obtained much attention in basic research and industrial applications

(24). EHs are widely distributed in nature within microbial communities (at least one EH is

found in each bacterial genome), they have been extensively examined with state-of-the-art

(meta)genomics techniques and investigated by functional screenings compared to many other

classes of enzymes. They also possess outstanding properties in terms of stability, reactivity,

and scalability that make them appropriate biocatalysts to improve competitiveness, innovation

capacity, and sustainability in a modern circular bio-economy (25). Recently, a large-scale

study on substrate promiscuity (P_{EH}, which denotes the number of esters hydrolyzed by an EH)

of 147 phylogenetically, environmentally, and structurally diverse microbial EHs was described

by Martínez-Martínez $et\ al.$ (19), in which all EHs were functionally assessed against a

customized library of 96 esters. As to mechanistic understanding, the authors related P_{EH} to a

structural parameter, the active site effective volume. However, the impact of enzyme flexibility

on P_{EH} was not assessed.

In our study, we thus ask the following questions: I) What is the relation between P_{EH} and EH

flexibility? II) Does this relation hold if experimentally determined EH thermostabilities are

used as proxies for enzyme flexibility? III) What is the relation between P_{EH} and EHs' specific

activities? IV) Is there a preference of promiscuous or specific EHs for a particular type of

esters. V) Can this preference be understood with respect to EHs flexibilities?

2. Materials and Methods

2.1.   Definition of data sets

The present study builds on the study from Martínez-Martínez *et al.* (19). In order to assess P_{EH}, the authors experimentally investigated 147 phylogenetically, environmentally, and structurally diverse microbial EHs (termed *experimental data set*) against a customized library of 96 different esters. Two commercial lipases, which have found wide biotechnological applications, CalA and CalB from *Pseudozyma aphidis* (formerly *Candida antarctica*), were included for comparison. For details on determining and classifying P_{EH}, see **Supplemental Materials and Methods**. To validate that P_{EH} defines promiscuity of EHs in a quantitative manner, k_{cat} and K_m values were determined for ten expressed and purified EHs covering the entire P_{EH} range (see **section 2.9**) and a promiscuity index I (**see Supplemental Materials and Methods**, **Eq. S4**) computed and compared to P_{EH}. Finally, the similarity of the ester substrates was assessed by the maximum pairwise Tanimoto-Combo similarity score δ_{ij} for compound i versus j, accounting for shape and chemical complementary between 3D structures, and the mean maximum pairwise Tanimoto-Combo similarity score δ_i of a substrate i to all other substrates in the data set (see **Supplemental Methods and Materials**).

As our computational approach involves extensive molecular dynamics (MD) simulations for generating conformational ensembles **(see section 2.3)**, we selected 35 EHs from the *volume data set* (termed *flexibility data set*) for comparative modeling **(see section 2.2)**. The criteria for choosing EHs of the *flexibility data set* are explained in **section 3.1**.

2.2.   Comparative modelling and validations of the *flexibility data set*

Comparative models of the *flexibility data set* **(see section 2.1)** were generated using our in-house structure prediction meta-tool TopModel (26) that has been successfully applied in previous studies (27-30). TopModel uses multiple state-of-the-art threading and sequence/structure alignment tools to generate a large ensemble of models from different pairwise and multiple alignments of the top five highest ranked template structures. The TopModel software is available at https://cpclab.uni-duesseldorf.de/index.php/Software.

The quality of the homology models was assessed by our meta Model Quality Assessment Program (meta-MQAP) TopScore (31). TopScore uses deep neural networks (DNN) to

combine scores from 15 different primary MQAP to predict accurate residue-wise and whole-

protein error estimates. For details on model quality assessment by TopScore and validation,

see **Supplemental Materials and Methods**.

To test whether CARs of the homology models are accessible for substrates, we applied the

CAVER 3.0.3 PyMOL Plugin (32). Starting points for the computations were defined based on

the Cartesian coordinates of the CARs' center of mass (COM). Default values were used for

the probe radius (0.9 Å), shell radius (3.0 Å), and shell depth (4.0 Å).

2.3.   Generation of structural ensembles

Structural ensembles of EHs were generated by all-atom MD simulations of in total 5 μs

simulation time per EH. For details on starting structure preparation, parametrization, and

equilibration see **Supplemental Materials and Methods**.

All minimization, equilibration, and production simulations were performed with the

pmemd.cuda module (33) of Amber19 (34). During production simulations, we set the time step

for the integration of Newton's equation of motion to 4 fs following the hydrogen mass

repartitioning strategy (35). Coordinates were stored into a trajectory file every 200 ps. This

resulted in 5000 configurations for each production run that were considered for subsequent

analyses.

2.4.   Constraint Network Analysis

The flexibility analyses were performed with the Constraint Network Analysis (CNA) software

package (version 3.0) (20-23). CNA functions as front- and back-end to the graph theory-based

software Floppy Inclusions and Rigid Substructure Topography (FIRST) (36). Applying CNA

to biomolecules aims at identifying their composition of rigid clusters and flexible regions,

which can aid in the understanding of biomolecular structure, stability, and function (21-23).

As the mechanical heterogeneity of biomolecular structures is intimately linked to their diverse

biological functions, biomolecules generally show a hierarchy of rigidity and flexibility (20).

In CNA, biomolecules are modeled as constraint networks in a *body-and-bar* representation,

which has been described in detail by Hesphenheide *et al*. (37). A fast combinatorial algorithm,

the *pebble game*, counts the bond rotational degrees of freedom and floppy modes (internal,

independent degrees of freedom) in the constraint network (38). In order to monitor the

hierarchy of rigidity and flexibility of biomolecules, CNA performs thermal unfolding
simulations by consecutively removing non-covalent constraints (hydrogen bonds, including
salt bridges) from a network in increasing order of their strength (39-41). For details on thermal
unfolding simulations, see **Supplemental Materials and Methods**. To improve the robustness
and investigate the statistical uncertainty, we carried out CNA on ensembles of network
topologies (ENT^{MD}) generated from MD trajectories **(see section 2.3)** (42).

The CNA software is available under academic license at https://cpclab.uni-
duesseldorf.de/index.php/Software and the CNA web server is accessible at https://cpclab.uni-
duesseldorf.de/cna.

2.5.   Local and global indices

From the thermal unfolding simulations, CNA computes a comprehensive set of indices to
quantify biologically relevant characteristics of the protein's stability. *Global* indices are used
for determining the rigidity and flexibility at a macroscopic level; *local* indices determine the
rigidity and flexibility at a microscopic level of bonds (43). The cluster configuration entropy
H_{type2} is a *global* index that has been introduced by Radestock and Gohlke (20). As done
previously, we applied H_{type2} as a measure for global structural stability of proteins (20, 41, 44-
48). The stability map rc_{ij} is a *local* index that has been introduced by Radestock and Gohlke
(20). We applied rc_{ij} as a measure for local structural stability of proteins in previous studies
(45, 47, 48). For details on both indices, see **Supplemental Materials and Methods**.

2.6.   Root mean square fluctuations

The per-residue root-mean-square fluctuations were calculated for each EH ($RMSF_{EH}$) and for
its CARS ($RMSF_{CAR}$) based on the MD trajectories **(see section 2.3)**. Prior to the calculations,
the structures of each trajectory were superimposed onto the average structure using the 90%
least mobile residues of the respective EHs (49).

2.7.   Torsion angles

For each of the 96 esters, the number of freely rotatable bonds (torsion angles, TA) was

calculated based on the SMILES codes provided by Martínez-Martínez *et al.* (19).

To compare how many esters with a specific TA are hydrolyzed by each EH, we calculated the

normalized proportion of ester hydrolysis with a specific TA (*Norm*ester(TA)) as the number of

hydrolyzed esters with a specific TA (*Ester*hydrolysed (TA)) divided by the total number of esters

in the data set with this specific TA (*Ester*library(TA)) (**Eq. 1**).

$$Norm_{ester}(TA) \ [\%] = \frac{Ester_{hydrolyzed}(TA)}{Ester_{library}(TA)} * 100\% \qquad \textbf{Eq. 1}$$

2.8.   Circular dichroism spectroscopy

Eleven serine ester hydrolases, EH1 (Protein data Bank acc. nr. 5JD4), EH_2 (GenBank acc. nr.

KY483643), EH_3 (GenBank acc. nr. KY483645), EH_4 (GenBank acc. nr. KR107250), EH_6

(GenBank acc. nr. KP347751), EH_8 (GenBank acc. nr. WP_011587341.1), EH_9 (GenBank acc.

nr. KY483648), EH_{16} (GenBank acc. nr. KP347759), EH_{21} (GenBank acc. nr. KP347760), EH_{37}

(GenBank acc. nr. KR107248), EH_{43} (GenBank acc. nr. KP347758) from metagenomic origin,

were used in the present study to perform CD determinations. The vector pET46 Ek/LIC and

the host *Escherichia coli* MC1061 were the source of the His6-tag EH_1, EH_4, EH_8, and EH_{37},

the vector pBXNH3 and the host *E. coli* MC1061 were the source of the His6-tag EH_2, EH_3,

and EH_9, and the vector p15Tv-L and the host E. coli BL21(DE3) were the source of EH_6, EH_{16},

EH_{21}, and EH_{43}. Prior to analyses, the soluble His-tagged proteins were produced and purified

after binding to a Ni-NTA His-Bind resin (Sigma-Aldrich, MO, US) as described by Martínez-

Martínez (19). Purity was assessed as >98% using SDS-PAGE analysis in a Mini PROTEAN

electrophoresis system (Bio-Rad, Madrid, Spain) and subsequent staining with Coomassie

Brilliant Blue. A total of about 10-20 mg total purified recombinant protein was obtained on

average from 1-liter culture. All proteins were stored at -20ºC at a concentration of 10 mg/ml

in 40 mM (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer (pH 7.0), until

used. Circular dichroism (CD) spectra were acquired between 190 and 270 nm with a Jasco J-

720 spectropolarimeter equipped with a Peltier temperature controller, employing a 0.1 mm cell

at 25°C. Spectra were analyzed, and denaturation temperatures were determined at 220 nm

between 10 and 85°C at a rate of 30°C per hour, in 40 mM (4-(2-hydroxyethyl)-1-

piperazineethanesulfonic acid (HEPES) buffer, pH 7.0. A protein concentration of 1.0 mg ml^{-1}

was used. Denaturation temperatures were calculated by fitting the ellipticity (mdeg) at 220 nm

at each of the different temperatures using a 5-parameters sigmoid fit with Sigma Plot 14.0.

2.9.   Kinetic parameter determination

Ten serine ester hydrolases, EH_1 (Protein data Bank acc. nr. 5JD4), EH_3 (GenBank acc. nr.

KY483645), EH_5 (GenBank acc. nr. KR107271), EH_7 (GenBank acc. nr. KY483644), EH_{12}

(GenBank acc. nr. KR107263), EH_{17} (GenBank acc. nr. KR107278), EH_{37} (GenBank acc. nr.

KR107248), EH_{102} (Protein data Bank acc. nr. 5JD3), EH_{114} (GenBank acc. nr. KR107274) and

EH_{127} (GenBank acc. nr. KR107253) from metagenomic origin, were used in the present study

to perform kinetic determinations (k_{cat} and K_m). The vector pET46 Ek/LIC and the host

Escherichia coli MC1061 were the source of the His$_6$-tag EH_1, EH_5, EH_{12}, EH_{17}, EH_{37}, EH_{102},

EH_{114}, and EH_{127}, and the vector pBXNH3 and the host *E. coli* MC1061 was the source of the

His$_6$-tag EH_3. The soluble His-tagged proteins was produced and purified as described by

Martínez-Martínez (19). For details see above.

The kinetic parameters were calculated at 550 nm using a continuous pH indicator (Phenol red;

$\varepsilon_{550\ nm}$ = 8450 M^{-1} cm^{-1}) assay at 550 nm in 384-well plates as previously described (Alonso et

al, 2020). Briefly, to 40 µl of 5 mM 4-(2-hydroxyethyl)-1-piperazinepropanesulfonic acid

(EPPS) buffer (pH 8.0) 2 µl of a stock ester solution was added to achieve the desired

concentration of each ester. Finally, 2 µl of stock protein solution was immediately added to

each well, to achieve the desired protein concentration, using an Eppendorf Repeater M4 pipette

(Eppendorf, Hamburg, Germany). The total reaction volume was 44 µl. Ester hydrolysis was

measured at 30°C in a Synergy HT Multi-Mode Microplate Reader in continuous mode at 550

nm over 24 h. One unit (U) of enzyme activity was defined as the amount of free enzyme or

enzyme bound to the carrier required to transform 1 µmol of substrate in 1 min under the assay

conditions using the reported extinction coefficient (Phenol red at 550 nm = 8450 M^{-1} cm^{-1}).

For K_m determination - [protein]: 4.5 µg ml^{-1}; [ester]: 0-100 mM; reaction volume: 44 µl; T:

30°C; pH: 8.0. For k_{cat} determination - [protein]: 0-270 µg ml^{-1}; [ester]: 100 mM; reaction

volume: 44 µl; T: 30°C; pH: 8.0. All values, in triplicates, were corrected for non-enzymatic

transformation. Kinetic parameters were calculated by fitting the data fit with Sigma Plot 14.0.

3. Results

3.1. Definition of data sets

To understand the structural origin of and develop a method to predict P_{EH}, the present study

builds on large-scale data from Martínez-Martínez *et al.* (19). The authors experimentally

investigated P_{EH} of 147 EHs (termed *experimental data set*) **(see section 2.1)**. In doing so,

compromises needed to be made regarding the measurement of catalytic activity, i.e., specific

activity was measured using enzymes expressed in *E. coli* without subsequent purification, only

a single concentration of wet cells expressing enzymes (0.4 mg per ester) was used to measure

activity, and the substrates were tested at a single concentration of circa 7 mM on average (19).

To validate that the P_{EH} derived from the measured activities define promiscuity of the enzymes

in a quantitative manner, k_{cat} and K_m values were determined now for ten expressed and purified

EH covering the entire P_{EH} range **(see section 2.9)**. k_{cat}/K_m ideally serves as the kinetic

parameter in enzyme promiscuity studies for comparison (50, 51). From k_{cat}/K_m values of an

enzyme towards a defined set of substrates, a quantitative index of promiscuity *I* **(Eq. S4)** can

be calculated based on information entropy (51). *I* yields a very good and significant ($R^2 = 0.79$,

$p = 0.0003$) correlation with P_{EH}, indicating that P_{EH} relates to EH promiscuity in a quantitative

manner **(Figure S1),** although the range of *I* suggests that large P_{EH} may still be associated with

moderate promiscuity. Note that, although the promiscuity index is a functional parameter that

is defined for a specified set of substrates, promiscuity indices for different enzymes are

quantitatively comparable if they have been calculated using the same substrate set (51).

Furthermore, for the ten EH and using the colorimetric assay herein used, k_{cat}/K_m can be

measured with a standard error of the mean as low as 0.05 min^{-1} mM^{-1}, which corresponds to

k_{cat} and K_m fitting values 2-fold above the background signals under assay conditions for each

of the enzymes and esters. When $k_{cat}/K_m > 0.05$ min^{-1} mM^{-1} is used as a criterion to define that

a substrate is hydrolyzed, the resulting number of substrates for the ten EHs yields an excellent

and significant ($R^2 = 0.99$, $p < 10^{-4}$) correlation with P_{EH} **(Figure S2)**, again indicating that P_{EH}

relates to EH promiscuity in a quantitative manner. Finally, esters that are chemically similar

to each other are expected to be metabolized similarly by an EH; such correlations in the

substrate set would reduce the effective EH promiscuity. Therefore, similarity of the substrates

was assessed by the maximum pairwise Tanimoto-Combo similarity score δ_{ij} for compound *i*

versus *j*, which is bounded between 0 for dissimilar compounds and 2 for identical ones, and

the mean maximum pairwise Tanimoto-Combo similarity score δ_i of a substrate *i* to all other

substrates in the data set; the Tanimoto-Combo similarity score accounts for shape and chemical

complementary between 3D structures as determined by the Rapid Overlay of Chemical
Structures approach (52). Complete linkage clustering on the pairwise distance matrix
calculated for all 96 compounds from δ_{ij} yielded 20 clusters at a distance of 1.0 (**Figure S3**),
which is equivalent to $\delta_{ij} = 1.0$, indicating that on average less than five esters share a similarity
that is half-way between dissimilar and identical. The negatively skewed histogram of δ_i
furthermore shows that δ_i peaks at 1.0 and is below 1.2 (**Figure S4**), indicating that an ester
generally shares a similarity to all other esters that is half-way between dissimilar and identical,
or worse.

Additionally, Martínez-Martínez *et al.* ranked (classified) P_{EH} of 96 EHs (termed *volume data*
set) based on the active site effective volume (**see section 2.1**) (**Eq. S1**) (19), which will be used
here as a reference to compare the power of P_{EH} predictions. As our computational approach
involves extensive MD simulations for generating conformational ensembles (**see section 2.3**),
we selected 35 EHs from the *volume data set* based on the following criteria; they constitute
the *flexibility data set*. I) The data set contains eleven EHs with known crystal structures
(including the commercial EHs CalA and CalB) (**Figure 1A, Table S1**) and 24 EHs for which
no experimental structure is available but for which comparative models can be generated (**see**
section 3.2) (**Figure 1A, Table S2**). That way, we can probe to what extent the source of
structural information influences the outcome of our results. II) The chosen EHs of the data set
show high diversities as to P_{EH} and association to esterase families (F_{EH}, as defined by Arpigny
and Jaeger (53)), similar to those found for the *volume data set* (**Figures S1 and S2, Tables S3**
and S4). This resulted in P_{EH} ranging from 4 to 72 (**Figure 1A, Tables S1 and S2**). In the
following, we consider P_{EH} as *low* if the EH hydrolyzes ≤ 9 esters (11% of the data set), as
moderate if the EH hydrolyzes between 10 and 29 esters (49%), and as *high* if the EH
hydrolyzes ≥ 30 esters (40%) (**Figure S5, Table S3**). The data set covers eleven F_{EH} of which
F_{IV} (44% of the data set) and F_V (21%) are the best represented ones (**Figure S6, Table S4**).
This reflects the proportion of their presence in the *volume data set*. III) Only EHs with amino
acid sequence identities $\geq 25\%$ in comparison to an existing crystal structure were considered
(**see section 2.1**) in order to ensure a sufficient quality of generated comparative models.

Finally, in order to uniformly depict the results across the present study, six EHs were selected
as representatives of the *flexibility data set* based on P_{EH} (termed *representative data set*): EHs
with the lowest (EH115) or highest P_{EH} (EH001) and known crystal structures, EHs with the
lowest (EH127) or the highest P_{EH} (EH005) and unknown crystal structures, and commercial
EHs with the lowest (CalA) or highest P_{EH} (CalB) (**Figure 1A-D, Tables S1 and S2**).

3.2. Comparative models of EHs generated by TopModel show an overall and residue-wise good quality

To generate structural models of EHs as starting points for our investigations, we performed template-based modeling of the *flexibility data set* using TopModel (26) **(see section 2.2)**. In doing so, we also generated comparative models of the eleven EHs for which crystal structures are available. These structural models will be used to judge the quality of the comparative modeling.

The quality of the comparative models of the *flexibility data set* were assessed with TopScore (31), a meta Model Quality Assessment Program (meta-MQAP) **(see section 2.2)**. For the eleven Es with known crystal structure, the global TopScores range from 0.074 to 0.305 **(Figure 1A, Table S1)**. As the global TopScore describes whole-protein error estimates, this shows that the structures contain between 7.4 and 30.5% error. Notably, the global TopScores well and significantly correlate (R^2 = 0.61, p = 0.004) with values of 1 – lDDT computed from comparisons of the comparative models of EHs with known crystal structure against these experimental reference structures, indicating that global TopScores are well suited to assess the model quality of EHs **(Figure S7, Table S5)**. The global TopScore values of the comparative models of the other 24 EHs range from 0.087 to 0.269 **(Figure 1A, Table S2)**, indicating that these models are of equal quality than the ones for EHs with known crystal structure. The TopScore values of the *representative data set* lie in a comparable range **(Figure 1A, Tables S1 and S2)**. Moreover, the comparative models of the *flexibility data set* show low residue-wise TopScore values (31), indicating that all parts of a model are of good quality. We illustrate this for the residue-wise TopScore of the comparative models of the *representative data set* **(Figures 1B-D)**. This also applies to structural regions around CARs **(Figures 1B-D)**. That way, it was possible to confirm CARs in models of EHs with known crystal structures and to unambiguously identify CARs in models of EHs with unknown crystal structures **(Figures 1B-D, Tables S1 and S2)**.

Additionally, we validated that CARs in all models are accessible for substrates according to CAVER results (32) **(see section 2.2)**, i.e., that all models are in an open conformation: CARs are either located on the protein surface or are buried and connected with the surface by tunnels. We illustrate this for the comparative models of the *representative data set* **(Figure S8)**.

348　To conclude, comparative models were generated for 35 EHs of the *flexibility data set* using

349　TopModel. The models showed both an overall and residue-wise good structural quality.

350　Additionally, we validated that CARs in all models are accessible for substrates.

3.3.　　Promiscuous EHs are globally less flexible

Previous studies indicated that enzyme flexibility influences the substrate promiscuity of

enzymes (12-14). For gaining insights into how the flexibility of EHs is linked to P_{EH}, we

applied CNA (21, 23), a rigidity theory-based approach to analyze biomolecular statics (21-23),

to the *flexibility data set* **(see sections 2.4)**. To improve the robustness and investigate the

statistical uncertainty, for each of the comparative models we carried out CNA on ensembles

of network topologies (ENT^{MD}) generated from five MD trajectories of 1 µs length each (44)

(see sections 2.3 and 2.4). In order to investigate if the global flexibility of the EHs influences

P_{EH}, we predicted T_p, the phase transition temperature previously applied as a measure of

structural stability of a protein (20, 41, 44-48), for each EH **(see section 2.5)**. T_p was averaged

over five ensembles **(see sections 2.3 and 2.4)**, resulting in all but one case in SEM < 1.87 K

(Figure 2A, Tables S1 and S2).

T_p and P_{EH} of the *flexibility data set* are well and significantly correlated ($R^2 = 0.60$, $p = 5.4*10^{-8}$)

(Figure 2A). To validate the consistency of our approach, we considered EHs with known

or unknown crystal structures separately. In both cases, good and significant correlations

between T_p and P_{EH} were revealed (known crystal structures: $R^2 = 0.48$, $p = 0.019$; unknown

crystal structures: $R^2 = 0.73$, $p = 1.1*10^{-7}$), lending support to the quality of comparative models

predicted with TopModel and indicating that future predictions on EHs with unknown

experimental structures should be promising. Notably, EHs with high P_{EH} have a high T_p and

vice versa, i.e., promiscuous EHs are globally less flexible. Exemplarily, this is depicted for

EHs of the *representative dataset* with known crystal structures and lowest (EH115) or highest

P_{EH} (EH001), which showed T_p of 322.3 K and 357.2 K, with unknown crystal structures and

lowest (EH127) or highest P_{EH} (EH005), which showed T_p of 318.6 K and 351.1 K, and CalA

and CalB, which showed T_p of 346.2 K and 351.6 K **(Figure 2A, Tables S1 and S2)**. The

differences in global structural stability of these EHs are illustrated by the rigid cluster

decomposition at 332 K during the thermal unfolding simulations **(Figure 2B-D)**: promiscuous

EHs are globally more structurally stable at the elevated temperature as indicated by fewer, but

larger, rigid clusters.

The EH flexibility analyzed so far is a static property and describes the potential of motions in
a biomolecule (23). Yet, direct information on mobility within EHs is available from the
ensembles generated by MD simulations. We thus computed exemplarily $RMSF_{EH}$, a measure
for protein mobility **(see section 2.6)**, across the ensembles of EHs from the *representative data*
set. $RMSF_{EH}$, averaged over all residues and all five MD trajectories, and P_{EH} do not yield a
significant correlation ($p = 0.13$) **(Figure S9A, Table S6)**, in contrast to T_p and P_{EH} ($R^2 = 0.93$,
$p = 1.8*10^{-3}$) **(Figure S9B, Table S6)**. Still, as promiscuous EHs are globally less mobile, the
same trend is obtained as in the case of the flexibility analysis.

To conclude, a good and significant correlation between T_p and P_{EH} was found for the *flexibility*
data set ($R^2 = 0.60$, $p = 5.4*10^{-8}$). These findings demonstrate that promiscuous EHs are
globally less flexible. $RMSF_{EH}$ is less predictive for P_{EH}, although again promiscuous EHs are
characterized by a lower global mobility, mutually confirming either result.

3.4.   Promiscuous EHs are more thermostable

Previous studies indicated that thermodynamically more thermostable proteins frequently have
a higher structural stability (45, 48). In order to investigate if promiscuous EHs, which were
predicted to be less flexible **(see section 3.3)**, are also more thermostable, CD spectroscopy was
applied to determine the melting temperature T_d of the EHs **(see section 2.8)**. Note that only if
the unfolding of a protein is reversible, CD spectroscopy provides true thermodynamic
properties (54). However, even if the unfolding is irreversible, because the protein aggregates
at high temperatures, the method can still give information about relative stabilities (54). Hence,
to reduce the potential impact of different aggregation kinetics of structurally different proteins,
we applied CD spectroscopy to one F_{EH} family only. In particular, we used F_{IV} because it is the
largest F_{EH} **(Table S7)**.

Exemplarily, a CD spectrum for T_d determination is shown for EH001 **(Figure 3A)**; for each
EH, T_d determination was performed in triplicates with STD < 0.62 K. T_d and P_{EH} yield a fair
and significant correlation ($R^2 = 0.32$, $p = 0.033$, **Figure 3B**; a similarly fair and significant
correlation is found if the data point with highest T_d is omitted ($R^2 = 0.45$, $p = 0.016$)).

To conclude, promiscuous EHs are not only globally less flexible but also more thermostable.

3.5.   Promiscuous EHs have less flexible catalytically active residues

The good correlation of P_{EH} and T_p encouraged us to investigate if local flexibility
characteristics of CARs will provide an even better predictor of EH promiscuity. We thus
computed $Flex_{CAR}$ for the *flexibility data set*, i.e., the stability of rigid contacts between CARs
and other residues that are at most 5 Å apart from each other, based on the *local* index $rc_{ij,neighbor}$
(see section 2.5). For each EH, $Flex_{CAR}$ was averaged over five ensembles **(see sections 2.3**
and 2.4), resulting in SEM < 0.06 kcal mol^{-1} **(Figure S10A, Tables 1 and 2)**.

$Flex_{CAR}$ and P_{EH} of the *flexibility data set* yield a good and significant correlation ($R^2 = 0.51$, p
$= 1.7*10^{-6}$) **(Figure S10A)**. To validate again the consistency of our approach, we considered
EHs with known and unknown crystal structures separately. In both cases, good and significant
correlations between $Flex_{CAR}$ and P_{EH} were found (known crystal structures: $R^2 = 0.63$, $p =$
$3.7*10^{-3}$; unknown crystal structures: $R^2 = 0.47$, $p = 2.2*10^{-4}$), again lending support to the
quality of comparative models predicted with TopModel. Hence, EHs with high P_{EH} have low
$Flex_{CAR}$ and *vice versa*, i.e., promiscuous EHs have less flexible CARs. Exemplarily, this is
detailed for EHs of the *representative dataset* with known crystal structures and lowest (EH115)
or highest P_{EH} (EH001), which showed $Flex_{CAR}$ of -0.74 kcal mol^{-1} and -1.91 kcal mol^{-1}, with
unknown crystal structures and lowest (EH127) or highest P_{EH} (EH005), which showed $Flex_{CAR}$
of -1.10 kcal mol^{-1} and -1.86 kcal mol^{-1}, and CalA and CalB, which showed $Flex_{CAR}$ of -1.31
kcal mol^{-1} and -1.95 kcal mol^{-1} **(Figure S10A, Tables S1 and S2)**. The differences in local
structural stability of these EHs are illustrated by rigid contacts between CARs and other
residues that are at most 5 Å apart from each other **(Figure S10B-D)**: promiscuous EHs are
locally more structurally stable as indicated by more stable rigid contacts.

Finally, we exemplarily computed $RMSF_{CAR}$, a measure for the mobility of a protein's CARs
(see section 2.6), across the ensembles of EHs from the *representative data set*. Averaged
$RMSF_{CAR}$ and P_{EH} correlate worse ($R^2 = 0.74$, $p = 0.029$) **(Figure S11A, Table S6)** than $Flex_{CAR}$
and P_{EH} ($R^2 = 0.92$, $p = 2.4*10^{-3}$) **(Figure S11B, Table S6)**, paralleling the above results for
the global measures. Still, again, as promiscuous EHs have less mobile CARs, the same trend
is obtained as in the case of the flexibility analysis.

To conclude, a good and significant correlation between $Flex_{CAR}$ and P_{EH} was found for the
flexibility data set ($R^2 = 0.51$, $p = 1.7*10^{-6}$). Hence, promiscuous EHs have less flexible CARs.
$RMSF_{CAR}$ is less predictive for P_{EH}, although again promiscuous EHs are characterized by less
mobile CARs, mutually confirming either result.

3.6. Promiscuous EHs have an increased specific activity

In the study by Martínez-Martínez *et al.* (19), the *experimental data set* was screened against 96 esters in a kinetic pH indicator assay **(see section 2.1)**. Besides the average specific activity $Act_{average}$ given in U / (g wet cells), also the average maximum specific activity Act_{max} was determined. Motivated by the reactivity-selectivity principle (RSP) initially introduced for organic chemistry reactions (55), which states that a more reactive chemical compound is less selective in chemical reactions, we intended to probe if P_{EH} is related to Act_{max}. For this, we established an approximate linear free-energy relationship (LFER) (56) by relating $\log(Act_{max})$ and $\log(P_{EH})$ **(Figure S12, Table S8)**. In this analysis, the CalA and CalB preparations were excluded because Act_{max} was given in U / (g total protein) there.

$\log(Act_{max})$ and $\log(P_{EH})$ of the *experimental data set* yield a good and significant correlation ($R^2 = 0.50$, $p = 4.6*10^{-23}$) **(Figure S12A)**. Likewise, $\log(Act_{max})$ and $\log(P_{EH})$ of the *flexibility data set* yield a fair and significant correlation ($R^2 = 0.22$, $p = 0.6*10^{-2}$) **(Figure S12B)**. To validate whether the same trend emerges for EHs with known and unknown crystal structures, we considered both types of EHs separately. In both cases, fair and significant correlations between $\log(Act_{max})$ and $\log(P_{EH})$ were found (known crystal structures: $R^2 = 0.34$, $p = 0.099$; unknown crystal structures: $R^2 = 0.23$, $p = 0.019$).

To conclude, good to fair and significant correlations between $\log(Act_{max})$ and $\log(P_{EH})$ of the *experimental data set* ($R^2 = 0.50$, $p = 4.6*10^{-23}$) and the *flexibility data set* ($R^2 = 0.22$, $p = 0.6*10^{-2}$) were found. Hence, promiscuous EHs have higher maximum specific activities.

3.7. Specific EHs prefer to hydrolyze large and flexible esters

Next, we investigated, which of the 96 esters was preferentially hydrolyzed by EHs with different P_{EH}. As a criterion, we chose the number of freely rotatable bonds of an ester, TA **(see section 2.7)**. We did so because TA is a combined measure for an ester's size and conformational dynamics (57). To account for the uneven distribution of esters in our data set with respect to TA, we calculated $Norm_{ester}(TA)$, i.e., the number of hydrolyzed esters with a specific TA ($Ester_{hydrolysed}(TA)$) divided by the total number of esters in the data set with this specific TA ($Ester_{library}(TA)$) **(see section 2.7) (Eq. 1)**.

According to TA, the esters were classified into 17 groups that ranged from small esters with

no rotatable bond to large esters with 56 rotatable bonds **(Figure 4, Table S9)**. Esters with three

(24% of the ester library) and four (16% of the ester library) rotatable bonds are most frequent.

The analysis of the *experimental data set* revealed that promiscuous EHs have high $Norm_{ester}$

values irrespective of TA, i.e., promiscuous EHs accept a large variety of esters with different

sizes and degrees of conformational dynamics **(Figure 4A, Table S9)**. In contrast, specific EHs

only have high $Norm_{ester}$ values regarding esters with high TA, i.e., specific EHs preferentially

hydrolyze (very) large and flexible esters **(Figure 4A, Table S9)**. The same tendency was

observed for the *flexibility data set* **(Figure 4B, Table S9)**.

To conclude, promiscuous EHs accept a large variety of esters with different sizes and degrees

of conformational dynamics whereas specific EHs preferentially hydrolyze (very) large and

flexible esters.

4. Discussion

The main outcomes of our analyses are I) that promiscuous EHs are significantly globally less flexible and have less flexible catalytically active residues than specific ones, II) that promiscuous EHs are significantly more thermostable, III) that promiscuous EHs have a significantly increased specific activity, and IV) that specific EHs prefer to hydrolyze large and flexible esters.

We established these relations using one of the still few experimental large-scale datasets where a diverse set of EHs was functionally assessed against a customized library of dissimilar esters (19). Functional promiscuity may arise from several conditions, including the environment of the enzyme or the concentration of a substrate, which may complicate the analysis of the molecular mechanism underlying promiscuity (5). Still, functional promiscuity ultimately is a result of recognition promiscuity (5); here, we therefore focused on substrate promiscuity (18), i.e., an enzyme carries out its typical catalytic function using non-canonical substrates, in that experimental conditions had been kept constant for the assessment of the different esterase/ester combinations (19). Almost all of the EHs were unambiguously assigned to one of the F_{EH} of the Arpigny and Jaeger classification, which is based mainly on a comparison of amino acid sequences (53). Except for classes with a few members only (cyclase-like EHs and the yeast family), all other classes cover at least two of the three P_{EH} ranges such that P_{EH} cannot be assigned based on the EH's class affiliation **(Figure S13, Table S10)** and, hence, amino acid sequence information. Even family F_{IV}, which contains a higher proportion of substrate-promiscuous EHs, also contains EHs with a small substrate range.

For scrutinizing the mechanism underlying esterase promiscuity at the atomistic level, we needed to apply comparative models of EHs, since only for ~7% of the experimentally assessed EHs crystal structures were available. Restricting the generation of esterase models to sequence identities ≥ 25% with respect to available targets yielded generally good structural models both globally and locally, as also validated against cases where crystal structures are known. Throughout our study, we probed for the consistency of our analyses between subsets of EHs for which either crystal structures are known or not; we only found quantitative differences, but no qualitative ones. One of the reasons is likely that rigidity analyses were based on structural ensembles generated by multiple and µs-long MD simulations, which has been shown to improve both global and local protein structure (58, 59) to the level of approaching experimental accuracy (60) and markedly increases the robustness of the results (42). We furthermore showed that results are consistent irrespective of whether EH flexibility

characteristics were assessed globally or only for CARs, and that mobility characteristics computed directly from MD trajectories show the same trend, although the correlation with P_{EH} is insignificant. Finally, we used experimental melting temperatures of EHs as indicators for enzyme flexibility (45, 48), which yielded the same relation with P_{EH} as computed flexibility characteristics. The partial use of comparative models rather than crystal structures throughout this study may lead to concern. Yet, our consistent and robust findings indicate that when applying this workflow to novel EHs, including to those for which no crystal structure exists but a structural homolog with a sequence identity $\geq$ 25%, it should be possible to discover enzymes with 'sufficient' substrate promiscuity to serve as a starting point for further exploration in biotechnology and synthetic organic chemistry. In that respect, the flexibility characteristics of EHs analyzed here have a notably stronger predictive power than the active site effective volume introduced earlier (19) (**Figure S14, Tables S11 and S12**).

The finding that promiscuous EHs are significantly globally *less* flexible and have *less* flexible catalytically active residues than specific EHs is in stark contrast to the general view of the role of structural flexibility for promiscuity (4, 5): Besides the examples of CYP and β-lactamase mentioned above, the possibility of dynamically restructuring active sites has also been recognized for other systems as underlying their promiscuity (61-64). Finally, interactions between antibodies and antigens are likely the quintessential example of the canonical relationship between flexibility and binding promiscuity: As antibodies mature to become more specific, their flexibility is decreased (5).

It has been recognized that conformational changes may not always be necessary for promiscuity if a variety of substrates can be bound by partial recognition or the presence of multiple binding sites (5). However, these cases do not seem to be relevant reasons for EH promiscuity because partial recognition often is associated with catalytic inefficiency (1), which is contrary to our observation that P_{EH} correlates with EH activity, and the presence of multiple binding sites that could give rise to promiscuity is controverted by the finding that promiscuous EHs have large active site effective volumes (19), i.e., large pockets with few subpockets. Inversely, our findings of rigid promiscuous EHs may be consistent with the idea that multiple ligands can be accommodated in a single site by exploiting diverse interacting residues (**Figure 5**).

Our results as to *specific but flexible* EHs may be reconciled with a model according to which conformational changes may have been selected in EH evolution for their ability to enhance specificity in recognition (**Figure 5**), resulting in what has been termed conformational

proofreading (16). In the case of specific EHs, flexibility may help to overcome a structural

mismatch between the enzyme and its substrate existing when both are in their ground states,

that way enhancing recognition specificity. This view is corroborated by our finding that

specific EHs prefer to hydrolyze large and flexible substrates: Larger substrates can form more

interactions with the enzyme, that way helping to overcome the deformation energy required

by the enzyme to optimizing the correct binding probability over the incorrect one; flexible

substrates can tolerate higher strains and thus can be expected to participate in more binding

events (65, 66) **(Figure 5)**. In that respect, the relation between structural flexibility of EHs and

promiscuity found here is more causative than that between active site effective volume and

promiscuity (19), because small active site effective volumes found for specific EHs cannot

rationalize why specific EHs prefer to hydrolyze large and flexible substrates.

In summary, the combined large-scale analysis of experimental EH promiscuity and computed

EH flexibility reveals that promiscuous EHs are significantly less flexible than specific ones.

This result is counterintuitive at first but may be reconciled with a model according to which

multiple ligands can be accommodated in a single active site of promiscuous EHs by exploiting

diverse interacting residues, whereas structural flexibility in the case of specific EHs serves for

conformational proofreading. Our results furthermore signify that EH sequence space, charted,

e.g., by (meta)genomics studies, can be screened by rigidity analyses for promiscuous EHs that

may serve as starting points for further exploration in biotechnology and synthetic organic

chemistry.

5. Acknowledgements

CN is funded through a grant ("Vernetzungsdoktorand") provided by the Forschungszentrum Jülich. Parts of the study were supported by the German Federal Ministry of Education and Research (BMBF) through funding number 031B0837A "LipoBiocat" to HG and KEJ and through funding number 031L0182 "InCelluloProtStruct" to HG, the German Research Foundation (DFG) through funding no. INST 208/704-1 FUGG to HG and INST 208/654-1 FUGG to KEJ, as well as the state of North Rhine Westphalia (NRW) and the European Regional Development Fund (EFRE) through funding no. 34-EFRE-0300096 "CLIB-Kompetenzzentrum Biotechnologie (CKB)" to HG. HG is grateful for computational support and infrastructure provided by the "Zentrum für Informations- und Medientechnologie" (ZIM) at the Heinrich Heine University Düsseldorf. HG gratefully acknowledges the computing time granted by the John von Neumann Institute for Computing (NIC) and provided on the supercomputer JUWELS at Jülich Supercomputing Centre (JSC) (user IDs: HKF7; protil (project ID: 15956)) (67). HG is grateful to OpenEye for an academic license. MF acknowledges the grant 'INMARE' from the European Union's Horizon 2020 (grant agreement no. 634486) and BIO2017-85522-R from the Ministerio de Ciencia, Innovación y Universidades (MCIU), Agencia Estatal de Investigación (AEI), Fondo Europeo de Desarrollo Regional (FEDER) and European Union (EU). CC thanks the Ministerio de Economía y Competitividad and FEDER for a PhD fellowship (Grant BES-2015-073829). The authors are grateful to David Almendral and Ruth Matesanz for their support of CD analysis.

6. Authors contributions

HG and KEJ conceived the study. CN analyzed the experimental data, performed structure prediction, MD simulations and CNA computations, analyzed the computational data, and wrote the manuscript together with HG. DM initially contributed to the structure prediction. BD performed similarity analysis of esters. MF and CC measured and analyzed melting temperatures and determined kinetic parameters. HG supervised and managed the project. All authors reviewed and approved the manuscript.

7. Conflict of interest

The authors declare no financial and non-financial competing interests.

8. Figures

Figure 1: Comparative modeling of EHs. (A) Based on sequence data provided by a large-scale study from Martínez-Martínez *et al.* (19), comparative models were generated for 35 EHs with known (left, 11 EHs) and unknown (right, 24 EHs) crystal structures using TopModel (26). These EHs constitute the *flexibility data set*. The EHs vary in P_{EH} (left ordinate, bars) and global TopScores (right ordinate, diamonds). Six EHs were selected as representatives of the *flexibility data set* (termed *representative data set*) as indicated by magenta arrows. The quality of the comparative models of **(B)** EHs with known crystal structures and lowest (EH115) or highest P_{EH} (EH001), **(C)** EHs with unknown crystal structures and lowest (EH127) or highest P_{EH} (EH005), and **(D)** commercial EHs with highest (CalA) or lowest P_{EH} (CalB) was evaluated by TopScore (31). For each comparative model the residue-wise TopScore is shown: A good (bad) homology model shows a low (high) residue-wise TopScore (see color scale at the bottom). Insets depict CARs (spheres) within an EH. For clarity the position of CARs is indicated by magenta stars.

Figure 2: Correlation of T_p *versus* P_{EH}. (A) Correlation between predicted T_p based on the global index H_{type2} and P_{EH} for the *flexibility data set*. Data points colored grey (black) represent comparative models of EHs with (un)known crystal structures. The *representative data set* is indicated by magenta crosses. Error bars show the SEM over five independent MD simulations of 1 µs length each. Rigid cluster decomposition at 332 K during the thermal unfolding simulation of **(B)** EHs with known crystal structures and lowest (EH115) or highest P_{EH} (EH001), **(C)** EHs with unknown crystal structures and lowest (EH127) or highest P_{EH} (EH005), and **(D)** commercial EHs with lowest (CalA) or highest P_{EH} (CalB). Rigid clusters are represented as uniformly colored blue, green, pink, cyan, and magenta bodies in the descending order of their sizes.

Figure 3: Determination of T_d via CD spectroscopy. (A) Exemplary CD spectrum of EH001. The ellipticity changes in mdeg at 220 nm was plotted against the temperature, resulting in a sigmoidal curve. The inflection point was used to obtain the T_d value (dotted line). **(B)** Correlation between T_d and P_{EH} for 12 EHs of F_{IV}.

Figure 4: Relation between the number of esters' TA and P_{EH}. Relation between $Ester_{norm}$, i.e., the relative proportion of the number of hydrolyzed esters with a specific TA, and P_{EH} of **(A)** the *experimental data set* and **(B)** the *flexibility data set* containing EHs with known crystal structures (left), EHs with unknown crystal structures (right), and EHs constituting the *representative data set* (indicated by magenta arrows). TA was calculated based on SMILES codes of 96 esters provided by Martínez-Martínez *et al.* (19). A blue (red) color indicates that the EH hydrolyzes many (few) esters with a specific TA relative to the total number of esters in the data set with this specific TA (see color scale on the right); the total number of esters with a specific TA is given in brackets on the y-axis. P_{EH} is defined as *low* if the EH hydrolyzes ≤ 9 esters, as *moderate* if the EH hydrolyzes between 10 and 29 esters, and as *high* if the EH hydrolyzes ≥ 30 esters.

Figure 5: Mechanistic model of EH flexibility, ligand size and conformational dynamics affecting P_{EH}. Impact of esters with **(A)** many or **(B)** few TA on specific, and hence more flexible (left), and promiscuous, and hence more rigid (right), EHs. Ligand parts connected by TA are represented as blue circles. Specific EHs and large ligands with many TA can mutually adapt (panel **A**, left), and promiscuous EH can bind large ligands (panel **A**, right) and small ligands (panel **B**, right) exploiting different interaction partners. Small (and/or rigid) ligands are not able to lead to a structural adaptation of specific EHs (panel **B**, left), though, resulting in conformational proofreading. The red bars indicate the flexibility of the EHs. A green tick (red cross) indicates that ester cleavage is (not) catalyzed.

9. References

1.   Copley SD (2017) Shining a light on enzyme promiscuity. *Curr. Opin. Struct. Biol.* 47:167-175.
2.   Chen R, *et al.* (2017) Molecular insights into the enzyme promiscuity of an aromatic prenyltransferase. *Nat. Chem. Biol.* 13(2):226-234.
3.   Huang H, *et al.* (2015) Panoramic view of a superfamily of phosphatases through substrate profiling. *Proc. Natl. Acad. Sci. U.S.A.* 112(16):E1974-E1983.
4.   Khersonsky O & Tawfik DS (2010) Enzyme promiscuity: a mechanistic and evolutionary perspective. *Annu. Rev. Biochem.* 79:471-505.
5.   Nobeli I, Favia AD, & Thornton JM (2009) Protein promiscuity and its implications for biotechnology. *Nat. Biotechnol.* 27(2):157-167.
6.   Eisenmesser EZ, Bosco DA, Akke M, & Kern D (2002) Enzyme dynamics during catalysis. *Science* 295(5559):1520-1523.
7.   Henzler-Wildman KA, *et al.* (2007) A hierarchy of timescales in protein dynamics is linked to enzyme catalysis. *Nature* 450(7171):913-916.
8.   Holliday MJ, Camilloni C, Armstrong GS, Vendruscolo M, & Eisenmesser EZ (2017) Networks of dynamic allostery regulate enzyme function. *Structure* 25(2):276-286.
9.   Goodey NM & Benkovic SJ (2008) Allosteric regulation and catalysis emerge via a common route. *Nat. Chem. Biol.* 4(8):474-482.
10.   Saavedra HG, Wrabl JO, Anderson JA, Li J, & Hilser VJ (2018) Dynamic allostery can drive cold adaptation in enzymes. *Nature* 558(7709):324-328.
11.   Tzeng SR & Kalodimos CG (2011) Protein dynamics and allostery: an NMR view. *Curr. Opin. Struct. Biol.* 21(1):62-67.
12.   Skopalík J, Anzenbacher P, & Otyepka M (2008) Flexibility of human cytochromes P450: molecular dynamics reveals differences between CYPs 3A4, 2C9, and 2A6, which correlate with their substrate preferences. *J. Phys. Chem. B.* 112(27):8165-8173.
13.   Hendrychová T, *et al.* (2011) Flexibility of human cytochrome P450 enzymes: molecular dynamics and spectroscopy reveal important function-related variations. *Biochim. Biophys. Acta.* 1814(1):58-68.
14.   Ekroos M & Sjögren T (2006) Structural basis for ligand promiscuity in cytochrome P450 3A4. *Proc. Natl. Acad. Sci. U.S.A.* 103(37):13682-13687.
15.   Zou T, Risso VA, Gavira JA, Sanchez-Ruiz JM, & Ozkan SB (2015) Evolution of conformational dynamics determines the conversion of a promiscuous generalist into a specialist enzyme. *Mol. Biol. Evol.* 32(1):132-143.
16.   Savir Y & Tlusty T (2007) Conformational proofreading: the impact of conformational changes on the specificity of molecular recognition. *PloS One* 2(5):e468.
17.   Ferrer M, *et al.* (2016) Estimating the success of enzyme bioprospecting through metagenomics: current status and future trends. *Microb. Biotechnol.* 9(1):22-34.
18.   Hult K & Berglund P (2007) Enzyme promiscuity: mechanism and applications. *Trends. Biotechnol.* 25(5):231-238.
19.   Martinez-Martinez M, *et al.* (2017) Determinants and prediction of esterase substrate promiscuity patterns. *ACS Chem. Biol.* 13(1):225-234.
20.   Radestock S & Gohlke H (2008) Exploiting the link between protein rigidity and thermostability for data-driven protein engineering. *Eng. Life Sci.* 8(5):507-522.
21.   Pfleger C, Rathi PC, Klein DL, Radestock S, & Gohlke H (2013) Constraint Network Analysis (CNA): a Python software package for efficiently linking biomacromolecular structure, flexibility,(thermo-) stability, and function. *J. Chem. Inf. Model.* 53:1007-1015.

22. Krüger DM, Rathi PC, Pfleger C, & Gohlke H (2013) CNA web server: rigidity theory-based thermal unfolding simulations of proteins for linking structure,(thermo-) stability, and function. *Nucleic Acids Res.* 41(W1):W340-W348.

23. Hermans SM, Pfleger C, Nutschel C, Hanke CA, & Gohlke H (2017) Rigidity theory for biomolecules: concepts, software, and applications. *Comput. Mol. Sci.* 7(4):e1311.

24. de Godoy Daiha K, Angeli R, de Oliveira SD, & Almeida RV (2015) Are lipases still important biocatalysts? A study of scientific publications and patents for technological forecasting. *PloS One* 10(6):e0131624.

25. Ferrer M, *et al.* (2015) Biodiversity for biocatalysis: a review of the α/β-hydrolase fold superfamily of esterases-lipases discovered in metagenomes. *Biocatal. Biotranfor.* 33(5-6):235-249.

26. Mulnaes D, *et al.* (2020) TopModel: Template-based protein structure prediction at low sequence identity using top-down consensus and deep neural networks. *J. Chem. Theory. Comput.* 16:1953-1967.

27. Gohlke H, *et al.* (2013) Binding region of alanopine dehydrogenase predicted by unbiased molecular dynamics simulations of ligand diffusion. *J. Chem. Inf. Model.* 53(10):2493-2498.

28. Widderich N, *et al.* (2014) Molecular dynamics simulations and structure-guided mutagenesis provide insight into the architecture of the catalytic core of the ectoine hydroxylase. *J. Mol. Biol.* 426(3):586-600.

29. Zhang Z, *et al.* (2016) Determinants of FIV and HIV Vif sensitivity of feline APOBEC3 restriction factors. *Retrovirology* 13(1):46.

30. Milić D, *et al.* (2018) Recognition motif and mechanism of ripening inhibitory peptides in plant hormone receptor ETR1. *Sci. Rep.* 8(1):3890.

31. Mulnaes D & Gohlke H (2018) TopScore: Using Deep Neural Networks and Large Diverse Data Sets for Accurate Protein Model Quality Assessment. *J. Chem. Theory Comput.* 14(11):6117-6126.

32. Chovancova E, *et al.* (2012) CAVER 3.0: a tool for the analysis of transport pathways in dynamic protein structures. *PLoS Comput. Biol.* 8(10):e1002708.

33. Salomon-Ferrer R, Götz AW, Poole D, Le Grand S, & Walker RC (2013) Routine microsecond molecular dynamics simulations with AMBER on GPUs. 2. Explicit solvent particle mesh Ewald. *J. Chem. Theory Comput.* 9(9):3878-3888.

34. Case DA, *et al.* (2019) AMBER 2019University of California, San Francisco).

35. Hopkins CW, Le Grand S, Walker RC, & Roitberg AE (2015) Long-time-step molecular dynamics through hydrogen mass repartitioning. *J. Chem. Theory Comput.* 11(4):1864-1874.

36. Jacobs DJ, Rader AJ, Kuhn LA, & Thorpe MF (2001) Protein flexibility predictions using graph theory. *Proteins* 44(2):150-165.

37. Hespenheide B, Jacobs D, & Thorpe M (2004) Structural rigidity in the capsid assembly of cowpea chlorotic mottle virus. *J. Condens. Matter Phys.* 16(44):S5055.

38. Jacobs DJ & Thorpe MF (1995) Generic rigidity percolation: the pebble game. *Phys. Rev. Lett.* 75(22):4051.

39. Rader AJ, Hespenheide BM, Kuhn LA, & Thorpe MF (2002) Protein unfolding: rigidity lost. *Proc. Natl. Acad. Sci. U.S.A.* 99(6):3540-3545.

40. Livesay DR & Jacobs DJ (2006) Conserved quantitative stability/flexibility relationships (QSFR) in an orthologous RNase H pair. *Proteins* 62(1):130-143.

41. Radestock S & Gohlke H (2011) Protein rigidity and thermophilic adaptation. *Proteins* 79(4):1089-1108.

42. Pfleger C & Gohlke H (2013) Efficient and robust analysis of biomacromolecular flexibility using ensembles of network topologies based on fuzzy noncovalent constraints. *Structure* 21(10):1725-1734.

43. Pfleger C, Radestock S, Schmidt E, & Gohlke H (2013) Global and local indices for characterizing biomolecular flexibility and rigidity. *J. Comput. Chem.* 34(3):220-233.

44. Rathi PC, Radestock S, & Gohlke H (2012) Thermostabilizing mutations preferentially occur at structural weak spots with a high mutation ratio. *J. Biotechnol.* 159(3):135-144.

45. Rathi PC, Jaeger K-E, & Gohlke H (2015) Structural rigidity and protein thermostability in variants of lipase A from Bacillus subtilis. *PloS One* 10(7):e0130289.

46. Dick M*, et al.* (2016) Trading off stability against activity in extremophilic aldolases. *Sci. Rep.* 6:17908.

47. Rathi PC, Fulton A, Jaeger K-E, & Gohlke H (2016) Application of rigidity theory to the thermostabilization of lipase A from Bacillus subtilis. *PLoS Comput. Biol.* 12(3):e1004754.

48. Nutschel C*, et al.* (2020) Systematically scrutinizing the impact of substitution sites on thermostability and detergent tolerance for Bacillus subtilis lipase A. *J. Chem. Inf. Model.* 60:1568-1584.

49. Gohlke H, Kuhn LA, & Case DA (2004) Change in protein flexibility upon complex formation: analysis of Ras-Raf using molecular dynamics and a molecular framework approach. *Proteins* 56(2):322-337.

50. Eisenthal R, Danson MJ, & Hough DW (2007) Catalytic efficiency and kcat/KM: a useful comparator? *Trends Biotechnol.* 25(6):247-249.

51. Nath A & Atkins WM (2008) A quantitative index of substrate promiscuity. *Biochemistry* 47(1):157-166.

52. Hawkins PC, Skillman AG, & Nicholls A (2007) Comparison of shape-matching and docking as virtual screening tools. *J. Med. Chem.* 50(1):74-82.

53. Arpigny JL & Jaeger K-E (1999) Bacterial lipolytic enzymes: classification and properties. *Biochem. J.* 343(1):177-183.

54. Greenfield NJ (2006) Using circular dichroism collected as a function of temperature to determine the thermodynamics of protein unfolding and binding interactions. *Nat. Protoc.* 1(6):2527-2535.

55. Mayr H & Ofial AR (2006) The reactivity–selectivity principle: an imperishable myth in organic chemistry. *Angew. Chem. Int. Ed. Engl.* 45(12):1844-1854.

56. Wells PR (1963) Linear Free Energy Relationships. *Chem. Rev.* 63(2):171-219.

57. Böhm H-J (1994) The development of a simple empirical scoring function to estimate the binding constant for a protein-ligand complex of known three-dimensional structure. *J. Comput. Aided Mol. Des.* 8(3):243-256.

58. Feig M (2017) Computational protein structure refinement: Almost there, yet still so far to go. *Wiley Interdiscip Rev Comput Mol Sci* 7(3).

59. Heo L & Feig M (2018) Experimental accuracy in protein structure refinement via molecular dynamics simulations. *Proc. Natl. Acad. Sci. U. S. A.* 115(52):13276-13281.

60. Heo L, Arbour CF, & Feig M (2019) Driven to near-experimental accuracy by refinement via molecular dynamics simulations. *Proteins* 87(12):1263-1275.

61. Seibert CM & Raushel FM (2005) Structural and catalytic diversity within the amidohydrolase superfamily. *Biochemistry* 44(17):6383-6391.

62. Oppermann U*, et al.* (2003) Short-chain dehydrogenases/reductases (SDR): the 2002 update. *Chem. Biol. Interact.* 143-144:247-253.

63. Fushinobu S, Nishimasu H, Hattori D, Song H-J, & Wakagi T (2011) Structural basis for the bifunctionality of fructose-1, 6-bisphosphate aldolase/phosphatase. *Nature* 478(7370):538-541.

64. Du J, Say RF, Lü W, Fuchs G, & Einsle O (2011) Active-site remodelling in the bifunctional fructose-1, 6-bisphosphate aldolase/phosphatase. *Nature* 478(7370):534-537.

65. Stockwell GR & Thornton JM (2006) Conformational diversity of ligands bound to proteins. *J. Mol. Biol.* 356(4):928-944.

66. Perola E & Charifson PS (2004) Conformational analysis of drug-like molecules bound to proteins: an extensive study of ligand reorganization upon binding. *J. Med. Chem.* 47(10):2499-2510.

67. Krause D (2019) JUWELS: Modular Tier-0/1 Supercomputer at the Jülich Supercomputing Centre. *JLSRF* 5:A135.