1

Promiscuous esterases counterintuitively are less flexible than specific ones

Christina Nutschel[1], Daniel Mulnaes[2], Cristina Coscolín[3], Manuel Ferrer[3], Karl-Erich Jaeger[4,5], Holger Gohlke[1,2,*]

[1] John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC) and Institute of Biological Information Processing (IBI-7: Structural Biochemistry), Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

[2] Institute for Pharmaceutical and Medicinal Chemistry, Heinrich Heine University Düsseldorf, 40225 Düsseldorf, Germany

[3] Institute of Catalysis, Consejo Superior de Investigaciones Científicas, 28049 Madrid, Spain

[4] Institute of Molecular Enzyme Technology, Heinrich Heine University Düsseldorf, 52425 Jülich, Germany

[5] Institute of Bio- and Geosciences IBG-1: Biotechnology, Forschungszentrum Jülich GmbH, 52425 Jülich, Germany

Running title: Flexibility and promiscuity of esterases

Keywords: Esterase promiscuity, structural flexibility, thermostability, Constraint Network Analysis, TopModel, TopScore, conformational proofreading

*Corresponding author:

John von Neumann Institute for Computing (NIC), Jülich Supercomputing Centre (JSC), and

Institute of Biological Information Processing (IBI-7: Structural Biochemistry)

Forschungszentrum Jülich GmbH

Wilhelm-Johnen-Straße

52425 Jülich

Germany

Email: gohlke@uni-duesseldorf.de or h.gohlke@fz-juelich.de

Abstract

Understanding mechanisms of promiscuity is increasingly important from a fundamental and application point of view. As to enzyme structural dynamics, more promiscuous enzymes generally have been recognized to also be more flexible. However, examples for the opposite received much less attention. Here, we exploit comprehensive experimental information on the substrate promiscuity of 147 esterases tested against 96 esters together with computationally efficient rigidity analyses to understand the molecular origin of the observed promiscuity range. Unexpectedly, our data reveal that promiscuous esterases are significantly less flexible than specific ones, are significantly more thermostable, and have a significantly increased specific activity. These results may be reconciled with a model according to which structural flexibility in the case of specific esterases serves for conformational proofreading. Our results signify that esterase sequence space can be screened by rigidity analyses for promiscuous esterases as starting points for further exploration in biotechnology and synthetic chemistry.

1. Introduction

Enzymes involved in primary metabolism typically exquisitely discriminate against other metabolites. Yet, evolution of specificity is only pushed by nature to the point at which 'unauthorized' reactions do not impair the fitness of the organism (1). As a result, the universe of promiscuous activities available in nature has been suggested to be enormous (2, 3). Understanding mechanisms of promiscuity has thus become increasingly important for the fundamental understanding of molecular recognition and how enzyme function has evolved over time(4) but also to optimize enzyme engineering applications (5). A particular challenge in the latter case is the ability to discover a suitable enzyme with 'sufficient' promiscuous activity to serve as a starting point for further exploration (1).

Enzyme structural dynamics, besides its role in catalysis (6, 7) and allosteric regulation (8-11), has been recognized as likely the single most important mechanism by which promiscuity can be achieved (5). Prominent examples are human cytochrome P450 (CYP) enzymes, for which crystallographic studies and molecular simulations demonstrated that more promiscuous CYPs show larger structural plasticity and mobility (12-14), or TEM-1 β-lactamase and a resurrected progenitor, for which molecular simulations show that the pocket of the ancestral, and more promiscuous, enzyme fluctuates to a greater extent (15). However, examples for the opposite, i.e., conformational changes selected in evolution such that they enhance specificity in molecular recognition (16), have received much less attention in the context of enzyme promiscuity.

A clear limitation for scrutinizing the link between enzyme structural dynamics and substrate promiscuity is the general lack of large-scale data on one enzyme (super)family tested against a multitude of ligands (17) (cf. ref. (1) for notable exceptions). Likewise, acquiring information on enzyme dynamics at the atomistic level by experimental techniques or classical molecular dynamics (MD) simulations is burdensome. Here, we exploit comprehensive experimental information on the substrate promiscuity (18) of esterases (abbreviated EHs, for "Ester Hydrolases") (19) together with computationally efficient rigidity analyses (20-23) of comparative models of EHs to understand the molecular origin of the observed promiscuity range. Enzyme rigidity, or its opposite flexibility, are static properties that denote the *impossibility*, or *possibility*, *of motions* in an enzyme under force, without giving information about directions and magnitudes of movements (23). Enzyme flexibility, thus, should not be confused with enzyme mobility, which describes *actual motions* in an enzyme. Rigidity analysis results do not rely on the correct description of the

time-dependency of processes (23), which makes them valuable in cases where timescales
over multiple orders of magnitude may govern such processes, like in enzyme dynamics (6,
7).

In recent years, EHs have obtained much attention in basic research and industrial
applications (24). EHs are widely distributed in nature within microbial communities (at least
one EH is found in each bacterial genome), they have been extensively examined with state-
of-the-art (meta)genomics techniques and investigated by functional screenings compared to
many other classes of enzymes. They also possess outstanding properties in terms of stability,
reactivity, and scalability that make them appropriate biocatalysts to improve
competitiveness, innovation capacity, and sustainability in a modern circular bio-economy
(25). Recently, a large-scale study on substrate promiscuity (P_{EH}, which denotes the number
of esters hydrolyzed by an EH) of 147 phylogenetically, environmentally, and structurally
diverse microbial EHs was described by Ferrer *et al.* (19), in which all EHs were functionally
assessed against a customized library of 96 esters. As to mechanistic understanding, the
authors related P_{EH} to a structural parameter, the active site effective volume. However, the
impact of enzyme flexibility on P_{EH} was not assessed.

In our study, we thus ask the following questions: I) What is the relation between P_{EH} and EH
flexibility? II) Does this relation hold if experimentally determined EH thermostabilities are
used as proxies for enzyme flexibility? III) What is the relation between P_{EH} and EHs'
specific activities? IV) Is there a preference of promiscuous or specific EHs for a particular
type of esters. V) Can this preference be understood with respect to EHs flexibilities?

2. Materials and Methods

2.1. Definition of data sets

The present study builds on the study from Ferrer *et al.* (19). In order to assess P_{EH}, the authors experimentally investigated 147 phylogenetically, environmentally, and structurally diverse microbial EHs (termed *experimental data set*) against a customized library of 96 different esters. Two commercial lipases, which have found wide biotechnological applications, CalA and CalB from *Pseudozyma aphidis* (formerly *Candida antarctica*), were included for comparison. For details on determining and classifying P_{EH}, see **Supplemental Materials and Methods**.

As our computational approach involves extensive molecular dynamics (MD) simulations for generating conformational ensembles **(see section 2.3)**, we selected 35 EHs from the *volume data set* (termed *flexibility data set*) for comparative modeling **(see section 2.2)**. The criteria for choosing EHs of the *flexibility data set* are explained in **section 3.1**.

2.2. Comparative modelling and validations of the *flexibility data set*

Comparative models of the *flexibility data set* **(see section 2.1)** were generated using our in-house structure prediction meta-tool TopModel (26) that has been successfully applied in previous studies (27-30). TopModel uses multiple state-of-the-art threading and sequence/structure alignment tools to generate a large ensemble of models from different pairwise and multiple alignments of the top five highest ranked template structures. The TopModel software is available at https://cpclab.uni-duesseldorf.de/index.php/Software.

The quality of the homology models was assessed by our meta Model Quality Assessment Program (meta-MQAP) TopScore (31). TopScore uses deep neural networks (DNN) to combine scores from 15 different primary MQAP to predict accurate residue-wise and whole-protein error estimates. For details on model quality assessment by TopScore and validation, see **Supplemental Materials and Methods**.

To test whether CARs of the homology models are accessible for substrates, we applied the CAVER 3.0.3 PyMOL Plugin (32). Starting points for the computations were defined based on the Cartesian coordinates of the CARs' center of mass (COM). Default values were used for the probe radius (0.9 Å), shell radius (3.0 Å), and shell depth (4.0 Å).

2.3.   Generation of structural ensembles

Structural ensembles of EHs were generated by all-atom MD simulations of in total 5 µs

simulation time per EH. For details on starting structure preparation, parametrization, and

equilibration see **Supplemental Materials and Methods**.

All minimization, equilibration, and production simulations were performed with the

pmemd.cuda module (33) of Amber19 (34). During production simulations, we set the time

step for the integration of Newton's equation of motion to 4 fs following the hydrogen mass

repartitioning strategy (35). Coordinates were stored into a trajectory file every 200 ps. This

resulted in 5000 configurations for each production run that were considered for subsequent

analyses.

2.4.   Constraint Network Analysis

The flexibility analyses were performed with the Constraint Network Analysis (CNA)

software package (version 3.0) (20-23). CNA functions as front- and back-end to the graph

theory-based software Floppy Inclusions and Rigid Substructure Topography (FIRST) (36).

Applying CNA to biomolecules aims at identifying their composition of rigid clusters and

flexible regions, which can aid in the understanding of biomolecular structure, stability, and

function (21-23). As the mechanical heterogeneity of biomolecular structures is intimately

linked to their diverse biological functions, biomolecules generally show a hierarchy of

rigidity and flexibility (20). In CNA, biomolecules are modeled as constraint networks in a

body-and-bar representation, which has been described in detail by Hesphenheide *et al.* (37).

A fast combinatorial algorithm, the *pebble game*, counts the bond rotational degrees of

freedom and floppy modes (internal, independent degrees of freedom) in the constraint

network (38). In order to monitor the hierarchy of rigidity and flexibility of biomolecules,

CNA performs thermal unfolding simulations by consecutively removing non-covalent

constraints (hydrogen bonds, including salt bridges) from a network in increasing order of

their strength (39-41). For details on thermal unfolding simulations, see **Supplemental**

Materials and Methods. To improve the robustness and investigate the statistical

uncertainty, we carried out CNA on ensembles of network topologies (ENT^{MD}) generated

from MD trajectories **(see section 2.3)** (42).

The CNA software is available under academic license at https://cpclab.uni-
duesseldorf.de/index.php/Software and the CNA web server is accessible at https://cpclab.uni-
duesseldorf.de/cna.

2.5.   Local and global indices

From the thermal unfolding simulations, CNA computes a comprehensive set of indices to
quantify biologically relevant characteristics of the protein's stability. *Global* indices are used
for determining the rigidity and flexibility at a macroscopic level; *local* indices determine the
rigidity and flexibility at a microscopic level of bonds (43). The cluster configuration entropy
H_{type2} is a *global* index that has been introduced by Radestock and Gohlke (20). As done
previously, we applied H_{type2} as a measure for global structural stability of proteins (20, 41,
44-48). The stability map rc_{ij} is a *local* index that has been introduced by Radestock and
Gohlke (20). We applied rc_{ij} as a measure for local structural stability of proteins in previous
studies (45, 47, 48). For details on both indices, see **Supplemental Materials and Methods**.

2.6.   Root mean square fluctuations

181 The per-residue root-mean-square fluctuations were calculated for each EH ($RMSF_{EH}$) and for
182 its CARS ($RMSF_{CAR}$) based on the MD trajectories **(see section 2.3)**. Prior to the calculations,
183 the structures of each trajectory were superimposed onto the average structure using the 90%
184 least mobile residues of the respective EHs (49).

2.7.   Torsion angles

For each of the 96 esters, the number of freely rotatable bonds (torsion angles, TA) was
calculated based on the SMILES codes provided by Ferrer *et al.* (19).

To compare how many esters with a specific TA are hydrolyzed by each EH, we calculated
the normalized proportion of ester hydrolysis with a specific TA ($Norm_{ester}$(TA)) as the
number of hydrolyzed esters with a specific TA ($Ester_{hydrolysed}$ (TA)) divided by the total
number of esters in the data set with this specific TA ($Ester_{library}$(TA)) **(Eq. 1)**.

$$Norm_{ester}(TA) \ [\%] = \frac{Ester_{hydrolyzed}(TA)}{Ester_{library}(TA)} * 100\%$$

Eq. 1

2.8.   Circular dichroism spectroscopy

Prior to analyses, soluble His-tagged proteins were produced and purified after binding to a

Ni-NTA His-Bind resin as described by Ferrer *et al.* (19). Circular dichroism (CD) spectra

were acquired between 190 and 270 nm with a Jasco J-720 spectropolarimeter equipped with

a Peltier temperature controller, employing a 0.1 mm cell at 25°C. Spectra were analyzed, and

denaturation temperatures were determined at 220 nm between 10 and 85°C at a rate of 30°C

per hour, in 40 mM (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer, pH

7.0. A protein concentration of 1.0 mg ml^{-1} was used. Denaturation temperatures were

calculated by fitting the ellipticity (mdeg) at 220 nm at each of the different temperatures

using a 5-parameters sigmoid fit with Sigma Plot 13.0.

3. Results

3.1. Definition of data sets

To understand the structural origin of and develop a method to predict P_{EH}, the present study builds on large-scale data from Ferrer *et al.* (19). The authors experimentally investigated P_{EH} of 147 EHs (termed *experimental data set*) **(see section 2.1)**. Additionally, they ranked (classified) P_{EH} of 96 EHs (termed *volume data set*) based on the active site effective volume **(see section 2.1) (Eq. S1)**, which will be used here as a reference to compare the power of P_{EH} predictions.

As our computational approach involves extensive MD simulations for generating conformational ensembles **(see section 2.3)**, we selected 35 EHs from the *volume data set* based on the following criteria; they constitute the *flexibility data set*. I) The data set contains eleven EHs with known crystal structures (including the commercial EHs CalA and CalB) **(Figure 1A, Table S1)** and 24 EHs for which no experimental structure is available but for which comparative models can be generated **(see section 3.2) (Figure 1A, Table S2)**. That way, we can probe to what extent the source of structural information influences the outcome of our results. II) The chosen EHs of the data set show high diversities as to P_{EH} and association to esterase families (F_{EH}, as defined by Arpigny and Jaeger (50)), similar to those found for the *volume data set* **(Figures S1 and S2, Tables S3 and S4)**. This resulted in P_{EH} ranging from 4 to 72 **(Figure 1A, Tables S1 and S2)**. In the following, we consider P_{EH} as *low* if the EH hydrolyzes ≤ 9 esters (11% of the data set), as *moderate* if the EH hydrolyzes between 10 and 29 esters (49%), and as *high* if the EH hydrolyzes ≥ 30 esters (40%) **(Figure S1, Table S3)**. The data set covers eleven F_{EH} of which F_{IV} (44% of the data set) and F_V (21%) are the best represented ones **(Figure S2, Table S4)**. This reflects the proportion of their presence in the *volume data set*. III) Only EHs with amino acid sequence identities $\geq$ 25% in comparison to an existing crystal structure were considered **(see section 2.1)** in order to ensure a sufficient quality of generated comparative models.

Finally, in order to uniformly depict the results across the present study, six EHs were selected as representatives of the *flexibility data set* based on P_{EH} (termed *representative data set*): EHs with the lowest (EH115) or highest P_{EH} (EH001) and known crystal structures, EHs with the lowest (EH127) or the highest P_{EH} (EH005) and unknown crystal structures, and commercial EHs with the lowest (CalA) or highest P_{EH} (CalB) **(Figure 1A-D, Tables S1 and S2)**.

3.2.   Comparative models of EHs generated by TopModel show an overall and residue-wise good quality

To generate structural models of EHs as starting points for our investigations, we performed template-based modeling of the *flexibility data set* using TopModel (26) **(see section 2.2)**. In doing so, we also generated comparative models of the eleven EHs for which crystal structures are available. These structural models will be used to judge the quality of the comparative modeling.

The quality of the comparative models of the *flexibility data set* were assessed with TopScore (31), a meta Model Quality Assessment Program (meta-MQAP) **(see section 2.2)**. For the eleven Es with known crystal structure, the global TopScores range from 0.074 to 0.305 **(Figure 1A, Table S1)**. As the global TopScore describes whole-protein error estimates, this shows that the structures contain between 7.4 and 30.5% error. Notably, the global TopScores well and significantly correlate ($R^2 = 0.61$, $p = 0.004$) with values of $1 - $ lDDT computed from comparisons of the comparative models of EHs with known crystal structure against these experimental reference structures, indicating that global TopScores are well suited to assess the model quality of EHs **(Figure S3, Table S5)**. The global TopScore values of the comparative models of the other 24 EHs range from 0.087 to 0.269 **(Figure 1A, Table S2)**, indicating that these models are of equal quality than the ones for EHs with known crystal structure. The TopScore values of the *representative data set* lie in a comparable range **(Figure 1A, Tables S1 and S2)**. Moreover, the comparative models of the *flexibility data set* show low residue-wise TopScore values (31), indicating that all parts of a model are of good quality. We illustrate this for the residue-wise TopScore of the comparative models of the *representative data set* **(Figures 1B-D)**. This also applies to structural regions around CARs **(Figures 1B-D)**. That way, it was possible to confirm CARs in models of EHs with known crystal structures and to unambiguously identify CARs in models of EHs with unknown crystal structures **(Figures 1B-D, Tables S1 and S2)**.

Additionally, we validated that CARs in all models are accessible for substrates according to CAVER results (32) **(see section 2.2)**, i.e., that all models are in an open conformation: CARs are either located on the protein surface or are buried and connected with the surface by tunnels. We illustrate this for the comparative models of the *representative data set* **(Figure S4)**.

To conclude, comparative models were generated for 35 EHs of the *flexibility data set* using

TopModel. The models showed both an overall and residue-wise good structural quality.

Additionally, we validated that CARs in all models are accessible for substrates.

3.3. Promiscuous EHs are globally less flexible

Previous studies indicated that enzyme flexibility influences the substrate promiscuity of

enzymes (12-14). For gaining insights into how the flexibility of EHs is linked to P_{EH}, we

applied CNA (21, 23), a rigidity theory-based approach to analyze biomolecular statics (21-

23), to the *flexibility data set* **(see sections 2.4)**. To improve the robustness and investigate the

statistical uncertainty, for each of the comparative models we carried out CNA on ensembles

of network topologies (ENT^{MD}) generated from five MD trajectories of 1 μs length each (44)

(see sections 2.3 and 2.4). In order to investigate if the global flexibility of the EHs

influences P_{EH}, we predicted T_p, the phase transition temperature previously applied as a

measure of structural stability of a protein (20, 41, 44-48), for each EH **(see section 2.5)**. T_p

was averaged over five ensembles **(see sections 2.3 and 2.4)**, resulting in all but one case in

SEM < 1.87 K **(Figure 2A, Tables S1 and S2)**.

T_p and P_{EH} of the *flexibility data set* are well and significantly correlated ($R^2 = 0.60$, $p =$

$5.4*10^{-8}$) **(Figure 2A)**. To validate the consistency of our approach, we considered EHs with

known or unknown crystal structures separately. In both cases, good and significant

correlations between T_p and P_{EH} were revealed (known crystal structures: $R^2 = 0.48$,

$p = 0.019$; unknown crystal structures: $R^2 = 0.73$, $p = 1.1*10^{-7}$), lending support to the quality

of comparative models predicted with TopModel and indicating that future predictions on

EHs with unknown experimental structures should be promising. Notably, EHs with high P_{EH}

have a high T_p and *vice versa*, i.e., promiscuous EHs are globally less flexible. Exemplarily,

this is depicted for EHs of the *representative dataset* with known crystal structures and lowest

(EH115) or highest P_{EH} (EH001), which showed T_p of 322.3 K and 357.2 K, with unknown

crystal structures and lowest (EH127) or highest P_{EH} (EH005), which showed T_p of 318.6 K

and 351.1 K, and CalA and CalB, which showed T_p of 346.2 K and 351.6 K **(Figure 2A,**

Tables S1 and S2). The differences in global structural stability of these EHs are illustrated

by the rigid cluster decomposition at 332 K during the thermal unfolding simulations **(Figure**

2B-D): promiscuous EHs are globally more structurally stable at the elevated temperature as

indicated by fewer, but larger, rigid clusters.

The EH flexibility analyzed so far is a static property and describes the potential of motions in

a biomolecule (23). Yet, direct information on mobility within EHs is available from the

ensembles generated by MD simulations. We thus computed exemplarily $RMSF_{EH}$, a measure

for protein mobility **(see section 2.6)**, across the ensembles of EHs from the *representative*

data set. $RMSF_{EH}$, averaged over all residues and all five MD trajectories, and P_{EH} do not

yield a significant correlation ($p = 0.13$) **(Figure S5A, Table S6)**, in contrast to T_p and P_{EH}

($R^2 = 0.93$, $p = 1.8*10^{-3}$) **(Figure S5B, Table S6)**. Still, as promiscuous EHs are globally less

mobile, the same trend is obtained as in the case of the flexibility analysis.

To conclude, a good and significant correlation between T_p and P_{EH} was found for the

flexibility data set ($R^2 = 0.60$, $p = 5.4*10^{-8}$). These findings demonstrate that promiscuous

EHs are globally less flexible. $RMSF_{EH}$ is less predictive for P_{EH}, although again promiscuous

EHs are characterized by a lower global mobility, mutually confirming either result.

3.4.   Promiscuous EHs are more thermostable

Previous studies indicated that thermodynamically more thermostable proteins frequently

have a higher structural stability (45, 48). In order to investigate if promiscuous EHs, which

were predicted to be less flexible **(see section 3.3)**, are also more thermostable, CD

spectroscopy was applied to determine the melting temperature T_d of the EHs **(see section**

2.8). Note that only if the unfolding of a protein is reversible, CD spectroscopy provides true

thermodynamic properties (51). However, even if the unfolding is irreversible, because the

protein aggregates at high temperatures, the method can still give information about relative

stabilities (51). Hence, to reduce the potential impact of different aggregation kinetics of

structurally different proteins, we applied CD spectroscopy to one F_{EH} family only. In

particular, we used F_{IV} because it is the largest F_{EH} **(Table S7)**.

Exemplarily, a CD spectrum for T_d determination is shown for EH001 **(Figure 3A)**; for each

EH, T_d determination was performed in triplicates with STD < 0.62 K. T_d and P_{EH} yield a fair

and significant correlation ($R^2 = 0.40$, $p = 0.027$) **(Figure 3B)**.

To conclude, promiscuous EHs are not only globally less flexible but also more thermostable.

3.5. Promiscuous EHs have less flexible catalytically active residues

The good correlation of P_{EH} and T_p encouraged us to investigate if local flexibility characteristics of CARs will provide an even better predictor of EH promiscuity. We thus computed $Flex_{CAR}$ for the *flexibility data set*, i.e., the stability of rigid contacts between CARs and other residues that are at most 5 Å apart from each other, based on the *local* index $rc_{ij,neighbor}$ **(see section 2.5)**. For each EH, $Flex_{CAR}$ was averaged over five ensembles **(see sections 2.3 and 2.4)**, resulting in SEM < 0.06 kcal mol^{-1} **(Figure S6A, Tables 1 and 2)**.

$Flex_{CAR}$ and P_{EH} of the *flexibility data set* yield a good and significant correlation ($R^2 = 0.51$, $p = 1.7*10^{-6}$) **(Figure S6A)**. To validate again the consistency of our approach, we considered EHs with known and unknown crystal structures separately. In both cases, good and significant correlations between $Flex_{CAR}$ and P_{EH} were found (known crystal structures: $R^2 = 0.63$, $p = 3.7*10^{-3}$; unknown crystal structures: $R^2 = 0.47$, $p = 2.2*10^{-4}$), again lending support to the quality of comparative models predicted with TopModel. Hence, EHs with high P_{EH} have low $Flex_{CAR}$ and *vice versa*, i.e., promiscuous EHs have less flexible CARs. Exemplarily, this is detailed for EHs of the *representative dataset* with known crystal structures and lowest (EH115) or highest P_{EH} (EH001), which showed $Flex_{CAR}$ of -0.74 kcal mol^{-1} and -1.91 kcal mol^{-1}, with unknown crystal structures and lowest (EH127) or highest P_{EH} (EH005), which showed $Flex_{CAR}$ of -1.10 kcal mol^{-1} and -1.86 kcal mol^{-1}, and CalA and CalB, which showed $Flex_{CAR}$ of -1.31 kcal mol^{-1} and -1.95 kcal mol^{-1} **(Figure S6A, Tables S1 and S2)**. The differences in local structural stability of these EHs are illustrated by rigid contacts between CARs and other residues that are at most 5 Å apart from each other **(Figure S6B-D)**: promiscuous EHs are locally more structurally stable as indicated by more stable rigid contacts.

Finally, we exemplarily computed $RMSF_{CAR}$, a measure for the mobility of a protein's CARs **(see section 2.6),** across the ensembles of EHs from the *representative data set*. Averaged $RMSF_{CAR}$ and P_{EH} correlate worse ($R^2 = 0.74$, $p = 0.029$) **(Figure S7A, Table S6)** than $Flex_{CAR}$ and P_{EH} ($R^2 = 0.92$, $p = 2.4*10^{-3}$) **(Figure S7B, Table S6)**, paralleling the above results for the global measures. Still, again, as promiscuous EHs have less mobile CARs, the same trend is obtained as in the case of the flexibility analysis.

To conclude, a good and significant correlation between $Flex_{CAR}$ and P_{EH} was found for the *flexibility data set* ($R^2 = 0.51$, $p = 1.7*10^{-6}$). Hence, promiscuous EHs have less flexible

362 CARs. $RMSF_{CAR}$ is less predictive for P_{EH}, although again promiscuous EHs are characterized

363 by less mobile CARs, mutually confirming either result.

3.6.   Promiscuous EHs have an increased specific activity

In the study by Ferrer *et al.* (19), the *experimental data set* was screened against 96 esters in a

kinetic pH indicator assay **(see section 2.1)**. Besides the average specific activity $Act_{average}$

given in U / (g wet cells), also the average maximum specific activity Act_{max} was determined.

Motivated by the reactivity-selectivity principle (RSP) initially introduced for organic

chemistry reactions (52), which states that a more reactive chemical compound is less

selective in chemical reactions, we intended to probe if P_{EH} is related to Act_{max}. For this, we

established an approximate linear free-energy relationship (LFER) (53) by relating log(Act_{max})

and log(P_{EH}) **(Figure S8, Table S8)**. In this analysis, the CalA and CalB preparations were

excluded because Act_{max} was given in U / (g total protein) there.

Log(Act_{max}) and log(P_{EH}) of the *experimental data set* yield a good and significant correlation

($R^2 = 0.50$, $p = 4.6*10^{-23}$) **(Figure S8A)**. Likewise, log(Act_{max}) and log(P_{EH}) of the *flexibility*

data set yield a fair and significant correlation ($R^2 = 0.22$, $p = 0.6*10^{-2}$) **(Figure S8B)**. To

validate whether the same trend emerges for EHs with known and unknown crystal structures,

we considered both types of EHs separately. In both cases, fair and significant correlations

between log(Act_{max}) and log(P_{EH}) were found (known crystal structures: $R^2 = 0.34$, $p = 0.099$;

unknown crystal structures: $R^2 = 0.23$, $p = 0.019$).

To conclude, good to fair and significant correlations between log(Act_{max}) and log(P_{EH}) of the

experimental data set ($R^2 = 0.50$, $p = 4.6*10^{-23}$) and the *flexibility data set* ($R^2 = 0.22$, $p =$

$0.6*10^{-2}$) were found. Hence, promiscuous EHs have higher maximum specific activities.

3.7.   Specific EHs prefer to hydrolyze large and flexible esters

Next, we investigated, which of the 96 esters was preferentially hydrolyzed by EHs with

different P_{EH}. As a criterion, we chose the number of freely rotatable bonds of an ester, TA

(see section 2.7). We did so because TA is a combined measure for an ester's size and

conformational dynamics (54). To account for the uneven distribution of esters in our data set

with respect to TA, we calculated $Norm_{ester}(TA)$, i.e., the number of hydrolyzed esters with a

specific TA (*Ester*_{hydrolysed} (TA)) divided by the total number of esters in the data set with this

specific TA (*Ester*_{library}(TA)) **(see section 2.7) (Eq. 1)**.

According to TA, the esters were classified into 17 groups that ranged from small esters with

no rotatable bond to large esters with 56 rotatable bonds **(Figure 4, Table S9)**. Esters with

three (24% of the ester library) and four (16% of the ester library) rotatable bonds are most

frequent. The analysis of the *experimental data set* revealed that promiscuous EHs have high

*Norm*_{ester} values irrespective of TA, i.e., promiscuous EHs accept a large variety of esters with

different sizes and degrees of conformational dynamics **(Figure 4A, Table S9)**. In contrast,

specific EHs only have high *Norm*_{ester} values regarding esters with high TA, i.e., specific EHs

preferentially hydrolyze (very) large and flexible esters **(Figure 4A, Table S9)**. The same

tendency was observed for the *flexibility data set* **(Figure 4B, Table S9)**.

To conclude, promiscuous EHs accept a large variety of esters with different sizes and

degrees of conformational dynamics whereas specific EHs preferentially hydrolyze (very)

large and flexible esters.

4. Discussion

The main outcomes of our analyses are I) that promiscuous EHs are significantly globally less flexible and have less flexible catalytically active residues than specific ones, II) that promiscuous EHs are significantly more thermostable, III) that promiscuous EHs have a significantly increased specific activity, and IV) that specific EHs prefer to hydrolyze large and flexible esters.

We established these relations using one of the still few experimental large-scale datasets where a diverse set of EHs was functionally assessed against a customized library of dissimilar esters (19). Functional promiscuity may arise from several conditions, including the environment of the enzyme or the concentration of a substrate, which may complicate the analysis of the molecular mechanism underlying promiscuity (5). Still, functional promiscuity ultimately is a result of recognition promiscuity (5); here, we therefore focused on substrate promiscuity (18), i.e., an enzyme carries out its typical catalytic function using non-canonical substrates, in that experimental conditions had been kept constant for the assessment of the different esterase/ester combinations (19). Almost all of the EHs were unambiguously assigned to one of the F_{EH} of the Arpigny and Jaeger classification, which is based mainly on a comparison of amino acid sequences (50). Except for classes with a few members only (cyclase-like EHs and the yeast family), all other classes cover at least two of the three P_{EH} ranges such that P_{EH} cannot be assigned based on the EH's class affiliation (**Figure S9, Table S10**) and, hence, amino acid sequence information. Even family F_{IV}, which contains a higher proportion of substrate-promiscuous EHs, also contains EHs with a small substrate range.

For scrutinizing the mechanism underlying esterase promiscuity at the atomistic level, we needed to apply comparative models of EHs, since only for ~7% of the experimentally assessed EHs crystal structures were available. Restricting the generation of esterase models to sequence identifies ≥ 25% with respect to available targets yielded generally good structural models both globally and locally, as also validated against cases where crystal structures are known. Throughout our study, we probed for the consistency of our analyses between subsets of EHs for which either crystal structures are known or not; we only found quantitative differences, but no qualitative ones. One of the reasons is likely that rigidity analyses were based on structural ensembles generated by multiple and μs-long MD simulations, which markedly increases the robustness of the results (42). We furthermore showed that results are consistent irrespective of whether EH flexibility characteristics were assessed globally or only for CARs, and that mobility characteristics computed directly from

MD trajectories show the same trend, although the correlation with P_{EH} is insignificant.

Finally, we used experimental melting temperatures of EHs as indicators for enzyme

flexibility (45, 48), which yielded the same relation with P_{EH} as computed flexibility

characteristics. Overall, these consistent and robust findings indicate that when applying this

workflow to novel EHs, it should be possible to discover enzymes with 'sufficient' substrate

promiscuity to serve as a starting point for further exploration in biotechnology and synthetic

organic chemistry. In that respect, the flexibility characteristics of EHs analyzed here have a

notably stronger predictive power than the active site effective volume introduced earlier (19)

(**Figure S10, Tables S11 and S12**).

The finding that promiscuous EHs are significantly globally *less* flexible and have *less*

flexible catalytically active residues than specific EHs is in stark contrast to the general view

of the role of structural flexibility for promiscuity (4, 5): Besides the examples of CYP and β-

lactamase mentioned above, the possibility of dynamically restructuring active sites has also

been recognized for other systems as underlying their promiscuity (55-58). Finally,

interactions between antibodies and antigens are likely the quintessential example of the

canonical relationship between flexibility and binding promiscuity: As antibodies mature to

become more specific, their flexibility is decreased (5).

It has been recognized that conformational changes may not always be necessary for

promiscuity if a variety of substrates can be bound by partial recognition or the presence of

multiple binding sites (5). However, these cases do not seem to be relevant reasons for EH

promiscuity because partial recognition often is associated with catalytic inefficiency (1),

which is contrary to our observation that P_{EH} correlates with EH activity, and the presence of

multiple binding sites that could give rise to promiscuity is controverted by the finding that

promiscuous EHs have large active site effective volumes (19), i.e., large pockets with few

subpockets. Inversely, our findings of rigid promiscuous EHs may be consistent with the idea

that multiple ligands can be accommodated in a single site by exploiting diverse interacting

residues (**Figure 5**).

Our results as to *specific but flexible* EHs may be reconciled with a model according to which

conformational changes may have been selected in EH evolution for their ability to enhance

specificity in recognition (**Figure 5**), resulting in what has been termed conformational

proofreading (16). In the case of specific EHs, flexibility may help to overcome a structural

mismatch between the enzyme and its substrate existing when both are in their ground states,

that way enhancing recognition specificity. This view is corroborated by our finding that

specific EHs prefer to hydrolyze large and flexible substrates: Larger substrates can form
more interactions with the enzyme, that way helping to overcome the deformation energy
required by the enzyme to optimizing the correct binding probability over the incorrect one;
flexible substrates can tolerate higher strains and thus can be expected to participate in more
binding events (59, 60) **(Figure 5)**.

In summary, the combined large-scale analysis of experimental EH promiscuity and
computed EH flexibility reveals that promiscuous EHs are significantly less flexible than
specific ones. This result is counterintuitive at first but may be reconciled with a model
according to which multiple ligands can be accommodated in a single site of promiscuous
EHs by exploiting diverse interacting residues, whereas structural flexibility in the case of
specific EHs serves for conformational proofreading. Our results furthermore signify that EH
sequence space, charted, e.g., by (meta)genomics studies, can be screened by rigidity analyses
for promiscuous EHs that may serve as starting points for further exploration in biotechnology
and synthetic organic chemistry.

5. Acknowledgements

CN is funded through a grant ("Vernetzungsdoktorand") provided by the Forschungszentrum Jülich. Parts of the study were supported by Bundesministerium für Bildung und Forschung (BMBF) through funding number 031B0837A "LipoBiocat" to HG and KEJ as well as the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through funding no. INST 208/704-1 FUGG to HG and INST 208/654-1 FUGG to KEJ. HG is grateful for computational support and infrastructure provided by the "Zentrum für Informations- und Medientechnologie" (ZIM) at the Heinrich Heine University Düsseldorf. HG gratefully acknowledges the computing time granted by the John von Neumann Institute for Computing (NIC) and provided on the supercomputer JUWELS at Jülich Supercomputing Centre (JSC) (user IDs: HKF7; protil (project ID: 15956)) (61). MF acknowledges the grant 'INMARE' from the European Union's Horizon 2020 (grant agreement no. 634486) and BIO2017-85522-R from the Ministerio de Ciencia, Innovación y Universidades (MCIU), Agencia Estatal de Investigación (AEI), Fondo Europeo de Desarrollo Regional (FEDER) and European Union (EU). CC thanks the Ministerio de Economía y Competitividad and FEDER for a PhD fellowship (Grant BES-2015-073829). The authors are grateful to David Almendral and Ruth Matesanz for their support of CD analysis.

6. Authors contributions

HG and KEJ conceived the study. CN analyzed the experimental data, performed structure prediction, MD simulations and CNA computations, analyzed the computational data, and wrote the manuscript together with HG. DM initially contributed to the structure prediction. MF and CC measured and analyzed melting temperatures. HG supervised and managed the project. All authors reviewed and approved the manuscript.

7. Conflict of interest

The authors declare no financial and non-financial competing interests.

8. Figures

Figure 1: Comparative modeling of EHs. (A) Based on sequence data provided by a large-scale study from Ferrer *et al.* (19), comparative models were generated for 35 EHs with known (left, 11 EHs) and unknown (right, 24 EHs) crystal structures using TopModel (26). These EHs constitute the *flexibility data set*. The EHs vary in P_{EH} (left ordinate, bars) and global TopScores (right ordinate, diamonds). Six EHs were selected as representatives of the *flexibility data set* (termed *representative data set*) as indicated by magenta arrows. The quality of the comparative models of **(B)** EHs with known crystal structures and lowest (EH115) or highest P_{EH} (EH001), **(C)** EHs with unknown crystal structures and lowest (EH127) or highest P_{EH} (EH005), and **(D)** commercial EHs with highest (CalA) or lowest P_{EH} (CalB) was evaluated by TopScore (31). For each comparative model the residue-wise TopScore is shown: A good (bad) homology model shows a low (high) residue-wise TopScore (see color scale at the bottom). Insets depict CARs (spheres) within an EH. For clarity the position of CARs is indicated by magenta stars.

Figure 2: Correlation of T_p versus P_{EH}. **(A)** Correlation between predicted T_p based on the global index H_{type2} and P_{EH} for the *flexibility data set*. Data points colored grey (black) represent comparative models of EHs with (un)known crystal structures. The *representative data set* is indicated by magenta crosses. Error bars show the SEM over five independent MD simulations of 1 µs length each. Rigid cluster decomposition at 332 K during the thermal unfolding simulation of **(B)** EHs with known crystal structures and lowest (EH115) or highest P_{EH} (EH001), **(C)** EHs with unknown crystal structures and lowest (EH127) or highest P_{EH} (EH005), and **(D)** commercial EHs with lowest (CalA) or highest P_{EH} (CalB). Rigid clusters are represented as uniformly colored blue, green, pink, cyan, and magenta bodies in the descending order of their sizes.

Figure 3: Determination of T_d via CD spectroscopy. (A) Exemplary CD spectrum of EH001. The ellipticity changes in mdeg at 220 nm was plotted against the temperature, resulting in a sigmoidal curve. The inflection point was used to obtain the T_d value (dotted line). **(B)** Correlation between T_d and P_{EH} for 12 EHs of F_{IV}.

Figure 4: Relation between the number of esters' TA and P_{EH}. Relation between *Ester*norm, i.e., the relative proportion of the number of hydrolyzed esters with a specific TA, and P_{EH} of **(A)** the *experimental data set* and **(B)** the *flexibility data set* containing EHs with known crystal structures (left), EHs with unknown crystal structures (right), and EHs constituting the *representative data set* (indicated by magenta arrows). TA was calculated based on SMILES codes of 96 esters provided by Ferrer *et al.* (19). A blue (red) color indicates that the EH hydrolyzes many (few) esters with a specific TA relative to the total number of esters in the data set with this specific TA (see color scale on the right); the total number of esters with a specific TA is given in brackets on the y-axis. P_{EH} is defined as *low* if the EH hydrolyzes ≤ 9 esters, as *moderate* if the EH hydrolyzes between 10 and 29 esters, and as *high* if the EH hydrolyzes ≥ 30 esters.

Figure 5: Mechanistic model of EH flexibility, ligand size and conformational dynamics affecting P_{EH}. Impact of esters with **(A)** many or **(B)** few TA on specific, and hence more flexible (left), and promiscuous, and hence more rigid (right), EHs. Ligand parts connected by TA are represented as blue circles. Specific EHs and large ligands with many TA can mutually adapt (panel **A**, left), and promiscuous EH can bind large ligands (panel **A**, right) and small ligands (panel **B**, right) exploiting different interaction partners. Small (and/or rigid) ligands are not able to lead to a structural adaptation of specific EHs (panel **B**, left), though, resulting in conformational proofreading. The red bars indicate the flexibility of the EHs. A green tick (red cross) indicates that ester cleavage is (not) catalyzed.

9. References

1. S. D. Copley, Shining a light on enzyme promiscuity. *Curr. Opin. Struct. Biol.* **47**, 167-175 (2017).
2. R. Chen *et al.*, Molecular insights into the enzyme promiscuity of an aromatic prenyltransferase. *Nat. Chem. Biol.* **13**, 226-234 (2017).
3. H. Huang *et al.*, Panoramic view of a superfamily of phosphatases through substrate profiling. *Proc. Natl. Acad. Sci. U.S.A.* **112**, E1974-E1983 (2015).
4. O. Khersonsky, D. S. Tawfik, Enzyme promiscuity: a mechanistic and evolutionary perspective. *Annu. Rev. Biochem.* **79**, 471-505 (2010).
5. I. Nobeli, A. D. Favia, J. M. Thornton, Protein promiscuity and its implications for biotechnology. *Nat. Biotechnol.* **27**, 157-167 (2009).
6. E. Z. Eisenmesser, D. A. Bosco, M. Akke, D. Kern, Enzyme dynamics during catalysis. *Science* **295**, 1520-1523 (2002).
7. K. A. Henzler-Wildman *et al.*, A hierarchy of timescales in protein dynamics is linked to enzyme catalysis. *Nature* **450**, 913-916 (2007).
8. M. J. Holliday, C. Camilloni, G. S. Armstrong, M. Vendruscolo, E. Z. Eisenmesser, Networks of dynamic allostery regulate enzyme function. *Structure* **25**, 276-286 (2017).
9. N. M. Goodey, S. J. Benkovic, Allosteric regulation and catalysis emerge via a common route. *Nat. Chem. Biol.* **4**, 474-482 (2008).
10. H. G. Saavedra, J. O. Wrabl, J. A. Anderson, J. Li, V. J. Hilser, Dynamic allostery can drive cold adaptation in enzymes. *Nature* **558**, 324-328 (2018).
11. S. R. Tzeng, C. G. Kalodimos, Protein dynamics and allostery: an NMR view. *Curr. Opin. Struct. Biol.* **21**, 62-67 (2011).
12. J. Skopalík, P. Anzenbacher, M. Otyepka, Flexibility of human cytochromes P450: molecular dynamics reveals differences between CYPs 3A4, 2C9, and 2A6, which correlate with their substrate preferences. *J. Phys. Chem. B.* **112**, 8165-8173 (2008).
13. T. Hendrychová *et al.*, Flexibility of human cytochrome P450 enzymes: molecular dynamics and spectroscopy reveal important function-related variations. *Biochim. Biophys. Acta.* **1814**, 58-68 (2011).
14. M. Ekroos, T. Sjögren, Structural basis for ligand promiscuity in cytochrome P450 3A4. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 13682-13687 (2006).
15. T. Zou, V. A. Risso, J. A. Gavira, J. M. Sanchez-Ruiz, S. B. Ozkan, Evolution of conformational dynamics determines the conversion of a promiscuous generalist into a specialist enzyme. *Mol. Biol. Evol.* **32**, 132-143 (2015).
16. Y. Savir, T. Tlusty, Conformational proofreading: the impact of conformational changes on the specificity of molecular recognition. *PloS One* **2**, e468 (2007).
17. M. Ferrer *et al.*, Estimating the success of enzyme bioprospecting through metagenomics: current status and future trends. *Microb. Biotechnol.* **9**, 22-34 (2016).
18. K. Hult, P. Berglund, Enzyme promiscuity: mechanism and applications. *Trends. Biotechnol.* **25**, 231-238 (2007).
19. M. Martinez-Martinez *et al.*, Determinants and prediction of esterase substrate promiscuity patterns. *ACS Chem. Biol.* **13**, 225-234 (2017).
20. S. Radestock, H. Gohlke, Exploiting the link between protein rigidity and thermostability for data-driven protein engineering. *Eng. Life Sci.* **8**, 507-522 (2008).
21. C. Pfleger, P. C. Rathi, D. L. Klein, S. Radestock, H. Gohlke, Constraint Network Analysis (CNA): a Python software package for efficiently linking biomacromolecular structure, flexibility,(thermo-) stability, and function. *J. Chem. Inf. Model.* **53**, 1007-1015 (2013).

22. D. M. Krüger, P. C. Rathi, C. Pfleger, H. Gohlke, CNA web server: rigidity theory-based thermal unfolding simulations of proteins for linking structure,(thermo-) stability, and function. *Nucleic Acids Res.* **41**, W340-W348 (2013).

23. S. M. Hermans, C. Pfleger, C. Nutschel, C. A. Hanke, H. Gohlke, Rigidity theory for biomolecules: concepts, software, and applications. *Comput. Mol. Sci.* **7**, e1311 (2017).

24. K. de Godoy Daiha, R. Angeli, S. D. de Oliveira, R. V. Almeida, Are lipases still important biocatalysts? A study of scientific publications and patents for technological forecasting. *PloS One* **10**, e0131624 (2015).

25. M. Ferrer *et al.*, Biodiversity for biocatalysis: a review of the α/β-hydrolase fold superfamily of esterases-lipases discovered in metagenomes. *Biocatal. Biotranfor.* **33**, 235-249 (2015).

26. D. Mulnaes *et al.*, TopModel: Template-based protein structure prediction at low sequence identity using top-down consensus and deep neural networks. *J. Chem. Theory. Comput.* **16**, 1953-1967 (2020).

27. H. Gohlke *et al.*, Binding region of alanopine dehydrogenase predicted by unbiased molecular dynamics simulations of ligand diffusion. *J. Chem. Inf. Model.* **53**, 2493-2498 (2013).

28. N. Widderich *et al.*, Molecular dynamics simulations and structure-guided mutagenesis provide insight into the architecture of the catalytic core of the ectoine hydroxylase. *J. Mol. Biol.* **426**, 586-600 (2014).

29. Z. Zhang *et al.*, Determinants of FIV and HIV Vif sensitivity of feline APOBEC3 restriction factors. *Retrovirology* **13**, 46 (2016).

30. D. Milić *et al.*, Recognition motif and mechanism of ripening inhibitory peptides in plant hormone receptor ETR1. *Sci. Rep.* **8**, 3890 (2018).

31. D. Mulnaes, H. Gohlke, TopScore: Using Deep Neural Networks and Large Diverse Data Sets for Accurate Protein Model Quality Assessment. *J. Chem. Theory Comput.* **14**, 6117-6126 (2018).

32. E. Chovancova *et al.*, CAVER 3.0: a tool for the analysis of transport pathways in dynamic protein structures. *PLoS Comput. Biol.* **8**, e1002708 (2012).

33. R. Salomon-Ferrer, A. W. Götz, D. Poole, S. Le Grand, R. C. Walker, Routine microsecond molecular dynamics simulations with AMBER on GPUs. 2. Explicit solvent particle mesh Ewald. *J. Chem. Theory Comput.* **9**, 3878-3888 (2013).

34. D. A. Case *et al.* (2019) AMBER 2019. (University of California, San Francisco).

35. C. W. Hopkins, S. Le Grand, R. C. Walker, A. E. Roitberg, Long-time-step molecular dynamics through hydrogen mass repartitioning. *J. Chem. Theory Comput.* **11**, 1864-1874 (2015).

36. D. J. Jacobs, A. J. Rader, L. A. Kuhn, M. F. Thorpe, Protein flexibility predictions using graph theory. *Proteins* **44**, 150-165 (2001).

37. B. Hespenheide, D. Jacobs, M. Thorpe, Structural rigidity in the capsid assembly of cowpea chlorotic mottle virus. *J. Condens. Matter Phys.* **16**, S5055 (2004).

38. D. J. Jacobs, M. F. Thorpe, Generic rigidity percolation: the pebble game. *Phys. Rev. Lett.* **75**, 4051 (1995).

39. A. J. Rader, B. M. Hespenheide, L. A. Kuhn, M. F. Thorpe, Protein unfolding: rigidity lost. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 3540-3545 (2002).

40. D. R. Livesay, D. J. Jacobs, Conserved quantitative stability/flexibility relationships (QSFR) in an orthologous RNase H pair. *Proteins* **62**, 130-143 (2006).

41. S. Radestock, H. Gohlke, Protein rigidity and thermophilic adaptation. *Proteins* **79**, 1089-1108 (2011).

42. C. Pfleger, H. Gohlke, Efficient and robust analysis of biomacromolecular flexibility using ensembles of network topologies based on fuzzy noncovalent constraints. *Structure* **21**, 1725-1734 (2013).

43. C. Pfleger, S. Radestock, E. Schmidt, H. Gohlke, Global and local indices for characterizing biomolecular flexibility and rigidity. *J. Comput. Chem.* **34**, 220-233 (2013).

44. P. C. Rathi, S. Radestock, H. Gohlke, Thermostabilizing mutations preferentially occur at structural weak spots with a high mutation ratio. *J. Biotechnol.* **159**, 135-144 (2012).

45. P. C. Rathi, K.-E. Jaeger, H. Gohlke, Structural rigidity and protein thermostability in variants of lipase A from Bacillus subtilis. *PLoS One* **10**, e0130289 (2015).

46. M. Dick *et al.*, Trading off stability against activity in extremophilic aldolases. *Sci. Rep.* **6**, 17908 (2016).

47. P. C. Rathi, A. Fulton, K.-E. Jaeger, H. Gohlke, Application of rigidity theory to the thermostabilization of lipase A from Bacillus subtilis. *PLoS Comput. Biol.* **12**, e1004754 (2016).

48. C. Nutschel *et al.*, Systematically scrutinizing the impact of substitution sites on thermostability and detergent tolerance for Bacillus subtilis lipase A. *J. Chem. Inf. Model.* **60**, 1568-1584 (2020).

49. H. Gohlke, L. A. Kuhn, D. A. Case, Change in protein flexibility upon complex formation: analysis of Ras-Raf using molecular dynamics and a molecular framework approach. *Proteins* **56**, 322-337 (2004).

50. J. L. Arpigny, K.-E. Jaeger, Bacterial lipolytic enzymes: classification and properties. *Biochem. J.* **343**, 177-183 (1999).

51. N. J. Greenfield, Using circular dichroism collected as a function of temperature to determine the thermodynamics of protein unfolding and binding interactions. *Nat. Protoc.* **1**, 2527-2535 (2006).

52. H. Mayr, A. R. Ofial, The reactivity–selectivity principle: an imperishable myth in organic chemistry. *Angew. Chem. Int. Ed. Engl.* **45**, 1844-1854 (2006).

53. P. R. Wells, Linear Free Energy Relationships. *Chem. Rev.* **63**, 171-219 (1963).

54. H.-J. Böhm, The development of a simple empirical scoring function to estimate the binding constant for a protein-ligand complex of known three-dimensional structure. *J. Comput. Aided Mol. Des.* **8**, 243-256 (1994).

55. C. M. Seibert, F. M. Raushel, Structural and catalytic diversity within the amidohydrolase superfamily. *Biochemistry* **44**, 6383-6391 (2005).

56. U. Oppermann *et al.*, Short-chain dehydrogenases/reductases (SDR): the 2002 update. *Chem. Biol. Interact.* **143-144**, 247-253 (2003).

57. S. Fushinobu, H. Nishimasu, D. Hattori, H.-J. Song, T. Wakagi, Structural basis for the bifunctionality of fructose-1, 6-bisphosphate aldolase/phosphatase. *Nature* **478**, 538-541 (2011).

58. J. Du, R. F. Say, W. Lü, G. Fuchs, O. Einsle, Active-site remodelling in the bifunctional fructose-1, 6-bisphosphate aldolase/phosphatase. *Nature* **478**, 534-537 (2011).

59. G. R. Stockwell, J. M. Thornton, Conformational diversity of ligands bound to proteins. *J. Mol. Biol.* **356**, 928-944 (2006).

60. E. Perola, P. S. Charifson, Conformational analysis of drug-like molecules bound to proteins: an extensive study of ligand reorganization upon binding. *J. Med. Chem.* **47**, 2499-2510 (2004).

61. D. Krause, JUWELS: Modular Tier-0/1 Supercomputer at the Jülich Supercomputing Centre. *JLSRF* **5**, A135 (2019).