ECOLOGICAL DIFFERENTIATION AND INCIPIENT SPECIATION IN THE FUNGAL PATHOGEN CAUSING

RICE BLAST

Maud THIERRY[1,2,3], Florian CHARRIAT[1], Joëlle MILAZZO[1,2], Henri ADREIT[1,2], Sébastien RAVEL[1,2],

Sandrine CROS-ARTEIL[1], Sonia BORRON[1], Violaine SELLA[3], Thomas KROJ[1], Renaud IOOS[3], Elisabeth

FOURNIER[1], Didier THARREAU[1,2,4], Pierre GLADIEUX[1,4]

[1]PHIM Plant Health Institute, Univ Montpellier, INRAE, CIRAD, Institut Agro, IRD, Montpellier, France

[2]CIRAD, UMR PHIM, 34090 Montpellier, France.

[3]ANSES Plant Health Laboratory, Mycology Unit, Domaine de Pixérécourt, Bâtiment E, F-54220 Malzéville,

France

[4]pierre.gladieux@inrae.fr; didier.tharreau@cirad.fr

**ABSTRACT**

Many invasive fungal species coexist as multiple populations on the same host, but the factors behind

the origin and maintenance of population structure remain largely unknown. Here, we analyzed

genetic and phenotypic diversity in isolates of the rice blast fungus (*Pyricularia oryzae*) covering a

broad geographical range. We showed that the four lineages of *P. oryzae* were found in areas that

differ in terms of prevailing environmental conditions and types of rice grown. Pathogenicity tests

revealed that specialization to rice subspecies contributed to niche separation between lineages, and

differences in repertoires of putative pathogenicity effectors were consistent with differences in host

range. Crossing experiments revealed that female sterility and early post-mating genetic

incompatibilities acted as strong barriers to gene flow between these lineages. Our results

demonstrate that the spread of a pathogen across heterogeneous habitats and divergent populations

of a crop species can lead to niche separation and incipient speciation.

Short title: Incipient speciation in the rice blast pathogen

## INTRODUCTION

Understanding and controlling natural variation in fungal plant pathogens is of great importance for maintaining food security and ecosystem health. In addition to the overwhelming phylogenetic diversity of pathogens (Burdon 1993), many pathogens also harbor substantial diversity at the species scale (Taylor & Fisher 2003). Most fungal pathogens of plants form distinct populations (Taylor *et al.* 2006), and this population structure is a compound outcome of migration, selection and drift, acting at multiple scales, and patterned by factors such as time (Ali *et al.* 2014; Pagliaccia *et al.* 2018), admixture (Robert *et al.* 2015), geography (Ali *et al.* 2014; Marin *et al.* 2009; Yang *et al.* 2018), mode of reproduction (Ali *et al.* 2014; Bueker *et al.* 2016; Gibson *et al.* 2012; Ropars *et al.* 2016) and environmental conditions (Walker *et al.* 2015). The identity of the host plants with which fungal pathogens interact is widely thought to be the environmental factor structuring the pathogen population with the strongest impact. Many plant pathogens reproduce within or on their host, resulting in assortative mating on the basis of host use and a strong association between adaptation to the host and reproductive isolation (Giraud *et al.* 2010; Servedio *et al.* 2011). However, more generally, population structure can result from limited dispersal (i.e. limited gene flow due to distance) or limited adaptation (i.e. limited gene flow because of differences in the capacity to exploit resources), both of which may be due to a wealth of potential factors largely unexplored in plant pathogens. Developing an understanding of how population structure emerges and is maintained in fungal plant pathogens is a major goal in evolutionary microbiology, because it will improve our comprehension of both disease emergence and the origin of fungal biodiversity.

*Pyricularia oryzae* (Ascomycota; syn. *Magnaporthe oryzae*) is a widespread model plant pathogen displaying population subdivision. Several lineages, each associated mostly with a single main cereal/grass host, have been characterized within *P. oryzae* (Gladieux *et al.* 2018a). The lineage infecting Asian rice (*Oryza sativa*) has been characterized in detail. Previous population genomic studies of population structure in the rice-infecting lineage revealed genetic subdivision into three (Saleh *et al.* 2014; Tharreau *et al.* 2009; Zhong *et al.* 2018) to six (Gladieux *et al.* 2018b) lineages,

estimated to have diverged approximately 1000 years ago (Gladieux *et al.* 2018b; Latorre *et al.* 2020;

Zhong *et al.* 2018). *Pyricularia oryzae* is a heterothallic fungus with a sexual cycle involving mating

between individuals of opposite mating types, with at least one of the partners capable of producing

female structures (i.e. "female-fertile"). One single lineage (lineage 1, prevailing in East and Southeast

Asia) displayed a genome-wide signal of recombination, a balanced ratio of mating type alleles and a

high frequency of fertile females, consistent with sexual reproduction (Gladieux *et al.* 2018b; Saleh *et*

*al.* 2014; Saleh *et al.* 2012b). All the other lineages had clonal population structures, highly biased

frequencies of mating types and few fertile females, suggesting strictly asexual reproduction

(Gladieux *et al.* 2018b; Saleh *et al.* 2014). In theory, gene flow remains possible between the majority

of the lineages, because there are rare fertile strains and some lineages have fixed opposite mating

types. However, while resequencing studies have revealed signatures of recent shared ancestry

between clonal lineages and the recombinant lineage from South-West Asia, this is not the case

between the different clonal lineages, consistent with a lack of recent gene flow between lineages of

opposite mating types and despite their wide and overlapping distributions (Gladieux *et al.* 2018b).

The loss of female fertility is probably a key factor in the emergence of clonal lineages, inducing a

shift towards strictly asexual reproduction, and protecting populations adapting to new conditions

from maladaptive gene flow from ancestral, recombining populations (Giraud *et al.* 2010). This raises

questions about the nature of the factors underlying the emergence of different rice-infecting

lineages in *P. oryzae*, and contributing to their maintenance, because, according to the competitive

exclusion principle, different lineages should not be able to co-exist on the same host (Abbate *et al.*

2018; Giraud *et al.* 2017; Hardin 1960). Previous studies have helped to cement the status of *P. oryzae*

as a model for studies of the population biology of fungal pathogens, but most efforts to understand

the population structure of the pathogen have been unable to provide a large-scale overview of the

distribution of rice-infecting lineages and of the underlying phenotypic differences, because the

number of genetic markers was limited (Saleh *et al.* 2014; Tharreau *et al.* 2009), the number of

isolates was relatively small (Gladieux *et al.* 2018b; Zhong *et al.* 2018) and because few phenotypic

traits were scored (Saleh *et al.* 2014; Tharreau *et al.* 2009; Zhong *et al.* 2018).

We provide here a detailed genomic and phenotypic overview of natural variation in rice-

infecting *P. oryzae* isolates sampled across all rice-growing areas of the world. With a view to

inferring and understanding the global population structure of rice-infecting *P. oryzae*, we analyzed

genetic variation, using genotyping data with a high resolution in terms of genomic markers and

geographic -coverage, and measured phenotypic variation for a representative subset of the sample

set. We also characterized the genetic variability and content of repertoires of effectors –secreted

proteins that modulate host responses – using whole-genome resequencing data for a representative

set of isolates. We addressed the following specific questions: (i) How many different lineages of *P.*

*oryzae* infect rice? (ii) Do lineages of *P. oryzae* display footprints of recombination? (iii) Do lineages of

*P. oryzae* present evidence of recent admixture? (iv) Are the different lineages sexually competent

and compatible? (v) Are the different lineages differentially adapted to host and temperature? (vi)

Can we find differences in number, identity and molecular evolution of putative effectors between

lineages? (vii) Can we find differences in the geographic and climatic distribution of lineages?

**RESULTS**

**The rice blast pathogen comprises one recombining, admixed lineage and three clonal, non-**

**admixed lineages**

We characterized the global genetic structure of the rice blast pathogen, using an Illumina genotyping

beadchip to score 5,657 SNPs distributed throughout the genome of 886 *P. oryzae* isolates collected

from cultivated Asian rice in 51 countries (Supplementary file 1). The final dataset included 3,686

SNPs after the removal of positions with missing data, which identify 264 distinct multilocus

genotypes out of 886 isolates. Clustering analyses based on sparse nonnegative matrix factorization

algorithms, as implemented in the sNMF program (Frichot et al., 2014), revealed four clusters,

hereafter referred to as "lineages" (Figure 1C). The model with $K$=4 clusters was identified as the best

model on the basis of the cross-entropy criterion, as models with $K$>4 induced only a small decrease

in cross-entropy, suggesting that $K$=4 captured the deepest subdivisions in the dataset (Figure 1 –

figure supplement 1). The neighbor phylogenetic network inferred with SPLITSTREE also supported the

subdivision into four widely distributed lineages, with long branches separating three peripheral

lineages branching off within a central lineage (Figure 1A; 1B). A comparison with previous findings

revealed that the central lineage corresponded to the combination of recombining lineage 1 and

lineages 5 and 6, represented by two and one individuals, respectively, in (Gladieux *et al.* 2018b)

(Supplementary file 2). This central lineage is referred to hereafter as lineage 1. The three peripheral

lineages in the network corresponded to the previously described lineages 2, 3 and 4 (Gladieux *et al.*

2018b) with lineages 2 and 3 similar to lineages B and C, respectively, identified on the basis of

microsatellite data in a previous study (Saleh *et al.* 2014)(Supplementary file 2). Genetic

differentiation between the four lineages was strong and significant (Weir and Cockerham's $F_{ST}$>

0.54), indicating strong barriers to gene flow between the lineages. All genotypes from lineages 2 to 4

had high proportions of membership $q$ in a single cluster ($q$ > 0.89), whereas shared ancestry with the

other three lineages was detected in lineage 1, with 32% of genotypes having $q$ > 0.10 in lineages 2-4

(Figure 1C; Figure 1 – source data 3). Admixture may account for the lower $F_{ST}$ observed in

comparisons between lineage 1 and the other lineages. We detected no footprints of recombination

in lineages 2, 3 and 4 with the pairwise homoplasy index (PHI) test and a null hypothesis of clonality

(Table 1), confirming previous findings (Gladieux *et al.* 2018b; Latorre *et al.* 2020; Saleh *et al.* 2014;

Zhong *et al.* 2018).

**Geographic structure and admixture within recombining lineage 1**

Lineage 1 was the only lineage for which there was population genetic evidence of recombination in

the PHI test (Table 1). Most of the lineage 1 isolates were collected in Asia (79%), but the lineage was

present in all continents in which the pathogen was sampled (Europe: one isolate; North America: 10,

Central and South America: 7, Africa: 22) (Figure 2). Clustering analysis on this single lineage with

sNMF detected four clusters with different geographic distributions (Figure 2). Estimates of

differentiation were lower between the clusters within lineage 1 ($F_{ST} < 0.49$) than between the main

lineages ($F_{ST} > 0.54$), consistent with a longer history of restricted gene flow between the main

lineages and a more recent history of admixture between clusters within lineage 1 (Supplementary

file 3; Figure 1C).

Two clusters within lineage 1 (referred to hereafter as *Baoshan* and *Yule*) consisted mostly of

isolates sampled from two different sites in the Yunnan province of China (Baoshan and Yule,

respectively, located about 370 km apart). The third cluster consisted mostly of isolates from Laos

and South China (referred to hereafter as *Laos*), and the fourth brought together 95% of the isolates

from lineage 1 collected outside Asia (referred to hereafter as *International*). In Asia, the *International*

cluster was found mostly in the Yunnan province of China, India and Nepal (Figure 2). PHI tests for

recombination rejected the null hypothesis of clonality in all four clusters (Table 1). Admixture was

widespread in lineage 1 (Figure 2), with most of the isolates (78%) displaying membership

proportions $q > 0.10$ for two or more clusters (Figure 2). Only five genotypes were detected in

multiple countries (genotype ID [number of countries]: 2 [6], 18 [2], 58 [2], 98 [3] and 254 [3],

corresponding to 41 isolates in total). All these genotypes belonged to the *International* cluster.

Table 1. PHI test for recombination (null hypothesis: clonality) and distribution of mating types and fertile females in

lineages, and in clusters within lineage 1. PHI tests were carried out using source files Figure 1 – source data 1 and Figure 2 –

source data 1. Information about mating types and female fertility is available in Supplementary file 1.

| Lineage/*Cluster* | *p*-value PHI test | Mat1.1/Mat1.2 (%) | Female-fertile isolates (%) |
|---|---|---|---|
| 1 | <0.0001 | 53/47 | 43 |
| *Baoshan* | <0.0001 | 69/31 | 67 |
| *International* | <0.0001 | 55/45 | 4 |
| *Laos* | <0.0001 | 68/32 | 37 |
| *Yule* | <0.0001 | 40/60 | 79 |
| 2 | 0.25 | 97/3 | 0 |
| 3 | 0.18 | 3/97 | 7 |
| 4 | 0.22 | 97/3 | 0 |

**Reproductive barriers caused by female sterility and postmating genetic incompatibilities**

Our analysis of genotyping data revealed the existence of a large, ancient, recombinant population

from which several clonal lineages have more recently emerged. This raised the question as to why

these populations became clonal, with no further gene flow. We addressed this question by

determining the capacity of the different lineages to engage in sexual reproduction, by characterizing

the distribution of mating types and the ability to produce female structures (i.e. female fertility).

Using *in vitro* crosses with tester strains, we found that lineages 2, 3 and 4 were composed almost

exclusively of a single mating type (97% of lineage 2 and 4 isolates tested carried the *Mat1.1* allele;

97% of lineage 3 isolates carried the *Mat1.2* allele), and had only small proportions (0-7%) of isolates

with female fertility (Table 1). By contrast to these clonal lineages, the mating type ratio was more

balanced in lineage 1 (52% of *Mat1.1*; Table 1), with *Mat1.1/Mat1.2* ratios ranging from 40/60 in the

*Yule* cluster to 69/31 in the *Baoshan* cluster. Female fertility rates were higher in three of the clusters

in lineage 1 (*Yule*: 79%; *Baoshan*: 67%; *Laos*: 37%; Table 1), but that in the International cluster (4%

female fertility) was as low as those in the clonal clusters (Table 1). These observations indicate that

there is a potential for sexual reproduction between lineages with compatible mating types and,

albeit rare, fertile females, but not between lineages 2 and 4, due to the low female fertility rates

and highly biased mating type ratios.

We further assessed the likelihood of sexual reproduction within and between lineages, by

evaluating the formation of sexual structures (perithecia), the production of asci (i.e. meiotic octads

within the perithecia) and the germination of ascospores (i.e. meiospores in meiotic octads) in

crosses between randomly selected isolates of opposite mating types from all four lineages (Figure 3).

This experiment revealed a marked heterogeneity in the rate of perithecium formation across

lineages. Three clusters in lineage 1 had the highest rates of perithecia formation, with isolates in the

*Yule* cluster, in particular, forming perithecia in 93% of crosses with isolates from the same cluster,

and in more than 46% of crosses with isolates from other lineages (Figure 3A). Isolates from lineages

2, 3 and 4 could only be crossed with isolates from other lineages, given their highly biased mating

type ratios, and the percentage of these crosses producing perithecia was highly variable (ranging

from 0% to 83%, depending on the lineages involved; Figure 3A). The rate of perithecium formation

was similar in crosses involving the *International* cluster of lineage 1 and in crosses involving clonal

lineages 2-4, and none of the crosses attempted between isolates from the *International* cluster led

to perithecium formation (Figure 3A). Perithecium dissection for a subset of crosses involving some of

the most fertile isolates revealed that most inter-lineage crosses produced perithecia that did not

contain asci or contained asci with low ascospore germination rates (Figure 3B). Whereas 100% of

crosses between isolates of the *Yule* cluster produced numerous germinating ascospores, ascospore

germination rates were only 33%, 56% and 7% in *Yule* x lineage 2, *Yule* x lineage 3 and *Yule* x lineage 4

crosses, respectively. Together, these results indicate that the clonal structure of lineages 2-4 is

caused by highly biased mating types and female sterility, and that the three clonal lineages and the

*International* cluster of lineage 1 are isolated from each other by strong pre- and early-postmating

barriers, including breeding system isolation (differences in mating type and female sterility), and a

combination of gametic incompatibility, zygotic mortality or hybrid non-viability. However, three of

the clusters in lineage 1 had biological features consistent with an ability to reproduce sexually,

suggesting possible sexual compatibility with clonal lineages 2-4 and the *International* cluster.

**Specialization to temperature conditions and rice subspecies**

Specialization can further reduce gene flow, and may facilitate niche separation between lineages,

which should favor the emergence of new lineages and their maintenance on the same host species

in conditions of competitive exclusion. The strong reproductive barriers demonstrated in our analysis

of crossing experiments would favor the establishment of such specialization, because reproductive

isolation facilitates adaptation to new conditions, by preventing maladaptive gene flow from

ancestral, non-adapted populations (Giraud *et al.* 2010). Given the very broad distribution of *P.*

*oryzae* and the strong environmental heterogeneity with which it is confronted in terms of the

nature of its hosts and the climate in which it thrives, we evaluated the variation of pathogen fitness

over different types of rice and temperature conditions.

We measured growth rate and sporulation rate of representative isolates cultured at different

temperatures to test the hypothesis of adaptation to temperature. Experiments were conducted with

41 isolates (lineage 1 [yule cluster]: 11; lineage 2: 10; lineage3: 10; lineage 4: 10). For all lineages,

mycelial growth rate increased with incubation temperature, although this trend was more visible

from 10°C to 15°C (increased mean mycelial growth of +2.22 mm/day) than from 25°C to 30°C (+0.05

mm/day) (growth curves and full statistical treatment of data are presented in Supplementary file 4;

data are reported in Supplementary file 5). Fitting a linear mixed-effects model with incubation time,

experimental replicate, and lineage of origin as fixed effects and isolate as a random effect, revealed a

significant lineage effect at each incubation temperature [10°C: $F_{(1,364)}=7988$, $p<0.001$; 15°C:

$F_{(1,419)}=33161$, $p<0.001$; 20°C: $F_{(1,542)}=40335$, $p<0.001$; 25°C: $F_{(1,413)}=30156$, $p<0.001$; 30°C:

$F_{(1,870)}=52681$, $p<0.001$]. Comparing least-squares means resulting from linear mixed-effects

models at each temperature revealed a significantly lower growth rate of lineage 4 at 10°C compared

to other lineages and a significantly higher growth rate of lineage 1 at 15°C, 20°C, 25°C and 30°C

compared to other lineages. As regards sporulation rates, sporulation was nearly absent at 10°C after

20 days of culture, with the few spores observed being not completely formed and divided by only

one septum instead of two septa in mature conidia (sporulation curves and full statistical treatment

of data are presented in Supplementary file 6; data are reported in Supplementary file 7). For all

lineages, sporulation rates increased with temperature from 15 to 25 °C and dropped at 30°C,

although isolates were cultured 7 days only at this latter temperature instead of 10 days at other

temperatures. Isolates were cultured 7 days only at 30°C because at this stage the mycelium had

reached the edges of the plates. Significant lineage effects were observed at 10°C, 15°C and 30°C

(Kruskal-Wallis tests; 10°C: H(3)=9.61, p=0.022; 15°C: H(3)=16.8, p<0.001; 30°C: H(3)=8.96, p=0.030).

Pairwise non-parametric comparisons using Dunn's test of lineages revealed significant differences

between lineages 1 and 2 at 10°C, between lineage 1 and lineages 3 and 4 at 15°C, and between

lineage 1 and 4 at 30°C.  Together, measurements of mycelial growth and sporulation at different

temperatures revealed differences in performance between lineages, but no clear pattern of

temperature specialization.

We assessed the importance of adaptation to the host, by challenging 45 rice varieties

representative of the five main genetic groups of Asian rice (i.e. the temperate japonica, tropical

japonica, aus, aromatic and indica subgroups of *Oryza sativa*) with 70 isolates representative of the

four lineages of *P. oryzae* and the four clusters within lineage 1 (Supplementary file 8). Interaction

phenotypes were assessed qualitatively by scoring resistance (from full to partial resistance: scores 1

to 3) or disease symptoms (from weak to full susceptibility: scores 4 to 6) and analysed by fitting a

proportional-odds model, followed by an analysis of variance (full statistical treatment of data is

presented in Supplementary file 9). This analysis revealed significant differences between groups of

isolates ($\chi^2$(6)=100, p<0.001), and between rice genetic groups ($\chi^2$(4)=161, p<0.001), and a significant

interaction between these two variables ($\chi^2$(24)=97, p<0.001). The finding of a significant interaction

between groups of isolate (lineage or cluster) and rice types indicates that the effect of the group of

isolates on the proportion of compatible interactions differed between rice types, suggesting

adaptation to the host. This is also supported by the specific behaviour of certain lineages. Isolates

from lineage 2 showed e.g. much lower symptom scores than the other lineages on all rice types

except temperate japonica and the isolates of the *Yule* cluster possess particularly high virulence on

indica varieties (Figure 4A; Supplementary file 8). In comparisons of rice genetic groups, significantly

higher symptom scores were observed on temperate japonica rice than on the other types of rice,

whereas the varieties of the aromatic genetic group were significantly more resistant to rice blast

(Figure 4B). Together, these experiments therefore revealed significant differences in host range

between lineages, but this specialization to the host is not strict since host ranges overlap (Figure 4C-

G). Adaptation to the host or small differences in temperature optima could, nonetheless, further

increase niche separation and reduce gene flow between lineages, via pre-mating barriers such as

immigrant non-viability (i.e., lower rates of contact between potential mates due to the mortality of

immigrants; (Gladieux *et al.* 2011; Nosil *et al.* 2005)), and post-mating barriers, such as ecological

hybrid non-viability (i.e., lower survival of ill-adapted hybrid offspring).

**Differences in number and genetic variability of putative effector genes among *P. oryzae* lineages**

Small secreted proteins named effectors are the most prominent class of fungal virulence factors.

They act by manipulating host cellular pathways and are key determinants of the host range and

fitness on compatible hosts (Liao *et al.* 2016; Yoshida *et al.* 2016). We therefore determined the

differences of effector repertoires in *P. oryzae* lineages in terms of presence/absence and nucleotide

polymorphism, and analyzed whether it is a particularly dynamic fraction of the gene space. For this

aim, we used whole-genome sequencing data for 123 isolates, including 29 isolates sequenced in this

study and 94 publicly available genomes (Gladieux *et al.* 2018a; Pordel *et al.* 2020; Zhong *et al.* 2018),

of which 33 were also genotyped using our Infinium genotyping beadchip (Supplementary file 10;

Appendix 1). Clustering analysis indicated that this dataset covered the four lineages of *P. oryzae*, and

three of the clusters of lineage 1 (Laos, International and Baoshan; Appendix 1; Supplementary file

10). Assembly lengths for the 123 sequenced isolates varied from 36.7 to 39.6 Mb, with an average of

38.1 Mb (Supplementary file 10). The number of assembled contigs longer than 500bp ranged from

1010 to 2438, and the longest contig was 1.1 Mb (Supplementary file 10).

Gene prediction identified 11,684 to 12,745 genes per genome, including 1,813 to 2,070

genes predicted to code for effector proteins. We found a significant effect of the lineage of origin on

both the mean number of putative effector genes (ANOVA, F=5.54, p=0.0014), and the mean number

of non-effector genes (ANOVA, F=7.95, p<0.001). Multiple comparisons revealed that mean numbers

of putative effectors and non-effector genes were only significantly different between genomes of

lineage 3 and lineages 2 (Tukey's HSD test; p<0.001; Supplementary file 11). Orthology analysis

identified 14,573 groups of orthologous sequences (i.e. orthogroups), and we applied a rarefaction

approach to the table of orthology relationships to estimate the size of accessory and core genomes

in lineages while accounting for differences in sample size. For both putative effectors and the

remaining of the gene space, we found that the number of core and accessory genes did not plateau

and changed almost linearly with the number of genomes resampled, indicating that the number of

core genes is likely substantially smaller, and the number of accessory genes substantially higher, than

estimated from 123 genomes (Supplementary file 12). With pseudo-samples of size n=30 per lineage

(i.e. excluding lineage 4, due to small sample size), the gene content was highly variable within

lineages with only 61-71% of all predicted effectors, and 68-73% of the remaining gene space, being

conserved in lineages 1-3 (Figure 5A). Despite extensive variation in gene content within lineages,

clustering of isolates based on presence/absence variation for both putative effectors and the

remaining of the gene space was highly similar to clustering based on 3,868 SNPs (Supplementary file

13), indicating that variation in gene content and SNP allelic variation reflect similar genealogical

processes.

To identify putative effectors that may be involved in host specialization of rice-infecting

lineages of *P. oryzae*, we first identified orthogroups with distinct patterns of presence/absence

across lineages. Principal component analysis of presence/absence data identified 72 orthogroups of

putative effectors that explained 95% of the variance of the three principal components that

differentiate the four lineages (PC1, PC2 and PC6; Supplementary file 13; Figure 5B), and may

contribute to the differences in host range among lineages. Interestingly, two among them, AvrRmg8

(=OG0011611) and PWL3 (=OG0011928), are host range determinants in wheat-infecting *P. oryzae*

since they trigger immunity in wheat varieties carrying the resistance genes *Rmg8* (Anh *et al.* 2018) or

*PWT3* (Inoue *et al.* 2017; Kang *et al.* 1995).

As host specialization can also involve sequence divergence at effector proteins, we also

scanned the corresponding genes for signatures of diversifying selection, i.e. an excess of non-

synonymous nucleotide diversity (ratio of non-synonymous to synonymous nucleotide diversity $\pi_N/\pi_S$

>1). We identified 185 orthogroups with $\pi_N/\pi_S$ > 1 in at least one lineage, including 164 orthogroups

with $\pi_N/\pi_S$ > 1 in only one lineage (lineage 1: 131 orthogroups; lineage 2: 18; lineage 3: 10; lineage 4:

5), and twelve, seven and two orthogroups with $\pi_N/\pi_S$ > 1 in two, three or four lineages, respectively

(Supplementary file 14). None of these orthogroups corresponded to effectors previously

characterized as being involved in *P. oryzae* virulence against rice or other *Poaceae* hosts.

**Geographic and climatic differentiation in the distribution of *P. oryzae* lineages**

Our tests of adaptation to host and temperature under controlled conditions showed that

specialization was not strict, but it remained possible that fitness differences between lineages would

be sufficient under natural conditions to induce separation in different ecological niches, and/or that

our experimental conditions did not capture the full extent of the phenotypic differences. Here, we

tested the hypothesis that lineages thrive in geographically and/or environmentally different

conditions, which would provide indirect evidence for specialization to different habitats. We tested

this hypothesis by collecting geographic and climatic data for all isolates, approximating to regions or

the nearest cities when the exact GPS position of the sampling area was not available (Supplementary

file 1). At a larger scale, clonal lineages 2 and 3 were both widespread, with lineage 2 found on all

continents, and lineage 3 on all continents except Europe. Lineage 2 was the only lineage sampled in

Europe (with the exception of a single isolate from lineage 1), whereas lineage 3 was more

widespread in intertropical regions. Lineage 4 was mostly found in South Asia (India, Bangladesh,

Nepal), and in the USA and Africa (Benin, Tanzania). The recombining lineage, lineage 1, was found in

all continents, but was mostly found in Asia (79% of all isolates and 94% of all genotypes). At a

smaller scale, two, or even three different lineages were sampled in the same year in the same city in

11 different countries from all continents. However, in only one instance did two isolates from

different lineages have identical GPS positions (isolates US0106 and US0107 sampled from the same

field). Thus, lineages had different, but partially overlapping distributions, including some overlap

with the distribution of the sexually reproducing lineage (lineage 1).

We then investigated whether differences in the geographic range of lineages were

associated with differences in climatic variables. By plotting sampling locations onto a map of the

major climate regions (Kottek *et al.* 2006) we were able to identify clear differences in the climatic

distributions of lineages 2 and 3, with lineage 2 mostly found in warm temperate climates and lineage

3 found in equatorial climates (Figure 6; Source code 1). We used the outlying mean index (OMI),

which measures the distance between the mean habitat conditions used by a lineage and the mean

habitat conditions used by the entire species, to test the hypothesis that different lineages are

distributed in regions with different climates. Using information for 19 climatic variables (biomes)

from the WorldClim bioclimatic variables database (Fick & Hijmans 2017) for all sampling locations,

we obtained statistically significant results, in a permutation test on OMI values, for lineages 2, 3 and

4 (Source code 2; permutation test: lineage 1: OMI=2.14, $p$=0.620; lineage 2: OMI=6.54, $p$<0.001;

lineage 3: OMI=2.08, $p$<0.001; lineage 4: OMI=13.73, $p$=0.017). The OMI analysis, in which the first

two axes accounted for 69% and 25% of the variability, respectively, revealed that lineage 2 was more

frequent in regions with a high annual temperature range (biome 7) or a high degree of seasonality

(biome 4); lineage 4 was associated with regions with high levels of seasonal precipitation (biomes

13, 16 and 18), and lineage 3 was more frequent in regions with high temperatures (biomes 1, 6, 10

and 11) and high levels of isothermality (biome 3), characteristic of tropical climes (Source code 2;

Figure 6; Supplementary file 15).

**DISCUSSION**

We describe here the population structure of the rice blast fungus, and the eco-evolutionary factors

underlying the emergence and maintenance of multiple divergent lineages in this widespread

pathogen. Our analysis of SNP genotyping data showed that *P. oryzae* could be subdivided into four

lineages. Two of the lineages previously detected in an analysis of whole-genome data, and

represented by only a few individuals (lineages 5 and 6; (Gladieux *et al.* 2018b)), were assigned to

lineage 1 in our analysis, and may correspond to a subdivision of lineage 1. Differences in the number

of lineages detected may reflect an ascertainment bias, given that our SNP-genotyping beadchip was

designed with a set of genomic sequences that include few representatives of lineages 5 and 6.

Alternatively, the sampling used in the previous study (Gladieux *et al.* 2018b) may not have been

dense enough to reveal the lack of differentiation between lineages 1, 5 and 6, making them appear

to be different entities. Consistent with previous findings, our analysis of the genealogical

relationships between isolates revealed three clonal groups connected by long branches to a central

recombining group. The finding of an internationally distributed cluster in lineage 1 sheds light on the

early evolutionary changes contributing to the emergence of clonal lineages in this pathogen. The

*International* cluster of lineage 1 displayed genetic footprints of recombination and sexual

reproduction (in the form of an excess of homoplasious variants or relatively balanced mating types),

but the signal of recombination detected here may be purely historical, because we also found that

the clonal fraction and the frequency of sterile females were very high in this cluster. The loss of

female fertility may, therefore, be an early and major cause of the shift towards asexual reproduction.

Some genotypes assigned to the international cluster of lineage 1 may, in the future, come to

resemble other clonal lineages in analyses of genealogical relationships, and form a long branch

stemming from the central lineage 1, particularly for genotypes already displaying clonal propagation,

such as genotypes 2, 18, 58, 98 and 254. However, according to the principle of competitive

exclusion, competition between clonal lineages should only allow hegemonic expansion of the most

competitive clonal groups, unless they separate into different ecological niches. It may be this process

that ultimately leads to the fixation of a single mating type, as in clonal lineages 2-4.

We found that female sterility and intrinsic genetic incompatibilities represented strong

barriers to gene flow between the clonal lineages, potentially accounting for their maintenance over

time, without fusion, despite the compatibility of mating-types (at least between lineages 2 and 4

[*Mat1.1*] on the one hand, and lineage 3 [*Mat1.2*] on the other). These barriers to gene flow may

have contributed to the establishment of lineage specialization (because reproductive isolation

facilitates adaptation by limiting the flow of detrimental ancestral alleles in populations adapting to a

new habitat), accounting for the maintenance of the clonal lineages on the same host species, when

faced with competitive exclusion. Our analyses also show that, despite widely overlapping large-scale

distributions, the different lineages are essentially found in different regions, with different climatic

characteristics, when their distributions are observed at a finer scale. Lineage 1 was mostly sampled

in Southeast Asia, lineage 4 predominated in India, and lineages 2 and 3 had global distributions, but

the geographic ranges of all these lineages overlapped. However, an analysis of climatic data

indicated that lineage 2 predominated in temperate climates, where temperate japonica rice is

grown, lineage 3 predominated in tropical climates, in which indica rice is grown, and lineage 4

predominated in regions with high levels of seasonal precipitation in which the indica and aromatic

rice types are the principal rice types grown. Despite the finding of separation in different climatic

regions, our experiments revealed no strong differences between lineages in terms of sporulation

and mycelial growth on synthetic media at different temperatures. This suggests that, if adaptation to

temperature occurs in this pathogen, it was not measured by our experiments, either because it

involves traits other than sporulation and hyphal growth, or because the *in vitro* conditions were not

suitable to demonstrate differences. The host range varied across lineages, but all host ranges

overlapped, indicating that host specialization was not strict, or was not fully captured by our

experiments. Adaptation to the host may, nevertheless, decrease gene flow between populations

even further. For instance, lineage 2 had a narrow host range, potentially limiting the opportunities

for encounters and mating with other groups.

Effector repertoires play a key role in pathogen specialization, in fungal pathogens in general

(Schulze-Lefert & Panstruga 2011), and *P. oryzae* in particular (Asuke *et al.* 2020; Couch *et al.* 2005;

Liao *et al.* 2016; Sweigard *et al.* 1995; Takabayashi *et al.* 2002). Previous work suggested that clonal

lineages of *P. oryzae* possess smaller effector repertoires (Latorre *et al.* 2020) and that the clonal

lineage associated with japonica rice (lineage 2) possess more effectors, and in particular Avr-

effectors, than other clonal lineages (Latorre *et al.* 2020; Liao *et al.* 2016) . Our analyzes on a larger

set of putative effectors (ca. 2000 in our study, vs 13 Avr-effectors in (Liao *et al.* 2016) and 178

known and candidate effectors in (Latorre *et al.* 2020)) confirm this trend, since lineage 2 associated

with temperate japonica exhibits significantly more putative effectors than lineage 3 associated with

indica. However, the number of non-effector genes is also greater in lineage 2 than in lineage 3, and

thus it remains possible that the larger effector complement of lineage 2 is a mere consequence of a

larger number of genes. Besides differences between lineages in richness of putative effector

repertoires, we show that patterns of presence / absence variation mirror patterns of population

subdivision based on SNP allelic variation, and thus that the differential sorting of both nucleotide

polymorphism and gene content across lineages reflect similar genealogical processes. Our

multivariate analyzes identify 72 effectors that most strongly contribute to the differentiation of the

4 lineages in terms of presence / absence. It is possible that the frequency differences observed at

these 72 effectors were due to chance events during bottlenecks at the onset of lineage formation,

or alternatively that their differential loss was important for the initial adaptation of lineages to new

rice populations or subspecies. Nucleotide diversity at synonymous and non-synonymous sites also

identify 185 effectors with signatures of diversifying selection in one or more lineage, possibly

mediated by coevolutionary interactions with host molecules. In the future, it will be interesting to

determine the molecular targets of these effectors and decipher the relationship between their
polymorphism and their mode of action.

Our results also indicate that lineage 1 may pose a major threat to rice production. Most
effectors with signature of diversifying selection were identified in lineage 1, highlighting the higher
ability of this lineage to rapidly fix advantageous mutations. Eleven of the 12 most multivirulent
isolates (lesion type > 2 on more than 40 varieties tested) belonged to lineage 1, and the *Yule* cluster,
in particular, was highly pathogenic on indica varieties. The propagation of such highly pathogenic
genotypes, belonging to a lineage that has both mating types and fertile females, should be
monitored closely.  The monitoring of lineage 1 is all the more critical because this lineage has an
intermediate geoclimatic distribution that overlaps largely with the distributions of the other three
lineages, and some of the attempted crosses between clonal lineage 2-4 isolates and recombining
lineage 1 isolates produced viable progeny. This finding confirms the possibility of gene flow into this
lineage, as previously demonstrated on the basis of admixture mapping (Gladieux *et al.* 2018b), but it
also raises the question of the threat posed by gene flow from lineage 1, which is highly pathogenic,
to the other lineages.

In conclusion, our study of genetic and phenotypic variation within and between clonal
lineages, and within the recombinant lineage of *P. oryzae* suggests a scenario for the emergence of
widespread clonal lineages. The loss of female fertility may be a potent driver of the emergence of
asexually reproducing groups of clones. The reproductive isolation generated by the loss of sex and
the accumulation of mutations due to the absence of sexual purging would facilitate the
specialization of some of these clonal groups, leading to the competitive exclusion of the least
efficient clonal groups, and, finally, to the propagation of clonal lineages fixed for a single mating
type. Our results, thus, demonstrate that the spread of a pathogen across heterogeneous habitats
and divergent populations of a crop species can lead to niche separation and incipient speciation in
the pathogen.

**MATERIALS AND METHODS**

**Biological material**

We chose 886 *P. oryzae* isolates collected on Asian rice between 1954 and 2014 as representative of

the global genetic diversity of the fungus. Isolates were selected on the basis of microsatellite data, to

maximize the number of multilocus genotypes represented ((Saleh *et al.* 2014) and unpublished

data), or based on geographic origin in the absence of genotypic data, to maximize the number of

countries represented in the dataset.

Sixty-eight isolates were selected for experimental measurements of reproductive success,

adaptation to host, and growth and sporulation experiments at different temperatures

(Supplementary file 1). This subset of isolates included 10 isolates from each of the three clonal

lineages (lineages 2-4), 27 isolates from the various clusters within lineage 1 [*Baoshan* (9 isolates),

*International* (10), *Laos* (8), and *Yule* (11)].

Forty-four varieties were chosen as representative of the five main genetic subgroups of

Asian rice (Garris *et al.* 2005): indica (Chau, Chiem chanh, DA11, De abril, IR8, JC120, Pappaku), aus

(Arc 10177, Baran boro, Black gora, DA8, Dholi boro, Dular, FR13 A, JC148, Jhona 26, Jhona 149,

Kalamkati, T1, Tchampa, Tepi boro), temperate japonica (Aichi asahi, Kaw luyoeng, Leung pratew,

Maratelli, Nep hoa vang, Nipponbare, Sariceltik, Som Cau 70A), tropical japonica (Azucena,

Binulawan, Canella de ferro, Dholi boro, Gogo lempuk, Gotak gatik, Trembese) and aromatic (Arc

10497, Basmati lamo, Dom zard, Firooz, JC1, Kaukkyisaw, N12). The Maratelli (temperate japonica)

and CO39 (indica) varieties were used as susceptible controls.

**Genotyping**

*Pyricularia oryzae* isolates were genotyped at 5,657 genomic positions with an Illumina Infinium

beadchip microarray carrying single-nucleotide polymorphisms identified in 49 genomes of rice- and

barley-infecting *P. oryzae* isolates previously characterized (Gladieux et al., 2018b). The final dataset

included 3,686 biallelic SNPs without missing data.

**Whole genome sequencing**

Twenty-nine isolates were sequenced using Illumina HiSeq 3000 (Supplementary file 10). Isolates

were grown on rice flour-agar medium for mycelium regeneration, then in liquid rice flour medium

(Adreit *et al.* 2007). Genomic DNA extraction was carried out using >100 mg of fresh mycelium from

liquid culture. Fresh mycelium dried on Miracloth paper was crushed in liquid nitrogen. Nucleic acids

were subsequently extracted using an extraction buffer (2 % CTAB – 1.4 M NaCl – 0.1 M Tris-HCl pH 8

– 20 mM EDTA pH 8 added before use with 1 % final of $Na_2SO_3$), then purified with a

chloroform:isoamyl alcohol 24:1 treatment, precipited overnight in isopropanol, and rinsed with 70%

ethanol. The extracted nucleic acids were further treated with Rnase A (0.2mg/mL final) and purified

with another chloroform:isoamyl alcohol 24:1 treatment followed by an overnight ethanol

precipitation. The concentration of extracted genomic DNA was assessed on Qubit® using the dsDNA

HS Assay Kit. The purity of extracted DNA was checked by verifying that the 260/280 and 260/230

absorbance ratios measured with NanoDrop were between 1.8 and 2.0. Preparation of TruSeq nano

libraries and HiSeq3000 sequencing (150 nucleotide reads, and 500 bp insert size) was carried out at

GeT-PlaGe (INRAE, Toulouse, France).

**Genome assembly, gene prediction, orthology analysis, effector prediction, and summary statistics**

**of nucleotide variation**

For the 123 sequenced isolates included in the dataset, low-quality reads were removed using the

software CUTADAPT (Martin 2011). Reads were assembled using ABYSS 2.2.3 (Jackman *et al.* 2017;

Simpson *et al.* 2009) using different K-mer sizes and for each isolate we chose the assembled

sequence with the highest N50 for further analyses. Genes were predicted with BRAKER 2.1.5 (Hoff *et*

*al.* 2015; Hoff *et al.* 2019) using RNAseq data (Pordel *et al.* 2020) as extrinsic evidence for model

refinement. Genes were also predicted with AUGUSTUS 3.4.0 (Stanke & Morgenstern 2005)(training

set=*Magnaporthe grisea*) and gene models that did not overlap with gene models identified with

BRAKER were added to the GFF file generated with the latter. Repeated regions were masked using

REPEATMASKER 4.1.0 (http://www.repeatmasker.org/). The completeness of genome assemblies and

gene predictions was assessed using BUSCO (Simão *et al.* 2015), and completeness in BUSCO genes

was greater than 93.7% (Supplementary file 10). Putative effector genes were identified as genes

encoding proteins predicted to be secreted by at least two methods among three [SIGNALP 4.1

(Nielsen 2017), TARGETP (Emanuelsson *et al.* 2000) and PHOBIUS (Käll *et al.* 2004)], without predicted

transmembrane domain based on TMHMM analysis (Krogh *et al.* 2001), without predicted motif of

retention in the endoplasmic reticulum based on PS-SCAN (Bhagwat & Aravind 2007), and without

CAZy annotation based on DBSCAN V7 (Yin *et al.* 2012). Differences in numbers of putative effectors

and non-effectors among lineages (ANOVA and Tukey's HSD test) were tested using using the SCIPY

1.6.0 package in PYTHON. Homology relationships among predicted genes were established using

ORTHOFINDER v2.4.0 (Emms & Kelly 2015). Principal component analysis of presence/absence

variation of putative effectors and non-effector proteins was carried out using R package PRCOMP. We

used the get_pca_var() in PRCOMP to extract the results for variables (i.e. effectors) and identify

effectors whose contribution to principal components was the greatest, defined as effectors with a

contribution to loadings of PC1, PC2 and PC6 greater than 1%. To estimate the numbers of core and

accessory genes, we used a rarefaction approach to account for differences in sample size across

lineages. For each pseudo-sample size, we estimated the size of the core and accessory genome in

each lineage using a maximum of 2000 pseudosamples. Sequences for each orthogroup were aligned

and cleaned with TRANSLATORX (Abascal *et al.* 2010) using default parameters. The ratio of non-

synonymous to synonymous diversity $\pi_N/\pi_S$ was calculated for each orthogroup for lineages with

sample size greater than or equal to four using EGGLIB 3 ([https://www.egglib.org](https://www.egglib.org)).

### Population subdivision and recombination

Among the 886 *P. oryzae* isolates genotyped, we identified 264 different multilocus genotypes, which

were used for the analysis of population subdivision. We used the SNMF program to infer individual

ancestry coefficients in *K* ancestral populations. This program is optimized for the analysis of large

datasets and does not assume Hardy-Weinberg equilibrium. It is, therefore, more appropriate to deal

with inbred or clonal lineages (Frichot *et al.* 2014) . We used SPLITSTREE version 4 (Huson & Bryant

2006) to visualize relationships between genotypes in a phylogenetic network, with reticulations to

represent the conflicting phylogenetic signals caused by homoplasy. We also used the pairwise

homoplasy index (PHI) test implemented in SPLITSTREE to test the null hypothesis of clonality. We used

the SNMF software to investigate population subdivision further within the cluster for which the null

hypothesis of clonality could be rejected. Weir and Cockerham's $F_{ST}$ was calculated with the R package

HIERFSTAT, by the WC84 method (Goudet 2005).

### Experimental measurement of reproductive success and female fertility

*Pyricularia oryzae* is a heterothallic fungus with two mating types (*Mat1.1* and *Mat1.2*). Sexual

reproduction between strains of opposite mating type can be observed in laboratory conditions, and

results in the production of ascospores within female sexual structures called perithecia (Saleh *et al.*

2012a). On synthetic media, perithecia are formed at the contact zone between parental mycelia.

Crosses were carried out on a rice flour agar medium (20 g rice flour, 2.5 g yeast extract, 15 g agar

and 1 L water, supplemented with 500 000 IU penicillin G after autoclaving for 20 min at 120°C), as

described by (Saleh *et al.* 2012a). We assessed reproductive success by determining the production of

perithecia produced after three weeks of culture at 20°C under continuous light. Each cross was

carried out twice and we computed the average number of perithecia across repeats. Perithecia can

be formed by the two interacting partners or by one partner only. Isolates forming perithecia are

described as female-fertile. We measured female fertility for 220 isolates (listed in **Supplementary file**

1), by monitoring perithecium production in crosses involving the tester strains CH0997 (Mat1.2) and

CH0999 (Mat1.1). We further assessed the presence of asci and germinating ascospores in perithecia

for a subset of crosses, by excising perithecia with a scalpel and counting the number of germinated

filaments for each individual ascus after overnight incubation on water agar. The subset of crosses

included 10 Mat1.1 isolates (lineage 1: CH0999, CH1065, CH1076; lineage 2: CH0092, MC0016,

SP0006; lineage 4: IN0017, IN0092, NP0070, CH0718) and 6 Mat1.2 isolates (lineage 1: CH0997;

CH1083, CH1120; lineage 3: BR0019, CH0549, MD0929).

**Pathogenicity tests**

Compatibility between *P. oryzae* isolates and rice plants from the five main genetic subgroups of rice

(indica, temperate japonica, tropical japonica, aus and aromatic) was measured in controlled

conditions. Seventy isolates were inoculated on 46 varieties. Inoculations were performed as

described by (Gallet *et al.* 2016). Conidial suspensions (25 000 conidia.mL$^{-1}$) in 0.5% gelatin were

sprayed onto three-week-old rice seedlings (> 6 plants/variety). The inoculated plants were incubated

for 16 hours in the dark at 27°C and 100% humidity and then for seven days with a day/night

alternation (13 hours at 27°C/11 hours at 21°C), before scoring symptoms. Lesion type was rated from

1 to 6 (Gallet *et al.* 2016) and the symptom type was assessed visually on leaves. Each interaction was

assessed in three independent experiments, and the maximum of the three scores was used in

calculations.

**Mycelial growth and sporulation rate at different temperatures**

Mycelial growth and sporulation rate were measured for 41 isolates at five different temperatures

(10°C, 15°C, 20°C, 25°C and 30°C). For each isolate, Petri dishes containing PDA medium were

inoculated with mycelial plugs placed at the center, and incubated in a growth chamber with a fixed

temperature. Mycelium diameter was estimated as the average of two measurements made along

two perpendicular axes at different time points. At the end of the experiment, conidia were collected

by adding 5 mL of water supplemented with 0.01% Tween 20 to the Petri dish and rubbing the

surface of the mycelium. Conidia were counted with a hemocytometer. Three or four independent

experiments were performed for each isolate, at each temperature. We had initially planned to carry

out only three independent experiments. However, because some isolates did not grow in the first

experiment, and because some cultures were invaded by mites and had to be discarded in the second

and third experiment, a fourth experiment was eventually performed for some temperature

conditions.

**Statistical analyses of climatic and phenotypic data**

<u>Ecological niche separation:</u> OMI (outlying mean index), or marginality, is used to study niche

separation and niche breadth. OMI gives the same weight to all samplings, whether rich or poor in

species and individuals, and is particularly suitable in cases in which sampling is not homogeneous.

OMI measures the deviation between the mean environmental conditions used by one lineage and

the mean environmental conditions used by all lineages. OMI analysis then places the lineages in

environmental conditions maximizing their OMI. Environmental values, consisting of 19 biome values

(WorldClim bioclimatic variables (Fick & Hijmans 2017)), were retrieved for all sampling locations.

Climatic values were normalized before the analysis. A contingency table was generated to associate

the number of isolates from each lineage with each sampling location. Only isolates with a precise

location (known region, city or GPS position of sampling) were included. A random permutation test,

with 10000 permutations, was used to assess the statistical significance of marginality for each

lineage.

<u>Pathotyping:</u> Ordinal symptoms scores were analyzed with a proportional-odds model

accounting for ordered categorical responses with a clm() function implemented in the ordinal R

package (version 2018.8-25). The significance of the factors was analyzed by ANOVA. Pairwise

comparisons of significant factors were performed after computing least-squares means with LSMEANS

version 2.30-0 in R, with Tukey adjustment.

<u>Mycelium growth:</u> Mycelium growth was analyzed separately for each temperature, with a

linear mixed-effects model. The significance of the factors was analyzed by ANOVA. Post-hoc pairwise

comparisons were performed after computing least-squares means with LSMEANS version 2.30-0 in R.

<u>Sporulation:</u> The median number of spores calculated for the various repetitions of the

experiment was analyzed with a Kruskal-Wallis test. Post-hoc pairwise comparisons were performed

using Dunn's non-parametric multiple comparison test.

**ACKNOWLEDGMENT**

We thank Tatiana Giraud for useful suggestions. We thank all our colleagues who shared strains or

participated in the collection of rice blast samples.

**REFERENCES**

Abascal F, Zardoya R, Telford MJ (2010) TranslatorX: multiple alignment of nucleotide sequences
guided by amino acid translations. *Nucleic Acids Research* **38**, W7-W13.
Abbate JL, Gladieux P, Hood ME*, et al.* (2018) Co-occurrence among three divergent plant-castrating
fungi in the same Silene host species. *Molecular ecology* **27**, 3357-3370.
Adreit H, Santoso, Andriantsimialona D*, et al.* (2007) Microsatellite markers for population studies of
the rice blast fungus, Magnaporthe grisea. *Molecular Ecology Notes* **7**, 667-670.
Ali S, Gladieux P, Leconte M*, et al.* (2014) Origin, migration routes and worldwide population genetic
structure of the wheat yellow rust pathogen Puccinia striiformis f. sp. tritici. *PLoS pathogens*
**10**, e1003903.
Anh VL, Inoue Y, Asuke S*, et al.* (2018) Rmg8 and Rmg7, wheat genes for resistance to the wheat blast
fungus, recognize the same avirulence gene AVR-Rmg8. *Molecular Plant Pathology* **19**, 1252-
1256.

Asuke S, Tanaka M, Hyon G-S, *et al.* (2020) Evolution of an Eleusine-specific subgroup of Pyricularia
oryzae through a gain of an avirulence gene. *Molecular Plant-Microbe Interactions* **33**, 153-
165.
Bhagwat M, Aravind L (2007) Psi-blast tutorial. In: *Comparative genomics*, pp. 177-186. Springer.
Bueker B, Eberlein C, Gladieux P, *et al.* (2016) Distribution and population structure of the anther
smut M icrobotryum silenes-acaulis parasitizing an arctic–alpine plant. *Molecular ecology* **25**,
811-824.
Burdon JJ (1993) The structure of pathogen populations in natural plant communities. *Annual review*
*of phytopathology* **31**, 305-323.
Couch BC, Fudal I, Lebrun M-H, *et al.* (2005) Origins of Host-Specific Populations of the Blast
Pathogen *Magnaporthe oryzae* in Crop Domestication With Subsequent Expansion of
Pandemic Clones on Rice and Weeds of Rice. *Genetics* **170**, 613-630.
Emanuelsson O, Nielsen H, Brunak S, Von Heijne G (2000) Predicting subcellular localization of
proteins based on their N-terminal amino acid sequence. *Journal of molecular biology* **300**,
1005-1016.
Emms DM, Kelly S (2015) OrthoFinder: solving fundamental biases in whole genome comparisons
dramatically improves orthogroup inference accuracy. *Genome Biology* **16**, 157.
Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land
areas. *International journal of climatology* **37**, 4302-4315.
Frichot E, Mathieu F, Trouillon T, Bouchard G, François O (2014) Fast and efficient estimation of
individual ancestry coefficients. *Genetics* **196**, 973-983.
Gallet R, Fontaine C, Bonnot F, *et al.* (2016) Evolution of compatibility range in the rice–
magnaporthe oryzae system: an uneven distribution of R genes between rice subspecies.
*Phytopathology* **106**, 348-354.
Garris AJ, Tai TH, Coburn J, Kresovich S, McCouch S (2005) Genetic structure and diversity in Oryza
sativa L. *Genetics* **169**, 1631-1638.
Gibson AK, Hood ME, Giraud T (2012) Sibling competition arena: selfing and a competition arena can
combine to constitute a barrier to gene flow in sympatry. *Evolution: International Journal of*
*Organic Evolution* **66**, 1917-1930.
Giraud T, Gladieux P, Gavrilets S (2010) Linking the emergence of fungal plant diseases with
ecological speciation. *Trends in ecology & evolution* **25**, 387-395.
Giraud T, Koskella B, Laine AL (2017) Introduction: microbial local adaptation: insights from natural
populations, genomics and experimental evolution. *Molecular ecology* **26**, 1703-1710.
Gladieux P, Condon B, Ravel S, *et al.* (2018a) Gene Flow between Divergent Cereal- and Grass-Specific
Lineages of the Rice Blast Fungus Magnaporthe oryzae. *mBio* **9**.
Gladieux P, Guerin F, Giraud T, *et al.* (2011) Emergence of novel fungal pathogens by ecological
speciation: importance of the reduced viability of immigrants. *Molecular ecology* **20**, 4521-
4532.
Gladieux P, Ravel S, Rieux A, *et al.* (2018b) Coexistence of multiple endemic and pandemic lineages of
the rice blast pathogen. *MBio* **9**.
Goudet J (2005) Hierfstat, a package for R to compute and test hierarchical F-statistics. *Molecular*
*Ecology Notes* **5**, 184-186.
Hardin G (1960) The competitive exclusion principle. *Science* **131**, 1292-1297.
Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M (2015) BRAKER1: unsupervised RNA-Seq-
based genome annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics* **32**, 767-769.
Hoff KJ, Lomsadze A, Borodovsky M, Stanke M (2019) Whole-genome annotation with BRAKER. In:
*Gene Prediction*, pp. 65-95. Springer.
Huson DH, Bryant D (2006) Application of Phylogenetic Networks in Evolutionary Studies. *Molecular*
*Biology and Evolution* **23**, 254-267.
Inoue Y, Vy TTP, Yoshida K, *et al.* (2017) Evolution of the wheat blast fungus through functional losses
in a host specificity determinant. *Science* **357**, 80-83.

Jackman SD, Vandervalk BP, Mohamadi H, *et al.* (2017) ABySS 2.0: resource-efficient assembly of
large genomes using a Bloom filter. *Genome Research* **27**, 768-777.
Käll L, Krogh A, Sonnhammer ELL (2004) A combined transmembrane topology and signal peptide
prediction method. *Journal of molecular biology* **338**, 1027-1036.
Kang S, Sweigard Ja Fau - Valent B, Valent B (1995) The PWL host specificity gene family in the blast
fungus Magnaporthe grisea. *Molecular Plant Microbe Interactions*.
Kottek M, Grieser J, Beck C, Rudolf B, Rubel F (2006) World map of the Köppen-Geiger climate
classification updated.
Krogh A, Larsson B, Von Heijne G, Sonnhammer ELL (2001) Predicting transmembrane protein
topology with a hidden Markov model: application to complete genomes. *Journal of*
*molecular biology* **305**, 567-580.
Latorre SM, Reyes-Avila CS, Malmgren A, *et al.* (2020) Differential loss of effector genes in three
recently expanded pandemic clonal lineages of the rice blast fungus. *BMC biology* **18**, 1-15.
Liao J, Huang H, Meusnier I, *et al.* (2016) Pathogen effectors and plant immunity determine
specialization of the blast fungus to rice subspecies. *Elife* **5**, e19377.
Marin M, Preisig O, Wingfield BD, Kirisits T, Wingfield MJ (2009) Single sequence repeat markers
reflect diversity and geographic barriers in Eurasian populations of the conifer pathogen
Ceratocystis polonica. *Forest Pathology* **39**, 249-265.
Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
*EMBnet. journal* **17**, 10-12.
Nielsen H (2017) Predicting secretory proteins with SignalP. In: *Protein function prediction*, pp. 59-73.
Springer.
Nosil P, Vines TH, Funk DJ (2005) Perspective: reproductive isolation caused by natural selection
against immigrants from divergent habitats. *Evolution; international journal of organic*
*evolution* **59**, 705-719.
Pagliaccia D, Urak RZ, Wong F, *et al.* (2018) Genetic structure of the rice blast pathogen
(Magnaporthe oryzae) over a decade in North Central California rice fields. *Microbial ecology*
**75**, 310-317.
Pordel A, Ravel S, Charriat F, *et al.* (2020) Tracing the origin and evolutionary history of Pyricularia
oryzae infecting maize and barnyard grass. *Phytopathology®*, PHYTO-09.
Robert S, Zapater M-F, Carlier J, Abadie C, Ravigné V (2015) Multiple introductions and admixture at
the origin of the continental spread of the fungal banana pathogen Mycosphaerella fijiensis
in Central America: a statistical test using approximate Bayesian computation. *Revue*
*d'écologie*.
Ropars J, Lo YC, Dumas E, *et al.* (2016) Fertility depression among cheese-making Penicillium
roqueforti strains suggests degeneration during domestication. *Evolution; international*
*journal of organic evolution* **70**, 2099-2109.
Saleh D, Milazzo J, Adreit H, Fournier E, Tharreau D (2014) South-East Asia is the center of origin,
diversity and dispersion of the rice blast fungus, Magnaporthe oryzae. *New Phytologist* **201**,
1440-1456.
Saleh D, Milazzo J, Adreit H, Tharreau D, Fournier E (2012a) Asexual reproduction induces a rapid and
permanent loss of sexual reproduction capacity in the rice fungal pathogen Magnaporthe
oryzae: results of in vitro experimental evolution assays. *BMC evolutionary biology* **12**, 1-16.
Saleh D, Xu P, Shen Y, *et al.* (2012b) Sex at the origin: an Asian population of the rice blast fungus
Magnaporthe oryzae reproduces sexually. *Molecular ecology* **21**, 1330-1344.
Schulze-Lefert P, Panstruga R (2011) A molecular evolutionary concept connecting nonhost
resistance, pathogen host range, and pathogen speciation. *Trends in plant science* **16**, 117-
125.
Servedio MR, Van Doorn GS, Kopp M, Frame AM, Nosil P (2011) Magic traits in speciation: 'magic' but
not rare? *Trends in ecology & evolution* **26**, 389-397.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM (2015) BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*
**31**, 3210-3212.
Simpson JT, Wong K, Jackman SD, *et al.* (2009) ABySS: a parallel assembler for short read sequence
data. *Genome Research* **19**, 1117-1123.
Stanke M, Morgenstern B (2005) AUGUSTUS: a web server for gene prediction in eukaryotes that
allows user-defined constraints. *Nucleic Acids Research* **33**, W465-W467.
Sweigard JA, Carroll AM, Kang S, *et al.* (1995) Identification, Cloning, and Characterization of
*PWL2*, a Gene for Host Species Specificity in the Rice Blast Fungus. *Plant Cell* **7**, 1221-
1233.
Takabayashi N, Tosa Y, Oh HS, Mayama S (2002) A Gene-for-Gene Relationship Underlying the
Species-Specific Parasitism of Avena/Triticum Isolates of Magnaporthe grisea on Wheat
Cultivars. *Phytopathology* **92**, 1182-1188.
Taylor JW, Fisher MC (2003) Fungal multilocus sequence typing—it's not just for bacteria. *Current*
*opinion in microbiology* **6**, 351-356.
Taylor JW, Turner E, Townsend JP, Dettman JR, Jacobson D (2006) Eukaryotic microbes, species
recognition and the geographic limits of species: examples from the kingdom Fungi.
*Philosophical Transactions of the Royal Society B: Biological Sciences* **361**, 1947-1963.
Tharreau D, Fudal I, Andriantsimialona D, *et al.* (2009) World population structure and migration of
the rice blast fungus, Magnaporthe oryzae. In: *Advances in genetics, genomics and control of*
*rice blast disease*, pp. 209-215. Springer.
Walker A, Gladieux P, Decognet V, *et al.* (2015) Population structure and temporal maintenance of
the multihost fungal pathogen *Botrytis cinerea*: causes and implications for disease
management. *Environmental microbiology* **17**, 1261-1274.
Yang N, Ma G, Chen K, Wu X (2018) The population genetics of Alternaria tenuissima in four regions
of China as determined by microsatellite markers obtained by transcriptome sequencing.
*Frontiers in microbiology* **9**, 2904.
Yin Y, Mao X, Yang J, *et al.* (2012) dbCAN: a web resource for automated carbohydrate-active enzyme
annotation. *Nucleic Acids Research* **40**, W445-W451.
Yoshida K, Saunders DGO, Mitsuoka C, *et al.* (2016) Host specialization of the blast fungus
Magnaporthe oryzae is associated with dynamic gain and loss of genes linked to transposable
elements. *Bmc Genomics* **17**, 1.
Zhong Z, Chen M, Lin L, *et al.* (2018) Population genomic analysis of the rice blast fungus reveals
specific events associated with expansion of three main clades. *The ISME journal* **12**, 1867-
1878.

## DATA AVAILABILITY

All raw sequencing data are deposited under accession code PRJEB42377.

The following data sets were generated:

1. European Nucleotide Archive

M Thierry, F Charriat, J Milazzo, H Adreit, S Ravel, S Cros-Arteil, S Borron, V Sella, T Kroj, R Ioos, E Fournier, D Tharreau, P

Gladieux

ID PRJEB42377 Whole genome sequencing of Pyricularia oryzae fungi isolated from rice.

2. Zenodo

M Thierry, F Charriat, J Milazzo, H Adreit, S Ravel, S Cros-Arteil, S Borron, V Sella, T Kroj, R Ioos, E Fournier, D Tharreau, P

Gladieux

Single-nucleotide polymorphisms, genome assemblies, genome annotations, and gene predictions of Pyricularia oryzae

isolates from rice

doi 10.5281/zenodo.4561581

The following previously published data sets were used:

1. European Nucleotide Archive

A Pordel, S Ravel, F Charriat, P Gladieux, S Cros-Arteil, J Milazzo, H Adreit, M Javan-Nikkhah, A Mirzadi-Gohari, A Moumeni,
D Tharreau
ID PRJEB41186 Origin and evolutionary history of Pyricularia oryzae fungal pathogens infecting maize and barnyard grass in
Iran
779
780 2. European Nucleotide Archive

781 Z Zhong, M Chen, L Lin, Y Han, J Bao, W Tang, L Lin, Y Lin, R Somai, L Lu, W Zhang, J Chen, Y Hong, X Chen, B Wang, WC Shen,
782 G Lu, J Norvienyeku, DJ Ebbole, Z Wang
783 ID PRJNA354675 Population Genomic Analysis of the Rice Blast Fungus Reveals Specific Events Associated With Expansion
784 of Three Main Clades
785

**ADDITIONAL FILES**

**Source code 1.** R code used to prepare maps in Figure 6.

**Source code 2.** R code used for analyses based on the Outlying Mean Index, including Principal Component

Analysis presented in Figure 6.

**Supplementary file 1.** Isolates of Pyricularia oryzae

**Supplementary file 2.** Assignment of genotypes to clusters identified in this study, in Gladieux et al. 2018b (A)

and in Saleh et al. 2014 (B and C).

**Supplementary file 3.** $F_{ST}$ estimates between clusters within lineage 1.

**Supplementary file 4.** Analysis of mycelial growth rates.

**Supplementary file 5.** Mycelial growth rates.

**Supplementary file 6.** Analysis of sporulation rates.

**Supplementary file 7.** Sporulation rates.

**Supplementary file 8.** Matrix of compatibility between four lineages of *P. oryzae* isolates and five subspecies of

rice plants, as determined by pathogenicity tests in controlled conditions. Numbers in the matrix represent the

maximum symptom score across three replicate experiments. Numbers at the right and bottom margins

represent the number of compatible interactions observed (symptom score >2) for varieties and isolates,

respectively.

**Supplementary file 9.** Analysis of pathogenicity data.

**Supplementary file 10.** Sequenced isolates.

**Supplementary file 11.** Violin plots showing the number of putative effectors and non-effector proteins

detected in each lineage. Asterisks indicate significant differences in gene content (Tukey's HSD test; p<0.001).

**Supplementary file 12.** Estimating the size of the core and accessory genome using a rarefaction approach. For
each lineage, genomes were resampled in <=2000 combinations of N-1 genomes (N being the sample size). (A)
Non-effector genes. (B) Putative effectors.

**Supplementary file 13.** Principal component analysis based on the presence/absence of accessory putative
effectors and non-effector proteins. Panels A and B show principal component PC1 against PC2, and PC1
against PC6, respectively, for effectors. Panels C and D show principal component PC1 against PC2, and PC1
against PC6, respectively, for non-effectors. Between parentheses is the proportion of variance represented by
each principal component.

**Supplementary file 14.** Summary statistics of non-synonymous and synonymous polymorphism in four lineages
of P. oryzae. For each lineage, only orthogroups with sample size >=4 were included in calculations. Lineage:
lineage of origin of genomes used in calculations for each orthogroup. Orthogroup: orthogroup ID. N: sample
size. numNS: number of non-synonymous sites. numS: number of synonymous sites. SNS: number of non-
synonymous segregating sites. SS: number of synonymous segregating sites. PiNS: non-synonymous nucleotide
diversity. PiS: synonymous nucleotide diversity.

**Supplementary file 15.** ecological niches of the four major lineages (referred to as L1 to L4) considering each of
the 19 biomes individually (referred to as Bio1 to Bio19). The x-axis represents the outlying mean index (OMI),
which measures the distance between the mean habitat conditions used by a lineage and the mean habitat
conditions used by the entire species, to test the hypothesis that different lineages are distributed in regions
with different climates.

**Figure 1 – source data 1.** Multilocus genotypes for neighbor-net phylogenetic inference.

**Figure 1 – source data 2.** Ancestry proportions in K=4 sNMF clusters.

**Figure 2 – source data 1.** Multilocus genotypes for neighbor-net phylogenetic inference.

**Figure 2 – source data 2.** Ancestry proportions in K=4 sNMF clusters.

**Figure 2 – source data 3.** Geographic distribution of clones (i.e. multilocus genotypes repeated multiple times).

**Figure 3 – Source data 1.** Production of perithecia.

**Figure 3 – Source data 2.** Production of germinating ascospores

**Figure 4 – source data 1.** Symptom scores for *P. oryzae* isolates inoculated onto rice varieties

**Figure 5 – source data 1.** Size of core and accessory genomes for putative effectors and non-effector proteins

**Figure 5 – source data 2.** Presence and absence of putative effectors in *P. oryzae* genomes

**Figure 6 – source data 1.** GPS position of isolates

Figure 1. Rice-infecting *P. oryzae* populations are divided into four major lineages. Population subdivision was inferred from 264 distinct *P. oryzae* genotypes, representing 886 isolates, and the four lineages were represented in different colors. A: Neighbor-net phylogenetic network estimated with SPLITSTREE; reticulations indicate phylogenetic conflicts caused by homoplasy. B: Geographic distribution of the four lineages identified with SPLITSTREE and sNMF, with disk area proportional to sample size. C: Ancestry proportions in $K$=4 clusters, as estimated with sNMF software; each multilocus genotype is represented by a vertical bar divided into four segments indicating membership in $K$=4 clusters.

Figure 1 – figure supplement 1. Cross-entropy (CE) as a function of the number of clusters K modeled in sNMF analyses of

population subdivision.

Figure 2. Population subdivision in lineage 1. A: Neighbor-net phylogenetic network estimated with SPLITSTREE; reticulations indicate phylogenetic conflicts caused by homoplasy. B: Geographic distribution of the four clusters identified with sNMF, with disk area proportional to number of isolates. C: Ancestry proportions in four clusters, as estimated with sNMF; each multilocus genotype is represented by a vertical bar divided into four segments, indicating membership in $K$=4 clusters. D: Number of isolates and their geographic origin for each multilocus genotype of lineage 1. Panels C and D share an $x$-axis (each vertical bar represents a different multilocus genotype).

## (A) Production of perithecia

## (B) Production of germinating ascospores

Figure 3. Success of crosses between lineages 2-4 and clusters within lineage 1 with (A) proportion of crosses producing at least one perithecium, (B) scoring of ascus formation and ascospore germination for a subset of crosses.

Figure 4: Compatibility between 70 *P. oryzae* isolates and 44 rice varieties, representing five types of rice. A: Proportions of symptom scores as a function of the lineage of origin of isolates, or cluster of origin for isolates from lineage 1; B: Proportions of symptom scores as a function of the type of rice; C-G: Proportions of symptom scores as a function of the lineage of origin of isolates, for each type of rice. Abbreviations: Y, *Yule*; I, *International*; B, *Baoshan*; L, *Laos*; 2, lineage 2; 3, lineage 3; 4, lineage 4; Ind, indica; TeJ, temperate japonica; TrJ, tropical japonica; Aro, aromatic. Small capitals indicate significant differences. All interactions were assessed in three independent experiments, and the highest of the three symptom scores was used in calculations.

Figure 5. Pangenome analyses of 123 genomes of *P. oryzae,* representing four rice-infecting lineages. (A) Size of core and accessory genomes for putative effectors and non-effector proteins estimated using a rarefaction approach with a pseudo-sample size of n=30 genomes. Lineage 4 was not included due to small sample size. (B) Presence and absence in lineage 1-4 of the 72 putative effectors with the highest contribution to principal components 1, 2 and 6, in a principal component analysis of presence/absence data (Supplementary file 13).

Figure 6. Geographic distribution of four lineages of *P. oryzae* and corresponding climatic data. (A, B) Pie charts representing the distribution of the four lineages in the world (A) and in South, East and Southeast Asia (B), keeping only isolates for which the sampling position was precisely known (i.e., for which the region, city or GPS position was documented). Background map is from OpenStreetMap (CC-BY-SA) and represents the updated Köppen-Geiger climate classification of main climates at world scale as described in (Kottek *et al.* 2006). (C, D) Outlying Mean Index analysis, testing the separation of ecological niches among lineages, with (C) canonical weights of the 19 environmental variables included in the analysis, and (D) site coordinates (dots) and realized niches of lineages represented as ellipses. Variable bio13 co-localizes with bio16, and variables bio1 and bio3 co-localize with bio11. The 11 temperate variables included in the analysis were (bio1) Annual Mean Temperature, (bio2) Mean Diurnal Range, (bio3) Isothermality [100*(bio2/bio7)], (bio4) Temperature Seasonality [standard deviation*100], (bio5) Max Temperature of Warmest Month, (bio6) Min Temperature of Coldest Month, (bio7) Temperature Annual Range [bio5-bio6], (bio8) Mean Temperature of Wettest Quarter, (bio9) Mean Temperature of Driest Quarter, (bio10) Mean Temperature of Warmest Quarter, (bio11) Mean Temperature of Coldest Quarter. The 8 precipitation variables included in the analysis were (bio12) Annual Precipitation, (bio13) Precipitation of Wettest Month, (bio14) Precipitation of Driest Month, (bio15) Precipitation Seasonality (Coefficient of

Variation), (bio16) Precipitation of Wettest Quarter, (bio17) Precipitation of Driest Quarter, (bio18) Precipitation of Warmest Quarter, (bio19) Precipitation of Coldest Quarter.