ECOLOGICAL DIFFERENTIATION AND INCIPIENT SPECIATION IN THE FUNGAL PATHOGEN CAUSING

RICE BLAST

Maud THIERRY[1,2,3], Joëlle MILAZZO[1,2], Henri ADREIT[1,2], Sébastien RAVEL[1,2,4], Sonia BORRON[1], Violaine

SELLA[3], Renaud IOOS[3], Elisabeth FOURNIER[1], Didier THARREAU[1,2,5], Pierre GLADIEUX[1,5]

[1]UMR BGPI, Université de Montpellier, INRAE, CIRAD, Institut Agro, F-34398 Montpellier, France

[2] CIRAD, UMR BGPI, F-34398 Montpellier, France.

[3]ANSES Plant Health Laboratory, Mycology Unit, Domaine de Pixérécourt, Bâtiment E, F-54220 Malzéville, France

[4] South Green Bioinformatics Platform, Alliance Bioversity-international CIAT, CIRAD, INRAE, IRD, Montpellier,

France.

[5]pierre.gladieux@inrae.fr; didier.tharreau@cirad.fr

ABSTRACT

Natural variation in plant pathogens has an impact on food security and ecosystem health. The rice

blast fungus *Pyricularia oryzae,* which limits rice production in all rice-growing areas, is structured into

multiple lineages. Diversification and the maintenance of multiple rice blast lineages have been

proposed to be due to separation in different areas and differential adaptation to rice subspecies.

However, the precise world distribution of rice blast populations, and the factors controlling their

presence and maintenance in the same geographic areas, remain largely unknown. We used

genotyping data for 886 isolates from more than 185 locations in 51 countries to show that *P. oryzae* is

structured into one recombining and three clonal lineages, each with broad geographic distributions.

No evidence was found for admixture in clonal lineages, and crossing experiments revealed that female

sterility and early postmating genetic incompatibilities acted as strong barriers to gene flow between

these lineages. An analysis of climatic and geographic data indicated that the four lineages of *P. oryzae*

were found in areas differing in terms of the prevailing environmental conditions and types of rice

grown. Pathogenicity tests with representatives of the five main rice subspecies revealed differences

in host range between pathogenic lineages, highlighting a contribution of specialization to niche

separation between lineages, despite co-existence on the same host species. Our results demonstrate

that the spread of a pathogen across heterogeneous habitats and divergent populations of a crop

species can lead to niche separation and incipient speciation in the pathogen.

Short title: Incipient speciation in the rice blast pathogen

INTRODUCTION

Understanding and controlling natural variation in fungal plant pathogens is of great importance for

maintaining food security and ecosystem health. In addition to the overwhelming phylogenetic

diversity of pathogens (Burdon, 1993), many pathogens also harbor substantial diversity at the species

scale (Taylor and Fisher, 2003). Most fungal pathogens of plants form distinct populations (Taylor et al.,

2006), and this population structure is a compound outcome of migration, selection and drift, acting

at multiple scales, and patterned by factors such as time (Ali et al., 2014; Pagliaccia et al., 2018),

admixture (Robert et al., 2015), geography (Ali et al., 2014; Amend et al., 2009; Marin et al., 2009; Sork

and Werth, 2014; Yang et al., 2018), mode of reproduction (Ali et al., 2014; Büker et al., 2013; Gibson

et al., 2012; Ropars et al., 2016) and environmental conditions (Walker et al., 2015). The identity of the

host plants with which fungal pathogens interact is widely thought to be the the environmental factor

structuring the pathogen population with the strongest impact. Many plant pathogens reproduce

within or on their host, resulting in assortative mating on the basis of host use and a strong association

between adaptation to the host and reproductive isolation (Giraud et al., 2010; Servedio et al., 2011).

However, more generally, population structure can result from limited dispersal (i.e. limited gene flow

because of distance) or limited adaptation (i.e. limited gene flow because of differences in the capacity

to exploit resources), both of which may be due to a wealth of potential factors largely unexplored in

plant pathogens. Developing an understanding of how population structure emerges and is maintained

50 in fungal plant pathogens is a major goal in evolutionary microbiology, because it will improve our

51 comprehension of both disease emergence and the origin of fungal biodiversity.

52  *Pyricularia oryzae* (Ascomycota; syn. *Magnaporthe oryzae*) is a model widespread fungal plant

53 pathogen displaying population subdivision. Several lineages, each associated mostly with a single main

54 cereal/grass host, have been characterized within *P. oryzae* (Gladieux et al. 2018a). The lineage

55 infecting Asian rice (*Oryza sativa*) has been characterized in detail. Previous population genomic studies

56 of population structure in the rice-infecting lineage revealed genetic subdivision, with three (Saleh et

57 al., 2014; Tharreau et al., 2009; Zhong et al., 2018) to six (Gladieux et al. 2018b) lineages, with different

58 geographic patterns. The lineages were estimated to have diverged approximately 1000 years ago

59 (Gladieux et al. 2018b; Zhong et al. 2018), and pathogenicity tests suggested the local adaptation of

60 some lineages to the indica or japonica subspecies of rice (Gallet et al., 2016; Liao et al., 2016; Gladieux

61 et al., 2018b). *Pyricularia oryzae* is a heterothallic fungus with a sexual cycle involving mating between

62 individuals of opposite mating-types, with at least one of the partners capable of producing female

63 structures (i.e. "female-fertile"). A single lineage (lineage 1, prevailing in Southeast Asia) displayed a

64 genome-wide signal of recombination, a balanced ratio of mating-type alleles and a high frequency of

65 fertile females, consistent with sexual reproduction (Saleh et al. 2012a; Saleh et al. 2014; Gladieux et

66 al. 2018b). The other lineages had clonal population structures and highly biased frequencies of mating

67 types or fertile females, suggesting strictly asexual reproduction (Saleh et al. 2014; Gladieux et al.

68 2018b). Gene flow remains possible between lineages, because there are rare fertile strains and some

69 lineages have fixed opposite mating types. However, resequencing studies have revealed signatures of

70 recent shared ancestry between clonal lineages and the recombinant lineage from South-West Asia,

71 but not between the different clonal lineages, consistent with a lack of recent gene flow between

72 widespread lineages of opposite mating types (Gladieux et al. 2018b). The loss of female fertility has

73 probably been a key factor in the emergence of clonal lineages, inducing a shift towards strictly asexual

74 reproduction, and protecting populations adapting to new conditions from maladaptive gene flow from

75 ancestral, recombining populations (Giraud et al., 2010). This raises questions about the nature of the

factors underlying the emergence of different rice-infecting lineages in *P. oryzae*, and contributing to

their maintenance today, because, according to the competitive exclusion principle, different lineages

should not be able to co-exist on the same host (Hardin 1960; Giraud et al. 2017; Abbate et al. 2018).

Previous studies have helped to cement the status of *P. oryzae* as a model for studies of the population

biology of fungal pathogens, but most efforts to understand the population structure of the pathogen

have been unable to provide a large-scale overview of the distribution of rice-infecting lineages and of

the underlying phenotypic differences, because the availability of sets of genetic markers was limited

(Saleh et al., 2014; Tharreau et al., 2009), the number of isolates was relatively small (Gladieux, et al.

2018b; Zhong et al. 2018) or because few phenotypic traits were scored (Saleh et al., 2014; Tharreau

et al., 2009; Zhong et al., 2018).

We provide here a detailed genomic and phenotypic overview of natural variation in rice-

infecting *P. oryzae* pathogens sampled across all rice-growing areas of the world. With a view to

inferring and understanding the population structure of rice-infecting *P. oryzae*, we analyzed genetic

variation, using data with a high resolution in terms of genomic markers and geographic coverage, and

measured phenotypic variation for representative isolates. We addressed the following specific

questions: (i) How many different lineages of *P. oryzae* infect rice? (ii) Do lineages of *P. oryzae* display

footprints of recombination? (iii) Do lineages of *P. oryzae* present evidence of recent admixture? (iv)

Are the different lineages sexually competent and compatible? (v) Are the different lineages

differentially adapted to host and temperature? (vi) Can we find differences in the geographic and

climatic distribution of lineages?

RESULTS

The rice blast pathogen comprises one recombining, admixed lineage and three clonal, non-admixed

lineages

We characterized the global genetic structure of the rice blast pathogen, using an Illumina genotyping

beadchip to score 5,657 SNPs distributed throughout the genome of 886 *P. oryzae* isolates collected

from cultivated Asian rice in 51 countries. The final dataset included 3,686 SNPs after the removal of positions for which quality was low or data were missing. Clustering analyses based on sparse nonnegative matrix factorization algorithms, as implemented in the sNMF program (Frichot et al., 2014), revealed four clusters, hereafter referred to as "lineages" (Figure 1C). The model with K=4 clusters was identified as the best model on the basis of the cross-entropy criterion, as models with K>4 induced only a small decrease in cross-entropy, suggesting that K=4 captured the deepest subdivisions in the dataset (Supplementary figure 1). The neighbor phylogenetic network inferred with SPLITSTREE also supported the subdivision into four widely distributed lineages, with long branches separating three peripheral lineages branching off within a central lineage (Figure 1A; 1B). A comparison with previous findings revealed that the three peripheral lineages in the network corresponded to the previously described lineages 2, 3 and 4 (Gladieux et al. 2018b), whereas the central lineage corresponded to the combination of recombining lineage 1 and lineages 5 and 6, represented by only a few individuals in Gladieux et al. (2018b). The central lineage is referred to hereafter as lineage 1, for the sake of simplicity. Lineages 2 and 3 were similar to lineages B and C, respectively, identified on the basis of microsatellite data in a previous study (Saleh et al., 2014 - Supplementary figure 2). Genetic differentiation between the four lineages was strong and significant (Weir and Cockerham's F_{ST}> 0.54), indicating strong barriers to gene flow between the lineages. All genotypes from lineages 2 to 4 had high proportions of membership q in a single cluster (q > 0.89), whereas shared ancestry with the other three lineages was detected in lineage 1, with 32% of genotypes having q > 0.10 in lineages 2-4 (Figure 1C). Admixture may account for the lower F_{ST} observed in comparisons between lineage 1 and the other lineages. We detected no footprints of recombination in lineages 2, 3 and 4 with the pairwise homoplasy index (PHI) test and a null hypothesis of clonality (Table 1), confirming previous findings (Saleh et al. 2014; Gladieux et al. 2018b; Zhong et al. 2018). The rooted genealogies of the clonal lineages suggested that they were of Asian origin, and the patterns of shared multilocus genotypes were consistent with multiple introductions and intense movements of biological material over long distances (Supplementary text 1).

Figure 1. Population subdivision inferred from 264 *P. oryzae* genotypes, representing 886 isolates, with four identified lineages represented in different colors. A: Neighbor-net phylogenetic network estimated with SPLITSTREE; reticulations indicate phylogenetic conflicts caused by homoplasy. B: Geographic distribution of the four lineages identified with SPLITSTREE and sNMF, with disk area proportional to sample size. C: Ancestry proportions in K=4 clusters, as estimated with sNMF software; each multilocus genotype is represented by a vertical bar divided into four segments indicating membership in K=4 clusters.

Geographic structure and admixture within recombining lineage 1

Lineage 1 was the only lineage for which there was population genetic evidence of recombination in the PHI test (Table 1). Most of the lineage 1 isolates were collected in Asia (79%), but the lineage was present in all continents in which the pathogen was sampled (Europe: one isolate; North America: 10, Central and South America: 7, Africa: 22) (Figure 2). Clustering analysis on this single lineage with sNMF detected four clusters with different geographic distributions (Figure 2). Estimates of differentiation were lower between the clusters within lineage 1 (F_{ST} < 0.49) than between the main lineages (F_{ST} > 0.54), consistent with a longer history of restricted gene flow between the main lineages and a more recent history of admixture between clusters within lineage 1 (Supplementary table 1; Figure 1C).

Two clusters within lineage 1 (referred to hereafter as *Baoshan* and *Yule*) consisted mostly of isolates sampled from two different sites in Yunnan (Baoshan and Yule, respectively) located about 700 km apart. The third cluster consisted mostly of isolates from Laos and South China (referred to

hereafter as *Laos*), and the fourth brought together 95% of the isolates from lineage 1 collected outside

Asia (referred to hereafter as *International*). In Asia, the *International* cluster was found mostly in the

Yunnan province of China, India and Nepal (Figure 2). PHI tests for recombination rejected the null

hypothesis of clonality in all four clusters (Table 1). Admixture was widespread in lineage 1 (Figure 2),

with most of the isolates (78%) displaying membership proportions $q > 0.10$ for two or more clusters

(Figure 2). Only five genotypes were detected in multiple countries (genotype ID [number of countries]:

2 [6], 18 [2], 58 [2], 98 [3] and 254 [3], corresponding to 41 isolates in total). All these genotypes

belonged to the *International* cluster.

Figure 2. Population subdivision in lineage 1. A: Neighbor-net phylogenetic network estimated with SPLITSTREE; reticulations

indicate phylogenetic conflicts caused by homoplasy. B: Geographic distribution of the four clusters identified with sNMF,

with disk area proportional to sample size. C: Ancestry proportions in four clusters, as estimated with sNMF; each multilocus

genotype is represented by a vertical bar divided into four segments, indicating membership in K=4 clusters. D: Number of

160   isolates and their geographic origin for each multilocus genotype of lineage 1. Panels C and D share an *x*-axis (each vertical

161   bar represents a different multilocus genotype).

162   Table 1. PHI test for recombination (null hypothesis: clonality) and distribution of mating types and fertile

163   females in lineages, and in clusters within lineage 1.

Lineage/*Cluster*	*p*-value PHI test	Mat1.1/Mat1.2 (%)	Female fertility (%)
1	<0.0001	53/47	43
Baoshan	<0.0001	69/31	67
International	<0.0001	55/45	4
Laos	<0.0001	68/32	37
Yule	<0.0001	40/60	79
2	0.25	97/3	0
3	0.18	3/97	7
4	0.22	97/3	0

Reproductive barriers caused by female sterility and postmating genetic incompatibilities

Our analysis of genotyping data revealed the existence of a large, ancient, recombinant population

from which several clonal lineages have more recently emerged. This raised the question as to why

these populations became clonal, with no further gene flow. We addressed this question by

determining the capacity of the different lineages to engage in sexual reproduction, by characterizing

the distribution of mating types and the ability to produce female structures (i.e. female fertility). Using

in vitro crosses with tester strains, we found that lineages 2, 3 and 4 were composed almost exclusively

of a single mating type (97% of lineage 2 and 4 isolates tested carried the *Mat1.1* allele; 97% of lineage

3 isolates carried the *Mat1.2* allele), and had only small proportions (0-7%) of isolates with female

fertility (Table 1). By contrast to these clonal lineages, the mating type ratio was more balanced in

lineage 1 (52% of *Mat1.1*; Table 1), with *Mat1.1*/*Mat1.2* ratios ranging from 40/60 in the *Yule* cluster

to 69/31 in the *Baoshan* cluster. Female fertility rates were higher in three of the clusters in lineage 1

(*Yule*: 79%; *Baoshan*: 67%; *Laos*: 37%; Table 1), but that in the International cluster (4% female fertility) was as low as those in the clonal clusters (Table 1). These observations indicate that there is a potential for sexual reproduction between lineages with compatible mating-types and, albeit rare, fertile females, but not between lineages 2 and 4, due to the low female fertility rates and highly biased mating-type ratios.

(A) Production of perithecia

(B) Production of germinating ascospores

Figure 3. Success of crosses between lineages 2-4 and clusters within lineage 1 with (A) proportion of crosses producing at least one perithecium, (B) scoring of ascus formation and ascospore germination for a subset of crosses.

We further assessed the likelihood of sexual reproduction within and between lineages, by

evaluating the formation of sexual structures (perithecia), the production of asci (i.e. meiotic octads

within the perithecia) and the germination of ascospores (i.e. meiospores in meiotic octads) in crosses

between randomly selected isolates of opposite mating types from all four lineages (Figure 3A). This

experiment revealed a marked heterogeneity in the rate of perithecium formation across lineages.

Three clusters in lineage 1 had the highest rates of perithecia formation, with isolates in the *Yule* cluster,

in particular, forming perithecia in 93% of crosses with isolates from the same cluster, and in more than

46% of crosses with isolates from other lineages (Figure 3A). Isolates from lineages 2, 3 and 4 could

only be crossed with isolates from other lineages, given their highly biased mating-type ratios, and the

percentage of these crosses producing perithecia was highly variable (ranging from 0% to 83%,

depending on the lineages involved; Figure 3A). The rate of perithecium formation was similar in

crosses involving the *International* cluster of lineage 1 and in crosses involving clonal lineages 2-4, and

none of the crosses attempted between isolates from the *International* cluster led to perithecium

formation (Figure 3A). Perithecium dissection for a subset of crosses involving some of the most fertile

isolates revealed that most inter-lineage crosses produced perithecia that did not contain asci or

contained asci with low ascospore germination rates (Figure 3B). Whereas 100% of crosses between

isolates of the *Yule* cluster produced numerous germinating ascospores, ascospore germination rates

were only 33%, 56% and 7% in *Yule* x lineage 2, *Yule* x lineage 3 and *Yule* x lineage 4 crosses, respectively.

Together, these results indicate that the clonal structure of lineages 2-4 is caused by highly biased

mating types and female sterility, and that the three clonal lineages and the *International* cluster of

lineage 1 are isolated from each other by strong pre- and early-postmating barriers, including breeding

system isolation (differences in mating type and female sterility), and a combination of gametic

incompatibility, zygotic mortality or hybrid non-viability. However, three of the clusters in lineage 1 had

biological features consistent with an ability to reproduce sexually, suggesting possible sexual

compatibility with clonal lineages 2-4 and the *International* cluster.

Specialization to temperature conditions and rice subspecies

Specialization can further reduce gene flow, and may facilitate niche separation between lineages,

which should favor the emergence of new lineages and their maintenance on the same host species in

conditions of competitive exclusion. The strong reproductive barriers demonstrated in our analysis of

crossing experiments would favor the establishment of such specialization, because reproductive

isolation facilitates adaptation to new conditions, by preventing maladaptive gene flow from ancestral,

non-adapted populations (Giraud et al., 2010). Given the very broad distribution of *P. oryzae* and the

strong environmental heterogeneity with which it is confronted in terms of the nature of its hosts and

the climate in which it thrives, we evaluated the variation of pathogen fitness over different types of

rice and temperature conditions. Our experiments measuring growth and sporulation rates at different

temperatures revealed limited differences in performance between lineages, and no pattern

consistent with strict specialization according to temperature (Supplementary text 2). We therefore

report only the results of pathogenicity tests below.

We assessed the importance of adaptation to the host, by challenging 45 rice varieties

representative of the five main genetic groups of Asian rice (i.e. the temperate japonica, tropical

japonica, aus, aromatic and indica subgroups of *Oryza sativa*) with 70 isolates representative of the

four lineages of *P. oryzae* and the four clusters within lineage 1 (Supplementary Figure 3). Qualitative

symptoms were noted and analyzed with a proportional-odds model, revealing significant differences

between lineages (*p*-value = 2.2×10^{-16}), and between rice genetic groups (*p*-value=2.2×10^{-16}), and a

significant interaction between these two variables (*p*-value=2.0×10^{-10}). In comparisons between

lineages, lineage 2 differed from the other lineages in generally producing smaller lesions (Figure 4A).

In comparisons of rice genetic groups, significantly strong symptoms were observed on temperate

japonica rice than on the other types of rice, whereas the varieties of the aromatic genetic group were

significantly more resistant to rice blast (Figure 4B). The finding of a significant interaction between

lineage and rice genetic group indicates that the effect of the lineage on the proportion of compatible

interactions differed between rice types, suggesting adaptation to the host. In particular, an analysis

of interactions also revealed that the isolates of the *Yule* cluster were highly pathogenic on most indica

varieties (Supplementary Figure 3), with all isolates from this cluster pathogenic to all indica varieties except IR8. Together, these experiments revealed significant differences in host range between lineages, but with overlaps between these host ranges, indicating that specialization to the host was not strict (Figure 4C-G). Adaptation to the host or small differences in temperature optima could, nonetheless, further increase niche separation and reduce gene flow between lineages, via pre-mating barriers such as immigrant non-viability (i.e., lower rates of contact between potential mates due to the mortality of immigrants; Gladieux et al., 2011; Nosil et al., 2005), and post-mating barriers, such as ecological hybrid non-viability (i.e., lower survival of ill-adapted hybrid offspring).

Figure 4: Compatibility between 70 *P. oryzae* isolates and 45 rice varieties, representing five subgroups of rice. A: Proportions of symptom scores as a function of the lineage of origin of isolates, or cluster of origin for isolates from lineage 1; B: Proportions of symptom scores as a function of the type of rice; C-G: Proportions of symptom scores as a function of the lineage of origin of isolates, for each type of rice. Abbreviations: Y, *Yule*; I, *International*; B, *Baoshan*; L, *Laos*; 2, lineage 2; 3,

250 lineage 3; 4, lineage 4; Ind, indica; TeJ, temperate japonica; TrJ, tropical japonica; Aro, aromatic. Small capitals indicate

251 significant differences.

Geographic and climatic differentiation in the distribution of *P. oryzae* lineages

Our tests of adaptation to host and temperature under controlled conditions showed that specialization was not strict, but it remained possible that fitness differences between lineages would be sufficient under natural conditions to induce separation in different ecological niches, and/or that our experimental conditions did not capture the full extent of the phenotypic differences. Here, we tested the hypothesis that lineages thrive in geographically and/or environmentally different conditions, which would provide indirect evidence for specialization to different habitats. We tested this hypothesis by collecting geographic and climatic data for all isolates, approximating to regions or the nearest cities when the exact GPS position of the sampling area was not available (Supplementary table 2). At a larger scale, clonal lineages 2 and 3 were both widespread, with lineage 2 found on all continents, and lineage 3 on all continents except Europe. Lineage 2 was the only lineage sampled in Europe (with the exception of a single isolate from lineage 1), whereas lineage 3 was more widespread in intertropical regions. Lineage 4 was mostly found in South Asia (India, Bangladesh, Nepal), and in the USA and Africa (Benin, Tanzania). The recombining lineage, lineage 1, was found in all continents, but was mostly found in Asia (79% of all isolates and 94% of all genotypes). At a smaller scale, two, or even three different lineages were sampled in the same year in the same city in 11 different countries from all continents. However, in only one instance did two isolates from different lineages have identical GPS positions (isolates US0106 and US0107 sampled from the same field). Thus, lineages had different, but partially overlapping distributions, including some overlap with the distribution of the sexually reproducing lineage (lineage 1).

Figure 5. Geographic distribution of four lineages of *P. oryzae* and corresponding climatic data. (A, B) Pie charts representing

the distribution of the four lineages in the world (A) and in South, East and Southeast Asia (B), keeping only isolates for which

the sampling position was precisely known (i.e., for which the region, city or GPS position was documented). Background map

is from OpenStreetMap (CC-BY-SA) and represents the updated Köppen-Geiger climate classification of main climates at

world scale as described in Kottek et al. (2006). (C, D) Outlying Mean Index analysis, testing the separation of ecological niches

among lineages, with (C) canonical weights of the19 environmental variables included in the analysis, and (D) site coordinates

(dots) and realized niches of lineages represented as ellipses. Variable bio13 co-localise with bio16, and variables bio1 and

bio3 co-localize with bio11. The 11 temperate variables included in the analysis were (bio1) Annual Mean Temperature, (bio2)

Mean Diurnal Range, (bio3) Isothermality [100*(bio2/bio7)], (bio4) Temperature Seasonality [standard deviation*100], (bio5)

Max Temperature of Warmest Month, (bio6) Min Temperature of Coldest Month, (bio7) Temperature Annual Range [bio5-

bio6], (bio8) Mean Temperature of Wettest Quarter, (bio9) Mean Temperature of Driest Quarter, (bio10) Mean Temperature

of Warmest Quarter, (bio11) Mean Temperature of Coldest Quarter. The 8 precipitation variables included in the analysis

were (bio12) Annual Precipitation, (bio13) Precipitation of Wettest Month, (bio14) Precipitation of Driest Month, (bio15)

Precipitation Seasonality (Coefficient of Variation), (bio16) Precipitation of Wettest Quarter, (bio17) Precipitation of Driest

Quarter, (bio18) Precipitation of Warmest Quarter, (bio19) Precipitation of Coldest Quarter.

We then investigated whether differences in the geographic range of lineages were associated

with differences in climatic variables. By plotting sampling locations onto a map of the major climate

regions (Kottek et al. 2006) we were able to identify clear differences in the climatic distributions of

lineages 2 and 3, with lineage 2 mostly found in warm temperate climates and lineage 3 found in

equatorial climates (Figure 5). We used the outlying mean index (OMI), which measures the distance

between the mean habitat conditions used by a lineage and the mean habitat conditions used by the

entire species, to test the hypothesis that different lineages are distributed in regions with different

climates. Using information for 19 climatic variables (biomes) from the WorldClim bioclimatic variables

database (Fick and Hijmans 2017) for all sampling locations, we obtained marginally significant results,

in a statistical permutation test on OMI values, for lineages 2, 3 and 4 (p-values: L1, p=0.6213; L2,

p=0.0003; L3, p=0.0001; L4, p=0.0176). The OMI analysis, in which the first two axes accounted for 69%

and 25% of the variability, respectively, revealed that lineage 2 was more frequent in regions with a

high annual temperature range (biome 7) or a high degree of seasonality (biome 4); lineage 4 was

associated with regions with high levels of seasonal precipitation (biomes 13, 16 and 18), and lineage

3 was more frequent in regions with high temperatures (biomes 1, 6, 10 and 11) and high levels of

isothermality (biome 3), characteristic of tropical climes (Figure 5; Supplementary figure 3).

DISCUSSION

We describe here the population structure of the rice blast fungus, and the eco-evolutionary factors

underlying the emergence and maintenance of multiple divergent lineages in this widespread

pathogen. Our analysis of SNP genotyping data showed that *P. oryzae* could be subdivided into four

lineages. Two of the lineages previously detected in an analysis of whole-genome data (lineages 5 and

6; Gladieux et al., 2018b), and represented by only a few individuals in our dataset, were assigned to

lineage 1 in our analysis, and may correspond to a subdivision of lineage 1. Differences in the number

of lineages detected may reflect an ascertainment bias, given that our SNP-genotyping beadchip was designed with a set of genomic sequences that did not include representatives of lineages 5 and 6. Alternatively, the sampling used in the previous study (Gladieux et al., 2018b) may not have been dense enough to reveal the lack of differentiation between lineages 1, 5 and 6, making them appear to be different entities. Consistent with previous findings, our analysis of the genealogical relationships between isolates revealed three clonal groups connected by long branches to a central recombining group. The finding of an internationally distributed cluster in lineage 1 sheds light on the early evolutionary changes contributing to the emergence of clonal lineages in this pathogen. The *International* cluster of lineage 1 displayed genetic footprints of recombination and sexual reproduction (in the form of an excess of homoplasious variants or relatively balanced mating types), but the signal of recombination detected here may be purely historical, because we also found that the clonal fraction and the frequency of sterile females were very high in this cluster. The loss of female fertility may, therefore, be an early and major cause of the shift towards asexual reproduction. Some genotypes assigned to lineage 1 may, in the future, come to resemble other clonal lineages in analyses of genealogical relationships, and form a long branch stemming from the central lineage 1, particularly for genotypes already displaying clonal propagation, such as genotypes 2, 18, 58, 98 and 254. However, according to the principle of competitive exclusion, competition between clonal lineages should only allow hegemonic expansion of the most competitive clonal groups, unless they separate into different ecological niches. It may be this process that ultimately leads to the fixation of a single mating type, as in clonal lineages 2-4.

We found that female sterility and intrinsic genetic incompatibilities represented strong barriers to gene flow between the clonal lineages, potentially accounting for their maintenance over time, without fusion, despite the compatibility of mating types (at least between lineages 2 and 4 [*Mat1.1*] on the one hand, and lineage 3 [*Mat1.2*] on the other). These barriers to gene flow may have contributed to the establishment of lineage specialization (because reproductive isolation facilitates adaptation by limiting the flow of detrimental ancestral alleles in populations adapting to a new

habitat), accounting for the maintenance of the clonal lineages on the same host species, when faced

with competitive exclusion. Our analyses also show that, despite widely overlapping large-scale

distributions, the different lineages are essentially found in different regions, with different climatic

characteristics, when their distributions are observed at a finer scale. Lineage 1 was mostly sampled in

Southeast Asia, lineage 4 predominated in India, and lineages 2 and 3 had global distributions, but the

geographic ranges of all these lineages overlapped. However, an analysis of climatic data indicated that

lineage 2 predominated in temperate climates, where temperate japonica rice is grown, lineage 3

predominated in tropical climates, in which indica rice is grown, and lineage 4 predominated in regions

with high levels of seasonal precipitation in which the indica and aromatic rice types are the principal

rice types grown. Despite the finding of separation in different climatic regions, our experiments

revealed no strong differences between lineages in terms of sporulation and mycelial growth on

synthetic media at different temperatures. This suggests that, if adaptation to temperature occurs in

this pathogen, it was not measured by our experiments, either because it involves traits other than

sporulation and hyphal growth, or because the *in vitro* conditions were not suitable to demonstrate

differences. The host range varied across lineages, but all host ranges overlapped, indicating that host

specialization was not strict, or was not fully captured by our experiments. Adaptation to the host may,

nevertheless, decrease gene flow between populations even further. For instance, lineage 2 had a

narrow host range, potentially limiting the opportunities for encounters and mating with other groups.

Our results also indicate that lineage 1 may pose a major threat to rice production. Eleven of

the 12 most multivirulent isolates (lesion type > 2 on more than 40 varieties tested) belonged to lineage

1, and the *Yule* cluster, in particular, was highly pathogenic on indica varieties. The propagation of such

highly pathogenic genotypes, belonging to a lineage that has both mating types and fertile females,

should be monitored closely. The monitoring of lineage 1 is all the more critical because this lineage

has an intermediate geoclimatic distribution that overlaps largely with the distributions of the other

three lineages, and some of the attempted crosses between clonal lineage 2-4 isolates and recombining

lineage 1 isolates produced viable progeny. This finding confirms the possibility of gene flow into this

lineage, as previously demonstrated on the basis of admixture mapping (Gladieux et al., 2018b), but it

also raises the question of the threat posed by gene flow from lineage 1, which is highly pathogenic, to

the other lineages.

In conclusion, our study of genetic and phenotypic variation within and between clonal

lineages, and within the recombinant lineage of *P. oryzae* suggests a scenario for the emergence of

widespread clonal lineages. The loss of female fertility may be a potent driver of the emergence of

asexually reproducing groups of clones. The reproductive isolation generated by the loss of sex and the

accumulation of mutations due to the absence of sexual purging would facilitate the specialization of

some of these clonal groups, leading to the competitive exclusion of the least efficient clonal groups,

and, finally, to the propagation of clonal lineages fixed for a single mating type. Our results, thus,

demonstrate that the spread of a pathogen across heterogeneous habitats and divergent populations

of a crop species can lead to niche separation and incipient speciation in the pathogen.

MATERIALS AND METHODS

Biological material

We chose 886 *P. oryzae* isolates collected on Asian rice between 1954 and 2014 as representative of

the global genetic diversity of the fungus. Isolates were selected on the basis of microsatellite data, to

maximize the number of multilocus genotypes represented (Saleh et al. 2014 and unpublished data),

or based on geographic origin in the absence of genotypic data, to maximize the number of countries

represented in the dataset.

Sixty-eight isolates were selected for experimental measurements of reproductive success,

adaptation to host, and growth and sporulation experiments at different temperatures (Supplementary

table 2). This subset of isolates included 10 isolates from each of the three clonal lineages (lineages 2-

4), 27 isolates from the various clusters within lineage 1 [*Baoshan* (9 isolates), *International* (10), *Laos*

(8), and *Yule* (10)] and isolate CH0999, a reference female-fertile strain of the Mat1.1 mating-type

(Saleh et al., 2012a).

Forty-six varieties were chosen as representative of the five main genetic subgroups of Asian

rice (Garris et al., 2005): indica (Chau, Chiem chanh, DA11, De abril, IR8, JC92, JC120, Pappaku), aus

(Arc 10177, Baran boro, Black gora, DA8, Dholi boro, Dular, FR13 A, JC148, Jhona 26, Jhona 149,

Kalamkati, T1, Tchampa, Tepi boro), temperate japonica (Aichi asahi, Kaw luyoeng, Leung pratew,

Maratelli, Nep hoa vang, Nipponbare, Sariceltik, Som Cau 70A), tropical japonica (Azucena, Binulawan,

Canella de ferro, Dholi boro, Gogo lempuk, Gotak gatik, Moroberekan, Trembese) and aromatic (Arc

10497, Basmati lamo, Dom zard, Firooz, JC1, Kaukkyisaw, N12). The Maratelli (temperate japonica) and

CO39 (indica) varieties were used as susceptible controls.

Genotyping

Pyricularia oryzae isolates were genotyped at 5,657 genomic positions with an Illumina Infinium

beadchip microarray carrying single-nucleotide polymorphisms identified in 25 previously sequenced

P. oryzae genomes (Gladieux et al., 2018b). According to our clustering analysis, the 25 isolates

sequenced belonged to lineage 1 (*n*=2 isolates), lineage 2 (*n*=9), lineage 3 (*n*=8) or lineage 4 (*n*=3), and

three isolates were not characterized here. The final dataset included 3,686 biallelic SNPs without

missing data.

Population subdivision and recombination

Among the 886 *P. oryzae* isolates genotyped, we identified 264 different multilocus genotypes, which

were used for the analysis of population subdivision. We used the sNMF program to infer individual

ancestry coefficients in *K* ancestral populations. This program is optimized for the analysis of large

datasets and does not assume Hardy-Weinberg equilibrium. It is, therefore, more appropriate to deal

with inbred or clonal lineages (Frichot et al., 2014). We used SPLITSTREE version 4 to visualize relationships between genotypes in a phylogenetic network, with reticulations to represent the conflicting phylogenetic signals caused by homoplasy. We also used the pairwise homoplasy index (PHI) test implemented in SPLITSTREE to test the null hypothesis of clonality. We then inferred a maximum likelihood genealogy, to investigate further the relationships between genotypes within the clusters for which clonality could not be rejected, and we used the sNMF software to investigate population subdivision further within the cluster for which the null hypothesis of clonality could be rejected. Weir and Cockerham's F_{ST} was calculated with the R package HIERFSTAT, by the WC84 method (Goudet, 2005).

Experimental measurement of reproductive success and female fertility

Pyricularia oryzae is a heterothallic fungus with two mating types (*Mat1.1* and *Mat1.2*). Sexual reproduction between strains of opposite mating type can be observed in laboratory conditions, and results in the production of ascospores within female sexual structures called perithecia (Saleh et al. 2012). On synthetic media, perithecia are formed at the contact zone between parental mycelia. Crosses were carried out on a rice flour agar medium (20 g rice flour, 2.5 g yeast extract, 15 g agar and 1 L water, supplemented with 500 000 IU penicillin G after autoclaving for 20 min at 120°C), as described by Saleh et al. (2012b). We assessed reproductive success by determining the production of perithecia produced after three weeks of culture at 20°C under continuous light. Perithecia can be formed by the two interacting partners or by one partner only. Isolates forming perithecia are described as female-fertile. We measured female fertility for 220 isolates (listed in Supplementary table 2), by monitoring perithecium production in crosses involving the tester strains CH0997 (Mat1.2) and CH0999 (Mat1.1). We further assessed the presence of asci and germinating ascospores in perithecia for a subset of crosses, by excising perithecia with a scalpel and counting the number of germinated filaments for each individual ascus after overnight incubation on water agar. The subset of crosses included 10 Mat1.1 isolates (lineage 1: CH0999, CH1065, CH1076; lineage 2: CH0092, MC0016, SP0006;

lineage 4: IN0017, IN0092, NP0070, CH0718) and 6 Mat1.2 isolates (lineage 1: CH0997; CH1083,

CH1120; lineage 3: BR0019, CH0549, MD0929).

Pathogenicity tests

Compatibility between *P. oryzae* isolates and rice plants from the five main genetic subgroups of rice

(indica, temperate japonica, tropical japonica, aus and aromatic) was measured in controlled

conditions. Seventy isolates were inoculated on 46 varieties. Inoculations were performed as described

by Gallet et al. (2016). Conidial suspensions (25 000 conidia.mL^{-1}) in 0.5% gelatin were sprayed onto

three-week-old rice seedlings (> 6 plants/variety). The inoculated plants were incubated for 8 hours at

27°C and 100% humidity and then for seven days with a day/night alternation (13 hours at 27°C/11

hours at 21°C), before scoring symptoms. Inoculation and symptom scoring were performed three

times, and the maximum of the three scores obtained was used in calculations. Lesion type was rated

from 1 to 6 (Gallet et al. 2016) and the disease area was assessed visually on leaves.

Mycelial growth and sporulation rate at different temperatures

Mycelial growth and sporulation rate were measured for 42 isolates at five different temperatures

(10°C, 15°C, 20°C, 25°C and 30°C). For each isolate, Petri dishes containing PDA medium were

inoculated with mycelial plugs placed at the center, and incubated in a growth chamber with a fixed

temperature. Mycelium diameter was measured along two perpendicular axes at different time points.

At the end of the experiment, conidia were collected by adding 5 mL of water supplemented with 0.01%

Tween 20 to the Petri dish and rubbing the surface of the mycelium. Conidia were counted in a

hemocytometer. Three biological replicates were performed for each isolate, at each temperature.

Statistical analyses of climatic and phenotypic data

<u>Ecological niche separation:</u> OMI (outlying mean index), or marginality, is used to study niche

separation and niche breadth. OMI gives the same weight to all samplings, whether rich or poor in

species and individuals, and is particularly suitable in cases in which sampling is not homogeneous.

OMI measures the deviation between the mean environmental conditions used by one lineage and the

mean environmental conditions used by all lineages. OMI analysis then places the lineages in

environmental conditions maximizing their OMI. Environmental values, consisting of 19 biome values

(WorldClim bioclimatic variables – Fick and Hijmans 2017), were retrieved for all sampling locations.

Climatic values were normalized before the analysis. A contingency table was generated to associate

the number of isolates from each lineage with each sampling location. Only isolates with a precise

location (known region, city or GPS position of sampling) were included. A random permutation test,

with 10000 permutations, was used to assess the statistical significance of marginality for each lineage.

Pathotyping: Ordinal symptoms scores were analyzed with a proportional-odds model

accounting for ordered categorical responses with a clm() function implemented in the ordinal R

package (version 2018.8-25). The significance of the factors was analyzed by ANOVA. Pairwise

comparisons of significant factors were performed after computing least-squares means with LSMEANS

version 2.30-0 in R, with Tukey adjustment.

Mycelium growth: Mycelium growth was analyzed separately for each temperature, with a

linear mixed-effects model. The significance of the factors was analyzed by ANOVA. Pairwise

comparisons of significant factors were performed after computing least-squares means with LSMEANS

version 2.30-0 in R.

Sporulation: The median number of spores calculated for the various repetitions of the

experiment was analyzed with a negative binomial generalized linear model. Pairwise comparisons of

significant factors were performed by computing least-squares means.

ACKNOWLEDGMENT

We thank Tatiana Giraud for useful suggestions. SNP data were obtained in the framework of a Bayer

Crop Science-CIRAD project. We thank all our colleagues who shared strains or participated in the

collection of rice blast samples.

Bibliography

Abbate, J. L., Gladieux, P., Hood, M. E., de Vienne, D. M., Antonovics, J., Snirc, A., & Giraud, T. (2018).
Co-occurrence among three divergent plant-castrating fungi in the same *Silene* host species. *Molecular*
Ecology, *27*(16), 3357-3370.

Adreit, H., Andriantsimialona, D., Utami, D.W., Notteghem, J.L., Lebrun, M.H., and Tharreau, D. (2007).
Microsatellite markers for population studies of the rice blast fungus, *Magnaporthe grisea*. Mol. Ecol.
Notes *7*, 667–670.

Ali, S., Gladieux, P., Rahman, H., Saqib, M.S., Fiaz, M., Ahmad, H., Leconte, M., Gautier, A., Justesen,
A.F., Hovmøller, M.S., et al. (2014). Inferring the contribution of sexual reproduction, migration and
off-season survival to the temporal maintenance of microbial populations: a case study on the wheat
fungal pathogen *Puccinia striiformis* f.sp. *tritici*. Mol. Ecol. *23*, 603–617.

Amend, A.S., Garbelotto, M., Fang, Z., and Keeley, S.C. (2009). Isolation by landscape in populations of
a prized edible mushroom *Tricholoma matsutake*. Conserv. Genet. *11*, 795–802.

Büker, B., Petit, E., Begerow, D., and Hood, M.E. (2013). Experimental hybridization and backcrossing
reveal forces of reproductive isolation in *Microbotryum*. BMC Evol. Biol. *13*, 224.

Burdon, J.J. (1993). The structure of pathogen populations in natural plant communities. Annu. Rev.
Phytopathol. *31*, 305–323.

Fick, S.E., and Hijmans, R.J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global
land areas. Int. J. Climatol. *37*, 4302–4315.

Frichot, E., Mathieu, F., Trouillon, T., Bouchard, G., and François, O. (2014). Fast and efficient
estimation of individual ancestry coefficients. Genetics *196*, 973–983.

Gallet, R., Fontaine, C., Bonnot, F., Milazzo, J., Tertois, C., Adreit, H., Ravigné, V., Fournier, E., and

Tharreau, D. (2016). Evolution of compatibility range in the rice− *Magnaporthe oryzae* system: an

uneven distribution of R genes between rice subspecies. Phytopathology *106*, 348–354.

Garris, A.J., Tai, T.H., Coburn, J., Kresovich, S., and McCouch, S. (2005). Genetic structure and diversity

in *Oryza sativa* L. Genetics *169*, 1631–1638.

Gibson, A.K., Hood, M.E., and Giraud, T. (2012). Sibling competition arena: selfing and a competition

arena can combine to constitute a barrier to gene flow in sympatry. Evolution *66*, 1917–1930.

Giraud, T., and Lambert, N. (2017). Reproducible Cartography. In Advances in Cartography and

GIScience, M.P. Peterson, ed. (Springer International Publishing), pp. 173–183.

Giraud, T., Gladieux, P., and Gavrilets, S. (2010). Linking the emergence of fungal plant diseases with

ecological speciation. Trends Ecol. Evol. *25*, 387–395.

Giraud, T., Koskella, B., & Laine, A. L. (2017). Introduction: microbial local adaptation: insights from

natural populations, genomics and experimental evolution. *Molecular Ecology*, *26*(7), 1703-1710.

Gladieux, P., Guérin, F., Giraud, T., Caffier, V., Lemaire, C., Parisi, L., ... & Le Cam, B. (2011). Emergence

of novel fungal pathogens by ecological speciation: importance of the reduced viability of

immigrants. *Molecular Ecology*, *20*(21), 4521-4532.

Gladieux, P., Condon, B., Ravel, S., Soanes, D., Maciel, J.L.N., Nhani, A., Chen, L., Terauchi, R., Lebrun,

M.-H., Tharreau, D., et al. (2018a). Gene flow between divergent cereal- and grass-specific lineages of

the rice blast fungus *Magnaporthe oryzae*. MBio *9*, e01219-17.

Gladieux, P., Ravel, S., Rieux, A., Cros-Arteil, S., Adreit, H., Milazzo, J., Thierry, M., Fournier, E., Terauchi,

R., and Tharreau, D. (2018b). Coexistence of multiple endemic and pandemic lineages of the rice blast

pathogen. Mbio *9*, e01806-17.

Goudet, J. (2005). hierfstat, a package for R to compute and test hierarchical F-statistics. Mol. Ecol.

Notes *5*, 184–186.

Hardin, G. (1960). The competitive exclusion principle. *Science*, *131*(3409), 1292-1297.

Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F. (2006). World map of the Köppen-Geiger

climate classification updated. Meteorol. Z. *15*, 259–263.

Liao JJ, Huang HC, Meusnier I, Adreit H, Ducasse A, Bonnot F, Pan L, He XH, Kroj T, Fournier E,

Tharreau D, Gladieux P, Morel JB. 2016. Pathogen effectors and plant immunity determine

specialization of the blast fungus to rice subspecies. e-Life 5:e19377. DOI:10.7554/eLife.19377

Marin, M., Preisig, O., Wingfield, B.D., Kirisits, T., and Wingfield, M.J. (2009). Single sequence repeat

markers reflect diversity and geographic barriers in Eurasian populations of the conifer pathogen

Ceratocystis polonica. For. Pathol. *39*, 249–265.

Nosil, P., Vines, T. H., & Funk, D. J. (2005). Reproductive isolation caused by natural selection against

immigrants from divergent habitats. *Evolution*, *59*(4), 705-719.

Pagliaccia, D., Urak, R.Z., Wong, F., Douhan, L.I., Greer, C.A., Vidalakis, G., and Douhan, G.W. (2018).

Genetic structure of the rice blast pathogen (*Magnaporthe oryzae*) over a decade in North Central

California rice fields. Microb. Ecol. *75*, 310–317.

Robert, S., Zapater, M., Carlier, J., Abadie, C., and Ravigné, V. (2015). Multiple introductions and

admixture at the origin of the continental spread of the fungal banana pathogen *Mycosphaerella*

fijiensis in Central America: a statistical test using approximate Bayesian computation. Revue

d'Ecologie (Terre et Vie). *70*, 127-138

Ropars, J., Lo, Y., Dumas, E., Snirc, A., Begerow, D., Rollnik, T., Lacoste, S., Dupont, J., Giraud, T., and

López-Villavicencio, M. (2016). Fertility depression among cheese-making *Penicillium roqueforti* strains

suggests degeneration during domestication. Evol. Int. J. Org. Evol. *70*, 2099–2109.

Saleh, D., Xu, P., Shen, Y., Li, C., Adreit, H., Milazzo, J., Ravigné, V., Bazin, E., Nottéghem, J.-L., Fournier,

E., et al. (2012a). Sex at the origin: an Asian population of the rice blast fungus *Magnaporthe oryzae*

reproduces sexually. Mol. Ecol. *21*, 1330–1344.

Saleh, D., Milazzo, J., Adreit, H., Tharreau, D., and Fournier, E. (2012b). Asexual reproduction induces

a rapid and permanent loss of sexual reproduction capacity in the rice fungal pathogen *Magnaporthe*

oryzae: results of *in vitro* experimental evolution assays. BMC Evol. Biol. *12*, 42.

Saleh, D., Milazzo, J., Adreit, H., Fournier, E., and Tharreau, D. (2014). South-East Asia is the center of

origin, diversity and dispersion of the rice blast fungus, *Magnaporthe oryzae*. New Phytol. *201*, 1440–

1456.

Servedio, M.R., Van Doorn, G.S., Kopp, M., Frame, A.M., and Nosil, P. (2011). Magic traits in speciation:

"magic" but not rare? Trends Ecol. Evol. *26*, 389–397.

Sork, V.L., and Werth, S. (2014). Phylogeography of *Ramalina menziesii*, a widely distributed lichen-

forming fungus in western North America. Mol. Ecol. *23*, 2326–2339.

Taylor, J.W., and Fisher, M.C. (2003). Fungal multilocus sequence typing — it's not just for bacteria.

Curr. Opin. Microbiol. *6*, 351–356.

Taylor John W, Turner Elizabeth, Townsend Jeffrey P, Dettman Jeremy R, and Jacobson David (2006).

Eukaryotic microbes, species recognition and the geographic limits of species: examples from the

kingdom Fungi. Philos. Trans. R. Soc. B Biol. Sci. *361*, 1947–1963.

Tharreau, D., Fudal, I., Andriantsimialona, D., Santoso, Utami, D., Fournier, E., Lebrun, M.-H., and

Notteghem, J.-L. (2009). World Population Structure and Migration of the Rice Blast Fungus,

Magnaporthe oryzae (Dordrecht: Springer).

Walker, A.-S., Gladieux, P., Decognet, V., Fermaud, M., Confais, J., Roudet, J., Bardin, M., Bout, A.,

Nicot, P.C., Poncet, C., et al. (2015). Population structure and temporal maintenance of the multihost

fungal pathogen *Botrytis cinerea*: causes and implications for disease management. Environ. Microbiol.

17, 1261–1274.

Yang, N., Ma, G., Chen, K., and Wu, X. (2018). The population genetics of *Alternaria tenuissima* in four

regions of china as determined by microsatellite markers obtained by transcriptome sequencing.

Front. Microbiol. *9*.

Zhong, Z., Chen, M., Lin, L., Han, Y., Bao, J., Tang, W., Lin, L., Lin, Y., Somai, R., Lu, L., et al. (2018).

Population genomic analysis of the rice blast fungus reveals specific events associated with

expansion of three main clades. ISME J. *12*, 1867-1878.