Title: *Anopheles albimanus* natural microbiota is altered within one generation of laboratory

colonization

Authors:

Nsa Dada[a,d,#], Ana Cristina Benedict[b], Francisco López[b], Juan C. Lol[b], Mili Sheth[c], Nicole

Dzuris[a], Norma Padilla[b] and Audrey Lenhart[a]

Author affiliations

[a] Entomology Branch, Division of Parasitic Diseases and Malaria, Center for Global Health,

United States Centers for Diseases Control and Prevention, Atlanta, GA, United States of

America

[b] Grupo de Biología y Control de Vectores, Centro de Estudios en Salud, Universidad del Valle

de Guatemala, Guatemala

[c] Biotechnology Core Facility Branch, Division of Scientific Resources, National Center for

Emerging & Zoonotic Infectious Diseases, United States Centers for Disease Control and

Prevention, Atlanta, GA, United States of America

[d] American Society for Microbiology, Washington, DC, United States of America

#Address correspondence to Nsa Dada: nsadada@yahoo.com or nsa.dada@nmbu.no

Abstract

Research on mosquito-microbe interactions may lead to new tools for mosquito and mosquito-

borne disease control. To date, such research has largely utilized laboratory-reared mosquitoes

that typically lack the microbial diversity of wild populations. A logical progression in this area

involves working under controlled settings using field-derived mosquitoes or, in most cases, their

progeny. Thus, an understanding of how laboratory colonization affects the assemblage of

mosquito microbiota would aid in advancing mosquito microbiome studies and their applications

beyond laboratory settings. Using high throughput 16S rRNA amplicon sequencing, we

characterized the internal and cuticle surface microbiota of F_1 progeny of wild-caught

adult *Anopheles albimanus* from four locations in Guatemala. A total of 132 late instar larvae

and 135 2-5day old, non-blood-fed virgin adult females that were reared under identical

laboratory conditions, were pooled (3 individuals/pool) and analyzed. F_1 larvae from different

maternal sites showed different microbial compositions (p=0.001; F = 9.5), but these differences

were absent at the adult stage (p=0.12; F =1.6). These results suggest that field-acquired

mosquito microbiota may be lost in as early as the first generation of colonization under normal

laboratory conditions, thus, requiring adjustments to maintain field-derived microbiota. This is

the first time that the microbiota of F_1 progeny of wild-caught mosquitoes has been characterized

in relation to maternal collection site. Our findings provide a comprehensive background for

studying how parentage and environmental conditions differentially or concomitantly affect

mosquito microbiome composition, and how this can be exploited in advancing mosquito

microbiome studies and their applications beyond laboratory settings.

Keywords: Mosquito microbiota, *Anopheles albimanus*, laboratory colonization, mosquito

microbiome, next generation sequencing, 16S rRNA gene amplicon sequencing

Background

Mosquitoes contain microbes that inhabit various tissues, such as the alimentary canal,

reproductive organs, and cuticle surface [1]. These microbes are thought to be principally

obtained from the mosquito habitat during larval development, and from food sources at the

adult stage [1]. In addition to acquisition from larval habitats and/or adult food sources,

transovarial bacterial transmission from adult females to their eggs, and transstadial

transmission across different immature stages, and into the adult stage, have been

demonstrated [2, 3]. As a key component of the mosquito microbiota, environmentally-acquired

microbes have been shown to affect mosquito life history traits such as the rate of pupation and

adult body size [4]. The mosquito microbiota has also been shown to affect mosquito immunity

to human pathogens [5], reproduction [6], insecticide resistance [7, 8], and ultimately vector

competence—the mosquito's ability to acquire, maintain and transmit pathogens [5]. These

effects on multiple physiological characteristics are being leveraged to develop novel

approaches to fight mosquito-borne diseases [9].

The use of next generation molecular biology tools has resulted in extensive characterization of

mosquito microbiota, with the initial focus on bacterial and archaeal components now expanding

to eukaryotic microbes [10, 11] and viruses [12, 13]. These advances in mosquito microbiota

research have led to field applications of mosquito symbionts for mosquito control. For *Aedes*

aegypti, the principal vector of dengue, Zika, chikungunya and yellow fever viruses, mosquito-

derived symbionts are now being used to suppress mosquito populations [14] and also being

considered to control the spread of pathogens [15]. However, studies exploring mosquito

symbionts for malaria control have largely remained at the laboratory stage [16, 17]. Similarly,

the microbiota of mosquito vectors in some geographical regions are well characterized and

studied compared to those from other regions. In malaria vectors for example, studies on the

microbiota have largely focused on Sub-Saharan African species—in particular, *Anopheles*

gambiae—and to a lesser extent on those from Southeast Asia [18]. In contrast, the microbiota

of Latin American malaria vectors have only recently been comprehensively characterized [7, 8,

19-21], with these studies describing associations between *An. albimanus* microbiota and

insecticide resistance [7, 8], and the factors that shape the composition of *An. darlingi*, *An.*

albimanus, *An. nunetzovari*, *An. rangeli,* and *An. triannulatus* microbiota [19-21].

To exploit the mosquito microbiota for malaria and malaria vector control, research must

successfully advance from laboratory to field settings, a transition which can be fraught with

challenges. For example, some malaria vectors such as *An. darlingi, An. vestitipennis,* and *An.*

gambiae, breed in sites that are small, temporary and often difficult to find and/or access [22-26],

making it hard to obtain sufficient immature field mosquitoes for experiments. Where larval

habitats are plentiful and easy to find and/or access, the subsequent rearing of field-collected

mosquitoes to obtain uniform characteristics can pose additional challenges [27, 28].

Additionally, some malaria vectors belong to species complexes whose members are

morphologically indistinguishable [29-32], constituting another layer of complexity that needs to

be considered in elucidating mosquito-microbe interactions in malaria vectors.

These challenges, which are common to research on mosquito ecology and control, are often

not reported or discussed in mosquito microbiome studies. Several failed attempts at collecting

and rearing sufficient immature mosquitoes from the field for our previous study on the role of

mosquito microbiota in insecticide resistance resulted in ultimately using either wild-caught

adults [7] or F_1 progeny derived from field-collected adult mosquitoes [8, 33]. While field-caught

adult mosquitoes or their F_1 progeny may offer insights into mosquito-microbe interactions in

field scenarios, obtaining adult field-collected mosquitoes with uniform and/or controlled

physiological characteristics is usually not feasible. Although geographically associated

heterogeneity in microbiota of field-collected mosquitoes has been previously described [34, 35],

there is limited information on the fate of field-acquired microbiota after laboratory colonization of

field-collected mosquitoes. However, it has previously been observed that upon eclosion, newly-

emerged laboratory-reared adult mosquitoes show a reduction in bacterial diversity in contrast to

earlier developmental stages [36]. In addition, a recent study on the fate of field-acquired

microbiota in laboratory-colonized *An. gambiae s.l.* showed a reduction in bacterial diversity of

the F_5 progeny (the first point of measurement) that were reared in dechlorinated tap water in

contrast to F_0 [37].　Another study of *Ae. aegypti* from different geographical locations showed

no geographically associated heterogeneity microbiota composition after several generations of

laboratory colonization [38]. At what point the microbiota of laboratory-colonized mosquitoes

become homogenous and whether the microbiota of F_1 laboratory progeny represent their

parental origin—and thus could be used for symbiont-based translational studies—remains

largely undescribed.

Here, we present a comprehensive characterization of the microbiota of laboratory-reared F_1

progeny from field-caught adult *An. albimanus*. These represent unanticipated findings from a

larger study that was aimed at characterizing the effects of insecticide exposure and resistance

on *An. albimanus* microbiota [8]. We expected little or no immediate loss of geographical

heterogeneity in microbial composition upon initial laboratory colonization based on evidence of

this type of heterogeneity in field-derived populations [34, 35]. However, our data showed

geographical heterogeneity in F_1 larval internal and cuticle surface microbiota vs only in F_1 adult

cuticle surface microbiota. These findings lay the foundations for studying how parentage and

environmental conditions differentially or concomitantly affect mosquito microbiome composition,

and how this can be exploited in advancing mosquito microbiome studies and their applications

beyond laboratory settings.

Results

Using high throughput deep sequencing of the universal bacterial and archaeal 16S rRNA gene,

we characterized the microbiota of the internal and cuticle surface microbial niches of laboratory

reared *An. albimanus* F_1 larvae (n=132) and adult (n=135) progeny originating from mothers that

were collected from four locations in south central Guatemala (Fig 1). Mosquitoes were

processed as pools (3 individuals per pool), resulting in a total of 44 larval and 45 adult pools. F_1

larvae were from mothers collected in El Tererro, Las Cruces 3 and Las Cruces 4, while F_1

adults were from mothers collected in Las Cruces 1, Las Cruces 3 and Las Cruces 4 (Table 1).

Following quality control of the resulting sequencing reads, amplicon sequence variants (ASVs)

were identified and used in downstream analysis. Since adult mosquito microbiota is distinct

from that of immature stages [39, 40], ASVs from larvae and adults were analyzed separately,

as were ASVs from each microbial niche.

Internal and cuticle surface microbiota of laboratory colonized F_1 *An. albimanus* larvae

differed by maternal collection site

Non-pairwise Bray-Curtis distance comparison showed significant differences in internal

(p=0.001) and cuticle surface (p=0.001) microbiota between F_1 larvae from different maternal

collection sites. Thus, irrespective of microbial niche, the microbial community structure

(composition and relative abundance of ASVs) of F_1 laboratory-colonized larvae differed by

maternal collection location. Pairwise PERMANOVA comparison of Bray-Curtis distances further

showed significant differences in microbial community structure in larval internal (q<0.01) and

cuticle surface microbiota (q<0.01) between every pair of maternal collection site (Table 2a).

This location-driven heterogeneity in microbial community structure was further demonstrated by

principal coordinate analysis (PCoA), where F_1 larval internal and cuticle surface microbiota

clustered distinctly by maternal collection site (Fig. 2).

Non-pairwise Shannon diversity comparisons showed significant differences in internal

(p=0.009) but not cuticle surface (p=0.09) microbiota of F_1 laboratory-colonized larvae from

different maternal collection sites, indicating that there was inter-sample variation in the diversity

of internal but not cuticle surface microbiota of larvae when all maternal collection sites were

taken into consideration. A pairwise Kruskal-Wallis comparison of Shannon diversity indices

showed that the inter-sample variation in diversity of larval internal microbiota held true when

every pair of maternal collection sites was considered except between Las Cruces 3 and 4

(Table 2b and Suppl. 1). Larvae originating from Las Cruces 3 had the highest internal

microbiota diversity, followed by Las Cruces 4 and El Terrero (Suppl. 1)

Cuticle surface, but not internal, microbiota of laboratory colonized F_1 adult *An.*

***albimanus* differed by maternal collection site**

Non-pairwise Bray-Curtis diversity comparisons showed significant differences in cuticle surface

(p=0.001), but not internal microbiota (p=0.12) between adult F_1 mosquitoes from different

maternal collection sites, suggesting a loss of location-driven heterogeneity in microbial

community structure in internal but not cuticle surface microbial niche of laboratory-colonized F_1

adults. Pairwise PERMANOVA comparisons of Bray-Curtis distances also showed significant

differences in microbial community structure of F_1 adult cuticle surface microbiota (q<0.01)

between every pair of maternal collection sites (Table 2a). These results were corroborated by

PCoA which showed that F_1 adult cuticle surface microbiota, but not internal microbiota,

clustered distinctly by maternal collection site (Fig. 2).

Non-pairwise Shannon diversity comparisons showed no differences in the internal (p=0.42) or

cuticle surface (p=0.4) microbiota of F_1 adults from different maternal collection sites, indicating

that there was little or no inter-sample variation in diversity of F_1 adult microbiota when all

maternal collection sites were taken into consideration. Pairwise Kruskal-Wallis comparisons of

Shannon diversity indices also detected no inter-sample variation in diversity of F_1 adult cuticle

surface or internal microbiota when every pair of maternal collection site was considered (Table

2b and Suppl. 1).

Laboratory-colonized F_1 *An. albimanus* larvae comprised a rich and diverse microbiota

that differed by maternal collection site

Overall, ASVs from larval internal microbiota were assigned to 180 bacterial taxa, and cuticle

surface microbiota to 194 bacterial taxa (suppl 2.). A majority of these taxa across all locations

(ranging from 118-139 taxa) were shared between the internal and cuticle surface microbiota

(Fig 3a), as well as across maternal collection sites (n=110 for cuticle surface and n=117 for

internal microbiota) (Fig 3b). While a majority of the identified microbial taxa were shared

between both microbial niches, their abundance was generally higher in internal (Fig 4a)

compared to cuticle surface (Fig 4b) microbiota. Although a majority of identified microbial taxa

in both internal and cuticle surface microbiota were shared across all locations, their abundance

differed by maternal location (Fig 4a, 4b and 5).

In general, larval internal microbiota was dominated by ASVs identifed as an uncharacterized

Enterobacteriaceae, Leucobacter, Thorsellia, and *Chryseobacterium* (Fig 4a), together making

up over 50% of ASVs (Suppl. 2). In contrast, *Acidovorax,* unchracterized *Comamonadaceae,*

and *Paucibacter* (Fig 4b) made up over 50% of ASVs detected on the larval cuticle surface

(Suppl. 2).

A few predominant bacterial taxa were present in the larval internal microbial niche across all

three maternal collection sites: unclassified *Enterobacteriaceae, Thorsellia, Rhizobium,*

Xantobacter, Acidovorax and *Pirellula* (Fig 4a.), with remaining taxa showing different patterns of

abundance between collection sites (Fig 5 and Suppl. 2). For example, ASVs assigned to the

genus *Azoracus* were predominant in larvae from El Terrero, while *Singulisphaera, Paucibacter,*

Ancylobacter, Gemmobacter, and *Rayranella* were predominant in those from Las Cruces 3.

Predominant in larvae from Las Cruces 3 and Las Cruces 4 were *Azospirillum, Bosea* and

Microbacterium; and in those from El Terrero and Las Cruces 3 were *Terrimicrobium* and

Legionella (Fig 5). Although outside of the cut off limit set for differential abundance, ASVs

assigned to the bacterial genera *Leucobacter* were predominant in the internal microbiota of

larvae from Las Cruces 4 and El Terrero, but predominant in only three of the 14 sample

replicates from Las Cruces 3 (Fig 4a). Similarly, *Chryseobacterium* was predominant in Las

Cruces 4 and 3, but only predominant in six of the 16 pools of larvae from El Terrero. Bacterial

taxa that were unique to each location comprised <8% of all taxa in larval internal microbiota

(Suppl. 3b), and were below the threshold for inclusion in the heatmap and differential

abundance testing (Suppl. 2).

Unlike the internal microbial niche, no microbial taxa was predominant in larval cuticle surface

microbiota across all three collection sites. However, some taxa showed notable patterns of

abundance between locations (Fig 5). These included the genus *Azoarcus,* which was detected

at low to moderate frequencies in 13 of 15 pools of larvae from El Terrero, at low frequency in a

single pool of larvae from Las Cruces 3, and was not detected at all in Las Cruces 4 (Fig 4b and

6). Similarly, ASVs assigned to the genus *Spirosoma* were detected at moderate frequencies in

all pools of larvae from Las Cruces 4, but only in a few pools from the other two locations. ASVs

assigned to the genus *Paucibacter* were present at relatively higher abundance in larvae from

both Las Cruces 3 and El Terrero compared to those from Las Cruces 4. Those assigned to the

genus *Acidovorax* were predominant in larvae from Las Cruces 3 and Las Cruces 4 in contrast

to El Terrero. ASVs assigned to *Microbacterium, Bdellovibrio* and *Pelomonas* were present at

moderate frequencies in larvae from both Las Cruces 3 and Las Cruces 4 but were not detected

in El Terrero(Fig. 5). Bacterial taxa that were unique to each maternal collection site comprised

<8% of larval cuticle surface microbiota (Fig 3b), and were below the threshold for inclusion in

the heatmap and differential abundance testing (Suppl. 2).

Laboratory-colonized adult F$_1$ *An. albimanus* were comprised of sparse internal and

cuticle surface microbiota that were dominated by ASVs assigned to the genus *Asaia*

ASVs from adult internal microbiota were assigned to 62 microbial taxa and cuticle surface

microbiota were assigned to 106 microbial taxa. Two of these ASVs which were only present in

the cuticle surface microbiota were classified as archaea, while all other remaining ASVs were

classified as bacteria (Suppl. 2). Unlike larval microbiota, less than half of the assigned taxa

across all locations (ranging from 19-37 taxa) were shared between internal and cuticle surface

microbiota (Fig 3a), and only 18 taxa on the cuticle surface and 19 internal taxa were shared

across all maternal collection sites (Fig 3b).

Overall, ASVs assigned to the bacterial genus *Asaia* dominated both adult internal and cuticle

surface microbiota (Fig 6), constituting at least 70% of taxa in each microbial niche (Suppl. 2). A

majority of identified taxa in adult internal, but not cuticle surface, microbiota was detected

across all three maternal collection sites, with a few of these taxa present in high abundance

across all collection sites (Fig. 6). Across all three maternal collection sites, ASVs assigned to

the genera *Acinetobacter, Gluconobacter, Pantoea* and *Pseudomonas* were present in

moderate to high abundance in adult internal microbiota in addition to *Asaia* (Fig 6).

After excluding low abundance taxa (Suppl. 2), the number of remaining taxa did not meet the

requirements for identifying collection site-specific microbial taxa in adult cuticle surface

microbiota. This was compounded by dominance (>70%) of ASVs that were assigned to the

bacterial genus *Asaia* (Suppl. 2). In addition, the cuticle surface microbiota of adults originating

from Las Cruces 4 comprised 43% of all adult cuticle surface microbial taxa (Fig 3b), although a

majority were of low abundance.

Discussion

The scientific community is increasingly investigating the role of mosquito microbiota in fighting mosquito-borne diseases [9]. The successful transition of mosquito microbiome research from laboratory to field requires a comprehensive understanding of the dynamics underlying the composition of the microbiota of field-collected mosquitoes and their progeny. We provide a comprehensive characterization of the internal and cuticle surface microbiota of laboratory-reared F_1 progeny from field-caught adult *An. albimanus* that were collected from different locations. Our results show that while location-driven heterogeneity in the microbial community structure of both internal and cuticle surface microbiota was present in the F_1 larvae, this heterogeneity was only evident in the cuticle surface microbiota of adult progeny from the same generation. This work provides comprehensive fundamental data for studying how parentage and environmental conditions differentially or concomitantly affect mosquito microbiome composition.

Previous studies on other mosquito species have also shown a loss of field-acquired internal mosquito microbiota following several generations of laboratory colonization [37, 38]. Until now, however, the laboratory generation at which this loss occurs was largely undescribed. Our study represents an initial step in filling this knowledge gap and provides further details regarding the dynamics of changes in both the internal and cuticle surface microbial niches. The homogeneity in internal microbiota of F_1 adults is suggestive of a loss in field-acquired microbiota, which could pose an additional set of challenges for field or semi-field mosquito microbiome studies—particularly studies that rely on laboratory progeny in lieu of field populations to circumvent the challenges involved [25, 27, 28] in obtaining sufficient numbers of wild mosquitoes of uniform physiological characteristics. Previous findings showed that rearing mosquitoes in water from the field could preserve the field-derived internal microbiota in laboratory-colonized adult progeny of

An. gambiae s. l. for several generations [37]. In addition, other studies, have shown that the

microbiota in larval habitat water significantly influences the internal microbiota of emerging adult

mosquitoes [40, 41]. Thus, to preserve wild-type microbiota, care should be taken to ensure that

the first laboratory generation are reared in water collected from larval habitats in the field. This

first generation would be most suitable for studies that require progeny from field populations.

Maternal egg-smearing has been proposed as a mechanism through which adult female

mosquitoes and other insects transfer microbes to their progeny [42-45]. This egg-smearing

could explain the heterogeneity observed in internal and cuticle surface microbiota between F_1

larvae with different maternal origins. On the other hand, a lack of this heterogeneity in the

internal microbiota of adult progeny could be attributed to the physiological changes that occur

during metamorphosis and adult eclosion, whereby elimination of the larval meconial peritrophic

membrane and meconium (midgut and midgut content), along with ingestion of exuvial fluid—

which is said to be bactericidal—results in sterile or nearly sterile midguts in newly emerged

adults [39, 46]. This, in addition to mechanisms that regulate the composition of adult mosquito

internal microbiota [47-49], could explain the observed homogeneity in F_1 adult internal

microbiota. Conversely, the heterogeneity in cuticle surface microbiota between adult progeny

originating from different maternal sites, suggests that maternally derived microbes in the rearing

trays may have colonized adult cuticle surfaces during emergence. The mechanisms underlying

the assemblage of the mosquito cuticle surface microbiome are largely undescribed, and thus

require further investigation.

The low inter-sample variation in microbial diversity observed in this study has largely been

described in laboratory mosquito colonies [38, 50]. The microbial composition of laboratory-

reared larvae is typically less diverse [50, 51] compared to those of field-derived larvae, but our

laboratory-reared larvae exhibited a rich microbial composition that was comparable to those of

field populations [39, 40]. In contrast, our adult progeny had a less diverse microbial composition

that was reflective of typical laboratory-reared adult mosquitoes [36, 52]. These findings further

suggest that field-acquired microbiota, although transferred to laboratory progeny, may be lost

within one generation of laboratory colonization—particularly at the adult stage.

In this study, we detected microbial taxa that have previously been identified in *Anopheles* and

other mosquito genera [7, 41, 53, 54]. While a majority of the taxa in F_1 larvae were shared

between both the internal and cuticle surface microbial niches, a greater abundance of microbial

taxa was detected in the internal microbial niche compared to the cuticle surface. The cuticle

surface microbiota of mosquitoes and other hematophagous insects are largely uncharacterized

and the mechanisms underlying their assemblage remain unknown. As such, we hypothesize

that although both internal and cuticle surface niches are exposed to the same water from which

the microbiota is derived, a more conducive and protected internal environment could allow for

greater proliferation of colonizing bacteria. In F_1 adults however, less than half of the detected

microbial taxa were shared between the internal and cuticle surface microbial niches, suggesting

differences in physiological conditions that favor microbial colonization, and corroborating

findings that point toward a microbial regulatory mechanisms within the mosquito midgut [47-49].

This minimal overlap of taxa between internal and cuticle surface microbial niches also suggests

a difference in the fate of maternally derived microbes between both niches. Although a few

microbial taxa overlapped between adult internal and cuticle surface microbial niches and the

most abundant taxa were shared, many of the unshared taxa have been previously detected in

adult mosquitoes including *Anopheles* [1, 54, 55], indicating that the cuticle surface microbiota

characterized in this study are inherently associated with mosquitoes. Like the larval microbiota,

there was a higher abundance of microbial taxa in the adult internal microbial niche compared to

the cuticle surface, further supporting the hypothesis of a more conducive and protected internal

environment permitting greater proliferation of colonizing bacteria.

With the exception of the adult cuticle surface microbial niche, a majority of all detected microbial

taxa overlapped between maternal collection sites in both F_1 larvae and adults, albeit with

differing abundances. This reflects restrictions imposed by controlled laboratory environments in

the development of mosquito microbiota. In both microbial niches of both larvae and adults,

microbial taxa that were specific to maternal collection sites were low in abundance, compared

to the moderate to high abundance of those that were shared across all locations. This was

particularly true for *Asaia*—notorius for rapidly colonizing laboratory mosquitoes [56]—which

constituted at least 70% of both adult internal and cuticle surface microbiota from progeny

across all maternal collection sites. These results suggest that field-acquired mosquito

microbiota may be lost in as early as the first generation of laboratory colonization.

We recognize that not having the microbial community profiles of the mothers from which the F_1

progeny were derived is a limitation of this study. However, these are unanticipated findings that

arose from a separate study whose focus was on field-derived F_1 progeny reared under

controlled settings [8]. The findings herein provide empirical data for further exploring the role of

parentage and environmental conditions on the assemblage of the mosquito microbiome, and

the fate of field-derived microbes upon laboratory colonization. This is critical for advancing

mosquito microbiome studies and their applications beyond laboratory settings.

Methods

The findings presented here extend those of a larger study [8]. Thus, the mosquito collection,

processing and sequencing procedures have previously been described in detail [8].

Mosquito collections and laboratory generation of F$_1$ progeny

Gravid and/or blood-fed adult female *An. albimanus* were sampled across four field sites in the

villages of Las Cruces and El Terrero, in La Gomera, department of Escuintla, Guatemala

(Figure 1). Field-collected mosquitoes from each location were held in separate paper cups,

sustained on 10% sucrose solution, and transported to the insectary at Universidad del Valle de

Guatemala in Guatemala City for species identification, oviposition and subsequent rearing of F$_1$

progeny. Mosquitoes that were morphologically identified as *An. albimanus* following

identification keys [57] (approximately 300 in total) were subjected to oviposition, and a

subsample of the resulting F$_1$ progeny were used for molecular verification of species identity as

described below. A modified oviposition procedure [58] was employed, wherein no more than 70

gravid females from the same location were placed in quart size paper ice cream containers with

distilled water to a depth of at least 2 cm. The containers were covered with fine mesh fabric that

were secured with rubber bands prior to the introduction of gravid field-caught adult *An.*

albimanus. The mesh was topped with cotton balls soaked in 10% sucrose to sustain the

mosquitoes, and subsequently covered with a thick piece of black plastic bag to keep the

containers dark and trap in moisture. The oviposition chambers were held under the following

insectary conditions; $27\pm2°C$, $80\pm10\%$ relative humidity, and 12-h light-dark cycle. After at least

48 h, the adult females were removed and discarded as they were not needed for the original

study. The F$_1$ eggs were collected, pooled by maternal location and reared separately under

identical ambient conditions (as described above) and the following larval feeding regimen. Eggs

were washed into 18 x 14 x 3 inch plastic larval trays (approximately 200 eggs per tray)

containing distilled water to a depth of at least 2 cm, and 3-4 drops of 10% yeast solution.

Hatched larvae were sustained on finely ground Koi fish food (Foster & Smith, Inc. Rhinelander,

WI) until pupation. At the third to fourth larval instar stage (L3-L4), half of the larvae were

separated and used for bioassays in the original study and subsequently processed for

microbiota characterization (n=132). Using a stereo microscope, female pupae from the

remaining mosquitoes were separated into 8 oz. paper ice cream cups and placed into

cardboard cages for adult eclosion. The resulting F_1 adult virgin female mosquitoes were

sustained on 10% sucrose solution until they were 2-5 days old and also used for bioassays in

the parent study. These were subsequently processed for microbiota characterization (n=135).

Post bioassays, larval and adult samples were preserved in RNALater® solution (Applied

Biosystems, Foster City, CA), shipped on dry ice to the US Centers for Disease Control and

Prevention (CDC) in Atlanta, USA, and stored at -80 °C until further processing. Samples

originating from different geographic locations were handled, stored and processed separately.

Genomic DNA extraction from mosquito legs, whole mosquito samples, and mosquito

cuticle surfaces

Stored mosquito samples were thawed overnight at 4°C, and the thawed RNALater® solution

was discarded. This was followed by a rinse with nuclease-free water to remove any residual

RNALater®. Legs from a subsample of individual F_1 adults were removed using sterile forceps

and placed in individual sterile1.5 mL Eppendorf tubes containing 75 μL Extracta DNA Prep

solution (Quantabio, Beverly, MA) for DNA extraction following manufacturer's instructions. Both

larvae and adult (with and without legs removed) F_1 samples were pooled for whole body and

cuticle surface DNA extractions. Each pool comprised 3 individuals, resulting in 44 pools of

larvae and 45 pools of adults. Table 1 shows the number of pools (replicates) processed per life

stage and maternal collection site. To obtain cuticle surface samples, each pool was submerged

in 500 μL of nuclease free water and washed vigorously by agitating with a vortex mixer for at

least 15 seconds. The resulting wash water was transferred to new sterile tubes for DNA

extraction. The washed sample pools were further surface sterilized using two vigorous washes;

first in 70% ethanol, then nuclease free water, each with at least 15 seconds agitation on a

vortex mixer. This was followed by one gentle rinse with nuclease free water. Genomic DNA

from the surface-sterilized samples and wash water (referred to as internal and cuticle surface

respectively) was isolated using the TissueLyser II and DNeasy Blood and Tissue Kit (QIAGEN,

Hilden, Germany) as follows; in individual 2 mL sterile tubes, 180 μL of buffer ATL (QIAGEN)

and 5 mm diameter stainless steel beads (QIAGEN) were added to each sample pool. With the

following settings; 30 hz/s for 8 and 15 minutes for internal and cuticle surface samples

respectively, samples were homogenized using the TissueLyser (QIAGEN) fitted with two sets of

96-well adapter plates that held the tubes. The plates were rotated every minute during

homogenization, and afterwards, samples were transferred into new sterile 1.5 mL Eppendorf

tubes for the remaining part of the DNA extraction process. 20 μL of Proteinase K (QIAGEN)

was added to each homogenized sample and incubated overnight at 56 °C, after which 200 μL

of buffer AL (QIAGEN) was added and incubated for a further 2 h at the same temperature. The

remaining steps were performed according to QIAGEN's spin-column protocol for purification of

DNA from animal tissues, and the purified DNA was eluted in 70 μL of buffer AE (QIAGEN). Two

sets of two blank controls (without samples) were processed alongside the internal and cuticle

surface samples, and all steps, along with those described below, were performed under sterile

conditions. The purified genomic DNA from samples and blank controls were stored at -80 °C

until further processing.

Molecular species confirmation of *An. albimanus* and 16S rRNA amplicon sequencing

DNA samples from mosquito legs were used as templates to amplify the second internal

transcribed spacer region (ITS2) of the mosquito ribosomal DNA in order to verify morphological

identification of *An. albimanus*. This was achieved by conventional PCR using the universal ITS2

primers (ITS2 A: TGTGAACTGCAGGACACAT and ITS2 B: TATGCTTAAATTCAGGGGGT) for

distinguishing members of *Anopheles* complexes [59]. With a final reaction volume of 25 μL,

each PCR comprised $\geq$100 ng/μL DNA template, 15 μM of each primer, 12.5 μL of 2X AccuStart

II PCR SuperMix (Quantabio, Beverly, MA), and PCR grade water to final volume. With the

following conditions, initial denaturation at 94 °C for 4 min, then 35 cycles of 94 °C for 30 s, 53 °C

for 40 s, and 72 °C for 30 s, followed by a final extension for 10 min at 72 °C, the PCRs were

performed using a T100™ Thermal Cycler (Bio-Rad, USA). Using EtBr-stained agarose gel

electrophoresis, the amplified products sized ~500 bp, confirmed all samples as *An. albimanus*

[60].

DNA from internal and cuticle surface samples, along with those from blank controls, were used

as templates to amplify the universal bacterial and archaeal 16S rRNA gene. This was also

achieved by conventional PCR using universal primers targeting the v3-v4 region of the 16S

rRNA gene (341F:

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG, and 805R:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC), each

with Illumina® (San Diego, CA USA) adapters (bold typeface). With a final reaction volume of 25

μL, each PCR comprised; ≥20 ng/µL DNA template, 5 µM of each primer, 10 µL of 2× KAPA HiFi

HotStart PCR mix (Roche, Switzerland), and PCR grade water to final volume. Three negative

controls with PCR grade water substituted for DNA template, were processed along with the

samples. The PCRs were performed using the T100™ Thermal Cycler (Bio-Rad, USA) under

the following conditions; initial denaturation at 95 °C for 3 min, then 25 cycles of 95 °C, 55 °C,

and 72 °C for 30 s each, followed by a final extension for 5 min at 72 °C. The amplification

products, sized ~460 bp, were verified by electrophoreses as described above and quantified

using a NanoDrop™ spectrophotometer (Thermo Fisher Scientific, Waltham, MA). Amplicons,

including PCR products from blanks and negative controls, which yielded no bands following

electrophoresis, were purified using Agencourt AMPure XP beads (Beckman Coulter Inc.,

Indianapolis, IN, USA) at 0.7X (internal) or 0.875X (cuticle surface) sample volume, and eluted in

40 µL of 10 mM Tris buffer (pH 8.5).

All purified products, including those of blanks and negative controls, were used as templates for

sequencing library preparation. This was accomplished via index PCR using the Nextera XT

Index kit v2 sets, A, B and D (Illumina, San Diego, CA). With a final reaction volume of 50 µL,

each index PCR comprised; 25 µL NEBNext High-Fidelity 2X PCR master mix (New England

Biolabs Inc., Ipswich, MA), 5 µL of each index primer, 10 µL of purified PCR products (0–

20 ng/µL) as template, and PCR grade water to final volume. The PCRs were performed under

the following reaction conditions; 98 °C for 30 s, then 8 cycles of 98 °C for 10 s, 55 °C and 65 °C

for 30 s each, and a final extension at 65 °C for 5 min, and resulting libraries were also cleaned

using Agencourt AMPure XP beads (Beckman Coulter Inc., Indianapolis, IN, USA) at 1.2X

sample volume and eluted in 25 µL of 10 mM Tris buffer (pH 8.5). The libraries were

subsequently analyzed for size and concentration, normalized to 2 nM, pooled and denatured

following Illumina guidelines for loading onto flowcells. Sequencing was performed on an

Illumina HiSeq 2500 machine, using 2x250 cycle paired-end sequencing kits.

Preprocessing of sequencing reads

Sequencing outputs were demultiplexed and converted to the fastq format for downstream

analysis using the bcl2fastq (v2.19) conversion software (Illumina®). A total of 115,250,077

demultiplexed paired-end sequencing reads, with a maximum length of 250 bp were initially

imported into the 'quantitative insights into microbial ecology' pipeline, QIIME2 v.2017.7.0 [61],

and further sequencing read processing and analysis were performed in v.2018.2.0 of the

pipeline. Using the DADA2 plugin in QIIME 2 [62], the `denoise-paired` command with the

following options; `trunc_len_f: 244, trunc_len_r: 244, and n_reads_learn:`

`500000`, was used to correct errors, remove chimeras and merge paired-end reads. The

resulting amplicon sequence variants (ASVs; n=30,956,883) were further filtered to remove

ASVs with <100 counts and ASVs that were associated with blanks and negative controls. This

last filtering step resulted in 17,225,776 ASVs, ranging from 3,277 to 223,222 per sample (mean

96,774), that were used for downstream comparison of bacterial composition and taxonomic

analysis. Suppl. 3. shows sequencing reads and ASV summary statistics.

Diversity indices

Analysis of microbial diversity within (alpha diversity) and between (beta diversity) samples were

performed in QIIME2 using the Shannon diversity index and Bray-Curtis dissimilarity index,

respectively. The Shannon diversity indices were calculated using rarefied ASVs counts per

sample, in which ASVs per sample were selected randomly without replacement at an even

depth (Suppl. 4) for ten iterations. The resulting average Shannon indices are presented and

were compared between samples using pairwise Kruskal-Wallis tests with Benjamini-Hochberg

false discovery rate (FDR) corrections for multiple comparisons.

The Bray-Curtis dissimilarity indices were computed with or without rarefaction, and resulting

indices were compared between samples using pairwise PERMANOVA tests (999 permutations)

with FDR corrections. There were no discernable differences between results of rarefied and

non-rarefied data. Thus, results of Bray-Curtis dissimilarity indices using non-rarefied data were

visualized by Principal Co-ordinates Analysis (PCoA) plots in R [63] using the phyloseq R

package [64].

Significance for both pair-wise analyses was set to $q < 0.05$ (i.e. post FDR p-value corrections).

Taxonomic analysis and differentially abundant microbial taxa

Taxonomic analysis of ASVs was performed using QIIME2's pre-trained Naïve Bayes classifier

[65] and q2-feature-classifier plugin [66]. Prior to analysis, the classifier was trained on the

QIIME-compatible 16S SILVA reference (99% identity) database v.128 [67], and using the

extract-reads command of the `q2-feature-classifier` plugin, the reference sequences

were trimmed to the v3-v4 region (425 bp) of the 16S rRNA gene. The relative abundance of

annotated ASVs across samples were subsequently visualized using the `qiime feature-`

`table heatmap` plugin based on Bray-Curtis distances with the plugin's default clustering

method.

Differentially abundant microbial taxa across locations were identified using QIIME2's analysis of

composition of microbiomes (ANCOM) [68] plugin. The cut-off for differential abundance was set

to an effect size of log F≥20 and W≥20, i.e. a taxon was differentially abundant across maternal

collection sites if the ratio of its abundance to those of at least 20 other taxa (25% of all included

taxa) differed significantly across sites.

Only annotated ASVs with counts ≥ 2000 (larvae) or ≥ 1000 (adults) were included in the

heatmaps and ANCOM analysis. Prior to each analysis, ASV frequency data was normalized by

$\log_{10}$ transformation following the addition of pseudocounts of 1.

The outputs of data analyses were aesthetically formatted using Inkscape [69]**.**

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and material: The raw sequencing reads generated from this project, including those from negative controls (blanks), have been deposited in the National Center for Biotechnology Information (NCBI), Sequence Read Archive under the BioProject PRJNA512122.

Competing interests: The authors declare that they have no competing interests

Funding: This work was supported by the US Centers for Disease Control and Prevention (CDC) through the American Society for Microbiology's (ASM) Infectious Disease and Public Health Microbiology Postdoctoral Fellowship awarded to ND†, and the CDC's Advanced Molecular Detection (AMD) program.

Author contributions: ND† & AL conceptualized and designed the study; NP facilitated and provided facilities for field work; ND†, ACB, FL & JCL performed mosquito collections, and mass rearing; ND† & MS performed molecular analysis and sequencing; ND† analyzed the data; ND† and ND performed the data visualizations; ND† drafted the manuscript; all authors reviewed and approved the final version of the manuscript.

Acknowledgements

We thank the cattle coral owners in Escuintla, Guatemala for permission to conduct mosquito collections on their corals, without which this study would not have been possible; the Malaria Research and Reference Reagent Resource Center (MR4) for providing the ITS2 primers used

519   **Authors' information:** ND[†] present address: Faculty of Science and Technology, Norwegian

520   University of Life Sciences, Aas, Norway

521   [†]Nsa Dada

References

1. Guégan M, Zouache K, Démichel C, Minard G, Tran Van V, Potier P, et al. The mosquito holobiont: fresh insight into mosquito-microbiota interactions. Microbiome. 2018;6(1):49. doi: 10.1186/s40168-018-0435-2.

2. Coon KL, Brown MR, Strand MR. Mosquitoes host communities of bacteria that are essential for development but vary greatly between local habitats. Molecular Ecology. 2016;25(22):5806-26. doi: 10.1111/mec.13877.

3. Lindh JM, Borg-Karlson AK, Faye I. Transstadial and horizontal transfer of bacteria within a colony of Anopheles gambiae (Diptera: Culicidae) and oviposition response to bacteria-containing water. Acta Tropica. 2008;107(3):242-50. doi: http://dx.doi.org/10.1016/j.actatropica.2008.06.008.

4. Dickson LB, Jiolle D, Minard G, Moltini-Conclois I, Volant S, Ghozlane A, et al. Carryover effects of larval exposure to different environmental bacteria drive adult trait variation in a mosquito vector. Science Advances. 2017;3(8). doi: 10.1126/sciadv.1700585.

5. Dennison NJ, Jupatanakul N, Dimopoulos G. The mosquito microbiota influences vector competence for human pathogens. Current opinion in insect science. 2014;3:6-13. doi: 10.1016/j.cois.2014.07.004. PubMed PMID: PMC4288011.

6. Coon KL, Brown MR, Strand MR. Gut bacteria differentially affect egg production in the anautogenous mosquito Aedes aegypti and facultatively autogenous mosquito Aedes atropalpus (Diptera: Culicidae). Parasites & Vectors. 2016;9(1):375. doi: 10.1186/s13071-016-1660-9.

7. Dada N, Sheth M, Liebman K, Pinto J, Lenhart A. Whole metagenome sequencing reveals links between mosquito microbiota and insecticide resistance in malaria vectors. Scientific Reports. 2018;8(1):2084. Epub 2018/02/03. doi: 10.1038/s41598-018-20367-4. PubMed PMID: 29391526; PubMed Central PMCID: PMCPMC5794770.

8. Dada N, Lol JC, Benedict AC, López F, Sheth M, Dzuris N, et al. Pyrethroid exposure alters internal and cuticle surface bacterial communities in Anopheles albimanus. The ISME Journal. 2019. doi: 10.1038/s41396-019-0445-5.

9. Huang W, Wang S, Jacobs-Lorena M. Use of Microbiota to Fight Mosquito-Borne Disease. 2020;11(196). doi: 10.3389/fgene.2020.00196.

10. Luis P, Vallon L, Tran F-H, Hugoni M, Tran-Van V, Mavingui P, et al. Aedes albopictus mosquitoes host a locally structured mycobiota with evidence of reduced fungal diversity in invasive populations. Fungal Ecology. 2019;39:257-66. doi: https://doi.org/10.1016/j.funeco.2019.02.004.

11. Thongsripong P, Chandler JA, Green AB, Kittayapong P, Wilcox BA, Kapan DD, et al. Mosquito vector-associated microbiota: Metabarcoding bacteria and eukaryotic symbionts across habitat types in Thailand endemic for dengue and other arthropod-borne diseases. Ecology and evolution. 2018;8(2):1352-68. Epub 2018/01/30. doi: 10.1002/ece3.3676. PubMed PMID: 29375803; PubMed Central PMCID: PMCPMC5773340.

12.   Shi C, Beller L, Deboutte W, Yinda KC, Delang L, Vega-Rúa A, et al. Stable distinct core
eukaryotic viromes in different mosquito species from Guadeloupe, using single mosquito viral
metagenomics. Microbiome. 2019;7(1):121. doi: 10.1186/s40168-019-0734-2.

13.   Atoni E, Wang Y, Karungu S, Waruhiu C, Zohaib A, Obanda V, et al. Metagenomic
Virome Analysis of Culex Mosquitoes from Kenya and China. Viruses. 2018;10(1). Epub
2018/01/13. doi: 10.3390/v10010030. PubMed PMID: 29329230; PubMed Central PMCID:
PMCPMC5795443.

14.   Crawford JE, Clarke DW, Criswell V, Desnoyer M, Cornel D, Deegan B, et al. Efficient
production of male Wolbachia-infected Aedes aegypti mosquitoes enables large-scale
suppression of wild populations. Nature Biotechnology. 2020;38(4):482-92. doi:
10.1038/s41587-020-0471-x.

15.   Flores HA, Taneja de Bruyne J, O'Donnell TB, Tuyet Nhu V, Thi Giang N, Thi Xuan
Trang H, et al. Multiple Wolbachia strains provide comparative levels of protection against
dengue virus infection in Aedes aegypti. PLOS Pathogens. 2020;16(4):e1008433. doi:
10.1371/journal.ppat.1008433.

16.   Capone A, Ricci I, Damiani C, Mosca M, Rossi P, Scuppa P, et al. Interactions between
Asaia, Plasmodium and Anopheles: new insights into mosquito symbiosis and implications in
Malaria Symbiotic Control. 2013;6(1):182. doi: 10.1186/1756-3305-6-182.

17.   Wang S, Dos-Santos ALA, Huang W, Liu KC, Oshaghi MA, Wei G, et al. Driving
mosquito refractoriness to *Plasmodium falciparum* with engineered symbiotic bacteria.
2017;357(6358):1399-402. doi: 10.1126/science.aan5478 %J Science.

18.   Villegas LM, Pimenta PFP. Metagenomics, paratransgenesis and the Anopheles
microbiome: a portrait of the geographical distribution of the anopheline microbiota based on a
meta-analysis of reported taxa. Memórias do Instituto Oswaldo Cruz. 2014;109:672-84.

19.   Bascuñán P, Niño-Garcia JP, Galeano-Castañeda Y, Serre D, Correa MMJM. Factors
shaping the gut bacterial community assembly in two main Colombian malaria vectors.
2018;6(1):148. doi: 10.1186/s40168-018-0528-y.

20.   Galeano-Castañeda Y, Urrea-Aguirre P, Piedrahita S, Bascuñán P, Correa MM.
Composition and structure of the culturable gut bacterial communities in Anopheles albimanus
from Colombia. PLOS ONE. 2019;14(12):e0225833. doi: 10.1371/journal.pone.0225833.

21.   Prussing C, Saavedra MP, Bickersmith SA, Alava F, Guzmán M, Manrique E, et al.
Malaria vector species in Amazonian Peru co-occur in larval habitats but have distinct larval
microbial communities. PLOS Neglected Tropical Diseases. 2019;13(5):e0007412. doi:
10.1371/journal.pntd.0007412.

22.   Hiwat H, Bretas G. Ecology of Anopheles darlingi Root with respect to vector importance:
a review. Parasites & Vectors. 2011;4(1):177. doi: 10.1186/1756-3305-4-177.

23.   Villarreal-Treviño C, Penilla-Navarro RP, Vázquez-Martínez MG, Moo-Llanes DA, Ríos-
Delgado JC, Fernández-Salas I, et al. Larval habitat characterization of Anopheles darlingi from

its northernmost geographical distribution in Chiapas, Mexico. Malaria Journal. 2015;14(1):517.
doi: 10.1186/s12936-015-1037-0.

24.   Fillinger U, Sombroek H, Majambere S, van Loon E, Takken W, Lindsay SW. Identifying
the most productive breeding sites for malaria mosquitoes in The Gambia. Malaria Journal.
2009;8(1):62. doi: 10.1186/1475-2875-8-62.

25.   Gimnig JE, Ombok M, Kamau L, Hawley WA. Characteristics of Larval Anopheline
(Diptera: Culicidae) Habitats in Western Kenya. Journal of Medical Entomology. 2001;38(2):282-
8. doi: 10.1603/0022-2585-38.2.282 %J Journal of Medical Entomology.

26.   Grieco JP, Johnson S, Achee N, Masuoka P, Pope K, Rejmánková E, et al. Distribution
of *Anopheles albimanu*s, *Anopheles vestitipennis*, and *Anopheles crucians* associated with land
use in northern Belize. 2006;43 Journal of Medical Entomology(3):614-22, 9.

27.   Tchigossou G, Akoton R, Yessoufou A, Djegbe I, Zeukeng F, Atoyebi SM, et al. Water
source most suitable for rearing a sensitive malaria vector, Anopheles funestus in the laboratory.
Wellcome open research. 2017;2:109-. doi: 10.12688/wellcomeopenres.12942.2. PubMed
PMID: 29387806.

28.   Benedict MQ, Knols BGJ, Bossin HC, Howell PI, Mialhe E, Caceres C, et al. Colonisation
and mass rearing: learning from others. Malaria Journal. 2009;8(2):S4. doi: 10.1186/1475-2875-
8-S2-S4.

29.   Organization WH. Anopheline species complexes in south-east Asia: WHO Regional
Office for South-East Asia; 1998.

30.   Rosa-Freitas MG, Lourenço-de-Oliveira R, Carvalho-Pinto CJd, Flores-Mendoza C,
Silva-do-Nascimento TF. Anopheline Species Complexes in Brazil. Current Knowledge of Those
Related to Malaria Transmission %J Memórias do Instituto Oswaldo Cruz. 1998;93:651-5.

31.   Cohuet A, Simard F, Toto J-C, Kengne P, Coetzee M, Fontenille D. Species identification
within the *Anopheles funestus* group of malaria vectors in Cameroon and evidence for a new
species 2003;69(2):200-5. doi: https://doi.org/10.4269/ajtmh.2003.69.200.

32.   Stevenson JC, Norris DE. Implicating Cryptic and Novel Anophelines as Malaria Vectors
in Africa. Insects. 2016;8(1). Epub 2016/12/28. doi: 10.3390/insects8010001. PubMed PMID:
28025486; PubMed Central PMCID: PMCPMC5371929.

33.   Omoke DN, Mugendi E, Ochomo E, Kipsum M, Otieno S, Esalimba E, et al., editors.
Bacterial composition differs between permethrin-susceptible and -resistant Anopheles gambiae
sensu stricto in a site with intense pyrethroid resistance in western Kenya. American Journal of
Tropical Medicine and Hygiene 2019: American Society of Tropical Medicine and Hygiene 8000
Westpark DR, STE 130, MCLEAN, VA 22101 USA.

34.   Duguma D, Hall MW, Smartt CT, Debboun M, Neufeld JD. Microbiota variations in Culex
nigripalpus disease vector mosquito of West Nile virus and Saint Louis Encephalitis from
different geographic origins. PeerJ. 2019;6:e6168. doi: 10.7717/peerj.6168.

35.	Muturi EJ, Lagos-Kutz D, Dunlap C, Ramirez JL, Rooney AP, Hartman GL, et al.
Mosquito microbiota cluster by host sampling location. Parasites & Vectors. 2018;11(1):468. doi:
10.1186/s13071-018-3036-9.

36.	Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut microbiota for
development. Mol Ecol. 2014;23(11):2727-39. Epub 2014/04/29. doi: 10.1111/mec.12771.
PubMed PMID: 24766707; PubMed Central PMCID: PMCPMC4083365.

37.	Akorli J, Namaali PA, Ametsi GW, Egyirifa RK, Pels NAP. Generational conservation of
composition and diversity of field-acquired midgut microbiota in Anopheles gambiae (sensu lato)
during colonization in the laboratory. Parasites & Vectors. 2019;12(1):27. doi: 10.1186/s13071-
019-3287-0.

38.	Dickson LB, Ghozlane A, Volant S, Bouchier C, Ma L, Vega-Rúa A, et al. Diverse
laboratory colonies of Aedes aegypti harbor the same adult midgut bacterial microbiome.
Parasites & Vectors. 2018;11(1):207. doi: 10.1186/s13071-018-2780-1.

39.	Duguma D, Hall MW, Rugman-Jones P, Stouthamer R, Terenius O, Neufeld JD, et al.
Developmental succession of the microbiome of Culex mosquitoes. 2015;15(1):140. doi:
10.1186/s12866-015-0475-8.

40.	Gimonneau G, Tchioffo MT, Abate L, Boissière A, Awono-Ambéné PH, Nsango SE, et al.
Composition of Anopheles coluzzii and Anopheles gambiae microbiota from larval to adult
stages. Infection, Genetics and Evolution. 2014;28:715-24. doi:
https://doi.org/10.1016/j.meegid.2014.09.029.

41.	Saab SA, Dohna Hz, Nilsson LKJ, Onorati P, Nakhleh J, Terenius O, et al. The
environment and species affect gut bacteria composition in laboratory co-cultured Anopheles
gambiae and Aedes albopictus mosquitoes. Scientific Reports. 2020;10(1):3352. doi:
10.1038/s41598-020-60075-6.

42.	Damiani C, Ricci I, Crotti E, Rossi P, Rizzi A, Scuppa P, et al. Mosquito-Bacteria
Symbiosis: The Case of Anopheles gambiae and Asaia. Microbial Ecology. 2010;60(3):644-54.
doi: 10.1007/s00248-010-9704-8.

43.	Rossi P, Ricci I, Cappelli A, Damiani C, Ulissi U, Mancini MV, et al. Mutual exclusion of
Asaia and Wolbachia in the reproductive organs of mosquito vectors. Parasites & Vectors.
2015;8(1):278. doi: 10.1186/s13071-015-0888-0.

44.	Benedict MQ. Transgenic insects: techniques and applications: CABI; 2014.

45.	Crotti E, Rizzi A, Chouaia B, Ricci I, Favia G, Alma A, et al. Acetic Acid Bacteria, Newly
Emerging Symbionts of Insects. 2010;76(21):6963-70. doi: 10.1128/AEM.01336-10 %J Applied
and Environmental Microbiology.

46.	Moll RM, Romoser WS, Modrzakowski MC, Moncayo AC, Lerdthusnee K. Meconial
peritrophic membranes and the fate of midgut bacteria during mosquito (Diptera: Culicidae)
metamorphosis. J Med Entomol. 2001;38(1):29-32. Epub 2001/03/28. doi: 10.1603/0022-2585-
38.1.29. PubMed PMID: 11268687.

47.   Rodgers FH, Gendrin M, Wyer CAS, Christophides GK. Microbiota-induced peritrophic
matrix regulates midgut homeostasis and prevents systemic infection of malaria vector
mosquitoes. PLOS Pathogens. 2017;13(5):e1006391. doi: 10.1371/journal.ppat.1006391.

48.   Song X, Wang M, Dong L, Zhu H, Wang J. PGRP-LD mediates A. stephensi vector
competency by regulating homeostasis of microbiota-induced peritrophic matrix synthesis. PLoS
pathogens. 2018;14(2):e1006899-e. doi: 10.1371/journal.ppat.1006899. PubMed PMID:
29489896.

49.   Gao L, Song X, Wang J. Gut microbiota is essential in PGRP-LA regulated immune
protection against Plasmodium berghei infection. Parasites & vectors. 2020;13(1):3-. doi:
10.1186/s13071-019-3876-y. PubMed PMID: 31907025.

50.   Minard G, Tran F-H, Tran Van V, Fournier C, Potier P, Roiz D, et al. Shared larval
rearing environment, sex, female size and genetic diversity shape Ae. albopictus bacterial
microbiota. PLOS ONE. 2018;13(4):e0194521. doi: 10.1371/journal.pone.0194521.

51.   Short SM, Mongodin EF, MacLeod HJ, Talyuli OAC, Dimopoulos G. Amino acid
metabolic signaling influences Aedes aegypti midgut microbiome variability. PLoS Negl Trop Dis.
2017;11(7):e0005677. Epub 2017/07/29. doi: 10.1371/journal.pntd.0005677. PubMed PMID:
28753661; PubMed Central PMCID: PMCPMC5549995.

52.   Boissière A, Tchioffo MT, Bachar D, Abate L, Marie A, Nsango SE, et al. Midgut
microbiota of the malaria mosquito vector Anopheles gambiae and interactions with Plasmodium
falciparum infection. PLoS pathogens. 2012;8(5):e1002742-e. Epub 2012/05/31. doi:
10.1371/journal.ppat.1002742. PubMed PMID: 22693451.

53.   Hegde S, Khanipov K, Albayrak L, Golovko G, Pimenova M, Saldaña MA, et al.
Microbiome Interaction Networks and Community Structure From Laboratory-Reared and Field-
Collected Aedes aegypti, Aedes albopictus, and Culex quinquefasciatus Mosquito Vectors.
Frontiers in microbiology. 2018;9:2160-. doi: 10.3389/fmicb.2018.02160. PubMed PMID:
30250462.

54.   Villegas LM, Pimenta PF. Metagenomics, paratransgenesis and the Anopheles
microbiome: a portrait of the geographical distribution of the anopheline microbiota based on a
meta-analysis of reported taxa. Mem Inst Oswaldo Cruz. 2014;109(5):672-84. Epub 2014/09/04.
doi: 10.1590/0074-0276140194. PubMed PMID: 25185007; PubMed Central PMCID:
PMCPMC4156461.

55.   Dinparast Djadid N, Jazayeri H, Raz A, Favia G, Ricci I, Zakeri S. Identification of the
Midgut Microbiota of An. stephensi and An. maculipennis for Their Application as a
Paratransgenic Tool against Malaria. PLOS ONE. 2011;6(12):e28484. doi:
10.1371/journal.pone.0028484.

56.   Favia G, Ricci I, Damiani C, Raddadi N, Crotti E, Marzorati M, et al. Bacteria of the genus
Asaia stably associate with *Anopheles stephensi*, an Asian malarial mosquito vector.
2007;104(21):9047-51. doi: 10.1073/pnas.0610451104 %J Proceedings of the National
Academy of Sciences.

57.   Wilkerson RC, Strickman D, Litwak TR. Illustrated key to the female anopheline
mosquitoes of Central America and Mexico. Journal of the American Mosquito Control
Association. 1990;6(1):7-34. Epub 1990/03/01. PubMed PMID: 2324726.

58.   Choochote W, Saeung A. Systematic techniques for the recognition of *Anopheles*
species complexes. In: Manguin S, editor. *Anopheles* mosquitoes: IntechOpen; 2013.

59.   Beebe NW, Saul A. Discrimination of all members of the Anopheles punctulatus complex
by polymerase chain reaction--restriction fragment length polymorphism analysis. Am J Trop
Med Hyg. 1995;53(5):478-81. Epub 1995/11/01. PubMed PMID: 7485705.

60.   MR4. Amplification of the second internal transcribed spacer region (ITS2) in
Anophelines. Methods in *Anopheles* research: BEI Resources; 2015.

61.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2.
Nature Biotechnology. 2019. doi: 10.1038/s41587-019-0209-9.

62.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2:
High-resolution sample inference from Illumina amplicon data. Nature Methods. 2016;13:581.
doi: 10.1038/nmeth.3869

https://www.nature.com/articles/nmeth.3869#supplementary-information.

63.   Crawley MJ. The R book: John Wiley & Sons; 2012.

64.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. PLOS ONE. 2013;8(4):e61217. doi:
10.1371/journal.pone.0061217.

65.   Zhang H, editor The optimality of Naive Bayes. FLAIRS Conference; 2004.

66.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing
taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier
plugin. Microbiome. 2018;6(1):90. doi: 10.1186/s40168-018-0470-z.

67.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools. Nucleic
Acids Research. 2013;41(D1):D590-D6. doi: 10.1093/nar/gks1219.

68.   Mandal S, Van Treuren W, White RA, Eggesbo M, Knight R, Peddada SD. Analysis of
composition of microbiomes: a novel method for studying microbial composition. Microb Ecol
Health Dis. 2015;26:27663. Epub 2015/06/02. doi: 10.3402/mehd.v26.27663. PubMed PMID:
26028277; PubMed Central PMCID: PMCPMC4450248.

69.   team I. Inkscape: http://www.inkscape.org/ 2004.

Supporting Information

Suppl. 1. Shannon alpha diversity plots

Suppl. 2. Frequency and relative abundance of microbial taxa detected in internal and cuticle

surface microbiota of F_1 laboratory colonized *An. albimanus* larvae and adults

Suppl. 3. Summary statistics of sequencing reads and amplicon sequence variants (ASVs) used

for downstream analysis

Suppl. 4. Rarefaction depth and plots of sequencing reads

Figure 1: Map showing the geographical origins of F_1 *Anopheles albimanus*. F_1 laboratory-colonized mosquitoes were derived from gravid and/or blood fed females collected from each location. Mosquito icons show the geographical origin of the F_1 life stages studied.

Figure 2. Principal coordinate analysis (PCoA) ordinations of internal and cuticle surface microbiota from F$_1$ laboratory-colonized *An. albimanus*. The PCoA plots are based on Bray-Curtis distances between the microbiota of mosquitoes with differing maternal collection sites. Each point on the plot represents the microbial composition of a pool of three individuals, and mosquito pools are color-coded by their origin/maternal location. For larvae, the first two principal component (PC) axes captured 50% (internal) and 48% (cuticle surface) of the

variance in the data, with both internal and cuticle surface microbiota clustering distinctly by maternal collection site. For adults, the first two PC axes captured 59% (internal) and 47% (cuticle surface) of the variance in the data, with cuticle surface but not internal microbiota clustering distinctly by maternal collection site. PERMANOVA statistics are presented at the bottom of each plot.

Figure 3. Number of unique and shared microbial taxa between microbial niches (A); number of unique and shared microbial taxa between maternal collection sites (B). The number of taxa shown in the Venn diagram represent bacterial taxa except in the cuticle surface microbiota of adults from Las Cruces 4, where two archaeal taxa were present. n = pools of mosquito samples analyzed per location or microbial.

a.

b.

Figure 4. Frequency of ASVs from the internal (a) and cuticle surface (b) microbiota of laboratory-colonized *An. albimanus* F$_1$ larvae originating from different maternal collection sites. ASVs were annotated to the genus level or the lowest possible taxonomic level (in square brackets) and are clustered by the average nearest-neighbors chain algorithm. Only taxonomically annotated ASVs with frequencies ≥2000 are presented.

LARVAL INTERNAL MICROBIOTA

Significantly different taxa	W	ElTerrero	LasCruces3	LasCruces4
Singulisphaera	76		✔	
[Planctomycetaceae]	76		✔	
Paucibacter	76		✔	
Azoarcus	76	✔		
Azospirillum	72		✔	✔
Ancylobacter	64		✔	
Gemmobacter	59		✔	
Terrimicrobium	59	✔	✔	
Legionella	54	✔	✔	
[Oligoflexales]	55		✔	✔
Reyranella	53		✔	
[Acetobacteraceae]	50		✔	
Mycobacterium	47		✔	
[Saccharibacteria]	38		✔	✔
Bosea	31		✔	✔
Microbacterium	31		✔	✔

LARVAL CUTICLE SURFACE MICROBIOTA

Significantly different taxa	W	ElTerrero	LasCruces3	LasCruces4
Azoarcus	71	✔		
Paucibacter	68	✔	✔	
[Planctomycetaceae]	67		✔	
Ancylobacter	64		✔	
Microbacterium	41		✔	✔
Spirosoma	40			✔
Bdellovibrio	36		✔	✔
[Oligoflexales]	27			✔
Pelomonas	25		✔	✔
Acidovorax	21		✔	✔

Figure 5. Volcano plots of differentially abundant bacterial taxa in F₁ laboratory colonized *An. albimanus* larvae originating from three different maternal collection sites. The plots show results of analysis of composition of microbiomes (ANCOM) tests for differentially abundant microbial taxa between maternal collection site, with an effect size set to log F≥20,

and a cut-off of differential abundance set to W≥20 (i.e. a taxon was differentially abundant across maternal collection sites if the ratio of its abundance to those of at least 20 other taxa (25% of all included taxa) differed significantly across sites). Differentially abundant taxa are highlighted (blue shaded area) and the taxa names and locations in which they were most abundant are presented in the adjoining tables.

INTERNAL

CUTICLE SURFACE

Figure 6. Frequency of ASVs from the internal and cuticle surface microbiota of laboratory-colonized F₁ adult *An. albimanus* originating from different locations. ASVs were annotated to the genus level or the lowest possible taxonomic level (in square brackets) and are clustered by the average nearest-neighbors chain algorithm. Only taxonomically annotated ASVs with frequencies ≥1000 are presented.

Table 1. Descriptive statistics of laboratory colonized F_1 *An. albimanus* processed, per life stage and maternal collection site.

Maternal collection site	Number of mosquitoes (number of pools)	
	L3-L4 larvae	Adult ♀s
El Terrero	45 (15)	-
Las Cruces 1	-	27 (9)
Las Cruces 3	42 (14)	45 (15)
Las Cruces 4	45 (15)	63 (21)

Table 2. Pairwise alpha and beta diversity comparisons of laboratory colonized F₁ *An. albimanus* microbiota from different maternal collection sites. a. Pairwise beta (Bray-Curtis) diversity comparison showed significant differences in larval internal and cuticle surface microbiota between maternal collection sites. In contrast, only adult cuticle surface but not internal microbiota were significantly different across maternal collection sites. **b.** Pairwise alpha (Shannon) diversity comparison showed significant differences in larval internal but not cuticle surface microbiota between maternal collection sites (two of the three pairs). In contrast, there was no significant difference in adult internal or cuticle surface microbiota between maternal collection sites. Pairwise alpha and beta diversity comparisons were conducted using Kruskal-Wallis and PERMANOVA (999 permutations) tests respectively, with Benjamini-Hochberg FDR correction (q-value). Significance was determined at q < 0.05.

a.

	Group 1	Group 2	Internal			Cuticle surface		
			pseudo-F	p-value	q-value	pseudo-F	p-value	q-value
Adults	LasCruces1	LasCruces3	2.21	0.088	0.161	**5.80**	**0.001**	**0.003**
	LasCruces1	LasCruces4	1.92	0.107	0.161	**4.97**	**0.004**	**0.005**
	LasCruces3	LasCruces4	0.85	0.443	0.443	**3.07**	**0.005**	**0.005**
Larvae	ElTerrero	LasCruces3	**9.81**	**0.001**	**0.001**	4.48	0.001	0.001
	ElTerrero	LasCruces4	**7.98**	**0.001**	**0.001**	11.43	0.001	0.001
	LasCruces3	LasCruces4	**10.88**	**0.001**	**0.001**	11.37	0.001	0.001

b.

	Group 1	Group 2	Internal			Cuticle surface		
			H	p-value	q-value	H	p-value	q-value
Adults	LasCruces1	LasCruces3	0.00	0.98	0.98	1.09	0.30	0.41
	LasCruces1	LasCruces4	0.49	0.48	0.72	1.23	0.27	0.41
	LasCruces3	LasCruces4	1.77	0.18	0.55	0.67	0.41	0.41
Larvae	ElTerrero	LasCruces3	**6.63**	**0.01**	**0.02**	4.45	0.03	0.10
	ElTerrero	LasCruces4	**6.94**	**0.01**	**0.02**	1.87	0.17	0.26
	LasCruces3	LasCruces4	0.49	0.48	0.48	1.10	0.29	0.29