Title: Motor Protein MYO1C is Critical for Photoreceptor Opsin Trafficking and Vision

Authors: Ashish K. Solanki[1], Elisabeth Obert[2], Stephen Walterhouse[1], Ehtesham Arif[1], Bushra Rahman[1], Joshua H. Lipschutz[1,3], Jeffery Sundstrom[4], Barbel Rohrer[2,3], Shahid Husain[2], Glenn P. Lobo[1,2]* and Deepak Nihalani[1]*

Affiliations: [1]Department of Medicine, Medical University of South Carolina, Charleston, SC, 29425, USA. [2]Department of Ophthalmology, Medical University of South Carolina, Charleston, SC, 29425, USA. [3]Ralph H. Johnson VA Medical Center, Division of Research, Charleston, SC 29420, USA. [4]Penn State Eye Center, 200 Campus Dr., Suite 800, Hershey, PA 17033. USA.

*Corresponding Authors
*Glenn P. Lobo, Ph.D.
Assistant Professor of Medicine
Department of Medicine
70 President Road
Drug Discovery Building DDB513
Charleston, SC 29425
Office: 843-876-2371
Fax: 843-792-8399
Email: lobo@musc.edu

*Deepak Nihalani, Ph.D.
Associate Professor of Medicine
Department of Medicine
70 President Road
Drug Discovery Building DDB514
Charleston, SC 29425
Office: 843-792-7659
Fax: 843-792-8399
Email: nihalani@musc.edu

Keywords: Motor protein, Myosin, MYO1C, Opsins, Photoreceptor, Outer Segments, Photoreceptor Cell Degeneration, Trafficking, Vision.

Abstract

Unconventional myosins linked to deafness are also proposed to play a role in retinal physiology. However, their direct role in photoreceptor structure and function remains unclear. Herein, we demonstrate that systemic loss of the unconventional myosin MYO1C in mice affected opsin trafficking and photoreceptor survival, leading to vision loss. Electroretinogram analysis of *Myo1c* knockout (*Myo1c*-KO) mice showed a progressive loss of rod and cone function. Immunohistochemistry and binding assays demonstrated MYO1C localization to photoreceptor inner and outer segments (OS) and identified a direct interaction of rhodopsin with the MYO1C cargo domain. In *Myo1c*-KO retinas, rhodopsin mislocalized to rod inner segments and the cell bodies, while cone opsins in OS showed punctate staining. In aged mice, the histological and ultrastructural examination of *Myo1c*-KO retinas showed a progressively degenerating photoreceptor OS phenotype. These results demonstrate that MYO1C is critical for opsin trafficking to the photoreceptor OS and normal visual function.

Introduction

Protein trafficking within the photoreceptors must occur efficiently and at high fidelity for photoreception, structural maintenance and overall retinal cell homeostasis. Additionally, it is well known that proper opsin trafficking is tightly coupled to photoreceptor cell survival and function [1-9]. However, the cellular events, including signaling pathways and trafficking events that participate in retinal injuries due to improper protein trafficking to the photoreceptor outer segments (OS), are not yet fully understood. While many proteins are known to play an essential role in retinal cell development and function, the involvement of motor proteins in eye biology is less understood. Identification of genetic mutations in the *Myo7a* gene associated with retinal degeneration in Usher syndrome suggests that unconventional myosins play a critical role in retinal pigmented epithelium (RPE) and photoreceptor cell function [10, 11]. Unconventional myosins are motor proteins that are proposed to transport membranous organelles along the actin filaments in an ATP-dependent manner, and additional roles are currently being uncovered [12-14]. The loss of *Myo7a* primarily affects RPE and OS phagocytosis leading to retinal cell degeneration. It is, however, believed that other, yet unidentified class I myosins, may participate more directly in photoreceptor function. Here we present compelling evidence for another unconventional actin-based motor protein, MYO1C, with its primary localization to photoreceptors, that plays a role in retinal cell structure and function via opsins trafficking to the photoreceptor OS.

Rhodopsin and cone pigments in photoreceptor OS mediate scotopic and photopic vision, respectively. They consist of opsin, the apo-protein, and 11-*cis* retinal, a

vitamin A-based chromophore. Light absorption by 11-*cis* retinal triggers a

conformational change in opsin [8]. This, in turn, initiates a G protein-coupled receptor

(GPCR) signaling pathway/phototransduction cascade, signaling the presence of light.

Thus, the presence of opsins in the photoreceptor OS is critical for vision [1-9]. Each

photoreceptor cell contains an OS housing the phototransduction machinery, an inner

segment (IS) where proteins are biosynthesized and energy is being produced, and a

synaptic terminal for signal transmission. One of the most fundamental steps in vision is

the proper assembly of signal-transducing membranes, including the transport and

sorting of protein components. A major cause of neurodegenerative and other inherited

retinal disorders is the improper localization of proteins. Mislocalization of the dim-light

photoreceptor protein rhodopsin is a phenotype observed in many forms of the blinding

disease, including retinitis pigmentosa. The proteins that participate in

phototransduction (including rhodopsin, transducin, phosphodiesterase [PDE6], or the

cyclic nucleotide gated channels [CNG], are synthesized in the IS and must be

transported through the connecting cilium (CC) to the OS. These proteins are either

transmembrane (TM) proteins or peripherally associated membrane proteins that are

attached to the membrane surface [1-9]. How the TM proteins (e.g., rhodopsin and

CNG) and peripherally associated proteins (e.g., transducin and PDE6) traffic through

the IS to incorporate eventually in the nascent disc membrane or the photoreceptor

outer membrane is not fully understood and constitutes an area of intense research, as

improper trafficking of these protein causes retinal cell degeneration and can lead to

blindness [1-9].

Genetic mutations in myosins that lead to hearing loss have also been

associated with retinal degeneration. Some of the essential genes known to be involved

in either or both of these functions belong to a family of unconventional motor proteins

and include MYO3A [15], MYO7A , MYO6 , MYO15 [16], and MYO5. A recent report

identified another unconventional myosin associated with deafness, MYO1C, where

mutations which affects its nucleotide-binding pocket and calcium binding ability [17].

Importantly, MYO1C was identified in the proteomic analysis of the retina and vitreous

fluid as part of a protein hub involved in oxidative stress [18-20]. MYO1C is an actin-

binding motor protein that is widely expressed in multiple cell types. It has been shown

to participate in a variety of cellular functions including protein trafficking and

translocation [12, 21-23]. As MYO1C has low tissue specificity based on mRNA and

protein expression, it remains unclear which cell type is most dependent on MYO1C

trafficking function and is affected by the loss of MYO1C.

In this study, we aimed to systematically analyze the function of the motor protein

MYO1C in protein trafficking in photoreceptors. Here, we find that a global genetic

deletion of *Myo1c* results in progressive mistrafficking of opsins to the OS. Using retinal

lysate from wild-type (WT) mice in co-immunoprecipitation assays, we show that

MYO1C and rhodopsin directly interact, indicating that opsin is a cargo for MYO1C.

Loss of MYO1C promoted a progressive OS degeneration concomitant with a reduction

in rod and cone photoreceptor cell function, suggesting that MYO1C is critical for retinal

structure and function. Our findings have significant clinical implications for

degenerative rod and cone diseases, as mutations in MYO1C or its interacting partners

could exert its effects on retinal health and vision by altering opsin trafficking to the

photoreceptor OS, a fundamental step for maintaining vision in humans.

Results

Construction and Validation of *Myo1c* null Mice*: We have previously generated

Myo1c floxed mice using the standard knockout strategy [24]. Systemic deletion of

Myo1c was achieved by crossing *Myo1c* floxed (*Myo1cfl/fl*) mice with Actin Cre+

(ActCre+) mice to generate *Myo1cfl/fl ActCre+/-* knockout mice (referred to as *Myo1c-*

KO mice in this manuscript). Western blotting of protein lysates from various tissues

including kidney, heart, and liver of *Myo1c*-KO mice showed complete loss of MYO1C,

thus confirming the systemic deletion of *Myo1c* (**Fig. S1a**). Additionally, immunostaining

analysis of these tissues further confirmed loss of MYO1C protein in *Myo1c*-KO mice

(**Figs. S2a-d**).

Genetic Deletion of *Myo1c* induces Visual Impairment in Mice:

Immunofluorescence analysis showed that MYO1C was enriched in the OS and IS of

WT mice and absent in *Myo1c*-KO retinas (**Fig. 1a**). Since mutations or deletion of the

motor protein MYO7A has been associated with retinal degeneration in Usher syndrome

or its animal model respectively, it prompted us to investigate whether *Myo1c* deletion

affects retinal function. Using electroretinograms (ERGs) [25, 26], we tested

photoreceptor cell function of *Myo1c*-KO and WT mice (*n*=8, per genotype and age-

group) under dark-adapted, scotopic and light-adapted, photopic conditions to

determine rod and cone function, respectively. In contrast to WT animals, we observed

reduced ERGs for *Myo1c*-KO mice at different ages. Two months old *Myo1c*-KO mice

showed significant reduction in the **a**-wave but not **b**-wave amplitudes ($P<0.0068$

and $P<0.098$, respectively) (**Figs. 1b-c**). Strikingly, ERG analysis of adult six months old

Myo1c-KO mice showed severe loss of retinal function, where a significant reduction in

both **a**- and **b**-waves was observed (38-45% of WT animals ($P<0.005$)) (**Figs. 1d-e**).

These analyses suggest the absence of key phototransduction components in the

photoreceptor OS of *Myo1c*-KO mice and/or photoreceptor OS degeneration, as

contributing factors.

Rod and Cone Visual Pigments in *Myo1c*-KO Mice: Since, the phototransduction

protein, rhodopsin, constitutes 85-90% of photoreceptor OS protein content [27], and as

the ERG responses were impaired in *Myo1c*-KO mice, we hypothesized that the loss of

MYO1C may affect opsin trafficking to the photoreceptor OS. To test this hypothesis, we

analyzed retinal sections from WT and *Myo1c*-KO mice (at 2 and 6 months of age; *n*=8

per genotype), probing for rhodopsin, two types of cone opsins (red/green (medium

wavelength, M-opsin and S-opsin (short wavelength)), rod-specific phosphodiesterase

6b (Pde6b) and a general cone marker, PNA lectin. In WT mice, rhodopsin localized

exclusively to the rod outer segments at 2 and 6 months of age. In contrast, in two

months old *Myo1c*-KO mouse retinas, while most rhodopsin trafficked to the OS, some

also mislocalized to the base of the rod IS and the cell bodies in the outer nuclear layer

(ONL), suggesting incomplete target transport/trafficking to photoreceptor OS in the

absence of MYO1C (**Fig. 2a;** white arrows; rhodopsin levels within individual retinal

layers are quantified and shown in **Fig. S3**). An even more severe mislocalization of

rhodopsin to the rod IS, and within the ONL was observed in the 6 months old *Myo1c*-

KO mice, suggesting a progressive retinal phenotype in the absence of MYO1C (**Fig.**

2a; rhodopsin expression within individual retinal layers are quantified and shown in

Fig. S4). In contrast, staining for Pde6b, a lipidated rod specific protein that does traffic

to OS independently of rhodopsin [28], showed normal trafficking to the rod OS in both

WT and *Myo1c*-KO retinas (**Fig. 2d**). Staining for the two cone opsins suggests that the

cone outer segments are shorter and swollen by 2 and increasingly so by 6 months of

age (**Figs. 2b-c**). This observation is supported by staining for PNA lectin, documenting

progressively shorter and fatter cone OS, indicative that cone OS degenerate as these

mice age in the absence of MYO1C (**Fig. 2e**). Cone visual arrestin in WT mice typically

outlines the entire cell, OS, IS, cell body, axon and cone pedicle. Staining for cone

arrestin in *Myo1c*-KO animals confirmed the short and swollen appearance of the cone

OS (**Fig. 3a,** red and white arrows, respectively). Although not apparent by

immunohistochemistry in mouse RPE, human RPE cells express functional MYO1C

[29], and low levels of *Myo1C* mRNA appear present in mouse RPE [30]. Since

elimination of the motor protein *Myo7a* in mouse leads to alterations in protein

localization in the RPE (RPE65) [31], we stained retinas of young and adult WT and

Myo1c-KO mice with anti-STRA6 antibody, another RPE-specific protein. This analysis

suggests that STRA6 expression and localization in the RPE was not affected in the

absence of MYO1C (**Fig. 4b**). Thus, since MYO1C is known primarily as a motor with

protein trafficking function [14, 18], its absence in photoreceptors of *Myo1c*-KO animals

may contribute specifically to the loss of opsin trafficking to the OS.

Myo1c-KO Mice demonstrate Photoreceptor Outer Segment Loss: To further evaluate if opsin mistrafficking is also associated with structural changes to the retina, the histological and TEM analysis of retinal sections of young and adult WT and _Myo1c_-KO mice was performed. In histological sections of H&E stained tissues, progressive shortening of rod photoreceptor OS was observed, where the OS of adult _Myo1c_-KO (6 months of age) mice appeared shorter than the younger _Myo1c_-KO mice (2 months of age), which in turn were shorter than those in WT mice (**Fig. 4a**). Also, in comparison to the WT mice, the photoreceptors in _Myo1c_-KO mice appeared to be less organized, especially in the older mice (**Fig. 4a**), suggesting that loss of MYO1C may affect progressively affect photoreceptor homeostasis. To further evaluate the structure of rod photoreceptors, ultrastructural analysis using TEM was performed. While the rod photoreceptor OS in the WT mice showed normal elongated morphology, they appeared shorter and wider in _Myo1c_-KO mice at two months of age (**Fig. 4b**). Specifically, comparing _Myo1c_-KO with WT mouse rod outer segment lengths at six months of age demonstrated that OS segment lengths in _Myo1c_ retinas were 35-48% shorter than those of WT mice ($P<0.005$; **Fig. 4c**). TEM analysis of _Myo1c_-KO mice retinas at six months of age for rod structure showed that they were significantly shorter in lengths ($P<0.05$; **Fig. 4d _vs_ 4e;** rod lengths quantified in **Fig. 4f**, at two months and **Fig. 4g**, at 6 months). Ultra-structurally, the cone outer segments in the _Myo1c_-KO mouse retina appeared shorter and had lost their typical cone shape, confirming the swollen shape identified by immunohistochemistry. Retinal lamination and nuclear cell layers in _Myo1c_-KO mice however, appeared normal (**Fig. 4a**). These results suggest

that the lack of MYO1C resulted in progressively severe opsin mislocalization (**Figs. 2**

and 3), shorter photoreceptor OS (**Fig. 4**), supporting the observed decrease in visual

function by ERG (**Figs. 1b-e**).

MYO1C Cargo Domain directly Interacts with Rhodopsin: Since the loss of MYO1C

resulted in retinal function defects with significant alterations in the localization of

opsins, we evaluated whether MYO1C exerts this effect through a physical interaction

with rhodopsin. Immunoprecipation analysis using WT and *Myo1c*-KO (n = 6 retinas

pooled per genotype) mice demonstrated that rhodopsin was pulled down using

MYO1C antibody and in a reciprocal fashion (**Figs. 5a** and **5b**). Using a baculovirus

produced purified recombinant mouse MYO1C protein, we further demonstrate that

MYO1C directly interacted with rhodopsin in an overlay assay, where opsin

immunoprecipitated either from mouse retinal lysate or HEK293 cells overexpressing

Myo1c, and separated on an SDS-PAGE gel and then probed with purified recombinant

full-length MYO1C (MYO1C FL) or GFP-MYO1C-790-1028 (tail domain, also known as

the cargo domain) [13]. The immunoblot analysis of the over-layered MYO1C showed

significant binding of both MYO1C proteins, full-length and tail, at the rhodopsin band

indicating a direct interaction between the two proteins (**Fig. 5**). Interestingly, the

interaction of MYO1C was noted with various multimers of rhodopsin, which further

indicates that opsin is a cargo for MYO1C (**Fig. 5**).

Genetic Deletion of *Myo1c* does not affect Systemic Organs in Mice: Finally, to

determine if the systemic deletion of *Myo1c* affects other organs, we harvested major

systemic organs, including liver, heart, and kidney of 6 months old *Myo1c*-KO and WT

mice (*n*=5 per genotype), and performed histology and H&E staining. Notably, both

young and adult *Myo1c*-KO mice developed and reproduced normally with no

observable differences between the control and *Myo1c*-KO genotypes (**Figs. S5a-c**).

Thus, except for the retinal phenotype, *Myo1c*-KO animals showed unremarkable

pathology of systemic organs.

Discussion:

Our work identifies for the first time an unconventional motor protein, MYO1C, as a

novel trafficking regulator of both rod and cone opsins to the photoreceptor OS in mice.

In this study, based on MYO1C localization within the IS and OS of photoreceptors, and

using a whole-body *Myo1c*-KO mouse model, we have functionally identified MYO1C as

a novel component of retinal physiology, where it was specifically found to be involved

in photoreceptor cell function. Retinal analysis of *Myo1c*-KO mice identified opsins as

novel cargo for MYO1C. In the absence of MYO1C, both young and adult *Myo1c*-KO

mice showed impaired opsin trafficking, where rhodopsin was retained in the

photoreceptor IS and the cell bodies. In contrast, cone opsins showed no retention in

the cell body or mistrafficking of other membranes, but staining pattern revealed

misformed cone OS. These two phenotypes manifested in a progressive decline of

visual responses in both the rod and cone ERGs and shorter photoreceptor OS as

Myo1c-KO animals aged, indicating a progressive phenotype. Significantly, only a

retinal phenotype was observed, as systemic organs of *Myo1c*-KO mice, including

heart, liver, and kidney, appeared unaffected. Our data thus points to a novel

mechanism by which MYO1C regulates opsins trafficking from the photoreceptor IS to

OS, which is a critical event for photoreceptors' function and long-term photoreceptor

cell survival. Our study identifies an unconventional motor protein MYO1C, as an

essential component of the mammalian photoreceptors, which plays a critical role in

promoting phototransduction and long-term photoreceptor cell survival.

MYO1C is Involved in Photoreceptors function via the Transport of opsins from

Inner segments to Outer Segments: Photoreceptor degeneration has devastating

consequences on visual function. Therefore, protein trafficking, in particular appropriate

trafficking of both rod and cone opsins within the photoreceptors must occur efficiently

and at high fidelity for normal photoreception, proper structural maintenance, and for

overall retinal cellular homeostasis. However, some of the molecular players remain

unknown, and therefore is an area of intense research. Herein, the identification of

mechanisms that contribute to opsin trafficking, and consequently to the integrity of

photoreceptors may help design protective/therapeutic strategies to restore

photoreceptor function in humans with inherited retinal degenerative diseases. The

observation that MYO1C is involved in opsin trafficking suggest that this protein could

be explored/targeted as a therapeutic modality to mitigate retinal degenerative

diseases, especially in patients with defective opsin trafficking.

MYO1C and other Opsin Trafficking Proteins: *Myo1c*-KO mice exhibited rhodopsin

mislocalization similar to *Rpgr*$^{-/-}$, *Myo7a*^{Sh1}, *Rp1*$^{-/-}$, *Kinesin II*$^{-/-}$, and *Tulp1*$^{-/-}$ mutant

mice [1-9]. And just like in these mutant mice, the *Myo1c*-KO mice showed opsin

mislocalization and shorter OS with little or no photoreceptor cell death up to the 6-

month time point of analysis. Since MYO1C primarily localized to photoreceptor IS and

OS, is known to be involved in protein trafficking and uses actin as a track [14, 18], we

hypothesized that MYO1C participates in the movement of opsins from IS to the OS of

photoreceptors. This hypothesis was supported by the observation that the rod opsins

were mislocalized to IS and cell bodies and defective assembly of cone OS in *Myo1c*-

KO mice suggests that this phenotype is caused by an aberrant protein transport and

that OS degeneration was likely a secondary event. Apparently, the normal

ultrastructure of photoreceptors in our *Myo1c*-KO mice suggests that the retinal

abnormalities in these animals are not due to structural defects in photoreceptors per

se, but instead are induced by an aberrant motor function leading to opsin

mislocalization.

MYO1C contributes to Phototransduction and Retinal Homeostasis: The opsin

molecules and other phototransduction proteins are synthesized in the cell body of the

photoreceptor [32, 33]. They are then transported to the distal IS [34], and subsequently

to the OS. Little is known about these transport processes and the molecular

components involved in this process [1-9]. The localization of MYO1C in the

photoreceptors' IS and OS, suggests that opsins may utilize this molecular motor for

transport to the OS. The immunohistochemical analysis of *Myo1c*-KO animals indicated

that while rod and cone opsins trafficked to the OS, significant mislocalization was noted

for rhodopsin in the IS and cell bodies in the ONL (**Fig. 2**). Since they represent plasma

membrane structural proteins, cone opsins presumably contribute to the cone OS

stability and rhodopsin to the rod OS formation and stability [7], and hence

photoreceptor OS degeneration in *Myo1c*-KO mice may be attributed, in large part, to

the mistrafficking of opsins to the IS or to a progressive reduction of opsins in the OS

membrane. Notably, the pattern of opsin mislocalization observed in *Myo1c*-KO mice

resembled closely with the retinal phenotype observed in our previously reported *Tulp1*-

KO, [4, 35], *Cnga3*$^{-/-}$ mice [5], *Lrat*$^{-/-}$ and *Rpe65*$^{-/-}$ [3, 8, 9], GC1-KO mice [1, 6], and, to

some extent in CFH (complement factor H) KO animals [2]. Importantly, in all these

studies, photoreceptor OS was unstable, and significant degeneration was noted.

However, because 85-90% of OS protein is rhodopsin, the mislocalization of other less

abundant proteins cannot be ruled out in the photoreceptors of *Myo1c*-KO mice.

Contributions from other Motor proteins in Opsin trafficking: Though the study

demonstrates mistrafficking of opsins due to a loss of MYO1C, the majority of opsin was

still correctly localized, suggesting that contribution or compensation from other myosins

cannot be ruled out. Nevertheless, the contributions from MYO1C are highly significant,

as its genetic deletion showed specific physiological defects in mouse retinas. It is likely

that some redundancy exists among molecular motors and several known candidates

might compensate for the lack of MYO1C in photoreceptor function. However, the qPCR

analysis of the retinas from WT and *Myo1c-KO* mice did not suggest compensation from

other family myosin 1 members (**Fig. S6**). However, compensation by other motor

proteins, including the members of kinesin superfamily [36, 37], myosin VIIa, and

conventional myosin (myosin II) [38, 39], which have also been detected in the RPE and

retina, cannot be ruled out and may need further investigation. Overall, results

presented herein support a role for MYO1C in opsin trafficking in the photoreceptor cells

of the retina and provide evidence that defective protein transport pathways may be a

pathologic mechanism responsible for OS degeneration and loss of vision, in these

mice.

Methods:

Materials: All chemicals, unless stated otherwise, were purchased from Sigma-Aldrich

(St. Louis, MO, USA) and were of molecular or cell culture grade quality.

Myo1c-**Knockout (***Myo1c*-**KO) Mouse Model**: We have previously generated *Myo1c*

transgenic mice (*Myo1cfl/fl*) in C57BL/6N-derived embryonic stem cells, flanking exons

5 to 13 of the mouse *Myo1c* gene, which has allowed us to specifically delete all *Myo1c*

isoforms in a cell-specific manner [24]. Here, a complete *Myo1c*-knockout was

generated by crossing *Myo1cfl/fl* mice with an F-actin Cre mouse strain (B6N.FVB-

Tmem163Tg(ACTB-cre)2Mrt/CjDswJ) obtained from Jackson labs. We will refer to the

Myo1cfl/fl x f-actin Cre cross as *Myo1c* knockout (*Myo1c*-KO) mice. For this study, the

Myo1c-KO mice were crossed onto a C57BL/6J background to avoid potential problems

with the *Rd8* mutation (found in C57BL/6N lines) [40].

Immunohistochemistry and Fluorescence Imaging: Eyes were fixed in 4%

paraformaldehyde buffered with 1X PBS for 2 hours at 4^0C using established protocols

[58]. After fixation, samples were washed in PBS and embedded in paraffin and

processed (MUSC histology core facility). Sections (10 μm) were cut and transferred

onto frost-free slides. Slide edges were lined with a hydrophobic marker (PAP pen) and

washed with PBS before blocking for 1-2 hr at RT. Blocking solution (1% BSA, 5%

normal goat serum, 0.2% Triton-X-100, 0.1% Tween-20 in 1X PBS) was applied for 2

hours in a humidified chamber. Primary antibodies were diluted in blocking solution as

follows: anti-rhodopsin (1:500, Millipore/Sigma, St. Louis, MO), anti-Myo1c (1:100),

cone-arrestin (1:250, Millipore/Sigma, St. Louis, MO), conjugated PNA-488 and PNA-

594 (1:2000, Molecular Probes, Eugene, OR), anti-red/green cone opsin (1:500;

Millipore, St. Louis, MO), anti-S-opsin (1:500, Millipore/Sigma, St. Louis, MO), and 4′,6-

diamidino-2-phenylendole (DAPI; 1:5000) was used to label nuclei. All secondary

antibodies (Alexa 488 or Alexa 594) were used at 1:5000 concentrations (Molecular

Probes, Eugene, OR). Optical sections were obtained with a Leica SP8 confocal

microscope (Leica, Germany) and processed with the Leica Viewer software.

Measurement of Photoreceptor Outer Nuclear Layer (ONL) Thickness and Outer

Segment (OS) Lengths: All fluorescently labeled retinal sections on slides were

analyzed by the program BIOQUANTNOVA (R & M Biometrics, Nashville). The length

of the photoreceptor OS and thickness of the ONL were measured at five consecutive

points in a 100 μm segment located 300-400 μm from the optic nerve. The ONL

thickness was measured from the base of the nuclei to the outer limiting membrane at a

90° angle. The OS length was measured from the base of the OS to the inner side of

the retinal pigment epithelium. Only sections in which columns of rod nuclei were

apparent were counted to ensure that sections were not oblique. Measurements from

the four quadrants (see above) were averaged; the SD was less than 10% of the

average value for each mouse.

Electroretinography (ERG) Analysis: Dark-adapted WT and *Myo1c*-KO mice (50:50 ratio of male and female*; n*=8 each genotype) at two months of age (young mice; early time-point), and between 6-7 months of age (end time-point) were anesthetized by intraperitoneal injection of a ketamine/xylene anesthetic cocktail (100 mg/kg and 20 mg/kg, respectively) and their pupils were dilated with 1% tropicamide and 2.5% phenylephrine HCl. Both scotopic and photopic ERGs were recorded with a computerized system (UTASE-3000; LKC Technologies, Inc., Gaithersburg, MD, USA), as previously described [41-43].

Transmission Electron Microscopy (TEM): Eyecups at the indicated time-points were harvested and fixed overnight at 4^0C in a solution containing 2% paraformaldehyde/2.5% glutaraldehyde (buffered in 0.1M cacodylate buffer). Samples were then rinsed in the buffer (0.1 M cacodylate buffer). Post-fixative 2% OsO_4/0.2 M cacodylate buffer 1 hour at 4^0C, followed by 0.1 M cacodylate buffer wash. The samples were dehydrated through a graded ethanol series and then embedded in epon (EMbed 812; EM Sciences). The cured blocks were cut at 0.5 microns and stained with 1% toluidine blue to orient the blocks to the required specific cell types. The blocks were then trimmed to the precise size needed for ultrathin sectioning. The blocks were cut at 70 nm and gathered on 1-micron grids. The grids were air-dried and then stained with uranyl acetate (15 minutes) and lead citrate (5 minutes) and rinsed in-between each stain. They were then allowed to dry and imaged with a JEOL 1010. Images were acquired with a Hamamatsu camera and software. All samples were processed by the

Electron Microscopy Resource Laboratory at the Medical University of South Carolina,

as previously described [53,70].

Western Blot Analysis and Densitometry: Total protein from cells or mouse tissues

(*n*=3 per genotype) were extracted using the M-PER protein lysis buffer

(ThermoScientific, Beverly, MA) containing protease inhibitors (Roche, Indianapolis, IN).

Approximately 25 μg of total protein was then electrophoresed on 4-12% SDS-PAGE

gels and transferred to PVDF membranes. Membranes were probed with primary

antibodies against anti-Myo1c (1:250), β-Actin or Gapdh (1:10,000, Sigma), in antibody

buffer (0.2% Triton X-100, 2% BSA, 1X PBS) [54,55,70]. HRP conjugated secondary

antibodies (BioRad, Hercules, CA) were used at 1:10,000 dilution. Protein expression

was detected using a LI-COR Odyessy system, and relative intensities of each band

were quantified (densitometry) using Image J software version 1.49 and normalized to

their respective loading controls.

Co-immunoprecipitation Assays: Co-immunoprecipitation of endogenously

expressed proteins (MYO1C and rhodopsin) was performed using mouse retinal

extracts. Six retinas of each genotype (WT and *Myo1c*-KO) were used for extraction of

retinal proteins in 250 μL of RIPA buffer (phosphate-buffered saline [PBS] containing

0.1% sodium dodecyl sulfate [SDS], 1% Nonidet P-40, 0.5% sodium deoxycholate, and

100 mM potassium iodide) with EDTA-free proteinase inhibitor mixture (Roche

Molecular Biochemicals). Lysates were cleared by centrifugation at 10000 rpm for 10

min at 4°C. The prepared lysates were incubated with anti-Myo1c, anti-rhodopsin, and

mouse/rabbit IgG overnight at 4°C and further with protein G-coupled agarose beads

(ROCHE) for 1-2 h. Beads were then collected by centrifugation at 3000 rpm for 5 min

at 4°C, extensively washed in 1X PBS, and resuspended in SDS gel loading buffer. The

proteins were separated on a 10% SDS-PAGE, transferred to a PVDF membrane, and

analyzed by immunoblotting with the corresponding antibodies.

Overlay direct binding assay: Rhodopsin protein was expressed in HEK293 cells

using transient transfection (pcDNA3 rod opsin construct, a gift from Robert Lucas

(Addgene plasmid # 109361, http://n2t.net/addgene:109361; RRID:Addgene_109361)

[44] and immunoprecipitated from the cell lysates using an anti-rhodopsin antibody (

Abcam). In parallel, rhodopsin was similarly immunoprecipitated from the mouse retinal

lysate. The immunoprecipitated complexes were separated on SDS-PAGE gel and

transferred to PVDF membrane. The membrane was then probed by overlaying it with 5

μg of baculovirus-produced and purified recombinant full-length MYO1C FL or GFP-

MYO1C-790-1028 (tail domain, also known as the cargo domain [13]) protein and

incubating at 4^0C for 4 h. Following incubation, the membranes were western blotted

with MYO1C antibody to detect the direct binding of MYO1C to the rhodopsin bands.

The location of rhodopsin on the membranes was marked by separately probing these

membranes with an anti-rhodopsin (1:500, Millipore Sigma) antibody.

Quantitative Real Time-PCR: RNA was isolated from retinas of WT and *Myo1c*-KO

animals using Trizol reagent, and processed as described previously [55,70]. One

microgram of total RNA was reverse transcribed using the SuperScript II cDNA

Synthesis Kit (Invitrogen, Eugene, OR). Quantitative Real-Time PCR (qRT-PCR) was

carried out using SYBR green 1 chemistry (BioRad, Hercules, CA). Samples for qRT-

PCR experiments were assayed in triplicate using the BioRad CFX96 Q-PCR machine.

Each experiment was repeated twice (n=6 reactions for each gene), using newly

synthesized cDNA.

Statistical Analysis: Data are expressed as means ± standard deviation by ANOVA in

the Statistica 12 software (StatSoft Inc., Tulsa, Oklahoma, USA). Differences between

means were assessed by Tukey's honestly significant difference (HSD) test. P-values

below 0.05 (P<0.05) were considered statistically significant. For western blot analysis,

relative intensities of each band were quantified (densitometry) using the Image J

software version 1.49 and normalized to the loading control β-Actin. QRT-PCR analysis

was normalized to 18S RNA, and the ΔΔCt method was employed to calculate fold

changes. Q-RTPCR data are expressed as mean ± standard error of mean (SEM).

Statistical analysis was carried out using PRISM 8 software-GraphPad.

DATA AVAILABILITY

The authors declare that all data supporting the finding of this study are available within

this article and its supplementary information files or from the corresponding author

upon request. The plasmids will be available from the corresponding author upon

reasonable request.

Disclosure

All the authors declared no competing interests.

Acknowledgements

This work was supported by grants from the NIH/NEI R21EY025034, R01EY030889,

Medical University of South Carolina (MUSC) start-up funds, and a Dialysis Clinic, Inc.

(award DCI-019898-001) research grant to G.P.L. 2R01DK087956-06A1, R56-

DK116887-01A1 and 1R03TR003038-01 to D.N. VA (Merit Award I01 BX000820), NIH

(P30DK074038), Dialysis Clinic, Inc. (research grant), and the American Heart

Association (AWRP Winter 2017 Collaborative Sciences Award) to J.H.L. This project

was also supported in part by the South Carolina Clinical and Translational Research

(SCTR) Institute at the Medical University of South Carolina, NIH/NCATS grant number

5UL1TR001450 to G.P.L.

Author Contributions

G.P.L., and D.N., designed the research studies. G.P.L., A.K.S., B. Rohrer., S.H., J.S.,

J.H.L., and D.N. wrote and edited the manuscript. G.P.L., A.K.S., E.O., D.N., E.A., S.W.,

S.H., and B.R. conducted experiments and acquired data. A.K.S., E.O., G.P.L., E.A., B.

Rohrer., S.H., J.S., and D.N., analyzed and interpreted the data. B. Rohrer., J.S., S.H.,

and J.H.L., supplied reagents and provided equipment for data analysis. All authors

have read and agreed to the published version of the manuscript.

 Figure Legends

 Fig. 1: MYO1C localizes to photoreceptors in mouse retina and its genetic
deletion induces vision defects: (**a**) Retina from adult wild-type (WT) and *Myo1c*-KO mice were harvested and immunostained with an anti-MYO1C antibody followed by secondary (Alexa 488) antibody staining. Dark-adapted scotopic ERGs were recorded in response to increasing light intensities in cohorts of two-month-old control (Wild-type/WT (white box)) and *Myo1c*-KO (black box) mice (**b, c**) and six months old (**d, e**) mice. Two-month-old *Myo1c* knock-out mice had lower dark-adapted *a*- and *b*-wave amplitudes compared with controls (post-hoc ANOVA: *a*-waves, $p < 0.0068$; *b*-waves, $p < 0.0098$), in particular at higher light intensities (20, 10, 6, 0 db). Six-month-old *Myo1c* knock-out mice had lower dark-adapted *a*- and *b*-wave amplitudes compared with controls (post-hoc ANOVA: *a*-waves, $p < 0.005$; *b*-waves, $p < 0.005$), in particular at higher light intensities (20, 10, 6, 0 db). Photoreceptor cell responses (*a*-waves), which drive the *b*-waves, were equally affected in *Myo1c*-KO animals (both reduced on average between 38-45% to WT animals). Data are expressed as mean ± S.E. (*Myo1c*-KO mice, $n = 8$; WT mice, $n = 8$). RPE, retinal pigmented epithelium; OS, outer segments; IS, inner segments; ONL, outer nuclear layer.

Fig. 2: Immunohistochemical analysis of wild-type/WT and *Myo1c* knock-out mice retina shows opsin trafficking defects: (**a**) Levels and localization of rhodopsin (Rho); **b***,* red/green (M-opsin); cone opsins (*green,* cone opsin); **c,** short wavelength (S-opsin); **d,** Pde6b; and **e,** PNA-488 in were analyzed from two and six months old WT and *Myo1c-KO* mice retinas. *Arrows* in ***a*** highlight rhodopsin mislocalization in *Myo1c* knock-out mouse retinas.

Fig. 3: Immunohistochemical analysis of protein trafficking in RPE and photoreceptors of wild-type/WT and *Myo1c* knock-out mice retinas: (**a**) Levels and localization of cone arresting (ARR); **b***,* STRA6, were analyzed from two- and six-months old WT and *Myo1c-KO* mice retinas to evaluate protein trafficking to photoreceptor outer segments or within the RPE. *Red Arrows* in ***a*** highlight cone photoreceptor nuclei, while *White Arrows* in ***a*** highlight cone photoreceptor OS in WT mouse retinas that were significantly reduced or shorter respectively in *Myo1c-KO* animals.

Fig. 4: Histological, H&E staining, and ultrastructural analysis shows reduced OS length of *Myo1c*-KO mice retinas: Retinas from two and six months old *Myo1c-KO* and WT mice (**a**) were sectioned and stained with H&E to evaluate pathological consequences of MYO1C loss. Ultrastructural analysis using transmission electron microscopy (TEM) of rod photoreceptors from two months old WT and *Myo1c-KO* mice (**b**), from six months old WT and *Myo1c-KO* mice (**c**) and cones from six months old mice for (**d, e**) are presented. Quantification of rod OS lengths (**f,** two months old mice; **g,** six months old). Rod (**b, c**) and cone (**d, e**) outer segment lengths were quantified based on the thickness of the respective outer segment areas. Scale bar=2um (**b, c**). Scale bar=800 nm (**d, e**).

542   RPE, retinal pigmented epithelium OS, outer segments; IS, inner segments;
543   ONL, outer nuclear layer; INL, inner nuclear layer; OPL, outer plexiform layer;
544   IPL, inner plexiform layer; GCL, ganglion cell layer.
545
Fig. 5: Rhodopsin is a direct cargo for MYO1C: (**a**) Using Rhodopsin antibody,
Rhodopsin (RHO) was immunoprecipitated either from mice retinal lysates or
from HEK293 cells where it was overexpressed, (**b**) and separated on SDS-
PAGE and transferred to PVDF membrane. The membrane was then incubated
with 5ug of purified recombinant active MYO1C-full length (**a**) or MYO1C C-
terminal cargo domain protein (**b**) generated from baculovirus expression
system. The blots were then western blotted with MYO1C antibody, which
showed direct binding of MYO1C to various Rhodopsin multimers present in the
retinal lysate and overexpressed HEK293 cells.

Supplementary Figures

Fig. S1: Western blot analysis of MYO1C in systemic tissues: Western blot analysis confirmed MYO1C absence in various systemic tissues of *Myo1c*-KO mice. Absence of MYO1C did not affect MYO1B or MYO1E expression. Actin was used as the protein loading control.

Fig. S2: MYO1C expression in mouse tissue by immunofluorescence: Expression of MYO1C in systemic tissues, kidney (**a**), liver (**b**), and heart (**c**) of WT and *Myo1c*-KO animals by immunofluorescence. WT, wild type; KO, knockout.

Fig. S3: Quantification of OS lengths and distribution of Opsins in *Myo1c* knock-out photoreceptors at two months of age: Cone lengths **(a)** and rhodopsin distribution within rod IS **(b)** and IS **(c)** were quantified based on the thickness of the respective outer segment areas. Approximately 500 cones in 5-6 retinal sections from each animal (n=8 animals; both genotypes) were sized. For quantification of rhodopsin distribution, 3-5 retinal sections from each animal (n=8 animals; both genotypes) were assayed. Mann-Whitney *U* test was used for statistical analysis and represented in Box-Whisker plots and considered significant *P*<0.05.

Fig. S4: Quantification and Distribution of Opsins in *Myo1c* knock-out photoreceptors at six months of age: Cone lengths **(a)** and rhodopsin distribution within rod IS **(b)** and IS **(c)** were quantified based on the thickness of the respective outer segment areas and fluorescence in different retinal layers. Approximately 500 cones in 5-6 retinal sections from each animal (n=8 animals; both genotypes) were sized. For quantification of rhodopsin distribution, 3-5 retinal sections from each animal (n=8 animals; both genotypes) were assayed. Mann-Whitney *U* test was used for statistical analysis and represented in Box-Whisker plots and considered significant *P*<0.05.

Fig. S5: Representative tissue histology and staining of systemic organs from WT and *Myo1c*-KO animals. Systemic organs, liver (**a**), kidney (**b**), and heart (**c**), of *Myo1c* and WT animals were sectioned and stained with H&E to evaluate any pathological consequences of systemic MYO1C loss.

Fig. S6: qPCR analysis of various MYO1C family members in mice retina: Retinas from WT and Myo1c-KO mice were isolated and processed for qPCR analysis using specific primers for various *Myo1c* family members including *Myo1b, d, e and f.*

References

1. Baehr, W., et al., *The function of guanylate cyclase 1 and guanylate cyclase 2 in rod and cone photoreceptors.* J Biol Chem, 2007. **282**(12): p. 8837-47.

2. Coffey, P.J., et al., *Complement factor H deficiency in aged mice causes retinal abnormalities and visual dysfunction.* Proc Natl Acad Sci U S A, 2007. **104**(42): p. 16651-6.

3. Fan, J., et al., *Rpe65-/- and Lrat-/- mice: comparable models of leber congenital amaurosis.* Invest Ophthalmol Vis Sci, 2008. **49**(6): p. 2384-9.

4. Grossman, G.H., et al., *Immunocytochemical evidence of Tulp1-dependent outer segment protein transport pathways in photoreceptor cells.* Exp Eye Res, 2011. **93**(5): p. 658-68.

5. Hüttl, S., et al., *Impaired channel targeting and retinal degeneration in mice lacking the cyclic nucleotide-gated channel subunit CNGB1.* J Neurosci, 2005. **25**(1): p. 130-8.

6. Karan, S., et al., *A model for transport of membrane-associated phototransduction polypeptides in rod and cone photoreceptor inner segments.* Vision Res, 2008. **48**(3): p. 442-52.

7. Pearring, J.N., et al., *Protein sorting, targeting and trafficking in photoreceptor cells.* Prog Retin Eye Res, 2013. **36**: p. 24-51.

8. Zhang, H., et al., *Trafficking of membrane-associated proteins to cone photoreceptor outer segments requires the chromophore 11-cis-retinal.* J Neurosci, 2008. **28**(15): p. 4008-14.

9. Zhang, T., et al., *Cone opsin determines the time course of cone photoreceptor degeneration in Leber congenital amaurosis.* Proc Natl Acad Sci U S A, 2011. **108**(21): p. 8879-84.

10. Abdelkader, E., et al., *Severe retinal degeneration at an early age in Usher syndrome type 1B associated with homozygous splice site mutations in MYO7A gene.* Saudi J Ophthalmol, 2018. **32**(2): p. 119-125.

11. Cheng, L., et al., *Identification of a novel MYO7A mutation in Usher syndrome type 1.* Oncotarget, 2018. **9**(2): p. 2295-2303.

12. Arif, E., et al., *Motor protein Myo1c is a podocyte protein that facilitates the transport of slit diaphragm protein Neph1 to the podocyte membrane.* Mol Cell Biol, 2011. **31**(10): p. 2134-50.

13. Nakamori, Y., et al., *Myosin motor Myo1c and its receptor NEMO/IKK-gamma promote TNF-alpha-induced serine307 phosphorylation of IRS-1.* J Cell Biol, 2006. **173**(5): p. 665-71.

14. Woolner, S. and W.M. Bement, *Unconventional myosins acting unconventionally.* Trends Cell Biol, 2009. **19**(6): p. 245-52.

15. Walsh, T., et al., *From flies' eyes to our ears: mutations in a human class III myosin cause progressive nonsyndromic hearing loss DFNB30.* Proc Natl Acad Sci U S A, 2002. **99**(11): p. 7518-23.

16. Rehman, A.U., et al., *Mutational Spectrum of MYO15A and the Molecular Mechanisms of DFNB3 Human Deafness.* Human mutation, 2016. **37**(10): p. 991-1003.

17. Adamek, N., M.A. Geeves, and L.M. Coluccio, *Myo1c mutations associated with hearing loss cause defects in the interaction with nucleotide and actin.* Cellular and molecular life sciences : CMLS, 2011. **68**(1): p. 139-150.

18. Adamek, N., M.A. Geeves, and L.M. Coluccio, *Myo1c mutations associated with hearing loss cause defects in the interaction with nucleotide and actin.* Cell Mol Life Sci, 2011. **68**(1): p. 139-50.

19. Lin, T., et al., *A hearing loss-associated myo1c mutation (R156W) decreases the myosin duty ratio and force sensitivity.* Biochemistry, 2011. **50**(11): p. 1831-8.

20. Skeie, J.M. and V.B. Mahajan, *Proteomic interactions in the mouse vitreous-retina complex.* PLoS One, 2013. **8**(11): p. e82140.

21. Boguslavsky, S., et al., *Myo1c binding to submembrane actin mediates insulin-induced tethering*
of GLUT4 vesicles. Mol Biol Cell, 2012. **23**(20): p. 4065-78.

22. Bose, A., et al., *Unconventional myosin Myo1c promotes membrane fusion in a regulated*
exocytic pathway. Mol Cell Biol, 2004. **24**(12): p. 5447-58.

23. Fan, Y., et al., *Myo1c facilitates G-actin transport to the leading edge of migrating endothelial*
cells. J Cell Biol, 2012. **198**(1): p. 47-55.

24. Arif, E., et al., *The motor protein Myo1c regulates transforming growth factor-beta-signaling and*
fibrosis in podocytes. Kidney Int, 2019.

25. Crosson, C.E., et al., *Inhibition of histone deacetylase protects the retina from ischemic injury.*
Invest Ophthalmol Vis Sci, 2010. **51**(7): p. 3639-45.

26. Husain, S., D.E. Potter, and C.E. Crosson, *Opioid receptor-activation: retina protected from*
ischemic injury. Invest Ophthalmol Vis Sci, 2009. **50**(8): p. 3853-9.

27. Krebs, W. and H. Kühn, *Structure of isolated bovine rod outer segment membranes.* Exp Eye Res,
1977. **25**(5): p. 511-26.

28. Pearring, J.N., et al., *Protein sorting, targeting and trafficking in photoreceptor cells.* Progress in
retinal and eye research, 2013. **36**: p. 24-51.

29. Brandstaetter, H., et al., *Loss of functional MYO1C/myosin 1c, a motor protein involved in lipid*
raft trafficking, disrupts autophagosome-lysosome fusion. Autophagy, 2014. **10**(12): p. 2310-23.

30. Mustafi, D., et al., *Photoreceptor phagocytosis is mediated by phosphoinositide signaling.* FASEB
journal : official publication of the Federation of American Societies for Experimental Biology,
2013. **27**(11): p. 4585-4595.

31. Lopes, V.S., et al., *The Usher 1B protein, MYO7A, is required for normal localization and function*
of the visual retinoid cycle enzyme, RPE65. Human molecular genetics, 2011. **20**(13): p. 2560-
2570.

32. Young, R.W., *The renewal of photoreceptor cell outer segments.* J Cell Biol, 1967. **33**(1): p. 61-72.

33. Young, R.W. and B. Droz, *The renewal of protein in retinal rods and cones.* J Cell Biol, 1968. **39**(1):
p. 169-84.

34. Findlay, J.B. and D.J. Pappin, *The opsin family of proteins.* Biochem J, 1986. **238**(3): p. 625-42.

35. Lobo, G.P., et al., *Involvement of Endoplasmic Reticulum Stress in TULP1 Induced Retinal*
Degeneration. PLoS One, 2016. **11**(3): p. e0151806.

36. Beech, P.L., et al., *Localization of kinesin superfamily proteins to the connecting cilium of fish*
photoreceptors. J Cell Sci, 1996. **109 (Pt 4)**: p. 889-97.

37. Muresan, V., et al., *Evidence for kinesin-related proteins associated with the axoneme of retinal*
photoreceptors. Exp Eye Res, 1997. **64**(6): p. 895-903.

38. Chaitin, M.H. and N. Coelho, *Immunogold localization of myosin in the photoreceptor cilium.*
Invest Ophthalmol Vis Sci, 1992. **33**(11): p. 3103-8.

39. Williams, D.S., M.A. Hallett, and K. Arikawa, *Association of myosin with the connecting cilium of*
rod photoreceptors. J Cell Sci, 1992. **103 (Pt 1)**: p. 183-90.

40. Mattapallil, M.J., et al., *The Rd8 mutation of the Crb1 gene is present in vendor lines of C57BL/6N*
mice and embryonic stem cells, and confounds ocular induced mutant phenotypes. Invest
Ophthalmol Vis Sci, 2012. **53**(6): p. 2921-7.

41. Lobo, G.P., et al., *The exocyst is required for photoreceptor ciliogenesis and retinal development.*
J Biol Chem, 2017. **292**(36): p. 14814-14826.

42. Husain, S., *Delta Opioids: Neuroprotective Roles in Preclinical Studies.* J Ocul Pharmacol Ther,
2018. **34**(1-2): p. 119-128.

43. Husain, S., et al., *PI3K/Akt Pathway: A Role in delta-Opioid Receptor-Mediated RGC*
Neuroprotection. Invest Ophthalmol Vis Sci, 2017. **58**(14): p. 6489-6499.

44. Bailes, H.J. and R.J. Lucas, *Human melanopsin forms a pigment maximally sensitive to blue light*
(λmax ≈ 479 nm) supporting activation of G(q/11) and G(i/o) signalling cascades. Proc Biol Sci,
2013. **280**(1759): p. 20122987.

Figure 1

693
694

Figure 3

Figure 4

  Figure 5

708
709
710 **Figure S1**
711

712
713

714　**Figure S2**

716
717

     Figure S3

     Figure S4

725 **Figure S5**
726

728   **Figure S6**
729

qPCR analysis from the eye sample isolated from wild-type and
Myo1c knockout mice