An alpaca nanobody neutralizes SARS-CoV-2 by blocking receptor interaction

Leo Hanke[1], Laura Vidakovics Perez[1], Daniel J. Sheward[1,4], Hrishikesh Das[2], Tim Schulte[3], Ainhoa

Moliner-Morro[1], Martin Corcoran[1], Adnane Achour[3], Gunilla B. Karlsson Hedestam[1], B. Martin

Hällberg[2, 5], Ben Murrell[1*] and Gerald M. McInerney[1*]

[1]Department of Microbiology, Tumor and Cell Biology, [2]Department of Cell and Molecular Biology and

[3]Science for Life Laboratory, Department of Medicine Solna, Karolinska Institutet, and Division of

Infectious Diseases, Karolinska University Hospital, Solna, Stockholm, Sweden. [4]Division of Virology,

Institute of Infectious Diseases and Molecular Medicine, Faculty of Health Sciences, University of Cape

Town, South Africa, [5]Centre for Structural Systems Biology, Notkestraße 85, 22607, Hamburg, Germany;

Karolinska Institutet VR-RÅC

* These authors contributed equally.

For correspondence: Martin Hällberg (martin.hallberg@ki.se), Ben Murrell (benjamin.murrell@ki.se) and

Gerald McInerney (gerald.mcinerney@ki.se).

Abstract

SARS-CoV-2 is the etiologic agent of COVID-19, currently causing a devastating pandemic for which

pharmacological interventions are urgently needed. The virus enters host cells through an interaction

between the spike glycoprotein and the angiotensin converting enzyme 2 (ACE2) receptor. Directly

preventing this interaction presents an attractive possibility for suppressing SARS-CoV-2 replication.

Here we report the isolation and characterization of an alpaca-derived single domain antibody fragment,

Ty1, that specifically targets the receptor binding domain (RBD) of the SARS-CoV-2 spike, directly

preventing ACE2 engagement. The nanobody binds with high affinity in the low nM range to the RBD,

occluding ACE2. A cryo-electron microscopy structure of the bound complex at 2.9 Å resolution reveals

that Ty1 binds to an epitope on the RBD accessible in both the 'up' and 'down' conformations and that Ty1

sterically hinders RBD-ACE2 binding. This 12.8 kDa nanobody does not need an Fc domain to neutralize

SARS-CoV-2, and can be expressed in high quantities in bacteria, presenting opportunities for

manufacturing at scale. Ty1 is therefore an excellent candidate as an intervention against COVID-19.

Main

SARS-CoV-2 emerged as the etiologic agent of COVID-19 in Wuhan, China in late 2019. In the comparatively short time since then, it has achieved pandemic status, causing more than 7.3 million cases, leading to at least 414,000 deaths and rising. Accordingly, the WHO declared the pandemic to be a public health emergency of international concern. A safe and effective vaccine is urgently needed, but requires time to develop. In the meantime, and indeed also in the post-vaccine era, highly specific and potent antiviral interventions are needed. Many generic or repurposed candidates are in trials, but so far results have been unremarkable. Since the virus is newly emerged, specifically designed drugs have not yet reached late phase trials. When available, specific antiviral drugs or antibody therapies will be used to protect individuals at risk and their widespread use will allow immunologically naïve populations to exit lockdowns more safely.

The virus is closely related to SARS-CoV-1, both being members of the lineage 2 betacoronaviruses. Cell entry of both viruses is achieved by first binding to the cell surface expressed receptor angiotensin-converting enzyme 2 (ACE2), followed by conformational changes in the viral spike glycoprotein trimer and subsequent membrane fusion. The affinity of SARS-CoV-2 receptor binding domain (RBD) for ACE2 is considerably higher than that for SARS-CoV-1[1,2], supporting efficient cell entry and likely contributing to pathogenesis. The RBD is a globular domain situated on the distal surface of the spike protein. Two conformations have been observed in the stabilized trimer. Specifically, one conformation where one RBD is ACE2-accessible while two are not and one conformation where all three RBDs are down, i.e. receptor inaccessible[2,3]. As the receptor-engaging part of the spike, the RBD is an attractive target for coronavirus neutralization, and a number of conventional neutralizing monoclonal antibodies that target the RBD and block receptor binding have already been isolated from convalescent patients[4–6].

Camelid-derived single domain antibody fragments, also called VHHs or nanobodies, offer several advantages over conventional antibodies as candidates for specific therapies. Despite being approximately one tenth of the size of a conventional antibody, they retain specificity and affinity similar to conventional antibodies while being far easier to clone, express and manipulate. They are readily expressed in bacteria in large quantities and show high thermal stability and solubility, making them easily scalable and extremely cost effective. Their modularity means that they can be oligomerized to increase avidity or to increase serum half-life[7]. Critical to their use as antivirals in humans, they can easily be humanized with existing protocols[8]. Importantly, they have proven to be highly potent inhibitors of viral infections in vivo, particularly respiratory infections[9,10].

We immunized one alpaca with SARS-CoV-2 S1-Fc and RBD in a 60-day immunization schedule. We

generated a phage display library and performed two consecutive rounds of phage display, followed by an

ELISA-based binding screen (Fig. 1a). We isolated one nanobody, Ty1, that binds specifically to the

RBD of the SARS-CoV-2 spike glycoprotein. In parallel we performed next generation sequencing

(NGS) on the baseline and post-enrichment libraries, and quantified variant frequency before and after

each enrichment step. Ty1 exhibited the greatest fold-change in frequency among all nanobody variants,

increasing over 10,000-fold from baseline to after the second enrichment round (Fig.1b). We report the

amino acid sequence of Ty1 in Figure 1c.

To determine whether Ty1 neutralized SARS-CoV-2 we employed an *in vitro* neutralization assay using

lentiviral particles pseudotyped with the SARS-CoV-2 spike protein. Ty1 neutralized SARS-CoV-2

pseudotyped viruses at an IC_{50} of 0.77 µg/ml (54 nM) (Fig. 2a). No neutralization of a lentivirus

pseudotyped with VSV-G by Ty1 was evident, and a control nanobody produced and purified in the same

way, but specific for influenza A virus nucleoprotein[11] showed no evidence of neutralization of SARS-

CoV-2 pseudotyped viruses. When Ty1 was expressed in mammalian cells as a Fc-fusion protein the

potent neutralization could be further increased to ~5 ng/ml (Fig. S1).

To confirm that Ty1 is directed specifically against the SARS-CoV-2 spike protein, we characterized the

specificity of Ty1 by flow cytometry. We site-specifically conjugated a fluorophore to the C-terminus of

Ty1 by means of a Sortase A reaction and copper-free click chemistry (Ty1-AS635P) and stained

untransfected cells and cells transiently transfected with SARS-CoV-2 spike under permeabilizing

conditions (Fig. 2b). While untransfected and unstained cells displayed similar signals, cells expressing

the viral spike protein showed a strong shift in fluorescence intensity when stained with Ty1-AS635P.

The apparent double peak likely reflected the varying efficiency of this transient transfection. To

determine if the same probe can be exploited to recognize the viral spike protein in immunofluorescence,

we infected Vero E6 cells with infectious SARS-CoV-2 at MOI 1 for 24 hours and stained the fixed and

permeabilized cells with Ty1-AS635P and anti-dsRNA antibody (Fig. 2c). While uninfected cells showed

no signal, infected cells were strongly labelled with both dsRNA antibody and Ty1-AS635P. Thus, Ty1

recognized the viral spike glycoprotein with high specificity in its native conformation in SARS-CoV-2-

infected cells. Importantly, the low background in both experiments also suggested that Ty1 is a highly

specific and suitable tool for research, diagnostics and therapy.

To understand the mechanism of neutralization, we evaluated the effect of Ty1 on RBD binding to ACE2.

We site-specifically conjugated a fluorophore to the C-terminus of the RBD (RBD-AS635P) and used this

probe to stain ACE2 expressing HEK293T cells (Fig. 2d). Preincubation of RBD-AS635P with unlabeled

Ty1 resulted in a significant reduction of ACE2 staining, while preincubation with the control nanobody
NP-VHH1 had no such effect. This result indicated that Ty1 directly prevents binding of SARS-CoV-2
RBD to its host cell receptor ACE2.

Specific and high-affinity binding of Ty1 to the RBD was also demonstrated in kinetic bio layer
interferometry (BLI) experiments. Dipping of surface-immobilized nanobodies into monomeric RBD
solutions at 550 nM yielded binding responses with fast association kinetics and amplitudes reaching 1.5
nm only for Ty1 but not NP-VHH1 (red and blue curves, respectively, in Figure 3a). Titration
experiments performed under normal (280 mM) and high salt (680 mM) conditions revealed
concentration-dependent kinetic response curves for binding of RBD to Ty1 (Figs. 3b and S2a,
respectively). The derived semi-log concentration-response curves revealed sigmoidal line-shapes with
fitted apparent K_D-values of 8 ± 1.5 and 13 ± 1.5 nm for binding at normal and high salt conditions,
respectively. Local fits to individual sensorgrams applying the standard 1:1 binding model appeared
reasonable for the association phases at lower to intermediate RBD concentrations, as well as for all
dissociation curves when fits were allowed to stay above zero (grey lines Figs. 3b and S2a 1[st] panel).
However, the model deviated from the observed data at higher RBD concentrations. Instead almost
perfect fits were obtained when the same data were analyzed in terms of a Bayesian two-dimensional
distribution of K_D and koff-rate constants to address heterogeneous ligand site populations on the sensor
surface[12–14]. For the two titrations at low and high salt conditions, distinct peaks at K_D- and k_{off}-rate values
of 5-9 nM and $4\text{-}5*10^{-3}$ s^{-1} were obtained (Fig. 3b and S2a 4[th] panel). In both conditions, a second
elevated plateau with K_D- and k_{off}-rate values of about 7 nM and $1\text{-}4*10^{-7}$ contributed significantly to the
observed sensorgrams. Since most high-affinity protein-protein interactions in the nM-range have
dissociation rates in the $1*10^{-3}$ s^{-1} range[15], we attribute the first defined peak as the relevant Ty1:RBD
interaction. The second broad plateau is likely caused by RBD competition and rebinding effects on the
sensor surface, as well as heterogeneous ligand populations[14,16,17]. The orthogonal biophysical method
isothermal titration calorimetry (ITC)[18] confirmed the high affinity binding of Ty1 to RBD with a K_D of 9
nM (with estimated bounds of 1 and 70 nM) characterized by an exothermic enthalpy of about -10 ± 0.5
kcal/mole (Fig. 3c, *left panel*). Exothermic binding was already evident from the three initial relatively
constant negative spikes that were caused by the injection of Ty1 to RBD (Fig. 3c, *right panel*). The
amplitude of the following three to four spikes returned to baseline demonstrating saturation of the
available RBD sites by Ty1 binding. Notably, return to baseline was accompanied by the appearance of
preceding positive spikes (Fig. 3c, *left panel* and Fig. S2b). These spikes were also detected when Ty1
was injected into the buffer (HBS) and thus treated as Ty1 dilution effects during data analysis. Injection
of NP-VHH1 into RBD did not cause any binding or dilution heat changes above background noise. It

should be noted that the ITC measurements were performed at the lowest possible protein concentrations

to derive K_D-values in the low nM range, while still being able to detect interaction heat above

background noise signals that were at about -0.17 ucal/s (maximum spike amplitude) and $\pm$ 0.05 ucal/s,

respectively. Altogether, we concluded from these results that RBD bound to surface-immobilized Ty1

with high affinity of about 5-10 nM.

To understand the structural basis underlying the potent neutralization of SARS-CoV-2 we performed a

cryo-EM structure determination of the prefusion-stabilized spike ectodomain in complex with Ty1. The

cryo-EM reconstruction reaches an overall resolution of 2.9 Å (Fig. 4a; 0.143 FSC) with strong variation

of estimated local resolution from high resolution in the core of the spike trimer to relatively low

resolution in the top of the spike. Nevertheless, the current reconstruction clearly shows that the spike

retains only one main conformation with one RBD 'up' and two RBDs 'down'. Importantly, all three

RBDs are decorated in their upper parts with a Ty1 nanobody. The nanobodies retain a similar binding

orientation to the RBD whether the RBD is found in the 'up' or 'down' conformation (Fig. 4a,b). Primary

interactions with the RBD are through the CDRs. Specifically, CDR1 interacts with RBD T470 and

V483-E484 and CDR3 interacts with RBD Y449, F490 and Q493. Interestingly, CDR2 does not form any

interactions with the RBD, instead it stabilizes the conformation of CDR1 in the RBD bound mode and

thereby acts indirectly to potentiate the Ty1-RBD binding.

Since ACE2 can only be bound by an RBD in the 'up' conformation, the current cryo-EM reconstruction

clearly shows that ACE2 binding is sterically hindered from two sides (Fig. 4c). Specifically, ACE2

binding is blocked both by the Ty1 nanobody bound to the RBD in the 'up' conformation and the

neighboring RBD in the 'down' conformation. Hence, ACE2 binding is sufficiently hindered with any

two of the available three binding RBD sites in the spike trimer.

The current coronavirus pandemic has drastic consequences for the world's population, and vaccines,

antibodies or antivirals are urgently needed. Neutralizing antibodies can block virus entry at an early step

of infection and potentially protect individuals that are at high risk of developing severe disease. We

report the identification and characterization of a SARS-CoV-2 RBD-specific single domain antibody

fragment (nanobody) termed Ty1 that potently neutralizes the virus. We identified Ty1 by binding assay

after two consecutive rounds of phage display, simultaneously monitoring sequence enrichment by NGS.

Although Ty1 exhibited the greatest fold-enrichment in the NGS analysis, multiple additional nanobodies

exhibited enrichment of varying extent across both rounds. As the correlation between phage display

enrichment and neutralization is likely imperfect, further analyses of our libraries will likely yield other

potent SARS-CoV-2 neutralizing nanobodies. In addition to neutralization activity, we also show that Ty1

can be used as a detection reagent in flow cytometry and immunofluorescence demonstrating its

suitability as a research tool and for diagnostics.

It should be noted that the nanobody Ty1 can be readily produced in bacteria at very high yield (in excess

of 30 mg/L culture), making it an excellent candidate for a low-cost, scalable antiviral agent against

SARS-CoV-2, and we provide the amino acid sequence, encouraging direct exploitation as such. While

nanobodies capable of binding SARS-CoV-2 spike have recently been isolated, these were generated after

SARS-CoV-1 spike immunization[19], or PCR maturation[20]. Also, in both cases a fusion to human Fc

domain is required for neutralization of SARS-CoV-2, precluding expression in bacterial culture. Naive

libraries of human single-domain antibodies (sdAbs) have also been screened to identify SARS-CoV-2

spike-specific nanobodies[21,22], but they lack detailed structural information. Ty1 represents the first

single-domain antibody isolated from an animal specifically immunized with a SARS-CoV-2 protein.

Future work will aim to improve the potency and potential efficacy of Ty1 through various strategies. For

example, mutational scanning may yield potency improvements to Ty1. Also, since Ty1 already

neutralizes as a monomeric protein, the generation of homodimeric or trimeric fusion constructs is

expected to further increase its neutralization activity. Indeed, fusion of Ty1 to a human IgG1-Fc

dramatically improved the IC_{50} of this molecule, to approximately 5 ng/ml (Fig. S2). Additional strategies

will explore linker-based constructs that chain multiple copies of Ty1 together, which may provide

similar improvements in potency while retaining the possibility of being expressed in bacteria. Ty1 may

additionally be a useful component of a bi-specific or tri-specific antibody, which could combine epitope

specificities to increase the mutational barrier to viral escape. Based on our work, we hope that Ty1 will

be investigated as a candidate for antiviral therapy.

Methods

Cells and virus. Vero E6 cells (ATCC-CRL-1586) and HEK293T cells (ATCC-CRL-3216) were

maintained in Dulbecco's Modified Eagle Medium (Gibco) supplemented with 10% fetal calf serum and

1% Penicillin-Streptomycin and cultured at 37°C in a humidified incubator with 5% CO_2. A HEK293T

cell line engineered to overexpress human ACE2 (HEK293T-ACE2) was generated by the lentiviral

transduction of HEK293T cells. Briefly, lentiviruses were produced by co-transfecting HEK293T cells

with a plasmid encoding VSV-G (Addgene cat#12259), a lentiviral Gag-Pol packaging plasmid (Addgene

cat#8455), and a human ACE2 transfer plasmid. Virions were harvested from the supernatant, filtered

through 0.45 µm filters, and used to transduce HEK293T cells. All cell lines used for experiments were

negative for *Mycoplasma* as determined by PCR. Infectious SARS-CoV-2[23] was propagated in Vero E6

cells and titrated by plaque assay.

Proteins and probes. The plasmid for expression of the SARS-CoV-2 prefusion-stabilized spike

ectodomain with a C-terminal T4 fibritin trimerization motif was obtained from[2]. The plasmid was used

to transiently transfect FreeStyle 293F cells using FreeStyle MAX reagent (Thermo Fisher Scientific).

The S ectodomain was purified from filtered supernatant on Streptactin XT resin (IBA Lifesciences),

followed by size-exclusion chromatography on a Superdex 200 in 5 mM Tris pH 8, 200 mM NaCl.

The RBD domain (RVQ – VNF) of SARS-CoV-2 was cloned upstream of an enterokinase cleavage site

and a human Fc. This plasmid was used to transiently transfect FreeStyle 293F cells using the FreeStyle

MAX reagent. The RBD-Fc fusion was purified from filtered supernatant on Protein G Sepharose (GE

Healthcare). The protein was cleaved using bovine enterokinase (GenScript) leaving a FLAG-tag at the

C-terminus of the RBD. Enzyme and Fc-portion was removed on HIS-Pur Ni-NTA resin (Thermo Fisher

Scientific) and Protein G sepharose (GE Healthcare) respectively, and the RBD was purified by size-

exclusion chromatography on a Superdex 200 in 50 mM Tris pH 8, 200 mM NaCl.

In addition, the RBD domain (RVQ – VNF) was cloned upstream of a Sortase A recognition site

(LPETG) and a 6xHIS tag and expressed in FreeStyle 293F cells as described above. RBD-HIS was

purified from filtered supernatant on His-Pur Ni-NTA resin, followed by size-exclusion chromatography

on a Superdex 200.

The nanobodies were cloned for expression in the pHEN plasmid with a C-terminal Sortase recognition

site (LPETG) and a 6xHIS tag. This plasmid was used to transform BL21 cells for periplasmic

expression. Expression was induced with 1 mM IPTG at OD600 = 0.6; cells were grown overnight at

30°C. Nanobodies were retrieved from the periplasm by osmotic shock and purified by Ni-NTA affinity

purification and size-exclusion chromatography.

For mammalian expression, the sequence encoding the nanobody Ty1 was cloned upstream of an human

IgG1. This plasmid was used to transiently transfect FreeStyle 293F cells using the FreeStyle MAX

reagent. The Ty1-Fc fusion was purified from filtered supernatant on Protein G Sepharose followed by

size-exclusion chromatography.

Biotinylated and fluorescent probes were generated using Sortase A as described here[24] and here[25]. In

brief, nanobodies were site-specifically biotinylated on the C-terminus using Sortase A 5M. Nanobody at

a concentration of 50 μM was incubated with sortase A 5M (5 μM), GGGK-biotin (200 μM) in 50 mM

Tris, pH 7.5, 150 mM NaCl, 10 mM CaCl$_2$, for 2 hours at 25°C. Unreacted nanobody and sortase was
removed with Ni-NTA resin and excess GGGK-biotin was removed using Zeba spin desalting columns
(0.5 mL, 7k MWCO, Thermo Fisher Scientific).

To generate the fluorescently labeled probes, first a dibenzocyclooctyne-amine (DBCO-amine, Sigma
Aldrich) was attached via sortase A to the nanobody or the RBD. (Reaction conditions: 50 µM RBD or
nanobody, 50 µM Sortase A 5M, 8 mM DBCO-amine in 50 mM Tris pH 7.5, 150 mM NaCl, 10 mM
CaCl$_2$, 2 hours, 25°C). Unreacted probe, sortase and excess DBCO-amine was removed using Ni-NTA
resin and PD-10 columns (GE Healthcare) respectively. Abberior Star 635P-azide (Abberior GMBH) was
attached to the DBCO-labeled proteins in a copper-free click chemistry reaction. Unreacted fluorophore
was removed on PD-10 column (RBD) or size-exclusion chromatography (nanobody).

Alpaca immunization. Alpaca immunization and phage display was performed similarly as described
here[26] and here[27]. In brief, the adult male alpaca Tyson at PreClinics, Germany, was immunized 4 times in
a 60-day immunization schedule. SARS-CoV-2 S1-sheep-Fc (Native Antigen Company, SKU:
REC31806) was used for the first two immunization, and SARS-CoV-2 RBD produced in FreeStyle 293F
cells was used for the last two immunizations.
246
Library generation and nanobody isolation. After the final boost, RNA was isolated from PBMCs
(RNA Plus mini kit, Qiagen). For cDNA synthesis, SuperScript III RT (Thermo Fisher Scientific) was
used with a combination of oligo(dT), random hexamers, or gene specific primers (AL.CH2,
ATGGAGAGGACGTCCTTGGGT and AL.CH2.2 TTCGGGGGGGAAGAYRAAGAC)[27]. Nanobody
sequences were PCR amplified and cloned into a phagemid vector for expression as pIII fusion. TG1 cells
(Lucigen) were transformed with this library by electroporation.

Cells were inoculated with VCSM13 helper phage, and the resulting phage was enriched in two
consecutive rounds of phage display on RBD immobilized on magnetic beads. After the second round of
phage display, individual bacterial colonies were picked in a 96 well format, grown until OD = 0.6 and
nanobody expression was induced by addition of 1 mM IPTG. After 16 hours incubation at 30°C,
bacterial supernatant was used as primary detection reagent in an ELISA coated with RBD or S
ectodomain. Bound nanobodies were detected with anti-E tag (Bethyl laboratories) secondary antibody.
Positive clones were sequenced and cloned into the pHEN expression vector for further characterization.

Amino acid sequence of Ty1

MAQVQLVETGGGLVQPGGSLRLSCAASGFTFSSVYMNWVRQAPGKGPEWVSRISPNSGNIGYT

DSVKGRFTISRDNAKNTLYLQMNNLKPEDTALYYCAIGLNLSSSSVRGQGTQVTVSS

Next generation sequencing (NGS) and analysis of nanobody libraries. Plasmids from nanobody

libraries before enrichment, and after each enrichment step, were amplified for 13 cycles using Q5 High-

Fidelity 2X Master Mix (NEB) according to manufacturer's instructions, using primers:

CACTCTTTCCCTACACGACGCTCTTCCGATCTCTCGCGGCCCAGCCGGCCATGG and

GGAGTTCAGACGTGTGCTCTTCCGATCTACCGGCGCACCACTAGTGCA, annealing at 72°C.

Illumina indexing primers were added using an additional 9 cycles, with Kapa HiFi. Amplicons were size

selected using Agencourt AMPure XP beads (bead ratio: 1:1), and were pooled at ratios of 6:2:1 for

pre:post-1:post-2 libraries, to account for the reduction in diversity expected during enrichment, and

sequenced on an Illumina MiSeq using the MiSeq Reagent Kit v3 (2x300) MS-102-3003.

Paired-end reads were merged using USEARCH11[28], and then processed in the Julia language, primarily

using the NextGenSeqUtils.jl package[29]. Briefly, reads are trimmed of primer sequences, and

deduplicated, maintaining read frequencies. Variant frequencies are calculated as combined frequency of

any reads matching a variant within 3% nucleotide divergence, using a kmer-based distance

approximation for rapid database search. Any reads with counts >3 from the second enrichment library

are searched for their variant frequencies across all databases. When calculating enrichment, to avoid

zeros due to sampling and to regularize against over-sensitivity to low-frequency baseline variants, all

frequencies are increased by the reciprocal of the size of the pre-enrichment database.

Neutralization Assay. Pseudotyped viruses were generated by the co-transfection of HEK293T cells with

plasmids encoding the SARS-CoV-2 spike protein harbouring an 18 amino acid truncation of the

cytoplasmic tail[30], a plasmid encoding firefly luciferase, and a lentiviral packaging plasmid (Addgene

cat#8455) using Lipofectamine 3000 (Invitrogen). Media was changed 12-16 hours after transfection, and

pseudotyped viruses were harvested at 48- and 72-hours post transfection, filtered through a 0.45 µm

filter, and stored at -80°C until use. Pseudotyped virus neutralization assays were adapted from protocols

previously validated to characterize the neutralization of HIV [31], but with the use of HEK293T-ACE2

cells. Briefly, pseudotyped viruses sufficient to generate ~100,000 RLUs were incubated with serial

dilutions of nanobodies for 60 min at 37°C. Approximately 15,000 HEK293T-ACE2 cells were then

added to each well and the plates were incubated at 37°C for 48 hours. Luminescence was then measured

using Bright-Glo (Promega) per the manufacturer's instructions on a GM-2000 luminometer (Promega)

with an integration time of 0.3s.

Flow Cytometry. Cells were trypsinized and fixed in 4% formaldehyde/PBS and stained with RBD-

AS635P under non-permeabilizing conditions or with Ty1-AS635P under permeabilizing conditions.

Fluorescence was quantified using a BD FACSCelesta and the FlowJo software package.

Immunofluorescence. Vero E6 cells were seeded onto coverslips in a 24-well plate and incubated

overnight at 37°C/5% CO_2. Cells were infected with SARS-CoV-2 at a MOI of 1 for 24 h. Cells were

fixed with 4% (v/v) formaldehyde, permeabilized in 0.5% Triton X-100 and blocked in 5% horse serum.

Cells were incubated with anti-dsRNA antibody (1:2000, J2 Scicons) for 1 hour at room temperature

followed by 1 hour staining with the secondary antibody anti-mouse-Alexa Fluor 488 (1:2000, Thermo

Fisher Scientific), Hoechst (1:1000, Invitrogen) and Ty1-AS635P (0.05 µg/mL). Coverslips were

mounted in mounting media and images were obtained using Zeiss Axiovert microscope and processed

using Adobe Photoshop.

Biophysical Biolayer interferometry (BLI) and isothermal titration calorimetry (ITC). BLI was

performed using single-use high-precision streptavidin biosensors (SAX) on an eight-channel Octet RED

instrument according to manufacturer's protocols (Fortebio)[16]. Assays were performed in 2xPBS

comprising 0,05% Tween-20 (PBST). Biotinylated nanobodies Ty1 and NP-VHH1 were loaded at

concentrations between 30 and 250 nM followed by quenching using biocytin to reach final sensor loads

of between 0.15 and 0.7 nm. For the comparative binding test, the eight sensors were divided into two

sets, each comprising double sample as well as single reference and single control sensors. Sample and

reference sensors were loaded with respective nanobodies. The SAX control was only quenched. Loading

of the two sets was performed consecutively to reach similar immobilization levels, while subsequent

association and dissociation phases were performed simultaneously. For association, the sample and

control sensors were dipped into RBD, while the reference sensor was dipped into PBST. For titration

experiments, all sensors were loaded simultaneously. During association one of the sensors was used as

reference and only dipped into PBST. Raw data were pre-processed, analyzed and fitted by applying the

1:1 binding model as implemented in the manufacturer's software. Bayesian analysis to obtain the two-

dimensional distribution of K_D and k_{off} -rate values were performed using Evilfit[12–14]. The shown titration

data were processed applying reference sensor subtraction and Savitzky-Golay filter operations.

For ITC, proteins were exchanged to 2xHBS-buffer (50 mM HEPES, 300 mM NaCl, pH 7.5) and isolated

as single peak populations by Superdex-200 HR10/300 size-exclusion chromatography. ITC

measurements were performed using an ITC200 calorimeter (GE Healthcare). The cell temperature was

set to 37ºC and the syringe stirring speed to 750 rpm. Before each experiment, the RBD and nanobodies

were loaded into the cell and syringe at concentrations of 4 μM and 75 μM, respectively. Data and

binding parameters were analyzed using the MicroCal PeakITC software (Malvern). The integrated heat

versus molar ratio plots of the Ty1:RBD interactions were obtained by subtracting the Ty1 dilution heat

uptake from the binding data. The NP-VHH1:RBD data were only baseline-corrected, since dilution

effects were not evident.

Raw and processed BLI/ITC data were imported into Rstudio for visualization and further analysis[32–34].

Data along with analysis R scripts will be made publicly available via Github and/or DataDryad.

Cryo-EM sample preparation and imaging. Spike trimer (0.7 mg/ml) and Ty1 (1.3 mg/ml) were mixed

in a 1:8 molar ratio and incubated on ice for 5 minutes. A 3-μl aliquot of the sample solution was applied

to glow-discharged CryoMatrix holey grids with amorphous alloy film (Zhenjiang Lehua Technology) in

a Vitrobot Mk IV (Thermo Fisher Scientific) at 4 degrees and 100% humidity (blot 10 s, blot force 3).

Cryo-EM data collection was performed with EPU 2.7 (Thermo Fisher Scientific) using a Krios G3i

transmission-electron microscope (Thermo Fisher Scientific) operated at 300 keV in the Karolinska

Institutet 3D-EM facility. Images were acquired in nanoprobe EFTEM mode with a slit width of 10 eV

using a GIF 967 energy filter (Ametek) and a K3 detector (Ametek) during 2.4 seconds with a dose rate of

4.1 e-/px/s resulting in a total dose of 38 e$^-$/Å^2 fractionated into 40 movie frames. Motion correction, CTF-

estimation, fourier binning (to 1.02 Å/px), picking and extraction in 428 pixel boxes were performed on

the fly using Warp [35].

A total of 13,589 micrographs were selected based on an estimated resolution cutoff of 4 Å and defocus

below 2 microns. and 573,036 particles were picked by Warp. Extracted particles were imported into

cryoSPARC v2.15.0[36] for 2D classification, 3D classification and non-uniform 3D refinement. The

particles were processed with C1 symmetry throughout. After 2D classification (300 classes) 354,678

particles were retained and used to build three ab-initio 3D reconstructions. These were further processed

for heterogeneous refinement that resulted in one reconstruction showing high-resolution structural

features in the core of the spike. One round of homogenous refinement followed by non-uniform

refinement resulted in a final reconstruction to an overall resolution of 2.9 Å (0.143 FSC) using 210,832

particles. To minimise the effects of loss of delocalised information due to tight boxing[37], the particles

were re-extracted with 600 pixel box size. In addition, localized reconstruction[38] were performed using

particles where all parts of the spike except the N-terminal domains, the RBDs and the nanobodies had

been subtracted[39]. The combined effects of these three approaches significantly increased the level of

density detail in the upper part of the spike.

Model building and structure refinement. A structure of the 2019-nCoV spike protein trimer[19] (PDB: 6VSB) was used as a starting model for model building. The model was extended and manually adjusted in COOT[40]. The Nanobody structure was homology modelled using SWISS-MODEL[41] taking PDB:5JMR[42] as a template. The missing regions of the RBD domains were built based on the RBD-ACE2 crystal structure (PDB: 6LZG)[43]. Structure refinement and manual model building were performed using COOT and PHENIX[44] in interspersed cycles with secondary structure and geometry restrained. All structure figures and all EM density-map figures were generated with UCSF ChimeraX[45].

Bibliography

1. Shang, J. *et al.* Cell entry mechanisms of SARS-CoV-2. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 11727–11734 (2020).

2. Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* **367**, 1260–1263 (2020).

3. Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* **181**, 281–292.e6 (2020).

4. Ju, B. *et al.* Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* (2020) doi:10.1038/s41586-020-2380-z.

5. Shi, R. *et al.* A human neutralizing antibody targets the receptor binding site of SARS-CoV-2. *Nature* (2020) doi:10.1038/s41586-020-2381-y.

6. Cao, Y. *et al.* Potent neutralizing antibodies against SARS-CoV-2 identified by high-throughput single-cell sequencing of convalescent patients' B cells. *Cell* (2020) doi:10.1016/j.cell.2020.05.025.

7. Bannas, P., Hambach, J. & Koch-Nolte, F. Nanobodies and Nanobody-Based Human Heavy Chain Antibodies As Antitumor Therapeutics. *Front. Immunol.* **8**, 1603 (2017).

8. Vincke, C. *et al.* General strategy to humanize a camelid single-domain antibody and identification of a universal humanized nanobody scaffold. *J. Biol. Chem.* **284**, 3273–3284 (2009).

9. Detalle, L. *et al.* Generation and Characterization of ALX-0171, a Potent Novel Therapeutic Nanobody for the Treatment of Respiratory Syncytial Virus Infection. *Antimicrob. Agents Chemother.* **60**, 6–13 (2016).

10. Cardoso, F. M. *et al.* Single-domain antibodies targeting neuraminidase protect against an H5N1 influenza virus challenge. *J. Virol.* **88**, 8278–8296 (2014).

11. Hanke, L. *et al.* The Antiviral Mechanism of an Influenza A Virus Nucleoprotein-Specific Single-Domain Antibody Fragment. *MBio* **7**, (2016).

12. Svitel, J., Balbo, A., Mariuzza, R. A., Gonzales, N. R. & Schuck, P. Combined affinity and rate
constant distributions of ligand populations from experimental surface binding kinetics and
equilibria. *Biophys. J.* **84**, 4062–4077 (2003).

13. Gorshkova, I. I., Svitel, J., Razjouyan, F. & Schuck, P. Bayesian Analysis of Heterogeneity in the
Distribution of Binding Properties of Immobilized Surface Sites. *Langmuir* **24**, 11577–11586 (2008).

14. Schuck, P. & Zhao, H. The Role of Mass Transport Limitation and Surface Heterogeneity in the
Biophysical Characterization of Macromolecular Binding Processes by SPR Biosensing. *Methods*
Mol. Biol. **627**, 15–54 (2010).

15. Pollard, T. D. A guide to simple and informative binding assays. *Mol. Biol. Cell* **21**, 4061–4067
(2010).

16. Abdiche, Y., Malashock, D., Pinkerton, A. & Pons, J. Determining kinetics and affinities of protein
interactions using a parallel real-time label-free biosensor, the Octet. *Anal. Biochem.* **377**, 209–217
(2008).

17. Kamat, V. & Rafique, A. Designing binding kinetic assay on the bio-layer interferometry (BLI)
biosensor to characterize antibody-antigen interactions. *Anal. Biochem.* **536**, 16–31 (2017).

18. Velazquez-Campoy, A., Leavitt, S. A. & Freire, E. Characterization of Protein-Protein Interactions
by Isothermal Titration Calorimetry. in *Protein-Protein Interactions* 35–54 (Humana Press, 2004).

19. Wrapp, D. *et al.* Structural Basis for Potent Neutralization of Betacoronaviruses by Single-Domain
Camelid Antibodies. *Cell* **181**, 1004–1015.e15 (2020).

20. Huo, J. *et al.* Structural characterisation of a nanobody derived from a naïve library that neutralises
SARS-CoV-2. (2020) doi:10.21203/rs.3.rs-32948/v1.

21. Chi, X. *et al.* Humanized Single Domain Antibodies Neutralize SARS-CoV-2 by Targeting Spike
Receptor Binding Domain. *bioRxiv* 2020.04.14.042010 (2020) doi:10.1101/2020.04.14.042010.

22. Wu, Y. *et al.* Identification of Human Single-Domain Antibodies against SARS-CoV-2. *Cell Host*
Microbe (2020) doi:10.1016/j.chom.2020.04.023.

23. Monteil, V. *et al.* Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using
Clinical-Grade Soluble Human ACE2. *Cell* **181**, 905–913.e7 (2020).

24. Antos, J. M. *et al.* Site-Specific Protein Labeling via Sortase-Mediated Transpeptidation. *Curr.*
Protoc. Protein Sci. **89**, 15.3.1–15.3.19 (2017).

25. Witte, M. D. *et al.* Production of unnaturally linked chimeric proteins using a combination of
sortase-catalyzed transpeptidation and click chemistry. *Nat. Protoc.* **8**, 1808–1819 (2013).

26. Sosa, B. A. *et al.* How lamina-associated polypeptide 1 (LAP1) activates Torsin. *Elife* **3**, e03239
(2014).

27. Maass, D. R., Sepulveda, J., Pernthaner, A. & Shoemaker, C. B. Alpaca (Lama pacos) as a

convenient source of recombinant camelid heavy chain antibodies (VHHs). *J. Immunol. Methods*

324, 13–25 (2007).

28.   Edgar, R. C. & Flyvbjerg, H. Error filtering, pair assembly and error correction for next-generation

sequencing reads. *Bioinformatics* **31**, 3476–3482 (2015).

29.   Kumar, V. *et al.* Long-read amplicon denoising. *Nucleic Acids Res.* **47**, e104 (2019).

30.   Rogers, T. F. *et al.* Rapid isolation of potent SARS-CoV-2 neutralizing antibodies and protection in

a small animal model. *bioRxiv* 2020.05.11.088674 (2020) doi:10.1101/2020.05.11.088674.

31.   Sarzotti-Kelsoe, M. *et al.* Optimization and validation of the TZM-bl assay for standardized

assessments of neutralizing antibodies against HIV-1. *J. Immunol. Methods* **409**, 131–146 (2014).

32.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer, 2016).

33.   Wickham, H. tidyverse: Easily Install and Load the 'Tidyverse. (2017).

34.   Team, R. C. R: A language and environment for statistical computing.

35.   Tegunov, D. & Cramer, P. Real-time cryo-electron microscopy data preprocessing with Warp. *Nat.*

Methods **16**, 1146–1152 (2019).

36.   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid

unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

37.   Rosenthal, P. B. & Henderson, R. Optimal Determination of Particle Orientation, Absolute Hand,

and Contrast Loss in Single-particle Electron Cryomicroscopy. *Journal of Molecular Biology* vol.

333 721–745 (2003).

38.   Ilca, S. L. *et al.* Localized reconstruction of subunits from electron cryomicroscopy images of

macromolecular complexes. *Nature Communications* vol. 6 (2015).

39.   Morais, M. C. *et al.* Bacteriophage φ29 scaffolding protein gp7 before and after prohead assembly.

Nature Structural & Molecular Biology vol. 10 572–576 (2003).

40.   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*

Crystallogr. D Biol. Crystallogr. **66**, 486–501 (2010).

41.   Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and complexes.

Nucleic Acids Res. **46**, W296–W303 (2018).

42.   Dahms, S. O. *et al.* The structure of a furin-antibody complex explains non-competitive inhibition by

steric exclusion of substrate conformers. *Sci. Rep.* **6**, 34303 (2016).

43.   Wang, Q. *et al.* Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. *Cell*

181, 894–904.e9 (2020).

44.   Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure

solution. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 213–221 (2010).

45.   Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis.

460  *Protein Sci.* **27**, 14–25 (2018).

461 46. Myszka, D. G., Jonsen, M. D. & Graves, B. J. Equilibrium analysis of high affinity interactions

462   using BIACORE. *Anal. Biochem.* **265**, 326–330 (1998).

Acknowledgements

SARS-CoV-2 virus was received from Jonas Klingström, KI. The authors thank James Voss and Deli Huang for providing reagents. We thank Jason McLellan for providing the expression plasmid of the stabilized SARS-CoV-2 spike protein. We thank PreClinics for their fast and efficient services as well as Gunnar Schulte (KI), Per-Åke Nygren (KTH Stockholm) and Tomas Nyman (Protein Science Facility at KI/SciLifeLabs) for providing access to BLI and ITC instruments. We also thank Fondation Dormeur, Vaduz for generous support of flow cytometry equipment. This work was supported by an EU grant (CoroNAb) to BM, GM and GBKH and by project grants from the Swedish Research Council to BM (2018-02381), BMH (2017-6702 and 2018-3808) and to GM (2018-03914 and 2018-03843).

Author Contributions

L.H., L.V.P., D.J.S., T.S., A.M.M., M.C. and B.M. performed experiments and analyzed data. H.D. and B.M.H. collected and processed cryo-EM data, built the model and performed the structural interpretation of the data with the corresponding paper sections. A.A. and G.B.K.H. gave critical advice. L.H., B.M. and G.M.M. conceived the study and wrote the initial draft. All authors contributed to the final draft.

Competing Interests Statement

The authors declare no conflicts of interest.

Figure 1 - Nanobody discovery.

a, Overview of the nanobody generation process. **b**, Variant frequencies quantified by NGS across successive enrichment steps. Identified using RBD bait, Ty1 exhibits the greatest total fold change of all nanobodies, increasing in proportion over 10,000-fold between initial and final libraries. **c**, Sequence of Ty1. Complementarity-determining regions (CDRs) are indicated. Color scheme according to ClustalX.

Figure 2 - Nanobody Ty1 neutralizes SARS-CoV-2 and specifically recognizes SARS-CoV-2 spike glycoprotein.

a, VSV G or SARS-CoV-2 spike pseudotyped lentivirus (PSV) was incubated with a dilution series of Ty1 or control nanobody (influenza NP-VHH1[11]). Infectivity relative to cells infected with pseudotyped virus in the absence of nanobody is shown. Neutralization by Ty1 was repeated in duplicate across 6 assays, and the error bars represent the standard deviation. **b**, Cells were transfected with a plasmid harboring the SARS-CoV-2 spike for 24 h. Cells were fixed, permeabilized and stained with Ty1-AS635P (black and red) or left unstained (grey). Cells were analyzed by flow cytometry. Cell counts are presented as % of max (representative histogram). **c**, Vero E6 cells were infected with SARS-CoV-2 at a MOI of 1 for 24 hours. Cells were fixed, permeabilized and stained for DNA (blue), dsRNA (green) and with Ty1-AS635P (red). Pictures were taken by fluorescence microscopy and representative examples are shown. **d**, ACE2 expressing HEK293T cells were trypsinized, fixed and stained with RBD-AS635P alone, or preincubated with IAV NP-VHH1 or Ty1. Cells were analyzed by flow cytometry.

Figure 3 - RBD bound to Ty1 with 5-10 nM affinities in biophysical BLI and ITC measurements

a, RBD bound to surface-immobilized Ty1 (red curves), but not to NP-VHH1 (blue curves). Almost equal

nanobody immobilization levels of about 0.7 nm were obtained by first loading Ty1 and then NP-VHH1.

In BLI, binding of molecules over time is recorded as sensorgrams recording the shift in wavelengths

(unit: nm) due to an increase in the optical thickness of the surface layer. **b**, (1st panel) RBD titration

sensorgrams obtained at high salt concentrations revealed concentration-dependent responses.

Sensorgrams are color-coded according to the log2 RBD concentration scale. Standard and Bayesian 1:1

binding models are shown as grey and black solid lines. (2nd panel) Pseudo-equilibrium response values

plotted against the logarithmic RBD concentration revealed sigmoidal binding curves that were fit to the

single-site interaction model [46] yielding K_D-values in the low nM range. K_D-values and standard

deviations are shown as solid and dotted lines, respectively. (3rd panel) Sensor immobilization levels are

shown as jittered box plots. (4th panel) Two-dimensional distribution of dissociation rate (koff) and

affinity (K_D) values obtained from the Bayesian and standard 1:1 model fits, visualized as densities

according to depicted normalized height scale and single black crosses, respectively. *See related Figure*

S2A. Plots for RBD/Ty1 titrations at low salt condition with same legends and scales. **c**, (1st panel) ITC

demonstrated high affinity binding of Ty1 to RBD with fitted K_D– and binding enthalpy (DH) mean

values of 9 nM (with estimated bounds of 1 and 70 nM) and -10 ± 0.5 kcal/mole, respectively, for two

binding experiments (red). Injection of NP-VHH1 into RBD yielded heat changes at background level

(blue). (2nd panel) Baseline-corrected heat changes plotted for two Ty1/RBD, a single Ty1/HBS buffer

and NP-VHH1/RBD titration experiments. Negative and positive heat changes are colored according to

the red-to-blue gradient. *See related Figure S2B:* Same figure on larger scale to highlight the Ty1 into

buffer dilution spikes.

a

Ty1-1　Ty1-3　Ty1-2

Side view

Viral membrane

Ty1-2　Ty1-3　Ty1-1

Side view

Viral membrane

180°

90°

Ty1-2

Ty1-1

Ty1-3

Top view

b

ACE2 interaction surface

Ty1-1

RBD1 (Up)

Ty1-3

Ty1-2

Ty1-2

CDR3

CDR2

CDR1

RBD

c

ACE2

Clash

Clash

Ty1-1

Ty1-3

Ty1-2

RBD1 (Up)

RBD3 (Down)

RBD2 (Down)

Figure 4 - Ty1 binds to the RBD in 'up' and 'down' conformation and prevents ACE2 engagement

a, Cryo-EM reconstruction to an overall resolution of 2.9 Å (0.143 FSC) of the spike trimer with three

bound molecules of Ty1. **b**, Atomic model (cartoon representation) of trimeric spike in complex with

three molecules of Ty1. Three chains of spike are shown in three different colors. The RBD of chain A

(light green) is present in 'up' conformation while the other two RBDs are captured in 'down'

conformation. The ACE2 interaction surface of RBD1 and the Ty1 interaction surface is highlighted.

Magnified view of RBD2 (in 'down' conformation) and Ty1 interaction is shown. CDR1,2 and 3 of Ty1

are shown in blue, green and red respectively. **c**, Ty1 shows a two-pronged inhibition of ACE2 receptor

binding through binding the RBD in the 'up' conformation and by binding to the neighbouring RBD in

the 'down' conformation. Binding of Ty1 to RBDs (both 'up' and 'down') would make ACE2 interaction

surface inaccessible for ACE2.

541
Supplemental Figure 1 - Increased neutralization of a Ty1-Fc construct

SARS-CoV-2 spike pseudotyped lentivirus (PSV) was incubated with a dilution series of Ty1 or Ty1-Fc.

Infectivity relative to cells infected with pseudotyped virus in the absence of nanobody is shown.

Neutralization was repeated in duplicate across 3 assays, and the error bars represent the standard

deviation

Supplemental Figure 2

a, related to Fig 3b, plots for RBD/Ty1 titrations at low salt condition with same legends and scales.

b, related to Fig 3c, Same data plotted on larger scale to highlight the Ty1 into buffer dilution spikes.

c, Samples after ITC were analyzed by SDS-PAGE.