An alpaca nanobody neutralizes SARS-CoV-2 by blocking receptor interaction

Leo Hanke[1], Laura Vidakovics Perez[1], Daniel J Sheward[1,4], Hrishikesh Das[2], Tim Schulte[3],

Ainhoa Moliner Morro[1], Martin Corcoran[1], Adnane Achour[3], Gunilla Karlsson Hedestam[1], B.

Martin Hällberg[2], Ben Murrell[1*] and Gerald M McInerney[1*]

[1]Department of Microbiology, Tumor and Cell Biology, [2]Department of Cell and Molecular

Biology and [3]Science for Life Laboratory, Department of Medicine Solna, Karolinska Institutet,

and Division of Infectious Diseases, Karolinska University Hospital, Solna, Stockholm, Sweden.

[4]Division of Virology, Institute of Infectious Diseases and Molecular Medicine, Faculty of Health

Sciences, University of Cape Town, South Africa.

* These authors contributed equally.

For correspondence: Ben Murrell (benjamin.murrell@ki.se), Martin Hällberg

(martin.hallberg@ki.se) and Gerald McInerney (gerald.mcinerney@ki.se).

Abstract

We report the isolation and characterization of an alpaca-derived, single domain antibody

fragment (nanobody) that specifically targets the receptor binding domain (RBD) of the SARS-

CoV-2 spike glycoprotein (spike) and potently neutralizes the virus. A cryo-electron microscopy

structure of the bound complex at 2.9 Å resolution reveals that the nanobody (Ty1) binds to an

epitope on the RBD accessible in both the 'up' and 'down' conformations and that Ty1 sterically

hinders RBD-ACE2 binding. Mechanistic characterization confirms that Ty1 directly interferes with

host cell receptor binding. This 12.8 kDa nanobody binds the SARS-CoV-2 spike with high

specificity and affinity, and can be produced in high quantities recombinantly thereby offering

potential as a potent and widely accessible SARS-CoV-2 antiviral agent.

Introduction

SARS-CoV-2 emerged as the etiologic agent of covid-19 in Wuhan, China in late 2019. In the

comparatively short time since then, it has achieved pandemic status, causing more than 5.5

million cases, leading to at least 378,000 deaths and rising. Accordingly, the WHO declared the

pandemic to be a public health emergency of international concern. A safe and effective vaccine

is urgently needed, but requires time to develop. In the meantime, and indeed also in the post-

vaccine era, highly specific and potent antiviral interventions are needed. Many generic or

repurposed candidates are in trials, but so far results have been unremarkable. Since the virus

is newly emerged, specifically designed drugs have not yet reached late phase trials. When

available, specific antiviral drugs or antibody therapies will be used to protect individuals at risk

and their widespread use will allow immunologically naïve populations to exit lockdowns more

safely.

The virus is closely related to SARS-CoV-1, both being members of the lineage 2

betacoronaviruses. Cell entry of both viruses is achieved by first binding to the cell surface

expressed receptor angiotensin-converting enzyme 2 (ACE2), followed by conformational

changes in the viral spike glycoprotein trimer and subsequent membrane fusion. The affinity of

SARS-CoV-2 receptor binding domain (RBD) for ACE2 is considerably higher than that for

SARS-CoV-1 [1,2], supporting efficient cell entry and likely contributing to pathogenesis. The

RBD is a globular domain situated on the distal surface of the spike protein. Two conformations

have been observed in the stabilized trimer. Specifically, one conformation where one RBD is

ACE2-accessible while two are not and one conformation where all three RBDs are down, i.e.

receptor inaccessible [2,3]. As the receptor-engaging part of the spike, the RBD is an attractive

target for coronavirus neutralization, and a number of conventional neutralizing monoclonal

antibodies that target the RBD and block receptor binding have already been isolated from

convalescent patients [4–6].

Camelid-derived single domain antibody fragments, also called VHHs or nanobodies, offer

several advantages over conventional antibodies as candidates for specific therapies. Despite

being approximately one tenth of the size of a conventional antibody, they retain specificity and

affinity similar to conventional antibodies while being far easier to clone, express and

manipulate. They are readily expressed in bacteria in large quantities and show high thermal

stability and solubility, making them easily scalable and extremely cost effective. Their

modularity means that they can be oligomerized to increase avidity or to increase serum half-life

[7]. Critical to their use as antivirals in humans, they can easily be humanized with existing

protocols [8]. Importantly, they have proven to be highly potent inhibitors of viral infections in

vivo, particularly respiratory infections [9,10].

Here, we describe the isolation, evaluation and molecular determination of an alpaca-derived

nanobody, Ty1, directed to the receptor-binding domain of the SARS-CoV-2 spike glycoprotein.

We demonstrate that the monomeric 12.8 kDa Ty1 molecule potently neutralizes SARS-CoV-2

spike pseudovirus. The nanobody binds with high affinity to the RBD in a manner that occludes

ACE2 interaction. We have also determined the mechanism of neutralization to be due to direct

interference with RBD binding to ACE2. Altogether, these results highlight the great potential of

Ty1 as a SARS-CoV-2 antiviral agent.

Results

<u>Generation of SARS-CoV-2 RBD-specific nanobodies</u>

We immunized one alpaca with SARS-CoV-2 S1-Fc and RBD in a 60-day immunization

schedule. We generated a phage display library and performed two consecutive rounds of

phage display, followed by an ELISA-based binding screen (Fig. 1A). We isolated one

nanobody, Ty1, that binds specifically to the RBD of the SARS-CoV-2 spike glycoprotein. In

parallel we performed next generation sequencing (NGS) on the baseline and post-enrichment

libraries, and quantified variant frequency before and after each enrichment step. Ty1 exhibited

the greatest fold-change in frequency among all nanobody variants, increasing over 10,000-fold

from baseline to after the second enrichment round (Fig.1B). We report the amino acid

sequence of Ty1 in Figure 1C.

<u>Ty1 neutralizes SARS-CoV-2 spike pseudotyped viruses</u>

To determine whether Ty1 neutralized SARS-CoV-2 we employed an *in vitro* neutralization

assay using lentiviral particles pseudotyped with the SARS-CoV-2 spike protein. Ty1 neutralized

SARS-CoV-2 pseudotyped viruses at an IC_{50} of 0.77 μg/ml (54 nM) (Fig. 2A). No neutralization

of a lentivirus pseudotyped with VSV-G by Ty1 was evident, and control nanobodies produced

and purified in the same way, but specific for either influenza A virus nucleoprotein [11] or GFP

[12], showed no evidence of neutralization of SARS-CoV-2 pseudotyped viruses.

<u>Ty1 specifically recognizes the SARS-CoV-2 spike protein in transfected and infected cells</u>

To confirm that Ty1 is directed against the SARS-CoV-2 spike protein, we characterized the

specificity of Ty1 by flow cytometry. We site-specifically conjugated a fluorophore to the C-

terminus of the Ty1 by means of a Sortase A reaction and copper-free click chemistry (Ty1-

AS635P), and stained untransfected cells and cells transiently transfected with SARS-CoV-2

spike under permeabilizing conditions (Fig. 2B). While untransfected and unstained cells

displayed similar signals, cells expressing the viral spike protein showed a strong shift in

fluorescence intensity when stained with Ty1-AS635P. The apparent double peak likely

reflected the varying efficiency of this transient transfection. To determine if the same probe can

be exploited to recognize the viral spike protein in immunofluorescence, we infected Vero E6

cells with infectious SARS-CoV-2 at MOI 1 for 24 hours and stained the fixed and permeabilized

cells with Ty1-AS635P and anti-dsRNA antibody (Fig. 2C). While uninfected cells showed no

signal, infected cells were strongly labelled with both dsRNA antibody and Ty1-AS635P. Thus,

Ty1 recognized the viral spike glycoprotein with high specificity in its native conformation in

SARS-CoV-2-infected cells. Importantly, the low background in both experiments also

suggested that Ty1 is a highly specific and suitable tool for research, diagnostics and therapy.

<u>Ty1 competes directly with ACE2 for binding to the RBD of SARS-CoV-2 spike</u>

To understand the neutralizing activity, we evaluated the effect of Ty1 on RBD binding to ACE2.

We site-specifically conjugated a fluorophore to the C-terminus of the RBD (RBD-AS635P) and

used this probe to stain ACE2 expressing HEK293T cells (Fig. 2D). Preincubation of RBD-

AS635P with unlabeled Ty1 resulted in a significant reduction of ACE2 staining, while

preincubation with the control nanobody NP-VHH1 had no such effect. This result indicated that

Ty1 directly prevents binding of SARS-CoV-2 RBD to its host cell receptor ACE2.

<u>Ty1 binds the RBD with high affinity</u>

The highly specific and high-affinity binding of Ty1 to the RBD was also confirmed in kinetic bio

layer interferometry (BLI) experiments. Monomeric RBD bound to surface-immobilized Ty1, but

not to NP-VHH1 or to biocytin-quenched Streptavidin (SAX) (Fig. 3A). Dipping of the two Ty1-

sensors into RBD at a concentration of 550 nM yielded binding responses with fast association

kinetics and amplitudes reaching 1.5 nm (red curves), while dipping of the two NP-VHH1-

sensors into the same RBD solutions yielded responses comparable to the reference sensors

(2x violet and blue as well as light-green curves, respectively). RBD did not bind to the SAX

control (green curves). Approximately half of RBD remained bound to Ty1 after 45 min (2700 s)

of dissociation.

Titration experiments performed under normal and high salt conditions revealed concentration-

dependent kinetic response curves for binding of RBD to Ty1 (Fig. 3B). The derived semi-log

concentration-response curves revealed sigmoidal line-shapes with fitted apparent K_D-values of

8 ± 1.5 and 13 ± 1.5 nm for binding at normal and high salt conditions, respectively.

Kinetic fits of each sensorgram applying a 1:1 binding model revealed reasonable fits for

association phases of curves obtained at low and intermediate RBD concentrations, as well as

for most dissociation curves (Fig. 3B and 3C, left panels). However, relatively large deviations of

the fits showed that the 1:1 interaction model did not fully explain the obtained data (residuals in

Fig. 3C). Nevertheless, most of the single-sensorgram fits yielded K_D-values in the 10-100 nM

range with association and dissociation constants of about 1-1.2 $*10^5$ (Ms)$^{-1}$ and 3.5–4.3$*10^{-3}$ s^{-1}

(k_{on} versus K_D plot in the lower panel of Fig. 3B). Most protein-protein interactions have

association rates in the 10^6 to 10^7 M^{-1}s^{-1} range, therefore high-affinity interactions with K_D-

values in the nM range have typically dissociation rates around 1$*10^{-3}$ s^{-1} [13].

Altogether, we concluded from these results that RBD bound to surface-immobilized Ty1 with

high affinity in the 5-50 nM range.

<u>Cryo-EM of prefusion spike in complex with Ty1</u>

To understand the structural basis underlying the potent neutralization of SARS-CoV-2 we

performed a preliminary cryo-EM structure determination of the prefusion-stabilized spike

ectodomain in complex with Ty1. The current cryo-EM reconstruction reaches an overall

resolution of 2.9 Å (Fig. 4A; 0.143 FSC) with strong variation of estimated local resolution from

high resolution in the core of the spike trimer to relatively low resolution in the top of the spike.

Nevertheless, the current reconstruction clearly shows that the spike retains only one main

conformation with one RBD 'up' and two RBDs 'down'. Importantly, all three RBDs are

decorated in their upper parts with a Ty1 nanobody. The nanobodies retain a similar binding

orientation to the RBD whether the RBD is found in the 'up' or 'down' conformation (Fig. 4B,C).

As ACE2 can only be bound by an RBD in the 'up' conformation, the current cryo-EM

reconstruction clearly shows that ACE2 binding is sterically hindered from two sides (Fig. 4D).

Specifically, ACE2 binding is blocked both by the Ty1 nanobody bound to the RBD in the 'up'

conformation and the neighboring RBD in the 'down' conformation. Hence, ACE2 binding is

sufficiently hindered with any two of the available three binding RBD sites in the spike trimer. As

the RBD-Ty1 interaction interface is poorly resolved in the present reconstruction, we currently

refrain from molecular modelling of the exact binding interactions.

Discussion

The current coronavirus pandemic has drastic consequences for the world's population, and

vaccines, antibodies or antivirals are urgently needed. Neutralizing antibodies can block virus

entry at an early step of infection and potentially protect individuals that are at high risk of

developing severe disease. We report the identification and characterization of a SARS-CoV-2

RBD-specific single domain antibody fragment (nanobody) termed Ty1 that potently neutralizes

the virus. We provide structural and mechanistic insights that demonstrate that Ty1 prevents

RBD binding to its host cell receptor ACE2, and thus prevents SARS-CoV-2 virions from

attaching to cells. Ty1 binds the RBD with a kD within the 5-50 nM range, and neutralizes

SARS-CoV-2 pseudotyped virus at an $IC_{50}$of 0.77ug/ml (54 nM), representing the most potent

SARS-CoV-2 specific nanobody reported to date.

We identified Ty1 by binding assay after consecutive rounds of phage display, simultaneously

monitoring sequence enrichment by NGS. Although Ty1 exhibited the greatest fold-enrichment

in the NGS analysis, multiple additional nanobodies exhibited enrichment of varying extent

across both rounds. As the correlation between phage display enrichment and neutralization is

likely imperfect, further analyses of our libraries will likely yield other potent SARS-CoV-2

neutralizing nanobodies. In addition to neutralization activity, we also show that Ty1 can be

used as a detection reagent in flow cytometry and immunofluorescence demonstrating its

suitability as a research tool and for diagnostics.

It should be noted that the nanobody Ty1 can be readily produced in bacteria at very high yield

(in excess of 30 mg/L culture), and shows high affinity, making it an excellent candidate as a

low-cost, scalable antiviral agent against SARS-CoV-2, and we provide the amino acid

sequence, encouraging direct exploitation as such. While a llama nanobody capable of binding

SARS-CoV-2 has recently been isolated, this molecule was elicited against SARS-CoV-1 and

Fc-fusion was required to mediate neutralization of SARS-CoV-2 [14], precluding expression in

bacterial culture. Since Ty1 already neutralizes as a monomeric protein, the generation of

homodimeric or trimeric fusion constructs is likely to further increase its neutralization activity.

Our structural results provide a template for the rational design of such oligomers that will

potentially 'lock' the spike in a receptor-inaccessible conformation. Based on our work, we hope

that Ty1 can be investigated as a candidate for antiviral therapy.

Methods

Cells and virus

Vero E6 cells and HEK293T cells (ATCC-CRL-3216) were maintained in Dulbecco's Modified

Eagle Medium (Gibco) supplemented with 10% fetal calf serum and 1% Penicillin-Streptomycin

and cultured at 37°C in a humidified incubator with 5% CO_2. A HEK293T cell line engineered to

overexpress human ACE2 (HEK293T-ACE2) was generated by the lentiviral transduction of

HEK293T cells. Briefly, lentiviruses were produced by co-transfecting HEK293T cells with a

plasmid encoding VSV-G (Addgene cat#12259), a lentiviral Gag-Pol packaging plasmid

(Addgene cat#8455), and a human ACE2 transfer plasmid. Virions were harvested from the

supernatant, filtered through 0.45 μm filters, and used to transduce HEK293T cells. All cell lines

used for experiments were negative for *Mycoplasma* as determined by PCR.

Infectious SARS-CoV-2 [15] was propagated in Vero E6 cells and titrated by plaque assay.

Proteins and probes.

The plasmid for expression of the SARS-CoV-2 prefusion-stabilized spike ectodomain with a C-

terminal T4 fibritin trimerization motif was obtained from [2]. The plasmid was used to transiently

transfect FreeStyle 293F cells using FreeStyle MAX reagent (Thermo Fisher Scientific). The S

ectodomain was purified from filtered supernatant on Streptactin XT resin (IBA Lifesciences),

followed by size-exclusion chromatography on a Superdex 200 in 5 mM Tris pH 8, 200 mM

NaCl.

The RBD domain (RVQ – VNF) of SARS-CoV-2 was cloned upstream of an enterokinase

cleavage site and a human FC. This plasmid was used to transiently transfect FreeStyle 293F

cells using the FreeStyle MAX reagent. The RBD-FC fusion was purified from filtered

supernatant on Protein G Sepharose (GE Healthcare). The protein was cleaved using bovine

enterokinase (GenScript) leaving a FLAG-tag at the C-terminus of the RBD. Enzyme and FC-

portion was removed on HIS-Pur Ni-NTA resin (Thermo Fisher Scientific) and Protein G

sepharose (GE Healthcare) respectively, and the RBD was purified by size-exclusion

chromatography on a Superdex 200 in 50 mM Tris pH 8, 200 mM NaCl.

In addition, the RBD domain (RVQ – VNF) was cloned upstream of a Sortase A recognition site

(LPETG) and a 6xHIS tag, and expressed in 293F cells as described above. RBD-HIS was

purified from filtered supernatant on His-Pur Ni-NTA resin, followed by size-exclusion

chromatography on a Superdex 200.

The nanobodies were cloned for expression in the pHEN plasmid with a C-terminal Sortase

recognition site (LPETG) and a 6xHIS tag. This plasmid was used to transform BL21 cells for

periplasmic expression. Expression was induced with 1 mM IPTG at OD600 = 0.6; cells were

grown overnight at 30°C. Nanobodies were retrieved from the periplasm by osmotic shock and

purified by Ni-NTA affinity purification and size-exclusion chromatography.

Biotinylated and fluorescent probes were generated using Sortase A as described here [16] and

here [17]. In brief, nanobodies were site-specifically biotinylated on the C-terminus using

Sortase A 5M. Nanobody at a concentration of 50 µM was incubated with sortase A 5M (5 µM),

GGGK-biotin (200 µM) in 50 mM Tris, pH 7.5, 150 mM NaCl, 10 mM CaCl$_2$, for 2 hours at 25°C.

Unreacted nanobody and sortase was removed with Ni-NTA resin and excess GGGK-biotin was

removed with Zeba spin desalting columns (0.5 mL, 7k MWCO, Thermo Fisher Scientific).

To generate the fluorescently labeled probes, first a dibenzocyclooctyne-amine (DBCO-amine,

Sigma Aldrich) was attached via sortase A to the nanobody or the RBD. (Reaction conditions:

50 µM RBD or nanobody, 50 µM Sortase A 5M, 8 mM DBCO-amine in 50 mM Tris pH 7.5, 150

mM NaCl, 10 mM CaCl$_2$, 2 hours, 25°C). Unreacted probe, sortase and excess DBCO-amine

was removed using Ni-NTA resin and PD-10 columns (GE Healthcare) respectively. Abberior

Star 635P-azide (Abberior GMBH) was attached to the DBCO-labeled proteins in a copper-free

click chemistry reaction. Unreacted fluorophore was removed on PD-10 column (RBD) or size-

exclusion chromatography (nanobody).

Alpaca immunization

Alpaca immunization and phage display was performed similarly as described here [18] and

here [19]. In brief, the adult male alpaca Tyson at PreClinics, Germany, was immunized 4 times

in a 60-day immunization schedule. SARS-CoV-2 S1-sheep-FC (Native Antigen Company,

SKU: REC31806) was used for the first two immunization, and SARS-CoV-2 RBD produced in

FreeStyle 293F cells was used for the last two immunizations.

Library generation and nanobody isolation

After the final boost, RNA was isolated from PBMCs (RNA Plus mini kit, Qiagen). For cDNA

synthesis, SuperScript III RT (Thermo Fisher Scientific) was used with a combination of

oligo(dT), random hexamers, or gene specific primers (AL.CH2,

ATGGAGAGGACGTCCTTGGGT and AL.CH2.2 TTCGGGGGGAAGAYRAAGAC) [19].

Nanobody sequences were PCR amplified and cloned into a phagemid vector for expression as

pIII fusion. This library was electroporated into TG1 cells (Lucigen).

Cells were inoculated with VCSM13 helper phage, and the resulting phage was enriched in two

consecutive rounds of phage display on RBD immobilized on magnetic beads. After the second

round of phage display, individual bacterial colonies were picked in a 96 well format, grown until

OD = 0.6 and nanobody expression was induced by addition of 1 mM IPTG. After 16 hours

incubation at 30°C, bacterial supernatant was used as primary detection reagent in an ELISA

coated with RBD or S ectodomain. Bound nanobodies were detected with anti-E tag (Bethyl

laboratories) secondary antibody. Positive clones were sequenced and cloned into the pHEN

expression vector for further characterization.

Amino acid sequence of Ty1

MAQVQLVETGGGLVQPGGSLRLSCAASGFTFSSVYMNWVRQAPGKGPEWVSRISPNSGNIG

YTDSVKGRFTISRDNAKNTLYLQMNNLKPEDTALYYCAIGLNLSSSSVRGQGTQVTVSS

Next generation sequencing (NGS) and analysis of nanobody libraries.

Plasmids from nanobody libraries before enrichment, and after each enrichment step, were

amplified for 13 cycles using Q5 High-Fidelity 2X Master Mix (NEB) according to manufacturer's

instructions, using primers:

CACTCTTTCCCTACACGACGCTCTTCCGATCTCTCGCGGCCCAGCCGGCCATGG and

GGAGTTCAGACGTGTGCTCTTCCGATCTACCGGCGCACCACTAGTGCA, annealing at 72°C.

Illumina indexing primers were added using an additional 9 cycles, with Kapa HiFi. Amplicons

were size selected using Agencourt AMPure XP beads (bead ratio: 1:1), and were pooled at

ratios of 6:2:1 for pre:post-1:post-2 libraries, to account for the reduction in diversity expected

during enrichment, and sequenced on an Illumina MiSeq using the MiSeq Reagent Kit v3

(2x300) MS-102-3003.

Paired-end reads were merged using USEARCH11 [20], and then processed in the Julia

language, primarily using the NextGenSeqUtils.jl package [21]. Briefly, reads are trimmed of

primer sequences, and deduplicated, maintaining read frequencies. Variant frequencies are

calculated as combined frequency of any reads matching a variant within 3% nucleotide

divergence, using a kmer-based distance approximation for rapid database search. Any reads

with counts >3 from the second enrichment library are searched for their variant frequencies

across all databases. When calculating enrichment, to avoid zeros due to sampling and to

regularize against over-sensitivity to low-frequency baseline variants, all frequencies are

increased by the reciprocal of the size of the pre-enrichment database.

Neutralization Assay

Pseudotyped viruses were generated by the co-transfection of HEK293T cells with plasmids

encoding the SARS-CoV-2 spike protein harboring an 18 amino acid truncation of the

cytoplasmic tail [22], a plasmid encoding firefly luciferase, and a lentiviral packaging plasmid

(Addgene cat#8455) using Lipofectamine 3000 (Invitrogen). Media was changed 12-16 hours

after transfection, and pseudotyped viruses were harvested at 48- and 72-hours post

transfection, filtered through a 0.45 µm filter, and stored at -80°C until use. Pseudotyped

neutralization assays were adapted from protocols previously validated to characterize the

neutralization of HIV [23], but with the use of HEK293T-ACE2 cells. Briefly, pseudotyped viruses

sufficient to generate ~100,000 RLUs were incubated with serial dilutions of nanobodies for 60

min at 37°C. Approximately 15,000 HEK293T-ACE2 cells were then added to each well and the

plates were incubated at 37°C for 48 hours. Luminescence was then measured using Bright-Glo

(Promega) per the manufacturer's instructions on a GM-2000 luminometer (Promega) with an

integration time of 0.3s.

Flow Cytometry

Cells were trypsinized and fixed in 4% formaldehyde/PBS and stained with RBD-AS635P under

non-permeabilizing conditions or with Ty1-AS635P under permeabilizing conditions.

Fluorescence was quantified using a BD FACSCelesta and the FlowJo software package.

Immunofluorescence

Vero E6 cells were seeded onto coverslips in a 24 well plate and incubated overnight at

37°C/5% CO_2. Cells were infected with SARS-CoV-2 at a MOI of 1 for 24 h. Cells were fixed

with 4% (v/v) formaldehyde, permeabilized in 0.5% Triton X-100 and blocked in 5% horse

serum. Cells were incubated with anti-dsRNA antibody (1:2000, J2 Scicons) for 1 hour at room

temperature followed by 1 hour staining with the secondary antibody anti-mouse-Alexa488

(1:2000, Thermo Fisher Scientific), Hoechst (1:1000, Invitrogen) and Ty1-AS635P (0.05 µg/mL).

Coverslips were mounted in mounting media and images were obtained using Zeiss Axiovert

microscope and processed using Adobe Photoshop.

Biolayer interferometry (BLI)

BLI was performed using single-use high-precision streptavidin biosensors (SAX) on an eight-

channel Octet RED instrument according to manufacturer's protocols (Fortebio)[24]. Assays

were performed in 2xPBS comprising 0,05% Tween-20 (PBST). Biotinylated nanobodies Ty1

and NP-VHH1 were loaded at concentrations between 30 and 250 nM followed by quenching

using biocytin to reach final sensor loads of between 0.15 and 0.7 nm.

For the comparative binding test, the eight sensors were divided into two sets, each comprising

double sample as well as single reference and single control sensors. Sample and reference

sensors were loaded with respective nanobodies. The SAX control was only quenched. Loading

of the two sets was performed consecutively to reach similar immobilization levels, while

subsequent association and dissociation phases were performed simultaneously. For

association, the sample and control sensors were dipped into RBD, while the reference sensor

was dipped into PBST. For titration experiments, all sensors were loaded simultaneously.

During association one of the sensors was used as reference and only dipped into PBST.

Raw data were pre-processed, analyzed and fitted by applying the 1:1 binding model as

implemented in the manufacturer's software[25] . The processed data were imported into

Rstudio for visualization and further analysis [26–28]. The comparative binding test shown in

Figure 3A comprises un-processed raw data. The titration data in Figure 3B were processed

applying reference sensor subtraction and Savitzky-Golay filter operations. Raw data and the

analysis visualization scripts will be uploaded to Dryad and/or Github.

Cryo-EM sample preparation and imaging

Spike trimer (0.7 mg/ml) and Ty1 (1.3 mg/ml) were mixed in a 1:8 molar ratio and incubated on

ice for 5 minutes. A 3-µl aliquot of the sample solution was applied to glow-discharged CryoMatrix

holey grids with amorphous alloy film (Zhenjiang Lehua Technology) in a Vitrobot Mk IV (Thermo

Fisher Scientific) at 4 degrees and 100% humidity (blot 10 s, blot force 3).

Cryo-EM data collection was performed with EPU 2.7 (Thermo Fisher Scientific) using a Krios G3i

transmission-electron microscope (Thermo Fisher Scientific) operated at 300 keV in the

Karolinska Institutet 3D-EM facility. Images were acquired in nanoprobe EFTEM mode with a slit

width of 10 eV using a GIF 967 energy filter (Ametek) and a K3 detector (Ametek) during 2.4

seconds with a dose rate of 4.1 e-/px/s resulting in a total dose of 38 $e^-/Å^2$ fractionated into 40

movie frames. Motion correction, CTF-estimation, fourier binning (to 1.02 Å/px), picking and

extraction were performed on the fly using Warp [29].

A total of 9,584 micrographs were and 350,022 particles were picked by Warp. Extracted particles

were imported into cryoSPARC v2.15.0 [30] for 2D classification, 3D classification and non-

uniform 3D refinement. The particles were processed with C1 symmetry throughout. After 2D

classification (300 classes) 219,491 particles were retained and used to build three ab-initio 3D

reconstructions. These were further processed for heterogenous refinement that resulted in one

reconstruction showing high-resolution structural features in the core of the spike. One round of

homogenous refinement followed by non-uniform refinement resulted in a final reconstruction to

an overall resolution of 2.9 Å (0.143 FSC) using 150,847 particles.

Model building and structure refinement

A structure of the 2019-nCoV spike protein trimer [14] (PDB: 6VSB) was used as a starting model

for model building. The model was extended and manually adjusted in COOT [31]. The Nanobody

structure was homology modelled using SWISS-MODEL [32] taking PDB:5JMR [33] as a

template. The missing regions of the RBD domains were built based on the RBD-Spike crystal

structure (PDB: 6LZG) [34]. Structure refinement and manual model building were performed

using COOT and PHENIX [35] in interspersed cycles with secondary structure and geometry

restrained. All structure figures and all EM density-map figures were generated with UCSF

ChimeraX [36].

Acknowledgements

SARS-CoV-2 virus was received from Jonas Klingström, KI. The authors thank James Voss

and Deli Huang for reagents. We thank PreClinics for their fast and efficient services and

Gunnar Schulte and Per-Åke Nygren for providing access to instruments for measuring binding

kinetics. We also thank Fondation Dormeur, Vaduz for generous support of flow cytometry

equipment.This work was supported by an EU grant (CoroNAb) to BM, GM and GHK and by

project grants from the Swedish Research Council to BM (2018-02381), BMH (2017-6702 and

2018-3808) and to GM (2018-03843).

The authors declare no conflict of interest

Bibliography

1.   Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, et al. Cell entry mechanisms of

SARS-CoV-2. Proc Natl Acad Sci U S A. 2020;117: 11727–11734.

2.   Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, et al. Cryo-EM

structure of the 2019-nCoV spike in the prefusion conformation. Science. 2020;367: 1260–

1263.

3.   Walls AC, Park Y-J, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and

Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell. 2020;181: 281–292.e6.

4.   Ju B, Zhang Q, Ge J, Wang R, Sun J, Ge X, et al. Human neutralizing antibodies elicited by

SARS-CoV-2 infection. Nature. 2020. doi:10.1038/s41586-020-2380-z

5.   Shi R, Shan C, Duan X, Chen Z, Liu P, Song J, et al. A human neutralizing antibody targets

the receptor binding site of SARS-CoV-2. Nature. 2020. doi:10.1038/s41586-020-2381-y

6.   Cao Y, Su B, Guo X, Sun W, Deng Y, Bao L, et al. Potent neutralizing antibodies against

SARS-CoV-2 identified by high-throughput single-cell sequencing of convalescent patients'

B cells. Cell. 2020. doi:10.1016/j.cell.2020.05.025

7.   Bannas P, Hambach J, Koch-Nolte F. Nanobodies and Nanobody-Based Human Heavy

Chain Antibodies As Antitumor Therapeutics. Front Immunol. 2017;8: 1603.

8.   Vincke C, Loris R, Saerens D, Martinez-Rodriguez S, Muyldermans S, Conrath K. General

strategy to humanize a camelid single-domain antibody and identification of a universal

humanized nanobody scaffold. J Biol Chem. 2009;284: 3273–3284.

9.   Detalle L, Stohr T, Palomo C, Piedra PA, Gilbert BE, Mas V, et al. Generation and

Characterization of ALX-0171, a Potent Novel Therapeutic Nanobody for the Treatment of

Respiratory Syncytial Virus Infection. Antimicrob Agents Chemother. 2016;60: 6–13.

10. Cardoso FM, Ibañez LI, Van den Hoecke S, De Baets S, Smet A, Roose K, et al. Single-

domain antibodies targeting neuraminidase protect against an H5N1 influenza virus

challenge. J Virol. 2014;88: 8278–8296.

11. Hanke L, Knockenhauer KE, Brewer RC, van Diest E, Schmidt FI, Schwartz TU, et al. The

Antiviral Mechanism of an Influenza A Virus Nucleoprotein-Specific Single-Domain

Antibody Fragment. MBio. 2016;7. doi:10.1128/mBio.01569-16

12. Kirchhofer A, Helma J, Schmidthals K, Frauer C, Cui S, Karcher A, et al. Modulation of

protein properties in living cells using nanobodies. Nat Struct Mol Biol. 2010;17: 133–138.

13. Pollard TD. A guide to simple and informative binding assays. Mol Biol Cell. 2010;21:

4061–4067.

14. Wrapp D, De Vlieger D, Corbett KS, Torres GM, Wang N, Van Breedam W, et al. Structural

Basis for Potent Neutralization of Betacoronaviruses by Single-Domain Camelid Antibodies.

Cell. 2020;181: 1004–1015.e15.

15. Monteil V, Kwon H, Prado P, Hagelkrüys A, Wimmer RA, Stahl M, et al. Inhibition of SARS-

CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human

ACE2. Cell. 2020;181: 905–913.e7.

16. Antos JM, Ingram J, Fang T, Pishesha N, Truttmann MC, Ploegh HL. Site-Specific Protein

Labeling via Sortase-Mediated Transpeptidation. Curr Protoc Protein Sci. 2017;89: 15.3.1–

15.3.19.

17. Witte MD, Theile CS, Wu T, Guimaraes CP, Blom AEM, Ploegh HL. Production of

unnaturally linked chimeric proteins using a combination of sortase-catalyzed

transpeptidation and click chemistry. Nat Protoc. 2013;8: 1808–1819.

18. Sosa BA, Demircioglu FE, Chen JZ, Ingram J, Ploegh HL, Schwartz TU. How lamina-

associated polypeptide 1 (LAP1) activates Torsin. Elife. 2014;3: e03239.

19. Alpaca (Lama pacos) as a convenient source of recombinant camelid heavy chain

antibodies (VHHs). J Immunol Methods. 2007;324: 13–25.

20. Edgar RC, Flyvbjerg H. Error filtering, pair assembly and error correction for next-

generation sequencing reads. Bioinformatics. 2015;31: 3476–3482.

21. Kumar V, Vollbrecht T, Chernyshev M, Mohan S, Hanst B, Bavafa N, et al. Long-read

amplicon denoising. Nucleic Acids Res. 2019;47: e104.

22. Rogers TF, Zhao F, Huang D, Beutler N, Burns A, He W-T, et al. Rapid isolation of potent

SARS-CoV-2 neutralizing antibodies and protection in a small animal model. bioRxiv. 2020.

p. 2020.05.11.088674. doi:10.1101/2020.05.11.088674

23. Sarzotti-Kelsoe M, Bailer RT, Turk E, Lin C-L, Bilska M, Greene KM, et al. Optimization and

validation of the TZM-bl assay for standardized assessments of neutralizing antibodies

against HIV-1. J Immunol Methods. 2014;409: 131–146.

24. Abdiche Y, Malashock D, Pinkerton A, Pons J. Determining kinetics and affinities of protein

457   interactions using a parallel real-time label-free biosensor, the Octet. Anal Biochem.

458   2008;377: 209–217.

459  25. Octet Systems Software \textbar ForteBio. Available:

460    https://www.fortebio.com/products/octet-systems-software

461  26. Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer; 2016.

462  27. Wickham H. tidyverse: Easily Install and Load the 'Tidyverse. R package version 1.2.1;

463    2017. Available: http://CRAN.R-project.org/package=tidyverse

464  28. Team RC. R: A language and environment for statistical computing. Vienna, Austria: R

465    Foundation for Statistical Computing; Available: http://www.R-project.org/

466  29. Tegunov D, Cramer P. Real-time cryo-electron microscopy data preprocessing with Warp.

467    Nat Methods. 2019;16: 1146–1152.

468  30. Punjani A, Rubinstein JL, Fleet DJ, Brubaker MA. cryoSPARC: algorithms for rapid

469    unsupervised cryo-EM structure determination. Nat Methods. 2017;14: 290–296.

470  31. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. Acta

471    Crystallogr D Biol Crystallogr. 2010;66: 486–501.

472  32. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al. SWISS-

473    MODEL: homology modelling of protein structures and complexes. Nucleic Acids Res.

474    2018;46: W296–W303.

475  33. Dahms SO, Creemers JWM, Schaub Y, Bourenkov GP, Zögg T, Brandstetter H, et al. The

476    structure of a furin-antibody complex explains non-competitive inhibition by steric exclusion

477    of substrate conformers. Sci Rep. 2016;6: 34303.

34. Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, et al. Structural and Functional Basis of

SARS-CoV-2 Entry by Using Human ACE2. Cell. 2020;181: 894–904.e9.

35. Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, et al. PHENIX: a

comprehensive Python-based system for macromolecular structure solution. Acta

Crystallogr D Biol Crystallogr. 2010;66: 213–221.

36. Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al. UCSF

ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci. 2018;27:

14–25.

37. Myszka DG, Jonsen MD, Graves BJ. Equilibrium analysis of high affinity interactions using

BIACORE. Anal Biochem. 1998;265: 326–330.

Figure 1 - Nanobody discovery.

A - Overview of nanobody generation process.

B - Variant frequencies quantified by NGS across successive enrichment steps. Identified using

RBD bait, Ty1 exhibits the greatest total fold change of all nanobodies, increasing in proportion

over 10,000-fold between initial and final libraries.

C - Sequence of Ty1. Complementarity-determining regions (CDRs) are indicated. Color

scheme according to ClustalX.

Figure 2 - Nanobody Ty1 neutralizes SARS-CoV-2 and specifically recognizes SARS-CoV-

2 spike glycoprotein.

A - VSV G or SARS-CoV-2 spike pseudotyped lentivirus was incubated with a dilution series of

Ty1 or control nanobody (influenza NP-VHH1 [11]). Infectivity relative to cells infected with

pseudotyped virus in the absence of nanobody is shown. Neutralization by Ty1 was repeated in

duplicate across 6 assays, and the error bars represent the standard deviation.

B - Cells were transfected with a plasmid harboring the SARS-CoV-2 spike for 24 h. Cells were

fixed, permeabilized and stained with Ty1-AS635P (black and red) or left unstained (grey). Cells

were analyzed by flow cytometry. Cell counts are presented as % of max (representative

histogram).

C - Vero E6 cells were infected with SARS-CoV-2 at a MOI of 1 for 24 hours. Cells were fixed,

permeabilized and stained for DNA (blue), dsRNA (green) and with Ty1-AS635P (red). Pictures

were taken by fluorescence microscopy and representative examples are shown.

D - ACE2 expression 293T cells were trypsinized, fixed and stained with RBD-AS635P alone, or

preincubated with IAV NP-VHH1 or Ty1. Cells were analyzed by flow cytometry.

533
534 **Figure 3 - RBD bound to surface-immobilized Ty1 with high affinity in the 5-50 nM range**
535 **in Biolayer Interferometry**
536
A - RBD bound to surface-immobilized Ty1 (red curves), but not to NP-VHH1 (violet curves) or to

biocytin-quenched Streptavidin (SAX, green curves). Almost equal nanobody immobilization

levels of about 0.7 nm were obtained by first loading Ty1 and then NP-VHH1. The SAX control

surface (green) was quenched, but not loaded with nanobodies. During association, nanobody

sample and control SAX sensors (red, violet and green) were dipped into RBD, while the reference

nanobody sensors were dipped into PBST (blue and light-green). For dissociation, all sensors

were dipped into PBST. Binding of molecules over time is recorded as sensorgrams recording

the shift in wavelengths (unit: nm) due to an increase in the optical thickness of the surface layer.

B - *(left)* RBD titration sensorgrams reveal concentration-dependent responses with an

association phase comprising a fast and a slow component. Dissociation seems to follow a single-

exponential decay but does not return to baseline over the measurement time. Fitted 1:1 binding

models are shown as thin lines. *(middle)* Pseudo-equilibrium response values extracted from the

final association phase were plotted against the logarithmic Ty1 concentration. Single-site

interaction fits of the sigmoidal binding curves according to [37] yielded K_D-values in the low nM

range. K_D-values and standard deviations are shown as solid and dotted lines, respectively. *(right)*

Sensor immobilization levels are shown as jittered box plots.

C - *(left)* Residuals of the 1:1 model fits to the interaction sensorgrams revealed that the simple

1:1 interaction model did not fully explain the obtained data, especially association phases at

higher RBD concentrations. Color coding as in the sensorgram above. *(right)* Plot of fitted

association rate constants (kon) versus affinity (K_D) values obtained from the kinetic 1:1 fits. Most

of the fits grouped together at K_D values of 10-100 nM with association and dissociation constants

of about 1-1.2 $*10^5$ (Ms)$^{-1}$ and 3.5–4.3$*10^{-3}$ s^{-1}, respectively.

Figure 4 - Structure of the SARS-CoV-2 spike in perfusion conformation in complex with

neutralizing nanobody Ty1

Ty1 binds to the RBD in 'up' and 'down' conformation and prevents ACE2 engagement

A - Cryo-EM reconstruction to an overall resolution of 2.9 Å of the spike trimer with bound Ty1

B - Surface representation of the spike trimer with three molecules of Ty1 bound. In our

reconstruction, one RBD takes the 'up' and two RBDs have the 'down' conformation. Ty1 is

bound on all three RBDs.

D - Cartoon representation of the spike trimer in complex with three molecules of Ty1.

E - Ty1 shows a two-pronged inhibition of ACE2 receptor binding through binding the RBD in

the 'up' conformation and by binding to the neighboring RBD in the 'down' conformation.