Resuscitation of the microbial seed bank alters plant-soil interactions

Venus Kuo, Brent K. Lehmkuhl, Jay T. Lennon[*]

Affiliations

Department of Biology, Indiana University, Bloomington, IN 47405

*Correspondence to: lennonj@indiana.edu

Key words: *Brassica rapa*, dormancy, plant-microbe interactions, microbiome, resuscitation promoting factor.

ABSTRACT

While microorganisms are recognized for driving belowground processes that influence

12 the productivity and fitness of plant populations, the vast majority of bacteria and fungi in soil

belong to a "seed bank" made up of dormant individuals. Still, plant performance may be

14 affected by microbial dormancy through its effects on the activity, abundance, and diversity of

soil microorganisms. To test how microbial seed banks influence plant-soil interactions, we

16 purified recombinant resuscitation promoting factor (Rpf), a bacterial protein that terminates

dormancy. Then, in a factorially designed experiment, we applied the Rpf to soil containing field

18 mustard (*Brassica rapa*), an agronomically important plant species. We found that plant biomass

was ~33 % lower in the Rpf treatment compared to plants growing in the presence of an

20 unmanipulated microbial seed bank. In addition, Rpf reduced soil respiration, decreased bacterial

abundance, and increased fungal abundance. These effects of Rpf on plant performance were

22 accompanied by shifts in bacterial community composition, which may have diluted mutualists

or resuscitated pathogens. Our findings suggest that changes in microbial seed banks may

24 influence the magnitude and direction of plant-soil feedbacks in ways that affect above- and

below-ground biodiversity and function.

INTRODUCTION

Belowground soil microbial communities play a critical role in determining the productivity and fitness of aboveground plants. Plant roots are intimately associated with thousands of bacterial and fungal taxa that are involved in soil processes such as decomposition, nutrient cycling, and pathogen suppression (Bardgett & van der Putten, 2014; Berendsen, Pieterse, & Bakker, 2012; Wagg, Bender, Widmer, & van der Heijden, 2014). In addition to containing mutualistic symbionts, soils also harbor pathogenic microorganisms that reduce plant performance (Mansfield et al., 2012). Together, belowground microbial communities are responsible for generating plant-soil feedbacks that can influence the diversity and composition of plant communities (Bever, 2003). However, the direction and magnitude of soil microbial effects on plants is highly variable (Hoeksema et al., 2010; Kulmatiski, Beard, Stevens, & Cobbold, 2008) and is likely influenced by the complexity of belowground communities (van der Heijden, de Bruin, Luckerhoff, van Logtestijn, & Schlaeppi, 2016; Wagg et al., 2014).

One belowground feature that is commonly overlooked when considering plant-microbe interactions is the metabolic heterogeneity of soil microbial communities. Many microorganisms are capable of entering a reversible state of reduced metabolic activity (i.e., dormancy) when challenged by suboptimal environmental conditions (Lennon & Jones, 2011). A large proportion ($\geq$90 %) of microorganisms are dormant in soils (Alvarez, Alvarez, Grigera, & Lavado, 1998; Blagodatsky, Heinemeyer, & Richter, 2000; Lennon & Jones, 2011). Collectively, these inactive individuals create a "seed bank", which not only maintains biodiversity, but also affects ecosystem functioning. The incorporation of microbial dormancy into ecosystem experiments and models has been shown to improve predictions for soil microbial activity and nutrient

cycling (Salazar, Lennon, & Dukes, 2019; Wang et al., 2015). Few studies, however, have explored how soil microbial seed banks influence plant-soil interactions.

Microbial seed banks may prevent local extinctions when exposed to fluctuating and stressful conditions that are typical in soil habitats (Shoemaker & Lennon, 2018). If microbial mutualists can persist by entering a seed bank, then this may promote beneficial soil functions such as disease-suppression, induction of plant immune response, stimulation of root growth, and biofertilization, which together, can increase plant performance (Lugtenberg & Kamilova, 2009; Mendes et al., 2011; Peralta, Sun, McDaniel, & Lennon, 2018; Wagg et al., 2014). In contrast, seed banks may contain pathogenic microorganisms that can reduce plant performance. For example, it is well known that destructive pathogens can reside in soil for months to years after the demise of plants, rendering potentially arable soil unusable (Koike, Subbarao, Davis, & Turini, 2003; Peeters, Guidot, Vailleau, & Valls, 2013). Although microbial seed banks likely contain both mutualistic and pathogenic taxa, the net effect of dormancy transitions on plant performance remains to be determined.

Resuscitation is an essential process that regulates microbial seed-bank dynamics. Dormancy is often terminated via the interpretation of environmental cues or communication signals that are produced by other microorganisms (Dworkin & Shah, 2010). One such signal is a bacterial protein called resuscitation promoting factor (Rpf), a muralytic enzyme that degrades the β-(1,4) glycosidic bond in peptidoglycan, which is a major cell-wall component of virtually all bacteria (Mukamolova, Kaprelyants, Young, Young, & Kell, 1998). Rpf genes are broadly distributed among the G+C-rich Gram-positive Actinobacteria (Ravagnani, Finan, & Young, 2005; Schroeckh & Martin, 2006). In some soil samples, Rpf homologs can be found in up to 25 % of all genomes (Lennon & Jones, 2011). Rpf has cross-species effects that stimulate the

72 growth of closely related dormant bacteria (Puspita, Kitagawa, Kamagata, Tanaka, & Nakatsu,

2015; Schroeckh & Martin, 2006) either through its direct effects on cell wall integrity or

74 through the release of small peptide fragments that cross-link peptidoglycan, which may serve as

signaling molecules (Jonathan Dworkin, 2014). Under laboratory conditions, Rpf can stimulate

76 the activity of some microbial taxa, but it has also been shown to inhibit the growth of other

microorganisms (Mukamolova et al., 1998; Mukamolova et al., 2002), which is not surprising

78 given that Rpf is homologous to lysozyme (Cohen-Gonsaud et al., 2005), an antimicrobial

enzyme produced by many eukaryotic organisms. Overall, Rpf may have complex and

80 interactive effects that influence plant-soil interactions.

In this study, we explored how seed-bank dynamics affect plant-microbe interactions by

82 manipulating the process of resuscitation. Throughout the lifespan of a host plant (*Brassica

rapa*), we applied purified recombinant Rpf to soils containing complex microbial communities.

84 The experiment was guided by a conceptual model that considers how active and inactive

microorganisms interact and potentially create feedbacks with plants (Fig. 1). We hypothesized

86 that resuscitating the microbial seed bank with Rpf would alter plant performance through

changes in the abundance, activity, and composition of soil microbial communities. On the one

88 hand, Rpf treatment could enhance plant performance by waking up mutualistic taxa in the soil

microbial community. On the other hand, resuscitating dormant microorganisms could

90 negatively influence plant performance by diluting the benefits of mutualistic taxa or through the

recruitment of pathogenic microorganisms. Here, we test these hypotheses while also

92 considering other effects of Rpf on plants, microbes, and their interactions.

METHODS

96 ***Experimental design*** — We conducted a growth chamber experiment to test the effect of Rpf on

plant-soil interactions using *Brassica rapa* L. (Brassicaceae), an economically important plant

98 that consists of a variety of cultivated subspecies. We sowed 16 *B. rapa* seeds into individual

pots in June 2016. Seeds used in this experiment were obtained from Wisconsin Fast Plants[TM]

100 (Standard stock seeds, Wisconsin Fast Plants Program, University of Wisconsin, Madison, WI,

USA). While *B. rapa* is non-mycorrhizal, it still associates with a diverse microbial community

102 that influences plant growth (Lau & Lennon, 2012). For our experiment, we used plastic pots (17

cm diameter, 13 cm height) filled with autoclaved (121 °C, 15 PSI, 16 h) soil medium (1 part

104 greenhouse metro-mix: 1 part vermiculite), which was inoculated with live field soil collected

from Indiana University Research and Teaching Preserve (DMS 39 °11'56.5"N 86 °30'40.5"W).

106 For the +Rpf treatment, we added 1 mL of recombinant Rpf protein (10 µM) to the soil surface

each week in three equidistant locations 5 cm away from main plant stem. The same was done

108 for the -Rpf treatment (negative control), except we used protein buffer (20 mM Tris-HCl, 100

mM NaCl) instead of recombinant Rpf. Our experiment was conducted in a Percival model

110 PGC-15 growth chamber with high efficiency light lamps (Philips series 700 32 watts 4100K

F32T8/TL741 AltoII). The growth chambers were set to constant and full light intensity (1100

112 $\mu mol\ m^{-2}\ s^{-1}$) with controlled humidity (60 %) and temperature (28 °C). Pots were arranged in a

fully randomized design where each plant was assigned to a resuscitation treatment (i.e., -Rpf vs.

114 +Rpf). All pots were watered with equal amounts of filtered and de-ionized water every other

day.

Recombinant Rpf — To manipulate microbial seed banks, we overexpressed Rpf from

118   *Micrococcus luteus* sp. KBS0714, a bacterial strain isolated from agricultural soil at the W.K.

Kellogg Biological Station, Hickory Corners, MI (MSU) (Kuo, Shoemaker, Muscarella, &

120   Lennon, 2017; Lennon, Aanderud, Lehmkuhl, & Schoolmaster Jr, 2012). *Micrococcus luteus* sp.

KBS0714 is a close relative of *M. luteus* NCTC 2662 (99 % sequence 16S rRNA sequence

122   similarity, NCBI CP001628.1), a model organism used for studying Rpf (Mukamolova et al.,

1998). To create a Rpf expression host, we amplified and cloned the *rpf* gene from KBS0714

124   into the expression vector pET15b. First, we extracted KBS0714 genomic DNA from pure

culture using a Microbial DNA isolation kit (MoBio, Carlsbad, CA) for PCR amplification of the

126   open reading frame of the *rpf* gene using two primers, Upper-F 5' GCC CAT ATG GCC ACC

GTG GAC ACC TG 3' and Lower-R 5' GGG GAT CCG GTC AGG CGT CTC AGG 3', with

128   incorporated restriction sites EcoRI, NdeI (forward primer) and BamHI (reverse primer)

(Koltunov et al., 2010; Mukamolova et al., 1998). We amplified the *rpf* gene sequence with the

130   following PCR conditions: initial: 95 °C for 5 m, 30 cycles of 95 °C for 30 s, 55 °C for 30 s, 72

°C for 1 m, and one final extension at 72 °C for 7 m. The *rpf* gene amplicon was ligated into

132   pET15b (Invitrogen) as an EcoRI/BamHI fragment and transformed into *Escherichia coli* TOP10

(a non-expression host). We then identified positive clones with correct *rpf* sequence and

134   orientation by Sanger sequencing at the Indiana University Center of Genomics and

Bioinformatics (IU-CGB). Next, we extracted purified pET15b plasmids with the *rpf* gene from

136   transformed TOP10 *E. coli* using the QIAprep Spin Miniprep Kit (Qiagen) following the

manufacturer's protocol. Finally, we transformed the purified pET15b expression vector with the

138   *rpf* gene into the *E. coli* Origami BL21 (DE3) expression host incorporated with a polyhistidine-

tag on the N-terminus of the recombinant protein.

To overexpress Rpf, we grew the *E. coli* expression host in Lysogeny Broth (LB) with appropriate antibiotics (ampicillin 100 µg/mL, kanamycin 15 µg/mL, and tetracycline 12.5

µg/mL). During logarithmic growth, recombinant protein production was induced with Isopropyl β-D-1-thiogalactopyranoside (IPTG) (100 µM final concentration). We confirmed

overexpression of Rpf with Western blots (Fig. S1). Cells were then collected by centrifugation, lysed by sonication, and filtered through a Ni-NTA Purification System (Invitrogen) using a 10

mL gravity fed column with a 2 mL resin bed to purify recombinant Rpf with the N-terminus polyhistidine-tag. Recombinant Rpf protein was washed with 5 mM imidazole buffer (300 mM

NaCl, 50 mM Tris-HCl, 5 mM imidazole) and then eluted with 125 mM imidazole buffer. Rpf protein was purified by buffer exchange using a 10 mL Zeba Spin Desalting Columns (Thermo

Fisher) with protein buffer (20 mM Tris-HCl, 100 mM NaCl) following manufacturer's instructions and then passed through a 0.2 µm syringe filter before adding to soil as described

above.

<u>Plant responses</u> — We censused the plants every day and recorded germination and flowering date. After most plants had finished flowering, we documented flower number, plant height, and

specific leaf area (SLA = leaf area/dry mass) as previously described (Lau & Lennon, 2012). All open flowers in each experimental unit were hand-pollinated by other open flowers in the same

treatment group with a soft paint brush, which was cleaned with 30 % isopropyl alcohol between treatment groups to prevent gene flow. Plants and seeds were harvested at the end of the six-

week growing period when most individuals ceased flowering and had begun to senesce. As an estimate of female fitness, we removed plant seed pods and counted the number of seeds

produced. To estimate male fitness, we counted the number of flowers per plant. In addition, we

measured total, aboveground, and belowground biomass of each plant after drying plants at 65

164 °C for 2 d. We performed two-tailed Student's *t*-test to evaluate the effect of Rpf on total

biomass, aboveground (i.e., shoot) biomass, root biomass, root : shoot ratio, SLA, and plant

166 fitness (i.e., flower and seed numbers).

168 ***Soil microbial activity*** —We measured soil respiration on a weekly basis to determine the effect

of Rpf treatment on belowground microbial activity. For each pot, we transferred 1 g of soil into

170 glass vials with a silicon membrane septum cap. After incubating the soil samples at 25 °C in the

dark for 24 h, we measured the CO_2 concentration (ppm) from 1 mL of headspace gas using an

172 infrared gas analyzer (IRGA) (LI-COR Environmental). We calculated of CO_2 concentration in

our samples based on values generated from a standard curve of known CO_2 concentrations. We

174 performed repeated-measures ANOVA (RM-ANOVA) using an AR(1) covariance structure to

test for the main effects and interaction of Rpf and time on rates of soil respiration.

Microbial abundance — We estimated bacterial and fungal abundance using quantitative PCR

178 (qPCR) on DNA extracted from soil samples. For each pot, we collected 1 g of soil pooled from

three subsamples and immediately stored the soil samples at -80 °C. We extracted the DNA from

180 1 g of soil using a PowerSoil® Total RNA Isolation Kit with DNA Elution Accessory Kit

(MoBio) following manufacturer's protocol, and the DNA was subsequently quantified using a

182 Take5 *Synergy* microplate reader (BioTek). We then conducted qPCR assays (Lau & Lennon,

2011) that consisted of 20 μL reactions containing 1 μL of DNA template (2.5 ng μL^{-1}), 1 μL of

184 each primer (10 μmol L^{-1}), 7 μL of molecular-grade water, and 10 μL iQTM SYBR Green

SuperMix (Bio-Rad Laboratories Hercules, CA, USA). For bacteria we used Eub338 (forward)

186   and Eub518 (reverse) primers to amplify the 16S rRNA gene, and for fungi we used ITS1f

(forward) and 5.8S (reverse) primers to amplify the ITS gene region (Fierer, Jackson, Vilgalys,

188   & Jackson, 2005). qPCR assays were performed using Eppendorf Mastercycler Realplex system

using previously reported thermal cycle conditions (Fierer et al., 2005). The coefficients of

190   determination (r^2) of our assay ranged from 0.95 and 1, while amplification efficiencies fell

between 0.93 and 0.99. Based on melting curve analyses, we found no evidence for primer

192   dimers. We estimated fungal and bacterial abundance based on the estimated gene copy number

from their respective standard curves generated from bacterial and fungal isolates as described

194   elsewhere (Lau & Lennon, 2012). We performed a two-tailed Student's *t*-test to determine the

effect of the Rpf treatment on soil bacterial (16S rRNA) and fungal (ITS) gene copy abundances,

196   as well as the fungal to bacterial ratio (F : B).

198   ***Microbial structure*** — To account for the variation in metabolic activity, we characterized soil

bacterial communities using RNA and DNA. DNA is a relatively stable molecule contained in

200   intact cells irrespective of their metabolic status. Accordingly, we interpret 16S rRNA sequences

recovered from the DNA pool as the "total" community. In contrast, RNA is a more ephemeral

202   molecule that is required for protein synthesis by actively growing cells. Therefore, we interpret

16S rRNA sequences recovered from the RNA pool after complementary DNA (cDNA)

204   synthesis as the "active" community (Jones & Lennon, 2010). Following protocols outlined

elsewhere (Lennon, Placella, & Muscarella, 2017), we characterized the effects of Rpf on soil

206   bacterial communities using high-throughput sequencing. We amplified the V4 region of the 16S

rRNA gene from bacterial DNA and cDNA to characterize total and active communities,

208   respectively, using barcoded primers (515F and 806R) designed to work with the Illumina MiSeq

platform (Caporaso et al., 2012). We removed residual DNA from RNA samples using DNase 1

210   (Invitrogen) following manufacturer's protocol and synthesized cDNA by reverse transcribing

RNA using iScript Reverse Transcription Supermix for RT-qPCR (Bio-Rad). We cleaned the

212   sequence libraries using the Agencourt AMPure XP purification kit (Beckman Coulter, Brea,

CA, USA), quantified the resulting product using the QuanIt PicoGreen kit (Invitrogen), and

214   pooled libraries at equal molar ratios (final concentration: 10 ng per library). We then sequenced

the pooled libraries with the Illumina MiSeq platform using paired end reads (Illumina Reagent

216   Kit v2, 500 reaction kit) at the IU-CGB. Paired-end raw 16S rRNA sequence reads were

assembled into contigs using the Needleman algorithm. After quality trimming with a moving

218   average quality score (window 50 bp, minimum quality score 35), we aligned the sequences to

the Silva Database (v 128) using the Needleman algorithm. Chimeric sequences were removed

220   using the UCHIME algorithm (Edgar, Haas, Clemente, Quince, & Knight, 2011). We created

operational taxonomic units (OTUs) by first splitting the sequences based on taxonomic class

222   (using the RDP taxonomy) and then binning sequences into OTUs based on 97 % sequence

similarity. All sequence processing was completed using the software package *mothur* (version

224   1.39.5) (Schloss et al., 2009).

     Using the 16S rRNA sequences, we determined the effect of Rpf treatment on bacterial

226   richness (i.e., the number of OTUs in a sample) using a resampling approach that subsampled

1,000 sequence observations per sample, resampled 999 additional times, and then calculated the

228   average number of OTUs to estimate per sample richness ($\pm$ SEM) (Muscarella, Jones, &

Lennon, 2016). Similarly, we estimated the effect of Rpf treatment on taxa evenness (i.e., the

230   equitability in relative abundance of OTUs in a sample) using the same resampling approach

with Simpson's index. We then determined the effect of Rpf treatment on the total and active

232   bacterial pools using multivariate statistical procedures. Specifically, we conducted a Principal

Coordinates Analysis (PCoA) with Bray-Curtis dissimilarity metric using $\log_{10}$-transformed

234   relative abundances to visualize the effects of Rpf on bacterial communities. To test the

hypothesis that Rpf affected the structure of the total (DNA) and active (RNA) bacterial pools,

236   we used permutational multivariate analysis of variance (PERMANOVA) with Bray-Curtis

dissimilarity metric implemented with the 'adonis' function in the *vegan* package (Oksanen et

238   al., 2011) in the R statistics environment (v 3.2.3). In the PERMANOVA, we coded pot number

as a factor to account for paired-sample design given that DNA and RNA were co-extracted from

240   the same soil sample within each experimental unit. To identify influential taxa driving structural

changes in response to the Rpf treatments, we calculated the correlation coefficient of each OTU

242   along PCoA axes using the 'add.spec.scores' function in the *BiodiversityR* package (Kindt, 2016)

and implemented a coefficients cut-off ($> |0.7|$) (Fowler, Cohen, & Jarvis, 2013). We then

244   performed a permutation test (n = 999) to evaluate the correlations of each OTU along PCoA

axes using the 'envfit' function on the *vegan* package (Oksanen et al., 2011).

RESULTS

248   ***Plant responses*** — Plants in +Rpf treatment had less total ($t_7 = 2.84$, $P = 0.013$), aboveground (t_7

$= 2.70$, $p = 0.017$), and belowground biomass ($t_7 = 2.30$, $P = 0.049$) than plants in the -Rpf

250   treatment (Fig. 2). While recombinant protein addition led to a ~33 % reduction in root biomass,

Rpf no effect on other plant traits including SLA ($t_7 = 0.80$, $P = 0.442$), root : shoot ratio ($t_7 =$

252   -1.20, $P = 0.250$), and shoot height ($t_7 = 1.11$, $P = 0.286$). Similarly, there was no effect of Rpf

on the number of seeds produced per plant ($t_7 = 0.83$, $P = 0.421$), but we detected a marginal

254 decrease in the number of flowers produced per plant in the +Rpf treatment ($t_7 = 1.79$, $P = 0.097$).

__Microbial activity__ — Rpf altered the activity of the belowground microbial community based on

258 measurements of soil respiration. We detected a significant main effect of Rpf ($F_{1, 84} = 9.68$, $P = 0.002$) and time ($F_{5, 84} = 21.60$, $P < 0.001$), but no interaction ($F_{5, 84} = 1.22$, $P = 0.308$), on soil

260 respiration (Fig. 3). While soil respiration increased throughout the experiment in both

treatments, based on marginal means, soil respiration was 24 % lower in the +Rpf treatment than

262 in the -Rpf treatment.

264 *__Microbial abundance__* — The Rpf treatment significantly altered the abundance of soil bacteria

($t_8 = 2.71$, $P = 0.016$) and fungi ($t_8 = -3.27$, $P = 0.007$) relative to the control (Fig. 4). Bacterial

266 abundance estimated as 16S rRNA gene copy number decreased by ~30 % in the +Rpf treatment

relative to the -Rpf treatment. In contrast, fungal abundance estimated as the ITS gene copy

268 number increased 2.8-fold in the +Rpf treatment. Consequently, soil F : B ratio increased 4.8-

fold under +Rpf treatment relative to the control treatment ($t_8 = -2.84$, $P = 0.018$).

__Bacterial structure__ — Since Rpf production is thought to be restricted to the Actinobacteria, we

272 hypothesized that Rpf treatment would disproportionally affect the diversity and composition of

taxa belonging to this group. Therefore, we estimated alpha- and beta-diversity of 16S rRNA

274 sequences belonging to the Actinobacteria phylum. Rpf had no effect on the OTU richness of

Actinobacteria regardless of whether sequences came from the total ($F_{1, 8} = 0.80$, $p = 0.397$) or

276 active community ($F_{1, 8} = 0.14$, $p = 0.722$) (Table S1). Similarly, Rpf had no effect on the

evenness of the total ($F_{1, 8} = 0.69$, $p = 0.429$) or active ($F_{1, 8} = 0.41$, $p = 0.538$) actinobacterial

taxa (Table S1). We detected a small and marginally significant reduction in the relative

abundance of actinobacterial sequences in the +Rpf treatment relative to the -Rpf treatment ($t_9 =$

1.36, $P = 0.097$) (Fig. A2). However, PERMANOVA revealed that actinobacterial composition

was significantly affected by metabolic status (i.e., active vs total) ($F_{1, 19} = 16.08$, $r^2 = 0.33$, $P =$

0.001) and Rpf treatment ($F_{1, 19} = 3.25$, $r^2 = 0.07$, $p = 0.010$). The PERMANOVA also indicated

that actinobacterial composition was marginally influenced by an interaction between metabolic

status and Rpf treatment ($F_{1, 19} = 1.96$, $r^2 = 0.04$, $P = 0.055$). This interaction reflected an Rpf

effect on the active community (i.e., RNA) that was not evident in total (i.e., DNA) community

(Fig. 5). Based on this trend, we identified 17 actinobacterial taxa, all of which belonged to a

single order (*Solirubrobacterales*), in the active community that had positive ($r > 0.7$) and

significant ($p < 0.001$) correlations with both PCoA axis 1 and PCoA axis 2 (Table A2).

DISCUSSION

Dormancy is an important life-history trait that influences the diversity, composition, and

function of soil microbial communities. After generating recombinant protein with a gene from

an environmental isolate, we applied resuscitation promoting factor (Rpf) to soils to test the

hypothesis that microbial seed-bank dynamics alter plant growth and fitness traits by driving

changes in the belowground microbial community (Fig. 1). The Rpf treatment decreased plant

biomass (Fig. 2) most likely by altering the activity (Fig. 3), abundance (Fig. 4), and composition

(Fig. 5) of soil microbial communities. Our findings are consistent with the view that soil

microbial seed banks can influence plant performance by disrupting interactions with beneficial

microorganisms or through the recruitment of pathogens. In the following sections, we discuss

300   these findings while exploring other potential ways in which Rpf may affect plant-microbe

interactions in soil environments.

Rpf indirectly affects plant performance — Resuscitation of microbial seed banks led to a 33 %

304   reduction in plant biomass (Fig. 2). For the following reasons, we infer that Rpf effects on plant

performance is likely to be indirect. Rpf hydrolyzes glyosidic linkages in peptidoglycan (Cohen-

306   Gonsaud et al., 2005; Mukamolova et al., 2006).While this polymer is found in the cell walls of

virtually all bacteria, it is absent from plant tissues. Nevertheless, we conducted an experiment

308   with *Arabidopsis thaliana*, a close relative of *B. rapa*, to evaluate whether Rpf can directly affect

plant performance. We grew seedlings on microbe-free agar while being exposed to recombinant

310   Rpf at concentrations similar to those in our plant-soil experiment with *B. rapa*. A lack of an

effect (Fig. S5) led us to conclude that Rpf influenced plants primarily via its effect on the

312   activity, abundance, or composition of the soil microbial community.

314   ***Rpf altered fungal-bacterial interactions*** — Rpf may have altered plant performance by

modifying fungal-bacterial interactions. We observed a nearly three-fold increase in fungal

316   abundance in response to Rpf additions (Fig. 4). As a result, soil F : B ratio increased under Rpf

treatment (Fig. 4), which may also explain the reduction in soil microbial activity (Fig. 3) given

318   that there are often differences in carbon-use efficiency between bacteria and fungi (Sakamoto &

Oba, 1994; Whitaker et al., 2014). Shifts in F : B may also reflect the way that Rpf influences

320   microorganisms with different cell-wall properties. Like plants, fungi do not contain

peptidoglycan, the substrate which Rpf acts upon. Instead, the fungal cell wall is primarily

322   composed of chitin. Therefore, an increase in fungal abundance could arise if Rpf decreased the

competitive ability of soil bacteria in our system. In principle, the muralytic activity of Rpf may

324   have even lysed some bacterial cells. If this occurred, the resulting necromass could be

scavenged by fungi to meet their metabolic demands (Bradley, Amend, & LaRowe, 2018). Our

326   results, however, do not support this hypothesis. We found that bacterial abundance from three

different soils increased with Rpf concentration up to 4 µM g^{-1} soil. At higher concentrations (8

328   µM g^{-1}), we no longer observed a growth-stimulating effect, but bacterial abundance never

dropped below levels observed in control soils without Rpf (Fig. S6). Rpf was much lower (40

330   nM g^{-1} soil) in our *Brassica* experiment, yet above concentrations shown to resuscitate dormant

bacteria (Cohen-Gonsaud et al., 2005; Mukamolova et al., 1998). Based on these findings, it

332   seems unlikely that reductions in bacterial abundance (Fig. 4) were due to lysis or direct

inhibition of the soil microbial community.

__Rpf altered microbial structure__ – While seed banks are known to promote biodiversity by

336   buffering ecological and evolutionary dynamics, microbial dormancy may have complex effects

that influence plant-soil dynamics. The influence of Rpf on communities of interacting species is

338   likely dependent on the enzyme specificity. Very few studies have investigated Rpf effects on a

broad range of microorganisms, but some evidence suggests cross-species resuscitation using

340   recombinant Rpf generated from actinobacteria belonging to *Mycobacterium* and *Brevibacterium*

(Mukamolova et al., 1998; Puspita et al., 2015). In our study, we found that Rpf treatment had

342   broad-scale effects on bacterial communities. The composition of Actinobacteria (Fig. 5), Gram-

positive bacteria (Fig. A3), and the entire domain of sequenced Bacteria (Fig. A4) were affected

344   by the six-week application of recombinant Rpf. Treatment effects were also apparent when we

compared the composition of active Acintobacteria represented by sequences from the RNA

346   pool, with those recovered from the DNA pool, which includes individuals spanning the full

spectrum of metabolic activity (i.e., active, dormant, dead) (Lennon et al., 2018; Fig. 5). This

348   divergence in actinobacterial composition was due to changes in the relative abundance of 17

influential OTUs belonging to the *Solirubrobacterales* (Table A2). Members of this group are

350   aerobic, non-spore-forming, and non-motile bacteria that are recovered in forest and agricultural

soils (Kim et al., 2007; Reddy & Garcia-Pichel, 2009; Seki, Matsumoto, Ōmura, & Takahashi,

352   2015). Little is known about the role of *Solirubrobacterales* in the context of plant-microbe

interactions, but some representatives assimilate simple organic compounds (e.g., glucose,

354   maltose, sucrose, xylose, and arginine) that are commonly found in the rhizosphere (Reddy &

Garcia-Pichel, 2009). Interestingly, the *Solirubrobacterales* are only distantly related to

356   *Micrococcus* KBS0714, the soil isolate containing the Rpf gene that we used for producing the

purified recombinant protein. This suggests that Rpf functionality is not tightly restricted to close

358   kin. Additional work is needed to unveil the network of interactions between Rpf producers and

responders, in addition to whether or not the evolutionary history of Rpf genes maps onto

360   phylogenetic relationships of Actinobacteria.

362   ***Implications of seed-bank dynamics for plant-microbe interactions*** – Our study demonstrates

that resuscitation promoting factor (Rpf) modified plant performance through its effects on soil

364   microbial communities. Reductions in plant growth were accompanied by shifts in soil microbial

community properties, which may have included the recruitment of pathogens. Although we did

366   not observe root lesions or other clear signs of infection, we cannot rule out the possibility the

seed banks harbor pathogenic or non-mutualistic taxa that could decrease plant performance.

368   Such a view is consistent with reports of microbial pathogens persisting in soil for long periods

of time, likely in a state of reduced metabolic activity (Sharma & Reynnells, 2016). More

370 explicit tests would include the targeting of known pathogens combined perhaps with

measurements of plant immune responses to altered microbial seed banks. Alternatively, our

372 findings could be explained by a dilution of mutualists in the soil or a general disruption of

beneficial plant-microbe interactions, which would likely be reflected in the profiling of plant

374 exudates and microbial metabolites.

While our study provides strong empirical support for the notion that microbial seed

376 banks are important for plant performance, it also raises a number of questions for future

exploration. We observed that a one protein from a single strain of bacteria can have cascading

378 effects on plant-microbe interactions. However, some microorganisms may be responsive to

certain families of Rpf, while others may not. This likely is due to the mechanisms by which Rpf

380 operates in a community context, which is a topic that received little attention to date. More

generally, metabolic transitions into and out of dormancy are influenced by mechanisms besides

382 Rpf (Lennon & Jones, 2011) many of which are tied to the interpretation of environmental

triggers (e.g., soil rewetting) that are known to resuscitate microorganisms (Aanderud, Jones,

384 Fierer, & Lennon, 2015). Finally, there is a need to explore interactions with other plants.

Generalities should be sought by examining the effects of Rpf outside of the Brassicaceae given

386 that these plants tend to lack mycorrhizal symbionts, which are important mutualists found

among most flowering plants. It is not difficult to imagine that more complex outcomes could

388 emerge when altered microbial seed banks are considered in diverse plant communities (Fig. 1).

Nevertheless, our study highlights a potentially important but overlooked component of plant-

390 microbe interactions. It is estimated that soil microbial communities can be dominated by

dormant taxa, yet changes in activity can be fast, suggesting that seed banks may be an important

392 factor contributing to plant-soil feedbacks, which is thought to be an important mechanism

maintaining landscape patterns of biodiversity (Bever, 2003).

ACKNOWLEDGEMENTS

396 We acknowledge Peyton Thomas for assistance in the laboratory and members of the Lennon lab

for critical feedback on an earlier version of the manuscript. This work was supported by the

398 National Science Foundation 1442246 (JTL) and the US Army Research Office Grant

(W911NF- 14-1-0411 JTL). Data and code used in this manuscript can be found on GitHub

400 (https://github.com/LennonLab/BrassicaRpf), Zenodo (DOI: 10.XXXX/zenodo.XXXXX), and

NCBI (BioProject PRJNA504042).

REFERENCES

404 Aanderud, Z. T., Jones, S. E., Fierer, N., & Lennon, J. T. (2015). Resuscitation of the rare
biosphere contributes to pulses of ecosystem activity. *Frontiers in Microbiology, 6*, 24.

406 Alvarez, C., Alvarez, R., Grigera, M., & Lavado, R. (1998). Associations between organic matter
fractions and the active soil microbial biomass. *Soil Biology Biochemistry, 30*(6), 767-
408 773.

Bardgett, R. D., & van der Putten, W. H. (2014). Belowground biodiversity and ecosystem
410 functioning. *Nature, 515*(7528), 505-511.

Berendsen, R. L., Pieterse, C. M., & Bakker, P. A. (2012). The rhizosphere microbiome and
412 plant health. *Trends in Plant Science, 17*(8), 478-486.

Bever, J. D. (2003). Soil community feedback and the coexistence of competitors: conceptual
414 frameworks and empirical tests. *New Phytologist, 157*(3), 465-473.

Blagodatsky, S. A., Heinemeyer, O., & Richter, J. (2000). Estimating the active and total soil
416 microbial biomass by kinetic respiration analysis. *Biology Fertility of Soils, 32*(1), 73-81.

Bradley, J. A., Amend, J. P., & LaRowe, D. E. (2018). Necromass as a limited source of energy
418 for microorganisms in marine sediments. *Journal of Geophysical Research:
Biogeosciences, 123*(2), 577-590.

420 Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., . . .
Bauer, M. (2012). Ultra-high-throughput microbial community analysis on the Illumina
422 HiSeq and MiSeq platforms. *The ISME Journal, 6*(8), 1621-1624.

Cohen-Gonsaud, M., Barthe, P., Bagnéris, C., Henderson, B., Ward, J., Roumestand, C., & Keep,
424 N. H. (2005). The structure of a resuscitation-promoting factor domain from
Mycobacterium tuberculosis shows homology to lysozymes. *Nature Structural Molecular*
426 *Biology, 12*(3), 270.

Dworkin, J. (2014). The medium is the message: interspecies and interkingdom signaling by
428 peptidoglycan and related bacterial glycans. *Annual Review of Microbiology, 68*, 137-
154.

430 Dworkin, J., & Shah, I. M. (2010). Exit from dormancy in microbial organisms. *Nature Reviews*
Microbiology, 8(12), 890-896.

432 Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME improves
sensitivity and speed of chimera detection. *Bioinformatics, 27*(16), 2194-2200.

434 Fierer, N., Jackson, J. A., Vilgalys, R., & Jackson, R. B. (2005). Assessment of soil microbial
community structure by use of taxon-specific quantitative PCR assays. *Applied*
436 *Environmental Microbiology, 71*(7), 4117-4120.

Fowler, J., Cohen, L., & Jarvis, P. (2013). *Practical statistics for field biology*. Chichester, UK:
438 John Wiley & Sons.

Hoeksema, J. D., Chaudhary, V. B., Gehring, C. A., Johnson, N. C., Karst, J., Koide, R. T., . . .
440 Moore, J. C. (2010). A meta☐analysis of context☐dependency in plant response to
inoculation with mycorrhizal fungi. *Ecology Letters, 13*(3), 394-407.

442 Jones, S. E., & Lennon, J. T. (2010). Dormancy contributes to the maintenance of microbial
diversity. *Proceedings of the National Academy of Sciences, 107*(13), 5881-5886.

444 Kim, M. K., Na, J.-R., Lee, T.-H., Im, W.-T., Soung, N.-K., Yang, D.-C. J. I. j. o. s., &
microbiology, e. (2007). *Solirubrobacter soli* sp. nov., isolated from soil of a ginseng
446 field. *57*(7), 1453-1455.

Kindt, R. (2016). Biodiversity R: package for community ecology and suitability analysis. R
448 version 2.7.2.

Koike, S., Subbarao, K., Davis, R. M., & Turini, T. (2003). *Vegetable diseases caused by*
450 *soilborne pathogens*: UCANR Publications.

Koltunov, V., Greenblatt, C. L., Goncharenko, A. V., Demina, G. R., Klein, B. Y., Young, M., &
452 Kaprelyants, A. S. (2010). Structural changes and cellular localization of resuscitation-
promoting factor in environmental isolates of *Micrococcus luteus*. *Microbial Ecology,*
454 *59*(2), 296-310.

Kulmatiski, A., Beard, K. H., Stevens, J. R., & Cobbold, S. M. (2008). Plant-soil feedbacks: a
456 meta-analytical review. *Ecology Letters, 11*(9), 980-992.

Kuo, V., Shoemaker, W., Muscarella, M., & Lennon, J. (2017). Whole-genome sequence of the
458 soil bacterium *Micrococcus* sp. KBS0714. *Genome Announcements, 5*(32), e00697-
00617.

460   Lau, J. A., & Lennon, J. T. (2011). Evolutionary ecology of plant–microbe interactions: soil microbial structure alters selection on plant traits. *New Phytologist, 192*(1), 215-224.

462   Lau, J. A., & Lennon, J. T. (2012). Rapid responses of soil microorganisms improve plant fitness in novel environments. *Proceedings of the National Academy of Sciences of the United*
464   *States of America, 109*(35), 14058-14062. doi:10.1073/pnas.1202319109

        Lennon, J. T., Aanderud, Z. T., Lehmkuhl, B., & Schoolmaster Jr, D. R. (2012). Mapping the
niche space of soil microorganisms using taxonomy and traits. *Ecology, 93*(8), 1867-1879.

Lennon, J. T., & Jones, S. E. (2011). Microbial seed banks: the ecological and evolutionary implications of dormancy. *Nature Reviews Microbiology, 9*(2), 119-130.

Lennon, J. T., Placella, S. A., & Muscarella, M. E. (2018). Relic DNA contributes minimally to estimates of microbial diversity. *mBio 9:e00637-18*.

Lugtenberg, B., & Kamilova, F. (2009). Plant-growth-promoting rhizobacteria. *Annual Review of Microbiology, 63*, 541-556.

Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., . . . Machado, M. A. (2012). Top 10 plant pathogenic bacteria in molecular plant pathology. *Molecular*
Plant Pathology, 13(6), 614-629.

        Mendes, R., Kruijt, M., de Bruijn, I., Dekkers, E., van der Voort, M., Schneider, J. H., . . .
Bakker, P. A. (2011). Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science, 332*, 1097-1100.

Mukamolova, G. V., Kaprelyants, A. S., Young, D. I., Young, M., & Kell, D. B. (1998). A bacterial cytokine. *Proceedings of the National Academy of Sciences, 95*(15), 8916-8921.

Mukamolova, G. V., Murzin, A. G., Salina, E. G., Demina, G. R., Kell, D. B., Kaprelyants, A. S., & Young, M. (2006). Muralytic activity of *Micrococcus luteus* Rpf and its
relationship to physiological activity in promoting bacterial growth and resuscitation. *Molecular Microbiology, 59*(1), 84-98.

Mukamolova, G. V., Turapov, O. A., Kazarian, K., Telkov, M., Kaprelyants, A. S., Kell, D. B., & Young, M. (2002). The *rpf* gene of *Micrococcus luteus* encodes an essential secreted
growth factor. *Molecular Microbiology, 46*(3), 611-621.

        Muscarella, M. E., Jones, S. E., & Lennon, J. T. (2016). Species sorting along a subsidy gradient
490   alters bacterial community stability. *Ecology, 97*(8), 2034-2043.

        Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R., . . . Wagner, H.
492   (2011). R package '*vegan*': Community ecology package. *R package version 2*, 117-118.

        Peeters, N., Guidot, A., Vailleau, F., & Valls, M. (2013). *Ralstonia solanacearum*, a widespread
bacterial plant pathogen in the post☐genomic era. *Molecular Plant Pathology, 14*(7), 651-662.

Peralta, A. L., Sun, Y. M., McDaniel, M. D., & Lennon, J. T. (2018). Crop rotational diversity increases disease suppressive capacity of soil microbiomes. *Ecosphere, 9*(5).

Puspita, I. D., Kitagawa, W., Kamagata, Y., Tanaka, M., & Nakatsu, C. H. (2015). Increase in bacterial colony formation from a permafrost ice wedge dosed with a *Tomitella biformata*

recombinant resuscitation-promoting factor protein. *Microbes Environments, 30*(2), 151-156.

Ravagnani, A., Finan, C. L., & Young, M. (2005). A novel Firmicute protein family related to the Actinobacterial resuscitation-promoting factors by non-orthologous domain

displacement. *BMC Genomics, 6*(1), 39-53.

Reddy, G. S., & Garcia-Pichel, F. (2009). Description of *Patulibacter americanus* sp. nov.,

506   isolated from biological soil crusts, emended description of the genus *Patulibacter* Takahashi *et al.* 2006 and proposal of *Solirubrobacterales* ord. nov. and

508   *Thermoleophilales* ord. nov. *International Journal of Systematic Evolutionary Microbiology, 59*(1), 87-94.

510   Sakamoto, K., & Oba, Y. (1994). Effect of fungal to bacterial biomass ratio on the relationship between CO_2 evolution and total soil microbial biomass. *Biology Fertility of Soils, 17*(1),

512   39-44.

Salazar, A., Lennon, J. T., & Dukes, J. S. (2019). Microbial dormancy improves predictability of

514   soil respiration at the seasonal time scale. *Biogeochemistry, 144*(1), 103-116.

Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., . . .

516   Robinson, C. J. (2009). Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities.

518   *Applied Environmental Microbiology, 75*(23), 7537-7541.

Schroeckh, V., & Martin, K. (2006). Resuscitation-promoting factors: distribution among

520   Actinobacteria, synthesis during life-cycle and biological activity. *Antonie van Leeuwenhoek, 89*(3-4), 359-365.

522   Seki, T., Matsumoto, A., Ōmura, S., & Takahashi, Y. J. T. J. o. a. (2015). Distribution and isolation of strains belonging to the order Solirubrobacterales. *68*(12), 763.

524   Sharma, M., & Reynnells, R. (2016). Importance of soil amendments: survival of bacterial pathogens in manure and compost used as organic fertilizers. *Microbiology Spectrum,*

526   *4*(4). PFS-0010-2015

Shoemaker, W. R., & Lennon, J. T. (2018). Evolution with a seed bank: The population genetic

528   consequences of microbial dormancy. *Evolutionary Applications, 11*(1), 60-75.

van der Heijden, M. G., de Bruin, S., Luckerhoff, L., van Logtestijn, R. S., & Schlaeppi, K.

530   (2016). A widespread plant-fungal-bacterial symbiosis promotes plant biodiversity, plant nutrition and seedling recruitment. *The ISME Journal, 10*(2), 389-399.

532   Wagg, C., Bender, S. F., Widmer, F., & van der Heijden, M. G. (2014). Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the*

534   *National Academy of Sciences, 111*(14), 5266-5270.

Wang, G., Jagadamma, S., Mayes, M. A., Schadt, C. W., Steinweg, J. M., Gu, L., & Post, W. M.

536   (2015). Microbial dormancy improves development and experimental validation of ecosystem model. *The ISME Journal, 9*(1), 226-237.

538   Whitaker, J., Ostle, N., Nottingham, A. T., Ccahuana, A., Salinas, N., Bardgett, R. D., . . . McNamara, N. P. (2014). Microbial community composition explains soil respiration

540   responses to changing carbon inputs along an Andes⬜to⬜Amazon elevation gradient. *Journal of Ecology, 102*(4), 1058-1071.

FIGURE CAPTIONS

546   **Fig. 1.** Conceptual framework depicting the role of the soil microbial seed bank in local plant-microbe interactions. Members of the soil microbial community can transition between the

548   metabolically active pool (M_a) and the seed bank (M_d) through initiation into dormancy and resuscitation into an active state. Both M_a and M_d pools have associated levels of microbial

550   abundance (N) and local diversity (α). While only members in M_a can reproduce, the baseline rate of mortality (m_d) in M_d is reduced relative to the rate for M_a (m_a). Active microorganisms

552   (M_a) and plants (P) can directly interact with one another through a diverse suite of mechanisms, such as mutualism, parasitism, root colonization, and secretion of phytochemicals. We envision

554   that the M_d pool has minimal effects on plants but is likely influenced by the release of compounds via plant rhizodeposition.

    Fig. 2. Influence of resuscitation promoting factor (Rpf) on (A) *Brassica rapa* shoot biomass,

558   (B) root biomass, and (C) flower number produced per plant. We compared plant traits from individuals that were exposed to weekly additions of recombinant Rpf (+Rpf) to those exposed to

560   a protein buffer control (-Rpf). Black symbols represent the mean $\pm$ 1 $SE_{\bar{x}}$. Grey symbols represent the individual observations.

    Fig. 3. Effect of resuscitation promoting factor (Rpf) on soil microbial activity. Soil respiration

564   was measured after applying Rpf (+Rpf) or protein buffer control (-Rpf) to soils on a weekly basis. Symbols represent the mean $\pm$ 1 $SE_{\bar{x}}$.

568 **Fig. 4.** Effect of resuscitation promoting factor (Rpf) on (A) soil bacterial 16S rRNA copy

number, (B) fungal ITS copy number, and (C) the fungi : bacteria gene copy ratio. Gene copy

570 number was measured from soil after six weekly applications of recombinant Rpf (+Rpf) or

protein buffer control (-Rpf) treatment. Black symbols represent the mean $\pm$ 1 $SE_{x\square}$. Grey

572 symbols represent the individual observations.

574 **Fig. 5**. Principal Coordinate Analysis (PCoA) plot depicting composition of active (i.e., RNA)

and total (i.e., DNA) Actinobacteria from soil that were exposed to +Rpf and -Rpf treatments at

576 the end of a six-week experiment. The ellipses were generated by 'ordiellipse' function using the

standard deviation of PCoA point scores to visualize the spread of each treatment.

580   **Fig. 1**

Fig. 2

Fig. 3

Fig. 4

Fig. 5