Resuscitation of the microbial seed bank alters plant-soil interactions

2   Venus Kuo, Brent K. Lehmkuhl, Jay T. Lennon[*]

4   **Affiliations**

Department of Biology, Indiana University, Bloomington, IN 47405

6   *Correspondence to: lennonj@indiana.edu

8   **Key words**: *Brassica rapa*, dormancy, biodiversity, microbiome, resuscitation promoting factor

ABSTRACT

While microorganisms are recognized for driving belowground processes that influence the productivity and fitness of plant populations, the vast majority of bacteria and fungi in soil

belong to a seed bank consisting of dormant individuals. Still, plant performance may be affected by microbial dormancy through its effects on the activity, abundance, and diversity of soil

microorganisms. To test how microbial seed banks influence plant-soil interactions, we purified recombinant resuscitation promoting factor (Rpf), a bacterial protein that terminates dormancy.

Then, in a factorially designed experiment, we applied the Rpf to soil containing field mustard (*Brassica rapa*), an agronomically important plant species. Plant biomass was ~33 % lower in

the Rpf treatment compared to plants grown with an unmanipulated microbial seed bank. In addition, Rpf reduced soil respiration, decreased bacterial abundance, and increased fungal

abundance. These effects of Rpf on plant performance were accompanied by shifts in bacterial community composition, which may have diluted mutualists or resuscitated pathogens. Our

findings suggest that changes in microbial seed banks may influence the magnitude and direction of plant-soil feedbacks in ways that affect above- and below-ground biodiversity and function.

INTRODUCTION

Belowground soil microbial communities play a critical role in determining the productivity and fitness of aboveground plants. Plant roots are intimately associated with

thousands of bacterial and fungal taxa that are involved in soil processes such as decomposition, nutrient cycling, and pathogen suppression (Bardgett & van der Putten, 2014; Berendsen,

Pieterse, & Bakker, 2012; Wagg, Bender, Widmer, & van der Heijden, 2014). In addition to containing mutualistic symbionts, soils harbor pathogenic microorganisms that reduce plant

performance (Mansfield et al., 2012). Together, belowground microbial communities are responsible for generating plant-soil feedbacks that can influence the diversity and composition

of plant communities (Bever, 2003). However, the direction and magnitude of soil microbial effects on plants is variable (Hoeksema et al., 2010; Kulmatiski, Beard, Stevens, & Cobbold,

2008) and likely influenced by the complexity of belowground communities (van der Heijden, de Bruin, Luckerhoff, van Logtestijn, & Schlaeppi, 2016; Wagg et al., 2014).

One belowground feature that is commonly overlooked when considering plant-microbe interactions is the metabolic heterogeneity of soil microbial communities. Many microorganisms

are capable of entering a reversible state of reduced metabolic activity (i.e., dormancy) when challenged by suboptimal environmental conditions (Lennon & Jones, 2011). A large proportion

($\geq$90 %) of microorganisms are dormant in soils (Alvarez, Alvarez, Grigera, & Lavado, 1998; Blagodatskaya & Kuzyakov, 2013; Blagodatsky, Heinemeyer, & Richter, 2000; Lennon & Jones,

2011). Collectively, these inactive individuals create a seed bank, which not only maintains biodiversity, but also affects ecosystem functioning. The incorporation of microbial dormancy

into ecosystem experiments and models has been shown to improve predictions for soil

microbial activity and nutrient cycling (Salazar, Lennon, & Dukes, 2019; Wang et al., 2015).

48   Yet, the influence of microbial seed banks on plant-soil interactions remains to be determined.

Microbial seed banks may prevent local extinctions when exposed to fluctuating and

50   stressful conditions that are typical in soil habitats (Shoemaker & Lennon, 2018). If microbial

mutualists can persist by entering a seed bank, then this may promote beneficial soil functions

52   such as disease-suppression, induction of plant immune response, stimulation of root growth, and

biofertilization, which together, can increase plant performance (Lugtenberg & Kamilova, 2009;

54   Mendes et al., 2011; Peralta, Sun, McDaniel, & Lennon, 2018; Wagg et al., 2014). In contrast,

seed banks may contain pathogenic microorganisms that can reduce plant performance. For

56   example, it is well known that destructive pathogens can reside in soil for months to years after

the demise of plants, rendering potentially arable soil unusable (Koike, Subbarao, Davis, &

58   Turini, 2003; Peeters, Guidot, Vailleau, & Valls, 2013). Although microbial seed banks likely

contain both mutualistic and pathogenic taxa, the net effect of dormancy transitions on plant

60   performance is unknown.

Resuscitation is an essential process that regulates microbial seed-bank dynamics

62   (Lennon, den Hollander, M., & Blath, 2020). Dormancy is often terminated via the interpretation

of environmental cues or communication signals that are produced by other microorganisms

64   (Dworkin & Shah, 2010). One such signal is a bacterial protein called resuscitation promoting

factor (Rpf), a muralytic enzyme that degrades the β-(1,4) glycosidic bond in peptidoglycan,

66   which is a major cell-wall component of virtually all bacteria (Mukamolova, Kaprelyants,

Young, Young, & Kell, 1998). Rpf genes are broadly distributed among the G+C-rich Gram-

68   positive Actinobacteria (Ravagnani, Finan, & Young, 2005; Schroeckh & Martin, 2006). In some

soil samples, Rpf homologs can be found in up to 25 % of all genomes (Lennon & Jones, 2011).

70   Rpf has cross-species effects that stimulate the growth of closely related dormant bacteria

      (Puspita, Kitagawa, Kamagata, Tanaka, & Nakatsu, 2015; Schroeckh & Martin, 2006) either

72   through its direct effects on cell wall integrity or through the release of small peptide fragments

      that cross-link peptidoglycan, which may serve as signaling molecules (Dworkin, 2014). Under

74   laboratory conditions, Rpf can stimulate the activity of some microbial taxa at picomolar

      concentrations, but it has also been shown to inhibit the growth of other microorganisms

76   (Mukamolova et al., 1998; Mukamolova et al., 2002), which is not surprising given that Rpf is

      homologous to lysozyme (Cohen-Gonsaud et al., 2005), an antimicrobial enzyme produced by

78   many eukaryotic organisms. Overall, Rpf may have complex and interactive effects that

      influence plant-soil interactions.

80   In this study, we explored how seed-bank dynamics affect plant-microbe interactions by

      manipulating the process of resuscitation. Throughout the lifespan of a host plant (*Brassica*

82   *rapa*), we applied purified recombinant Rpf to soils containing a complex microbial community.

      The experiment was guided by a conceptual model that considers how active and inactive

84   microorganisms interact and potentially create feedbacks with plants (Fig. 1). We hypothesized

      that resuscitating the microbial seed bank with Rpf would alter plant performance through

86   changes in the abundance, activity, and composition of soil microbial communities. On the one

      hand, Rpf treatment could enhance plant performance by waking up mutualistic taxa in the soil

88   microbial community. On the other hand, resuscitating dormant microorganisms could

      negatively influence plant performance by diluting the benefits of mutualistic taxa or through the

90   recruitment of pathogenic microorganisms. Here, we evaluate these outcomes while considering

      other effects of Rpf on plants, microbes, and their interactions.

METHODS

Experimental design — We conducted a growth chamber experiment to test the effect of Rpf on

96   plant-soil interactions using *Brassica rapa* L. (Brassicaceae), an economically important plant

that consists of a variety of cultivated subspecies. We sowed 16 *B. rapa* seeds into individual

98   pots, which were obtained from Wisconsin Fast Plants™ (Standard stock seeds, Wisconsin Fast

Plants Program, University of Wisconsin, Madison, WI, USA). While *B. rapa* is non-

100   mycorrhizal, it still associates with a diverse microbial community that influences plant growth

(Lau & Lennon, 2012). For our experiment, we used plastic pots (17 cm diameter, 13 cm height)

102   filled with an autoclaved (121 °C, 15 PSI, 16 h) substrate consisting of, on a volumetric basis,

one part Metro Mix, one part Vermiculite, and one part soil. Based on a previous study, this

104   mixture prevented compaction and promoted *B. rapa* growth (Lau & Lennon, 2011). The soil

added to the potting mixture was collected from Indiana University Research and Teaching

106   Preserve (DMS 39 °11'56.5"N 86 °30'40.5"W). We collected this silt-loam soil from the surface

(0-5 cm) of a mesic, mid-successional site that was dominated by oaks (*Quercus velutina* and

108   *Quercus rubra*) and sugar maple (*Acer saccharum*) within the Norman Upland bedrock

physiographic unit (Schneider, 1966). Chemical and physical characteristics of the soil are listed

110   in Table S1.

For the +Rpf treatment, we added 1 mL of recombinant Rpf protein (10 µM) to the surface of

112   the soil substrate each week in three equidistant locations 5 cm away from main plant stem. The

same was done for the -Rpf treatment (negative control), except we used protein buffer (20 mM

114   Tris-HCl, 100 mM NaCl) instead of recombinant Rpf. Our experiment was conducted in a

Percival model PGC-15 growth chamber with high efficiency light lamps (Philips series 700 32

116   watts 4100K F32T8/TL741 AltoII). The growth chambers were set to constant and full light

       intensity (1100 µmol m^{-2} s^{-1}) with controlled humidity (60 %) and temperature (28 °C). Pots

118   were arranged in a fully randomized design where each plant was assigned to a resuscitation

       treatment (i.e., -Rpf vs. +Rpf). All pots were watered with equal amounts of filtered and de-

120   ionized water every other day.

122   ***Recombinant Rpf*** — To manipulate microbial seed banks, we overexpressed Rpf from

       Micrococcus luteus sp. KBS0714, a bacterial strain isolated from agricultural soil at the W.K.

124   Kellogg Biological Station, Hickory Corners, MI (MSU) (Kuo, Shoemaker, Muscarella, &

       Lennon, 2017; Lennon, Aanderud, Lehmkuhl, & Schoolmaster Jr, 2012). *Micrococcus luteus* sp.

126   KBS0714 is a close relative of *M. luteus* NCTC 2662 (99 % sequence 16S rRNA sequence

       similarity, NCBI CP001628.1), a model organism used for studying Rpf (Mukamolova et al.,

128   1998). To create a Rpf expression host, we amplified and cloned the *rpf* gene from KBS0714

       into the expression vector pET15b. First, we extracted KBS0714 genomic DNA from pure

130   culture using a Microbial DNA isolation kit (MoBio, Carlsbad, CA) for PCR amplification of the

       open reading frame of the *rpf* gene using two primers, Upper-F 5' GCC CAT ATG GCC ACC

132   GTG GAC ACC TG 3' and Lower-R 5' GGG GAT CCG GTC AGG CGT CTC AGG 3', with

       incorporated restriction sites EcoRI, NdeI (forward primer) and BamHI (reverse primer)

134   (Koltunov et al., 2010; Mukamolova et al., 1998). We amplified the *rpf* gene sequence with the

       following PCR conditions: initial: 95 °C for 5 min, 30 cycles of 95 °C for 30 s, 55 °C for 30 s, 72

136   °C for 1 min, and one final extension at 72 °C for 7 min. The *rpf* gene amplicon was ligated into

       pET15b (Invitrogen) as an EcoRI/BamHI fragment and transformed into *Escherichia coli* TOP10

138   (a non-expression host). We then identified positive clones with correct *rpf* sequence and

orientation by Sanger sequencing at the Indiana University Center of Genomics and

140 Bioinformatics (IU-CGB). Next, we extracted purified pET15b plasmids with the *rpf* gene from

transformed TOP10 *E. coli* using the QIAprep Spin Miniprep Kit (Qiagen) following the

142 manufacturer's protocol. Finally, we transformed the purified pET15b expression vector with the

rpf gene into the *E. coli* Origami BL21 (DE3) expression host incorporated with a polyhistidine-

144 tag on the N-terminus of the recombinant protein.

To overexpress Rpf, we grew the *E. coli* expression host in Lysogeny Broth (LB) with

146 appropriate antibiotics (ampicillin 100 µg mL^{-1}, kanamycin 15 µg mL^{-1}, and tetracycline 12.5 µg

mL^{-1}). During logarithmic growth, recombinant protein production was induced with Isopropyl

148 β-D-1-thiogalactopyranoside (IPTG) (100 µM final concentration). We confirmed

overexpression of Rpf with Western blots (Fig. S1). Cells were then collected by centrifugation,

150 lysed by sonication, and filtered through a Ni-NTA Purification System (Invitrogen) using a 10

mL gravity fed column with a 2 mL resin bed to purify recombinant Rpf with the N-terminus

152 polyhistidine-tag. Recombinant Rpf protein was washed with 5 mM imidazole buffer (300 mM

NaCl, 50 mM Tris-HCl, 5 mM imidazole) and then eluted with 125 mM imidazole buffer. Rpf

154 protein was purified by buffer exchange using a 10 mL Zeba Spin Desalting Columns (Thermo

Fisher) with protein buffer (20 mM Tris-HCl, 100 mM NaCl) following manufacturer's

156 instructions and then passed through a 0.2 µm syringe filter before adding to soil substrate as

described above.

Plant responses — We censused the plants every day and recorded germination and flowering

160 date. After most plants had finished flowering, we documented flower number, plant height, and

specific leaf area (SLA = leaf area/dry mass) as previously described (Lau & Lennon, 2012). All

162 open flowers in each experimental unit were hand-pollinated by other open flowers in the same

treatment group with a soft paint brush, which was cleaned with 30 % isopropyl alcohol between

164 treatment groups to prevent gene flow. Plants and seeds were harvested at the end of the six-

week growing period when most individuals ceased flowering and had begun to senesce. As an

166 estimate of female fitness, we removed plant seed pods and counted the number of seeds

produced. To estimate male fitness, we counted the number of flowers per plant. In addition, we

168 measured total, aboveground, and belowground biomass of each plant after drying plants at 65

°C for 48 h. We performed two-tailed Student's t-test to evaluate the effect of Rpf on total

170 biomass, aboveground (i.e., shoot) biomass, root biomass, shoot : root ratio, SLA, and plant

fitness (i.e., flower and seed numbers).

172   In addition to our main study, we conducted a smaller scale experiment with *Arabidopsis*

thaliana, a relative of *B. rapa* that also belongs to the Brassicaceae. Because this species is

174 amenable to being grown axenically in the absence of soil, we were able to test for the direct

effect of recombinant Rpf on plant performance. We placed seeds on sterile Murashige-Skoog

176 (MS) agar plates containing either Rpf protein (final concentration: 1.6 µmol L^{-1}) (+Rpf) or

protein buffer control (-Rpf). For five weeks, we maintained eight seedlings in a Percival model

178 PGC-15 growth chamber equipped with high efficiency lighting (Philips series 700 32 watts

4100K F32T8/TL741 AltoII). The growth chamber was set to constant and full light intensity

180 (1100 µmol m^{-2} s^{-1}) with controlled humidity (60 %) and temperature (28 °C). At the end of the

experiment, we assayed plant biomass based on estimates of leaf surface area by taking

182 standardized digital images which were analyzed with ImageJ software and a two-tailed

Student's t-test.

Soil microbial activity —We measured respiration of the soil substrate on a weekly basis to

determine the effect of Rpf treatment on belowground microbial activity. For each pot, we

transferred 1 g of soil substrate into glass vials with a silicon membrane septum cap. After

incubating at 25 °C in the dark for 24 h, we measured the CO_2 concentration (ppm) from 1 mL of

headspace gas using an infrared gas analyzer (IRGA) (LI-COR Environmental). We then

estimated CO_2 concentration in our samples based on values generated from a standard curve of

known CO_2 concentrations. We performed repeated-measures ANOVA (RM-ANOVA) using an

AR(1) covariance structure to test for the main effects and interaction of Rpf and time on rates of

respiration.

Microbial abundance — We estimated bacterial and fungal abundance using quantitative PCR

(qPCR) on DNA extracted from soil substrate. For each pot, we collected 1 g of soil substrate

pooled from three subsamples and immediately stored them at -80 °C. We extracted the DNA

from the sample using a PowerSoil® Total RNA Isolation Kit with DNA Elution Accessory Kit

(MoBio) following manufacturer's protocol, and the DNA was subsequently quantified using a

Take5 *Synergy* microplate reader (BioTek). We then conducted qPCR assays described in greater

detail elsewhere (Fierer, Jackson, Vilgalys, & Jackson, 2005; Lau & Lennon, 2011). Briefly,

each 20 μL reaction contained 1 μL of DNA template (2.5 ng μL^{-1}), 1 μL of each primer (10

μmol L^{-1}), 7 μL of molecular-grade water, and 10 μL iQTM SYBR Green SuperMix (Bio-Rad

Laboratories Hercules, CA, USA). For bacteria we used the Eub338 forward primer

(ACTCCTACGGGAGGCAGCAG) (Lane, 1991) and the Eub518 reverse primer

(ATTACCGCGGCTGCTGG) (Muyzer, Dewaal, & Uitterlinden, 1993) to amplify the 16S

rRNA gene, and for fungi we used the ITS1f forward primer (TCCGTAGGTGAACCTGCGG)

208   (Gardes & Bruns, 1993) and the 5.8S reverse primer (CGCTGCGTTCTTCATCG) (Vilgalys &

Hester, 1990) to amplify the ITS gene region. qPCR assays were performed using Eppendorf

210   Mastercycler Realplex system using the previously reported thermal cycle conditions: 15 min at

95 °C, followed by 40 cycles of 95 °C for 1 min, 30 s at 53 °C for annealing, followed by 72 °C

212   for 1 min (Fierer et al., 2005). The coefficients of determination (r^2) of our assay ranged from

0.95 and 1, while amplification efficiencies fell between 0.93 and 0.99. Based on melting curve

214   analyses, we found no evidence for primer dimers. We estimated fungal and bacterial abundance

based on the estimated gene copy number from their respective standard curves generated from

216   bacterial and fungal isolates as described elsewhere (Lau & Lennon, 2012). We performed a two-

tailed Student's *t*-test to determine the effect of the Rpf treatment on soil bacterial (16S rRNA)

218   and fungal (ITS) gene copy abundances, as well as the fungal to bacterial ratio (F : B).

220   ***<u>Microbial diversity</u>*** — To account for the variation in metabolic activity, we characterized

bacterial communities from the soil substrate using pools of RNA and DNA. DNA is a relatively

222   stable molecule contained in intact cells irrespective of their metabolic status. Accordingly, we

interpret 16S rRNA sequences recovered from the DNA pool as the "total" community. In

224   contrast, RNA is a more ephemeral molecule that is required for protein synthesis by actively

growing cells. Therefore, we interpret 16S rRNA sequences recovered from the RNA pool after

226   complementary DNA (cDNA) synthesis as the "active" community (Jones & Lennon, 2010). We

assume that dormant individuals can create discrepancies between the active and total

228   composition of a given sample, but we do not attempt to use RNA and DNA to directly

characterize dormant taxa. Following previously described protocols (Lennon, Placella, &

230   Muscarella, 2017), we characterized the effects of Rpf on soil bacterial communities using high-

throughput sequencing. First, we extracted nucleic acids using the PowerSoil Total RNA Extraction

232   Kit with DNA Elution Accessory Kit (MoBio, Carlsbad, CA) followed by cleaning via ethanol

precipitation. We removed residual DNA from RNA samples using DNase 1 (Invitrogen)

234   following manufacturer's protocol and synthesized cDNA by reverse transcribing RNA using

iScript Reverse Transcription Supermix for RT-qPCR (Bio-Rad). After ensuring that there was

236   no product in our no-template negative controls, we amplified the V4 region of the 16S rRNA

gene from bacterial DNA and cDNA using the 515F primer (GTGYCAGCMGCCGCGGTAA)

238   and the 806R primer (GGACTACNVGGGTWTCTAAT) that included a unique barcode for

each sample (Caporaso et al., 2012). Thermal cycle conditions for the PCR reaction consisted of

240   3 min at 94 °C, followed by 30 cycles of 94 °C for 45 s, 30 s at 50 °C, 90 s at 72 °C, followed by

72 °C for 10 min. We cleaned the sequence libraries using the Agencourt AMPure XP

242   purification kit (Beckman Coulter, Brea, CA, USA), quantified the resulting products using the

QuanIt PicoGreen kit (Invitrogen), and pooled libraries at equal molar ratios (final concentration:

244   10 ng per library). We then sequenced the pooled libraries with the Illumina MiSeq platform

using 250 x 250 bp paired end reads (Illumina Reagent Kit v2, 500 reaction kit) at the IU-CGB.

246   Paired-end raw 16S rRNA sequence reads were assembled into contigs using the Needleman

algorithm (Needleman & Wunsch, 1970). We then trimmed the resulting sequences using a

248   moving average quality score (window = 50 bp, score = 35), in addition to removing long

sequences, ambiguous base calls, and sequences that matched Archaea, chloroplasts, and other

250   non-bacteria. We also removed chimeric sequences that were detected using the UCHIME

algorithm (Edgar, Haas, Clemente, Quince, & Knight, 2011). After this filtering, there was a

252   total of 3,217,419 quality reads. We aligned the sequences to the Silva Database (v 128) using

the Needleman algorithm (Needleman & Wunsch, 1970). We created operational taxonomic

254   units (OTUs) by first splitting the sequences based on taxonomic class (using the RDP

taxonomy) and then binning sequences into OTUs based on 97 % sequence similarity using the

256 OptiClust algorithm (Westcott & Schloss, 2017). We removed all OTUs with less than two

occurrences in the data set. Altogether, this led to a high degree of coverage across samples

258 (minimum Good's coverage = 0.97). All sequence processing was completed using the software

package *mothur* (version 1.39.5) (Schloss et al., 2009).

260 First, we used the 16S rRNA sequences to test how the Rpf treatment affected measures

of alpha diversity with samples. We estimated bacterial richness (i.e., the number of OTUs in a

262 sample) using a resampling with replacement approach that subsampled 1,000 sequence

observations per sample, resampled 999 additional times, and then calculated the average

264 number of OTUs to estimate per sample richness ($\pm$ SEM) (Gotelli & Colwell, 2011; Muscarella,

Jones, & Lennon, 2016). Similarly, we estimated the effect of Rpf treatment on taxa evenness

266 (i.e., the equitability in relative abundance of OTUs in a sample) using the same resampling

approach with the Evar index (Smith & Wilson, 1996). We tested for the effect of Rpf on these

268 measures of diversity using ANOVA.

Second, we used the sequence data to test how the Rpf treatment affected beta diversity

270 among samples. Given that relative abundances spanned order of magnitudes, we

$\log_{10}$-transformed data to prevent undue weight of dominant species. Following this, we

272 conducted Principal Coordinates Analyses (PCoA) using the Bray-Curtis dissimilarity metric to

visualize the effects of Rpf on bacterial communities. To test the hypothesis that Rpf affected the

274 composition of the total (DNA) and active (RNA) bacterial pools, we used permutational

multivariate analysis of variance (PERMANOVA) with Bray-Curtis dissimilarity metric

276 implemented with the `adonis` function in the *vegan* package (Oksanen et al., 2011) in the R

statistics environment (v 3.2.3). We coded pot number as a factor to account for paired-sample

278   design given that DNA and RNA were co-extracted from the same soil-substrate sample within

an experimental unit. Last, we conducted indicator species analyses to identify influential taxa

280   driving compositional changes in response to the Rpf treatment. Specifically, we calculated

Pearson's phi coefficients of association with taxa abundance data using the `multipatt` function

282   in the *indicspecies* R package (De Cáceres & Legendre, 2009). We filtered the output from the

indicator species analyses so as to only consider associations between taxa and treatments where

284   P-values were ≤ 0.05 and correlations were $\geq |0.7|$.

286   **RESULTS**

Plant responses — Plants in +Rpf treatment had less total ($t_7 = 2.84$, $P = 0.013$), aboveground (t_7

288    $= 2.70$, $p = 0.017$), and belowground biomass ($t_7 = 2.30$, $P = 0.049$) than plants in the -Rpf

treatment (Fig. 2). While recombinant protein led to a ~33 % reduction in root biomass, Rpf had

290   no effect on other plant traits including SLA ($t_7 = 0.80$, $P = 0.442$), shoot : root ratio ($t_7 = -1.20$,

$P = 0.250$), shoot height ($t_7 = 1.11$, $P = 0.286$), or the number of seeds produced per plant ($t_7 =$

292   0.83, $P = 0.421$) (Fig. S2). However, we did detect a marginal decrease in the number of flowers

produced per plant in the +Rpf treatment ($t_7 = 1.79$, $P = 0.097$). In the *Arabidopsis* experiment

294   where individuals were grown in the absence of soil-substrate and an accompanying

microbiome, there was no significant effect of Rpf on estimates of plant biomass ($t_3 = -1.11$, $P =$

296   0.320, Fig. S3).

298   ***Microbial activity*** — Rpf altered the activity of the belowground microbial community based on

respiration of the soil substrate. We detected a significant main effect of Rpf ($F_{1, 84} = 9.68$, $P =$

300   0.002) and time ($F_{5, 84} = 21.60$, $P < 0.001$), but no interaction ($F_{5, 84} = 1.22$, $P = 0.308$), on

respiration (Fig. 3). While respiration increased throughout the experiment in both treatments,

302   based on marginal means, soil respiration was 24 % lower in the +Rpf treatment than in the -Rpf

treatment.

__Microbial abundance__ — The Rpf treatment significantly altered the abundance of soil bacteria

306   (t_8 = 2.71, P = 0.016) and fungi (t_8 = -3.27, P = 0.007) relative to the control (Fig. 4). Bacterial

abundance estimated as 16S rRNA gene copy number decreased by ~30 % in the +Rpf treatment

308   relative to the -Rpf treatment. In contrast, fungal abundance estimated as the ITS gene copy

number increased 2.8-fold in the +Rpf treatment. Consequently, soil F : B ratio increased 4.8-

310   fold under +Rpf treatment relative to the control treatment (t_8 = -2.84, P = 0.018).

312   *__Bacterial diversity__* — Bacteria in the soil-substrate were diverse and reflected compositions that

were not unlike many natural communities. Based on 16S rRNA sequencing, we recovered a

314   total 32,055 taxa across the experimental units. The bacterial community was dominated by

OTUs belonging to the following phyla: Proteobacteria (49 %), Acidobacteria (14 %),

316   Verrucomicrobia (7 %), Planctomycetes (7 %), Bacteroidetes (5 %), Actinobacteria (4 %),

Chlorofexi (2 %), and Firmicutes (2 %).

To evaluate broad-scale treatment effects on microbial diversity, we first examined changes

320   among the major groups of bacteria. Rpf additions had no effect on bacterial richness regardless

of whether sequences came from the total ($F_{1, 8}$ = 0.44, P = 0.526) or active community ($F_{1, 8}$ =

322   0.047, P = 0.835) (Table S2). Similarly, Rpf had no effect on the evenness of the total ($F_{1, 8}$ =

0.527, P = 0.489) or active ($F_{1, 8}$ = 0.108, P = 0.751) community (Table S2). In contrast, bacterial

324   composition was significantly affected by metabolic status (i.e., active vs total) ($F_{1, 19} = 7.24$, r^2

= 0.23, $P = 0.001$) and the Rpf treatment ($F_{1, 19} = 1.5$, $r^2 = 0.06$, $P = 0.032$), which can be

326   visualized in Fig. 5. Indicator species analysis revealed that there were 137 taxa belonging to the

Proteobacteria and Bacteroidetes that were associated with the + Rpf treatment. Meanwhile, 195

328   taxa in the Planctomycetes, Proteobacteria, Chlorflexi, Acidobacteria, and Actinobacteria were

significantly associated with the -Rpf treatment.

        Since the production of Rpf is thought to be restricted to the Actinobacteria, we also

332   evaluated how treatments affected the diversity of taxa within this phylum. Rpf had no effect on

the richness (total community: $F_{1, 8} = 0.80$, $P = 0.397$; active community: $F_{1, 8} = 0.14$, $P = 0.722$),

334   nor did it influence evenness (total community: $F_{1, 8} = 0.69$, $P = 0.429$; active community: $F_{1, 8} =$

0.41, $P = 0.538$) (Table S3). There was a small and marginally significant reduction in the

336   relative abundance of actinobacterial sequences in the +Rpf treatment compared to the -Rpf

treatment ($t_9 = 1.46$, $P = 0.083$) (Fig. S4). Based on PERMANOVA, actinobacterial composition

338   was significantly affected by both metabolic status (i.e., active vs total) ($F_{1, 19} = 16.08$, $r^2 = 0.33$,

$P = 0.001$) and the Rpf treatment ($F_{1, 19} = 3.25$, $r^2 = 0.07$, $p = 0.010$) (Fig. S5). Indicator species

340   analysis revealed that OTUs belonging to the *Acidothermus*, *Catenulispora*, and *Mycobacterium*

were associated with the -Rpf treatment, while only a single taxon belonging to the

342   Solirubrobacterales had a significant association with the +Rpf treatment (Fig. S6)

344   **DISCUSSION**

        Dormancy is an important life-history trait that influences the diversity, composition, and

346   function of soil microbial communities. After generating recombinant protein with a gene from

an environmental isolate, we applied resuscitation promoting factor (Rpf) to a soil community to

348   test the hypothesis that microbial seed-bank dynamics alter plant growth and fitness traits by

driving changes in the belowground microbial community (Fig. 1). The Rpf treatment decreased

350   plant biomass (Fig. 2) most likely by altering the activity (Fig. 3), abundance (Fig. 4), and

composition (Fig. 5) of soil microbial communities. These findings are consistent with the view

352   that soil microbial seed banks can influence plant performance, perhaps by disrupting

interactions with beneficial microorganisms or through the recruitment of pathogens. In the

354   following sections, we discuss these findings while exploring other potential ways in which Rpf

may affect plant-microbe interactions in soil environments.

Rpf indirectly affects plant performance — Resuscitation of microbial seed banks led to a 33 %

358   reduction in plant biomass (Fig. 2). For the following reasons, we infer that Rpf effects on plant

performance are likely to be indirect. First, it is known that Rpf activity involves the hydrolysis

360   of glyosidic linkages in peptidoglycan (Cohen-Gonsaud et al., 2005; Mukamolova et al.,

2006).While this polymer is found in the cell walls of virtually all bacteria, it is absent from plant

362   tissue. Second, we conducted an experiment with *Arabidopsis thaliana*, a fairly close relative of

B. rapa, to evaluate whether Rpf can directly affect plant performance. We detected no

364   significant treatment effect after maintaining a relatively small number of seedlings on

microbe-free agar for five weeks. However, there was a trend of reduced plant growth (18 %) for

366   individuals in the +Rpf treatment. While additional experiments may be warranted, for the

purposes of this study, we cautiously conclude that Rpf influenced plants primarily via its effect

368   on the activity, abundance, or composition of the soil microbial community.

370 ***Rpf altered fungal-bacterial interactions*** — Rpf may have altered plant performance by

modifying fungal-bacterial interactions. We observed a nearly three-fold increase in fungal

372 abundance in response to Rpf additions (Fig. 4). As a result, soil F : B ratio increased under Rpf

treatment (Fig. 4), which may also explain the reduction in soil microbial activity (Fig. 3) given

374 that there are often differences in carbon-use efficiency between bacteria and fungi (Sakamoto &

Oba, 1994; Whitaker et al., 2014). Shifts in F : B may also reflect the way that Rpf influences

376 microorganisms with different cell-wall properties. Like plants, fungi do not contain

peptidoglycan, the substrate which Rpf acts upon. Instead, the fungal cell wall is primarily

378 composed of chitin. Therefore, an increase in fungal abundance could arise if Rpf decreased the

competitive ability of soil bacteria in our system. In principle, the muralytic activity of Rpf may

380 have even lysed some bacterial cells. If this occurred, the resulting necromass could be

scavenged by fungi to meet their metabolic demands (Bradley, Amend, & LaRowe, 2018). Our

382 results, however, do not support this hypothesis. We found that bacterial abundance from three

contrasting soils increased with Rpf concentration up to 4 μM g^{-1} soil. At higher concentrations

384 (8 μM g^{-1}), we no longer observed a growth-stimulating effect, but bacterial abundance never

dropped below levels observed in control soils without Rpf (Fig. S6). Rpf was much lower (40

386 nM g^{-1} soil) in our *Brassica* experiment, yet above concentrations shown to resuscitate dormant

bacteria (Cohen-Gonsaud et al., 2005; Mukamolova et al., 1998). In addition to demonstrating

388 the generality of our recombinant protein on different soils, these findings suggest that it is

unlikely that reductions in bacterial abundance (Fig. 4) were due to lysis or direct inhibition of

390 the soil microbial community.

392 ***Rpf altered microbial diversity*** – While seed banks are known to promote biodiversity by

buffering ecological and evolutionary dynamics, microbial dormancy may have complex effects

394 that influence plant-soil dynamics. The influence of Rpf on communities of interacting species is

likely dependent on enzyme specificity. Very few studies have investigated Rpf effects on a

396 broad range of microorganisms, but some evidence suggests cross-species resuscitation using

recombinant Rpf generated from actinobacterial strains belonging to *Mycobacterium* and

398 *Brevibacterium* (Mukamolova et al., 1998; Puspita et al., 2015).

In our study, we found that Rpf had broad-scale effects on bacterial communities. Results

400 from an indicator species analysis suggest that some taxa may have been inhibited by the

treatment, while others belonging to the Proteobacteria and Bacteroidetes were favored by Rpf

402 additions. These findings imply that Rpf from a single species may resuscitate a wide range of

taxa, consistent with reports of a conserved catalytic site within the peptidoglycan substrate

404 (Mukamolova et al., 2002). Alternatively, Rpf may only stimulate growth in a smaller set of taxa,

but their metabolism leads to a cascade of community change. Our indicator species point to one

406 such group of taxa within the Actinobacteria, which are the only known producers of Rpf.

Specifically, OTUs belonging to the *Solirubrobacterales* were associated with the +Rpf

408 treatment. Members of this group are aerobic, non-spore-forming, and non-motile bacteria that

are often recovered in forest and agricultural soils (Kim et al., 2007; Reddy & Garcia-Pichel,

410 2009; Seki, Matsumoto, Ōmura, & Takahashi, 2015). Little is known about the role of

Solirubrobacterales in the context of plant-microbe interactions, but some representatives

412 assimilate simple organic compounds (e.g., glucose, maltose, sucrose, xylose, and arginine) that

are commonly found in the rhizosphere (Reddy & Garcia-Pichel, 2009). Interestingly, the

414 *Solirubrobacterales* are only distantly related to *Micrococcus* KBS0714, the soil isolate

containing the Rpf gene that we used for producing the purified recombinant protein. Again, this

416   suggests that Rpf functionality may not be restricted to close kin. Additional work is needed to

unveil the network of interactions among Rpf producers and responders, in addition to whether

418   or not Rpf genes map onto the phylogeny of Actinobacteria.

420   ***Implications of seed-bank dynamics for plant-microbe interactions*** – Our study demonstrates

that resuscitation promoting factor (Rpf) modified plant performance most likely through its

422   effects on soil microbial communities. Reductions in plant growth were accompanied by shifts in

soil microbial community properties, which may have included the recruitment of pathogens.

424   Although we did not observe root lesions or other clear signs of infection, we cannot rule out the

possibility the seed banks harbor pathogenic or non-mutualistic taxa that could decrease plant

426   performance. Such a view is consistent with reports of microbial pathogens persisting in soil for

long periods of time, likely in a state of reduced metabolic activity (Sharma & Reynnells, 2016).

428   More explicit tests would include the targeting of known pathogens combined perhaps with

measurements of plant immune responses to altered microbial seed banks. Alternatively, our

430   findings could be explained by a dilution of mutualists in the soil or a general disruption of

beneficial plant-microbe interactions, which would likely be reflected in the profiling of plant

432   exudates and microbial metabolites.

While our study provides empirical support for the notion that microbial seed banks can

434   affect plant performance, it also raises a number of questions for future exploration. We observed

that a one protein from a single strain of bacteria can have broad effects on interacting plants and

436   microorganisms. However, some bacteria may be responsive to certain families of Rpf, while

others may not. This likely is due to the mechanisms by which Rpf operates in a community

438 context, which is a topic that has received little attention to date. More generally, metabolic

transitions into and out of dormancy are influenced by mechanisms besides Rpf (Lennon &

440 Jones, 2011) many of which are tied to the interpretation of environmental triggers (e.g., soil

rewetting) that are known to resuscitate microorganisms (Aanderud, Jones, Fierer, & Lennon,

442 2015). Finally, there is a need to scale up manipulative experiments to explore microbial seed

banks in more complex communities. Generalities could be assessed by examining the effects of

444 Rpf outside of the Brassicaceae given that these plants tend to lack mycorrhizal symbionts,

which are important mutualists found among most flowering plants. It is not difficult to imagine

446 that more complex outcomes could emerge when altered microbial seed banks are considered in

diverse plant communities (Fig. 1). Nevertheless, our study highlights a potentially important but

448 overlooked component of plant-microbe interactions. It is estimated that soil microbial

communities can be dominated by dormant taxa, yet changes in activity can be fast, suggesting

450 that seed banks may be an important factor contributing to plant-soil feedbacks, which is thought

to be an important mechanism maintaining landscape patterns of biodiversity (Bever, 2003).

ACKNOWLEDGEMENTS

454 We acknowledge Peyton Thomas for assistance in the laboratory and members of the Lennon lab

for critical feedback on an earlier version of the manuscript. This work was supported by the

456 National Science Foundation (1442246 JTL, 1934554 JTL), the US Army Research Office Grant

(W911NF- 14-1-0411 JTL), and the National Aeronautics and Space Administration

458 (80NSSC20K0618 JTL).

AUTHOR CONTRIBUTIONS

V.K. and J.T.L. designed the study; V.K. and B.K.L performed the experiments; V.K. and J.T.L analyzed the data; V.K. and J.T.L. wrote the paper.

DATA AVAILABILITY STATEMENT

All DNA sequences can be downloaded from the NCBI BioProject PRJNA504042. Other data and code are available from the Dryad Digital Repository (https ://doi.org/10.XXX/XXXXX) and GitHub (https://github.com/LennonLab/BrassicaRpf)

REFERENCES

Aanderud, Z. T., Jones, S. E., Fierer, N., & Lennon, J. T. (2015). Resuscitation of the rare biosphere contributes to pulses of ecosystem activity. *Frontiers in Microbiology, 6*, 24.

Alvarez, C., Alvarez, R., Grigera, M., & Lavado, R. (1998). Associations between organic matter fractions and the active soil microbial biomass. *Soil Biology Biochemistry, 30*(6), 767-773.

Bardgett, R. D., & van der Putten, W. H. (2014). Belowground biodiversity and ecosystem functioning. *Nature, 515*(7528), 505-511.

Berendsen, R. L., Pieterse, C. M., & Bakker, P. A. (2012). The rhizosphere microbiome and plant health. *Trends in Plant Science, 17*(8), 478-486.

Bever, J. D. (2003). Soil community feedback and the coexistence of competitors: conceptual frameworks and empirical tests. *New Phytologist, 157*(3), 465-473.

Blagodatskaya, E., & Kuzyakov, Y. (2013). Active microorganisms in soil: Critical review of

482   estimation criteria and approaches. *Soil Biology & Biochemistry, 67*, 192-211.

Blagodatsky, S. A., Heinemeyer, O., & Richter, J. (2000). Estimating the active and total soil

484   microbial biomass by kinetic respiration analysis. *Biology Fertility of Soils, 32*(1), 73-81.

Bradley, J. A., Amend, J. P., & LaRowe, D. E. (2018). Necromass as a limited source of energy

486   for microorganisms in marine sediments. *Journal of Geophysical Research:*

Biogeosciences, 123(2), 577-590.

488   Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., . . .

Bauer, M. (2012). Ultra-high-throughput microbial community analysis on the Illumina

490   HiSeq and MiSeq platforms. *The ISME Journal, 6*(8), 1621-1624.

Cohen-Gonsaud, M., Barthe, P., Bagnéris, C., Henderson, B., Ward, J., Roumestand, C., & Keep,

492   N. H. (2005). The structure of a resuscitation-promoting factor domain from

Mycobacterium tuberculosis shows homology to lysozymes. *Nature Structural Molecular*

494   *Biology, 12*(3), 270.

De Cáceres, M., & Legendre, P. (2009). Associations between species and groups of sites:

496   indices and statistical inference. http://sites.google.com/site/miqueldecaceres/.

Dworkin, J. (2014). The medium is the message: interspecies and interkingdom signaling by

498   peptidoglycan and related bacterial glycans. *Annual Review of Microbiology, 68*, 137-

154.

500   Dworkin, J., & Shah, I. M. (2010). Exit from dormancy in microbial organisms. *Nature Reviews*

Microbiology, 8(12), 890-896.

502   Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., & Knight, R. (2011). UCHIME improves

sensitivity and speed of chimera detection. *Bioinformatics, 27*(16), 2194-2200.

504   Fierer, N., Jackson, J. A., Vilgalys, R., & Jackson, R. B. (2005). Assessment of soil microbial

community structure by use of taxon-specific quantitative PCR assays. *Applied*

506   *Environmental Microbiology, 71*(7), 4117-4120.

Gardes, M., & Bruns, T. D. (1993). ITS primers with enhanced specificity for Basidiomycetes:

508   application to the identification of mycorrhizae and rusts. *Molecular Ecology, 2*(2), 113-

118.

510   Gotelli, N. J., & Colwell, R. K. (2011). Estimating species richness. In A. E. Magurran & B. J.

McGill (Eds.), *Biological Diversity: Frontiers in Measurement and Assessment* (pp. 39-

512   54). United Kingdom: Oxford University Press.

Hoeksema, J. D., Chaudhary, V. B., Gehring, C. A., Johnson, N. C., Karst, J., Koide, R. T., . . .

514   Moore, J. C. (2010). A meta-analysis of context-dependency in plant response to

inoculation with mycorrhizal fungi. *Ecology Letters, 13*(3), 394-407.

516   Jones, S. E., & Lennon, J. T. (2010). Dormancy contributes to the maintenance of microbial

diversity. *Proceedings of the National Academy of Sciences, 107*(13), 5881-5886.

518   Kim, M. K., Na, J.-R., Lee, T.-H., Im, W.-T., Soung, N.-K., Yang, D.-C. (2007). *Solirubrobacter*

soli sp. nov., isolated from soil of a ginseng field. *International Journal of Systematic and*

520   *Evolutionary Microbiology*, *57*(7), 1453-1455.

Koike, S., Subbarao, K., Davis, R. M., & Turini, T. (2003). *Vegetable diseases caused by*

522   *soilborne pathogens*: UCANR Publications.

Koltunov, V., Greenblatt, C. L., Goncharenko, A. V., Demina, G. R., Klein, B. Y., Young, M., &

524   Kaprelyants, A. S. (2010). Structural changes and cellular localization of resuscitation-

promoting factor in environmental isolates of *Micrococcus luteus*. *Microbial Ecology,*

526   *59*(2), 296-310.

Kulmatiski, A., Beard, K. H., Stevens, J. R., & Cobbold, S. M. (2008). Plant-soil feedbacks: a

528   meta-analytical review. *Ecology Letters, 11*(9), 980-992.

Kuo, V., Shoemaker, W., Muscarella, M., & Lennon, J. (2017). Whole-genome sequence of the

530   soil bacterium *Micrococcus* sp. KBS0714. *Genome Announcements, 5*(32), e00697-

        00617.

532 Lane, D. (1991). 16S/23S rRNA sequencing. In E. Stackebrandt & M. Goodfellow (Eds.),

        Nucleic acid techniques in bacterial systematics. (pp. 115-175). West Sussex, United

534   Kingdom: JohnWiley & Sons.

Lau, J. A., & Lennon, J. T. (2011). Evolutionary ecology of plant–microbe interactions: soil

536   microbial structure alters selection on plant traits. *New Phytologist, 192*(1), 215-224.

Lau, J. A., & Lennon, J. T. (2012). Rapid responses of soil microorganisms improve plant fitness

538   in novel environments. *Proceedings of the National Academy of Sciences of the United

        States of America, 109*(35), 14058-14062.

540 Lennon, J. T., Aanderud, Z. T., Lehmkuhl, B., & Schoolmaster Jr, D. R. (2012). Mapping the

        niche space of soil microorganisms using taxonomy and traits. *Ecology, 93*(8), 1867-

542   1879.

Lennon, J. T., den Hollander, F., M., W.-B., & Blath, J. (2020). Principles of seed banks:

544   complexity emerging from dormancy. *arXiv*, arXiv:2012.00072.

Lennon, J. T., & Jones, S. E. (2011). Microbial seed banks: the ecological and evolutionary

546   implications of dormancy. *Nature Reviews Microbiology, 9*(2), 119-130.

Lennon, J. T., Placella, S. A., & Muscarella, M. E. (2017). Relic DNA contributes minimally to

548   estimates of microbial diversity. *mBio,* 9:e00637-18.

Lugtenberg, B., & Kamilova, F. (2009). Plant-growth-promoting rhizobacteria. *Annual Review of*

550   *Microbiology, 63*, 541-556.

Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., . . . Machado, M.

552   A. (2012). Top 10 plant pathogenic bacteria in molecular plant pathology. *Molecular*

        Plant Pathology, 13(6), 614-629.

554 Mendes, R., Kruijt, M., de Bruijn, I., Dekkers, E., van der Voort, M., Schneider, J. H., . . .

        Bakker, P. A. (2011). Deciphering the rhizosphere microbiome for disease-suppressive

556   bacteria. *Science, 332*, 1097-1100.

Mukamolova, G. V., Kaprelyants, A. S., Young, D. I., Young, M., & Kell, D. B. (1998). A

558   bacterial cytokine. *Proceedings of the National Academy of Sciences, 95*(15), 8916-8921.

Mukamolova, G. V., Murzin, A. G., Salina, E. G., Demina, G. R., Kell, D. B., Kaprelyants, A.

560   S., & Young, M. (2006). Muralytic activity of *Micrococcus luteus* Rpf and its

        relationship to physiological activity in promoting bacterial growth and resuscitation.

562   *Molecular Microbiology, 59*(1), 84-98.

Mukamolova, G. V., Turapov, O. A., Kazarian, K., Telkov, M., Kaprelyants, A. S., Kell, D. B.,

564   & Young, M. (2002). The *rpf* gene of *Micrococcus luteus* encodes an essential secreted

        growth factor. *Molecular Microbiology, 46*(3), 611-621.

566 Muscarella, M. E., Jones, S. E., & Lennon, J. T. (2016). Species sorting along a subsidy gradient

        alters bacterial community stability. *Ecology, 97*(8), 2034-2043.

568 Muyzer, G., Dewaal, E. C., & Uitterlinden, A. G. (1993). Profiling of complex microbial

        populations by denaturing gradient gel electrophoresis analysis of polymerase chain

570   reactioin-amplified genes coding for 16S ribosomal. *Applied and Environmental*

        Microbiology, 59(3), 695-700.

572   Needleman, S. B., & Wunsch, C. D. (1970). A general method applicable to the search for

similarities in the amino acid sequence of two proteins. *Journal of Moleciular Biology,*

574   *48*, 443-453.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R., . . . Wagner, H.

576   (2011). R package '*vegan*': Community ecology package. *R package version 2*, 117-118.

Peeters, N., Guidot, A., Vailleau, F., & Valls, M. (2013). *Ralstonia solanacearum*, a widespread

578   bacterial plant pathogen in the post-genomic era. *Molecular Plant Pathology, 14*(7), 651-

662.

580   Peralta, A. L., Sun, Y. M., McDaniel, M. D., & Lennon, J. T. (2018). Crop rotational diversity

increases disease suppressive capacity of soil microbiomes. *Ecosphere, 9*(5).

582   Puspita, I. D., Kitagawa, W., Kamagata, Y., Tanaka, M., & Nakatsu, C. H. (2015). Increase in

bacterial colony formation from a permafrost ice wedge dosed with a *Tomitella biformata*

584   recombinant resuscitation-promoting factor protein. *Microbes Environments, 30*(2), 151-

156.

586   Ravagnani, A., Finan, C. L., & Young, M. (2005). A novel Firmicute protein family related to

the Actinobacterial resuscitation-promoting factors by non-orthologous domain

588   displacement. *BMC Genomics, 6*(1), 39-53.

Reddy, G. S., & Garcia-Pichel, F. (2009). Description of *Patulibacter americanus* sp. nov.,

590   isolated from biological soil crusts, emended description of the genus *Patulibacter*

Takahashi *et al*. 2006 and proposal of *Solirubrobacterales* ord. nov. and

592   *Thermoleophilales* ord. nov. *International Journal of Systematic Evolutionary*

Microbiology, 59(1), 87-94.

594   Sakamoto, K., & Oba, Y. (1994). Effect of fungal to bacterial biomass ratio on the relationship

between CO_2 evolution and total soil microbial biomass. *Biology Fertility of Soils, 17*(1),

596   39-44.

Salazar, A., Lennon, J. T., & Dukes, J. S. (2019). Microbial dormancy improves predictability of

598   soil respiration at the seasonal time scale. *Biogeochemistry, 144*(1), 103-116.

Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., . . .

600   Robinson, C. J. (2009). Introducing mothur: open-source, platform-independent,

community-supported software for describing and comparing microbial communities.

602   *Applied Environmental Microbiology, 75*(23), 7537-7541.

Schneider, A. F. (1966). Physiography. In A. A. Lindseay (Ed.), *Natural Features of Indiana*.

604   Indianapolis, Indiana: Indiana Academy of Science, Indiana State Library.

Schroeckh, V., & Martin, K. (2006). Resuscitation-promoting factors: distribution among

606   Actinobacteria, synthesis during life-cycle and biological activity. *Antonie van

Leeuwenhoek, 89*(3-4), 359-365.

608   Seki, T., Matsumoto, A., Ōmura, S., & Takahashi, Y. (2015). Distribution and isolation of strains

belonging to the order Solirubrobacterales. *The Journal of Antibiotics, 68*(12), 763.

610   Sharma, M., & Reynnells, R. (2016). Importance of soil amendments: survival of bacterial

pathogens in manure and compost ussed as organic fertilizers. *Microbiology Spectrum,

612   4*(4).

Shoemaker, W. R., & Lennon, J. T. (2018). Evolution with a seed bank: The population genetic

614   consequences of microbial dormancy. *Evolutionary Applications, 11*(1), 60-75.

Smith, B., & Wilson, J. B. (1996). A consumer's guide to evenness indices. *Oikos*, 70-82.

616   van der Heijden, M. G., de Bruin, S., Luckerhoff, L., van Logtestijn, R. S., & Schlaeppi, K.

(2016). A widespread plant-fungal-bacterial symbiosis promotes plant biodiversity, plant

618   nutrition and seedling recruitment. *The ISME Journal, 10*(2), 389-399.

Vilgalys, R., & Hester, M. (1990). Rapid genetic identification and mapping of enzymatically

620   amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology,*

172(8), 4238-4246.

622   Wagg, C., Bender, S. F., Widmer, F., & van der Heijden, M. G. (2014). Soil biodiversity and soil

community composition determine ecosystem multifunctionality. *Proceedings of the*

624   *National Academy of Sciences, 111*(14), 5266-5270.

Wang, G., Jagadamma, S., Mayes, M. A., Schadt, C. W., Steinweg, J. M., Gu, L., & Post, W. M.

626   (2015). Microbial dormancy improves development and experimental validation of

ecosystem model. *The ISME Journal, 9*(1), 226-237.

628   Westcott, S. L., & Schloss, P. D. (2017). OptiClust, an improved method for assigning amplicon-

based sequence data to operational taxonomic units. *mSphere*, 2:e00073-00017.

630   Whitaker, J., Ostle, N., Nottingham, A. T., Ccahuana, A., Salinas, N., Bardgett, R. D., . . .

McNamara, N. P. (2014). Microbial community composition explains soil respiration

632   responses to changing carbon inputs along an Andes-to-Amazon elevation gradient.

Journal of Ecology, 102(4), 1058-1071.

 FIGURE CAPTIONS

Fig. 1. Conceptual framework depicting the role of the soil microbial seed bank in local plant-microbe interactions. Members of the soil microbial community can transition between the

metabolically active pool (M_a) and the seed bank (M_d) through initiation into dormancy and resuscitation into an active state. Both M_a and M_d pools have associated levels of microbial

abundance (N) and diversity (α). While only members in M_a can reproduce, the baseline rate of mortality (m_d) in M_d is reduced relative to the rate for M_a (m_a). Active microorganisms (M_a) and

plants (P) can directly interact in the soil with one another through a suite of mechanisms, such as mutualism, parasitism, root colonization, as well as the secretion of phytochemicals or other

signaling compounds. We envision that the M_d pool has minimal effects on plants. In contrast, this pool of inactive microorganisms may be resuscitated via processes including the release of

root exudates, which may vary in space and time.

Fig. 2. Influence of resuscitation promoting factor (Rpf) on (A) *Brassica rapa* shoot biomass, (B) root biomass, and (C) flower number produced per plant. We compared plant traits from

individuals that were exposed to weekly additions of recombinant Rpf (+Rpf) to those exposed to a protein buffer control (-Rpf). Black symbols represent the mean ± 95 % confidence intervals.

Grey symbols represent the individual observations.

Fig. 3. Effect of resuscitation promoting factor (Rpf) on soil microbial activity. Soil respiration was measured after applying Rpf (+Rpf) or protein buffer control (-Rpf) to soils on a weekly

basis. Symbols represent the mean ± 1 SE$_{\bar{x}}$.

Fig. 4. Effect of resuscitation promoting factor (Rpf) on (A) soil bacterial 16S rRNA copy

662   number, (B) fungal ITS copy number, and (C) the fungi : bacteria gene copy ratio. Gene copy

number was measured from soil after six weekly applications of recombinant Rpf (+Rpf) or

664   protein buffer control (-Rpf) treatment. Black symbols represent the mean ± 95 confidence

intervals. Grey symbols represent the individual observations.

Fig. 5. Principal Coordinate Analysis (PCoA) plot depicting composition of active (i.e., RNA)

668   and total (i.e., DNA) Actinobacteria from soil that were exposed to +Rpf and -Rpf treatments at

the end of a six-week experiment. The ellipses were generated by 'ordiellipse' function using the

670   standard deviation of PCoA point scores to visualize the spread of each treatment.

    FIGURES

Fig. 1

 Fig. 2

682 **Fig. 5**