Title: Antibiotic tolerance, persistence, and resistance of the evolved minimal cell, *Mycoplasma mycoides* JCVI-Syn3B.

Authors: Tahmina Hossain[1], Heather S. Deter[2], Eliza J. Peters[1], and Nicholas C. Butzin[1]*

Affiliations:
[1]Department of Biology and Microbiology; South Dakota State University; Brookings, SD, 57006; USA.
[2]Department of Bioengineering; University of Illinois Urbana-Champaign; Urbana, IL, 61801; USA.
*Corresponding author. Email: *nicholas.butzin@gmail.com*

Summary
Antibiotic resistance is a growing problem, but bacteria can evade antibiotic treatment via tolerance and persistence. Antibiotic persisters are a small subpopulation of bacteria that tolerate antibiotics due to a physiologically dormant state. Hence, persistence is considered a major contributor to the evolution of antibiotic-resistant and relapsing infections. Here, we used the synthetically developed minimal cell *Mycoplasma mycoides* JCVI-Syn3B to examine essential mechanisms of antibiotic survival. The minimal cell contains <500 genes, most genes are essential. Its reduced complexity helps to reveal hidden phenomenon and fundamental biological principles can be explored because of less redundancy and feedback between systems compared to natural cells. We found that Syn3B evolves antibiotic resistance to different types of antibiotics expeditiously. The minimal cell also tolerates and persists against multiple antibiotics, and contains a few already identified persister-related genes although lacking many systems previously linked to persistence (e.g. toxin-antitoxin systems, ribosome hibernation genes, etc.).

Keywords
Minimal cell, evolution, antibiotic tolerance, antibiotic persistence, antibiotic resistance

Introduction
   The evolution of antibiotic resistance is a pressing public health concern in the 21st century; antibiotic resistance results from one or more genetic mutations that counteract an antibiotic (Van den Bergh et al., 2016). Resistance is regarded as the primary culprit of antibiotic treatment failure, but bacteria also employ less publicized strategies to evade antibiotic treatment, namely antibiotic tolerance and persistence (Fisher et al., 2017a; Harms et al., 2016; Lewis, 2010). Tolerance can be hereditary or non-hereditary phenomena, where bacteria are able to outlive prolonged periods of antibiotic exposure by having a reduced killing rate (Balaban et al., 2019a). Persisters are a subpopulation of tolerant cells, which can sustain longer against antibiotic treatment in comparison to slow-growing tolerant cells by entering a metabolically repressed state (non-multiplying cells) (Cabral et al., 2018; Fisher et al., 2017b). Most antibiotics are only effective against growing cells, and by not growing, the persister population can survive longer even without being genetically resistant. Though the difference between tolerance and persistence is still debated, for this article, we used the definition based on a consensus statement released after a discussion panel with 121 researchers, which defined antibiotic persistence as a tolerant subpopulation of cells that result in a distinct phase of population decay (Balaban et al., 2019b). The difference between tolerance and persistence can be seen in a biphasic death curve,

where the tolerant population dies off quickly until persisters remain (and viable but nonculturable cells (VBNC); not explored in this work), dying at a slower rate than tolerant cells (Balaban et al., 2019a; Deter et al., 2019a) (Fig. 1. B). Two types of persisters may exist, triggered persisters (formed by environmental stimuli) and spontaneous persisters (generated stochastically) (Balaban et al., 2019a; Balaban et al., 2004; Sulaiman and Lam, 2020; Uruén et al., 2021), although spontaneous persister formation is controversial and sparse (Keren et al., 2004; Kim and Wood, 2016; Orman and Brynildsen, 2013). What makes persisters medically relevant is that they can revive and give rise to a new progeny after antibiotic treatment; the new progeny are genetically identical to the original susceptible kin, and this process plays a pivotal role in recurring infections (Wilmaerts et al., 2019b) (Fig. 1. A). Furthermore, evolutionary studies have determined that repeated exposure to antibiotics over many generations rapidly increases tolerance leading to antibiotic resistance (Balaban and Liu, 2019; Cohen et al., 2013; Fridman et al., 2014; Liu et al., 2020; Schumacher et al., 2015; Sulaiman and Lam, 2020; Van den Bergh et al., 2016).

The precise molecular mechanisms underlying persistence are still debated (multiple mechanisms are likely to exist), albeit several genes have been implicated (Wilmaerts et al., 2019a; Wu et al., 2015). Toxin-antitoxin (TA) systems have been implicated in persistence and were previously thought to be the key players for persistence (Lewis, 2010, 2012; Wang and Wood, 2011a). Particularly important is the HipAB TA system, because a toxin mutant, *hipA7,* increases persistence by impeding cellular growth in the absence of its cognate antitoxin *hipB* (Moyed and Bertrand, 1983). Subsequent studies also report that overexpression of other toxins from TA systems increased persistence (Correia et al., 2006; Kim and Wood, 2010; Korch and Hill, 2006). In contrast, new research found that the deletion of 10 TA systems (~22% of TA system in the genome) in *E. coli* does not have an apparent effect on persistence both at the population and single-cell levels (Goormaghtigh et al., 2018), though *E. coli* can have more than 45 TA systems (Horesh et al., 2018; Karp et al., 2014; Xie et al., 2018). Though this work supports that TA systems are not controlling persistence, the remaining 35 TA systems (~78%) could be controlling this phenomenon. Mauricio *et al.* studied persistence on Δ12TA *Salmonella enterica* and demonstrated that TA systems are dispensable (Pontes and Groisman, 2019), although this strain has 18 predicted TA systems based on TAfinder tool (Xie et al., 2018). Another knockout study on *Staphylococcus aureus* deleted three copies of type II TA system from Newman strain (Conlon et al., 2016). But further studies identified several type-I (sprG1/sprF1, sprG2/sprF2, sprG4/sprF4) (Riffaud et al., 2019) and type-II (SavRS) (Wen et al., 2018) TA system in HG003 and NCTC-8325 strains, respectively. These are the parental strains of Newman, and these TA genes are also found in Newman strain (Sassi et al., 2015). Moreover, a recent study showed that antitoxin sprF1 mediates translation inhibition to promote persister formation in *S. aureus* (Pinel-Marie et al., 2021). All these findings raise many questions about the implication of TA systems in bacterial persistence (Goormaghtigh et al., 2018; Kim and Wood, 2016; Pontes and Groisman, 2019; Tsilibaris et al., 2007). However, there are major limitations of studying TA systems due to their high abundance in most bacterial genomes; they often have redundant and overlapping functions and can have interdependencies within network clusters (Harms et al., 2018; Ronneau and Helaine, 2019; Wang et al., 2013). Additionally, TA systems respond to a variety of stresses (e.g. bacteriophage infection, oxidative stress, etc.), which creates a hurdle to probe their connection with any phenomenon related to stress response, namely antibiotic tolerance and persistence (Harms et al., 2018; Kang et al., 2018; Ronneau and Helaine, 2019). TA systems are also involved in other mechanisms in the cell that respond to

stress, such as the stringent and SOS responses (Ronneau and Helaine, 2019), virulence (De la Cruz et al., 2013), and the regulation of pathogen intracellular lifestyle in varied host cell types (Lobato-Marquez et al., 2015). Furthermore, new types of TA systems may be yet unidentified. These challenges could be resolved using a strain that lacks any TA systems (as we did in this work), but TA systems are naturally abundant and large-scale knockouts are both error-prone and labor-intensive.

Several other mechanisms have also been considered in persister research including SOS response, oxidative stress response, etc. (Trastoy et al., 2018; Wilmaerts et al., 2019a; Wilmaerts et al., 2019b). Two extensively studied phenomenon relates to persistence is cellular ATP levels and (p)ppGpp levels. The stress-related persistence mechanism is controlled by the stringent response, which is mediated by the accumulation of (p)ppGpp (stress sensing alarmone) (Harms et al., 2016). (p)ppGpp regulates many bionetworks (such as ribosome dimerization) that can cause cells to go into dormancy (Gaca et al., 2015; Song and Wood, 2020; Wood and Song, 2020). Another well-studied model, *ATP depletion increase persistence*, has drawn much attention in persister research (Conlon et al., 2016). This finding is consistent with REF (Pu et al., 2019), which demonstrated that lower ATP levels lead to protein aggregation and increased tolerance. This result is also coherent with our recently published data in which we established that interfering with protein degradation by forming a proteolytic queue at ClpXP will increase tolerance levels dramatically (Deter et al., 2019a) and transcriptomic analysis of queuing-tolerant population showed upregulation of genes related to metabolism and energy (Deter et al., 2020). However, another study reported that ppGpp and reduction of ATP are not essential for persistence (Pontes and Groisman, 2019). These contradictory studies are common in persister research, and we hypothesize these inconsistencies are due to the interconnection of gene networks surrounding persistence. One goal of our study is to clarify mechanisms using a minimal, simpler system. Research over the last several years has resulted in a lot of discussion concerning genes essential for persistence (Pontes and Groisman, 2019; Ronneau and Helaine, 2019; Wilmaerts et al., 2019a; Wilmaerts et al., 2019b). Since persistence and tolerance are present in phylogenetically diversified bacterial species (Meylan et al., 2018; Wilmaerts et al., 2019b), it is feasible that an underlying genetic mechanism is conserved in evolutionarily related microorganisms, and the most likely candidates are essential genes or the disruption of crucial networks.

In our recent work, we demonstrate that antibiotic tolerance in *Escherichia coli* is a whole-cell response and can occur through multiple pathways or networks, which may work simultaneously and cooperatively to survive against antibiotics (Deter et al., 2020). Our strategy to study the underlying mechanisms of persistence was to use a bacterial species that contains mainly essential genes and networks with reduced complexity and fewer networks. The minimal system can help us reveal hidden phenomenon (Glass et al., 2017) of antibiotic survival; for example, genes and networks that have previously been identified can be eliminated as causal if they are lacking from the genome, while genes present in Syn3B that were previously identified in other organisms can become the focus of the work. The reduced complexity has its limits as there are likely several method microbes use to survive antibiotics, and not all methods will be in a minimal system. With these limitations well-understood, we explored antibiotic tolerance, persistence, and resistance in the minimal cell *Mycoplasma mycoides* JCVI-Syn3B (called Syn3B throughout), a synthetic genetic background that contains the least number of genes and smallest genome of any known organism (Syn3B contains 473 genes and its genome is ~531 Kbp long, while *E. coli* contains >4,000 genes and its genome is ~4,600 Kbp long). Syn3B

genome consists predominantly of essential and a few non-essential and quasi-essential genes
(Hutchison et al., 2016). This microbe was designed to have the minimal number of genes and
networks with the expectation that many of the first principles of cellular life could be explored
in the simplest biological systems (Glass et al., 2017). We show that Syn3B populations contain
both persister and tolerant cells, and its genome contains a few previously identified persister-
related genes. This work establishes that many systems previously shown to be related to
bacterial persistence, such as TA systems and ribosome dimerization, are not essential for
persistence in the minimal cell, and demonstrates the effectiveness of using the minimal cell to
study antibiotic survival.
148
Result
Whole-genome analysis of the evolved minimal cell. Although the minimal genome (Syn3B)
was designed for ease of genetic manipulation, Syn3B grows slowly and to a low cell density.
We adjusted the original SP4 media (Tully et al., 1979) to address these limitations. The new
media, SP16, allows for faster cell growth and higher yields in liquid cultures. We noticed that
Syn3B growth was slightly better after every subculture. Thus, we did a cyclic subculture of
Syn3B in our optimized media by passing cells during logarithmic growth. After 26 passages, we
observed better growth and isolated a single colony named Syn3B P026 (Pass 26). This strain
has a shorter lag phase and an increased growth rate; the average doubling time of P026 is
approximately 2.5 hours compared to the ancestral Syn3B doubling time of ~6 hours under the
same conditions (Fig. S1.A-B). We performed a whole-genome analysis of Syn3B P026 to
examine the genetic basis of these changes. Most of the mutations (9 of 11) in P026 are
intergenic except one synonymous (*fakA*) and one non-synonymous (*dnaA*) (Table 1). *dnaA* is a
positive regulator of DNA replication initiation. Considering that bacterial growth rate is
dependent on the frequency of DNA replication and that *dnaA* mutants have been known to
result in over-replication (Skarstad and Boye, 1994), we hypothesize that the mutation of *dnaA* in
P026 is responsible for the higher growth rate. *fakA* is a fatty acid kinase, and there is less
evidence to suggest a direct connection to substantial alterations in bacterial growth. It is not
surprising that the DNA replication machine is not optimized in Syn3B because its genome was
built in a laboratory.

Table 1. A. Mutation of *Mycoplasma mycoides* JCVI-Syn3B and Whole Genome Sequence
(WGS) analysis identified the mutations (bold and underlined). All antibiotic-resistant mutant
strains were named based on the number of passes (P) in a specific antibiotic (K: Ksg, S: Strep,
C: Cip, SC: Strep-Cip). All antibiotic-resistant mutants were selected from two separate
evolutionary lineages, and named L1 for lineage 1 or L2 for linage 2. **B.** Annotation of mutated
genes.

A. Syn3B Strains	Intergenic mutation[a]	Mutation positions[b]	Genotype change	Amino acid substitution	Gene
Parent*					
P026	9	547	**G**CA → **A**CA	Ala → Thr	*dnaA*
		274928	TA**C** → TA**T**	Tyr → Tyr	*fakA[c]*
Ksg^R					
PK07-L1	8	479174	**C**CT → **A**CT	Pro → Thr	*rpoC*
		3322	G**G**A → G**A**A	Gly → Glu	*ksgA*
PK07-L2	8	479174	**C**CT → **A**CT	Pro → Thr	*rpoC*
		3322	G**G**A → G**A**A	Gly → Glu	*ksgA*
Strep^R					
PS04-L1	4	101452	A**A**A → A**G**A	Lys → Arg	*rpsL*
		33652	**G**GA → **A**GA	Gly → Arg	*CDS_6*
PS04-L2	3	101452	A**A**A → A**G**A	Lys → Arg	*rpsL*
		147803	A**G**C → A**T**C	Ser → Ile;	*rpsD*
		33652	**G**GA → **A**GA	Gly → Arg	*CDS 6*
Cip^R					
PC06-L1	4	7735	**G**AA → **A**AA	Glu→ Lys	*gyrA*
		305613	**G**AT → **A**AT	Asp→ Asn	*parC*
PC06-L2	4	7735	**G**AA → **A**AA	Glu→ Lys	*gyrA*
		305613	**G**AT → **A**AT	Asp→ Asn	*parC*
		304707	G**A**T → G**C**T	Asp→ Ala	*parE*
		33652	**G**GA → **A**GA	Gly → Arg	*CDS_6*
Strep^R-Cip^R					
PSC09-L1	4	7725	AG**T** → AG**G**	Ser→ Arg	*gyrA*
		101452	A**A**A → A**G**A	Lys → Arg	*rpsL*
		33652	**G**GA → **A**GA	Gly → Arg	*CDS_6*
		305602	A**G**T → A**T**T	Ser→ Ile	*parC*
PSC09-L2	4	7725	AG**T** → AG**G**	Ser→ Arg	*gyrA*
		101452	A**A**A → A**G**A	Lys → Arg	*rpsL*
		33652	**G**GA → **A**GA	Gly → Arg	*CDS_6*
		305602	A**G**T → A**T**T	Ser→ Ile	*parC*

*Parent: High growth rate mutant parent to all of the strains in table.
[a.]Mutation in the non-coding DNA sequences located between genes.
[b.]Genomic position numberings correspond to *Mycoplasma mycoides* JCVI-Syn3B.
[c.]Synonymous mutation: mutation does not change in the encoded amino acid sequence.

B. Mutated Gene	Annotation
dnaA	Chromosomal replication initiator protein DnaA
fakA	Dihydroxyacetone kinase
rpoC	DNA-directed RNA polymerase subunit beta
ksgA	16S rRNA (adenine(1518)-N(6)/adenine(1519)-N(6))- dimethyltransferase
rpsL	30S ribosomal protein S12
CDS_6	Unknown
rpsD	30S ribosomal protein S4
gyrA	DNA gyrase subunit A
parC	DNA topoisomerase 4 subunit A
parE	DNA topoisomerase 4 subunit B

Syn3B and evolved minimal cell P026 displays antibiotic tolerance & persistence. We assessed tolerance and persistence of Syn3B (ancestor strain) and Syn3B P026 cultures using two bactericidal antibiotics, ciprofloxacin (a fluoroquinolone) and streptomycin (an aminoglycoside). Ciprofloxacin inhibits DNA replication by targeting the activity of DNA gyrase and topoisomerase IV (Sanders, 1988), whereas streptomycin blocks protein synthesis by irreversibly binding to the 16s rRNA of the 30S ribosomal subunit (Luzzatto et al., 1968). Syn3B and P026 were grown to stationary phase and diluted into fresh media containing the antibiotics to observe population decay (see Methods). Both parent strain and P026 cultures showed a typical biphasic death curve from stationary phase; the death rate became slower at ~20-24 h treatment with both antibiotics compared to the earlier stage of population decay, which indicates Syn3B displays both tolerance and persistence (Fig. 1. C-D). It appears that the survival was higher in Syn3B ancestor strain than P026 for both antibiotic treatments. We posit that it could be due to the slower growth rate of ancestor strain, which is consistent with the literature, as it is already established by several research groups that growth rate has a strong correlation with antibiotic susceptibility (Abshire and Neidhardt, 1993; Lee et al., 2018; Pontes and Groisman, 2019; Tuomanen et al., 1986). As we observed tolerance and persistence in both the ancestor strain and Syn3B P026, we decided to do further analysis with Syn3B P026, because it was much easier to work.

Persister cells have been identified in native bacterial species in both stationary and exponential phase cultures, and we tested if this was also true for the minimal cell. P026 was grown to exponential phase and treated with antibiotics to determine whether the minimal genome showed similar biphasic death curve in exponential phase. As expected, we observed a similar biphasic killing curve in exponential phase with slightly lower survival in exponential phase compared to stationary phase for both antibiotics (Fig. 1. E). Moreover, we performed resistance assays through the course of this work to rule out the possibility of resistance instead of tolerance or persistence (see Methods), and no resistant colonies were identified. We then tested if the surviving population had increased persister levels after antibiotic treatment. After 48 h of antibiotic treatment, the culture was passed into fresh media, and when in stationary phase exposed to the same antibiotic under the same condition and, as expected, no significant difference between the two death curves was observed (Fig. S2).

Fig. 1. Population decay during antibiotic treatment shows tolerance and persistence of Syn3B. A. Persister cells survive antibiotic treatment, reestablish the population when antibiotics are removed and increase the odds of gaining antibiotic resistance (through mutation) that allows for growth in the presence of the antibiotic. **B.** A simplified model for defining tolerant and persister states. Antibiotic tolerance and persistence are defined in this study by using a biphasic death curve, where the first phase (faster death rate) is tolerance and the second phase (slower death rate) is persistence. **C.** Population decay of Syn3B (parent strain). Overnight cultures of Syn3B (parent strain) were grown to stationary phase, diluted to 1:10 and treated with streptomycin (100 µg/mL) or ciprofloxacin (1 µg/mL) and sampled over time. Error bars represent SEM (n ≥ 3). **D-E.** Population decay of Syn3B P026. Syn3B P026 cells were treated with streptomycin (100 µg/mL) or ciprofloxacin (1 µg/mL) and sampled over time. (**D**) Stationary phase cells were diluted 1/10 into fresh media containing antibiotics. Error bars represent SEM (n ≥ 6). (**E**) Exponential phase cells were treated streptomycin (100 µg/mL) or ciprofloxacin (1 µg/mL) and sampled over time. Error bars represent SEM (n ≥ 3). There is 100% survival at time zero, because percent survival is determined by the surviving CFU/ml compared to the CFU/ml at time zero.

At this point, we have demonstrated that tolerance and persistence can be detected in Syn3B cultures during both stationary and exponential phase, and this is consistent with native bacterial species. Another well-known phenotype of antibiotic survival is that the tolerant population increases with co-treatment of a bacteriostatic and bactericidal antibiotics (Lewin and Smith, 1988) (pre-treatment is another method (Kwan et al., 2013)). To use the minimal cell as a model of antibiotic survival, this phenomenon should also be observed. Bacteriostatic antibiotics can counteract the killing activity of the bactericidal antibiotic by arresting growth of the rapidly growing cells, which increases tolerance to the antibiotics (Kwan et al., 2013). Chloramphenicol, a bacteriostatic antibiotic inhibits translation by binding to bacterial ribosomes and inhibiting protein synthesis, thereby inhibiting bacterial growth (Volkov et al., 2019). We co-treated with the bacteriostatic antibiotic chloramphenicol and either streptomycin or ciprofloxacin. As expected, the overall percent survival with chloramphenicol co-treatment increased compared to streptomycin or ciprofloxacin alone after 24 h and 48 h (Fig. 2). This result supports that inhibition of translation by co-treating the cell with chloramphenicol increases tolerance in P026.

Fig. 2. Co-treatment of bactericidal antibiotic (streptomycin (Strep) or ciprofloxacin (Cip)) with a bacteriostatic antibiotic (chloramphenicol (Cm)) shows increased survival of cells for 24 h (left) and 48 h (right) in Syn3B P026. Error bars represent SEM ($n \geq 3$). *$p<0.05$. **$p<0.01$.

The minimal genome contains previously identified genes related to tolerance and persistence. The mechanisms of persister formation are complex, but recent studies identify several genes involved in this process (Cameron et al., 2018; Wilmaerts et al., 2019a), and a few of these genes are present in the minimal genome. We looked for known and predicted genes related to tolerance and persistence in the Syn3B genome (Table 2). During nutrient limitation, bacteria generally switch their gene expression profile from rapid growth to a survival state by (p)ppGpp levels (a hallmark of the stringent response), which is regulated by the RelA protein in *E. coli* (Boutte and Crosson, 2013). Another important gene in the stringent response is *phoU*, a global negative regulator, which is deactivated upon inorganic phosphate (Pi) starvation. Mutation or deletion of this gene in other microorganisms leads to a metabolically hyperactive state and results in decreased persistence (Li and Zhang, 2007; Zhang and Li, 2010). The SOS response is another signaling pathway that upregulates DNA repair functions, which appears to be linked to bacterial persistence. Genes related to the SOS response have been found upregulated in *E. coli* persister (e.g. *uvrA, uvrB*) (Keren et al., 2004), and other mutants (such as *xseB*) have lost their ability to induce detectable levels of persistence when pretreated with rifampicin, a bacteriostatic antibiotic (Cui et al., 2018).

Other literature advocates the connection of persistence with energy metabolism, although the outcomes are often inconsistent between models (Wilmaerts et al., 2019a). For example, the deletion of *atpA* (encoding for ATP synthase subunit alpha) decreases the persister fraction (Lobritz et al., 2015), but deletion of genes encoding other parts of the ATP synthase (*atpC and atpF*) increases the persister fraction (Girgis et al., 2012; Kiss, 2000). Several heat shock proteins, mainly proteases (e.g. *lon* (Pu et al., 2016; Wu et al., 2015)) and chaperons (e.g. *dnaK* (Wu et al., 2015)*, dnaJ* (Hansen et al., 2008)*, clpB* (Wu et al., 2015)), are related to persistence considering that deletion of those genes decreases persistence.

Another survival mechanism connected to persistence is trans-translation, which allows bacteria to recycle stalled ribosomes and tag unfinished polypeptides for degradation, which in *E. coli* requires tmRNA (encoded by *ssrA*) and a small protein (smpB). The deletion of these genes causes persister-reduction in *E.coli* (Liu et al., 2017; Wu et al., 2015; Yamasaki et al., 2020). Additional genes (*glyA* (Cui et al., 2018)*, galU* (Girgis et al., 2012), *trmE* (Cui et al., 2018), *efp* (Cui et al., 2018)*, ybeY* (Cui et al., 2018) etc.; see Table 2) are indirectly related to stress response or metabolism and have been reported to affect persistence. Overall, this short list (Table 2 compare to Table S1) of genes demonstrates that there are far fewer genes to explore in the minimal cell than any other known free-living microorganism on the planet. However, further exploration is needed to test if there is a relationship between genes in Table 2 to antibiotic survival in the minimal cell.

Table 2. Syn3B contains genes previously shown to be related to *E.coli* persistence/tolerance.

Gene	Protein/RNA	Major function	REF
dnaK, dnaJ, clpB	Chaperon	Global regulator	(Hansen et al., 2008; Wu et al., 2015)
phoU	Phosphate-specific transport system accessory protein	Global regulator	*(Li and Zhang, 2007)*
xseB	Exodeoxyribonuclease	DNA mismatch repair and recombination	(Cui et al., 2018)
uvrA, uvrB	Endonuclease	SOS response	(Cui et al., 2018; Keren et al., 2004; Wu et al., 2015)
relA	GTP pyrophosphokinase	Stringent response	(Wu et al., 2015)
lon	Protease	Protease	(Wu et al., 2015)
glyA	Serine hydroxymethyltransferase	Metabolism	(Cui et al., 2018)
atpA, atpC, atpF	ATP synthase	ATP synthesis	(Girgis et al., 2012; Kiss, 2000; Lobritz et al., 2015)
galU	UTP--glucose-1-phosphate uridylyltransferase	Lipopolysaccharide precursor synthesis	(Girgis et al., 2012)
trmE; efp; ybeY	tRNA modification GTPase; Elongation factor P; Endoribonuclease	Translation	(Cui et al., 2018)
ssrA; smpB	Transfer messenger RNA (tmRNA); Small Protein B	Trans-translation	(Wu et al., 2015; Yamasaki et al., 2020)

Evolution of the minimal cell against different types of antibiotic and whole-genome analysis of the resistant strain. Up until this point, we have focused on antibiotic tolerance and persistence. For the minimal system to be also useful as a model of antibiotic resistance, it must be able to evolve resistance without horizontal gene transfer (i.e. isolated from other organisms).

We hypothesize that due to lack of complexity and less control of mutation, simpler cells likely
evolve faster than more complex organisms; a simpler, slimmed-down genome allows for rapid
evolution to selective pressures. We selected different classes of bactericidal antibiotics
including streptomycin (Strep); ciprofloxacin (Cip); a combination of streptomycin and
ciprofloxacin (Strep-Cip), and one bacteriostatic antibiotic-kasugamycin (Ksg), to study the
evolution of antibiotic resistance in the minimal genome. We applied cyclic antibiotic treatments
where we repeatedly regrew and retreated the culture with the lethal dose of the same antibiotic
for a few cycles with two biological replicates (Fig. S3. B) to make resistant mutants. In seven
passes or less (nine passes with two antibiotics), the minimal genome rapidly evolved resistance
to all antibiotics tested. We isolated single colonies (two separate evolutionary lineages named
L1 for lineage 1 or L2 for linage 2) from each resistant mutants named Syn3B PS04 (4 Passes in
Streptomycin), PC06 (6 Passes in Ciprofloxacin), PSC09 (9 Passes in Streptomycin and
Ciprofloxacin), and PK07 (7 Passes in Kasugamycin). We then analyzed the whole genome for
mutations. Most mutations were in intergenic regions of the resistant mutants, similar to Syn3B
P026 (Table 1).
We observed Syn3B P026 quickly evolved against streptomycin, as it only took 4 passes to
become resistant. Only two non-synonymous mutations were found in both lineages. The first
one is in *rpsL* gene (encoding the S12 ribosomal protein), which is frequently identified in
streptomycin-resistant strains (Villellas et al., 2013), and the other one is *CDS_6* gene, the
function of which is not yet known. In PS04_L2, we also observed another mutation in the *rpsD*
gene (encoding the S4 ribosomal protein). We found that Syn3B P026 took 6 and 9 cycles to
gain resistance against ciprofloxacin and a combination of both streptomycin and ciprofloxacin,
respectively. In both replicates of PC06 we observed mutations in *gyrA* (encoding the DNA
gyrase subunit A) and *parC* (encoding DNA topoisomerase 4 subunit A). In PC06_L2, we
observed another two mutations, one in *parE* (encoding DNA topoisomerase 4 subunit B) and
another in *CDS_6* gene. For the combined streptomycin-ciprofloxacin resistant strains, both
lineages showed similar mutations in *rpsL, gyrA*, *parC*, and *CDS_6* genes. In the kasugamycin
resistant strain PK07, only two nonsynonymous mutations were found. The first one is in the
ksgA gene, which encodes a methylase that modifies 16S rRNA and inhibition of this protein by
the antibiotic kasugamycin causes susceptibility. Therefore, when *ksgA* is inactivated, cells
became kasugamycin-resistant (Mariscal et al., 2018; van Gemen et al., 1987). We reasoned that
the nonsynonymous mutation in *ksgA* in PK07 makes the protein inactive and able to confer
kasugamycin resistance. The other detected non-synonymous mutation is in *rpoC* (RNA
polymerase subunit).
328
Discussion
Overall, we have demonstrated that the minimal cell contains both antibiotic tolerant and
persistent subpopulations, and it can quickly evolve to gain resistance. We have successfully
generated an evolved strain of the minimal cell Syn3B P026, which has a better growth rate than
the ancestral strain and overcomes one of the major difficulties of working with the minimal cell.
We show that Syn3B can be used as a model for studying antibiotic survival, especially when we
consider that the minimal genome's core proteins are present in many other microorganisms
including human pathogens. Syn3B has over 50% similarity to human pathogens that have been
identified as a concern with respect to the growing antibiotic-resistance problem (Table
S2)(Centers for Disease Control and Preventions, 2019). For example, out of 455 protein-
encoding genes of Syn3B, 338 of them (74%) have homologs to *S. aureus* NCTC 8325 proteins.

S. aureus causes meningitis and pneumonia, and antibiotic resistance is a major problem of this
organism (Foster, 2017).
Furthermore, we also find that Syn3B genome lacks homologs of several genes and pathways
reported in earlier research to be related to tolerance and persistence (Table S1). For instance,
TA systems are often implicated for persistence and are still referenced as causing persistence
(Moreno-del Álamo et al., 2020; Riffaud et al., 2020; Shenkutie et al., 2020), although recent
research provided evidence of a lack of causation between persister formation and TA system
(Conlon et al., 2016; Pontes and Groisman, 2019). In this study, we identify no known TA
systems or homologous genes based on the results from TADB 2.0 (a database designed to
search for TA systems based on homology) (Xie et al., 2018) in Syn3B. Our study clearly shows
that TA systems are not required for tolerance or persistence in the minimal cell, and it seems
likely that most bacteria do not require them either. It is not surprising that minimal genome does
not contain TA systems because the genome was designed by researchers and not subject to the
natural environment. Natural TA systems are often described as "addictive" systems that are hard
to lose once acquired; they often have overlapping toxin and antitoxin genes making mutations
less likely to be selected. Curiously, we observed some overlapping genes that are not at all
similar to TA systems (they are also much longer in sequence than traditional TA systems, and
are not homologous to any known TA system), and whose functions are not yet defined (Table
S3).
Recently, the Wood's group (Song and Wood, 2020; Wood and Song, 2020) proposed the
ppGpp ribosome dimerization persister (PRDP) model for entering and exiting the persister state
where ppGpp induces *hpf, rmf*, and *raiA*, which converts active ribosomes into their inactive
state (such as 100s ribosome) to reduce translation and consequently cells enter into persistence.
Upon removal of antibiotic stress, cAMP level is reduced and *HflX* production is stimulated,
which dissociates inactive 100S ribosomes into active 70S ribosomes and growth resumes.
However, we did not detect homologs to the required genes, *raiA, rmf, hpf, and hflX*, for
ribosome dimerization in the minimal cell genome, which means ribosome dimerization is not an
essential mechanism for this organism (or another unknown mechanism for ribosome
dimerization exists in Syn3B). (p)ppGpp may still play a role in Syn3B persistence, the genome
contains *relA* (converts ATP and GTP to (p)ppGpp; Table 2) and a potentially fruitful study
would be to study the effects on survival by altering (p)ppGpp of Syn3B cultures.
In this strain, it is evident that TA systems and ribosome hibernation genes are not required
for bacterial tolerance or persistence. Moreover, there are far fewer genes (less than 20 genes)
present in Syn3B, which have been shown related to persistence. Thus, if specific genes or
regulons are responsible for persistence, then Syn3B should be very useful in identifying them in
future studies because it has a minimal number of genes. It is possible that persister formation
and maintenance results from slowed or disrupted cellular networks (this is the hypothesis we
most prescribe to), rather than the activity of specific genes or regulons. If tolerance/persistence
results from slowed or disrupted cellular networks, then Syn3B is an ideal model organism to use
because it has a minimal number of networks. While different genes and networks are likely
responsible for persistence depending on the antibiotic, strain, or bacterial species, identifying
genes in this minimal system should be applicable to a number of other microorganisms
including the pathogen *Mycoplasma mycoides* from which Syn3B was derived. Regardless of
how cells enter the dormant state, Syn3B provides a new model to study the genes and networks
that allow cells to survive antibiotic treatment and could pave the way of finding new drugs that
target the persister and tolerant subpopulations.

Limitations of the study

We observed that despite controlling methodology to the greatest extent possible, persister levels varied considerably (far more than observed in our previous work with *E. coli*) (Deter et al., 2019a) between experiments for both streptomycin and ciprofloxacin treatments (Table S4). We hypothesize that this variability might be due to the stochastic fluctuations (noise) in gene expression levels, which results in protein level variations even among genetically identical cells in a similar environment (Soltani et al., 2016). We expect that gene expression and protein production to be more erratic in Syn3B compared to natural microorganisms because this cell has a designed genome lacking many control mechanisms and did not evolve to achieve some level of internal cellular "equilibrium" like native cells have.

Methods

Microbial strains and media. *Mycoplasma mycoides* JCVI-Syn3B (Hutchison et al., 2016) and its derivatives were used in this study. Syn3B was a gift from Dr. John I. Glass from J. Craig Venter Institute, La Jolla, CA, USA. For evolution and antibiotic survival assays, cells were cultured at 37°C in SP16 media (57.5% 2X P1, 10.0% P4, 17.0% FBS, tetracycline 0.4% and vitamin B1 0.5%) (see Table S5 for details), which was developed based on SP4 media (Tully et al., 1979). All cultures were plated in SP4 agarose media (0.55% agarose) for colony counts. Note that agar is not used because it inhibits growth.

Evolution by serial passage. Syn3B cultures were grown in 3 mL tubes at 37°C overnight in SP16 media. Overnight exponential cultures were serially passaged after each cycle of growth by transferring 30 µL of culture into 3 mL fresh media to initiate the next cycle of growth, and the cycles continued until a satisfactory growth rate was observed (Fig. S3. A). The culture was plated after 26 passages and a single colony was isolated, P026.

Genome extraction, whole-genome sequencing, and identification of mutations. For whole-genome sequencing (WGS), a single colony of the evolved strains Syn3B P026 and all the resistant mutants (PK07_L1, PK07_L2, PS04_L1, PS04_L2, PC06_L1, PC06_L2, PSC09_L1, PSC09_L2) were isolated and inoculated into SP16 media for 24 h at 37°C. Next, genomic DNA was harvested and purified using Genomic DNA Purification Kit (ThermoFisher) in accordance with the manufacturer's instructions. For quality checks, DNA purity and concentration were assessed by gel electrophoresis and Qubit Fluorimeter prior to sending for sequencing. Novogene Ltd. sequenced the genomes using paired-end Illumina sequencing at 2×150 bp read length and 350 bp insert size. A total amount of ~1 µg of DNA per sample was used as input material for the DNA sample preparation. Sequencing libraries were generated from the qualified DNA samples using the NEB Next Ultra DNA Library Prep Kit for Illumina (NEB, USA) following the manufacturer's protocol. For data processing, the original sequence data were transformed into raw sequenced reads by CASAVA base calling and stored in FASTQ (fq) format. For subsequent analysis, the raw data were filtered off the reads containing adapter and low-quality reads to obtain clean data. The resequencing analysis was based on reads mapping to reference genome of Syn3B by BWA software (Li and Durbin, 2009). SAMTOOLS (Li et al., 2009) was used to detect single nucleotide polymorphism (SNP) and InDels.

We also used Oxford Nanopore Technologies (ONT) MinION long-read sequencer to search for large insertions or gene duplication in all the evolved strains. ONT libraries were prepared using *Ligation kit* (SQK-LSK109) according to the manufacturer's instructions. R 9.5 flow cell

432   (FLO-MIN107, ONT) was used for sequencing based on manufacture protocol. The flow cell
433   was mounted on a MinION Mk 1B device (ONT) for sequencing with the MinKNOW versions
434   19.12.5_Sequencing_Run_FLO-MIN107_SQK-LSK109 script. Then, reads were mapped
435   against reference genome Syn3B using Geneious Prime software version 2020.1.
436   (*https://www.geneious.com*). No large insertions were found in the sequenced genomes.

Minimum inhibitory concentration (MIC) tests. Overnight cultures were serially diluted and
439   plated onto SP4 agarose plates containing different concentrations of antibiotics (Strep, Cip, Cm,
440   and Ksg) to determine the MIC of each antibiotic. Plates were incubated 4-5 days at 37°C before
441   colony counts. MIC values were defined in this study as the lowest antibiotic concentration that
442   inhibit the growth of Syn3B (See Fig. S4).

Antibiotic survival assays. The schematic of the antibiotic survival assay from a stationary
phase culture is shown in Fig. S3. C. Briefly, an overnight culture was diluted 1:100 into pre-
warmed media and grown to exponential phase (OD_{600} 0.1-0.3). Next, the culture was separated
into three flasks (for three biological replicates) and grown at 37˚C until it reached stationary
phase (OD 0.45-0.55, which takes ~3-6 h). After that, each culture was diluted 1:10 into 100
µg/mL streptomycin (10X MIC) or 1 µg/mL (10X MIC) ciprofloxacin containing pre-warmed
media and kept at 37˚C shaking at 250 rpm for 72 h. Samples were taken at different time points
until 72 h for the time-kill assays. To remove the antibiotic before plating, 100 µl of each sample
was washed with 1.9 mL ice-cold SP16 media and collected by centrifugation (16,000 rpm for 3
min at 4˚C). Cells were then resuspended and serially diluted into ice-cold SP16 media and
plated to count the colony-forming units (CFU). Persisters were quantified by comparing CFUs
per milliliter (CFU/ml) before antibiotic treatment to CFU/ml after antibiotic treatment. Plates
were incubated at 37°C for 4-5 days, then scanned using a flatbed scanner (Datla et al., 2017;
Deter et al., 2019a; Levin-Reisman et al., 2017). Custom scripts were used to identify and count
bacterial colonies (Deter et al., 2019a; Deter et al., 2019b) used to calculate CFU/ml and
persister frequency. Over 200 colonies were streaked periodically into antibiotic-containing
plates to test for antibiotic-resistant mutants. Also, antibiotic-treated culture was washed after 48
h, regrew to stationary phase, and exposed the culture again in the same antibiotic treatment for
48 h and plated after 24 h and 48 h to observe the difference between the first (Strep/Cip) and
second death curve (Strep re-exposed/Cip re-exposed) in both antibiotic treatment.

Evolution through cyclic antibiotic treatment. Stationary phase culture was exposed to 100
µg/mL streptomycin (∼10× MIC), 1 µg/mL ciprofloxacin (∼10× MIC), a combination of
streptomycin (100 µg/mL) and ciprofloxacin (1 µg/mL) and 300 µg/mL kasugamycin (∼10×
MIC) antibiotic for 24 h, then antibiotic-containing medium was removed by washing twice with
SP16 medium (10 min of centrifugation at 7000 g at 4°C). Finally, the culture was resuspended
in 10 mL of fresh SP16 media and grown overnight at 37 °C. After every cycle of antibiotic
treatment, the tolerance phenotype was observed. Finally, we isolated single colony from
evolved populations from streptomycin (PS04), ciprofloxacin (PC06), combination of
streptomycin and ciprofloxacin (PSC09) and kasugamycin (PK07) treatment after four, six, nine
and seven cycle, respectively, and then their genomes were sequenced. Two different
evolutionary lineages were used for all evolved populations.

Determination of growth and doubling times. Overnight cultures were diluted into OD 0.1 (measured in SpectronicTM 200E) and 30 µL of diluted cultures were inoculated into individual wells containing 270 µL of SP16 media in a 96-Well Optical-Bottom Plate with Polymer Base (ThermoFisher) to measure OD at 600 nm using FLUOstar Omega microplate reader. Doubling time was determined by the linear regression of the natural logarithm of the OD over time during exponential growth as described in REF (Widdel, 2007).

Statistical analysis. All data is presented in the manuscript as mean ± SEM of at least three independent biological replicates. Statistical significance was assessed using an f-test to determine variance ($p < 0.05$ was considered to have significant variance), followed by a two-tailed t-test with unequal variances (if F statistic > F critical value) or equal variances (if F statistic < F critical value).

Acknowledgments
Mycoplasma mycoides JCVI-Syn3B was a generous gift from Dr. John I. Glass from J. Craig Venter Institute (JCVI), La Jolla, CA, USA. This work is supported by the National Science Foundation award Numbers 1922542 and 1849206, and by a USDA National Institute of Food and Agriculture Hatch project grant number SD00H653-18/project accession no. 1015687.

Author contributions
T.H. wrote the manuscript and performed most experiments. H.S.D. repeated streptomycin persister assay and developed custom code for colony counting, growth rate, and statistical analyses. E.P. performed cyclic antibiotic treatment experiments. N.C.B. planned and directed the project. All authors contributed to discussing and editing the manuscript.

Declaration of interests
The authors declare no competing interests.

Data and materials availability
All data that supports the findings of this study are available from the corresponding author upon request. Whole genomes data of P026, PK07_L1, PK07_L2, PS04_L1, PS04_L2, PC06_L1, PC06_L2, PSC09_L1 and PSC09_L2 strains has been deposited on the NCBI Genome Bank in BioProject PRJNA635211. The code used for colony counting is available on GitHub at *https://github.com/hdeter/CountColonies.*

References
Abshire, K.Z., and Neidhardt, F.C. (1993). Growth rate paradox of Salmonella typhimurium within host macrophages. Journal of bacteriology *175*, 3744-3748.
Balaban, N.Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D.I., Brynildsen, M.P., Bumann, D., Camilli, A., Collins, J.J*., et al.* (2019a). Definitions and guidelines for research on antibiotic persistence. Nature Reviews Microbiology *17*, 441-448.
Balaban, N.Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D.I., Brynildsen, M.P., Bumann, D., Camilli, A., Collins, J.J*., et al.* (2019b). Definitions and guidelines for research on antibiotic persistence. Nat. Rev. Microbiol.

Balaban, N.Q., and Liu, J. (2019). Evolution Under Antibiotic Treatments: Interplay Between
Antibiotic Persistence, Tolerance, and Resistance. In Persister Cells and Infectious Disease, pp.
1-17.
Balaban, N.Q., Merrin, J., Chait, R., Kowalik, L., and Leibler, S. (2004). Bacterial persistence as
a phenotypic switch. Science *305*, 1622-1625.
Boutte, C.C., and Crosson, S. (2013). Bacterial lifestyle shapes stringent response activation.
Trends in microbiology *21*, 174-180.
Cabral, D.J., Wurster, J.I., and Belenky, P. (2018). Antibiotic Persistence as a Metabolic
Adaptation: Stress, Metabolism, the Host, and New Directions. Pharmaceuticals (Basel) *11*.
Cameron, D.R., Shan, Y., Zalis, E.A., Isabella, V., and Lewis, K. (2018). A genetic determinant
of persister cell formation in bacterial pathogens. Journal of bacteriology *200*, e00303-00318.
Centers for Disease Control and Preventions (2019). Biggest threats and data.
Christensen, S.K., Maenhaut-Michel, G., Mine, N., Gottesman, S., Gerdes, K., and Van
Melderen, L. (2004). Overproduction of the Lon protease triggers inhibition of translation in
Escherichia coli: involvement of the yefM-yoeB toxin-antitoxin system. Molecular microbiology
51, 1705-1717.
Cohen, N.R., Lobritz, M.A., and Collins, J.J. (2013). Microbial persistence and the road to drug
resistance. Cell host & microbe *13*, 632-642.
Conlon, B.P., Rowe, S.E., Gandt, A.B., Nuxoll, A.S., Donegan, N.P., Zalis, E.A., Clair, G.,
Adkins, J.N., Cheung, A.L., and Lewis, K. (2016). Persister formation in Staphylococcus aureus
is associated with ATP depletion. Nature microbiology *1*, 1-7.
Correia, F.F., D'Onofrio, A., Rejtar, T., Li, L., Karger, B.L., Makarova, K., Koonin, E.V., and
Lewis, K. (2006). Kinase activity of overexpressed HipA is required for growth arrest and
multidrug tolerance in Escherichia coli. Journal of bacteriology *188*, 8360-8367.
Cui, P., Niu, H., Shi, W., Zhang, S., Zhang, W., and Zhang, Y. (2018). Identification of genes
involved in bacteriostatic antibiotic-induced persister formation. Frontiers in microbiology *9*,
413.
Datla, U.S., Mather, W.H., Chen, S., Shoultz, I.W., Täuber, U.C., Jones, C.N., and Butzin, N.C.
(2017). The spatiotemporal system dynamics of acquired resistance in an engineered
microecology. Scientific reports *7*, 1-9.
De la Cruz, M.A., Zhao, W., Farenc, C., Gimenez, G., Raoult, D., Cambillau, C., Gorvel, J.P.,
and Meresse, S. (2013). A toxin-antitoxin module of Salmonella promotes virulence in mice.
PLoS pathogens *9*, e1003827.
Deter, H.S., Abualrahi, A.H., Jadhav, P., Schweer, E.K., Ogle, C.T., and Butzin, N.C. (2019a).
Proteolytic queues at ClpXP increase antibiotic tolerance. ACS Synthetic Biology.
Deter, H.S., Dies, M., Cameron, C.C., Butzin, N.C., and Buceta, J. (2019b). A Cell
Segmentation/Tracking Tool Based on Machine Learning. In Computer Optimized Microscopy
(Springer), pp. 399-422.
Deter, H.S., Hossain, T., and Butzin, N.C. (2020). Antibiotic tolerance is associated with a broad
and complex transcriptional response in E. coli. bioRxiv, 2020.2008.2027.270272.
Dörr, T., Vulić, M., and Lewis, K. (2010). Ciprofloxacin causes persister formation by inducing
the TisB toxin in Escherichia coli. PLoS Biol *8*, e1000317.
Fisher, R.A., Gollan, B., and Helaine, S. (2017a). Persistent bacterial infections and persister
cells. Nature Reviews Microbiology *15*, 453.
Fisher, R.A., Gollan, B., and Helaine, S. (2017b). Persistent bacterial infections and persister
cells. Nat. Rev. Microbiol. *15*, 453-464.

Foster, T.J. (2017). Antibiotic resistance in Staphylococcus aureus. Current status and future
prospects. FEMS microbiology reviews *41*, 430-449.
Fridman, O., Goldberg, A., Ronin, I., Shoresh, N., and Balaban, N.Q. (2014). Optimization of lag
time underlies antibiotic tolerance in evolved bacterial populations. Nature *513*, 418-421.
Gaca, A.O., Colomer-Winter, C., and Lemos, J.A. (2015). Many means to a common end: the
intricacies of (p) ppGpp metabolism and its control of bacterial homeostasis. Journal of
bacteriology *197*, 1146-1156.
Garcia Rodriguez, G., Talavera Perez, A., Konijnenberg, A., Sobott, F., Michiels, J., and Loris,
R. (2020). The Escherichia coli RnlA–RnlB toxin–antitoxin complex: production,
characterization and crystallization. Acta Crystallographica Section F: Structural Biology
Communications *76*.
Girgis, H.S., Harris, K., and Tavazoie, S. (2012). Large mutational target size for rapid
emergence of bacterial persistence. Proceedings of the National Academy of Sciences *109*,
12740-12745.
Glass, J.I., Merryman, C., Wise, K.S., Hutchison, C.A., 3rd, and Smith, H.O. (2017). Minimal
Cells-Real and Imagined. Cold Spring Harb Perspect Biol *9*.
Goormaghtigh, F., Fraikin, N., Putrins, M., Hallaert, T., Hauryliuk, V., Garcia-Pino, A., Sjodin,
A., Kasvandik, S., Udekwu, K., Tenson, T.*, et al.* (2018). Reassessing the Role of Type II Toxin-
Antitoxin Systems in Formation of Escherichia coli Type II Persister Cells. mBio *9*.
Hansen, S., Lewis, K., and Vulić, M. (2008). Role of global regulators and nucleotide
metabolism in antibiotic tolerance in Escherichia coli. Antimicrobial agents and chemotherapy
52, 2718-2726.
Hansen, S., Vulić, M., Min, J., Yen, T.-J., Schumacher, M.A., Brennan, R.G., and Lewis, K.
(2012). Regulation of the Escherichia coli HipBA toxin-antitoxin system by proteolysis. PloS
one *7*, e39185.
Harms, A., Brodersen, D.E., Mitarai, N., and Gerdes, K. (2018). Toxins, Targets, and Triggers:
An Overview of Toxin-Antitoxin Biology. Mol Cell *70*, 768-784.
Harms, A., Maisonneuve, E., and Gerdes, K. (2016). Mechanisms of bacterial persistence during
stress and antibiotic exposure. Science *354*.
Horesh, G., Harms, A., Fino, C., Parts, L., Gerdes, K., Heinz, E., and Thomson, N.R. (2018).
SLING: a tool to search for linked genes in bacterial datasets. Nucleic Acids Res.
Hu, Y., Kwan, B.W., Osbourne, D.O., Benedik, M.J., and Wood, T.K. (2015). Toxin YafQ
increases persister cell formation by reducing indole signalling. Environmental microbiology *17*,
1275-1285.
Hutchison, C.A., 3rd, Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Ellisman,
M.H., Gill, J., Kannan, K., Karas, B.J., Ma, L.*, et al.* (2016). Design and synthesis of a minimal
bacterial genome. Science *351*, aad6253.
Kang, S.M., Kim, D.H., Jin, C., and Lee, B.J. (2018). A Systematic Overview of Type II and III
Toxin-Antitoxin Systems with a Focus on Druggability. Toxins (Basel) *10*.
Karp, P.D., Weaver, D., Paley, S., Fulcher, C., Kubo, A., Kothari, A., Krummenacker, M.,
Subhraveti, P., Weerasinghe, D., Gama-Castro, S.*, et al.* (2014). The EcoCyc Database. EcoSal
Plus *6*, ESP-0009-2013: 0001-0013.
Keren, I., Shah, D., Spoering, A., Kaldalu, N., and Lewis, K. (2004). Specialized persister cells
and the mechanism of multidrug tolerance in Escherichia coli. Journal of bacteriology *186*, 8172-
8180.

Kim, J.-S., and Wood, T.K. (2016). Persistent persister misperceptions. Frontiers in microbiology *7*, 2134.

Kim, Y., and Wood, T.K. (2010). Toxins Hha and CspD and small RNA regulator Hfq are involved in persister cell formation through MqsR in Escherichia coli. Biochemical and biophysical research communications *391*, 209-213.

Kiss, P. (2000). Results concerning products and sums of the terms of linear recurrences. Acta Acad. Agriensis Sectio Math *27*, 1-7.

Korch, S.B., Henderson, T.A., and Hill, T.M. (2003). Characterization of the hipA7 allele of Escherichia coli and evidence that high persistence is governed by (p)ppGpp synthesis. Mol Microbiol *50*, 1199-1213.

Korch, S.B., and Hill, T.M. (2006). Ectopic overexpression of wild-type and mutant hipA genes in Escherichia coli: effects on macromolecular synthesis and persister formation. Journal of bacteriology *188*, 3826-3836.

Kwan, B.W., Valenta, J.A., Benedik, M.J., and Wood, T.K. (2013). Arrested protein synthesis increases persister-like cell formation. Antimicrob Agents Chemother *57*, 1468-1473.

Lee, A.J., Wang, S., Meredith, H.R., Zhuang, B., Dai, Z., and You, L. (2018). Robust, linear correlations between growth rates and β-lactam–mediated lysis rates. Proceedings of the National Academy of Sciences *115*, 4069-4074.

Levin-Reisman, I., Ronin, I., Gefen, O., Braniss, I., Shoresh, N., and Balaban, N.Q. (2017). Antibiotic tolerance facilitates the evolution of resistance. Science *355*, 826-830.

Lewin, C.S., and Smith, J.T. (1988). Bactericidal mechanisms of ofloxacin. J Antimicrob Chemother *22 Suppl C*, 1-8.

Lewis, K. (2010). Persister cells. Annual review of microbiology *64*, 357-372.

Lewis, K. (2012). Persister cells: molecular mechanisms related to antibiotic tolerance. Handb Exp Pharmacol, 121-133.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. bioinformatics *25*, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009). The sequence alignment/map format and SAMtools. Bioinformatics *25*, 2078-2079.

Li, Y., and Zhang, Y. (2007). PhoU is a persistence switch involved in persister formation and tolerance to multiple antibiotics and stresses in Escherichia coli. Antimicrobial agents and chemotherapy *51*, 2092-2099.

Liu, J., Gefen, O., Ronin, I., Bar-Meir, M., and Balaban, N.Q. (2020). Effect of tolerance on the evolution of antibiotic resistance under drug combinations. Science *367*, 200-204.

Liu, S., Wu, N., Zhang, S., Yuan, Y., Zhang, W., and Zhang, Y. (2017). Variable persister gene interactions with (p) ppGpp for persister formation in Escherichia coli. Frontiers in microbiology *8*, 1795.

Lobato-Marquez, D., Moreno-Cordoba, I., Figueroa, V., Diaz-Orejas, R., and Garcia-del Portillo, F. (2015). Distinct type I and type II toxin-antitoxin modules control Salmonella lifestyle inside eukaryotic cells. Sci Rep *5*, 9374.

Lobritz, M.A., Belenky, P., Porter, C.B., Gutierrez, A., Yang, J.H., Schwarz, E.G., Dwyer, D.J., Khalil, A.S., and Collins, J.J. (2015). Antibiotic efficacy is linked to bacterial cellular respiration. Proceedings of the National Academy of Sciences *112*, 8173-8180.

Luzzatto, L., Apirion, D., and Schlessinger, D. (1968). Mechanism of action of streptomycin in
E. coli: interruption of the ribosome cycle at the initiation of protein synthesis. Proceedings of
the National Academy of Sciences of the United States of America *60*, 873.
Mariscal, A.M., Kakizawa, S., Hsu, J.Y., Tanaka, K., Gonzalez-Gonzalez, L., Broto, A., Querol,
E., Lluch-Senar, M., Pinero-Lambea, C., Sun, L.*, et al.* (2018). Tuning Gene Activity by
Inducible and Targeted Regulation of Gene Expression in Minimal Bacterial Cells. ACS Synth
Biol *7*, 1538-1552.
Meylan, S., Andrews, I.W., and Collins, J.J. (2018). Targeting antibiotic tolerance, pathogen by
pathogen. Cell *172*, 1228-1238.
Moreno-del Álamo, M., Marchisone, C., and Alonso, J.C. (2020). Antitoxin ε Reverses Toxin ζ-
Facilitated Ampicillin Dormants. Toxins *12*, 801.
Moyed, H.S., and Bertrand, K.P. (1983). hipA, a newly recognized gene of Escherichia coli K-12
that affects frequency of persistence after inhibition of murein synthesis. Journal of bacteriology
155, 768-775.
Orman, M.A., and Brynildsen, M.P. (2013). Dormancy is not necessary or sufficient for bacterial
persistence. Antimicrobial agents and chemotherapy *57*, 3230-3239.
Pinel-Marie, M.-L., Brielle, R., Riffaud, C., Germain-Amiot, N., Polacek, N., and Felden, B.
(2021). RNA antitoxin SprF1 binds ribosomes to attenuate translation and promote persister cell
formation in Staphylococcus aureus. Nature Microbiology, 1-12.
Pontes, M.H., and Groisman, E.A. (2019). Slow growth determines nonheritable antibiotic
resistance in Salmonella enterica. Sci Signal *12*.
Pu, Y., Li, Y., Jin, X., Tian, T., Ma, Q., Zhao, Z., Lin, S.Y., Chen, Z., Li, B., Yao, G.*, et al.*
(2019). ATP-Dependent Dynamic Protein Aggregation Regulates Bacterial Dormancy Depth
Critical for Antibiotic Tolerance. Mol. Cell *73*, 143-156 e144.
Pu, Y., Zhao, Z., Li, Y., Zou, J., Ma, Q., Zhao, Y., Ke, Y., Zhu, Y., Chen, H., and Baker, M.A.
(2016). Enhanced efflux activity facilitates drug tolerance in dormant bacterial cells. Molecular
cell *62*, 284-294.
Riffaud, C., Pinel-Marie, M.-L., and Felden, B. (2020). Cross-Regulations between Bacterial
Toxin–Antitoxin Systems: Evidence of an Interconnected Regulatory Network? Trends in
Microbiology.
Riffaud, C., Pinel-Marie, M.-L., Pascreau, G., and Felden, B. (2019). Functionality and cross-
regulation of the four SprG/SprF type I toxin–antitoxin systems in Staphylococcus aureus.
Nucleic acids research *47*, 1740-1758.
Ronneau, S., and Helaine, S. (2019). Clarifying the link between toxin-antitoxin modules and
bacterial persistence. Journal of molecular biology.
Sanders, C.C. (1988). Ciprofloxacin: in vitro activity, mechanism of action, and resistance.
Reviews of infectious diseases *10*, 516-527.
Sassi, M., Augagneur, Y., Mauro, T., Ivain, L., Chabelskaya, S., Hallier, M., Sallou, O., and
Felden, B. (2015). SRD: a Staphylococcus regulatory RNA database. Rna *21*, 1005-1017.
Schumacher, M.A., Balani, P., Min, J., Chinnam, N.B., Hansen, S., Vulić, M., Lewis, K., and
Brennan, R.G. (2015). HipBA–promoter structures reveal the basis of heritable multidrug
tolerance. Nature *524*, 59-64.
Shenkutie, A.M., Yao, M.Z., Siu, G.K.-h., Wong, B.K.C., and Leung, P.H.-m. (2020). Biofilm-
Induced Antibiotic Resistance in Clinical Acinetobacter baumannii Isolates. Antibiotics *9*, 817.
Skarstad, K., and Boye, E. (1994). The initiator protein DnaA: evolution, properties and
function. Biochimica et Biophysica Acta (BBA)-Gene Structure and Expression *1217*, 111-130.

Soltani, M., Vargas-Garcia, C.A., Antunes, D., and Singh, A. (2016). Intercellular Variability in Protein Levels from Stochastic Expression and Noisy Cell Cycle Processes. PLoS Comput Biol *12*, e1004972.

Song, S., and Wood, T.K. (2020). ppGpp ribosome dimerization model for bacterial persister formation and resuscitation. Biochemical and Biophysical Research Communications *523*, 281-286.

Sulaiman, J.E., and Lam, H. (2020). Proteomic Investigation of Tolerant Escherichia coli Populations from Cyclic Antibiotic Treatment. J Proteome Res.

Sun, C., Guo, Y., Tang, K., Wen, Z., Li, B., Zeng, Z., and Wang, X. (2017). MqsR/MqsA toxin/antitoxin system regulates persistence and biofilm formation in Pseudomonas putida KT2440. Frontiers in microbiology *8*, 840.

Trastoy, R., Manso, T., Fernandez-Garcia, L., Blasco, L., Ambroa, A., Del Molino, M.P., Bou, G., Garcia-Contreras, R., Wood, T., and Tomas, M. (2018). Mechanisms of bacterial tolerance and persistence in the gastrointestinal and respiratory environments. Clinical Microbiology Reviews *31*.

Tripathi, A., Dewan, P.C., Barua, B., and Varadarajan, R. (2012). Additional role for the ccd operon of F-plasmid as a transmissible persistence factor. Proceedings of the National Academy of Sciences *109*, 12497-12502.

Tsilibaris, V., Maenhaut-Michel, G., Mine, N., and Van Melderen, L. (2007). What Is the Benefit to Escherichia coli of Having Multiple Toxin-Antitoxin Systems in Its Genome? Journal of Bacteriology *189*, 6101-6108.

Tully, J.G., Rose, D.L., Whitcomb, R., and Wenzel, R. (1979). Enhanced isolation of Mycoplasma pneumoniae from throat washings with a newly modified culture medium. Journal of Infectious Diseases *139*, 478-482.

Tuomanen, E., Cozens, R., Tosch, W., Zak, O., and Tomasz, A. (1986). The rate of killing of Escherichia coli byβ-lactam antibiotics is strictly proportional to the rate of bacterial growth. Microbiology *132*, 1297-1304.

Uruén, C., Chopo-Escuin, G., Tommassen, J., Mainar-Jaime, R.C., and Arenas, J. (2021). Biofilms as Promoters of Bacterial Antibiotic Resistance and Tolerance. Antibiotics *10*, 3.

Van den Bergh, B., Michiels, J.E., Wenseleers, T., Windels, E.M., Boer, P.V., Kestemont, D., De Meester, L., Verstrepen, K.J., Verstraeten, N., Fauvart, M.*, et al.* (2016). Frequency of antibiotic application drives rapid evolutionary adaptation of Escherichia coli persistence. Nat Microbiol *1*, 16020.

van Gemen, B., Koets, H.J., Plooy, C.A., Bodlaender, J., and Van Knippenberg, P.H. (1987). Characterization of the ksgA gene of Escherichia coli determining kasugamycin sensitivity. Biochimie *69*, 841-848.

Vega, N.M., Allison, K.R., Khalil, A.S., and Collins, J.J. (2012). Signaling-mediated bacterial persister formation. Nature chemical biology *8*, 431-433.

Villellas, C., Aristimuño, L., Vitoria, M.-A., Prat, C., Blanco, S., de Viedma, D.G., Domínguez, J., Samper, S., and Aínsa, J.A. (2013). Analysis of mutations in streptomycin-resistant strains reveals a simple and reliable genetic marker for identification of the Mycobacterium tuberculosis Beijing genotype. Journal of Clinical Microbiology *51*, 2124-2130.

Volkov, I.L., Seefeldt, A.C., and Johansson, M. (2019). Tracking of single tRNAs for translation kinetics measurements in chloramphenicol treated bacteria. Methods *162*, 23-30.

Wang, X., Lord, D.M., Hong, S.H., Peti, W., Benedik, M.J., Page, R., and Wood, T.K. (2013).
Type II toxin/antitoxin MqsR/MqsA controls type V toxin/antitoxin GhoT/GhoS. Environmental
microbiology *15*, 1734-1744.
Wang, X., and Wood, T.K. (2011a). Toxin-antitoxin systems influence biofilm and persister cell
formation and the general stress response. Appl Environ Microbiol *77*, 5577-5583.
Wang, X., and Wood, T.K. (2011b). Toxin-antitoxin systems influence biofilm and persister cell
formation and the general stress response. Applied and environmental microbiology *77*, 5577-
5583.
Wen, W., Liu, B., Xue, L., Zhu, Z., Niu, L., and Sun, B. (2018). Autoregulation and virulence
control by the toxin-antitoxin system SavRS in Staphylococcus aureus. Infection and immunity
86.
Wen, Y., Behiels, E., and Devreese, B. (2014). Toxin–Antitoxin systems: their role in
persistence, biofilm formation, and pathogenicity. Pathogens and disease *70*, 240-249.
Widdel, F. (2007). Theory and measurement of bacterial growth. Di dalam Grundpraktikum
Mikrobiologie *4*, 1-11.
Wilmaerts, D., Herpels, P., Michiels, J., and Verstraeten, N. (2019a). Genetic Determinants of
Persistence in Escherichia coli. In Persister Cells and Infectious Disease (Springer), pp. 133-180.
Wilmaerts, D., Windels, E.M., Verstraeten, N., and Michiels, J. (2019b). General Mechanisms
Leading to Persister Formation and Awakening. Trends Genet *35*, 401-411.
Wood, T.K., and Song, S. (2020). Forming and waking dormant cells: The ppGpp ribosome
dimerization persister model. Biofilm *2*, 100018.
Wu, N., He, L., Cui, P., Wang, W., Yuan, Y., Liu, S., Xu, T., Zhang, S., Wu, J., and Zhang, W.
(2015). Ranking of persister genes in the same Escherichia coli genetic background demonstrates
varying importance of individual persister genes in tolerance to different antibiotics. Frontiers in
microbiology *6*, 1003.
Xie, Y., Wei, Y., Shen, Y., Li, X., Zhou, H., Tai, C., Deng, Z., and Ou, H.Y. (2018). TADB 2.0:
an updated database of bacterial type II toxin-antitoxin loci. Nucleic Acids Res *46*, D749-D753.
Yamasaki, R., Song, S., Benedik, M.J., and Wood, T.K. (2020). Persister cells resuscitate using
membrane sensors that activate chemotaxis, lower cAMP levels, and revive ribosomes. Iscience
23, 100792.
Zhang, Y., and Li, Y. (2010). PhoU (PerF), A PERSISTENCE SWITCH INVOLVED IN
PERSISTER FORMATION AND TOLERANCE TO MULTIPLE ANTIBIOTICS AND
STRESSES AS A DRUG TARGET FOR PERSISTER BACTERIA. (Google Patents).
779
780
781

Supplementary materials

Fig. S1. Growth of the minimal cell. (**A**) Growth curve and (**B**) doubling time of evolved strain Syn3B P026 and parent strain Syn3B. Error bar represents SEM. n = 12 independent biological replicates. *p<0.05, **p<0.01, ***p<0.001, ****p < 0.0001.

Fig. S2. Death curve comparison of Syn3B P026 by re-exposing streptomycin and ciprofloxacin. Overnight cultures of Syn3B were grown to stationary phase (OD~0.3-0.35), diluted to 1:10 and treated with streptomycin (100 µg/mL) or ciprofloxacin (1 µg/mL) for 48 h and sampled after 24 h and 48 h. Antibiotic treated culture then washed twice through centrifugation, grew back to stationary phase and re-exposed to same antibiotic to make the second death curve. Error bars represent SEM (n ≥ 3).

A. Evolution of Syn3B

B. Evolution of Syn3B P026 through cyclic antibiotic treatment

C. Designed antibiotic survival assay for Syn3B

Fig. S3. A. Evolution of Syn3B P026. Cells were evolved in SP16 media by a serial passage from P01 to P09. From P09, exponential cultures were repeatedly diluted 1:100 into fresh SP16 media for higher growth yield up until the 26 passage. Then, a single colony called P026 was selected and send for sequencing. **B.** Evolution of Syn3B resistant mutants through cyclic antibiotic treatment. Stationary phase cultures of P026 were diluted 1:10 in SP16 media containing lethal doses of different types of antibiotic (Streptomycin, Ciprofloxacin, Streptomycin-Ciprofloxacin and Kasugamycin) for 24 h, then washed twice through centrifugation, regrew in the similar condition and re-exposed with same antibiotic. Finally, a single colony was selected for whole genome sequencing. **C.** Antibiotic survival assay was optimized based on traditional agar plate method. Overnight cultures were grown to stationary phase (OD>0.5), diluted to 1:10 in antibiotic-containing media. Percent surviving cells were calculated by the counting colony number before and after antibiotic treatment. Over 200 individual colonies were tested for bacterial resistance, and as expected, no resistant colonies were detected.

815
Fig. S4. Determination of Minimal Inhibitory Concentration (MIC). A. Streptomycin (Strep)
B. Ciprofloxacin (Cip). **C.** Chloramphenicol (Cm). **D.** Kasugamycin (Ksg). Exponential phase
cultures with different dilutions were plated on SP4 agarose plate with different concentrations
of antibiotics. The MIC was determined to be 10 µg/ml for streptomycin, 0.1 µg/mL for
ciprofloxacin, 0.5 µg/mL for chloramphenicol, and 30 µg/mL for kasugamycin. Error bars
represent the standard deviation and Ŧ represents data is out of detectable range.

Table S1. The Syn3B genome is lacking homologs to stress response genes that are linked with tolerance and persistence.

Gene name	Major function	Reference
rpoS, fis, hns, hnr, dksA, rob, rcnR	Global regulator	(Cui et al., 2018; Hansen et al., 2008; Wu et al., 2015)
priA, ruvA, recG, recC, recN, uvrD, ybaZ	DNA mismatch repair and recombination	(Cui et al., 2018; Girgis et al., 2012)
recA, lexA, umuD	SOS response	(Cui et al., 2018; Wu et al., 2015)
SpoT	Stringent response	(Korch et al., 2003)
hslU	Protease	(Girgis et al., 2012)
glp, plsB, tkt, sucB, ubiF, ubiE	Energy production	(Girgis et al., 2012)
lpcA, folB, mltC, apaH, surA, ygfA, yigB	Metabolism	(Cui et al., 2018; Hansen et al., 2008)
rfaE, rfaP, rfaQ	Lipopolysaccharide synthesis	(Cui et al., 2018; Girgis et al., 2012)
asmA, tolR	Assembly of outer membrane proteins; maintains outer membrane integrity	(Cui et al., 2018; Girgis et al., 2012)
livJ	Amino acid transporter	(Girgis et al., 2012)
rffM	Enterobacterial common antigen synthesis	(Girgis et al., 2012)
visC	ubiquinone biosynthetic process	(Girgis et al., 2012)
yacC	Exonuclease domain-containing protein	(Girgis et al., 2012)
cspD	DNA replication inhibitor	(Kim and Wood, 2010)
metG	Methionyl-tRNA synthetase	(Girgis et al., 2012)
pspF	Transcriptional activator for the phage shock protein (psp) operon	(Vega et al., 2012)
yiiS, yfcN, yhaC, yjbE, yceA, yagM, ybcK, ydhL, yibA	Unknown	(Cui et al., 2018; Girgis et al., 2012)
flgE, flgJ, fliG, flhB	Flagellar system	(Cui et al., 2018)
rrmJ	Translation	(Cui et al., 2018)
raiA, rmf, hpf, hflX	Ribosome dimerization	(Song and Wood, 2020; Wood and Song, 2020)
hscB, yfhJ	Iron sulfur Cluster	(Cui et al., 2018)
acrA, acrB, yfgL, yfbK	Transporter	(Cui et al., 2018)
oxyR	Antioxidant defense	(Wu et al., 2015)
mqsR-mqsA;dinJ-yafQ;hipA-hipB;yefM-yoeB;rnlB-rnlA;yafN-yafO;mazE-mazF;hicA-hicB;chpS-chpB;higA-higB;prlF-yhaV;relB-relE;vapB-vapC;hok-sok;ldrD-rdlD;tisB-istR;shoB-ohsC;symE-symR;ghoT-ghoS;dinQ-agrB;CbtA-CbtE;parD-parE;ccdA-ccdB;ralR-ralA;zorO-orzO;yeeU-yeeV; pndA-pndB;cptA-cptB;srnB-srnC	TA modules	(Christensen et al., 2004; Dörr et al., 2010; Garcia Rodriguez et al., 2020; Hansen et al., 2012; Hu et al., 2015; Sun et al., 2017; Tripathi et al., 2012; Wang and Wood, 2011b; Wen et al., 2014; Wilmaerts et al., 2019a)

Table S2. Comparison of % similarity (based on homologous proteins) of Syn3B with most threatening
 human pathogens.

Human pathogens	Related diseases	Homologous proteins No. in Syn3B*	% similarity with Syn3B
Escherichia coli O157	Severe intestinal infection	287	63
Staphylococcus aureus	Pneumonia, meningitis, osteomyelitis, endocarditis, toxic shock syndrome, bacteremia, sepsis, impetigo, boils, cellulitis, folliculitis and carbuncles etc.	338	74
Burkholderia cepacia	pneumonia in immunocompromised individuals	262	58
Pseudomonas aeruginosa	Urinary tract infections, respiratory system infections and dermatitis	273	60
Clostridium difficile	Diarrhea and inflammation of the colon	329	73
Klebsiella pneumoniae	Pneumonia and infection in the lungs	294	65
Acinetobacter baumannii	Infection in the blood, urinary tract, and lungs (pneumonia)	272	60
Mycobacterium tuberculosis (MTB)	Tuberculosis	275	60
Neisseria gonorrhoeae	Gonorrhea	265	58
Streptococcus pyogenes	Pharyngitis, tonsillitis, scarlet fever, cellulitis, erysipelas and rheumatic fever etc.	328	72

845 **Table S3.** Overlapping genes in Syn3B.

Gene name	Gene Length (bp)	Locus tag	Annotation
ieltA	1062	JCVISYN3A_0132	AAA family ATPase
ietS	2,274	JCVISYN3A_0133	hypothetical protein
CDS_33	681	JCVISYN3A_0281	hypothetical protein
proRS	1425	JCVISYN3A_0282	Proline--tRNA ligase
truB	879	JCVSYN2_00715	tRNA pseudouridine(55) synthase TruB
ribF	555	JCVSYN2_00720	FAD synthetase
CDS_50	741	JCVSYN2_00955	hypothetical protein
CDS_51	345	JCVSYN2_00960	hypothetical protein
trmK;\yqfN	678	JCVSYN2_01080	hypothetical protein
CDS_60	777	JCVSYN2_01085	dinuclear metal center protein, YbgI family
pncB	1062	JCVSYN2_01655	Nicotinate phosphoribosyltransferase
CDS_97	609	JCVSYN2_01660	Uncharacterized protein
CDS_124	681	JCVSYN2_02430	hypothetical protein
CDS_125	1539	JCVSYN2_02435	amino acid permease
CDS_99	363	JCVSYN2_01695	lipoprotein
CDS_100	1071	JCVSYN2_01700	hypothetical protein

846
847

848 **Table S4.** Experimental variation in population decay curve of Syn3B P026.

Exp. ID	Streptomycin treatment for 24 h	Streptomycin treatment for 48 h	Ciprofloxacin treatment for 24 h	Ciprofloxacin treatment for 48 h	Replicates (n)
1.	0.03±0.0014	0.0012±0.0003	0.09±0.0096	0.003±0.0005	3
2.	0.083±0.52	0.008±0.067	0.78±0.840	0.017±0.008	3
3.	0.012±0.013	0.006±0.002	3.1±0.93	0.06±0.003	3
4.	0.04±0.009	0.002±0.00012	0.4±0.14	0.007±0.005	3
5.	0.015±0.002	0.008±0.0004	0.5±0.2	0.02±0.014	3
6.	0.004±0.0006	0.002±0002	0.04±0.006	0.004±0006	12

850 **Table S5.** SP16 media composition.

SP16 media components			
Components	**Final**	**Supplier**	**Cat No.**
2XP1			
Mycoplasma Broth Base (PPLO Broth)	7 mg/ml	Thomas Scientific	BD 211458
Tryptone	20 mg/ml	Fisher	AC611841000
Peptone	10.6 mg/ml	Fisher	BP9725-500
Yeast extract solution (autoclaved)	14 mg/ml	Fisher	BP9727-500
TC Yeastolate (Autoclaved)	4 mg/ml	Thomas Scientific	BD 255772
P4			
D(+)-Glucose	0.50%	Fisher	Alfa Aesar A1682836
CMRL 1066 (with L-Glutamine, without Sodium bicarbonate)	0.25X	Thomas Scientific	C992B09 Mfr. No. AT110-1L
NaHCO3 (Sodium Bicarbonate)	1.1 mg/ml	Fisher	S233-3
Penicillin G	625 µg/ml (~1000 U/ml)	Fisher	MP Biomedicals 0210054380 (powder: 500-1700 u/mg)
L-Glutamine	146 µg/ml	Fisher	Alfa Aesar A1420118
FBS			
Fetal Bovine Serum (FBS), Heat inactivated	17%	Fisher	10-438-018 Gibco 10438018
Tetracycline	4 µg/ml	Fisher	BP912-100
Vit B1	5 µg/ml	Acros organic	148990100