CAMAP: Artificial neural networks unveil the role of codon arrangement in modulating MHC-I peptides presentation

Tariq Daouda[1,2,*], Maude Dumont-Lagacé[1,3,*], Albert Feghaly[1], Yahya Benslimane[1,3], Rébecca Panes[1,4], Mathieu Courcelles[1,5], Mohamed Benhammadi[1,3], Lea Harrington[1,3], Pierre Thibault[1,5], François Major[1,6], Yoshua Bengio[6], Étienne Gagnon[1,4], Sébastien Lemieux[1,2,6], Claude Perreault[1,3]

[1]Institute for Research in Immunology and Cancer; Université de Montréal; Montréal, Québec H3C 3J7, Canada

[2]Department of Biochemistry; Université de Montréal; Montréal, Québec H3C 3J7, Canada

[3]Department of Medicine; Université de Montréal; Montréal, Québec H3C 3J7, Canada

[4]Department of Microbiology, Infectiology and Immunology; Université de Montréal; Montréal, Québec H3C 3J7, Canada

[5]Department of Chemistry; Université de Montréal; Montréal, Québec H3C 3J7, Canada

[6]Department of Informatics and Operational Research; Université de Montréal; Montréal, Québec H3C 3J7, Canada

Tariq Daouda is now affiliated to: (1) Broad Institute of MIT and Harvard, Cambridge, United States; (2) Center for Cancer Research, Massachusetts General Hospital, Charlestown, United States; (3) Department of Medicine, Harvard Medical School, Boston, United States; and (4) Center for Immunology and Inflammatory Diseases, Massachusetts General Hospital, Charlestown, United States.

*Corresponding author: Tariq Daouda

Email: tdaouda@broadinstitute.org

*These authors contributed equally.

Abstract

MHC-I associated peptides (MAPs) play a central role in the elimination of virus-infected and neoplastic cells by CD8 T cells. However, accurately predicting the MAP repertoire remains difficult, because only a fraction of the transcriptome generates MAPs. In this study, we investigated whether codon arrangement (usage and placement) regulates MAP biogenesis. We developed an artificial neural network called Codon Arrangement MAP Predictor (CAMAP), predicting MAP presentation solely from mRNA sequences flanking the MAP-coding codons (MCCs), while excluding the MCC *per se*. CAMAP predictions were significantly more accurate when using original codon sequences than shuffled codon sequences which reflect amino acid usage. Furthermore, predictions were independent of mRNA expression and MAP binding affinity to MHC-I molecules and applied to several cell types and species. Combining MAP ligand scores, transcript expression level and CAMAP scores was particularly useful to increaser MAP prediction accuracy. Using an *in vitro* assay, we showed that varying the synonymous codons in the regions flanking the MCCs (without changing the amino acid sequence) resulted in significant modulation of MAP presentation at the cell surface. Taken together, our results demonstrate the role of codon arrangement in the regulation of MAP presentation and support integration of both translational and post-translational events in predictive algorithms to ameliorate modeling of the immunopeptidome.

50   Author summary

MHC-I associated peptides (MAPs) are small fragments of intracellular proteins presented at the

surface of cells and used by the immune system to detect and eliminate cancerous or virus-infected

cells. While it is theoretically possible to predict which portions of the intracellular proteins will

be naturally processed by the cells to ultimately reach the surface, current methodologies have

prohibitively high false discovery rates. Here we introduce an artificial neural network called

Codon Arrangement MAP Predictor (CAMAP) which integrates information from mRNA-to-

protein translation to other factors regulating MAP biogenesis (e.g. MAP ligand score and

transcript expression levels) to improve MAP prediction accuracy. While most MAP predictive

approaches focus on MAP sequences per se, CAMAP's novelty is to analyze the MAP-flanking

mRNA sequences, thereby providing completely independent information for MAP prediction.

We show on several datasets that the integration of CAMAP scores with other known factors

involved in MAP presentation (i.e. MAP ligand score and mRNA expression) significantly

improves MAP prediction accuracy, and further validate CAMAP learned features using an *in-*

vitro assay. These findings may have major implications for the design of vaccines against cancers

and viruses, and in times of pandemics could accelerate the identification of relevant MAPs of

viral origins.

Abbreviations: MHC-I: major histocompatibility complex class-I, MAP: MHC-I associated

peptides, CAMAP: Codon arrangement MAP predictor, DRiP: defective ribosomal product,

ANN: artificial neural network, MCC: MAP-coding codons, B-LCL: B-lymphoblastoid cell line,

KL: Kullback-Leibler, BS: binding score, OVA: ovalbumin protein, WT: wildtype, EP:

enhanced presentation, RP: reduced presentation.

Introduction

In jawed vertebrates, virtually all nucleated cells present at their surface major histocompatibility

complex class-I (MHC-I) associated peptides (MAPs), collectively referred to as the

immunopeptidome [1,2]. MAPs play a central role in shaping the adaptive immune system, as they

orchestrate the development, survival and activation of CD8 T cells [3]. Moreover, recognition of

abnormal MAPs is essential to the elimination of virus-infected and neoplastic cells [4]. Therefore,

systems-level understanding of MAP biogenesis and molecular composition remains a central

issue in immunobiology [5,6].

The generation of the immunopeptidome can be conceptualized in two main events: (a) the

generation of MAP candidates (i.e. peptides of appropriate length for MHC-I presentation) through

protein degradation, and (b) a subsequent filtering step through the binding of MAP candidates to

the available MHC-I molecules. Rules that regulate the second event have been well characterized

using artificial neural networks (ANN) and weighted matrix approaches [7,8]. However,

accurately predicting which peptides will ultimately reach MHC-I molecules following a multistep

processing in the cytosol and endoplasmic reticulum remains an open question [6]. Most efforts at

modeling MAP generation have focused on post-translational events and their regulation by the

amino acid sequence of MAPs and of directly adjacent residues (typically 10-mers at the N- and

C-termini). While the consideration of preferential sites of proteasome cleavage has proven useful

to enrich for MAP candidates [9], it remains insufficient for MAP prediction, due to prohibitive

false discovery rates [10–12].

A large body of evidence suggests that a substantial portion of MAPs are produced co-

translationally [13–15], deriving from defective ribosomal products (DRiPs), that is, polypeptides

that fail to achieve a stable conformation during translation and are consequently rapidly degraded.

97 This concept was initially supported by two observations: (i) viral MAPs can be detected within

98 minutes after viral infection, much earlier than their associated proteins half-life [16], and (ii) MAP

99 presentation correlates more closely with translation rate than with overall protein abundance

100 [17,18]. In addition, while all proteins contain peptides that are predicted to bind MHC-I

101 molecules, mass spectrometry analyses have revealed that the immunopeptidome is not a random

102 excerpt of the transcriptome or the proteome [1,19]. Indeed, proteogenomic analyses of 25,270

103 MAPs isolated from B lymphocytes of 18 individuals showed that 41% of expressed protein-

104 coding genes generated no MAPs [19]. These authors also provided compelling evidence that the

105 presentation of MAPs cannot be explained solely by their affinity to MHC-I alleles and their

106 transcript expression levels, while ruling out low mass spectrometry sensitivity as an explanation

107 for the non-presentation of the strong binders. Because (i) MAPs appear to preferentially derive

108 from DRiPs and (ii) codon usage influences both precision and efficiency of protein synthesis

109 [20,21], we hypothesized that codon usage in the vicinity of MAP-coding codons (MCCs) might

110 significantly contribute to the regulation of MAP biogenesis. We developed an artificial neural

111 network called Codon Arrangement MAP Predictor (CAMAP), trained to identify MCCs flanking

112 regions. We then used CAMAP to uncover key codon features that characterize mRNA sequences

113 encoding for MAPs (i.e. source) when compared to sequences that do not (i.e. non-source).

Results

Dataset description

117 We analyzed a previously published dataset consisting of MAPs presented on B lymphoblastoid

118 cell line (B-LCL) by a total of 33 MHC-I alleles from 18 subjects [19,22]. Because we were

searching for features that influence MAP generation and not the binding of MAP to MHC-I

molecules, we elected to analyze the MCC flanking sequences only and excluded the MCCs *per*

se from our positive (hits) and negative (decoys) sequences (Fig. 1A). To facilitate data analysis

and interpretation, we restricted our hit dataset to MAPs with a length of 9 amino acids, for a total

of 19,656 9-mer MAPs (which represents 78% of MAPs in this dataset). We next created a decoy

dataset from transcripts that generated no MAPs, by randomly selecting 98,290 9-mers from these

transcripts. Finally, we used pyGeno [23] to extract MCCs flanking regions corresponding to both

hit and decoy MAPs, which constituted our final dataset for CAMAP. Of note, each sequence in

the final dataset is unique and derives from the canonical reading frame. In addition, in order to

investigate the relative importance of codon vs. amino acid usage in MAP biogenesis, we

generated a dataset of shuffled sequences (for both positive and negative datasets) in which original

codon sequences were randomly replaced by synonymous codons according to their usage

frequency in the dataset (Fig. 1B). This transformation was performed to ensure that both neural

networks received the same number of parameters as input, preventing the introduction of a

favorable bias for the codon network. The random shuffling causes any codon-specific feature to

be shared among synonyms, thereby causing the shuffled codon distribution to reflect the amino

acid usage (see Materials and Methods for more details). Indeed, codon distributions in the

shuffled datasets more closely reflected those of their corresponding amino acid than in the original

dataset (Supplementary Figure S1), with 92% of codons in the shuffled dataset showing a strong

correlating ($R^2 > 0.95$) with the amino acid distribution, compared to only 69% in the original

dataset ($p < 2x10^{-16}$, Supplementary Figure S2). Importantly, this shuffling does not affect the

resulting amino acid sequence thereby preserving all potential amino acid-related motifs.

Distributions of each codons in the original VS shuffled dataset and compared to its corresponding

amino acid can be found in Supplementary Figure S3.

Figure 1. Construction of the dataset. (a) Transcripts expressed in B cells from 18 subjects were

considered as source or non-source transcripts depending on their match with at least one MAP.

Because we were searching for features that might influence MAP generation and not the binding

of MAP to MHC-I, we focused our attention on mRNA sequences adjacent to the nine MCCs (i.e.

up to 162 nucleotides on each side of MCCs). (b) Creation of the shuffled dataset. Codons were

randomly replaced by a synonymous codon according to their respective frequencies (i.e. codon

usage) in the dataset. The random shuffling causes any codon-specific feature to be shared among

synonyms, thereby causing the shuffled codon distribution to reflect the amino acid usage.

Importantly, both the original sequence and its shuffled version translates into the same amino

acids.

CAMAP links codon usage to MAP presentation

To assess the importance of codon usage in MAP biogenesis, we reasoned that if codons bear

important information that is operative at the translational rather than the post-translational level,

then: (i) CAMAP trained to identify MCCs flanking regions should consistently perform better

when trained on original codon sequences than on shuffled codon sequences (reflecting amino acid

sequences), and (ii) synonymous codons should have different effects on the prediction. To test

these hypotheses, CAMAP received as inputs MCCs flanking regions from hit and decoy

sequences from either the original or shuffled datasets. It was then trained to predict the probability

that individual input sequences were MCCs flanking regions (i.e. hit) rather than sequences from

the negative dataset (Supplementary Figure S4A).

We compared CAMAP performance when predicting MAP presentation from original codon

sequences, versus shuffled sequences representing amino acid arrangement. To evaluate the

robustness of our approach, 12 different CAMAPs were trained in parallel, with different train-

validation-test splits of the dataset. Our results show that predictions were consistently better when

CAMAP received the original codons rather than the shuffled sequences (Fig. 2A). CAMAPs

receiving information from both pre-MCCs and post-MCCs sequences (i.e. whole MCC flanking

context) also performed better than when receiving only pre- or post-MCCs context (Fig. 2A and

Supplementary Figure S4B-C), suggesting that pre- and post-MCCs context are not redundant.

Indeed, we found a weak correlation between the prediction scores of CAMAPs trained only with

pre- or post-MCCs sequences (Supplementary Fig. S5). In addition, CAMAPs receiving longer

sequences performed better than those receiving shorter sequences (Fig. 2B). Because sequences

located far upstream and downstream of the MCCs (i.e. in ranges exceeding the direct influence

of proteases) are informative regarding MAP presentation, it supports the existence of factors

unrelated to protein degradation modulating MAP presentation.

Figure 2. CAMAP predictions on MAP-flanking sequences. (A) Area under the curve (AUC)

score for CAMAPs trained with whole MCCs context, versus CAMAPs trained with only pre- or

post-MCCs context. All CAMAPs presented here were trained with a context size of 162

nucleotides. (B) AUC for CAMAPs trained with codon context sizes of 9, 27, 81 and 162

nucleotides (context here refer to mRNA sequences flanking the MCCs).

Both MAP binding affinity to the MHC-I molecule and the level of gene expression are predictive

of MAP presentation [19]. Because codon usage has been shown to be different in highly expressed

genes, we wanted to verify whether the codon-specific rules captured by CAMAP were associated

with potential biases in our positive dataset, which is enriched in highly expressed genes. We first

show that there is no correlation between gene expression levels and CAMAP scores in both the

positive and negative datasets (R < 0.1, Fig. 3A). This was true for both average expression levels

across our samples (Fig. 3A), and for samples individually (see Supplementary Fig. S6). Secondly,

we trained CAMAP networks using a decoy dataset that mirrored the positive dataset gene

expression level (Supplementary Fig. S7A) and showed similar results: CAMAP trained on

original codon sequence performed better than CAMAP trained on shuffled sequences

(Supplementary Fig. S7B). These results show that the codon-specific rules captured by CAMAP

trained on original sequences are independent of gene expression levels.

Figure 3. Correlation between CAMAP prediction score and (A) transcript expression levels

and (B) MAP binding affinity. CAMAP used here was trained on original codon sequences using

a context size of 162 nucleotides (both pre- and post-MCCs context).

We stipulate that the presence of MHC-I binding motifs in the MCCs in the positive dataset might

be associated with biases in the MAP-flanking regions, which could also influence CAMAP

training. Therefore, to evaluate the presence of this potential bias, we first evaluated the correlation

between CAMAP scores and MAPs binding affinity. Again, our result showed no correlation

between CAMAP scores and MAP binding affinity, both when considering the minimal binding

affinity of each MAP to the MHC-I alleles contained in our dataset (Fig. 3B) or when considering

each allele individually (Supplementary Fig. S8). Secondly, we trained CAMAP networks using a

decoy dataset that mirrored the positive dataset MAP binding affinities (Supplementary Fig. S9A).

Again, CAMAPs trained on original codon sequence performed better than CAMAPs trained on

shuffled sequences (Supplementary Fig. S9B). These results show that codon-specific rules

captured by CAMAP trained on original sequences are independent of MAP binding affinities and

of potential biases in codon usage of MAP-flanking sequences associated with the presence of an

MHC-I binding motif in the MCCs.

We next evaluated the possibility of biases associated with many MAPs originating from

conserved regions (e.g., found in multiple domains of the same domain family such as zinc fingers

or kinases). We first evaluated MAPs that could originate from different transcripts within the

transcriptome (i.e. transcripts with sufficient expression levels detected by RNA sequencing) as

they are likely to represent conserved regions in the genome. While 79.9% of MAP originated

from unique contexts (Supplementary Fig. S10A), 2.1% of MAPs had more than 3 possible origins,

which represented 11.7% of the hit dataset (Supplementary Fig. S10B). These MAPs with several

possible origins preferentially derived from zinc finger proteins, which are known to share

homologous regions (Supplementary Fig. S11). We therefore trained CAMAPs with datasets

excluding entries encoding for MAPs that had >3 or >10 possible origins and compared their

performance with that of CAMAPs trained without excluding these MAPs. Our results show that

whatever the dataset used, CAMAP trained with original sequences always significantly

outperformed CAMAP trained with shuffled sequences (Supplementary Fig. S12). Taken together,

these results suggest that the codon-specific rules captured by CAMAP are independent of

potential homologies in the hit dataset, as they do not appear to influence CAMAP performance.

We next validated our CAMAP trained on 9-mer MAPs derived from B-LCL using 5 datasets

derived from different human and mouse cell types. All the validation datasets were described

through proteogenomic analyses similarly to our B-LCL training datasets. However, all the

validation datasets included MAPs of 8-11 mers, in contrast with the training dataset that contained

only 9-mer MAPs. The validation datasets consisted of (i) our B-LCL dataset, this time including

all peptide lengths [19,22], (ii) a dataset of human peripheral blood mononucleated cells or PBMCs

[24], (iii) a dataset of B-lymphoblastoid cells expressing unique HLA alleles (B721.221 [11]), (iv)

murine colon carcinoma cell line (CT26) and (v) a murine lymphoma cell line (EL4, [24,25]). For

all datasets, we created hit and decoy datasets of original and shuffled sequences using the same

approach described above but including MAPs of 8-11 amino acids. Notably, CAMAPs trained on

human sequences encoding 9-mers MAPs from one human cell type (i.e. B-LCL) could also

predict presentation of 8-11 mers MAPs in other human cell types (Fig. 4), as well as from mouse

cell lines, albeit with lower performances (Fig. 4). Here again, CAMAPs trained on original

sequences consistently outperformed CAMAPs trained on shuffled sequences (Fig. 4). These

results show that the codon-specific rules derived by CAMAPs to predict MAP presentation are

valid across different cell types, and can even be applied to another species, albeit with slightly

lower performances. These results support a role for codons in the modulation of MAP

presentation.

Figure 4. Validation of CAMAP predictions on 5 datasets derived from human and murine

cell lines. CAMAP prediction score for different datasets derived from humans (i.e. B-LCL,

PBMCs and B721.221) or mouse (i.e. CT26 and EL4) cells. Of note, all CAMAPs were trained on

B-LCL-derived sequences encoding for 9-mer MAPs only with a context size of 162 nucleotides.

Results are reported for 8 to 11-mer MAPs derived from the 5 datasets. In all panels, 12 CAMAPs

trained with original or shuffled synonymous sequences were compared (significance assessed

using Student T test).

The lower performances of CAMAP trained with shuffled sequences (representing amino acid

distribution) suggests that amino acids in MAP-flanking sequences are less informative than

codons regarding MAP presentation. We formally quantified this difference in information using

the Kullback-Leibler (KL) divergence (see Materials and Methods for more details). Most codons

(47/61, 77%) showed greater KL divergence in the original dataset than the shuffled dataset,

indicating that codon distributions contained more information with regards to MAP presentation

than amino acid distributions (Supplementary Fig. S13). These results suggest that codons in

MAP-flanking regions play a role that is non-redundant with amino acids in MAP biogenesis.

We wondered whether some regions were more influential on MAP presentation than others. To

address this question, we retrieved the model preferences for each codon at each position. The

preferences correspond to the prediction score of our best model (trained with original codon

sequences for a context size of 162 nucleotides) when a single codon at a single position is

provided as input (all other positions being set at [0,0] coordinates in the embedding space). The

model's preferences are therefore a measure of each individual codon's propensity to increase or

decrease the model's output probability as a function of its position relative to the MCCs. A value

of 0.5 denotes a neutral preference, while negative and positive preferences correspond to values

below and above 0.5, respectively. Preferences were obtained by feeding CAMAP sequences in

which all codon values were masked, except for a single position that received a non-null codon

label.

Interestingly, while codons closest to the MCCs were the most influential on CAMAP scores,

some synonymous codons showed opposite effects, further demonstrating that codon usage does

not recapitulate amino acid usage (Fig. 5A-B and Supplementary Fig. S14). The use of embeddings

to encode codons has the advantage of arranging them into a semantic space, wherein codons with

similar influences are positioned close to each other. Interestingly, most synonymous codons did

not form clusters, with a notable exception being proline codons (Fig. 5C). This finding indicates

that for some codons, their effect on CAMAP prediction score may be closer to that of a non-

synonymous codon than to that of one of its synonyms.

Figure 5. CAMAP interpretation of codon impact on MAP biogenesis. Preferences for a

network trained on a context of 162 nucleotides (54 codons) for (A) serine, proline and tyrosine

codons, and (B) leucine codons. (C) Learned codon embeddings. Some synonymous codons,

such as those encoding for Isoleucine (I), Cysteine (C) or Arginine (R) are located far from one

another, while others tend to cluster together (e.g. Proline [P] and Glutamic acid [E]).

CAMAP increases MAP prediction accuracy

We next compared MAP prediction capacities of CAMAPs scores to that of MAP predicted ligand

score (ranks as predicted by NetMHCpan4.0) and mRNA transcript expression levels. We used

ligand scores as predicted by NetMHCpan4.0, which was shown to possess the best predictive

capacities for naturally processed peptides compared to other predictive algorithms [26]. Because

MAP binding to the MHC molecule is essential for its presentation at the cell surface, we elected

to only compare hits and decoys encoding potential binders, i.e. with a minimal ligand score of

1% for at least one allele in the B-LCL dataset. Using a linear regression model, we compared the

predictive capacity of each single parameter using Matthews correlation coefficient, which

measures the quality of binary classifications [27]. Of note, only the predictions on the test set

were used to evaluate the Matthew correlation coefficient in our different models.

Because only potential binders were analyzed here, the mRNA expression level had the highest

predictive capacity, then followed by ligand scores (second) and CAMAP scores (third, Fig. 6A).

As expected due to the multiplicative relationship between MAP ligand score and expression levels

in predicting naturally processed MAPs [11], combining both variables greatly increased

prediction performances (Fig. 6B). Importantly, adding CAMAP scores to the regression model

further increased predictive performances (Fig. 6B). We next computed how many predicted

peptides would need to be tested to capture 1, 5 or 10% of hits in the B-LCL dataset. Results

presented in Table 1 show that using only NetMHCpan4.0 ligand scores (ranks) leads to a very

high false positive rate (FPR) at 72.1% when targeting the top 1%. Adding the expression levels

greatly increased prediction accuracy and decreased the FPR to 32.8% for the top 1% hits. When

adding CAMAP scores as a third variable, the number of peptides needed to capture 1% of hits

greatly decreased, resulting in a very low FPR at 1.1%. Similar trends were observed when

targeting 5 or 10% of hits, although with higher FPR (see Table 1). Similarly, adding CAMAP

scores to expression levels and ligand scores also ameliorated prediction accuracies for the two

other human datasets introduced above (B721.221 and PBMCs, see Supplementary Table S2).

These results show that combining CAMAP scores with the MAP's ligand score (ranks) and its

corresponding transcript expression level significantly improves prediction of MAP and facilitate

identification of relevant epitopes through more accurate predictions.

Figure 6. CAMAP prediction score contributes to the prediction of MAPs. (A) Matthews

correlation coefficient for MAP prediction using a single variable. (B) Matthews correlation

coefficient for MAP prediction using multivariable regression models. The B-LCL dataset (all

MAP lengths) was filtered for MAP with a minimal ligand score (rank) of 1% (NetMHCpan4.0).

Table 1. Number of peptides needed to capture 1%, 5% or 10% of epitopes detected by mass

spectrometry. The lower the number of peptides needed to capture the respective number of

epitopes, the better the performance of the prediction model. This is also illustrated by the

percentage of false identification (false positive rate, FPR) reported here. Peptides were rank-

ordered according to regression scores, for a total of 490,297 unique peptides and 8,991 hits. Of

note, only the maximal regression score was kept for peptides with multiple potential origins.

Regression model	1% hits (n=90)	5% hits (n=450)	10% hits (n=899)

	n	FPR	n	FPR	n	FPR
NetMHCpan4.0	322 ± 18	72.05%	2183 ± 69	79.39%	4927 ± 136	81.75%
NetMHCpan4.0 + expression	134 ± 6	32.84%	601 ± 10	25.12%	1211 ± 16	25.76%
NetMHCpan4.0 + expression **+ CAMAP**	**91 ± 2**	**1.11%**	**524 ± 13**	**14.12%**	**1170 ± 18**	**23.16%**

Codon usage can modulate MAP presentation

To evaluate whether changing the codon arrangement in a MAP-coding sequence might directly

lead to modulation of MAP presentation, we generated three variants of the chicken ovalbumin

(OVA) protein containing the model MAP SIINFEKL [28]. One construct encoded the wild type

OVA (OVA-WT). For the other two constructs, we used CAMAP (trained on original human B-

LCL sequences; Fig. 2) to generate two OVA variants *in silico*, both encoding for the same OVA

protein but using different synonymous codons: one predicted to enhance SIINFEKL presentation

(OVA-EP), the other predicted to reduce it (OVA-RP). Accordingly, the respective CAMAP

scores for OVA-RP, OVA-WT and OVA-EP were: 0.03, 0.65, and 0.96 (Fig. 7A). All variants

encoded the same amino acid sequence but used different synonymous codons. Notably, the sole

difference between the three constructs were the 162 nucleotides flanking each side of the

SIINFEKL-coding codons (i.e. the RNA sequences coding for OVA$_{202-256}$ and OVA$_{265-319}$,

Supplementary Table S1 and Supplementary Figure S15).

Figure 7. Codon usage in MAP-flanking mRNA sequences can influence antigen

presentation and translation efficiency. (A) Design of the inducible Translation Reporter (iTR-

OVA) constructs and CAMAP scores for OVA-WT, OVA-EP and OVA-RP sequences. (B)

Schematic representation of possible translation events. When mRNA codon usage leads to

efficient (uninterrupted) translation, similar amounts of eGFP and Ametrine proteins would be

synthesized. When codon usage in the MAP-flanking regions enhances the frequency of

translation interruption, a lower Ametrine/eGFP ratio would be observed. (C) Kinetics of

SIINFEKL MAP presentation following induction of iTR-OVA constructs expression by

doxycycline, measured in a T-cell activation assay. To remove the influence of differential

expression levels on antigenic presentation and of varying proportion of transduced cells

between samples, T-cell activation levels were normalized to the average Ametrine fluorescence

intensity and to the proportion of eGFP+ cells (i.e. cells expressing the construct). (D)

Translation efficiency as measured by Ametrine/eGFP ratio following iTR-OVA construct

induction. For C and D, results are normalized over the WT sample from the same experiment

(n=4). Statistical differences at each time point were determined using bilateral paired Student T

tests. Significance for the comparison against WT are indicated with *, while comparison of EP

vs RP is indicated with †. N.B.: Each replicate is shown with a dot, while the line and shaded

area represent the average and 95% confidence interval, respectively.

Because codon usage affects translation efficiency, theoretically leading to DRiP formation

through premature translation arrest [20,21], we expected the variable regions of our construct to

affect both translation rates and SIINFEKL presentation in our variants. Therefore, each construct

also coded for two other proteins, eGFP and Ametrine, placed upstream and downstream of the

OVA coding sequence, respectively (Fig. 7A). While the Ametrine fluorescence intensity reflected

the translation rate of the whole construct, the ratio of Ametrine/eGFP fluorescence intensity was

informative regarding the translation efficiency of the whole construct. Indeed, efficient translation

of the full-length construct should produce equivalent quantities of Ametrine and eGFP proteins,

while inefficient/interrupted translation of the construct (i.e. leading to DRiP formation) should

decrease the Ametrine/eGFP ratio (Fig. 7B). The three protein coding sequences were separated

with P2A self-cleaving peptides [29], therefore allowing the co-synthesis of three separate

proteins, controlled by the doxycycline-inducible Tet-On promoter. Importantly, the three proteins

were tightly co-expressed because of the presence of only one start codon at the 5' end of the GFP

protein, as shown by the very high correlation between eGFP and Ametrine fluorescence for each

construct (R>0.97, see Supplementary Figure S16). As we assumed that CAMAP scores reflected

the probability of DRiP generation leading to increased MAP presentation, we expected the OVA-

RP construct to show both reduced SIINFEKL presentation and enhanced translation efficiency

compared to the OVA-EP and OVA-WT constructs. However, as both the OVA-EP and OVA-

WT have CAMAP scores above the neutral threshold of 0.5 and closer to one another (0.98 and

0.65, respectively) compared to the OVA-RP construct (0.03), we expected OVA-EP and OVA-

WT to behave more similarly.

We then used a SIINFEKL-H2-K^b specific T-cell activation assay [30] to measure SIINFEKL

presentation at the cell surface following doxycycline induction. Results for the T-cell activation

assay were normalized by both the Ametrine mean fluorescence intensity and the percentage of

transduced (eGFP+) cells in each specific sample, so that any difference in T-cell activation

observed between our constructs could only be ascribed to synonymous codon variants in the

SIINFEKL-flanking OVA codons. Two main findings emerged from our analyses. First, in

accordance with CAMAP predictions, variation in codon usage led to a 2.3-fold difference in

SIINFEKL presentation between the OVA-EP and OVA-RP variants, with OVA-WT in between

(Fig. 7C). Second, translation efficiency (Ametrine/eGFP ratio) was higher with OVA-RP than

with OVA-EP or OVA-WT, while OVA-EP showed similar translation efficiency compared to

ONA-WT (Fig. 7D). Hence, synonymous codon variations led to slightly divergent outcomes in

OVA-EP and OVA-RP: they modulated the levels of SIINFEKL presentation in both constructs,

but enhanced translation efficiency could only be detected for OVA-RP. These data show that

codon arrangement can modulate MAP presentation strength without any changes in the amino

acid sequence and support a role for translation efficiency and DRiP formation in the modulation

of MAP presentation.

Discussion

Our analyses of large datasets using artificial neural networks and other bioinformatics approaches

provide compelling evidence that codon usage regulates MAP biogenesis via both short- and long-

range effects. While most MAP predictive approaches focus on MAP sequences *per se*, CAMAP's

novelty is that it only receives the MAP-flanking mRNA sequences as input, and no information

on the MAP itself, thereby providing completely independent information for MAP prediction.

The better prediction accuracy of CAMAPs trained with original codons rather than with shuffled

synonyms supports the role of codon usage in modulating MAP biogenesis (Fig. 2). In addition,

we demonstrated that the codon-specific signal that is captured by CAMAP was independent of

transcript expression levels and MAP ligand scores, thereby providing complementary and non-

overlapping information regarding MAP presentation. Additionally, while CAMAP preferences

were more influential for codons located close to the MCCs (Fig. 5), the better performance of

CAMAP trained with longer context size pointed toward a long-range impact of codon usage on

MAP presentation.

The functional link between codon arrangement and MAP biogenesis was illustrated by our *in*

vitro analyses of SIINFEKL biogenesis, in which we were able to modulate SIINFEKL

presentation solely by substituting synonymous codons in mRNA regions flanking SIINFEKL

codons, without changing the protein sequence. While the experimental data derives from a single

model thus limiting the interpretability of our results, this points nonetheless to an interesting

mechanism that could be exploited to enhance antigenic presentation in peptide-bas4ed

immunotherapy (i.e. dendritic cells modified to express a specific MAP).

Further analyses will be needed to assess the full extent of codon arrangement's impact on both

classic MAPs (i.e. derived from canonical reading frames of coding sequences) and cryptic MAPs

(i.e. derived from non-canonical reading frames and non-coding sequences) [31,32], as well as the

potential contribution of codons in non-coding regions (e.g. 5'- or 3'-UTRs) on the regulation of

MAP presentation. However, our results show that the integration of CAMAP scores to the two

best predictive factors for naturally processed MAPs led to a significant increase in prediction

accuracy. Indeed, our regression model combining only transcript expression levels to MAP ligand

scores (ranks as predicted by NetMHCpan4.0), showed that a total of 134 peptides would need to

be tested in order to capture 1% of all presented MAPs (hits), leading to a false positive rate of

32.8%. In contrast, the addition of CAMAP to this model decreased the false positive rate to only

1.1%, leading to 90 correct identifications out of 91 MAPs tested. Although predictions were not

as accurate for the two other human datasets, adding CAMAP scores always resulted in improved

prediction accuracy. Our results therefore support the combined use of ligand scores, transcript

expression levels and CAMAP scores in MAP predictive algorithms. These results have important

practical implications for cancer immunotherapy and peptide-based vaccines, where discovery of

suitable target antigens remains a formidable challenge to this day [33,34].

Materials and methods

Dataset generation

We analyzed a previously published dataset consisting of MAPs presented on B lymphocytes by

a total of 33 MHC-I alleles from 18 subjects [19,22]. Since this dataset was assembled using older

versions of MHC-I binding prediction algorithms (i.e. using a combination of NetMHC3.4 for

common alleles and NetMHCcons1.1 for rare alleles), we verified that the majority of MAPs in

this dataset would also be predicted as binders using more recent algorithms (i.e. a rank $\leq$ 2.0%

using NetMHC4.0 or NetMHCpan4.0). We found an overlap of >92% between these methods (see

Supplementary Fig. S17), thereby validating this dataset for further analysis. In addition, we

reasoned that a transcript should be considered as a genuine positive or negative regarding MAP

biogenesis only if it was expressed in the cells. We therefore excluded from the dataset all

transcripts with very low expression (<1st percentile in terms of FPKM).

To facilitate data analysis and interpretation, we only included transcripts coding for MAPs with

a length of 9 amino acids, for a total of 19,656 9-mer MAPs (which represents 78% of MAPs in

this dataset). We then used pyGeno [23] to extract the mRNA sequences of transcripts coding for

these 9-mer MAPs, which constituted our source-transcripts (Fig. 1A). We next created a negative

(non-source) dataset from transcripts that generated no MAPs. Importantly, transcripts that

encoded for MAPs of any length (i.e. 8 to 11-mer) were excluded from the negative dataset. We

then randomly selected 98,290 non-MAP 9-mers from this negative dataset, and extracted their

coding sequences using pyGeno. Of note, both positive and negative datasets were derived from

the canonical reading frame of non-redundant transcripts.

We analyzed only the MAP context and excluded the MCCs *per se* from our positive (hits) and

negative (decoys) sequences (Fig. 1A). We limited our analyses of flanking sequences to 162

nucleotides (54 codons) on each side of MCCs, because longer lengths would entail the exclusion

of >25% of transcripts (Supplementary Fig. S18).

Creation of the shuffled synonymous codon dataset

To create the shuffled synonymous codon dataset, each sequence was re-encoded by replacing each codon with itself or with a random synonym according to the human transcriptome usage frequencies. These frequencies were calculated using the annotations provided by *Ensembl* for the human reference genome GRCh37.75. Thus, all codon-specific features differing between the positive and negative datasets was removed from the shuffled datasets. Because codons were replaced by their synonymous codons, the shuffled sequences directly reflected amino acid usage in the positive and negative datasets.

CAMAP architecture, sequence encoding and training

The first (input) layer received either MCCs flanking regions from the hit dataset or sequences of the same length contained in the decoy dataset (Fig. 1A). The second layer (Supplementary Fig. 4A) was a codon embedding layer similar to that introduced for a neural language model [35]. Embedding is a technique used in natural language processing to encode discrete words, and has been shown to greatly improve performances [36]. With this technique, the user defines a fixed number of dimensions in which words should be encoded. When the training starts, each word receives a random vector-valued position (its embedding coordinates) in that space. The network then iteratively adjusts the words' embedding vectors during the training phase and arranges them in a way that optimizes the classification task. Notably, embeddings have been shown to represent semantic spaces in which words of similar meanings are arranged close to each other [36]. In the present work, we treated codons as words: each codon received a set of random 2D coordinates that were subsequently optimized during training. The third (output) layer delivered the probability that the input sequence was a MCCs flanking region (rather than a sequence from the negative dataset).

CAMAPs were trained on sequences resulting from the concatenation of pre- and post-MCCs

regions. Before presenting sequences to our CAMAPs, we associated each codon to a unique

number ranging from 1 to 64 (we reserved 0 to indicate a null value) and used this encoding to

transform every sequence into a vector of integers representing codons. Neural networks were built

using the Python package Mariana [37] [https://www.github.com/tariqdaouda/Mariana]. The

Embedding layer of Mariana was used to associate each label superior to 0 to a set of 2D trainable

parameters; the 0 label represents a *null* (masking) embedding fixed at coordinates (0,0). As an

output layer, we used a *Softmax* layer with two outputs (positive / negative). Because negative

sequences are more numerous than positive ones, we used an oversampling strategy during

training. At each epoch, CAMAPs were randomly presented with the same number of positive and

negative sequences. All CAMAPs in this work share the same architecture (Supplementary Fig.

4A), number of parameters and hyper-parameter values: learning rate: 0.001; mini-batch size: 64;

embedding dimensions: 2; linear output without offset on the embedding layer; *Softmax* non-

linearity without offset on the output layer.

For each condition (e.g. context size), the positive and negative datasets were randomly divided

into three non-redundant subsets: (i) the training subsets containing 60% of the positive and

negative transcripts, (ii) the validation and (iii) the test subsets each containing 20% of the positive

and negative transcripts. Transcripts were assigned through a sequence redundancy removal

algorithm, thereby ensuring that no transcript was assigned to multiple subsets. We used an early

stopping strategy on validation sets to prevent over-fitting and reported average performances

computed on test sets. We trained 12 CAMAPs for each combination of conditions, each one using

a different random split of train/validation/test sets. To mask sequences either before or after the

MCCs, we masked either half with *null* value.

Kullback-Leibler divergence

The Kullback-Leibler (KL) divergence computes how well a given distribution is approximated

by another distribution. Its value can be either positive or 0, a null value indicating that the two

distributions are identical (see Materials and Methods for more details). Accordingly, a higher KL

divergence for codon distributions vs. amino acid distributions would indicate that codon

variations are not entirely accounted for by amino acid variations. KL divergence is not a metric,

as it is neither symmetric nor does it satisfy the triangle inequality. It is nevertheless an accurate

and most common way of comparing two probability distributions.

We defined the probability of having codon c at position i as a function of the number of

occurrences of c at position i, divided by the total number of occurrences of that same codon:

518
$$Q_{(c,y,s)}(i) = \frac{N_{c,y,s}(i)}{\sum_j N_{c,y,s}(j)}$$

519 Here Q is a probability, N is a number of occurrences, c is a codon, y is a class (positive or

520 negative), s indicates if codons have been randomized (true or false), i is a position in sequence.

521 For the remainder of the text we will use the following abbreviations:

522
$$P_c(i) = Q_{c,y=positive,s=false}(i)$$

523
$$D_c(i) = Q_{c,y=negative,s=false}(i)$$

524
$$PS_c(i) = Q_{c,y=positive,s=true}(i)$$

525
$$DS_c(i) = Q_{c,y=negative,s=true}(i)$$

526 We then used the KL divergence to compute how well P_c distributions approximate D_c

527 distributions and PS_c distributions approximate DS_c distributions.

528   The KL divergence was defined as:

529
$$D_{KL}(P||Q) = \sum_i P(i) \log \left(\frac{P(i)}{Q(i)} \right)$$

We performed this calculation for both the original and the shuffled dataset, which we then

compared together. If codons and amino acid distributions were equivalent, KL divergence

between hits and decoys would be the same for both original and shuffled sequences, and codons

would cluster along the diagonal.

Predicting MAP presentation with linear regressions

The prediction capacity of CAMAP, NetMHCpan-4.0 ligand score and transcription expression

(TPM) was tested in different combinations of those parameters (Ligand Score + Expression,

Ligand score + Expression + CAMAP score) using the *LogisticRegressionCV* function from the

python package *sklearn* (*sklearn.linear_mode*l, v0.22.1). In each case, the dataset containing hits

and decoy sequences was split into train and test datasets with a ratio of 0.7 to 0.3, respectively.

Values for CAMAP score, Ligand Score and TPM were each scaled to a range of 0-1 in the train

set using MinMaxScaler from *sklearn.preprocessing* and the same scaling model was applied to

the test set afterwards. Regression analysis was performed using *LogisticRegressionCV* with a 10x

cross-validation using the *lbfgs* solver with 1000 iterations. MCC scores were calculated using

matthews_corrcoef from *sklearn.metrics*. When a peptide had multiple sources (multiple

transcripts or genes), only the maximum value from its regression scores was kept.

In vitro assay – inducible translation reporter (iTR)-OVA construct design

An inducible translation reporter was generated by flanking the truncated chicken ovalbumin

(OVA) cDNA (amino acids 144-386) with EGFP-P2A (in 5') and P2A-Ametrine (in 3') cDNA

sequences. MCCs flanking contexts for the EP and RP construct were synthesized as gBlocks

(purchased from Integrated DNA Technologies). The fragments were amplified by PCR and joined

by Gibson assembly under a doxycycline-inducible Tet-ON promoter in a pCW backbone.

Synthetic variants of the OVA coding sequence were generated in silico by varying synonymous

codon usage in the MAP context regions (i.e. 162 nucleotides pre- and post-MCCs). Importantly,

the amino acid sequence was preserved between the different variants; only nucleotide sequences

in the MAP context (162 nucleotides on either side) differed. The sequences with the highest (EP)

and the lowest (RP) prediction scores were selected for further in vitro validation and swapped

into the iTR-OVA plasmid by Gibson assembly [38]. OVA-EP and OVA-RP sequences can be

found in Supplementary Table 1.

Important features of our inducible translation reporter construct and T cell activation assay were:

(i) No changes in amino acid sequence between the three variants: only co-translational events can

differ between the three variants, post-translational events being equivalent for the three

constructs; (ii) Only one start codon, at the beginning of the eGFP coding sequence: this is

important for the translation reporter aspect of our construct (i.e. Ametrine/eGFP ratio), to ensure

that translation can only start at the 5'-end of the whole construct, and not at the beginning of the

OVA or Ametrine coding sequences; (iii) Separation of the three proteins using P2A peptide:

allows the inducible synthesis of three separate proteins in a highly correlated manner; also, the

degradation of one protein will be independent from the others. As we hypothesized that codon

usage might lead to DRiP formation, we did not want the degradation of OVA-derived polypeptide

to induce degradation of attached eGFP or Ametrine, which would affect our translation reporter

assay (Ametrine/eGFP ratio); (iv) Because transcript expression level impacts MAP presentation,

we normalized T-cell activation results by both the number of transduced cells present in the

samples (% of eGFP+ cells) and the Ametrine mean fluorescence intensity of eGFP+ cells

(representing whole construct expression level). Because of these four features, any difference

between the three constructs could be ascribed solely to synonymous codon variants in the

SIINFEKL-flanking OVA codons.

Stable cell line generation

Wildtype and transduced Raw-K^b cells [39] were cultured in DMEM supplemented with 10% Fetal

Bovine Serum (FBS), penicillin (100 units/ml), and streptomycin (100mg/ml). B3Z cells [40] were

maintained in RPMI medium supplemented with 5% FBS, penicillin (100 units/ml), and

streptomycin (100mg/ml).

Lentiviral particles were produced from HEK293T cells by co-transfection of iTR-OVA WT, EP

or RP along with pMD2-VSVG, pMDLg/pRRE and pRSV-REV plasmids. Viral supernatants

were used for Raw-K^b transduction. Raw-K^b OVA-WT, Raw-K^b OVA-EP were sorted on

Ametrine and GFP double positive population after 24h of doxycycline treatment (1 mg/ml).

T-cell activation assay

Raw-K^b OVA-EP, OVA-RP and OVA-WT cells were plated at a density of 250,000 cells/well in

24 well-plates 24h prior to doxycycline treatment (1 mg/ml). After the corresponding treatment

duration, cells were harvested and fixed using PFA 1% for 10 minutes at room temperature and

washed using DMEM 10% FBS. Raw-K^b were then co-cultured (37°C, 5% CO_2) in triplicates with

the CD8 T cell hybridoma cell line B3Z cells at a 3:2 ratio for 16h (7.5 x 10^5 B3Z and 5 x 10^5

Raw-K^b) in 96 well-plates. Cells were lysed for 20 minutes at room temperature using 50 µl/well

of lysis solution (25mM Tris-Base, 0.2 mM CDTA, 10% glycerol, 0.5% Triton X-100, 0.3mM

DTT; pH 7.8). 170 µl/well CPRG buffer was added (0.15mM chlorophenol red-β-d-

galactopyranoside (Roche), 50mM $Na_2HPO_4•7H_2O$, 35mM $NaH_2PO_4•H_2O$, 9mM KCl, 0.9mM

$MgSO_4•7H_2O$). β-galactosidase activity was measured at 575 nm using SpectraMax® 190

Microplate Reader (Molecular Devices). In parallel, cells were analyzed by flow cytometry using

a BD FACS CantoII for eGFP and Ametrine fluorescence.

Data Availability

The datasets analyzed for this study can be found:

• Human B-LCL: RNA-Seq data can be accessed on the NCBI Bioproject database

(http://www.ncbi.nlm.nih.gov/bioproject/; accession PRJNA286122).

• Human PBMC: RNA-sequencing data for human PBMC were extracted from healthy

donors in Zucca et al (2019) [41] and can be accessed under the GEO accession number

GSE106443 and GSE115259, while MAPs were extracted from Murphy et al (2017) [24].

• Human B721.221: The B721.221 dataset was retrieved from Abelin et al (2017) [11]; RNA

sequencing data can be accessed under the GEO accession number GSE93315.

• Murine CT26: RNA-Seq data can be accessed under the GEO accession number

GSE111092. Mass spectrometry data can be found on the ProteomeXchange Consortium

via the PRIDE partner repository (human B-LCL: PXD004023 and murine CT26:

PXD009065 and 10.6019/PXD009065).

• Murine EL4: MAP dataset was extracted from Murphy et al (2017) [24] and EL4 RNA

sequencing dataset was extracted from Sidoli et al (2019) [42] and can be accessed under

the GEO accession number GSE125384.

All figures were generated using R's package "ggplot2". Source code for pyGeno

(https://github.com/tariqdaouda/pyGeno, doi: 10.12688/f1000research.8251.2) and Mariana

(https://github.com/tariqdaouda/Mariana, doi: [to be provided after acceptance]) are freely

available online.

Acknowledgements

This work was supported by grants from the Canadian Cancer Society (number 705604 and 705714), the Oncopole and the Leukemia & Lymphoma Society of Canada. Perreault' lab is supported in part by The Katelyn Bedard Bone Marrow Association. MDL was supported by a studentship from the Canadian Institute of Health Research. Y.Benslimane was supported by a fellowship from the Cole Foundation and a Canadian Institutes of Health Research operating grant (to LH: #13784). EG lab is supported by the Canadian Institute for Health Research operating grant (MOP-133726). The B3Z CD8+ T cell hybridoma cell line was a kind gift from Nilabh Shastri. RAW-Kb cells were kindly provided by Michel Desjardins.

Author contributions

TD designed all computational experiments. TD and AF performed computational experiments. TD wrote pyGeno and Mariana, contributed to design of the iTR-OVA construct, co-wrote the first draft of the paper. MDL contributed to data analysis, to design and synthesis of the iTR-OVA construct, performed flow cytometry analysis, with input of EG, co-wrote the first draft of the paper. AF contributed to data analysis, study design and performed computational experiments (validation on 5 datasets and regressions). Y.Benslimane contributed to design and synthesis of the iTR-OVA construct, with input from LH and EG. RP produced viruses for transduction of the iTR-OVA construct, transduced RAW cells, optimized and performed T-cell activation assay using mild fixation, with input from EG, and reviewed the manuscript. MC performed peptide affinity predictions. MB contributed to the optimization of culture conditions for the iTR-OVA

640 assay. PT reviewed the manuscript. Y.Bengio reviewed and contributed to the manuscript. SL and

641 CP contributed to study design, reviewed and contributed to the manuscript. All co-authors

642 reviewed the manuscript.

643 The authors declare no competing interests.

References

1. Granados DP, Laumont CM, Thibault P, Perreault C. The nature of self for T cells—a
systems-level perspective. Curr Opin Immunol. 2015;34: 1–8. doi:10.1016/j.coi.2014.10.012

2. Caron E, Espona L, Kowalewski DJ, Schuster H, Ternette N, Alpízar A, et al. An open-source
computational and data resource to analyze digital maps of immunopeptidomes. Chakraborty
AK, editor. eLife. 2015;4: e07661. doi:10.7554/eLife.07661

3. Davis MM, Krogsgaard M, Huse M, Huppa J, Lillemeier BF, Li Q. T Cells as a Self-
Referential, Sensory Organ. Annu Rev Immunol. 2007;25: 681–695.
doi:10.1146/annurev.immunol.24.021605.090600

4. Schumacher TN, Schreiber RD. Neoantigens in cancer immunotherapy. Science. 2015;348:
69–74. doi:10.1126/science.aaa4971

5. Caron E, Vincent K, Fortier M-H, Laverdure J-P, Bramoullé A, Hardy M-P, et al. The MHC
I immunopeptidome conveys to the cell surface an integrative view of cellular regulation.
Mol Syst Biol. 2011;7: 533. doi:10.1038/msb.2011.68

6. Neefjes J, Jongsma MLM, Paul P, Bakke O. Towards a systems understanding of MHC class
I and MHC class II antigen presentation. Nat Rev Immunol. 2011;11: 823–836.
doi:10.1038/nri3084

7. Bassani-Sternberg M, Gfeller D. Unsupervised HLA Peptidome Deconvolution Improves
Ligand Prediction Accuracy and Predicts Cooperative Effects in Peptide–HLA Interactions.
J Immunol. 2016;197: 2492–2499. doi:10.4049/jimmunol.1600808

8. Nielsen M, Andreatta M. NetMHCpan-3.0; improved prediction of binding to MHC class I
molecules integrating information from multiple receptor and peptide length datasets.
Genome Med. 2016;8. doi:10.1186/s13073-016-0288-x

9. Tenzer S, Peters B, Bulik S, Schoor O, Lemmel C, Schatz MM, et al. Modeling the MHC
class I pathway by combining predictions of proteasomal cleavage, TAP transport and MHC
class I binding. Cell Mol Life Sci CMLS. 2005;62: 1025–1037. doi:10.1007/s00018-005-
4528-2

10. Nielsen M, Lundegaard C, Lund O, Keşmir C. The role of the proteasome in generating
cytotoxic T-cell epitopes: insights obtained from improved predictions of proteasomal
cleavage. Immunogenetics. 2005;57: 33–41. doi:10.1007/s00251-005-0781-7

11. Abelin JG, Keskin DB, Sarkizova S, Hartigan CR, Zhang W, Sidney J, et al. Mass
Spectrometry Profiling of HLA-Associated Peptidomes in Mono-allelic Cells Enables More
Accurate   Epitope   Prediction.   Immunity.   2017;46:   315–326.
doi:10.1016/j.immuni.2017.02.007

12. Capietto A-H, Jhunjhunwala S, Delamarre L. Characterizing neoantigens for personalized
cancer immunotherapy. Curr Opin Immunol. 2017;46: 58–65. doi:10.1016/j.coi.2017.04.007

13. Antón LC, Yewdell JW. Translating DRiPs: MHC class I immunosurveillance of pathogens
and tumors. J Leukoc Biol. 2014;95: 551–562. doi:10.1189/jlb.1113599

14. Wei J, Kishton RJ, Angel M, Conn CS, Dalla-Venezia N, Marcel V, et al. Ribosomal Proteins
Regulate MHC Class I Peptide Generation for Immunosurveillance. Mol Cell. 2019;73:
1162-1173.e5. doi:10.1016/j.molcel.2018.12.020

15. Yewdell JW, Antón LC, Bennink JR. Defective ribosomal products (DRiPs): a major source
of antigenic peptides for MHC class I molecules? J Immunol. 1996;157: 1823–1826.

16. Croft NP, Smith SA, Wong YC, Tan CT, Dudek NL, Flesch IEA, et al. Kinetics of antigen
expression and epitope presentation during virus infection. PLoS Pathog. 2013;9: e1003129.
doi:10.1371/journal.ppat.1003129

17. Milner E, Barnea E, Beer I, Admon A. The turnover kinetics of major histocompatibility
complex peptides of human cancer cells. Mol Cell Proteomics MCP. 2006;5: 357–365.
doi:10.1074/mcp.M500241-MCP200

18. Hassan C, Kester MGD, de Ru AH, Hombrink P, Drijfhout JW, Nijveen H, et al. The human
leukocyte antigen-presented ligandome of B lymphocytes. Mol Cell Proteomics MCP.
2013;12: 1829–1843. doi:10.1074/mcp.M112.024810

19. Pearson H, Daouda T, Granados DP, Durette C, Bonneil E, Courcelles M, et al. MHC class
I–associated peptides derive from selective regions of the human genome. J Clin Invest.
2016;126: 4690–4701. doi:10.1172/JCI88590

20. Cannarozzi G, Schraudolph NN, Faty M, von Rohr P, Friberg MT, Roth AC, et al. A Role
for   Codon   Order   in   Translation   Dynamics.   Cell.   2010;141:   355–367.
doi:10.1016/j.cell.2010.02.036

21. Plotkin JB, Kudla G. Synonymous but not the same: the causes and consequences of codon
bias. Nat Rev Genet. 2011;12: 32–42. doi:10.1038/nrg2899

22. Granados DP, Rodenbrock A, Laverdure J-P, Côté C, Caron-Lizotte O, Carli C, et al.
Proteogenomic-based discovery of minor histocompatibility antigens with suitable features

for immunotherapy of hematologic cancers. Leukemia. 2016;30: 1344–1354.
doi:10.1038/leu.2016.22

23. Daouda T, Perreault C, Lemieux S. pyGeno: A Python package for precision medicine and
proteogenomics. F1000Research. 2016;5: 381. doi:10.12688/f1000research.8251.2

24. Murphy JP, Konda P, Kowalewski DJ, Schuster H, Clements D, Kim Y, et al. MHC-I Ligand
Discovery Using Targeted Database Searches of Mass Spectrometry Data: Implications for
T-Cell Immunotherapies. J Proteome Res. 2017;16: 1806–1816.
doi:10.1021/acs.jproteome.6b00971

25. Laumont CM, Vincent K, Hesnard L, Audemard É, Bonneil É, Laverdure J-P, et al.
Noncoding regions are the main source of targetable tumor-specific antigens. Sci Transl Med.
2018;10: eaau5516. doi:10.1126/scitranslmed.aau5516

26. Paul S, Croft NP, Purcell AW, Tscharke DC, Sette A, Nielsen M, et al. Benchmarking
predictions of MHC class I restricted T cell epitopes in a comprehensively studied model
system. PLoS Comput Biol. 2020;16: e1007757. doi:10.1371/journal.pcbi.1007757

27. Powers D. Evaluation: From Precision, Recall and F-Factor to ROC, Informedness,
Markedness & Correlation. Mach Learn Technol. 2008;2.

28. Dersh D, Yewdell JW, Wei J. A SIINFEKL-Based System to Measure MHC Class I Antigen
Presentation Efficiency and Kinetics. Methods Mol Biol Clifton NJ. 2019;1988: 109–122.
doi:10.1007/978-1-4939-9450-2_9

29. Kim JH, Lee S-R, Li L-H, Park H-J, Park J-H, Lee KY, et al. High Cleavage Efficiency of a
2A Peptide Derived from Porcine Teschovirus-1 in Human Cell Lines, Zebrafish and Mice.
PLOS ONE. 2011;6: e18556. doi:10.1371/journal.pone.0018556

30. Shastri N, Gonzalez F. Endogenous generation and presentation of the ovalbumin peptide/Kb
complex to T cells. J Immunol Baltim Md 1950. 1993;150: 2724–2736.

31. Laumont CM, Daouda T, Laverdure J-P, Bonneil É, Caron-Lizotte O, Hardy M-P, et al.
Global proteogenomic analysis of human MHC class I-associated peptides derived from non-
canonical reading frames. Nat Commun. 2016;7: 10238. doi:10.1038/ncomms10238

32. Zanker DJ, Oveissi S, Tscharke DC, Duan M, Wan S, Zhang X, et al. Influenza A Virus
Infection Induces Viral and Cellular Defective Ribosomal Products Encoded by Alternative
Reading Frames. J Immunol Baltim Md 1950. 2019;202: 3370–3380.
doi:10.4049/jimmunol.1900070

33. Ehx G, Perreault C. Discovery and characterization of actionable tumor antigens. Genome
Med. 2019;11: 29. doi:10.1186/s13073-019-0642-x

34. Sette A, Fikes J. Epitope-based vaccines: an update on epitope identification, vaccine design
and delivery. Curr Opin Immunol. 2003;15: 461–470. doi:10.1016/s0952-7915(03)00083-9

35.   Bengio Y, Ducharme R, Vincent P, Jauvin C. A Neural Probabilistic Language Model. J
Mach Learn Res. 2003;3: 1137–1155.

36.   LeCun Y, Bengio Y, Hinton G. Deep learning. Nature. 2015;521: 436–444.
doi:10.1038/nature14539

37.   Daouda T. Mariana: The Cutest Deep learning Framework. 2015. Available:
https://www.github.com/tariqdaouda/Mariana

38.   Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison Iii CA, Smith HO. Enzymatic
assembly of DNA molecules up to several hundred kilobases. Nat Methods. 2009;6: 343–
345. doi:10.1038/nmeth.1318

39.   Bell C, English L, Boulais J, Chemali M, Caron-Lizotte O, Desjardins M, et al. Quantitative
Proteomics Reveals the Induction of Mitophagy in Tumor Necrosis Factor-α-activated
(TNFα) Macrophages. Mol Cell Proteomics MCP. 2013;12: 2394–2407.
doi:10.1074/mcp.M112.025775

40.   Karttunen J, Sanderson S, Shastri N. Detection of rare antigen-presenting cells by the lacZ
T-cell activation assay suggests an expression cloning strategy for T-cell antigens. Proc Natl
Acad Sci U S A. 1992;89: 6020–6024.

41.   Zucca S, Gagliardi S, Pandini C, Diamanti L, Bordoni M, Sproviero D, et al. RNA-Seq
profiling in peripheral blood mononuclear cells of amyotrophic lateral sclerosis patients and
controls. Sci Data. 2019;6: 190006. doi:10.1038/sdata.2019.6

42.   Sidoli S, Lopes M, Lund PJ, Goldman N, Fasolino M, Coradin M, et al. A mass spectrometry-
based assay using metabolic labeling to rapidly monitor chromatin accessibility of modified
histone proteins. Sci Rep. 2019;9: 13613. doi:10.1038/s41598-019-49894-4

Supporting information captions

Supplementary Figures

Supplementary Figure S1. Codon distribution in the shuffled datasets more closely resembles

that of amino acids, compared to the original datasets. (A) Pearson correlation (R^2) factors and

(b) Kullback-Leibler (KL) divergence between positional distribution of codons and their

corresponding amino acid in the shuffled (y axis) VS original (x axis) datasets. For all codons, the

shuffled dataset showed greater correlations (A) and smaller KL divergence to their respective

amino acid distributions than the original datasets ($p < 1 \times 10^{-8}$, assessed using unilateral paired

Student T test).

Supplementary Figure S2. Distribution of Pearson's correlation factors calculated between

codons and amino acids positional distributions in the original (green) and shuffled (coral)

datasets. 92% of codons in the shuffled dataset reflecting the amino acids distribution with a R2

> 0.95, compared to only 69% in the original dataset ($p < 5 \times 10^{-5}$).

Supplementary Figure S3. Distribution of amino acid and codon usage per position in the

original VS shuffled datasets. (A) Alanine – A. (B) Cysteine – C. (C) Aspartic acid – D. (D)

Glutamic acid – E. (E) Phenylalanine – F. (F) Glycine – G. (G) Histidine – H. (H) Isoleucine – I.

(I) Lysine – K. (J) Leucine – L. (K) Asparagine – N. (L) Proline – P. (M) Glutamine – Q. (N)

Arginine – R. (O) Serine – S. (P) Threonine – T. (Q) Valine – V. (R) Tyrosine – Y.

Supplementary Figure S4. CAMAP architecture and detailed predictions. (A) Architecture of the

ANN used in this work. (B) Results for the AUC on all train, validation and test subsets. Grey

areas represent the 95% confidence intervals. (C) Distributions of output probabilities of CAMAPs

used to calculate correlations in Supplementary Figure S5.

Supplementary Figure S5. Correlation between CAMAP prediction score trained only with pre-
MCC or post-MCC sequences. For each sequence in the test set we calculated the average
prediction score given by CAMAPs in each condition, and calculated the Pearson correlation using
the R software. Densities were calculated on all points and drawn using ggplot2. Only a random
subset of the points is represented in the figures to limit their size.

Supplementary Figure S6. Absence of correlation between CAMAP prediction score and
transcript expression levels in 4 individual B-LCL samples (each derived from a different subject).

Supplementary Figure S7. Training of CAMAP on dataset selected to reflect positive
dataset's distribution in expression levels. (A) Distribution of transcript expression levels for
normal datasets (related to Figure 2) and the dataset used here to retrain CAMAP. As shown in
this figure, the decoy dataset was selected to mirror the distribution of transcript expression level
in the hit dataset. (B) CAMAP performance (measured by the AUC) when trained using the decoy
dataset that mirrors the transcript expression levels of the hit dataset. Significance was assessed
using bilateral paired Student T test ($p = 5.36$ x 10^{-7}).

Supplementary Figure S8. Absence of correlation between CAMAP prediction score and
binding affinities for individual alleles for decoys (A) and hits (B).

Supplementary Figure S9. Training of CAMAP on dataset selected to reflect positive
dataset's distribution in binding affinities. (A) Distribution of binding affinities for normal
datasets (related to Figure 2) and the corrected dataset used to retrain CAMAP. As shown in this
figure, the decoy dataset was selected to mirror the distribution of binding affinities in the hit
dataset. (B) CAMAP performance (measured by the AUC) when trained using the decoy dataset
that mirrors the binding affinities of the hit dataset. Significance was assessed using bilateral paired
Student T test ($p = 1.21$ x 10^{-9}).

Supplementary Figure S10. Evaluation of homology in hit dataset and its impact of CAMAP

performance. (A) Proportion of unique MAPs that can be ascribed to a single origin, 2-3, or >10

possible origins. (B) Proportion of entries in the hit dataset that encode for MAPs with a single

origin, 2-3, 4-10 or >10 possible origins

Supplementary Figure S11. Gene families overrepresented in hits with >3 possible origins.

Supplementary Figure S12. CAMAP performance (AUC) when trained using either all hits

(left), hits with 10 possible origins or less (center) or hits with 3 possible origins or less (right).

Supplementary Figure S13. Kullback-Leibler divergence between hit and decoy datasets in

original codon (y-axis) or shuffled synonymous codon sequences (x-axis). Shuffled sequences

represent amino acid usage, as codon-specific information are removed with synonymous codon

shuffling.

Supplementary Figure S14. Preferences per position for all codons for CAMAP trained with

original sequences. See Materials and Methods for more details.

Supplementary Figure S15. OVA-construct alignment, showing point mutations (red lines) in

the mRNA sequences flanking the SIINFEKL MCC. (A) Comparison of the OVA-EP nucleotide

sequence to the wildtype OVA sequence. The OVA-EP and OVA-WT sequences have 93.3%

nucleotide identity for a total of 78 modified nucleotides. (B) Comparison of the OVA-RP

nucleotide sequence to the wildtype OVA sequence. The OVA-EP and OVA-WT sequences have

92.6% nucleotide identity, for a total of 86 modified nucleotides. Mutations, shown in red, are

located only in the 162 nucleotide regions flanking the SIINFEKL coding codons. Of note, the

SIINFEKL coding codons (nucleotides 772-799) were not modified between the 3 constructs.

Supplementary Figure S16. Correlations between eGFP and Ametrine fluorescence intensity at

the single cell level. Single cell eGFP and Ametrine fluorescence intensities measured at 10 hours

post-induction are shown for the OVA-WT (A), OVA-EP (B) and OVA-RP (C) constructs. N.B.:

only transduced cells are shown (eGFP+ cells).

Supplementary Figure S17. Validation of MHC-I associated peptides (MAP) dataset from

Pearson H. *et al*. (2016) using the new versions of MAP binding affinity prediction algorithm

NetMHC4.0 (A) and NetMHCpan4.0 (B).

Supplementary Figure S18. Percentage of transcript ineligibility as a function of context size.

Transcript length corresponds to *C x 2 + 27*, where *C* is the context size in nucleotides and 27 the

length of the MCCs. Related to Figure 1A.

Supplementary Tables

Supplementary Table S1. Nucleotide sequences of the EP and RP constructs. SIINFEKL

MCCs are shown in bold, while the variant regions (pre- and post-MCCs flanking sequences,

context size of 162-nucleotides) are in blue and italics. Related to Fig. 7.

Supplementary Table S2. Number of peptides needed to capture 1%, 5 and 10% of epitopes

detected by mass spectrometry in B721.221 and PBMC cell lines. The lower the number of

peptides needed to capture the respective number of epitopes, the better the performance of the

prediction model. This is also illustrated by the percentage of false identification (false positive

rate, FPR) reported here. Peptides were rank-ordered according to regression scores. Of note, only

the maximal regression score was kept for peptides with multiple potential origins.

A iTR-OVA construct

rtTA

eGFP — OVA — Ametrine

SIINFEKL peptide
Variable context

CAMAP scores

RP	0.03
WT	0.65
EP	0.98

B

Uninterrupted translation:

mRNA transcript ≈ protein

Interrupted translation:

DRiPs >

Legend
Ribosome   eGFP   Ametrine

C Peptide presentation

D Translation reporter

Symbols	p values vs WT	p values EP vs RP
EP	* p < 0.05	† p < 0.05
RP	** p < 0.01	†† p < 0.01