A recently formed triploid *Cardamine insueta* inherits leaf vivipary and submergence tolerance traits of parents

Running title: Parental legacies in contemporary triploid

Jianqiang Sun[1]§, Rie Shimizu-Inatsugi[2]§, Hugo Hofhuis[3], Kentaro Shimizu[4], Angela Hay[3],

Kentaro K. Shimizu[2,5]*, Jun Sese[6,7]*

[1]Research Center for Agricultural Information Technology, National Agriculture and Food Research

Organization, 3-1-1 Kannondai, Tsukuba, Ibaraki, 305-8517, Japan

[2]Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, CH-8057, Zurich, Switzerland

[3]Max Planck Institute for Plant Breeding Research, Germany

[4]Department of Biotechnology, Graduate School of Agricultural and Life Sciences, The University of

Tokyo, Tokyo, 113-8657, Japan

[5]Kihara Institute for Biological Research (KIBR), Yokohama City University, 641-12 Maioka,

Totsuka-ward, Yokohama, 244-0813, Japan

[6]Artificial Intelligence Research Center, National Institute of Advanced Industrial Science and

Technology, 2-3-26 Aomi, Koto-ku, Tokyo 135-0064, Japan

[7]Humanome Lab, Inc., L-HUB 3F, 1-4, Shumomiyabi-cho, Shinjuku, Tokyo, 162-0822, Japan

§equal contributions

* Correspondence:

Kentaro Shimizu

Jun Sese

Keywords: allopolyploid, homeolog, RNA-Seq, meristem formation, ecological niche

Abstract

Contemporary speciation provides a unique opportunity to directly observe the traits and

environmental responses of a new species. *Cardamine insueta* is an allotriploid species that appeared

within the past 150 years in a Swiss village, Urnerboden. In contrast to its two progenitor species, *C.*

amara and *C. rivularis* that live in wet and open habitats, respectively, *C. insueta* is found in-between

their habitats with temporal water level fluctuation. This triploid species propagates clonally and

serves as a triploid bridge to form higher ploidy species. Although niche separation is observed in

field studies, the mechanisms underlying the environmental robustness of *C. insueta* are not clear. To

characterize responses to a fluctuating environment, we performed a time-course analysis of

homeolog gene expression in *C. insueta* in response to submergence treatment. For this purpose, the

two parental (*C. amara* and *C. rivularis*) genome sequences were assembled with a reference-guided

approach, and homeolog-specific gene expression was quantified by using HomeoRoq software. We

found that *C. insueta* and *C. rivularis* initiated vegetative propagation by forming ectopic meristems

on leaves, while *C. amara* did not. We examined homeolog-specific gene expression of three species

at nine time points during the treatment. The genome-wide expression ratio of homeolog pairs was

2:1 over the time-course, consistent with the ploidy number. By searching the genes with high

coefficient of variation of expression over time-course transcriptome data, we found many known

key transcriptional factors related to meristem development and formation upregulated in both *C.*

rivularis and *rivularis*-homeolog of *C. insueta*, but not in *C. amara*. Moreover, some *amara*-

homeologs of these genes were also upregulated in the triploid, suggesting trans-regulation. In turn,

Gene Ontology analysis suggested that the expression pattern of submergence tolerant genes in the

triploid was inherited from *C. amara*. These results suggest that the triploid *C. insueta* combined

advantageous patterns of parental transcriptomes to contribute to its establishment in a new niche

along a water-usage gradient.

1   Introduction

The molecular basis of speciation has been a central question in biology (Jerry A and H. Allen,

2004). Little is known still about how a new species obtains new traits to adapt to a distinct

environment. A major obstacle in studying this is that most speciation events occurred in the past,

and thus the traits and the environment at the time of speciation are not directly observable. The

difference in traits and environments between current species may represent evolution after

speciation rather than the changes that occurred at speciation. A unique opportunity to study

speciation in action is contemporary allopolyploid speciation (Soltis and Soltis, 2009; Abbott et al.,

2013). During the past 150 years, several cases of polyploid speciation have been documented, for

example in *Tragopogon, Senecio, Mimulus, Spartina*, and *Cardamine* (Urbanska et al., 1997; Abbott

and Andrew, 2004; Ainouche et al., 2004; Soltis et al., 2004). Because polyploid speciation

immediately confers complete or partial reproductive isolation between the new polyploid and

progenitor species, a new polyploid species must establish and propagate while surrounded by

individuals with different ploidy. To overcome this situation termed "minor cytotype disadvantage",

two traits are suggested to facilitate establishment (Comai, 2005). First, the distinct environmental

niche of a polyploid species would reduce competition with progenitor species. Second, clonal

vegetative propagation or self-fertilization would assure the persistence of new polyploids at the

initial stages because meiotic abnormality is common in newly formed polyploid species. This would

be critical for odd-ploidy species including triploids, which often contribute to the formation of

higher polyploids via a so-called triploid bridge (Bretagnolle and Thompson J. D., 1995; Ramsey and

Schemske, 1998; Mable, 2003; Husband, 2004; Tayalé and Parisod, 2013; Mason and Pires, 2015).

Despite the significance of these traits, the underlying molecular mechanisms are yet to be studied.

The contemporary polyploid *C. insueta* belongs to the genus *Cardamine,* which has long been

studied for ecological polyploid speciation (Howard, 1948; Hussein, 1948), and represents adaptive

radiation by recurrent polyploidization along water-usage gradients (Shimizu-Inatsugi et al., 2016;

Akiyama et al., 2019). A major advantage to studying *Cardamine* is that it is closely related to the

model plant *Arabidopsis thaliana*, and a reference genome assembly of *Cardamine hirsuta* (Gan et

al., 2016) is publicly available, thus functional and genomic data of these model species are readily

available. One allotriploid species in *Cardamine, C. insueta* ($2n = 3x = 24$; RRA), is a textbook

example of contemporary speciation discovered by Urbanska and Landolt in 1974 (Urbanska-

Worytkiewicz and Landolt, 1974b). It was formed by the hybridization of two progenitor diploids

Cardamine amara ($2n = 2x = 16$; AA) and *Cardamine rivularis* ($2n = 2x = 16$; RR, belonging to *C.

pratensis* complex sensu lato) approximately 100–150 years ago at the valley of Urnerboden in the

Swiss Alps (Urbaska-Worytkiewicz and Landolt, 1972; Urbanska-Worytkiewicz and Landolt, 1974b;

Urbanska et al., 1997; Mandáková et al., 2013; Zozomová-Lihová et al., 2014) (**Fig. S1A**). The two

diploid progenitors have distinct ecological habitats. While *C. amara* grows in and beside water

streams, *C. rivularis* inhabits slightly moist sites, avoiding permeable and fast drying soil (Urbanska-

Worytkiewicz and Landolt, 1974b, 1974a) (**Fig. S1*B***). Around the end of the 19th to the early 20th

centuries, the deforestation and land-use conversion to grazing induced the hybridization of these two

diploids to produce the triploid species *C. insueta*, which is abundant in manured hay-meadows

(Urbaska-Worytkiewicz and Landolt, 1972; Urbanska et al., 1997; Mandáková et al., 2013).

Cytogenetic studies suggested that *Cardamine insueta* served as a triploid bridge in the formation of

pentaploid and hexaploid *C. schulzii* by the further hybridization with autotetraploid *C. pratensis*

(sensu stricto, $2n = 2x = 30$; PPPP; hypotetraploid derived from a chromosomal fusion) in

Urnerboden (Mandáková et al., 2013).

The propagation of triploids mainly depends on vegetative propagation for two reasons, high male

sterility *per se* and hay cutting and grazing in flowering season (Urbanska et al., 1997). One of the

progenitor species, *C. rivularis*, can produce plantlets on the surface of leaves and nodes by ectopic

meristem formation, which is a common feature of the *C. pratensis* complex (Smith, 1825; Salisbury,

1965; Dickinson, 1978). This characteristic is inherited by *C. insueta*, enabling it to be a dominant

species at the site despite its ploidy level (Urbanska-Worytkiewicz and Landolt, 1974a; Urbanska et

al., 1997). This type of leaf vivipary is only found in a limited number of angiosperms and assumed

to contribute to population establishment in polyploids (Dickinson, 1978). In this sense, the trait of

leaf vivipary can be considered a key factor for the establishment of this triploid.

Another interesting aspect of *C. insueta* establishment is its ecological niche shift relative to its

progenitor species. Genus *Cardamine* is known to include many submergence tolerant species

including *C. amara* (Shimizu-Inatsugi et al., 2016; Akiyama et al., 2019). An allotetraploid *C.*

flexuosa, derived from *C. amara* and *C. hirsuta* diploid progenitors, was shown to inherit parental

traits and be successful in a wider soil moisture range (Shimizu-Inatsugi et al., 2016; Akiyama et al.,

2019). The transcriptomic response of *C. flexuosa* to submergence or drought stress was shown to be

combined although attenuated compared to its progenitor species, which could confer the wider

tolerance found in the polyploid. Even though the niche separation between *C. rivularis* and *C.*

insueta is not yet clearly illustrated, our field observations are consistent with this hypothesis.

In this study, we focused on the time-course gene expression pattern of the triploid *C. insueta* and its

two diploid progenitors during submergence treatment, which induces both water stress and ectopic

meristem formation on leaves. To study the time-course data of homeologs, we employed

bioinformatic methods of variably expressed genes because data points of a time-course are not

independent and serve partly as replicates (Yamaguchi et al., 2008; Shin et al., 2014). Here we

combined the time course analysis with subgenome-classification bioinformatic workflow of

HomeoRoq (Akama et al., 2014), and detected variably expressed homeologs (VEH) during the

treatment. We address the following specific questions:

(1) What is the expression rate and the ratio of homeologous genes in triploid species in response to

submergence, either genome-wide or between each homeologous gene pair?

(2) Which gene ontology categories are enriched in VEH? Do they reflect the phenotypic trait of each

progenitor species or the triploid? How does *C. insueta* combine the expression patterns of the two

progenitors?

2   Materials and Methods

2.1   Plant materials and RNA sequencing

Cardamine insueta, *C. amara*, and *C. rivularis* plants used in this study were collected from

Urnerboden. All plants were grown together in a plant cultivation room with 16 hr light and 8 hr dark

cycle. The plants were planted in single pots, placed on trays, and watered from below.

Submergence treatment was started in the morning at 07:00. Two mature leaves were detached and

submerged in water. We isolated RNA from the floating leaflets of the three species at nine time

points after the start of submergence treatment (0 hr, 2 hr, 4 hr, 8 hr, 12 hr, 24 hr, 48 hr, 72 hr, and 96

hr) using Qiagen RNeasy kit (Qiagen, Maryland, U.S.A.). RNA quality was assessed by Bioanalyser

Nanochip (Agilent, Santa Clara, U.S.A.) and libraries quantified by Qubit (ThermoFisher, Waltham,

U.S.A.). In total 27 libraries (3 species x 9 time points) were prepared according to NEBNext Ultra™

Directional RNA Library Prep Kit for Illumina (New England Biolabs, Ipswich, U.S.A.) followed by

paired end sequencing (100bp x 2) on a HiSeq2000 with a HiSeq Paired-End Cluster Generation Kit

and HiSeq Sequencing Kit (Illumina, San Diego, U.S.A.). Trimmomatic (ver. 0.36) (Bolger et al.,

2014) was used for discarding the low-quality reads with parameters of "PE -threads 4 -phred33

ILLUMINACLIP:adapters.fa:2:30:10 LEADING:20 TRAILING:20 SLIDINGWINDOW:4:20

MINLEN:50".

2.2   Reference sequence assembly

The reference sequences of A-genome and R-genome were assembled by SNP substitution at coding
regions from the *C. hirsuta* genome (Gan et al., 2016) with the following steps. To assemble the
reference sequence of A-genome, first, we pooled all RNA-Seq reads of the nine RNA-Seq samples
of *C. amara*. Second, we mapped the reads onto the reference sequence (i.e., H-genome) using STAR
(ver. 2.3.0e) (Dobin et al., 2013). Third, we detected single-nucleotide polymorphisms (SNPs) and
short indels from the mapping result using samtools (ver. 0.1.18) (Li et al., 2009). SNPs and indels.
were defined as the polymorphic loci where at least 80% of reads have the alternative nucleotides.
Fourth, we replaced the nucleotides on the reference with the alternative nucleotides, if the
alternative nucleotide was covered by at least five reads. Finally, the gene annotations of the
assembled sequence were converted from the H-genome annotations with the replacement
information. To improve the accuracy of sequence, we used the assembled sequence as a reference
sequence, and repeated steps two through five, nine times. The resulting A-genome was used for the
mapping of individual RNA-seq data from all three species. The R-genome was also reconstructed
with the same protocol. As a result, 1,496,561 and 1,484,186 SNP regions on the H-genome were
replaced for *C. amara* and *C. rivularis*, respectively.

2.3   Evaluation of HomeoRoq classification confidence using diploids

We used HomeoRoq (ver. 2.1) (Akama et al., 2014) to classify genomic origins of homeolog-specific
reads in the nine *C. amara* and *C. rivularis* samples. Following the HomeoRoq pipeline, for each *C.*
amara sample, we used STAR to map reads onto the A-genome and R-genome and used HomeoRoq
to classify reads as *A-origin*, *R-origin*, and *unclassified*. Then, we calculated the percentage of
misclassified reads (i.e., the reads that were classified as *R-origin*). Similarly, we used HomeoRoq to
calculate the percentage of misclassified reads (i.e., the reads that were classified as *A-origin*) in each
C. rivularis sample.

2.4   Homeolog expression quantification and A-origin ratio definition of triploid

We used HomeoRoq to analyze the nine *C. insueta* samples. For each *C. insueta* sample, we used
STAR to map reads onto A-genome and R-genome and used HomeoRoq to classify reads as *A-*
origin, *R-origin*, and *unclassified*. Then, we customized HTSeq (Planet et al., 2012) to count the
number of read pairs that mapped on homeolog region for *A-origin*, *R-origin*, and *unclassified* reads
of each *C. insueta* sample separately. In the customized HTSeq, if a read mapped on the region
overlapped by multiple homeologs, a read was divided by the number of homeologs.

172 To calculate the number of fragments per kilobase mapped (FPKM) for *C. insueta* samples (I_A and I_R

173 samples), we first allocated the *unclassified* reads into *A-origin* and *R-origin* reads with A-origin

174 ratio. A-origin ratio of homeolog h at the time point s was defined as $p_h^s = a_h^s/(a_h^s + r_h^s)$, where a_h^s

175 and r_h^s are the numbers of *A-origin* and *R-origin* reads of homeolog h at the time point s,

176 respectively. Thus, the number of *A-origin* reads after *unclassified* reads allocation ($a_h'^s$) was

177 calculated as $a_h'^s = a_h^s + u_h^s p_h^s$, where u_h^s is the number of *unclassified* reads of homeolog h in

178 sample s. Similarly, $r_h'^s = a_h^s + u_h^s(1 - p_h^s)$ for *R-origin* reads. Then, FPKM of *A-origin* reads of

179 homeolog h in sample s was calculated as $10^9 a_h'^s/(L_h^A A^s)$, where L_h^A is the length of homeolog h on

180 A-genome and A^s is the total number of *A-origin* reads in sample s; likewise, FPKM of *R-origin*

181 reads was calculated as $10^9 a_h'^s/(L_h^R R^s)$, where L_h^R is the length of homeolog h on R-genome and R^s

182 is the total number of *R-origin* reads in sample s.

183 In addition, FPKM of progenitors were calculated from the total number of reads (i.e., $a_h^s + u_h^s +$

184 r_h^s). Therefore, FPKM of *C. amara* and *C. rivularis* were calculated as $10^9 (a_h^s + u_h^s + s_h^s)/(L_h^A A^s)$

185 and $10^9 (a_h^s + u_h^s + s_h^s)/(L_h^A R^s)$, respectively.

2.5 Expressed homeologs and PCA analysis

187 An expressed homeolog was defined as a homeolog with FPKM > 1.0. A homeolog expressed in a

188 sample (i.e., either *amara*-derived in *C. insueta* (I_A), *rivularis*-derived in *C. rivularis* (I_R), *C. amara*

189 or *C. rivularis*) was defined as a homeolog with FPKM > 1.0 at least at one of the nine time points. In

190 total, 21,131 homeologs were expressed at least in one sample. PCA was performed against $\log_{10}$-

191 transformed FPKM of these 21,131 expressed homeologs. To avoid calculating $\log_{10}0$, the $\log_{10}$-

192 transfromed FPKM was truly calculated as $\log_{10}(\text{FPKM} + 1)$.

2.6 Identification of variably expressed homeologs (VEH) and Gene ontology (GO)
  enrichment analysis

195 Mean and coefficient of variation (CV) were calculated from $\log_{10}$-transformed FPKM over the nine

196 time points. VHE was defined as an homeolog satisfied the mean > 1.0 and the CV > 0.20. We

197 identified from I_A, I_R, *C. amara*, and *C. rivularis* samples, separately.

198 Gene ontology (GO) enrichment analysis was performed for the four variably expressed homeolog

199 (VEH) sets with R packages clusterProfiler (ver. 3.12.0) and org.At.tair.db (ver. 3.8.2) (Yu et al.,

200 2012). To remove redundancies of GO categories, only GO categories which are associated with 10–

500 *Cardamine* homeologs and below the third level in the GO category hierarchy were used. The

threshold FDR = 0.1 was used for cutoff of significantly enriched GO categories.

3   Results

3.1   Plantlet induction on *C. insueta* and *C. rivularis* leaves by submergence

At the field of Urnerboden valley, we could scarcely observe normal seed setting on *C. insueta*, but

small plantlets on leaves were frequently observed after flowering, as described previously

(Urbanska, 1977). We also observed small plantlets on the leaves of *C. rivularis*. In contrast, *C.*

amara does not form plantlets on leaves, rather adventitious roots and shoots were formed from

rhizomes. In the natural habitat, the plantlet formation of *C. rivularis* and *C. insueta* can be seen at

flowering to post-flowering season (Salisbury, 1965; Urbanska, 1977). It was also reported that *C.*

pratensis (which is closely related to *C. insueta* or considered the same species) tend to bear more

plantlets on the leaves in damper sites than in drier sites (Salisbury, 1965), implying that high

moisture could be the trigger for meristem formation. Thus, we tested plantlet induction by

submergence treatment using dissected leaves with this trio of species in the lab. We detached mature

leaves from mother plants propagated in a climate chamber and floated the leaves on water. Within

16 h, we observed the activation of dormant shoot meristems and initiation of ectopic root meristems,

which formed visible plantlets on *C. rivularis* leaves 96 hours after submergence (**Fig. S2, Dataset**

S1). Induction of ectopic plantlets followed a similar time-course in *C. insueta*. In contrast, plantlet

induction was not observed on the leaves of *C. amara*. In addition, during the 96-hr treatment, no

symptoms of necrosis appeared on any of the leaves, suggesting that all three species have some

submergence tolerance for at least 96 hr.

3.2   Gene annotation on the two diploid progenitor reference sequences

To detect how homeologous genes are expressed in plantlet induction and submergence treatment,

we harvested time-course RNA-Seq samples of *C. insueta* and diploid progenitor leaves at nine time

points after initial submergence (i.e., 0, 2, 4, 8, 12, 24, 48, 72, and 96 hr) (**Fig. S3***A*). We harvested

the first lateral leaflet pair in young leaves with no ectopic plantlets. To quantify homeolog-specific

gene expression, we assembled the genomes of *C. amara* (A-genome) and *C. rivularis* (R-genome),

respectively, using the same pipeline of a reference-guided approach using RNA-Seq reads (**Fig.**

S3*B*). The genome sequence of a close relative, *C. hirsuta* (H-genome) (Gan et al., 2016), was used

as a reference. The A-genome structure is reported to be almost perfectly collinear with that of H-

genome, except for one pericentric inversion at chromosome 1, by cytological studies (Mandáková et

al., 2013, 2014). The genome structures of the A-genome and R-genome are also similar to each

other (Mandáková et al., 2013). The length of assembled reference sequences of A-genome and R-

genome are 198,651,635 and 198,654,862 nucleotides, respectively, which are nearly the same as the

length of the original H-genome (198,654,690 nucleotides). We also annotated the orthologous genes

of *C. amara* and *C. rivularis* according to the information of *C. hirsuta* H-genome. In total, we found

23,995 and 24,115 genes covered by at least one read among the nine time points on the assembled

A-genome and R-genome, respectively. These gene sets, which correspond to 81.5% and 81.7% of

29,458 genes in H-genome, respectively, were defined as expressed and used for the following

analysis.

3.3   Expression ratio from each subgenome is consistent with the number of chromosomes

We applied the HomeoRoq analysis pipeline (Akama et al., 2014) to classify the origin of each RNA-

seq read of *C. insueta* samples to either *A-origin* (i.e., the genomic origin of the read is A-

subgenome) or *R-origin* (**Fig. S3*C***). After filtering for read quality, 10.6 million read pairs on

average among the nine samples could be classified as homeolog-specific read pairs (**Dataset S2**). Of

the total homeolog-specific read pairs in the *C. insueta* 0 hr sample, 27.3% and 56.7% of read pairs

were classified as *A-origin* and *R-origin*, respectively. The remaining 16.0% of read pairs could be

classified to neither *A-origin* nor *R-origin* (*unclassified*) due to the lack of SNPs or the identical

sequence on the correspondence region. As a whole genome, the ratio of *A-origin* to *R-origin* reads

was approximately 1:2. When we analyzed all samples from the other eight time points, we observed

a slight increase in the proportion of *A*-origin reads in correlation with the time point, from 1:2.07 at

0 hr to 1:1.90 at 96 hr (**Dataset S2**). Instead of this minor transition, the expression ratio between

subgenomes remained A:R ≈ 1:2 with *C. insueta* samples at all time points, indicating that the

expression ratio from each subgenome is consistent with the number of chromosome regardless of

the submergence treatment.

3.4   Most homeolog pairs were expressed in proportion to the subgenomes in *C. insueta*

To investigate the proportion of expression levels of homeolog pairs in *C. insueta*, we quantified the

expression level of each homeolog pair at each time point. We found that (i) the correlation between

the expression levels of homeolog pairs was higher than 0.81 at any time point (**Fig. 1*A* and Fig. S4**).

However, (ii) the expression levels of most homeologs expressed from the A-subgenome (A-

homeolog) were approximately half that of R-homeologs. To understand the proportion of expression

levels of homeolog pairs in detail, we calculated A-origin ratio—the proportion of A-homeolog

expression level to the total A-homeolog and R-homeolog expression levels—for all homeolog pairs

at each time point. We found that the distribution of A-origin ratios had a gentle peak at the position

of 0.33 at all time points (**Fig. 1*B* and Fig. S5**). This result suggests the expression ratio of the

majority of homeolog pairs is consistent with the copy number, i.e. the subgenome-set numbers of

the triploid. In addition, we found two sharp peaks at both edges, the positions of 0.0 and 1.0, of A-

origin ratio, which represent the homeologs only expressed in of either subgenome.

Additionally, to investigate whether the A-origin ratio changes during the submergence treatment, we

compared the A-origin ratio distributions between different time points. The patterns of all time

points were correlated to each other, with the least coefficiency (0.66) between 0 hr and 2 hr (**Fig. 1*B***

and Table S1). This result indicates that A-origin ratios did not change drastically in most homeolog

pairs by the submergence treatment, but a limited number of homeolog pairs change the expression

balance.

3.5   The whole genome expression pattern of each *C. insueta* subgenome is closer to that of its
        progenitor genome

To gain an overview of how homeologous gene expression varies at the whole genome level among

C. insueta and the progenitor species *C. amara* and *C. rivularis*, we conducted principal component

analysis (PCA). PCA was performed against the log_{10}-transformed FPKM of 21,131 expressed

homeologs (**Fig. 2**). We found that the first principal component (PC1) grouped samples into two

groups: the one with A-homeologs of *C. insueta* (I_A) and *C. amara* (*A*) samples and the other with R-

homeologs of *C. insueta* (I_R) and *C. rivularis* (*R*) samples. In addition, we also found that the second

principal component (PC2) grouped samples into two groups: one consisting of polyploid samples (I_A

and I_R samples, lower side of **Fig. 2A**) and the other consisting of diploid samples (*A* and *R* samples,

upper side of **Fig. 2A**). By PC1 and PC2, the samples were grouped into four clusters according to

the subgenome type. In contrast, by PC2 and the third principal component (PC3), we observed the

transition according to the treatment time, showing a characteristic transition from 0 to 12 hr, and the

recurrence of 24, 48, 72 and 96 hr samples towards 0 hr samples in each subgenome, which might

reflect the combined effect of submergence stress and circadian rhythm (**Fig. 2B**). The result of PC1

suggests that the majority of the homeologs of I_A and I_R should retain a similar expression pattern to

each parent, A and R. When we focus on PC2, the distance between R and I_R is slightly closer than

that between A and I_A. This might reflect the difference in the number of subgenome sets in the

triploid, $A:R$=1:2, implying a stronger effect from the pattern with more subgenome sets.

3.6   VEHs related to submergence and their GO enrichment analysis

To understand the difference among species in plantlet formation on the leaf and in submergence

response, we focused on the homeologs with a higher expression change during the treatment.

Standard tools to identify differentially expressed genes between different conditions are not directly

applicable to time-course data, in which expression levels of neighboring time points may be highly

correlated. We defined variably expressed homeologs (VEHs) according to the coefficient of

variation (CV) among the expression levels of the nine time points, since CV is used for identifying

variably expressed genes in various studies involving time-course analysis (Czechowski et al., 2005;

Yamaguchi et al., 2008; Shin et al., 2014; Zhao et al., 2017). We identified 1,194, 1,144, 1,030, and

1,063 VEHs from I_A, I_R, A and R genome/subgenome with the cutoff CV > 0.2 throughout the

treatment, respectively (**Dataset S3**). We visualized the patterns by focusing on two genes that were

expected to be affected (**Fig. S6**). The genes associated with ethylene-response such as *ERF1*

(AT3G23240) (Chao et al., 1997; Solano et al., 1998) and circadian rhythm such as *CCA1*

(AT2G46830) (Alabadí et al., 2001) were identified as VEHs in all samples, which should reflect the

ethylene-response to submergence and circadian rhythm response, respectively. The expression

pattern of these two homeologs were similar among all four VEH sets from I_A, I_R, A and R (**Fig. S6**).

In addition to these common VEHs, we also found more homeologs identified as VEHs only in one

to three samples (**Fig. S7**).

To investigate the biological processes of VEH sets of I_A, I_R, A and R, we performed gene ontology

(GO) enrichment analysis against the four VEH sets (**Table 1**, **Dataset S4**). The numbers of enriched

GO categories were 146, 155, 160, and 181, respectively for I_A, I_R, A and R. A value of negative

$\log10$(q-value) more than 1 was defined as significant, and a higher value indicates stronger

enrichment. We found that some of the GO categories related to water stress, including GO:0006066

(alcohol metabolic process), GO:0009723 (response to ethylene) and GO:0009414 (response to water

deprivation), were enriched in all four VEH sets (**Table 1**). As gas diffusion rates are restricted under

water, submergence of plants induces ethylene accumulation and low oxygen availability, which

could result in the reorganization of the ethylene-response pathway and fermentation pathway (e.g.

anaerobic respiration and alcohol metabolism). The enriched categories GO:0006066 and

GO:0009723 indicate that I_A, I_R, A, and R all respond to ethylene and hypoxia signals with the

submergence treatment. Two alcohol related categories (GO:0006066 alcohol metabolic process and

GO:0046165 alcohol biosynthetic process) were more strongly enriched in A and I_A, which was two

orders of magnitude higher than I_R and R (>2 difference in negative log10(q-value) in **Table 1**). In

addition, GO:0009414 (response to water deprivation), which encompasses the expression changes of

aquaporin genes and ethylene-responsive genes (**Dataset S5**), was enriched.

In contrast, some GO categories related to submergence stress were only above the significance

threshold in part of the four VEH sets with various combinations (**Table 1**). The two categories

related to ethylene metabolism, GO:0009873 (ethylene-activated signaling pathway) and

GO:0071369 (cellular response to ethylene stimulus), were not detected in I_R but all other three. All

these ethylene related GO categories were most strongly enriched in A, suggesting larger number of

genes are detected than other VEH sets. In addition, the categories related to abscisic acid signaling,

which is known to work antagonistically to ethylene, GO:0009738 (abscisic acid-activated signaling

pathway) and GO:0071215 (cellular response to abscisic acid stimulus), were also detected only in A

with many inactivated genes by treatment. In contrast, the categories related to oxidative stress

showed the strongest enrichment in R than others, suggesting higher intensity of oxidative stress in *C.*

rivularis than other species.

3.7   VEHs related to meristem and their GO enrichment analysis

Among the GO categories enriched in four VEH sets, three categories were related to meristem

activity: GO:0035266 (meristem growth), GO:0010075 (regulation of meristem growth), and

GO:0048509 (meristem development) (**Table 1**). They were enriched only in VEH sets of I_A and I_R,

but not in A and R, although *C. rivularis* can also produce ectopic meristems.

We analyzed the expression pattern of several known transcriptional factors which could be involved

in ectopic meristem formation and development in *Cardamine* (**Fig. 3** and **Fig. S8**). Class I Knotted1-

like homeobox (KNOX) transcription factors function to maintain shoot apical meristem activity in

many different plant species (Vollbrecht et al., 1991; Long et al., 1996; Hay and Tsiantis, 2010).

Importantly, the overexpression of *SHOOTMERISTEMLESS* (*STM*) and another KNOX gene,

Arabidopsis knotted 1-like gene (*KNAT1*) are known to cause ectopic meristem formation on the leaf

in *A. thaliana* (Chuck et al., 1996; Williams, 1998). Moreover, an *STM* ortholog is required for leaf

vivipary in *Kalanchoë daigremontiana* (Garcês et al., 2007), a clonal propagation trait that is also

observed in *C. rivularis* and *C. insueta*. As summarized in Fig. 3, orthologs of the four *A. thaliana*

KNOXI genes, *STM*, *KNAT1, KNAT2 and KNAT6*, showed upregulated expression in all or any of I_A,

I_R and R, but not in A. In addition, we also found that *PDF1* increased expression levels in *C. insueta*

(both I_A and I_R) and R, which is exclusively detected in the L1 layer of shoot apical meristem

throughout the shoot development of *Arabidopsis* (Abe et al., 1999). Three other transcription factor-

encoding genes, *CUC2*, *CUC3* and *LAS*, which contribute to ectopic shoot apical meristem in tomato

leaves (Rossmann et al., 2015), were induced in R and I_R but not in A.

The expression of genes related to root apical meristem maintenance and formation showed similar

patterns to those related to shoot apical meristem formation. Transcription factors with an AP2/ERF

domain for the maintenance of root apical meristem (*PLT1*, *PLT2* and *PLT3;* (Drisch and Stahl,

2015)), were scarcely expressed in A but induced in others. For genes with similar function, *SHR* and

SCR, expression level of *SHR* was increased in all four sets, while that of *SCR* was upregulated only

temporarily and reverted in 24 hr. The expression of *WOX5*, a key factor to maintain the root stem

cell (Sarkar et al., 2007), was very low in all sets, most probably due to the extremely limited

expression area only at the quiescent center.

Many of the above-mentioned transcription factors contributing to meristem formation and

maintenance are known to be related to or controlled by auxin, thus the transportation of auxin might

be also involved in ectopic meristem development in *C. insueta* and *C. rivularis*. One of the auxin

transporter genes, *PIN1* was induced by the treatment in all four sets soon after the start of

submergence, but after 24 hr the high expression level was only retained in I_A, I_R, and R. On the other

hand, the other auxin transporter genes *PIN3*, *PIN4*, and *PIN7* were temporarily induced by the

treatment, but soon decreased among all sets. The peaks of expression of these three genes were at 4–

12 hr in A, 12–24 hr in R, and 8–24 hr in I_A and I_R, suggesting involvement in meristem formation

and development. By the VEH enrichment analysis, GO:0060918 (auxin transport) and GO:0009926

(auxin polar transport) were enriched in I_A and I_R. In contrast, the other two auxin related GO

categories, GO:0009850 (auxin metabolic process) and GO:0009851 (auxin biosynthetic process),

were enriched in only two parents, A and R.

4   **Discussion**

4.1   Applicability of HomeoRoq to diverse ploidy levels

HomeoRoq was developed to classify genomic origins of RNA-Seq reads of allopolyploids

consisting of two subgenomes (Akama et al., 2014), and has already been applied to *Arabidopsis*

kamchatica ($2n = 4x = 32$; HHLL), an allotetraploid between two diploids of *Arabidopsis halleri* ($2n$

$= 2x = 16$; HH) and *Arabidopsis lyrata* ($2n = 2x = 16$; LL). Here, we successfully applied HomeoRoq

to another species with a different ploidy level. The average proportions of the reads mapped on the

wrong genome in *C. amara* and *C. rivularis* samples were $1.1 \pm 0.1\%$ and $1.2 \pm 0.1\%$, respectively

(**Dataset S2**). This high accuracy is comparable to the evaluation of the *A. kamchatica* data, 1.23%–

1.64% (Akama et al., 2014; Kuo et al., 2020).

The proportion of *unclassified* reads in this study, which has the same matching rates on both

parental genomes, was very close to that in the *A. kamchatica* study. In this study, $11.5 \pm 2.0\%$ of

reads in *C. insueta* samples were *unclassified* on average, compared to 11.0% in *A. kamchatica*

(Akama et al., 2014), suggesting a similar divergence level between subgenomes in the two cases.

Considering the percentage of unclassified reads and the low misclassification rate with diploid

progenitors, HomeoRoq can be applied to genomes of any ploidy level providing that the genome

consists of two types of subgenome.

4.2   Total gene expression level of each subgenome is consistent with the chromosome number

The ratio of *A-origin* to *R-origin* reads in *C. insueta* was approximately 1:2. This result is consistent

with the distribution of A-origin ratio showing a gentle peak at around 0.33 with a smooth decrease

toward the edges (**Fig. 1**). This distribution indicates that expression ratios of most homeologs

correlates with the copy number. A similar tendency could be found in other Brassicaceae

allotetraploids (2, 5, 6). In the analysis of triploid banana ($2n = 3x = 33$; ABB), a hybrid between

Musa acuminata ($2n = 2x = 22$; AA) and *M. balbisiana* ($2n = 2x = 22$; BB), the read proportion is

distributed around 0.66 for the B alleles by 155 homeologs with rather high expression level detected

by LC-MSMS as isoforms (van Wesemael et al., 2018). This could also be seen in hexaploid bread

wheat consisting of three subgenomes, where 70% of genes showed balanced expression among

homeologs (Ramírez-González et al., 2018). So far, this consistency between ploidy number and

expression ratio looks like a general rule in many species with some exceptions like tetraploid cotton

(Yoo et al., 2013).

In addition to the majority of genes that show balanced expression between homeologs, a limited

proportion of genes show significant differential expression. Even though a direct comparison among

studies is difficult due to different thresholding policies, the number of genes with unbalanced

homeolog expression tends to be the minor fraction in many quantitative studies. Further studies

should show whether a similar pattern is observed in even higher ploidy levels or other odd ploidies.

4.3   Limited number of homeolog pairs changed expression ratio in submergence condition

Though the number of homeologs with unbalanced expression is smaller than that with balanced

expression, they could play a significant role in speciation of polyploid species, especially for

achieving a combined trait from progenitors. A series of studies have reported that homeolog

expression ratios can be changed depending on external environments (Bardil et al., 2011; Dong and

Adams, 2011; Akama et al., 2014; Paape et al., 2016). Akama *et al*. evaluated the changes of the

homeolog expression ratio of *A. kamchatica* after cold treatment (Akama et al., 2014). They reported

that the homeolog expression ratios before and after cold treatment were highly correlated ($R_2 =$

0.87), and only 1.11% of homeolog pairs statistically significantly changed in expression ratios in

response to cold treatment (Akama et al., 2014). A similar result was reported for zinc treatment of *A.*

kamchatica. The correlation of homeolog expression ratios between zinc treatment and control

ranged from 0.89 to 0.94, and 0.3%–1.5% of homeologs significantly changed expression ratios after

Zn treatment (Paape et al., 2016).

In this study using another Brassicaceae species, *C. insueta*, the correlation coefficients of A-origin

ratios between 0 hr and the other time-points ranged from 0.68 to 0.82 (**Table S1**). The lowest

correlation occurring between 2 hr and other time points may suggest that the initial reaction to the

treatment had the strongest effect on gene expression. The overall high correlations among time

points indicate that the expression ratios of most homeologs do not change considerably in response

to treatment. Even though *C. rivularis* and *C. amara* show species-specific responses to

submergence, leaf vivipary and submergence tolerance respectively, no specific expression

preference or dominance of either progenitor was detected in the triploid. This suggests that

transcriptional changes in only a limited number of homeologs, rather than genome-wide, might be

responsible for the control of physiological change under submergence conditions.

4.4   Triploid inherited advantageous traits from progenitors

Only about 6% of the expressed genes were detected as VEH throughout the 96-hr treatment in each

genome and subgenome, suggesting the criteria were fairly conservative. Among enriched GO

categories are water stress related ones, particularly ethylene-response and fermentation.

Fermentation metabolism in plants is important for submergence stress. We found more VEH genes

in the fermentation-related categories in the diploid *C. amara* and the *amara*-derived subgenome of

C. insueta than counterparts (Table 1). This suggests that *C. insueta* inherited the fermentation ability

as a submergence response more largely from *C. amara* side. The ethylene signaling pathway should

be stimulated in all three species as many related GO categories are found enriched in all VEH genes.

However, the stress level seems to be variable according to the species as shown in the difference of

enriched GO categories. In all of these ethylene related GO categories, *C. amara* had the strongest

enrichment (i.e., highest number of VEH genes), and the enrichment in *amara*-derived subgenome

was stronger than in *rivularis*-derived subgenome in *C. insueta*. These enrichment intensities should

suggest that *C. amara* has higher acclimation ability to submergence through an activation of alcohol

metabolic pathway and alteration in hormone signaling pathway and thus suffer from less oxidative

stress as a result, as speculated by its habitat and a previous study (Shimizu-Inatsugi et al., 2016). In

addition, in *C. insueta*, the contribution to the stress response of I_A seems larger than that of I_R, found

as stronger enrichment in I_A than in I_R.

GO enrichment analysis with VEH genes also showed three GO categories related to meristem,

GO:0035266 (meristem growth), GO:0010075 (regulation of meristem growth) and GO:0048509

(meristem development). They were only enriched in the VEH sets of I_A and I_R, but not above the

significance threshold in two parents, despite the fact that *C. rivularis* also produces plantlets on the

leaf by the activation of ectopic meristems. This might imply that the ability to form ectopic plantlets

in response to submergence is enhanced in the triploid *C. insueta* compared to the diploid *C.*

rivularis. Considering the disadvantage in sexual reproduction due to the odd ploidy, effective

vegetative propagation through plantlets might have been critically important for *C. insueta*.

The expression pattern of known key regulatory genes that function to maintain meristem activity

showed two typical patterns, as shown in **Fig. 3** and **Fig S8**. Expression of these genes was

upregulated in *C. rivularis* (*R*, **Fig. 3**) but not in *C. amara* (*A*, **Fig. 3**) in response to submergence.

Expression of these genes was also upregulated in the *C. insueta* subgenome I_R, but followed two

different patterns in the I_A subgenome. These patterns could be categorized as either non-induced,

similar to *C. amara*, or induced, similar to *C. rivularis*, suggesting that non-induced homeologs could

be *cis*-regulated by I_A, and induced homeologs could be *trans*-regulated by I_R. One possibility is that

this difference reflects the developmental timing of gene expression during meristem formation. For

example, the *cis*-regulated genes *STM* and *CUC2* are expressed earlier during embryogenesis in *A.*

thaliana than the *trans*-regulated genes *KNAT6* and *KNAT1/BP* (Hay and Tsiantis, 2010). This

variation might imply a regulatory relationship among these genes in the gene regulatory network

controlling plantlet formation in *C. insueta* leaves. This type of information might provide insights

that warrant further study into the molecular mechanism of leaf vivipary in *C. rivularis* and *C.*

insueta.

5   Acknowledgments

482 We would like to thank Walter Brücker for the support in the fieldwork in Urnerboden, Martin
483 Lysak, Terezie Mandáková, Karol Marhold, Judita Zozomová-Lihová, Pamela Soltis and Douglas
484 Soltis for discussion. Data analysis was partially performed on National Institute of Genetics (NIG)
485 at Research Organization of Information and Systems (ROIS), Japan.

6   Author Contributions

487 JSun analysed data. JSun and RSI wrote the manuscript. RSI, AH, KKS and JSese refined the

488 manuscript. HH performed experiment. AH, KKS, JSese supervised the project. All authors read,

489 corrected, and approved the manuscript.

7   Conflict of Interest

491 The authors declare that the research was conducted in the absence of any commercial or financial

492 relationships that could be construed as a potential conflict of interest.

8   Contribution to the Field Statement

In the research of genome evolution, a newborn species offers a unique and interesting case to study

how the genome evolves during speciation and adaptation. A *Brassicaceae* plant, *Cardamine insueta*,

which was born in a small village in Swiss Alps in the 20th century, applies to this. *C. insueta* is

generated by the hybridization and genome duplication between two closely related progenitor

species, which owned their respective habitats. This plant has adapted to a new habitat covering the

intermediate area between the progenitor species. How could it achieve this in the course of

speciation? The usage of the genes inherited from two progenitors should have the clue to answer this

question. We studied the gene expression pattern of *C. insueta* to analyze how it regulates the two

types of the same gene inherited from its diploid progenitors, and found that it conserves the gene

expression pattern of the advantageous progenitor according to different traits. For the trait of

submergence tolerance, it exploits the pattern of one progenitor having the trait. For another trait,

clonal propagation, it exploits the pattern of the other progenitor having this trait. This result

contributes to our understanding of speciation, and how various genes are affected in a speciation

process.

9   Funding

This study was funded by the Swiss National Science Foundation to RSI (Marie-Heim Högtlin grant)

and KKS (31003A_182318); by University Research Priority Programs, Evolution in Action of the

University of Zurich to RSI and KKS; by the Human Frontier Science Program to KKS, AH, and

JSese; by the Japan Science and Technology Agency, Core Research for Evolutionary Science and

Technology grant number JPMJCR16O3, Japan; and by KAKENHI grant numbers 16H06469 to

JSese, KKS, and JSun.

10   References

Abbott, R., Albach, D., Ansell, S., Arntzen, J. W., Baird, S. J. E., Bierne, N., et al. (2013).
Hybridization and speciation. *J. Evol. Biol.* 26, 229–246. doi:10.1111/j.1420-9101.2012.02599.x

Abbott, R., and Andrew, J. (2004). Origins, establishment and evolution of new polyploid species:
Senecio cambrensis and S. eboracensis in the British Isles. *Biol. J. Linn. Soc.* 82, 467–474.
doi:10.1111/j.1095-8312.2004.00333.x

Abe, M., Takahashi, T., and Komeda, Y. (1999). Cloning and characterization of an L1 layer-specific
gene in *Arabidopsis thaliana*. *Plant Cell Physiol.* 40, 571–580.
doi:10.1093/oxfordjournals.pcp.a029579

Ainouche, M. L., Baumel, A., and Salmon, A. (2004). Spartina anglica C. E. Hubbard: a natural
model system for analysing early evolutionary changes that affect allopolyploid genomes. *Biol.*
J. Linn. Soc. 82, 475–484. doi:10.1111/j.1095-8312.2004.00334.x

Akama, S., Shimizu-Inatsugi, R., Shimizu, K. K., and Sese, J. (2014). Genome-wide quantification of
homeolog expression ratio revealed nonstochastic gene regulation in synthetic allopolyploid
Arabidopsis. *Nucleic Acids Res.* 42, 1–15. doi:10.1093/nar/gkt1376

Akiyama, R., Sun, J., Hatakeyama, M., Lischer, H. E. L., Briskine, R. V., Hay, A., et al. (2019).
Fine-scale ecological and transcriptomic data reveal niche differentiation of an allopolyploid
from diploid parents in *Cardamine*. *bioRxiv*, 600783. doi:10.1101/600783

Alabadí, D., Oyama, T., Yanovsky, M. J., Harmon, F. G., Más, P., and Kay, S. A. (2001). Reciprocal
regulation between TOC1 and LHY/CCA1 within the *Arabidopsis* circadian clock. *Science* 293,
880–883. doi:10.1126/science.1061320

Bardil, A., de Almeida, J. D., Combes, M. C., Lashermes, P., and Bertrand, B. (2011). Genomic
expression dominance in the natural allopolyploid *Coffea arabica* is massively affected by
growth temperature. *New Phytol.* 192, 760–774. doi:10.1111/j.1469-8137.2011.03833.x

Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* 30, 2114–2120. doi:10.1093/bioinformatics/btu170

Bretagnolle, F., and Thompson J. D. (1995). Gametes with the somatic chromosome number:
mechanisms of their formation and role in the evolution of autopolyploid plants. *New Phytol.*
129, 1–22. doi:10.1111/j.1469-8137.1995.tb03005.x

Chao, Q., Rothenberg, M., Solano, R., Roman, G., Terzaghi, W., and Ecker†, J. R. (1997). Activation
of the ethylene gas response pathway in Arabidopsis by the nuclear protein ETHYLENE-
INSENSITIVE3 and related proteins. *Cell* 89, 1133–1144. doi:10.1016/S0092-8674(00)80300-1

Chuck, G., Lincoln, C., and Hake, S. (1996). *KNAT1* induces lobed leaves with ectopic meristems
when overexpressed in Arabidopsis. *Plant Cell* 8, 1277–1289. doi:10.1105/tpc.8.8.1277

Comai, L. (2005). The advantages and disadvantages of being polyploid. *Nat. Rev. Genet.* 6, 836–
846. doi:10.1038/nrg1711

Czechowski, T., Stitt, M., Altmann, T., and Udvardi, M. K. (2005). Genome-wide identification and
testing of superior reference genes for transcript normalization. *Society* 139, 5–17.
doi:10.1104/pp.105.063743.1

Dickinson, T. A. (1978). Epiphylly in angiosperms. *Bot. Rev.* 44, 181–232. doi:10.1007/BF02919079

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., et al. (2013). STAR:
ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21.
doi:10.1093/bioinformatics/bts635

Dong, S., and Adams, K. L. (2011). Differential contributions to the transcriptome of duplicated
genes in response to abiotic stresses in natural and synthetic polyploids. *New Phytol.* 190, 1045–
1057. doi:10.1111/j.1469-8137.2011.03650.x

Drisch, R. C., and Stahl, Y. (2015). Function and regulation of transcription factors involved in root
apical meristem and stem cell maintenance. *Front. Plant Sci.* 6, 1–8.
doi:10.3389/fpls.2015.00505

Gan, X., Hay, A., Kwantes, M., Haberer, G., Hallab, A., Ioio, R. Dello, et al. (2016). The *Cardamine*
hirsuta genome offers insight into the evolution of morphological diversity. *Nat. plants* 2,
16167. doi:10.1038/nplants.2016.167

Garcês, H. M. P., Champagne, C. E. M., Townsley, B. T., Park, S., Malhó, R., Pedroso, M. C., et al.
(2007). Evolution of asexual reproduction in leaves of the genus *Kalanchoë. Proc. Natl. Acad.*
Sci. U. S. A. 104, 15578–15583. doi:10.1073/pnas.0704105104

Hay, A., and Tsiantis, M. (2010). KNOX genes: versatile regulators of plant development and
diversity. *Development* 137, 3153–3165. doi:10.1242/dev.030049

Howard, H. W. (1948). Chromosome number of *Cardamine pratensis*. *Nature* 161, 277–277.
doi:10.1038/161277a0

Husband, B. C. (2004). The role of triploid hybrids in the evolutionary dynamics of mixed-ploidy
populations. *Biol. J. Linn. Soc.* 82, 537–546. doi:10.1111/j.1095-8312.2004.00339.x

Hussein, F. (1948). Chromosome number of *Cardamine pratensis*. *Nature* 161, 1015–1015.
doi:10.1038/1611015a0

Jerry A, C., and H. Allen, O. (2004). Speciation. Sunderland, Mass: Sinauer Associates

Kuo, T. C. Y., Hatakeyama, M., Tameshige, T., Shimizu, K. K., and Sese, J. (2020). Homeolog
expression quantification methods for allopolyploids. *Brief. Bioinform.* 21, 395–407.
doi:10.1093/bib/bby121

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., et al. (2009). The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.
doi:10.1093/bioinformatics/btp352

Long, J. A., Moan, E. I., Medford, J. I., and Barton, M. K. (1996). A member of the KNOTTED class
of homeodomain proteins encoded by the *STM* gene of *Arabidopsis*. *Nature* 379, 66–69.
doi:10.1038/379066a0

Mable, B. K. (2003). Breaking down taxonomic barriers in polyploidy research. *Trends Plant Sci.* 8,
582–590. doi:10.1016/j.tplants.2003.10.006

Mandáková, T., Kovařík, A., Zozomová-Lihová, J., Shimizu-Inatsugi, R., Shimizu, K. K.,
Mummenhoff, K., et al. (2013). The More the Merrier: Recent Hybridization and Polyploidy in
Cardamine. *Plant Cell* 25, 3280–3295. doi:10.1105/tpc.113.114405

Mandáková, T., Marhold, K., and Lysak, M. A. (2014). The widespread crucifer species *Cardamine*
flexuosa is an allotetraploid with a conserved subgenomic structure. *New Phytol.* 201, 982–992.
doi:10.1111/nph.12567

Mason, A. S., and Pires, J. C. (2015). Unreduced gametes: Meiotic mishap or evolutionary
mechanism? *Trends Genet.* 31, 5–10. doi:10.1016/j.tig.2014.09.011

Paape, T., Hatakeyama, M., Shimizu-Inatsugi, R., Cereghetti, T., Onda, Y., Kenta, T., et al. (2016).
Conserved but attenuated parental gene expression in allopolyploids: constitutive zinc
hyperaccumulation in the allotetraploid *Arabidopsis kamchatica*. *Mol. Biol. Evol.* 33, 2781–
2800. doi:10.1093/molbev/msw141

Planet, E., Attolini, C. S. O., Reina, O., Flores, O., and Rossell, D. (2012). htSeqTools: High-
throughput sequencing quality control, processing and visualization in R. *Bioinformatics* 28,
589–590. doi:10.1093/bioinformatics/btr700

Ramírez-González, R. H., Borrill, P., Lang, D., Harrington, S. A., Brinton, J., Venturini, L., et al.

606 (2018). The transcriptional landscape of polyploid wheat. *Science* 361, eaar6089.
607 doi:10.1126/science.aar6089

608 Ramsey, J., and Schemske, D. W. (1998). Pathways, mechanisms, and rates of polyploid formation in
609 flowering plants. *Annu. Rev. Ecol. Syst.* 29, 467–501. doi:10.1146/annurev.ecolsys.29.1.467

610 Rossmann, S., Kohlen, W., Hasson, A., and Theres, K. (2015). Lateral suppressor and Goblet act in
611 hierarchical order to regulate ectopic meristem formation at the base of tomato leaflets. *Plant J.*
612 81, 837–848. doi:10.1111/tpj.12782

613 Salisbury, E. J. (1965). The reproduction of *Cardamine pratensis* L. and *Cardamine palustris*
614 Peterman particularly in relation to their specialized foliar vivipary, and its deflexion of the
615 constraints of natural selection. *Proc. R. Soc. London. Ser. B. Biol. Sci.* 163, 321–342.
616 doi:10.1098/rspb.1965.0072

617 Sarkar, A. K., Luijten, M., Miyashima, S., Lenhard, M., Hashimoto, T., Nakajima, K., et al. (2007).
618 Conserved factors regulate signalling in *Arabidopsis thaliana* shoot and root stem cell
619 organizers. *Nature* 446, 811–814. doi:10.1038/nature05703

620 Shimizu-Inatsugi, R., Terada, A., Hirose, K., Kudoh, H., Sese, J., and Shimizu, K. K. (2016). Plant
621 adaptive radiation mediated by polyploid plasticity in transcriptomes. *Mol. Ecol.* 26, 193-207.
622 doi:10.1111/mec.13738

623 Shin, H., Shannon, C. P., Fishbane, N., Ruan, J., Zhou, M., Balshaw, R., et al. (2014). Variation in
624 RNA-Seq transcriptome profiles of peripheral whole blood from healthy individuals with and
625 without globin depletion. *PLoS One* 9, 1–11. doi:10.1371/journal.pone.0091041

626 Smith, J. E. (1825). English Flora. III. Flora.

627 Solano, R., Stepanova, A., Chao, Q., and Ecker, J. R. (1998). Nuclear events in ethylene signaling: a
628 transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-
629 RESPONSE-FACTOR1. *Genes Dev.* 12, 3703–14. doi:10.1101/gad.12.23.3703

630 Soltis, D. E., Soltis, P. S., Pires, J. C., Kovarik, A., Tate, J. A., and Mavrodiev, E. (2004). Recent and
631 recurrent polyploidy in *Tragopogon* (Asteraceae): cytogenetic, genomic and genetic
632 comparisons. *Biol. J. Linn. Soc.* 82, 485–501. doi:10.1111/j.1095-8312.2004.00335.x

633 Soltis, P. S., and Soltis, D. E. (2009). The role of hybridization in plant speciation. *Annu. Rev. Plant*
634 *Biol.* 60, 561–588. doi:10.1146/annurev.arplant.043008.092039

635 Tayalé, A., and Parisod, C. (2013). Natural pathways to polyploidy in plants and consequences for
636 genome reorganization. *Cytogenet. Genome Res.* 140, 79–96. doi:10.1159/000351318

637 Urbanska-Worytkiewicz, K., and Landolt, E. (1974a). Biosystematic investigations in *Cardamine*
638 *pratensis* L.s.I. I. Diploid taxa from Central Europe and their fertility relationships. *Berichte des*
639 *Geobot. Instituts der ETH Stift. Rübel* 42, 42–139

640 Urbanska-Worytkiewicz, K., and Landolt, E. (1974b). Hybridation naturelle entre *Cardamine*
641 *rivularis* SCHUR et *C. amara* L., ses aspects cytologiques et écologiques. *Act. Soc. Helv. Sci.*
642 *Nat.*, 89–90

Urbanska, K. M. (1977). Reproduction in natural triploid hybrids (2n=24) between *Cardamine*
rivularis Schur and *C. amara* L. *er. Geobot. Inst. ETH Stift. Rubel* 44, 42–85

Urbanska, K. M., Hurka, H., Landolt, E., Neuffer, B., and Mummenhoff, K. (1997). Hybridization
and evolution in *Cardamine* (*Brassicaceae*) at Urnerboden, Central Switzerland: Biosystematic
and molecular evidence. *Plant Syst. Evol.* 204, 233–256. doi:10.1007/BF00989208

Urbaska-Worytkiewicz, K., and Landolt, E. (1972). Natürliche Bastarde zwischen *Cardamine amara*
L . und *C* . *rivularis* Schur aus den Schweizer Alpen. *Ber. Geobot. Inst. ETH. Stift. Rübel* 41,
88–101. doi:10.5169/seals-377675

van Wesemael, J., Hueber, Y., Kissel, E., Campos, N., Swennen, R., and Carpentier, S. (2018).
Homeolog expression analysis in an allotriploid non-model crop via integration of
transcriptomics and proteomics. *Sci. Rep.* 8, 1353. doi:10.1038/s41598-018-19684-5

Vollbrecht, E., Veit, B., Sinha, N., and Hake, S. (1991). The developmental gene *Knotted-1* is a
member of a maize homeobox gene family. *Nature* 350, 241–243. doi:10.1038/350241a0

Williams, R. W. (1998). Plant homeobox genes: many functions stem from a common motif.
BioEssays 20, 280–282. doi:10.1002/(SICI)1521-1878(199804)20:4<280::AID-
BIES2>3.0.CO;2-U

Yamaguchi, R., Imoto, S., Yamauchi, M., Nagasaki, M., Yoshida, R., Shimamura, T., et al. (2008).
Predicting differences in gene regulatory systems by state space models. *Genome Inform.* 21,
101–113. doi:10.11234/gi1990.21.101

Yoo, M.-J., Szadkowski, E., and Wendel, J. F. (2013). Homoeolog expression bias and expression
level dominance in allopolyploid cotton. *Heredity (Edinb).* 110, 171–180.
doi:10.1038/hdy.2012.94

Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. *Omi. A J. Integr. Biol.* 16, 284–287.
doi:10.1089/omi.2011.0118

Zhao, J., Yang, F., Feng, J., Wang, Y., Lachenbruch, B., Wang, J., et al. (2017). Genome-wide
constitutively expressed gene analysis and new reference gene selection based on transcriptome
data: a case study from poplar/canker disease interaction. *Front. Plant Sci.* 8, 1–13.
doi:10.3389/fpls.2017.01876

Zozomová-Lihová, J., Krak, K., Mandáková, T., Shimizu, K. K., Španiel, S., Vít, P., et al. (2014).
Multiple hybridization events in *Cardamine* (Brassicaceae) during the last 150 years: revisiting
a textbook example of neoallopolyploidy. *Ann. Bot.* 113, 817–830. doi:10.1093/aob/mcu012

11   Table

Table 1 The negative log10(q-value) of the enriched GO categories in each VEH set described in the manuscript. This data is the extract from the list of all enriched GOs (Dataset S4). ND means that the category was not detected as enriched by the threshold FDR = 0.1.

Keyword	Accession	VEH set				GO Name
		A	I_A	I_R	R	
Alcohol	GO:0006066	6.9	5.2	2.7	2.0	alcohol metabolic process
	GO:0046165	6.4	6.4	4.4	2.7	alcohol biosynthetic process
Water	GO:0009414	3.9	3.3	5.1	7.3	response to water deprivation
Ethylene	GO:0009723	7.9	3.6	2.8	7.5	response to ethylene
	GO:0009873	6.1	1.5	ND	2.5	ethylene-activated signaling pathway
	GO:0071369	6.1	1.3	ND	2.3	cellular response to ethylene stimulus
	GO:0009692	2.6	ND	ND	ND	ethylene metabolic process
	GO:0009693	2.6	ND	ND	ND	ethylene biosynthetic process
Abscisic acid	GO:0009738	1.4	ND	ND	ND	abscisic acid-activated signaling pathway
	GO:0071215	1.4	ND	ND	ND	cellular response to abscisic acid stimulus
Oxidative stress	GO:0006979	1.9	ND	ND	2.6	response to oxidative stress
	GO:0042743	ND	ND	ND	3.0	hydrogen peroxide metabolic process
	GO:2000377	ND	1.6	1.5	5.6	regulation of reactive oxygen species metabolic process
	GO:0010310	ND	1.3	1.1	5.1	regulation of hydrogen peroxide metabolic process
Meristem	GO:0035266	ND	3.1	1.9	ND	meristem growth
	GO:0010075	ND	3.4	2.0	ND	regulation of meristem growth
	GO:0048509	ND	2.8	1.8	ND	regulation of meristem development

12 Supplementary Material

Supplementary_Material.pdf (Table S1, Fig. S1 to Fig. S8)

SI dataset 1 Videos for visualizing of plantlet initiation of *C. insueta* Video file (MP4) visualizes the
plantlet initiation of *C. insueta* at the incubator under the standard condition with the 16 hr light and
8 hr dark. Leaflet were detached form *C. insueta* individuals and floated on the water in a beaker.
Photos were taken between May 16, 2018 and June 7, 2018, with 90 minutes intervals. Photos taken
at daylight were concatenated into video.

SI dataset 2 Statistics of RNA-Seq data processing An Excel format file that contains the statistics of
RNA-Seq data processing with HomeoRoq pipeline. Sheet 1: Number of read pairs before and after
quality controls with Trimmomatic; Sheet 2: Number of reads that were mapped onto A-genome and
R-genome, and number of reads that were classified into A-origin, R-origin, and unclassified reads
with HomeoRoq.

SI dataset 3 FPKM and CV of variably expressed homeologs An Excel format file that contains gene

names, averages of log10-transformed FPKM, and coefficient of variation (CV) of log10-

transformed FPKM of variably expressed homeologs (VEHs). Sheet 1: VEHs of IA samples; Sheet 2:

VEHs of IR samples; Sheet 3: VEHs of C. amara samples; Sheet 4: VEHs of C. rivularis samples.

SI dataset 4 Enriched GO terms of variably expressed homeologs An Excel format file that contains

gene ontology (GO) enrichment analysis results of VEHs. Sheet 1: Summarization of GO enrichment

analysis results. Values in cells represent negative log10(q-value); Sheet IA: GO enrichment analysis

result of VEHs of I_A samples; Sheet IR: GO enrichment analysis result of VEHs of I_R samples; Sheet

A: GO enrichment analysis result of VEHs of *C. amara* samples; Sheet R: GO enrichment analysis

result of VEHs of *C. rivularis* samples.

SI dataset 5 Relative gene expression levels of the genes of the category GO:0009414 (response to

water deprivation).

13   Data Availability Statement

The datasets generated for this study can be found in DNA Data Bank of Japan (DDBJ) Sequence

Read Archive (DRA), www.ddbj.nig.ac.jp [accession no. PRJDB9426].

14   Figure Legends

Figure 1 Comparison of homeolog expression from A- and R-subgenomes. **A** Expression ratio

between A- and R-homeologs before submergence treatment in the triploid *C. insueta*. Each dot

shows the relation between the log10-transformed A-origin and R-origin read of a homeolog pair at

0hr point. Only the homeolog pairs with FPKM > 1.0 in either I_A or I_R samples are shown. The red

line represents the ratio A:R=1:1, and the orange line represents the ratio A:R=1:2. **B** Comparison of

A-origin ratios between two time points, 0 hr and 2 hr, in the triploid *C. insueta*. Each point shows

the A-origin ratios of a homeolog pair at 0 hr and 2 hr. The orange lines represent the position of A-

origin=0.33 at each time point.

Figure 2 Principal component analysis of the expressed homeologs/genes in *C. insueta* (I_A), *C.*

insueta (I_R), *C. amara* (A) and *C. rivularis* (R) samples at 9 time points. PCA was performed against

log10-transformed FPKM of 21,131 expressed homeologs. The two plots show the relation between

PC1-PC2 (**A**) and PC2-PC3 (**B**). The colors represent genome/subgenome, and the numbers represent

the time points after the start of submergence treatment.

Figure 3 A schematic drawing of *cis*- and *trans*-regulation of key regulatory genes in meristem

formation. The expression patterns of I$_A$ homeologs are arbitrary categorized to *cis*- or *trans*-

regulated according to the expression pattern of each homeolog. See Figure S8 for their original

temporal expression patterns.