Title

Ultra-structural analysis and morphological changes during the differentiation of

trophozoite to cyst in *Entamoeba invadens*

Nishant Singh[1*†], Sarah Naiyer[†2] and Sudha Bhattacharya[3]

School of Environmental Sciences, Jawaharlal Nehru University, New Delhi, India-110067

2- Present Address

Dr. Sarah Naiyer

Department of Immunology and Microbiology

University of Illinois at Chicago, USA

Email: snaiyer@uic.edu, sara.naiyer@gmail.com

3- Present Address

Prof. Sudha Bhattacharya

Ashoka University, Rajiv Gandhi Education City, Sonipat, Haryana, India -131029

Email: sbjnu110@gmail.com

*-Corresponding author

1- Present Address

Dr. Nishant Singh

Department of Urology, University of Pittsburgh, Pittsburgh, PA-USA 15213

Email: nishant900@gmail.com, singhn5@upmc.edu

[†-] Share equal first author

Abstract

Entamoeba Histolytica, a pathogenic parasite, is the causative organism of amoebiasis and uses

human colon to complete its life cycle. It destroys intestinal tissue leading to invasive disease.

Since it does not form cyst in culture medium, a reptilian parasite *Entamoeba invadens* serves as

the model system to study encystation. Detailed investigation on the mechanism of cyst

formation, information on ultra-structural changes and cyst wall formation during encystation are

still lacking in *E. invadens*. Here, we used electron microscopy to study the ultrastructural

changes during cyst formation and showed that the increase in heterochromatin patches and

deformation of nuclear shape were early events in encystation. These changes peaked at ~20h

post induction, and normal nuclear morphology was restored by 72h. Two types of cellular

structures were visible by 16h. One was densely stained and consisted of the cytoplasmic mass

with clearly visible nucleus. The other consisted of membranous shells with large vacuoles and

scant cytoplasm. The former structure developed into the mature cyst while the latter structure

was lost after 20h, This study of ultra-structural changes during encystation in *E. invadens* opens

up the possibilities for further investigation into the mechanisms involved in this novel process.

Keywords: *Entamoeba invadens*; *Entamoeba histolytica*; nucleus; electron microscopy;

encystation; excystation; hetero chromatin.

Introduction

Entamoeba histolytica, a parasitic protist which causes amoebiasis in humans, is distributed

worldwide with greater prevalence in developing countries (Debnath et al., 2012). It has simple

life cycle consisting of dormant tetra-nucleate cyst which under favorable conditions converts

into the actively dividing uninucleate trophozoite. These differentiation stages can be reproduced

in the lab conditions in the reptilian parasite *E. invadens* since encystation has not yet been

achieved in axenically grown *E. histolytica* trophozoites (Sanchez et al., 1994). In encystation

process of *E. invadens* motile trophozoite rounds up and eventually becomes encapsulated within

a rigid, relatively impermeable cell wall (Chavez et al., 1978). The cyst wall components are

concentrated and transported in specific encystation specific vesicles (ESV) which contain a

fibrillar material (Chavez-Munguia et al., 2003). This material is digested on the plasma

membrane through the rupture of the vesicles. Similar type of ESVs are also found in

Acanthamoeba castellanii (Chavez-Munguia et al., 2005) and *Giardia* (Reiner et al., 1990). This

morphological evidence supports that the vesicles found in the amoeba and those identified in the

flagellated *G. lamblia* may correspond to a common biosynthetic mechanism of synthesis,

transport and deposition of cell wall components (Chavez-Munguia et al., 2008). Literature

indicates that *Entamoeba* undergoes sexual or parasexual reproduction at some stage but timing

is not known (Ali et al., 2008, Ramesh et al., 2005, Weedall et al., 2012, Weedall and Hall,

2015). Meiosis specific genes were found upregulated and homologous recombination was

observed during stage conversion or under stress conditions (Singh et al., 2013).

During encystation, nuclear division takes place in the absence of cytokinesis, such that the uni-

nucleated trophozoite transforms into a dormant tetra-nucleated cyst, enveloped by a protective

cyst wall. Detailed studies on the mechanism of cyst formation, information on ultra-structural

changes during encystation are lacking in *E. invadens*. In this study, we attempt to address the

mechanism of cyst formation by transmission electron microscopy (TEM) and fluorescence

microscopy.

Results

Nuclear dynamics during encystation

Nuclear shape and heterochromatin (HC) distribution of *E. invadens* was investigated by

transmission electron microscopy during encystation. In normal proliferating trophozoite,

condensed HC masses are arranged directly underneath the nuclear membrane and the nuclear

shape is spherical (Fig. 1) as reported previously in *E. histolytica* (Chavez-Munguia et al., 2008,

Dahl et al., 2008, Solis and Barrios, 1991). HC is densely packed chromatin, which usually

reflects modifications of DNA, histones and other DNA binding proteins, and is typically

transcriptionally inactive (John, 1988). In *Entamoeba*, peripheral HC contains the ribosomal

DNA circles and stains with nucleolar markers like fibrillarin (Jhingan et al., 2009). We looked

for changes in nuclear distribution of HC during encystation and found that large patches of HC

began to appear as early as 4h after induction of cyst formation, and by 20h the nuclei in 90%

cells had these patches of HC (Fig. 1, 3). At 24h and 48h, HC patches were seen not only at the

periphery but at internal locations as well. The nuclear shape began to change from spherical to

distorted within 2h of encystation, and by 16h, ~95% cells had distorted nuclei. By 20h most

cells had nuclei with HC patches and distorted shape (Fig. 3), after which there was a gradual

increase in nuclei with the normal pattern. In mature cysts (72h) the normal pattern of HC and

nuclear shape was restored (Fig. 1). Chromatoid bodies composed of poly ribosomes appeared in

the cytoplasm after 16h.

Morphological changes during cyst formation

Sequential ultra-structural changes occurring in the trophozoite during encystation were

investigated through TEM. Trophozoites were inoculated in encystation medium and samples

were fixed after each time point. Thin sections were examined for morphological changes

between 8h to 16h (Fig. 2). In high resolution electron microscopy at zero time point,

trophozoites showed normal circular nucleus and no chromatoid bodies. Plasma membrane was

in uniform contact with cytoplasm (Fig. 2, 0h). Remarkable changes were visible after 8h. In a

number of cells the cytoplasmic mass could be seen separated from the plasma membrane (Fig.

2A). Gradually the separation between plasma membrane and cytoplasmic material increased,

and in some cells the cytoplasmic mass was almost completely detached from the membrane

(Fig. 2, A-C). By 16h, ~60% cells showed a near complete separation of the plasma membrane

and cytoplasm (Fig. 3). Two types of structures were visible. One was densely stained and

consisted of the cytoplasmic mass with clearly visible nucleus. The other consisted of

membranous shells with large vacuoles and scant cytoplasm (Fig. 2, D-E). After 20h the latter

were lost and the former developed into mature cyst, with the spherical nuclei being restored at

72h.

We also observed the encysting cells in real time to further understand how the above mentioned

phenomenon took place. Cells were inoculated in encystation medium and monitored under

confocal microscope between 8h to 12h. It is well documented that cells which are getting ready

to encyst undergo clump formation and clumping is a prerequisite for encystation. Thus it was

not possible to focus on well-separated cells and follow them in real time. In addition, there was

rapid pseudopodial movement and constant movement of neighboring cells in and out of the

viewing field. Due to these reasons it was difficult to observe a single encysting cell in a

sequential manner over an extended time period. Our observations of cells 8-12 h post-induction

(Fig. 4) show that cells (marked by arrow in Fig. 4A) in a clump of trophozoites could be seen

rounded and with a clear zone in the center, with some cytoplasmic material on the periphery.

With time, structures became visible which had large vacuoles with little cytoplasmic material

and looked like hollow barrels or cups (marked by * in Fig. 4 F, G, J, K) symbol marked

(Mam).(marked by O in Fig. 4 B, C, D, E, G, H, I, J, K). Simultaneously, one could also see the

accumulation of round cytoplasmic masses (with almost the same diameter as the opening of the

barrel/cup), these are marked by ▲ in Fig. 4 B, C, D, F, H-N. Whether the cytoplasmic spheres

are extruded from the vacuolated cell or the two are independent entities needs to be resolved. At

later times (panels M-P) very few original trophozoites are visible, and the structures seen are

either the transparent vacuolated shells or the dense cytoplasmic spheres.

That the vacuolated shells were indeed plasma membrane bound was shown with a plasma

membrane specific dye ("cell mask" red cat no. C10046 from Invitrogen) which specifically

binds to plasma membrane. Membrane staining was also done with peroxiredoxin antibody

which has been shown to localize preferentially to the plasma membrane in *Entamoeba* (Choi et

al., 2005). In normal trophozoite the membrane specific reagents labeled the membrane, with

some internal diffusion (Fig. 5). The 16h sample showed very clear labeling of the vacuolated

shell with "cell mask" dye and the absence of nucleus was confirmed by lack of Hoechst

staining. The dense cytoplasmic spheres were uni-nucleate. Peroxiredoxin antibody also showed

the same pattern in the vacuolated shell, which lacked any nucleus.

Discussion

Our EM study showed that the nucleus of *E. invadens* in normal proliferative cells was typically

spherical or ellipsoid which was identical to *E. histolytica* (Chavez-Munguia et al., 2008, Solis

and Barrios, 1991). Heterochromatin was present at the periphery, which is also the site of

nucleolar localization (Jhingan et al., 2009). In this respect also *E. invadens* is identical with *E.*

histolytica. HC is generally associated with low activity of gene expression, and many specific

proteins and characteristic histone modifications present in heterochromatin are responsible for

gene silencing (Dahl et al., 2008, Sarma and Reinberg, 2005). Conversely, euchromatin is rich in

transcriptionally active genes and is often located at the nuclear interior in more open chromatin

structures. In *E. invadens*, the nuclear shape changed and larger patches of HC were visible

during encystation (Fig. 1). Our results showed that the increase in HC patches was an early

event during encystation. It could be related to up- or down regulation of stage specific genes

which takes place in cells committed for encystation (Ehrenkaufer et al., 2013, Manna et al.,

2014).

It has been observed that open euchromatin structures are more deformable than tightly packed

heterochromatin structures (Pajerowski et al., 2007), and any external or intracellular forces

could reorganize gene rich areas relatively easily. Even though the nucleus is the stiffest cellular

organelle and is 2- to 10-times stiffer than the surrounding cytoskeleton, (Caille et al., 2002,

Guilak et al., 2000), extracellular forces and strain result in clearly detectable nuclear

deformations (Guilak, 1995, Guilak et al., 2000, Maniotis et al., 1997). An abnormal nuclear

shape is frequently associated with cancer (Webster et al., 2009, Zink et al., 2004). It has been

speculated that change in nuclear shape leads to change in chromosome organization which in

turn can alter gene expression (He et al., 2008). Altered nuclear shape in cancer cells is thought

to facilitate metastasis because of reduced nuclear stiffness, which could increase the ability of

transformed cells to penetrate tissue (Dahl et al., 2008). Inactivation of some proteins which

associate with endoplasmic reticulum can affect nuclear shape (Higashio et al., 2000, Matynia et

al., 2002). Nuclear shape can also be affected by lipid synthesis. This has been shown in yeast

and *Caenorhabditis elegans*, where the inactivation of a lipid phosphatase that is homologous to

the mammalian lipin (Reue and Zhang, 2008) was shown to cause expansion of the ER

membrane and alteration in nuclear shape (Campbell et al., 2006, Golden et al., 2009).

Changes in nuclear structure and function could contribute both to increased cellular sensitivity

to stress and to altered transcriptional regulation (Lammerding et al., 2004). Some specialized

cells undergo dramatic changes in nuclear shape during differentiation and maturation. For

example, spermatids have extremely elongated nuclei (Dadoune, 2003) and neutrophils develop

extremely lobulated nuclei, which are associated with loss of lamin A/C (Yabuki et al., 1999)

and expression of lamin B receptor (Hoffmann et al., 2007). Thus changes in nuclear shape are

well-documented with respect to cellular differentiation, stress and disease. We speculate that

adaptations in nuclear shape and structure are directly related to the functionality of the *E.*

invadens cell and in the differentiation process.

In this study, we also documented the detailed observation of ultra-structural changes during

encystation in *E. invadens*. Encystation is similar to the well-known process of sporulation in

yeast and bacteria in that both are triggered by adverse growth conditions, and result in

production of a resting cell that is more resistant to environmental factors than the vegetative

cell. In *S. cerevisiae* during sporulation, a meiotic division takes place, resulting in the

production of four haploid nuclei which are then enveloped by de novo formed plasma

membranes within the cytoplasm of the mother cell to form immature spores or prospores

(Neiman, 2011). In *Bacillus subtilis*, a single spore (fore spore) is formed by a process of

engulfment following an asymmetrical division within the mother cell. Upon completion of

spore formation the mother cell lyses, releasing the mature spore (Higgins and Dworkin, 2012).

The major difference between the trophozoite and cyst in *E. invadens* is that the latter is about

half the average diameter of the former, and while the trophozoite is uni-nucleate, the cyst is tetra

nucleate. In addition, the cyst possesses a chitin wall. Since one trophozoite gives rise to only

one cyst it was assumed that once the differentiation process is set in motion in *E. invadens*, the

process would be relatively straightforward. Our TEM and confocal observations reveal a more

complex picture. We find that once the initial signal for differentiation is conveyed to the

trophozoite, the cytoplasmic mass begins to move away from the original plasma membrane. At

this stage the spherical cytoplasmic mass may be referred to as a pre-cyst and the chitin wall is

not yet visible by calcofluor staining. In addition, it contains only one nucleus. A second class of

structures is visible which consists of large vacuoles bound by plasma membrane and containing

little cytoplasm. Whether the pre-cyst is extruded as a spherical mass, leaving behind this empty

shell, or the two structures are unrelated needs to be understood. It is possible that the vacuolated

structures represent cells that have undergone apoptosis and would not contribute to cyst

formation. An altruistic possibility may also exist whereby the death of these cells could benefit

other cells to complete the differentiation process (Picazarri et al., 2008). In a study of

autophagosome-like structures in encysting *E. invadens* cells it was shown that the autophagy

gene, Atg8 was associated with structures that looked like vesicles/vacuoles as early as 2 h after

induction of encystation. Some of these had diameter >4 µm, and they peaked at 24 h when

>65% of cells contained these structures (Picazarri et al., 2008). It is possible that these could be

the precursors of large vacuolated cells reported in our study.

Our data show that *E. invadens* has evolved a unique process of cyst formation, the details of

which are not clearly understood. This work opens up the possibilities for further investigation

into the mechanisms involved, which might help to understand the significance of this novel

process.

Materials and Methods

Strains and cell culture

All experiments were carried out with *E.invadens* strain IP-1 was obtained from the American

Type Culture Collection and maintained at 25°C in TYI-S-33 containing 15% heat inactivated

adult bovine serum,125ml/100 mlstreptomycin/penicillin G and 2.0% vitamin mix (Singh et al.,

2012).

Cyst induction and excystation

This was done essentially as described and adapted in our lab (Singh et al., 2011). Briefly, log

phase trophozoites grown in 50 ml flasks were chilled on ice for 10 min to remove the cells from

the wall and harvested by centrifugation at 500g for 5 min at 4°C. $5x10^5$ trophozoites ml^{-1} were

transferred into induction medium (LG): TYI medium was prepared without glucose and diluted

to 2.12 times with water and completed with 5% heat inactivated adult bovine serum, 2.6%

vitamin mix and 125 mL/100 ml antibiotic. Cysts obtained in LG medium after three days were

harvested and treated with 0.05% Sarkosyl (Sigma) to destroy the trophozoites. Typical spherical

refractile cysts were observed by light microscopy and checked for staining with calcofluor and

observed 80–90% encystation efficiency. For excystation, the cysts were further washed with

phosphate-buffered saline (PBS), and inoculated in normal TYI-S-33 medium.

Electron microscopy

To examine cellular and nuclear morphology during encystation, *E. invadens* cells were grown

till late log phase and 5×10^5 cell/ml was transfer into LG medium for cyst induction for

different time points and fixed in 2.5% glutaraldehyde2.0% paraformaldehyde in 0.1M PBS, pH

7.2 for 2h at RT and overnight at 4°C. Washed thrice with 0.1M PBS. Specimens were post fixed

in 1% osmium tetroxide and 0.1M PBS, for 2h at 4°C, washed with 0.1M PBS. The specimens

were dehydrated with increasing concentrations of ethanol and then clearing for 2h in toluene at

RT, followed by infiltrate in a 1:1 mixture of toluene and Epoxy resin-Araldite (M) EY212

(TAAB laboratory equipment ltd.) overnight. Specimens were subsequently embedded in Epoxy

resin at 60ºC for 2 days. Semi- and ultra-thin sections were obtained using an ultra-microtome

Leica UC-6. Semi-thin sections (1µ) were dried on slides at 80ºC and stained with a mixture of 1.0% methylene blue. Ultra-thin section (70-90nm) on 200 mess Copper grids stained with saturated solution of uranyl acetate in 50% alcohol for 15 min followed by lead citrate (50 mg/12ml MQ) for 10 min at RT. Sections were scanned and photographs were taken with a JEOL- JEM-2100F transmission electron microscope at 120 kV and scanned images were processed using Adobe Photoshop software. Sample preparation and viewing was done at Advanced Instrumentation Research Facility (AIRF), Jawaharlal Nehru University, New Delhi, India.

Immuno-fluorescence microscopy

Cells were washed with PBS, fixed in cold methanol for 20 min at -20ºC, again washed twice with PBS buffer and permeabilized with 0.1% triton-X-100 for 5 min at room temperature, washed and blocked with 3% BSA at RT for 60 min. After blocking cells were washed once with PBS and incubated for 45 min at RT in monoclonal/polyclonal antibody generated in mouse/rabbit in suitable dilutions, washed with PBS and incubated for 30 min with anti-mouse/rabbit secondary antibody tagged with fluorochrome in suitable dilutions along with nuclear strain Hoechst 33342 (2.5 µg/ml). For positive control, polyclonal anti- rabbit/mouse anti- fibrillarin antibody (Jhingan et al., 2009) was used in each preparation in 1:500 dilutions. Trophozoites were then washed thrice with PBS and mounted on glass slide in 25% glycerol with anti-fed p-phenylenediamine (sigma). All steps were done in 1.5ml microfuge tubes, and centrifuged at 4ºC; 500g for 5 min. Phase contrast and fluorescent images were taken using a Zeiss Axio Imager.M1 microscope (Germany) with a 100x magnification

Acknowledgement

NS and SN were recipient of fellowship from CSIR, India. SB is a recipient of J.C. Bose

fellowship. We thank Prof. Alok Bhattacharya, School of Life Sciences Jawaharlal Nehru

University for his critical inputs and scientific discussions. We also thank Advanced

Instrumentation Research Facility (AIRF) and Dr. Gajender Saini, Jawaharlal Nehru University

for his help during sample preparation and Electron Microscopy.

Figure legends

Figure 1: Transmission electron micrograph showing the changes in the nuclear structure

and heterochromatin distribution in *E. invadens* during encystation. Samples were taken at

different time points i.e. 0h, 8h, 16h, 24h, 48h, and 72h, after transfer to LG medium. HC;

heterochromatin, N; nucleus, CB; chromatoid body.

Figure 2: Formation of cyst within plasma membrane showed by electron microscopy in *E.*

***invadens* during encystation** 0; normal proliferating trophozoite, A; 8h, B, C; 12h, D; 16h, E;

20h, F; 24h, G; 48h, H; 72h. N; Nucleus, PM; Plasma membrane.

Figure 3: Distribution of different cell type during encystation in *E. invadens* investigated

by electron microscopy. Blue; normal nucleus, Brown; distorted nucleus, green;

heterochromatin patches, purple; detached plasma membrane.

Figure 4: Live cell imaging of encysting *E. invadens* trophozoites. Cells were inoculated into

induction medium and viewed by microscopy between 8 to 12h. Arrow indicated the plasma

membrane of particular cell (ghost).

Figure 5: Staining with plasma membrane specific reagents. Cells were taken at 16h during

encystation. Peroxiredoxin antibody was kind gift from Sharon Reed (Choi et al., 2005) which

localized to plasma membrane of *E. invadens.* Green; anti mouse-peroxiredoxin primary antibody (1:250) and anti-mouse-secondary antibody tagged with alexa-488 (1:500), red; plasma membrane specific red dye "cell mask" cat no. C10046 from Invitrogen. Hoechst 33342 was used as nuclear stain.

References

ALI, I. K., SOLAYMANI-MOHAMMADI, S., AKHTER, J., ROY, S., GORRINI, C., CALDERARO, A., PARKER, S. K., HAQUE, R., PETRI, W. A., JR. & CLARK, C. G. 2008. Tissue invasion by Entamoeba histolytica: evidence of genetic selection and/or DNA reorganization events in organ tropism. *PLoS Negl Trop Dis,* 2**,** e219.

CAILLE, N., THOUMINE, O., TARDY, Y. & MEISTER, J. J. 2002. Contribution of the nucleus to the mechanical properties of endothelial cells. *J Biomech,* 35**,** 177-87.

CAMPBELL, J. L., LORENZ, A., WITKIN, K. L., HAYS, T., LOIDL, J. & COHEN-FIX, O. 2006. Yeast nuclear envelope subdomains with distinct abilities to resist membrane expansion. *Mol Biol Cell,* 17**,** 1768-78.

CHAVEZ, B., MARTINEZ-PALOMO, A. & DE LA TORRE, M. 1978. [Ultramicroscopic structure of the cyst wall of Entamoeba invadens, E. histolytica and E. coli]. *Arch Invest Med (Mex),* 9 Suppl 1**,** 113-6.

CHAVEZ-MUNGUIA, B., CRISTOBAL-RAMOS, A. R., GONZALEZ-ROBLES, A., TSUTSUMI, V. & MARTINEZ-PALOMO, A. 2003. Ultrastructural study of Entamoeba invadens encystation and excystation. *J Submicrosc Cytol Pathol,* 35**,** 235-43.

CHAVEZ-MUNGUIA, B., OMANA-MOLINA, M., GONZALEZ-LAZARO, M., GONZALEZ-ROBLES, A., BONILLA, P. & MARTINEZ-PALOMO, A. 2005. Ultrastructural study

of encystation and excystation in Acanthamoeba castellanii. *J Eukaryot Microbiol,* 52**,** 153-8.

CHAVEZ-MUNGUIA, B., TALAMAS-ROHANA, P., RIOS, A., GONZALEZ-LAZARO, M. & MARTINEZ-PALOMO, A. 2008. Entamoeba histolytica: fibrilar aggregates in dividing trophozoites. *Exp Parasitol,* 118**,** 280-4.

CHOI, M. H., SAJED, D., POOLE, L., HIRATA, K., HERDMAN, S., TORIAN, B. E. & REED, S. L. 2005. An unusual surface peroxiredoxin protects invasive Entamoeba histolytica from oxidant attack. *Mol Biochem Parasitol,* 143**,** 80-9.

DADOUNE, J. P. 2003. Expression of mammalian spermatozoal nucleoproteins. *Microsc Res Tech,* 61**,** 56-75.

DAHL, K. N., RIBEIRO, A. J. & LAMMERDING, J. 2008. Nuclear shape, mechanics, and mechanotransduction. *Circ Res,* 102**,** 1307-18.

DEBNATH, A., PARSONAGE, D., ANDRADE, R. M., HE, C., COBO, E. R., HIRATA, K., CHEN, S., GARCIA-RIVERA, G., OROZCO, E., MARTINEZ, M. B., GUNATILLEKE, S. S., BARRIOS, A. M., ARKIN, M. R., POOLE, L. B., MCKERROW, J. H. & REED, S. L. 2012. A high-throughput drug screen for Entamoeba histolytica identifies a new lead and target. *Nat Med,* 18**,** 956-60.

EHRENKAUFER, G. M., WEEDALL, G. D., WILLIAMS, D., LORENZI, H. A., CALER, E., HALL, N. & SINGH, U. 2013. The genome and transcriptome of the enteric parasite Entamoeba invadens, a model for encystation. *Genome Biol,* 14**,** R77.

GOLDEN, A., LIU, J. & COHEN-FIX, O. 2009. Inactivation of the C. elegans lipin homolog leads to ER disorganization and to defects in the breakdown and reassembly of the nuclear envelope. *J Cell Sci,* 122**,** 1970-8.

GUILAK, F. 1995. Compression-induced changes in the shape and volume of the chondrocyte
nucleus. *J Biomech,* 28**,** 1529-41.

GUILAK, F., TEDROW, J. R. & BURGKART, R. 2000. Viscoelastic properties of the cell
nucleus. *Biochem Biophys Res Commun,* 269**,** 781-6.

HE, S., DUNN, K. L., ESPINO, P. S., DROBIC, B., LI, L., YU, J., SUN, J. M., CHEN, H. Y.,
PRITCHARD, S. & DAVIE, J. R. 2008. Chromatin organization and nuclear
microenvironments in cancer cells. *J Cell Biochem,* 104**,** 2004-15.

HIGASHIO, H., KIMATA, Y., KIRIYAMA, T., HIRATA, A. & KOHNO, K. 2000. Sfb2p, a
yeast protein related to Sec24p, can function as a constituent of COPII coats required for
vesicle budding from the endoplasmic reticulum. *J Biol Chem,* 275**,** 17900-8.

HIGGINS, D. & DWORKIN, J. 2012. Recent progress in Bacillus subtilis sporulation. *FEMS
Microbiol Rev,* 36**,** 131-48.

HOFFMANN, K., SPERLING, K., OLINS, A. L. & OLINS, D. E. 2007. The granulocyte
nucleus and lamin B receptor: avoiding the ovoid. *Chromosoma,* 116**,** 227-35.

JHINGAN, G. D., PANIGRAHI, S. K., BHATTACHARYA, A. & BHATTACHARYA, S.
2009. The nucleolus in Entamoeba histolytica and Entamoeba invadens is located at the
nuclear periphery. *Mol Biochem Parasitol,* 167**,** 72-80.

JOHN, B. 1988. The biology of heterochromatin. In: Heterochromatin: Molecular and Structural
Aspects, Verma, R. S. (ed.) *Cambridge University Press, New York.*

LAMMERDING, J., SCHULZE, P. C., TAKAHASHI, T., KOZLOV, S., SULLIVAN, T.,
KAMM, R. D., STEWART, C. L. & LEE, R. T. 2004. Lamin A/C deficiency causes
defective nuclear mechanics and mechanotransduction. *J Clin Invest,* 113**,** 370-8.

MANIOTIS, A. J., CHEN, C. S. & INGBER, D. E. 1997. Demonstration of mechanical

connections between integrins, cytoskeletal filaments, and nucleoplasm that stabilize

nuclear structure. *Proc Natl Acad Sci U S A,* 94**,** 849-54.

MANNA, D., EHRENKAUFER, G. M. & SINGH, U. 2014. Regulation of gene expression in

the protozoan parasite Entamoeba invadens: identification of core promoter elements and

promoters with stage-specific expression patterns. *Int J Parasitol,* 44**,** 837-45.

MATYNIA, A., SALUS, S. S. & SAZER, S. 2002. Three proteins required for early steps in the

protein secretory pathway also affect nuclear envelope structure and cell cycle

progression in fission yeast. *J Cell Sci,* 115**,** 421-31.

NEIMAN, A. M. 2011. Sporulation in the budding yeast Saccharomyces cerevisiae. *Genetics,*

189**,** 737-65.

PAJEROWSKI, J. D., DAHL, K. N., ZHONG, F. L., SAMMAK, P. J. & DISCHER, D. E. 2007.

Physical plasticity of the nucleus in stem cell differentiation. *Proc Natl Acad Sci U S A,*

104**,** 15619-24.

PICAZARRI, K., NAKADA-TSUKUI, K. & NOZAKI, T. 2008. Autophagy during proliferation

and encystation in the protozoan parasite Entamoeba invadens. *Infect Immun,* 76**,** 278-88.

RAMESH, M. A., MALIK, S. B. & LOGSDON, J. M., JR. 2005. A phylogenomic inventory of

meiotic genes; evidence for sex in Giardia and an early eukaryotic origin of meiosis. *Curr*

Biol, 15**,** 185-91.

REINER, D. S., MCCAFFERY, M. & GILLIN, F. D. 1990. Sorting of cyst wall proteins to a

regulated secretory pathway during differentiation of the primitive eukaryote, Giardia

lamblia. *Eur J Cell Biol,* 53**,** 142-53.

REUE, K. & ZHANG, P. 2008. The lipin protein family: dual roles in lipid biosynthesis and
367   gene expression. *FEBS Lett,* 582**,** 90-6.

368 SANCHEZ, L., ENEA, V. & EICHINGER, D. 1994. Identification of a developmentally
369   regulated transcript expressed during encystation of Entamoeba invadens. *Mol Biochem*
370   *Parasitol,* 67**,** 125-35.

371 SARMA, K. & REINBERG, D. 2005. Histone variants meet their match. *Nat Rev Mol Cell Biol,*
372   6**,** 139-49.

373 SINGH, N., BHATTACHARYA, A. & BHATTACHARYA, S. 2013. Homologous
374   recombination occurs in Entamoeba and is enhanced during growth stress and stage
375   conversion. *PLoS One,* 8**,** e74465.

376 SINGH, N., BHATTACHARYA, S. & PAUL, J. 2011. Entamoeba invadens: dynamics of DNA
377   synthesis during differentiation from trophozoite to cyst. *Exp Parasitol,* 127**,** 329-33.

378 SINGH, N., OJHA, S., BHATTACHARYA, A. & BHATTACHARYA, S. 2012. Stable
379   transfection and continuous expression of heterologous genes in Entamoeba invadens.
380   *Mol Biochem Parasitol,* 184**,** 9-12.

381 SOLIS, F. J. & BARRIOS, R. 1991. Entamoeba histolytica: microtubule movement during
382   mitosis. *Exp Parasitol,* 73**,** 276-84.

383 WEBSTER, M., WITKIN, K. L. & COHEN-FIX, O. 2009. Sizing up the nucleus: nuclear shape,
384   size and nuclear-envelope assembly. *J Cell Sci,* 122**,** 1477-86.

385 WEEDALL, G. D., CLARK, C. G., KOLDKJAER, P., KAY, S., BRUCHHAUS, I., TANNICH,
386   E., PATERSON, S. & HALL, N. 2012. Genomic diversity of the human intestinal
387   parasite Entamoeba histolytica. *Genome Biol,* 13**,** R38.

WEEDALL, G. D. & HALL, N. 2015. Sexual reproduction and genetic exchange in parasitic

protists. *Parasitology,* 142 Suppl 1**,** S120-7.

YABUKI, M., MIYAKE, T., DOI, Y., FUJIWARA, T., HAMAZAKI, K., YOSHIOKA, T.,

HORTON, A. A. & UTSUMI, K. 1999. Role of nuclear lamins in nuclear segmentation

of human neutrophils. *Physiol Chem Phys Med NMR,* 31**,** 77-84.

ZINK, D., FISCHER, A. H. & NICKERSON, J. A. 2004. Nuclear structure in cancer cells. *Nat*

Rev Cancer, 4**,** 677-87.

Bright field **Hoechst** **Membrane specific** **Merge**

Plasma membrane specific dye

Normal

16h Encystation

Peroxiredoxin

Normal

16h Encystation