Clustering of loci controlling species differences in chemical bouquets of sympatric *Heliconius*
butterflies

Kelsey J. R. P. Byers[1,2,*], Kathy Darragh[1,2,a,*], Sylvia Fernanda Garza[2,b], Diana Abondano Almeida[2,c], Ian A.
Warren[1], Pasi M. A. Rastas[3], Richard M. Merrill[2,4], Stefan Schulz[5], W. Owen McMillan[2], Chris D. Jiggins[1,2,6]

[1] *Department of Zoology, University of Cambridge, Cambridge, United Kingdom*

[2] *Smithsonian Tropical Research Institute, Panama, Panama*

[3] *Institute of Biotechnology, University of Helsinki, Helsinki, Finland*

[4] *Division of Evolutionary Biology, Ludwig-Maximilians-Universität München, Munich, Germany*

[5] *Institute of Organic Chemistry, Department of Life Sciences, Technische Universität Braunschweig,*
Braunschweig, Germany

[6] *To whom correspondence should be addressed: University of Cambridge, Department of Zoology,*
Downing Street, Cambridge, CB2 3EJ, United Kingdom. c.jiggins@zoo.cam.ac.uk; Tel: (+44)(0)1223
769021; Fax: (+44)(0)1223 336676

[a] *Present address: Department of Evolution and Ecology, University of California Davis, Davis, California,*
USA

[b] *Present address: Department of Collective Behaviour, Max Planck Institute of Animal Behaviour,*
Konstanz, Germany & Centre for the Advanced Study of Collective Behaviour, University of Konstanz,
Konstanz, Germany

[c] *Present address: Institute for Ecology, Evolution and Diversity, Goethe Universität, Frankfurt, Germany*

* Kelsey J.R.P. Byers and Kathy Darragh should be considered joint first author

ORCID K.J.R.P.B.: 0000-0002-0923-8264
ORCID K.D.: 0000-0002-9215-014X
ORCID R.M.M.: 0000-0003-4527-9298
ORCID W.O.M.: 0000-0003-2805-2745
ORCID C.D.J.: 0000-0002-7809-062X

Keywords: pheromones; speciation; quantitative trait locus mapping; Lepidoptera; *Heliconius*

Running title: Clustering of pheromone loci in *Heliconius*

Author contributions:

KJRPB and KD should be considered joint first author. CDJ conceived and designed the study. KJRPB and
KD performed phenotyping and QTL mapping for wing and genital compounds respectively. KJRPB

performed downstream analyses. KD and RMM designed and performed QTL mapping crosses and KD
and IAW prepared sequencing libraries. KD, SFG, DAA, KJRPB, and RMM reared butterflies for mapping
experiments. SS provided mass spectrometry libraries and advice on compound identification. PMAR
analyzed raw sequencing data, produced phased sequence data, and constructed linkage maps. CDJ and
WOM contributed funding and resources and, with RMM, helped design the QTL mapping experiment.
KJRPB wrote the manuscript, and all authors contributed to the manuscript editing process. All authors
approved the final version of the manuscript.

Acknowledgments:

Permits for research and collection of butterfly stocks were provided by the government of Panama.
KJRPB, KD, IAW, and CDJ were funded by the European Research Council (FP7-IDEAS-ERC 339873); KD
was additionally funded by a Natural Environment Research Council Doctoral Training Partnership (Grant
No. NE/L002507/1) and a Smithsonian Tropical Research Institute Short Term Fellowship; PMAR was
funded by the Jane and Aatos Erkko Foundation; WOM was funded by the Smithsonian Tropical
Research Institute; and SS was funded by the Deutsche Forschungsgemeinschaft (grant DFG
Schu984/13-1).

Data availability:

Data and QTL analysis code have been deposited in Dryad under accession XXX. Sequencing reads
leading to linkage map construction have been deposited in the European Nucleotide Archive (ENA)
under project PRJEB34160.

Competing interest statement:

The authors declare no competing interests.

Abstract

The degree to which loci promoting reproductive isolation cluster in the genome – *i.e.* the genetic

architecture of reproductive isolation - can influence the tempo and mode of speciation. Tight linkage

between these loci can facilitate speciation in the face of gene flow. Pheromones play a role in

reproductive isolation in many Lepidoptera species, and their role as secondary metabolites decreases

the likelihood of pleiotropy associated with many barrier loci. *Heliconius* butterflies use male sex

pheromones to both court females (aphrodisiac wing pheromones) and ward off male courtship (male-

transferred anti-aphrodisiac genital pheromones), and it is likely that these compounds play a role in

reproductive isolation between *Heliconius* species. Using a set of backcross hybrids between *H.*

melpomene and *H. cydno*, we investigated the genetic architecture of putative pheromone compound

production. We found a set of 40 significant quantitative trait loci (QTL) representing 33 potential

pheromone compounds. QTL clustered significantly on two chromosomes, chromosome 8 for genital

compounds and chromosome 20 for wing compounds, and chromosome 20 was enriched for potential

pheromone biosynthesis genes. There was minimal overlap between pheromone QTL and known QTL

for mate choice and color pattern. Nonetheless, we did detect linkage between a QTL for wing

androconial area and *optix*, a color pattern locus known to play a role in reproductive isolation in these

species. This tight clustering of putative pheromone loci might contribute to coincident reproductive

isolating barriers, facilitating speciation despite ongoing gene flow.

Introduction

The genetic architecture of population differences can profoundly affect the evolution and maintenance
of new species, especially in the face of ongoing gene flow. For example, large effect loci are expected to
facilitate speciation (Via, 2012, Merrill et al., 2019) because they are less likely to be lost to drift (Kimura,
1983) and a single large effect locus can lead to substantial reproductive isolation on its own (e.g.
Bradshaw & Schemske, 2003). Similarly, tight physical linkage of barrier loci, *i.e.* those that contribute to
reproductive isolation, will promote speciation by impeding the breakdown of genetic associations
between traits that isolate emerging species (Felsenstein, 1981, Smadja & Butlin, 2011). As a result,
there is considerable interest in finding the genetic basis of traits that contribute to reproductive
isolation and understanding both the effect size and distribution of loci across the genome.

Physical linkage between barrier traits has been detected in a variety of taxa, including pea aphids
(Hawthorne & Via, 2001), *Laupala* crickets (Wiley et al., 2011), *Heliconius* butterflies (Merrill et al.,
2019), *Ficedula* flycatchers (Sæther et al., 2007), stickleback fish (Bay et al., 2017), and *Aquilegia*
columbines (Hodges et al., 2002). Barrier traits can include sex pheromones, chemical signals that
mediate intraspecific communication important for mating. Due to their critical role in mate attraction,
and their ability to convey information about species identity and male quality, pheromones can be
important for establishing and maintaining reproductive isolation through relatively simple changes in
chemical bouquets (Smadja & Butlin, 2009). As products of secondary metabolism (*i.e.* products not
required for survival), alterations in pheromones also have the potential to be both relatively simple at
the molecular level and, if such changes are relatively late in the pathway, likely avoid major pleiotropic
consequences.

Some of the best studied sex pheromones involved in speciation are those of female Lepidoptera, and in
some cases the genetic basis of variation in these pheromones is now well known. In *Ctenopseustis*, for
example, differential regulation of a desaturase drives differences in ratios of sex pheromone
components between species (Albre et al., 2012). Desaturases are also important in *Ostrinia* (Roelofs et
al., 2002, Sakai et al., 2009, Fujii et al., 2015) and *Helicoverpa* (Li et al., 2015, 2017). Fatty acyl-CoA
reductases (FARs) are responsible for species-specific pheromone alterations in *Ostrinia nubialis* strains
(Lassance et al., 2010, 2013). Within single species, desaturases and FARs have received the most
attention, being functionally characterized in a variety of Lepidoptera including *Yponomeuta* (Liénard et
al 2010), *Agrotis* (Ding & Löfstedt, 2015), *Manduca* (Buček et al., 2015), *Bombyx* (Moto et al., 2004), and
Antheraea (Wang et al., 2010). In some cases, variation in these female pheromones have also been

shown to contribute to reproductive isolation (e.g. Liebherr & Roelofs, 1975, Wu et al., 1999, Emelianov

et al., 2001; reviewed in Smadja & Butlin, 2009).

In contrast, male pheromones in Lepidoptera appear more varied, with a diverse array of compound

classes (Conner & Iyengar, 2016, Löfstedt et al., 2016). Many male Lepidopteran pheromones resemble

plant compounds and are thought to be diet-derived, especially terpenoids and pyrrolizine alkaloids, the

latter of which can be derived from either larval or adult feeding (Conner & Iyengar, 2016). In

comparison to female pheromones, the genetic basis of variation in male sex pheromones has received

less attention. In male *Bicyclus anynana*, a desaturase (*Ban-Δ11*) and two FARs, *Ban-wFAR1* and *Ban-*

wFAR2) produce pheromone components and precursors (Liénard et al., 2014), and in the moth *Ostrinia*

nubialis, males use the same Δ11-desaturase and Δ14-desaturase as females to produce pheromones

(Lassance & Löfstedt, 2009). However, the role of these genes in mediating differences between species

remains unclear.

Speciation is typically thought to rely on the accumulation of multiple reproductive barriers, involving

divergence in many different traits (Butlin & Smadja, 2017). In order to understand the extent of genetic

linkage that underlies species differences it will therefore be useful to study taxa in which multiple

different traits can be mapped. *Heliconius* butterflies have been well studied in the context of speciation

(Bates, 1861, Merrill et al., 2015, Jiggins, 2017), and in particular we know a great deal about the species

complex of *Heliconius melpomene* and its sister lineage *Heliconius cydno/timareta*. Genetic loci

underlying wing pattern and mate preference have been mapped in this group (Merrill et al., 2010,

Jiggins, 2017, Merrill et al., 2019). The additional role of chemical signaling in reproductive isolation has

long been suspected (Jiggins, 2008), but only recently studied in any detail. Chemical profiles of the male

wing androconia and genital regions (thought to act as aphrodisiacs and anti-aphrodisiacs, respectively)

differ between species (Estrada et al., 2011, Mann et al., 2017, Darragh et al., 2020), and are important

for mate choice (Mérot et al., 2015, Darragh et al., 2017, González-Rojas et al., 2020). Genital

pheromones play a more complex role in mate choice in *Heliconius*, as they are transferred by males to

females during mating and subsequently decrease advances by other males (Gilbert, 1976, Schulz et al.,

2007, Estrada, 2009).

One challenge of studying the entire chemical profile is determining which (combination of) compounds

are behaviorally active. The chemical bouquets of wings and genitals are complex, often consisting of

30-70 compounds (Mann et al., 2017, Darragh et al., 2017, 2020, Byers et al., 2020, this study), but it

remains unknown how many of these are important for signaling. A single wing compound in *H.*

melpomene, octadecanal, is known to be biologically active in *H. melpomene* and *H. cydno* (Byers et al.,

2020), while the genital compound (*E*)-β-ocimene acts as the main anti-aphrodisiac in *H. melpomene*

(Schulz et al., 2007). In addition, one bioactive genital compound in *H. cydno* (hexyl 3-methylbutyrate,

Estrada, 2009) is known. It seems likely that additional compounds in the complex bouquet (30-70

compounds) produced by each of these species are biologically active, but this remains to be tested.

The simple genetic control of pheromone components in other systems and their role as secondary

metabolism products suggests that they can be altered relatively simply without major pleiotropic

consequences; in combination with their importance in inter- and intraspecific mate choice, this makes

them ripe material for adaptation and speciation. We here conduct quantitative trait locus (QTL)

analyses for wing and genital compounds that differ between *Heliconius melpomene* L. and *H. cydno*

Doubleday (see also Byers et al., 2020, Darragh et al., 2020). We then investigated the patterns of QTL

distribution across the genome to test for clustering of loci across chromosomes.

Methods

Heliconius melpomene rosina and *H. cydno chioneus* (hereafter *H. melpomene* and *H. cydno*) and their

interspecific hybrids were reared in outdoor insectaries at the Smithsonian Tropical Research Institute in

Gamboa, Panama. Initial stocks were established from wild-caught individuals, and outbred stocks were

refreshed with more wild-caught individuals as necessary. Sexes were kept separately and virgin females

were kept separate from egg-laying females. To control for host plant effects, all larvae were fed on the

same host plant (*Passiflora platyloba* var. *williamsi*), and adults were maintained on a 20% sugar

solution and *Gurania eriantha*, *Psiguria triphylla*, *Psiguria warscewiczii*, and *Psychotria poeppigiana* as

pollen sources. Pheromones were collected from male individuals (both parental species and offspring

of mapping crosses) following (Darragh et al. 2017). Wing and genital data from *H. melpomene* and wing

data from *H. cydno* have previously been published in (Darragh et al., 2017, Darragh et al., 2019, Byers

et al., 2020). Briefly, males of 7-14 days post eclosion had their wings and genitals removed and soaked

in dichloromethane plus 1 ng/μL 2-tetradecyl acetate (DCM+IS) for one hour, after which the DCM+IS

was removed to a fresh vial and stored at -20°C before gas chromatography-mass spectrometry analysis.

Bodies were collected and preserved in 20% dimethyl sulfoxide (DMSO) and 0.25 M EDTA (pH 8.0) and

stored at −20°C for later DNA extraction.

Wing and genital extractions were run on an Agilent 7890B gas chromatograph coupled with an Agilent

5977 mass spectrometer (GC-MS) with electron ionization (Agilent Technologies, California, USA) with

an HP-5MS capillary column (30m length, 0.25mm inner diameter) and helium as the carrier gas (1.2

mL/min). Injection used an Agilent ALS 7693 autosampler and was splitless, with an inlet temperature of

250°C. The oven was first held isothermally at 50°C for five minutes, then increased at 5°C/minute to

320°C and again held isothermally for five minutes. Compounds were identified with a custom MS

library and quantified using the internal standard area. Only compounds that could be definitively

identified were included in the data set. To be included in the mapping experiment, compounds had to

be present in at least two thirds of the individuals from one of the parent species, i.e. they had to be

"typical" *H. melpomene* or *H. cydno* compounds. A limited set of wing compounds (three

methyloctadecanals and two henicosenes from *H. melpomene* and *H. cydno* respectively) were excluded

from the wing analysis due to close retention times in the GC-MS data preventing definitive

identification of the species phenotype in the mapping individuals. Absolute androconial area and

absolute hindwing area were measured in the GNU Image Manipulation Program (GIMP) as pixel counts,

then absolute androconial area was divided by hindwing area to produce the percentage of the

hindwing taken by the androconial region, the relative androconial area.

Mapping crosses consisted of backcrosses of either an *H. cydno* or *H. melpomene* mother and an F_1

father, as F_1 females are normally sterile. Ten families were constructed as backcrosses to *H. melpomene*

(representing 89 individuals for the wing phenotype, 81 individuals for the genital phenotype, and 78

individuals for the hindwing relative androconial area phenotype) and fifteen families were constructed

as backcrosses to *H. cydno* (127 individuals for the wing, 114 individuals for the genital phenotype, and

124 individuals for the hindwing relative androconial area phenotype) for a total of 216 and 195 hybrid

individuals for wing and genital studies respectively. DNA extraction, library preparation, and QTL map

construction are detailed in (Byers et al., 2020, Darragh et al., bioRxiv). Briefly, Qiagen DNeasy kits

(Qiagen) were used for DNA extraction, and individuals then genotyped either by RAD-seq or low-

coverage whole genome sequencing using nextera-based libraries (Picelli et al., 2014, Davey et al., 2017,

Merrill et al., 2019). Samples were sequenced by HiSeq 3000 (Illumina) by BGI (China). Linkage mapping

was conducted using the standard Lep-MAP3 (LM3) pipeline (Rastas, 2017). The initial linkage groups

and marker order were constructed based on the *H. melpomene* genome for 21 chromosomes. This

resulted in a linkage map with 447,818 SNP markers, which was then evenly thinned by a factor of ten to

44,782 markers to ease computation.

To determine which cross direction to use for each compound, we inspected the distribution of the

compound phenotype in the parents, F_1 individuals, and the backcross individuals. Compounds that

were present in equal amounts in both parent species (assessed using a Kruskal-Wallis test) were not

mapped. Once cross direction was determined, the remaining compounds were log-transformed to

approximate normality before being regressed against the linkage map using R/qtl2 (Broman et al.,

2018). Due to the family structure present in our crosses, we additionally included a kinship matrix

calculated by R/qtl2 using the LOCO (leave one chromosome out) method. Permutation testing was

used (with 1000 replicates) to determine QTL threshold. Compounds are likely to be correlated and our

data set represents multiple testing of the same mapping crosses, therefore, we used Bonferroni

correction separately for each cross direction to establish a second, more conservative, LOD threshold

for significance by dividing the threshold by the number of compounds (traits) mapped in the relevant

backcross direction. No correction was used for relative androconial area mapping as the trait was

mapped alone due to its general lack of correlation with individual volatile compounds. QTL confidence

intervals were obtained using bayes_int. To identify clustering of QTL, we first calculated the expected

number of QTL per chromosome, taking into account the chromosome's length in cM, then compared

that against the observed distribution of QTL per chromosome using a chi-squared test, computing p-

values with Monte Carlo permutation with B = 10,000 permutations. Examination of the residuals was

used to determine which chromosome(s) displayed clustering, with any residual over 2 considered

significant, as in (Erickson et al., 2016). We also repeated these tests taking into account chromosome

length in basepairs because the relationship between map length and physical length can differ across

chromosomes. The results were unchanged.

As our linkage map is based on whole genome sequencing data, we were able to recover the basepair

intervals corresponding to the QTL centimorgan (cM) confidence intervals (defined as the outermost

markers at those cM positions) as well as their peaks (defined as the first marker in at that cM position).

Lepbase (Challis et al., 2016) was queried to identify genes within the main interval on chromosome 20

for wing fatty acid-derived compound production (identified as the minimum overlapping window of all

Bonferroni-significant compound confidence intervals). Genes were searched against the nr (non-

redundant) protein database using BLASTp (Altschul et al., 1990) to obtain putative functional

annotations.

Results

We first looked at the distribution of compounds in *H. melpomene*, *H. cydno*, and their hybrids. Analysis

of the pheromones found a total of 31 compounds in the wings and 68 compounds in the genitals across

the two species that were possible to map (i.e. not unknown or overlapping in the gas chromatography-

mass spectrometry trace) (Figure 1, Table S1). There was limited overlap in compounds between the

wings and genitals, with only the straight-chain alkanes henicosane, docosane, tricosane, pentacosane,

and hexacosane (all of which may be part of the normal whole-body cuticular hydrocarbon profile), the

unsaturated aldehyde (*Z*)-11-icosenal, and the nitrogenous aromatic benzyl cyanide found in both body

regions. A subset of compounds (6 in the wings and 18 in the genitals) were not mapped as they did not

differ significantly between the parental species. The rest showed significant differences and thus had

adequate variation for QTL mapping to be feasible. Of the wing compounds, 11 showed segregation in

backcrosses to *H. cydno* and thus were mapped with those individuals; nine were mapped in

backcrosses to *H. melpomene*; and five showed segregation in both backcross directions and thus were

mapped separately in backcrosses to each species. For the genital compounds, 19 were mapped in

backcrosses to *H. cydno*; 17 in backcrosses to *H. melpomene*; and 14 in both directions.

Most compounds mapped did not produce significant QTL at either the $p = 0.05$ or the Bonferroni-

adjusted cutoff levels (the latter taking into account the number of compounds mapped) (Figures S1 and

S2). A total of 21 genital compounds and 14 wing compounds produced significant QTL, including three

peaks already published in (Byers et al., 2020, Darragh et al., bioRxiv) for octadecanal, 1-octadecanol,

and (*E*)-β-ocimene. Of these QTL, nine genital and ten wing compounds were significant at the more

conservative Bonferroni threshold (Tables 1 and 2; Table S2). The higher drop-out rate for genital

compounds after Bonferroni correction is likely the result of the more stringent correction due to a

higher number of compounds corrected for. We also found a single significant QTL on chromosome 18

for relative hindwing androconial area in backcrosses to *H. cydno* (Figure S3).

Compounds that showed significant QTL tended to be of specific chemical classes. For the wings, of the

14 wing compounds with significance at $p = 0.05$, 12 were fatty acid-derived compounds (FADs),

including alkanes as well as oxygenated alkanes and alkenes (aldehydes and alcohols). The remaining

two significant wing compounds were both nitrogenous aromatics. For the genitals, 10 compounds

producing significant QTL were macrolides and eight were FADs (two alkanes, four saturated esters, and

two unsaturated esters). The remaining three compounds were again nitrogenous aromatics (two

compounds) and one terpene. This is in general agreement with the distribution of compound types in

both body regions. The mapped wing compounds comprise 18 fatty acid-derived compounds, three

nitrogenous aromatics, two non-nitrogenous aromatics, and two terpenoids, so the finding of significant
QTL mostly for fatty acid-derived compounds is not surprising. For the genitals, the mapped compounds
consist of 22 fatty acid-derived compounds, 21 macrolides, four terpenoids, two nitrogenous aromatics,
and one lactone.

Significant quantitative trait loci showed some clustering on specific chromosomes (Figure S4). Notably,
these clusters did not overlap with known QTL for color pattern, mate choice, or other traits, which
generally tend not to be clustered in the genome, apart from a close association between mate choice
and a wing patterning gene (Figure 2). When either all QTL significant at $p < 0.05$ or Bonferroni-
corrected significant QTL were combined with known QTL, clustering was statistically significant (χ^2 =
64.036 for Bonferroni-significant QTL, χ^2 =78.524 for all significant QTL, $p < 0.001$ for both) for
chromosomes 8 and 20. In particular, all but one QTL for FAD compounds in the wings mapped to
chromosome 20 (the exception mapped to chromosome 1 as well as having a peak on chromosome 20).
The other significant wing compounds (all nitrogenous aromatics) mapped to chromosomes 10, 17, 19,
and the Z chromosome (chromosome 21). QTL on chromosome 20 were broadly overlapping, in part
due to our relatively small mapping populations producing broad Bayesian confidence intervals for QTL
location. In addition, the peaks themselves were strongly overlapping on the latter half of chromosome
20 (Figure 3). This clustering was statistically significant (χ^2 = 178.39 for all significant QTL, χ^2 = 113.02 for
Bonferroni-corrected significant QTL, $p < 0.001$ in both cases) and residual analysis identified
chromosome 20 as the sole outlier. When broken down by chemical class, only FAD compounds
clustered significantly, again on chromosome 20 (χ^2 = 232.49 for all significant QTL, $p < 0.001$), and this
was also significant when only Bonferroni-corrected significant QTL were tested (χ^2 = 137.55, $p < 0.001$).

Similar compound class-specific clustering was seen to a lesser extent for the genitals, where six of the
ten macrolides had QTL on chromosome 8, with the others more broadly dispersed on chromosomes 2,
7, 13, 14, and 20. The fatty acid-derived compounds (mostly esters) mapped more broadly across the
genome than in the wings, with QTL on chromosomes 1, 2, 3, 6, 8, 9, 14, and 20. Clustering was again
statistically significant (χ^2 = 46.037 for all significant QTL, χ^2 = 101.83 for Bonferroni-corrected significant
QTL, $p = 0.0017$ and $p < 0.001$ respectively) and residual analysis identified chromosomes 8 and 14 as
the outliers when all QTL were considered, and chromosome 8 when only Bonferroni-corrected QTL
were included. The clustering on chromosome 8 is solely due to the presence of macrolide QTL on this
chromosome, while chromosome 14 contains QTL for both a macrolide and three FAD compounds.
When broken down by chemical class, only macrolide compounds clustered significantly in the genitals,

again on chromosome 8 (χ^2 = 58.079 for all significant QTL, p < 0.001), and this was also significant when

only Bonferroni-corrected significant QTL were tested (χ^2 = 110.88, p < 0.001). Again, the QTL peaks

themselves were strongly overlapping in the middle of chromosome 8 (Figure 3) rather than being

dispersed across the chromosome. Two compounds had significant QTL in both wings and genitals:

henicosane and benzyl cyanide. Both data sets showed QTL on the same chromosomes (20 and 17

respectively, with additional QTL for benzyl cyanide in the wings on other chromosomes). The

henicosane QTL were within 1.5cM of one another, while the benzyl cyanide QTL on chromosome 17

were 2.3cM apart.

The strong clustering of QTL for wing fatty acid-derived compounds on chromosome 20 suggested a

common gene or set of linked genes responsible for fatty acid-derived compound metabolism or its

regulation. We searched the region between 43.11 - 56.37 cM (9,398,348 - 14,585,564 basepairs) for

potential candidate genes that could be responsible for this clustering. This region spanned the

confidence intervals for all Bonferroni-significant QTL apart from 1-octadecanol and contained 363

genes. Annotated proteins from this region (from Lepbase, Challis et al., 2016) were searched using

BLASTp against the NCBI non-redundant (nr) database, revealing a total of 20 genes potentially involved

in the biosynthesis of the wing compounds within this confidence interval on chromosome 20. These

genes include two acetyl-CoA carboxylases (involved in primary metabolism), fifteen fatty acyl-CoA

reductases (FARs), and two alcohol dehydrogenases (the latter two families involved in both primary

and secondary metabolism), none of which have been functionally characterized. Comparing our

candidates to those identified in (Liénard et al., 2014), none of our FARs fall within the pheromone gland

FAR (pgFAR) clade, whose members fall on chromosome 19. Inspecting the genome for additional

secondary metabolism fatty acid-derived metabolism genes, we identified clusters of FARs on

chromosomes 19 and 20 (Figure 4), with a more even distribution of alcohol dehydrogenases and

desaturases across the genome. Notably, we did not identify any QTL on chromosome 19 for fatty acid-

derived compound production in the wings.

Discussion

The genetic basis for traits contributing to reproductive isolation remains poorly studied. This is

especially the case for differences in chemical signaling between closely related non-model species, in

particular for species with complex chemical signatures where multiple components may play a role in

mate choice or species recognition. Here, we mapped a total of 40 QTL controlling the segregation of

wing and genital pheromone bouquet compounds in crosses between *Heliconius melpomene* and its

close relative *H. cydno*. There was significant evidence for non-random clustering of QTL, most notably

for wing compounds on chromosome 20, which showed broadly overlapping QTL with closely spaced

peaks. There could be a number of reasons for this clustering of loci. First, theory predicts that

mutations controlling species differences may accumulate, and persist, in regions already resistant to

gene flow, such that recombination is less likely to disrupt species differences (Smadja & Butlin, 2011).

Second, functionally similar genes are likely to occur in close linkage due to *e.g.* tandem gene

duplication, such that chemically similar compounds could be regulated by linked genes due to

functional similarities between their synthesis pathways (Osbourn, 2010). Finally, related compounds

could differ between species as a pleiotropic effect of the same mutation if they are downstream

products of the same biochemical pathway or in the case of a multifunctional enzyme. The end result of

all of these processes is linkage that can facilitate speciation by impeding recombination that will

otherwise disrupt the genetic associations between traits that characterize emerging species.

In the case of the compounds described here, it is likely that shared biosynthetic origins play an

important role in their chromosomal distributions. The majority of the wing compounds found in either

species are fatty acid-derived compounds (FADs; 19 of 31), and the majority of these FADs (12 of 19)

were associated with a significant QTL on chromosome 20. It is interesting that there was clustering of

one class of biosynthetic enzyme in this pathway, the fatty acyl-CoA reductases (FARs), on both

chromosomes 19 and 20, but no QTL on chromosome 19 for any wing or genital FAD compound,

suggesting the FARs on chromosome 19 are not involved in putative pheromone compound production.

Similarly, QTL for genital compounds exhibited slight clustering of loci underlying macrolide compound

production on chromosome 8, but otherwise were more evenly spread across the genome, likely due to

their greater biosynthetic diversity. The QTL for shared macrolide compounds may also be due to

shared biosynthetic origins of these compounds, but the biosynthesis of macrolides is less well

understood and consequently we were unable to identify candidate genes on this chromosome. We also

found a QTL on chromosome 3 for hexyl 3-methylbutyrate, a known bioactive component of the genital

pheromone of *H. cydno* (Estrada, 2009) whose genetic basis was previously unknown. Of note, the QTL

for relative androconial area roughly peaks on the gene *optix* (chromosome 18), a locus controlling red

color patterns in *Heliconius* (Reed et al., 2011, Van Belleghem et al., 2017). This locus is also known to

control male-specific scales in the basal heliconiine *Dryas iulia* and has an expression domain in the

hindwing androconia region (Martin et al., 2014). This overlap raises the possibility for coupled control

of color pattern and androconial pheromone production differences between *Heliconius* species.

Overall, we found no evidence for linkage of any of the chemical QTL clusters with other loci or QTL that

have been shown to play a difference in the strong assortative mating observed between *H. melpomene*

and *H. cydno*. The major wing patterning genes are located on chromosomes 1 (*aristalless*), 10 (*WntA*),

15 (*Cortex*), and 18 (*optix*), only chromosome 1 of which harbors a chemical QTL, but the peak does not

include the *aristalless* gene. In addition, major QTL for male mate choice map to chromosome 1, 17, and

18 (Merrill et al., 2019), but, similarly, the chemical QTL we identified on chromosomes 1 and 17 do not

overlap with the major QTL identified for male mate choice. We might expect an overlap between

chemical QTL and female mate choice QTL (as females likely assess males partially based on their wing

androconial chemistry), but the latter has not been mapped in this species pair. Nonetheless, the

clustering of wing compounds in particular might contribute to pheromone-based speciation,

particularly in the face of gene flow (Via, 2012). If more than one pheromone component that maps to

chromosome 20 is involved in reproductive isolation, and if genes responsible for their production are

tightly linked (or if pleiotropy occurs), this may also facilitate speciation (Felsenstein, 1981, Smadja &

Butlin, 2011, Merrill et al., 2019) and prevent the breakup of potentially adaptive pheromone bouquet

mixtures. Evidence suggests that mixture effects are important in pheromone processing (Riffell et al.

,2009, Clifford & Riffell, 2013, Lei et al., 2013), and these mixtures might be maintained through tight

genetic linkage between biosynthetic loci, facilitated by gene evolutionary events such as tandem

duplication. In contrast to the tight clustering seen on chromosome 20 for wing FAD compounds, we

saw more limited clustering for the more chemically diverse genital compounds, likely due to their more

diverse potential biosynthesis pathways. This might hamper any potential role in reproductive isolation.

Given their known function as anti-aphrodisiacs, genital compounds are not so likely to play a major role

in isolation between *Heliconius* species (Gilbert, 1976, Schulz et al., 2007, Estrada, 2009).

Some of the compounds found on wings and in genitalia might have a plant origin. Several compounds

in the wings and genitals did not differ between parental species, and many of these have been shown

to be affected by larval diet in *Heliconius melpomene*. Most of the invariant wing compounds are

aromatics, which are expected to be plant derived in *Heliconius* (Darragh et al., 2019). Notably, we

reared all mapping butterflies on the same larval and adult diets, thus effectively creating a dietary

common garden experiment, and so we do not expect larval or adult diet to affect the result of the QTL

mapping. Many male moth pheromone components are derived from host plant volatiles with minimal

alterations, potentially due to females' pre-existing biases for detecting these compounds during the

host plant search (Conner & Iyengar, 2016). However, many of our mapped *Heliconius* compounds do

appear to have a genetic basis (i.e. a QTL was found), in line with the biosynthesis of FAD and related

compounds by female moths. Breaking our compounds down into four classes (FADs, macrolides,

aromatics, and terpenoids), we failed to find QTL (at the $p = 0.05$ level) for 54%, 58%, 43%, and 83% of

these classes respectively across both wings and genitals. In particular, terpenoids that differ between

species may be the result of differential larval host plant use. It is of course possible that plant derived

compounds could show a genetic basis if there were species differences in the genes regulating their

uptake, and one example might be icosanal, which has been shown to be influenced by larval host plant

(Darragh et al., 2019) but which is also associated with a QTL in this study.

This study provides a first step towards better characterizing the genes causing chemical differences

between species. Fatty acid-derived compound biosynthesis candidate genes and QTL for FAD

compounds are both clustered on chromosome 20. These FAD compounds are produced from acetyl-

CoA followed by rounds of chain elongation, desaturation, fatty acyl-CoA reduction, oxidization, and

acetyltransferase activity to produce the final pheromone compounds. The fatty acyl-CoA reduction

step, key to pheromone biosynthesis, uses FARs to remove the CoA moiety and return an alcohol. More

than half of these FAR candidates are found in tandem duplications, as are the two alcohol

dehydrogenase candidates. These duplications could have provided raw material for the origin of new

biosynthetic function and thus increased pheromone diversity. FARs also cluster on chromosome 19,

and these FARs (unlike those on chromosome 20) fall within the pheromone gland FAR (pgFAR)

subfamily (Liénard et al., 2014, Löfstedt et al., 2016). However, these phylogenetic analyses of FARs in

Heliconius, *Bombyx*, *Danaus*, and *Bicyclus* do not generally show lineage-specific clustering of the

chromosome 20 FARs; instead, they are dispersed throughout the tree. This suggests an ancient

evolutionary origin of the *Heliconius* chromosome 20 FAR cluster, in contrast to *Heliconius*-specific

duplications of chromosome 19 FARs in the pgFAR subfamily which are clustered in the tree.

Single FAR and alcohol dehydrogenase genes are missing from *H. cydno* (Byers et al., 2020), suggesting

an obvious mechanism for the striking difference in FAD compounds between the two species; these

differences may also be due to e.g. differential regulation of FARs or other biosynthetic genes, or

mutations within these genes. Alternately, a single FAR gene or small set of genes may be responsible

for the biosynthesis of the majority of FAD compounds in *Heliconius*, similar to the situation found with

eloF in *Drosophila*, where knockdown of the gene produces widely pleiotropic effects on cuticular

hydrocarbon profiles (Combs et al., 2018). Similar effects are seen in *Spodoptera litura*, where RNAi of

FAR17 alone alters levels of the four major sex pheromone components (Lin et al., 2017). Another

option, though one we consider unlikely, is that this tight cluster of FARs on chromosome 20 serves as a

supergene, allowing multiple phenotypic traits (here individual chemical compounds) to co-segregate

and maintaining strong integration of the chemical phenotype. Supergenes have been described for

other phenotypic morphs in *Heliconius* (Joron et al., 2011), but we are unaware of mechanisms on

chromosome 20 that might contribute to supergene formation and maintenance, e.g. a chromosomal

inversion.

Information on potential biosynthetic gene clustering in other Lepidoptera species is largely lacking, as

most studies have produced transcriptomes of pheromone glands rather than linkage maps or

transcriptomes mapped to genomes. Additionally, as much work in Lepidoptera has focused on single

pheromone components and thus single biosynthetic genes (Groot et al., 2016), clustering of QTL and

candidate genes is rarely discussed. The lack of annotated genomes in many lepidopteran species also

can prevent in-depth assessment of clustering of candidate genes, requiring mapping QTL linkage groups

back to e.g. the *Bombyx* genome. However, such clustering has been seen in *Nasonia* wasps, where both

QTL for multiple cuticular hydrocarbon components and candidate desaturase genes overlap strongly

across the genome (Niehuis et al., 2011, 2013). Clusters of acetyltransferases and desaturases exist in

Heliothis (inferred from synteny with *Bombyx mori*), but the acetyltransferases do not appear to

underlie the QTL for sex pheromone differences between *H. virescens* and *H. subflexa* (Groot et al.,

2013, 2014). Quantitative trait loci for multiple compounds were investigated in *Heliothis*, finding

multiple chromosomes responsible for the production of the same compound classes (all FADs), with

more limited clustering (Sheck et al., 2006). This is similar in pattern to what we found in the genital

pheromone components, although the latter cover multiple compound classes.

Understanding the genetic architecture of complex traits can allow us to understand their potential role

in speciation. Determining whether the observed clustering QTL on e.g. chromosomes 8 and 20 reflect

tight linkage of multiple biosynthetic and/or regulatory genes or a single "master" locus responsible for

the biosynthesis or regulation of multiple pheromone components will contribute to our understanding

of both pheromone biosynthesis and pleiotropy and linkage in general. With genes in hand, CRISPR (a

technique that works in *Heliconius*, Livraghi et al., 2018) could be used to alter gene function, allowing

confirmation of the role of pheromone components via behavioral assays. As major effect loci (as these

QTL appear to be) and tight linkage (as potentially seen here) can both facilitate the speciation process,

especially in the face of gene flow (Felsenstein, 1981, Merrill et al., 2010, Smadja & Butlin, 2011, Via,

2012), the QTL identified here could be barrier or speciation genes in *Heliconius*. These specific QTL may

be involved in interspecific mate choice and reproductive isolation, and thus the genes underlying them

could contribute to the maintenance of species boundaries. We have here identified a total of 40 QTL

underlying species differences in 33 potential pheromone components, demonstrated clustering of

these QTL for wing and genital compounds on specific chromosomes not linked to known loci for species

differences, and identified candidate genes underlying the production of the major chemical class of

wing compounds. Together, these findings further our knowledge of chemical ecology, pheromone

genetics, and their potential roles in *Heliconius* butterfly mate choice and speciation.

References

Albre, J., Liénard, M. A., Sirey, T. M., Schmidt, S., Tooman, L. K., Carraher, C., ...Newcomb, R. D. (2012)

Sex Pheromone Evolution Is Associated with Differential Regulation of the Same Desaturase Gene in

Two Genera of Leafroller Moths. PLOS Genetics 8, e1002489. http://doi: 10.1371/journal.pgen.1002489

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990) Basic local alignment search tool.

Journal of Molecular Biology, 215, 403-410. http://doi: 10.1016/S0022-2836(05)80360-2

Bates, H. W. (1862) XXXII. Contributions to an Insect Fauna of the Amazon Valley. Lepidoptera:

Heliconidæ. Transactions of the Linnean Society of London, 23, 495-566. http://doi: 10.1111/j.1096-

3642.1860.tb00146.x

Bay, R. A., Arnegard, M. E., Conte, G. L., Best, J., Bedford, N. L., McCann, S. R., ...Peichel, C. L. (2017)

Genetic Coupling of Female Mate Choice with Polygenic Ecological Divergence Facilitates Stickleback

Speciation. Current Biology, 27, 3344-3349.e4. http://doi: 10.1016/j.cub.2017.09.037

Bradshaw, H. D., & Schemske, D. W. (2003) Allele substitution at a flower colour locus produces a

pollinator shift in monkeyflowers. Nature, 426, 176-178. http://doi: 10.1038/nature02106

Broman, K. W., Gatti, D. M., Simecek, P., Furlotte, N. A., Prins, P., Sen, Ś., ...Churchill, G. A. (2018) R/qtl2:

software for mapping quantitative trait loci with high-dimensional data and multi-parent populations.

Genetics, 211, 495-502. http://doi: 10.1534/genetics.118.301595

Buček, A., Matoušková, P., Vogel, H., Šebesta, P., Jahn, U., Weißflog., J., ...Pichová, I. (2015) Evolution of
moth sex pheromone composition by a single amino acid substitution in a fatty acid desaturase.
Proceedings of the National Academy of Sciences USA, 112, 12586-12591. http://doi:
10.1073/pnas.1514566112

Butlin, R. K., & Smadja, C. M. (2017) Coupling, reinforcement, and speciation. The American Naturalist,
191, 155-172. http://doi: 10.1086/695136

Byers, K. J. R. P., Darragh, K., Musgrove, J., Abondano Almeida, D., Fernanda Garza, S., Warren, I. A.,
...Jiggins, C. D. (2020) A major locus controls a biologically active pheromone component in *Heliconius*
melpomene. Evolution, 74, 349-364. http://doi: 10.1111/evo.13922

Challis, R. J., Kumar, S., Dasmahapatra, K. K., Jiggins, C. D., & Blaxter, M. (2016) Lepbase: the
Lepidopteran genome database. bioRxiv, 056994v1. http://doi: 10.1101/056994

Clifford, M. R., & Riffell, J. A. (2013) Mixture and odorant processing in the olfactory systems of insects: a
comparative perspective. Journal of Comparative Physiology A, 199, 911-928. http://doi:
10.1007/s00359-013-0818-6

Combs, P. A., Krupp, J. J., Khosla, N. M., Bua, D., Petrov, D. A., Levine, J. D., & Fraser, H. B. (2018) Tissue-
Specific cis-Regulatory Divergence Implicates *eloF* in Inhibiting Interspecies Mating in *Drosophila*.
Current Biology, 28, 3969-3975.e3. http://doi: 10.1016/j.cub.2018.10.036

Connor, W. E., & V. K. Iyengar (2016) Male pheromones in moths: reproductive isolation, sexy sons, and
good genes. In Pheromone Communication in Moths, J. D. Allison and R. Cardé, ed. Oakland, CA, USA:
University of California Press.

Darragh, K., Byers, K. J. R. P., Merrill, R. M., McMillan, W. O., Schulz, S., & Jiggins, C. D. (2019a) Male
pheromone composition depends on larval but not adult diet in *Heliconius melpomene*. Ecological
Entomology, 44, 397-405. http://doi: 10.1111/een.12716

Darragh, K., Montejo-Kovacevich, G., Kozak, K. M., Morrison, C. M., Figueiredo, C. M. E., Ready, J. S.,
...Jiggins, C. D. (2020) Species specificity and intraspecific variation in the chemical profiles of *Heliconius*
butterflies across a large geographic range. Ecology and Evolution. http://doi: 10.1002/ece3.6079

Darragh, K., Orteu, A., Byers, K. J. R. P., Szczerbowski, D., Warren, I. A., Rastas, P., ...Jiggins, C. D. (2019b)

A novel terpene synthase produces an anti-aphrodisiac pheromone in the butterfly *Heliconius*

melpomene. bioRxiv, 779678v1. http://doi: 10.1101/779678

Darragh, K., Vanjari, S., Mann, F., González-Rojas, M. F., Morrison, C. R., Salazar, C., ...Jiggins, C. D. (2017)

Male sex pheromone components in *Heliconius* butterflies released by the androconia affect female

choice. PeerJ, 5, e3953. http://doi: 10.7717/peerj.3953

Davey, J. W., Barker S. L., Rastas, P. M., Pinharanda, A., Martin, S. H., Durbin, R., ...Jiggins, C. D. (2017) No

evidence for maintenance of a sympatric *Heliconius* species barrier by chromosomal inversions.

Evolution Letters, 1, 138-154. http://doi: 10.1002/evl3.12

Ding, B.-J., & Löfstedt, C. (2015) Analysis of the *Agrotis segetum* pheromone gland transcriptome in the

light of sex pheromone biosynthesis. BMC Genomics, 16, 711. http://doi: 10.1186/s12864-015-1909-2

Emelianov, I., Drès, M., Baltensweiler, W., & Mallet, J. (2001) Host-induced assortative mating in host

races of the larch budmoth. Evolution, 55, 2002-2010. http://doi: 10.1111/j.0014-3820.2001.tb01317.x

Erickson, P. A., Glazer, A. M., Killingbeck, E. E., Agoglia, R. M., Baek, J., Carsanaro, S. M., ...Miller, C. T.

(2016) Partially repeatable genetic basis of benthic adaptation in threespine sticklebacks. Evolution, 70,

887-902. http://doi: 10.1111/evo.12897

Estrada, C. (2009) Sexual Behavior, Intraspecific Signaling and the Evolution of Mimicry among Closely

Related Species. PhD thesis, University of Texas Austin.

Estrada, C., Schulz, S., Yildizhan, S., & Gilbert, L. E. (2011) Sexual selection drives the evolution of

antiaphrodisiac pheromones in butterflies. Evolution, 65, 2843-2854. http://doi: 10.1111/j.1558-

5646.2011.01352.x

Felsenstein, J. (1981) Skepticism towards Santa Rosalia, or why are there so few kinds of animals?

Evolution, 35, 124-138. http://doi: 10.1111/j.1558-5646.1981.tb04864.x

Fujii, T., Yasukochi, Y., Rong, Y., Matsuo, T., & Ishikawa, Y. (2015) Multiple Δ11-desaturase genes

selectively used for sex pheromone biosynthesis are conserved in *Ostrinia* moth genomes. Insect

Biochemistry and Molecular Biology, 61, 62-68. http://doi: 10.1016/j.ibmb.2015.04.007

Gilbert, L. E. (1976) Postmating female odor in *Heliconius* butterflies: a male-contributed

antiaphrodisiac? Science, 193, 419-420. http://doi: 10.1126/science.935877

González-Rojas, M. F., Darragh, K., Robles, J., Linares, M., Schulz, S., McMillan, W. O., …Salazar, C. (2020)

Chemical signals act as the main reproductive barrier between sister and mimetic *Heliconius* butterflies.

Proceedings of the Royal Society B: Biological Sciences, 287, 20200587. http://doi:

10.1098/rspb.2020.0587

Groot, A. T., Dekker, T., & Heckel, D. G. (2016) The genetic basis of pheromone evolution in moths.

Annual Review of Entomology, 61, 99-117. http://doi: 10.1146/annurev-ento-010715-023638

Groot, A. T., Schöfl, G., Inglis, O., Donnerhacke, S., Classen, A., Schmalz, A., …Heckel, D. G. (2014) Within-

population variability in a moth sex pheromone blend: genetic basis and behavioral consequences.

Proceedings of the Royal Society B: Biological Sciences, 281, 20133054. http://doi:

10.1098/rspb.2013.3054

Groot, A. T., Staudacher, H., Barthel, A., Inglis, O., Schöfl, G., Santangelo, R. G., …Gould, F. (2013) One

quantitative trait locus for intra- and interspecific variation in a sex pheromone. Molecular Ecology, 22,

1065-1080. http://doi: 10.1111/mec.12171

Hawthorne, D. J., & Via, S. (2001) Genetic linkage of ecological specialization and reproductive isolation

in pea aphids. Nature, 412, 904-907. http://doi: 10.1038/35091062

Hodges, S. A., Whittall, J. B., Fulton, M., & Yang, J. Y. (2002) Genetics of Floral Traits Influencing

Reproductive Isolation between *Aquilegia formosa* and *Aquilegia pubescens*. American Naturalist, 159,

S51-S60. http://doi: 10.1086/338372

Jiggins, C. D. (2008) Ecological speciation in mimetic butterflies. BioScience, 58, 541-548. http://doi:

10.1641/B580610

Jiggins, C. D. (2017) The ecology and evolution of *Heliconius* butterflies. Oxford, UK: Oxford University

Press.

Joron, M., Frezal, L., Jones, R. T., Chamberlain, N. L., Lee, S. F., Haag, C. R., …ffrench-Constant, R. H.

(2011) Chromosomal rearrangements maintain a polymorphic supergene controlling butterfly mimicry.

Nature, 477, 203-206. http://doi: 10.1038/nature10341

Kimura, M. (1983) The neutral theory of molecular evolution. Cambridge, UK: Cambridge University
Press.

Lassance, J.-M., & Löfstedt, C. (2009) Concerted evolution of male and female display traits in the
European corn borer, *Ostrinia nubialis*. BMC Biology, 7, 10. http://doi: 10.1186/1741-7007-7-10

Lassance, J.-M., Groot, A. T., Liénard, M. A., Antony, B., Borgwardt, C., Andersson, F., ...Löfstedt, C.
(2010) Allelic variation in a fatty-acyl reductase gene causes divergence in moth sex pheromones.
Nature, 466, 486-489. http://doi: 10.1038/nature09058

Lassance, J.-M., Liénard, M. A., Antony, B., Qian, S., Fujii, T., Tabata, J., ...Löfstedt, C. (2013) Functional
consequences of sequence variation in the pheromone biosynthetic gene *pgFAR* for *Ostrinia* moths.
Proceedings of the National Academy of Sciences USA, 110, 3967-3972. http://doi:
10.1073/pnas.1208706110

Lei, H., Chiu, H.-Y., & Hildebrand, J. G. (2013) Responses of protocerebral neurons in *Manduca sexta* to
sex-pheromone mixtures. Journal of Comparative Physiology A, 199, 997-1014. http://doi:
10.1007/s00359-013-0844-4

Li, R.-T., Ning, C., Huang, L.-Q., Dong, J.-F., Li, X., & Wang, C.-Z. (2017) Expressional divergences of two
desaturase genes determine the opposite ratios of two sex pheromone components in *Helicoverpa*
armigera and *Helicoverpa assulta*. Insect Biochemistry and Molecular Biology, 90, 90-100. http://doi:
10.1016/j.ibmb.2017.09.016

Li, Z.-Q., Zhang, S., Luo, J.-Y., Wang, C.-Y., Lv, L.-M., Dong, S.-L., & Cui, J.-J. (2015) Transcriptome
comparison of the sex pheromone glands from two sibling *Helicoverpa* species with opposite sex
pheromone components. Scientific Reports, 5, 9324. http://doi: 10.1038/srep09324

Liebherr, J., & Roelofs, W. (1975) Laboratory Hybridization and Mating Period Studies Using Two
Pheromone Strains of *Ostrinia nubilalis*. Annals of the Entomological Society of America, 68, 305-309.
http://doi: 10.1093/aesa/68.2.305

Lin, X., Wang, B., & Du, Y. (2017) Key genes of the sex pheromone biosynthesis pathway in female moths
are required for pheromone quality and possibly mediate olfactory plasticity in conspecific male moths
in *Spodoptera litura*. Insect Molecular Biology, 27, 8-21. http://doi: 10.1111/imb.12335

Livraghi, L., Martin, A., Gibbs, M., Braak, N., Arif, S., & Breuker, C. J. (2018) Chapter Three - CRISPR/Cas9

as the Key to Unlocking the Secrets of Butterfly Wing Pattern Development and Its Evolution. Advances

in Insect Physiology, 54, 85-115. http://doi: 10.1016/bs.aiip.2017.11.001

Liénard, M. A., Hagström, Å. K., Lassance, J.-M., & Löfstedt, C. (2010) Evolution of multicomponent

pheromone signals in small ermine moths involves a single fatty-acyl reductase gene. Proceedings of the

National Academy of Sciences USA, 107, 10955-10960. http://doi: 10.1073/pnas.1000823107

Liénard, M. A., Wang, H.-L., Lassance, J.-M., & Löfstedt, C. (2014) Sex pheromone biosynthetic pathways

are conserved between moths and the butterfly *Bicyclus anynana*. Nature Communications, 5, 3957.

http://doi: 10.1038/ncomms4957

Löfstedt, C., Wahlberg, N., & Millar, J. G. (2016) Evolutionary patterns of pheromone diversity in

Lepidoptera. In Pheromone Communication in Moths, J. D. Allison and R. Cardé, ed. Oakland, CA, USA:

University of California Press.

Mann, F., Vanjari, S., Rosser, N., Mann, S., Dasmahapatra, K. K., Corbin, C., …Schulz, S. (2017) The scent

chemistry of *Heliconius* wing androconia. Journal of Chemical Ecology, 43, 843-857. http://doi:

10.1007/s10886-017-0867-3

Martin, A., McCulloch, K. J., Patel, N. H., Briscoe, A. D., Gilbert, L. E., & Reed, R. D. (2014) Multiple recent

co-options of *Optix* associated with novel traits in adaptive butterfly wing radiations. EvoDevo, 5, 7.

http://doi: 10.1186/2041-9139-5-7

Merrill, R. M., Dasmahapatra, K. K., Davey, J. W., Dell'Aglio, D. D., Hanly, J. J., Huber, B., …Yu, Q. (2015)

The diversification of *Heliconius* butterflies: what have we learned in 150 years? Journal of Evolutionary

Biology, 28, 1417-1438. http://doi: 10.1111/jeb.12672

Merrill, R. M., Rastas P., Martin S. H., Melo, M. C., Barker, S., Davey, J., …Jiggins, C. D. (2019) Genetic

dissection of assortative mating behavior. PLOS Biology 17:e2005902. http://doi:

10.1371/journal.pbio.2005902

Merrill, R. M., Van Schooten, B., Scott, J. A., & Jiggins, C. D. (2010) Pervasive genetic associations

between traits causing reproductive isolation in *Heliconius* butterflies. Proceedings of the Royal Society

of London B: Biological Sciences, 278, 511-518. http://doi: 10.1098/rspb.2010.1493

Moto, K., Suzuki, M. G., Hull, J. J., Kurata, R., Takahashi, S., Yamamoto, M., ...Matsumoto, S. (2004)

Involvement of a bifunctional fatty-acyl desaturase in the biosynthesis of the silkmoth, *Bombyx mori*, sex

pheromone. Proceedings of the National Academy of Sciences USA, 101, 8631-8636. http://doi:

10.1073/pnas.0402056101

Mérot, C., Frérot, B., Leppik, E., & Joron, M. (2015) Beyond magic traits: Multimodal mating cues in

Heliconius butterflies. Evolution, 69, 2891-2904. http://doi: 10.1111/evo.12789

Niehuis, O., Büllesbach, J., Gibson, J. D., Pothmann, D., Hanner, C., Mutti, N. S., ...Schmitt, T. (2013)

Behavioral and genetic analyses of *Nasonia* shed light on the evolution of sex pheromones. Nature, 494,

345-348. http://doi: 10.1038/nature11838

Niehuis, O., Büllesbach, J., Judson, A. K., Schmitt, T., & Gadau, J. (2011) Genetics of cuticular

hydrocarbon differences between males of the parasitoid wasps *Nasonia giraulti* and *Nasonia*

vitripennis. Heredity, 107, 61-70. http://doi: 10.1038/hdy.2010.157

Osbourn, A. (2010) Secondary metabolite gene clusters: evolutionary toolkits for chemical innovation.

Trends in Genetics, 26, 449-457. http://doi: 10.1016/j.tig.2010.07.001

Picelli S., Björklund, Å. K., Reinius, B., Sagasser, S., Winberg, G., & Sandberg, R. (2014) Tn5 transposase

and tagmentation procedures for massively scaled sequencing projects. Genome Research, 24, 2033-

2040. http://doi: 10.1101/gr.177881.114

Rastas, P. 2017. Lep-MAP3: robust linkage mapping even for low-coverage whole genome sequencing

data. Bioinformatics, 33, 3726-3732. http://doi: 10.1093/bioinformatics/btx494

Reed, R. D., Papa, R., Martin, A., Hines, H. M., Counterman, B. A., Pardo-Diaz, C., ...McMillan, W. O.

(2011) *optix* Drives the Repeated Convergent Evolution of Butterfly Wing Pattern Mimicry. Science, 333,

1137-1141. http://doi: 10.1126/science.1208227

Riffell, J. A., Lei, H., Christensen, T. A., & Hildebrand, J. G. (2009) Characterization and Coding of

Behaviorally Significant Odor Mixtures. Current Biology, 19, 335-340. http://doi:

10.1016/j.cub.2009.01.041

Roelofs, W. L., Liu, W., Hao, G., Jiao, H., Rooney, A. P., & Linn, C. E., Jr. (2002) Evolution of moth sex

pheromones via ancestral genes. Proceedings of the National Academy of Sciences USA, 99, 13621-

13626. http://doi: 10.1073/pnas.152445399

Sakai, R., Fukuzawa, M., Nakano, R., Tatsuki, S., & Ishikawa, Y. (2009) Alternative suppression of
transcription from two desaturase genes is the key for species-specific sex pheromone biosynthesis in
two *Ostrinia* moths. Insect Biochemistry and Molecular Biology, 39, 62-67. http://doi:
10.1016/j.ibmb.2008.10.001

Schulz, S., Estrada, C., Yildizhan, S., Boppré, M., & Gilbert L. E. (2007) An Antiaphrodisiac in *Heliconius*
melpomene Butterflies. Journal of Chemical Ecology, 34, 82-93. http://doi: 10.1007/s10886-007-9393-z

Sheck, A. L., Groot, A. T., Ward, C. M., Gemeno, C., Wang, J., Brownie, C., ...Gould, F. (2006) Genetics of
sex pheromone blend differences between *Heliothis virescens* and *Heliothis subflexa*: a chromosome
mapping approach. Journal of Evolutionary Biology, 19, 600-617. http://doi: 10.1111/j.1420-
9101.2005.00999.x

Smadja, C. M., & Butlin, R. K. (2009) On the scent of speciation: the chemosensory system and its role in
premating isolation. Heredity, 102, 77-97. http://doi: 10.1038/hdy.2008.55

Smadja, C. M., & Butlin, R. K. (2011) A framework for comparing processes of speciation in the presence
of gene flow. Molecular Ecology, 20, 5123-5140. http://doi: 10.1111/j.1365-294X.2011.05350.x

Sæther, S. A., Sætre, G.-P., Borge, T., Wiley, C., Svedin, N., Andersson, G., ...Qvarnström, A. (2007) Sex
Chromosome-Linked Species Recognition and Evolution of Reproductive Isolation in Flycatchers.
Science, 318, 95-97. http://doi: 10.1126/science.1141506

Van Belleghem, S. M., Rastas P., Papanicolaou, A., Martin, S. H., Arias, C. F., Supple, M. A., & Papa, R.
(2017) Complex modular architecture around a simple toolkit of wing pattern genes. Nature Ecology and
Evolution, 1, 0052. http://doi: 10.1038/s41559-016-0052

Via, S. (2012) Divergence hitchhiking and the spread of genomic isolation during ecological speciation-
with-gene-flow. Philosophical Transactions of the Royal Society of London B: Biological Sciences, 367,
451-460. http://doi: 10.1098/rstb.2011.0260

Wang, H.-L., Liénard, M. A., Zhao, C.-H., Wang, C.-Z., & Löfstedt, C. (2010) Neofunctionalization in an
ancestral insect desaturase lineage led to rare Δ6 pheromone signals in the Chinese tussah silkworm.
Insect Biochemistry and Molecular Biology, 40, 742-751. http://doi: 10.1016/j.ibmb.2010.07.009

Wiley, C., Ellison, C. K., & Shaw, K. L. (2011) Widespread genetic linkage of mating signals and

preferences in the Hawaiian cricket *Laupala*. Proceedings of the Royal Society of London B: Biological

Sciences, 279, 1203–1209. http://doi: 10.1098/rspb.2011.1740

Wu, W., Cottrell, C. B., Hansson, B. S., & Löfstedt, C. (1999) Comparative study of pheromone production

and response in Swedish and Zimbabwean populations of turnip moth, *Agrotis segetum*. Journal of

Chemical Ecology, 25, 177-196. http://doi: 10.1023/A:1020849419193

    Figures

671   Figure 1: Major compounds found in the wings (left) and genitals (right) of *Heliconius melpomene* and *H.*

672   *cydno*, arranged by compound type. Only those compounds comprising at least 1% of either species'

673   pheromone bouquet are included. n.s., not significant; ***: p < 0.005.

674
675

Figure 2: QTL for various traits across the *Heliconius melpomene* and *H. cydno* genomes. Wing

color/pattern loci represent the four major color loci (Jiggins, 2017); male mate choice are from (Merrill

et al., 2019); (relative) androconia area and Bonferroni-significant wing and genital compounds from this

study.

Figure 3: Chromosome-specific QTL plots for compounds with significant QTL (after Bonferroni

correction) on those chromosomes. Red: compound mapped in backcrosses to *H. melpomene*; blue:

compound mapped in backcrosses to *H. cydno*.

    Figure 4: Location of candidate genes for fatty acid-derived compound biosynthesis across the genome

    of *Heliconius melpomene*.

Tables

692 Table 1: Details of compounds with Bonferroni-significant QTL in *Heliconius* wing androconia. a: from

693 (Byers et al., 2020). c: in backcross to *H. cydno*. m: in backcross to *H. melpomene*.

Compound	Type	Mean (*H. melpomene*, ng)[a]	Mean (*H. cydno*, ng)[a]	Bonferroni-significant QTL on chromosome	QTL position on chromosome (bp)
(*Z*)-9-Octadecenal	Unsaturated aldehyde	10.94	Absent	1[m]	9955174[m]
Benzyl cyanide	Aromatic (nitrogenous)	Absent	1.813	17[m]	5596551[m]
Octadecanal[a]	Saturated aldehyde	783.4	0.569	20[c]; 20[m]	10947362[c]; 11725848[m]
Henicosane	Alkane	12.45	1152	20[m]	12126129[m]
Methyloctadecanol	Saturated alcohol	12.23	Absent	20[m]	12374765[m]
1-Octadecanol[a]	Saturated alcohol	345.6	Absent	20[m]	12652775[m]
(*Z*)-16-Methyl-9-octadecenol	Unsaturated alcohol	7.501	Absent	20[m]	12652775[m]
(*Z*)-13-Docosenal	Unsaturated aldehyde	62.56	0.276	20[c]	12917911[c]
(*Z*)-11-Icosenal	Unsaturated aldehyde	150.7	3.290	20[c]	13267611[c]
Methyl 3-indoleacetate	Aromatic (nitrogenous)	3.782	0.013	Z[c]	8478732[c]

695 Table 2: Details of compounds with Bonferroni-significant QTL in *Heliconius* genitals. a: from (Darragh et

696 al., 2020). c: in backcross to *H. cydno*. m: in backcross to *H. melpomene*.

Compound	Type	Mean (*H. melpomene*, ng)[a]	Mean (*H. cydno*, ng)	Bonferroni-significant QTL on chromosome	QTL position on chromosome (bp)
Hexyl 3-methylbutyrate	Saturated ester	Absent	421.1	3[m]	234227[m]
2-*sec*-Butyl-3-methoxypyrazine	Aromatic (nitrogenous)	32.66	3.569	5[c]	4427722[c]
12-Octadecanolide	Macrolide	Absent	105.4	8[c]	4512378[c]
(*Z*)-9-Octadecen-11-olide	Macrolide	Absent	7509	8[m]	5341821[m]
(*Z*9,*E*11)- 9,11-Octadecadien -13-olide	Macrolide	Absent	19714	8[m]	5341821[m]
Hexadecenolide	Macrolide	6.866	165.5	8[m]	5341821[m]
Hexyl octadecanoate	Saturated ester	121.8	24.21	8[m]	5987924[m]
(*Z*)-9-Octadecen-13-olide	Macrolide	7.282	5856	8[m]	6072562[m]
Isopropyl oleate	Unsaturated ester	50.22	2259	14[m]	7370836[m]

Supplemental Figures and Tables

Supplemental Figure 1: QTL plots for all mapped wing compounds. Dashed horizontal line: p = 0.05

significance threshold. Solid horizontal line: Bonferroni-corrected significance threshold. BC, backcross.

Supplemental Figure 2 (in two parts): QTL plots for all mapped genital compounds. Dashed horizontal

line: p = 0.05 significance threshold. Solid horizontal line: Bonferroni-corrected significance threshold.

BC, backcross.

Supplemental Figure 3: QTL plots for relative hindwing androconial area. Solid horizontal line: p = 0.05

significance threshold. HW, hindwing. BC, backcross.

Supplemental Figure 4: Whole-genome QTL plots for compounds with at least one significant QTL. Red:

compound mapped in backcrosses to *H. melpomene*; blue: compound mapped in backcrosses to *H.*

cydno. A: Compounds with at least one QTL significant at the p = 0.05 level shown. B: Compounds with

at least one QTL significant after Bonferroni correction shown.

Supplemental Table 1: Full list of compounds found in the wings and genitalia of *Heliconius melpomene*

and *H. cydno*, details of statistical comparisons of amounts in both species, and cross direction(s) the

compound was mapped in, if any.

Supplemental Table 2: Details of quantitative trait loci passing a significance threshold of either p = 0.05

or a Bonferroni-corrected threshold.