1

Tracking the origin of Baker's yeast: Seven new long-read genomes of highly divergent, wild

Saccharomyces cerevisiae **strains from East Asia**

Devin P Bendixsen*, Noah Gettle, Ciaran Gilchrist, Zebin Zhang, and Rike Stelkens

Division of Population Genetics, Department of Zoology, Stockholm University, Svante Arrheniusväg 18

B, 106 91 Stockholm, Sweden

* Corresponding author: Devin P Bendixsen, Department of Zoology Stockholm University;

email: devin.bendixsen@zoologi.su.se

Running title Tracking the origin of Baker's yeast

Keywords Saccharomyces cerevisiae, yeast, long-read, genome assembly, structural variation

Abstract

Comparative genome analyses have suggested East Asia to be the cradle of the domesticated microbe

Baker's yeast (*Saccharomyces cerevisiae*), used in the food and biotechnology industry worldwide. Here,

we provide seven new, high quality long read genomes of non-domesticated yeast strains isolated from

primeval forests and other natural environments in China and Taiwan. In a comprehensive analysis of our new

genome assemblies, along with all long read *Saccharomycetes* genomes available to date, we show that the

newly sequenced East Asian strains are among the closest living relatives of the ancestors of the global diversity

of Baker's yeast, confirming predictions made from short read genomic data. Three of these strains (termed the

East Asian Clade IX Complex here) share a recent ancestry and evolutionary history suggesting an early

divergence from other *S. cerevisiae* strains prior to the larger radiation of the species. Our genomic analyses

reveal that the wild East Asian strains contain elevated levels of structural variations (balanced and unbalanced),

including large inversions and non-reciprocal chromosomal translocations. Two strains have severely reduced

genomes, due to the loss of exon content rather than coding genes. The new genomic resources provided

here contribute to our understanding of the natural diversity of *S. cerevisiae*, expand the intraspecific

genetic variation found in this this heavily domesticated microbe, and provide a foundation for

understanding its origin and global colonization history.

Introduction

The history of the yeast *S. cerevisiae* is deeply interwoven with that of human migration, agriculture, and food technology. The sequencing of hundreds of *Saccharomyces* genomes has provided insight about the domestication of the species, its population genetic structure, and eukaryotic genome evolution (Scannell et al. 2011; Wang et al. 2012; Duan et al. 2018; Peter et al. 2018). The vast majority of these genomes have been analyzed using short-read sequencing, resulting in a focus on single nucleotide variants (SNVs). However, larger structural variations (SVs), such as inversions, deletions and gene duplications, are difficult to resolve using only short-read data (Goodwin et al. 2016). SVs are vital for evolutionary adaptation in yeast, especially to stressful conditions (Yona et al. 2012; Zheng et al. 2014; Jeffares et al. 2017; Zhang et al. 2020), and an increasing number of studies in other taxa support the role of inversions in adaptation and speciation (Hoffmann and Rieseberg 2008; Hooper and Price 2017) and the evolution of disease (Feder et al. 2014; Merker et al. 2018; Sanchis-Juan et al. 2018; Wellenreuther et al. 2019). To be able to dependably identify and resolve SVs, we need high quality long-read genome assemblies, spanning complex or repetitive genomic regions with a single continuous read. This approach has recently been applied to 15 *Saccharomyces* strains (Hebly et al. 2015; Yue et al. 2017; Naseeb et al. 2018; Macías et al. 2019) which, together with the annotation of individual genes and proteins, offers valuable insights into the functional genomics of the natural yeast diversity.

Among the known *Saccharomyces cerevisiae* strains are several highly divergent wild strains originating in East Asia (Wang et al. 2012; Duan et al. 2018; Peter et al. 2018). These wild lineages (termed CHN I-IX and Taiwanese) represent the most divergent *S. cerevisiae* strains described to date, with up to 1.1% genome-wide divergence, and have been suggested to represent the geographic origin of the species. Here, we applied long-read sequencing to seven of the highly divergent East Asian strains and one common laboratory strain (Y55, **Table 1**) and generated eight new high-quality genomes. Using these new genomes, we generated new phylogenetic reconstructions that place these East Asian strains at the

base of *S. cerevisiae* and provide further evidence for an out-of-China colonization history of this species. Furthermore, we find notable intraspecific sequence divergence among these newly sequenced East Asian strains and vast amounts of chromosomal SVs, including small and large inversions, translocations, chromosome fusions and aneuploidies, shedding light on the genomic history and dynamics of *S. cerevisiae*.

Table 1: Descriptions of the *Saccharomyces cerevisiae* strains sequenced in this study.

Lineage	Strain	Species	Source	Geographic Location
CHN I	HN1	*S. cerevisiae*	rotten wood, primeval forest	Diaoluo Mountain, Hainan, China
CHN II	SX2	*S. cerevisiae*	bark of a Fagaceae tree, primeval forest	Qinling Mountain, Shaanxi, China
CHN IV	HLJ1	*S. cerevisiae*	bark of Quercus mongokica, secondary forest	Jingbo lake, Heilongjiang, China
CHN VI	BJ4	*S. cerevisiae*	intestine of a butterfly, park	Haidian, Beijing, China
CHN IX	JXXY16.1	*S. cerevisiae*	bark, primeval forest	Xiangxiyuan, Hubei Province, China
CHN IX	XXYS1.4	*S. cerevisiae*	bark, primeval forest	Xiangxiyuan, Hubei Province, China
Taiwanese	EN14S01-3B	*S. cerevisiae*	soil	Sinyi, Nantou, Taiwan
West African+	Y55	*S. cerevisiae*	wine grapes	France

Results

Genome sequencing and assembly

We applied long-read PacBio sequencing to seven highly divergent and one common lab strain of *S. cerevisiae*, with an average per base genomic coverage of 235 (median 203). The nuclear and mitochondrial assemblies were highly complete with all chromosomes assembled into single contigs. Assessment of the completeness of the genome assembly and annotation using BUSCO found that six of our eight genomes had similarly high BUSCO scores (C > 97.2%). The remaining two genomes (BJ4 and HLJ1) had marginally lower scores (84.0% and 89.8%, respectively). Correspondingly, we found that BJ4 and HLJ1 had reduced genome sizes (11.42 and 11.37 Mbp compared to on average 11.93 Mbp in the other genomes sequenced here).

  Phylogenomics

Our newly constructed consensus species tree, based on 5847 genes, placed six of the newly assembled

East Asian strains in a basal position within the *S. cerevisiae* radiation (**Fig. 1**). Interestingly, three of these

strains (EN14S01-3B, XXYS1.4 and JXXY16.1), which were previously reported as being the most divergent

(CHN IX and Taiwanese clades), show early divergence from all other *S. cerevisiae* strains. The common

lab strain, Y55, clustered with two other domesticated strains (DBVPG6044 and SK1) within the West

African+ clade. Construction of an Alignment and Assembly-Free (AAF) phylogeny shows similar clustering

among the Far East strains and the common lab strain (**Fig. S1**) as well as a large amount of divergence of

the CHNIX and Taiwanese strains (hereon referred to as the Clade IX Complex) from the rest of *S.*

cerevisiae. Relative to the constructed consensus phylogeny, the AAF analysis places the seventh East

Asian strain (BJ4) closer to the other East Asian strains. AAF analysis comparing the long-read sequencing

data generated in this study and previous short-read data found a high-level of similarity (**Fig. S2**).

Figure 1: Maximum-likelihood phylogenetic tree of yeast long-read genomes. The tree was built from the alignment of all orthologous genes found in at least 5 strains (5847 genes). A maximum-likelihood single-tree inference was made for each locus and the consensus species tree is reported. Internal branches are labeled with gene concordance factor (gCF)/site concordance factor (sCF). gCF indicates the percent of gene trees supporting a given branch. sCF indicates the percentage of total aligned sites supporting a given branch. All internal branches were also supported with quartet scores equal to 1. For species with more than a single long-read genome assembly (*S. cerevisiae and S. paradoxus*), species clades are indicated in italics. *Saccharomyces* strains are colored according to their location of origin and branch tip shape indicates whether it is a domesticated (square) or wild (circle) strain. New long-read genome assemblies presented in this study are indicated in bold.

Comparative nuclear genomics

A comparison of our eight assembled *S. cerevisiae* genomes to each other and to previously published long-read genomes reveal a high-level of collinearity with the *S. cerevisiae* reference genome (S288C), with some notable deviations (**Fig. 2**, **Fig. S3-S4**). In particular, we found several large rearrangements within the Clade IX Complex. In all three member strains, we found a large terminal translocation from chromosome XI to chromosome XII (**Fig. 2 inset**). This ~80kb translocation is found conserved in these three strains but is not found in any other *Saccharomyces* strain. Another ~75kb inversion, was found in chromosome IV of the Taiwanese isolate EN14S01-3B but not in any other strain. We also found an apparent chromosomal fusion between chromosome VIII and chromosome XIII in strain JXXY16.1. We did not, however, find a loss or clear disruption of centromeric region from either chromosome. Other notable rearrangements are a large inversion in chromosome X of BJ4, three large translocations in chromosomes V, VII and X of HN1, and a large translocation and inversion in chromosome X of SX2 (**Fig. S3-S4**). The common lab strain, Y55, showed a high level of collinearity, with only minor deviations from homology.

Figure 2: *Saccharomyces cerevisiae* **long-read PacBio genome assemblies** Genome comparison of the reference strain, S288C, and three members of the East Asian Clade IX Complex: JXXY16.1, XXYS1.4 and EN14S01-3B. Sequence homology within the dot plots is indicated by red dots for forward matches and blue dots for reverse matches. Chromosomal paintings and chromosome intersections between genome assemblies are colored according to the S288C reference genome. Insets depict examples of deviations from homology: 1) a possible chromosome fusion between chromosome VIII and XIII in JXXY16.1, 2) a large translocation between XI and chromosome XII found conserved in JXXY16.1, XXYS1.4 and 3) EN14SO1-3B and a large inversion in chromosome IV of EN14S01-3B.

Nuclear genome content

With regards to genome content, our newly sequenced East Asian genomes largely resemble

those of Y55 and other assembled *S. cerevisiae* with a slightly lower range in gene counts (6095 – 6257

compared to 6289 – 6573) and 1.08 exons per gene. On average, 95.2% of these are directly orthologous

to genes identified in the reference strain, S288C. Similarly, numbers of ncRNA and rRNA regions resemble

those in other strains. To determine the source of the reduced genome sizes in strains BJ4 and HLJ1 we

also examined genomic feature content in terms of overall length. While these strains do not show

reduced numbers of genes, much of the differences in overall size are accounted for in loss of exon

content with regard to their total length rather than reductions in unannotated genome content that is

presumably non-coding (**Fig. S5-S7**). In addition to being outliers in exon content, BJ4 and HLJ1 also have

considerably more intronic content than other *S. cerevisiae* strains (**Fig. 3A**). Composite lengths in base

pairs of genomic features relative to overall feature mean for *S. cerevisiae* for total genomic length ($\bar{x}$ =

11.92 x 10^6; *s* = 2.26 x 10^5), exon regions ($\bar{x}$ = 8.46 x 10^6; *s* = 2.54 x 10^5), intron regions ($\bar{x}$ = 9.50 x 10^4; *s* =

1.47 x 10^4), and transposable elements (TEs) ($\bar{x}$ = 2.36 x 10^5; *s* = 9.68 x 10^4).

The amount of each genome composed of transposable elements (TEs) varied considerably

among strains, but does not appear to follow a clear geographical or phylogenetic pattern nor is it

correlated with genome size (**Fig. 3B**). In terms of TE make-up, all seven East Asian genomes, including the

Clade IX Complex, have fewer functional TY1 elements than other *S. cerevisiae* strains with relative

expansions in TY3 and TY5 elements, but clear patterns matching phylogenetic history are lacking.

Figure 3: Genomic feature composition of *Saccharomyces* (A) Composite lengths in base pairs of genomic features relative to overall feature mean for *S. cerevisiae* for total genomic length, exon regions, intron regions, and transposable elements (TEs). Strains are ordered according to the total genomic length. **(B)** Transposable element composition in total length subdivided by TY classification for *Saccharomyces sensu stricto* strains. Phylogram is taken from species tree described in **Fig. 1**. solo LTR = long terminal repeat components of TY elements without replicative machinery.

Unbalanced and balanced structural variations

In addition to the large rearrangements mentioned above, which are readily identified in a collinearity analysis, we performed a comprehensive analysis to quantify the amount of structural variations in our genomes. We assessed both unbalanced and balanced variations using pairwise comparisons between

the 15 *S. cerevisiae* strains with long-read assemblies. Unbalanced variations are those that change the

overall genomic content, such as insertions and deletions. Balanced variations are those that maintain

genomic content such as inversions and translocations. Unbalanced variations were largely composed of

repeat expansions/contractions (**Fig. S8**), whereas most of the balanced variations were translocations

(**Fig. S9**). Pairwise comparisons of all *S. cerevisiae* strains revealed that the three Clade IX Complex strains

(JXXY16.1, XXYS1.4, EN14S01-3B), showed significantly more balanced and unbalanced variation relative

to each other and the rest of the analyzed *S. cerevisiae* strains (**Fig. 4**). On average these strains had ~38

unbalanced and ~36 balanced variations, whereas the average for all other comparisons was ~23 for each

unbalanced and balanced. Interestingly, we also found large amounts of variation among the three

divergent strains. However, the largest number of structural rearrangements, especially unbalanced

variation, was found in the genomes of BJ4 and HLJ1, corresponding to their overall reduced genome size.

Figure 4: Structural variations within *Saccharomyces cerevisiae*. Pairwise comparisons among all *Saccharomyces cerevisiae* genome assemblies depicting the total **(A)** unbalanced and **(B)** balanced variations. Cladograms are adapted from the maximum-likelihood tree (**Fig. 1**). New long-read genome assemblies presented in this study are indicated in bold. For unbalanced variations, repeat expansions (RE) and contractions (RC), tandem expansions (TE) and contractions (TC), insertions (IS) and deletions (DL) were assessed. For balanced variations, inversions (IV), translocations (TR), fusions (FU) and fissions (FI) were assessed. Cartoons illustrate the different types of variation. Insets depict examples of pairwise comparisons. BJ4 and HLJ1 are plotted separately due to outlying counts of unbalanced SVs. The color scale is retained from the main heatmap with all values above the maximum of the main heatmap coloured dark blue. The maximum number of variants for either BJ4 or HLJ1 is shown for reference.

Comparative mitochondrial genomics

Overall, the mitochondrial genomes of the *S. cerevisiae* strains showed high-levels of collinearity (**Fig. S3-S4**). We did not, however, find a direct correspondence between our consensus gene phylogeny and that based on mitochondrial content (**Fig. 5**). Rather, we found a large division within *S. cerevisiae* strains with no clear geographical or evolutionary pattern. Five of the six *S. cerevisiae* strains that are domesticated were found on one side of the division. The nine wild strains, including the seven divergent East Asian strains along with a single domesticated strain (Y12) clustered on the other side of the division. Among the seven East Asian strains presented in this study, we found several interesting rearrangements including a duplication/inversion of COX1, ATP 6/8 in SX2, a duplication/inversion of 21S rRNA in JXXY16.1 and two duplications and an inversion of RPM1 in HN1. Most surprisingly, the 15S rRNA was not found in the mitochondrial assembly of BJ4 and a BLAST search of the raw sequencing data was not able to detect the gene. We also found that RPM1 was missing in XXYS1.4 and there was a severe reduction in size of VAR1.

Previous analyses have suggested that hybridization events can generate discordance between species and mitochondrial phylogenies (Peris et al. 2017; De Chiara et al. 2020). To investigate this, we also included other *Saccharomyces* species with available long read data in our mitochondrial phylogenetic analyses. For the most part, our mitochondrial phylogenies matched our species-level phylogeny with the notable exception of a strain of *S. jurei* (NCYC3947), a recently described European

species (Naseeb et al. 2018) that appears to share mitochondrial ancestry with a subgroup of European

strains of *S. paradoxus*. The mitochondrial genomes from this subgroup also contain large structural

variations (previously described in Yue et al. 2017) relative to other strains of *S. paradoxus* and *S. cerevisae*

further supporting their shared ancestry (**Fig. 5**).

Figure 5: Mitochondrial phylogenetics and genomic arrangements. Phylogenetic tree based on mitochondrial genomic content. Internal branches are labeled with bootstrap support. *Saccharomyces* strains are colored according to their location of origin and branch tip shape indicates whether it is a domesticated (square) or wild (circle) strain. New long-read genome assemblies presented in this study are indicated in bold. Major genomic elements found on yeast mitochondria are shown and colored according to guide elements at the top. Inverted elements appear on the underside of the line. Missing elements are indicated by dashed lines. Duplicated elements are indicated with asterisk.

Discussion

Comparative genomic analyses have provided clues about the origin of Baker's yeast (*S. cerevisiae*) and

have suggested an out-of-China origin (Peter et al. 2018). Here, we provide seven new, high quality long

read genomes of highly divergent wild strains recently isolated in East Asia. Phylogenomic analyses of the

long-read assemblies agree with previous findings that the wild East Asian strains (CHN, Taiwanese) are

basal relative to other *S. cerevisiae* strains (Duan et al. 2018; Peter et al. 2018) and, in the case of the CHN

IX and Taiwanese clades, show considerable divergence (**Fig. 1**). However, this is the first report where

representatives from these two clades are directly assessed and compared. We found that the two strains

from the CHN IX clade (JXXY16.1 and XXYS1.4) and the one strain from the Taiwanese clade (EN14S01-3B)

(collectively referred to as the Clade IX Complex) were highly divergent from other strains but share a

recent ancestry and evolutionary history. The placement of this clade relative to both the other East Asian

strains we describe, as well as other domesticated and lab strains from Africa and Europe, also indicates

that it diverged prior to the larger radiation of *S. cerevisiae* into other parts of the world.

In addition to providing insight into the evolutionary history of *S. cerevisiae*, our results in the

context of the seven previously assembled long-read genomes help reveal the extent of evolutionary

diversity in both genome structure and content. For example, BJ4 and HLJ1, belonging to CHNIV and CHVI

clades respectively, have genome sizes considerably smaller than not only all the *S. cerevisiae* strains

considered in our study (**Fig. 3A**) but also all *S. paradoxus*, *S. jurei*, *S. eubayanus*, and *S. jurei strains* (**Fig.**

S5). Contrary to our expectations that such size reductions would correspond to losses in non-coding

regions, we actually observed reduced amounts of genomic space taken up by coding regions (although

not in total number of genes) and an increase in intronic material in these shorter genomes (**Fig. 3A**).

The inclusion of the standard reference strain S288C in our analyses of genome structure and

content also adds to a growing consensus that, despite not being an outlier like BJ4 and HLJ1, S288C does

not reflect the average *S. cerevisiae* genome very well. One area where the use of the S288C genome as

representative appears problematic is the study of yeast transposable element evolution. Based on the

collective long read assemblies of divergent *S. cerevisiae* strains we analyzed here, we were able to

generate a broad scale map of *Ty* elements throughout this species (**Fig. 3B**). Previous conclusions based

largely on the S288C reference assembly have suggested that *Ty2* elements are a recent addition to *S.*

cerevisiae, arriving in the genome through horizontal gene transfer (HGT) from *S. mikatae* (Carr et al.

2012). While the lack of this element type in other closely related *Saccharomyces* species does provide

support for an HGT event, we found this element existing to varying degrees in even the most divergent

S. cerevisiae strains. This suggests an early origin of the *Ty2* element in the evolution of *S. cerevisiae.*

Without further genomic data available from diverse strains of *S. mikatae* it is difficult to delineate the

direction of this HGT. Previous findings have suggested that S288C may be a phenotypic extreme among

S. cerevisiae strains (Warringer et al. 2011). Overall, S288C may not be as representative of the species as

previously thought.

A similar level of diversity in TY composition among strains of *S. paradoxus*, *S. jurei*, *S. eubayanus*,

and *S. kudriavzevii* suggests a complex evolutionary history of TEs in *Saccharomyces*. TY2 and TY2p

elements were present in all *S. cerevisiae* strains but absent in other species, indicating their origin and

evolution only occurred after the divergence of *S. cerevisiae* from other Saccharomycotina.

Our comprehensive analyses of structural variations (SVs) further elucidates the evolutionary

history and intraspecific diversity of *S. cerevisiae*. Balanced and unbalanced SVs were identified for each

strain pair revealing patterns of genomic divergence. As expected, strains with reduced genome size (BJ4

and HLJ1) showed a large number of SVs dominated by unbalanced variations (**Fig. 4**). There is also a

noteworthy pattern of higher amounts of SVs in wild East Asian strains, especially the three strains within

the Clade IX Complex. As a species, *S. cerevisiae* has been shown to accumulate balanced variations at a

slower rate compared to *S. paradoxus* (Yue et al. 2017). This is likely due to the different selection histories

of these species. Many *S. cerevisiae* strains have long been associated with human activities where

domestication, cross-breeding and admixture have resulted in largely mosaic genomes (Liti et al. 2009;

Hyma and Fay 2013; Liti 2015), whereas *S. paradoxus* strains are recently isolated wild strains.

Interestingly, we found that wild East Asian strains accumulated both balanced and unbalanced SVs at a

high rate, more similar to rates normally seen in *S. paradoxus* (**Fig. 4**). It has been suggested that the

geographical isolation of some *S. paradoxus* subpopulations may have favored quick fixation of structural

rearrangements (Leducq et al. 2016). We may be witnessing similar patterns in the wild East Asian *S.*

cerevisiae strains here.

Overall, this study generates new, valuable genomic resources and expands our understanding of

the genetic variation and evolutionary history of this important microbial model system. This set of high-

quality genomes, encompassing both domesticated and wild populations from different ecological

backgrounds, provides a great testing field for open questions in evolutionary biology and fodder for

future research into the dynamics that govern eukaryotic genome evolution.

Methods

Yeast strain origins

We selected eight *Saccharomyces cerevisiae* strains for long-read sequencing and genome assembly

(**Table 1**). Seven of these strains are highly divergent strains originating from East Asia. Six strains were

isolated in China (Wang et al. 2012; Duan et al. 2018) from a variety of ecological niches and one in Taiwan

(Peter et al. 2018). The six Chinese strains cover many of the lineages (CHN I, II, IV, VI, and IX) previously

shown to be highly divergent from other *S. cerevisiae* strains based on short-read sequencing. The final

strain (Y55) is a common laboratory strain with a known mosaic genomic background originating in West

Africa, but first isolated in France. To place our analyses in context, we also utilized all currently available

Saccharomyces sensu stricto long read genome assemblies as well as assemblies from *Torulaspora*

delbrueckii and *Kluyveromyce lactis* (**Table S1**).

DNA preparation and long-read sequencing

Before sequencing, strains were sporulated and tetrads were dissected to allow for autodiploidization,

making strains homozygous across all loci. Strains were incubated at 30°C in 5ml YEPD (1% yeast extract,

2% peptone, 2% dextrose) in a shaking incubator for 24h before we harvested cells by centrifugation. We

extracted genomic DNA using NucleoSpin Microbial DNA extraction kit according to the manufacturer's

instructions (Macherey-Nagel). Genomic DNA for strain Y55 was extracted independently using the

QIAGEN Blood & Culture DNA Midi Kit. Samples were sequenced on PacBio Sequel and Sequel II platforms

at the NGI/Uppsala Genome Center (Science for Life Laboratory, Sweden) and the University of Minnesota

Sequencing Center (USA). In addition to these PacBio data, we also used publicly available paired-end

Illumina sequence data previously generated for each strain (**Table S1**).

Genome assembly and annotation

Contigs were assembled using raw PacBio reads with Flye (default settings) (Kolmogorov et al. 2019). We

used short read sequences for each strain to error-correct the long reads using FMLRC (Wang et al. 2018).

These corrected long reads and the short reads were subsequently used to polish the Flye assemblies

using Racon (Vaser et al. 2017) and POLCA (Zimin and Salzberg 2019) respectively. We further scaffolded

the contigs based on the reference S288C genome (GCA_000146045.2) using RaGOO (Alonge et al. 2019)

and filled any gaps this generated using multiple iterations of LR Gapcloser (Xu et al. 2019) and Gapcloser

(Luo et al. 2012). To account for any errors introduced by using long reads to fill gaps, we polished each

assembly once more using POLCA. We used REAPR (Hunt et al. 2013) to check for any misassemblies.

Assembly processing was conducted independently for each nuclear and mitochondrial genome. We

further adjusted the mitochondrial genomes by replacing large deletions of >500bp (relative to S288C)

with proportionate lengths of N sequence filler and repeated previous gap filling and polishing steps.

Completeness of the final genome assemblies was assessed using BUSCO v4.0.5 (Simão et al. 2015;

Waterhouse et al. 2018).

We annotated nuclear genes, mitochondrial genes, centromeres, transposable elements, core X

elements and Y-prime elements using pipelines within the LRSDAY package (Yue and Liti 2018). In addition

to our 8 newly assembled genomes, we also used the same method to annotate the other long read

assemblies publicly available for S. cerevisiae strains as well as related yeast species (**Table S2**). Nuclear

genes orthologous to annotated genes in the S. cerevisiae S288C reference genome were identified using

Proteinortho (Lechner et al. 2011). Genes for which no orthologous protein was found in the reference

were clustered based on orthology to each other.

Phylogenomic analysis

To place our 8 assembled genomes within the context of other *Saccharomyces* strains, we employed both

a consensus gene tree and Assembly and Alignment-Free (AAF) approach to phylogenetic tree

construction. For the consensus gene tree approach, we aligned all orthologous genes found in at least 5

strains (5847 genes) using MUSCLE (Edgar 2004) and performed maximum-likelihood single-tree inference

for each locus using RAxML-NG (Kozlov et al. 2019) with a discrete GAMMA model of rate heterogeneity.

We used Astral-III (Zhang et al. 2018) with these gene trees to generate a consensus species tree (**Fig. 1**).

Alternatively, AAF (Fan et al. 2015) was used with a k-mer size of 40 nucleotides and a threshold frequency

of 3 for each k-mer to be included in the analysis. First, AAF was used on the long-read sequencing data

generated in this study and compared to previously published Illumina short-read sequencing data for the

same strains (**Fig. S2, Table S1**). We then used a non-parametric bootstrap (n=100) to generate a

consensus tree using the N44 genome as an outgroup of the analysis resulting in 100% bootstrap support

across the tree. We then used AAF with the same parameters to build a phylogenetic network of the 8 *S.*

cerevisiae strains presented in this study along with 13 other *Saccharomyces* genomes that have

previously been sequenced using long-read sequencing (**Fig. S1, Table S1**).

To generate the mitochondrial phylogeny, we reoriented the start of each assembly based on the

position of the tRNA gene, trnP(ugg), then aligned these assemblies to each other using Mugsy (Angiuoli

and Salzberg 2011). This multiple sequence alignment was then used to create a maximum likelihood tree

using IQ-TREE (v2.0.5) (options: -m TPM2u+F+R3 -B 1000 -bnni -alrt 1000) (Nguyen et al. 2015; Hoang et

al. 2018). The model was determined using the ModelFinder component of IQ-Tree (Kalyaanamoorthy et

al. 2017).

Unbalanced and balanced structural variation

To identify the structural variations between strains within *Saccharomyces cerevisiae*, we performed

exhaustive pairwise comparisons between the 15 strains with long-read assemblies (210 comparisons).

To identify unbalanced variation, we used nucmer (Kurtz et al. 2004) to generate a whole-genome

alignment for each strain pair (options: -maxmatch -c 500). We then used Assemblytics (Nattestad and

Schatz 2016) to identify repeat expansions (RE) and contractions (RC), tandem expansions (TE) and

contractions (TC), insertions (IS) and deletions (DL) (options: unique sequence length required = 10000,

minimum variant size = 50, maximum variant size = 10000). We then filtered these variants to retain only

those that encompassed at least a single protein-coding gene (CDS) by intersecting our variants with our

genome annotations using bedtools intersect (Quinlan and Hall 2010). This method of identifying

structural variations results in two sets of identified variations within a pairwise comparison, one with

each genome acting as reference sequence and the other acting as query sequence. These sets are often

identical (albeit reversed), however some variants might be found in only one direction and missed by the

algorithm in the other. In these instances, the average of the two sets of variations is reported (sometimes

resulting in a half variation being reported).

In order to identify balanced variations, we first identified conserved synteny blocks for each

pairwise comparison using SynChro from the CHROnicle package (Drillon et al. 2013; Drillon et al. 2014).

The input files were prepared for SynChro by separating genome fasta files and GFF annotation files by

chromosome and converting them to EMBL format using EMBLmyGFF3 (Norling et al. 2018). We then

used ReChro from the CHROnicle package (Vakirlis et al. 2016), with synteny block stringency parameter

(delta) set to 0, to identify inversions (IV), translocations (TR), fusions (FU) and fissions (FI).

Data Access

All raw sequencing reads and complete genome assemblies from this study have been submitted to the

European Nucleotide Archive under accession number **XXXXXXXXX**.

Acknowledgments

We would like to thank Feng-Yan Bai and Gianni Liti for donating strains, Jia-Xing Yue for advice on the

LRSDAY pipeline, and Gianni Liti and Chris Wheat for discussion. We acknowledge the support of the

National Genomics Infrastructure (NGI)/Uppsala Genome Center and UPPMAX for providing assistance in

massive parallel sequencing and computational infrastructure. Work performed at NGI/Uppsala Genome

Center (project SNIC 2019/8-23) has been funded by RFI/VR and Science for Life Laboratory, Sweden. This

work was supported by grants from Stockholm University (Science for Life Laboratory sequencing grant

SU FV-2.2.2-1843-17 to RS), the Swedish Research Council (2017-04963 to RS), the Knut and Alice

Wallenberg Foundation (2017.0163 to RS), the Carl Trygger foundation (to ZZ), the Wenner-Gren

Foundations (UPD2018-0196, UPD2019-0110 to DPB), and The University of Minnesota Department of

Ecology, Evolution, and Behavior.

References

Alonge M, Soyk S, Ramakrishnan S, Wang X, Goodwin S, Sedlazeck FJ, Lippman ZB, Schatz MC.
2019. RaGOO: fast and accurate reference-guided scaffolding of draft genomes. *Genome*
biology **20**: 1-17.
Angiuoli SV, Salzberg SL. 2011. Mugsy: fast multiple alignment of closely related whole
genomes. *Bioinformatics* **27**: 334-342.
Carr M, Bensasson D, Bergman CM. 2012. Evolutionary genomics of transposable elements in
Saccharomyces cerevisiae. *PloS one* **7**.
De Chiara M, Friedrich A, Barré B, Breitenbach M, Schacherer J, Liti G. 2020. Discordant
evolution of mitochondrial and nuclear yeast genomes at population level. *BMC biology*
18: 1-15.
Drillon G, Carbone A, Fischer G. 2013. Combinatorics of chromosomal rearrangements based on
synteny blocks and synteny packs. *Journal of Logic and Computation* **23**: 815-838.
Drillon G, Carbone A, Fischer G. 2014. SynChro: a fast and easy tool to reconstruct and visualize
synteny blocks along eukaryotic chromosomes. *PLoS One* **9**.
Duan S-F, Han P-J, Wang Q-M, Liu W-Q, Shi J-Y, Li K, Zhang X-L, Bai F-Y. 2018. The origin and
adaptive evolution of domesticated populations of yeast from Far East Asia. *Nature*
communications **9**: 1-13.
Fan H, Ives AR, Surget-Groba Y, Cannon CH. 2015. An assembly and alignment-free method of
phylogeny reconstruction from next-generation sequencing data. *BMC genomics* **16**:
522.
Feder JL, Nosil P, Flaxman SM. 2014. Assessing when chromosomal rearrangements affect the
dynamics of speciation: implications from computer simulations. *Frontiers in genetics* **5**:
295.
Goodwin S, McPherson JD, McCombie WR. 2016. Coming of age: ten years of next-generation
sequencing technologies. *Nature Reviews Genetics* **17**: 333.
Hebly M, Brickwedde A, Bolat I, Driessen MR, de Hulster EA, van den Broek M, Pronk JT,
Geertman J-M, Daran J-M, Daran-Lapujade P. 2015. S. cerevisiae× S. eubayanus
interspecific hybrid, the best of both worlds and beyond. *FEMS Yeast Research* **15**.
Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: improving the
ultrafast bootstrap approximation. *Molecular biology and evolution* **35**: 518-522.
Hoffmann AA, Rieseberg LH. 2008. Revisiting the impact of inversions in evolution: from
population genetic markers to drivers of adaptive shifts and speciation? *Annual review*
of ecology, evolution, and systematics **39**: 21-42.
Hooper DM, Price TD. 2017. Chromosomal inversion differences correlate with range overlap in
passerine birds. *Nature ecology & evolution* **1**: 1526-1534.
Hunt M, Kikuchi T, Sanders M, Newbold C, Berriman M, Otto TD. 2013. REAPR: a universal tool
for genome assembly evaluation. *Genome biology* **14**: R47.
Hyma KE, Fay JC. 2013. Mixing of vineyard and oak-tree ecotypes of Saccharomyces cerevisiae
in North American vineyards. *Molecular ecology* **22**: 2917-2930.
Jeffares DC, Jolly C, Hoti M, Speed D, Shaw L, Rallis C, Balloux F, Dessimoz C, Bähler J, Sedlazeck
FJ. 2017. Transient structural variations have strong effects on quantitative traits and
reproductive isolation in fission yeast. *Nature communications* **8**: 1-11.

Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. *Nature methods* **14**: 587.
Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using
repeat graphs. *Nature biotechnology* **37**: 540-546.
Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004.
Versatile and open software for comparing large genomes. *Genome biology* **5**: R12.
Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ. 2011. Proteinortho: detection
of (co-) orthologs in large-scale analysis. *BMC bioinformatics* **12**: 124.
Leducq JB, Nielly-Thibault L, Charron G, Eberlein C, Verta JP, Samani P, Sylvester K, Hittinger CT,
Bell G, Landry CR. 2016. Speciation driven by hybridization and chromosomal plasticity
in a wild yeast. *Nat Microbiol* **1**: 15003.
Liti G. 2015. The fascinating and secret wild life of the budding yeast S. cerevisiae. *eLife*
4:e0585.
Liti G, Carter DM, Moses AM, Warringer J, Parts L, James SA, Davey RP, Roberts IN, Burt A,
Koufopanou V. 2009. Population genomics of domestic and wild yeasts. *Nature* **458**:
337-341.
Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y. 2012. SOAPdenovo2: an
empirically improved memory-efficient short-read de novo assembler. *Gigascience* **1**:
2047-2217X-2041-2018.
Macías LG, Morard M, Toft C, Barrio E. 2019. Comparative genomics between Saccharomyces
kudriavzevii and S. cerevisiae applied to identify mechanisms involved in adaptation.
Frontiers in Genetics **10**: 187.
Merker JD, Wenger AM, Sneddon T, Grove M, Zappala Z, Fresard L, Waggott D, Utiramerur S,
Hou Y, Smith KS. 2018. Long-read genome sequencing identifies causal structural
variation in a Mendelian disease. *Genetics in Medicine* **20**: 159-163.
Naseeb S, Alsammar H, Burgis T, Donaldson I, Knyazev N, Knight C, Delneri D. 2018. Whole
genome sequencing, de novo assembly and phenotypic profiling for the new budding
yeast species Saccharomyces jurei. *G3: Genes, Genomes, Genetics* **8**: 2967-2977.
Nattestad M, Schatz MC. 2016. Assemblytics: a web analytics tool for the detection of variants
from an assembly. *Bioinformatics* **32**: 3021-3023.
Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular biology*
and evolution **32**: 268-274.
Norling M, Jareborg N, Dainat J. 2018. EMBLmyGFF3: a converter facilitating genome
annotation submission to European Nucleotide Archive. *BMC research notes* **11**: 1-5.
Peris D, Arias A, Orlić S, Belloch C, Perez-Traves L, Querol A, Barrio E. 2017. Mitochondrial
introgression suggests extensive ancestral hybridization events among Saccharomyces
species. *Molecular phylogenetics and evolution* **108**: 49-60.
Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B, Freel K,
Llored A. 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates.
Nature **556**: 339.
Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**: 841-842.

Sanchis-Juan A, Stephens J, French CE, Gleadall N, Mégy K, Penkett C, Shamardina O, Stirrups K,
Delon I, Dewhurst E. 2018. Complex structural variants in Mendelian disorders:
identification and breakpoint resolution using short-and long-read genome sequencing.
Genome medicine **10**: 95.
Scannell DR, Zill OA, Rokas A, Payen C, Dunham MJ, Eisen MB, Rine J, Johnston M, Hittinger CT.
2011. The Awesome Power of Yeast Evolutionary Genetics: New Genome Sequences
and Strain Resources for the Saccharomyces sensu stricto Genus. *G3 (Bethesda)* **1**: 11-
25.
Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics **31**: 3210-3212.
Vakirlis N, Sarilar V, Drillon G, Fleiss A, Agier N, Meyniel J-P, Blanpain L, Carbone A, Devillers H,
Dubois K. 2016. Reconstruction of ancestral chromosome architecture and gene
repertoire reveals principles of genome evolution in a model yeast genus. *Genome*
research **26**: 918-932.
Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly from
long uncorrected reads. *Genome research* **27**: 737-746.
Wang JR, Holt J, McMillan L, Jones CD. 2018. FMLRC: Hybrid long read error correction using an
FM-index. *BMC bioinformatics* **19**: 50.
Wang QM, Liu WQ, Liti G, Wang SA, Bai FY. 2012. Surprisingly diverged populations of
Saccharomyces cerevisiae in natural environments remote from human activity.
Molecular ecology **21**: 5404-5417.
Warringer J, Zörgö E, Cubillos FA, Zia A, Gjuvsland A, Simpson JT, Forsmark A, Durbin R, Omholt
SW, Louis EJ. 2011. Trait variation in yeast is defined by population history. *PLoS*
genetics **7**.
Waterhouse RM, Seppey M, Simão FA, Manni M, Ioannidis P, Klioutchnikov G, Kriventseva EV,
Zdobnov EM. 2018. BUSCO applications from quality assessments to gene prediction
and phylogenomics. *Molecular biology and evolution* **35**: 543-548.
Wellenreuther M, Mérot C, Berdan E, Bernatchez L. 2019. Going beyond SNPs: the role of
structural genomic variants in adaptive evolution and species diversification. *Molecular*
ecology **28**: 1203-1209.
Xu G-C, Xu T-J, Zhu R, Zhang Y, Li S-Q, Wang H-W, Li J-T. 2019. LR_Gapcloser: a tiling path-based
gap closer that uses long reads to complete genome assembly. *GigaScience* **8**: giy157.
Yona AH, Manor YS, Herbst RH, Romano GH, Mitchell A, Kupiec M, Pilpel Y, Dahan O. 2012.
Chromosomal duplication is a transient evolutionary solution to stress. *Proceedings of*
the National Academy of Sciences **109**: 21010-21015.
Yue J-X, Li J, Aigrain L, Hallin J, Persson K, Oliver K, Bergström A, Coupland P, Warringer J,
Lagomarsino MC. 2017. Contrasting evolutionary genome dynamics between
domesticated and wild yeasts. *Nature genetics* **49**: 913.
Yue J-X, Liti G. 2018. Long-read sequencing data analysis for yeasts. *Nature protocols* **13**: 1213.
Zhang Z, Bendixsen DP, Janzen T, Nolte AW, Greig D, Stelkens R. 2020. Recombining your way
out of trouble: The genetic architecture of hybrid fitness under environmental stress.
Molecular biology and evolution **37**: 167-182.

Zheng D-Q, Chen J, Zhang K, Gao K-H, Li O, Wang P-M, Zhang X-Y, Du F-G, Sun P-Y, Qu A-M.
2014. Genomic structural variations contribute to trait improvement during whole-
genome shuffling of yeast. *Applied microbiology and biotechnology* **98**: 3059-3070.
Zimin AV, Salzberg SL. 2019. The genome polishing tool POLCA makes fast and accurate
corrections in genome assemblies. *bioRxiv*.
516