Information theoretic approaches to deciphering the neural code with functional fluorescence imaging

Jason R. Climer[1], PhD and Daniel A. Dombeck[1]*, Ph.D.

[1] Department of Neurobiology, Northwestern University, Evanston IL
*Lead Contact: d-dombeck@northwestern.edu

Abstract

Information theoretic metrics have proven highly useful to quantify the relationship between behaviorally relevant parameters and neuronal activity with relatively few assumptions. However, such metrics are typically applied to action potential recordings and were not designed for the slow timescales and variable amplitudes typical of functional fluorescence recordings (e.g. calcium imaging). Therefore, the power of information theoretic metrics has yet to be fully exploited by the neuroscience community due to a lack of understanding for how to apply and interpret the metrics with such fluorescence traces. Here, we used computational methods to create mock action potential traces with known amounts of information and from them generated fluorescence traces to examine the ability of different information metrics to recover the known information values. We provide guidelines for the use of information metrics when applied to functional fluorescence and demonstrate their appropriate application to GCaMP6f population recordings from hippocampal neurons imaged during virtual navigation.

Introduction

Neurons encode parameters important for animal behavior, at least in part, through the rate of production of action potentials. Evidence for this fundamental insight into the brain's language for coding different types of parameter spaces can be found from electrophysiological action potential recordings of orientation tuning in the visual system (Hubel and Wiesel, 2009), chemical sensing in the olfactory system (Leveteau and MacLeod, 1966; Wachowiak and Shipley, 2006), and spatial encoding in the hippocampus (O'Keefe, 1976). Defining metrics to quantify the relationship between a neuron's spiking rate and the behavioral parameter spaces has therefore become a key step in deciphering the neural code, but there are many metrics used for quantification. These metrics are often used to compare neural responses across conditions or in neurons with complex responses, but the underlying assumptions of the different metrics then become important factors to consider when determining which one to use.

Information theory is growing in popularity in the neuroscience community, largely because it provides a means to quantify rate coding with relatively few assumptions compared to other metrics. For example, information theory can be applied to neurons with widely varying response properties because it: 1. Is a nonlinear metric, and therefore does not require the linearity assumptions of correlation metrics (e.g. (Grubb and Thompson, 2006; Hinman et al., 2016; Jiaying Tang, 2015; Kropff et al., 2015), 2. Does not assume a response shape, as is often the case with Gaussian receptive field mapping metrics (e.g. Kraus et al., 2015; Soo et al., 2011) or metrics using curve fitting to an exponential or polynomial curve (Hinman et al., 2016; Jiaying Tang, 2015), and 3. Uses the full time trace or full shape of the mean response profile, rather than defining receptive fields as the portion that exceeds some threshold (e.g. Harvey et al., 2009; Niell and Stryker, 2008; Pastalkova et al., 2008). A useful information theoretic measure is mutual information (MI). Typically measured in bits per unit time, MI describes the increase in predictability of the neural response when behavioral parameters are known. However, MI can be nontrivial to estimate from neural and behavioral recordings and its estimation is an ongoing area of research (Belghazi et al., 2018; Gao et al., 2017; Kraskov et al., 2004; Timme and Lapish, 2018).

Here we focus on the most widely used estimator of MI used in neuroscience, the SMGM estimator which was developed by Skaggs, McNaughton, Gothard and Markus (Skaggs et al., 1993), though as a point of comparison, we also consider the Binned Estimator (Timme and Lapish, 2018) and a technique developed by Kraskov, Stogbauer and Grassberger (KSG, Kraskov et al., 2004). The Binned Estimator estimates the joint probability distribution using a 2D histogram of neural response vs. behavioral variable; this transforms continuous variables into discrete values. This estimator requires dense sampling of both neural activity and behavioral spaces and is dependent on the exact number of bins and binning method (Timme and Lapish, 2018). It has been used to estimate behavioral information contained in extracellularly recorded action potentials (García-Rosales et al., 2018; Linsenbardt et al., 2019), EEGs (Dauwels et al., 2010; Jeong et al., 2001; Min, 2003), and LFPs (Belitski et al., 2008). KSG estimates mutual information by examining the distance between data-points in the neural activity-behavioral parameter space. This technique is relatively more complex to use and has been benchmarked primarily on well-behaved, continuous random variables. The KSG estimator has been used to estimate behavioral information contained in continuous neural data, such as FMRI (Gómez-Verdejo et al., 2012), AP waveform shape (Chen et al., 2019), MEG (Robinson and Mandell, 2015), LFP (Manyakov and Van Hulle, 2008), and intracellular recordings (Sun et al., 2020; Wang et al., 2017).

The SMGM estimator, on the other hand, relies on the assumption that AP firing follows an inhomogeneous Poisson process. The SMGM estimator therefore requires binning of only the behavioral variable(s), in contrast to the Binned Estimator. The profile of firing rates vs. behavioral variable is then used to estimate the MI. The relative simplicity of the SMGM estimator has added to its popularity and widespread use in neuroscience applications for estimating behavioral information contained in single unit action potential recordings. This metric has proven useful in quantifying rate coding in place cells (Knierim et al., 1995; Lee et al., 2006; Markus et al., 1995; Poucet and Sargolini, 2013), complex spatial responses in neurons such as hippocampal interneurons (Frank et al., 2001; Wilent and Nitz, 2007), odor sequence cells (Allen et al., 2016), time cells (MacDonald et al., 2013), head direction cells (Stackman and Taube, 1998), speed cells (Fyhn et al., 2002), and face differential neurons (Nguyen et al., 2014, 2013), and has been used across multiple different species (Hazama and Tamura, 2019; Mankin et al., 2019; Yartsev and Ulanovsky, 2013). Furthermore, as a single neuron metric it provides statistical power for comparisons. Thus, it has been used to quantify differences in rate coding across different brain regions (Simonnet and Brecht, 2019) and has also been used to examine the effect on rate coding by experimental interventions such as lesions (Calton et al., 2003; Liu et al., 2004), inactivations (Brandon et al., 2011; Hok et al., 2013; Huang et al., 2009; Koenig et al., 2011), and applications of drugs (Newman et al., 2014; Robbe and Buzsáki, 2009). Further, it has been used to examine differences in encoding that arise during different behaviors (Aronov and Tank, 2014; Park et al., 2011; Zinyuk, 2000), and to demonstrate changes arising in disease models (Fu et al., 2017; Gerrard et al., 2008; Zhou et al., 2007). SMGM information is often normalized from measuring bits per unit time to measure bits per AP. This creates a measure sensitive only to the selectivity of a neuron, and not the average firing rate of the neuron.

The power of MI estimators has yet to be fully exploited by the neuroscience community. For example, the estimators have not yet been widely used to compare encoding properties of large numbers of genetically identified neurons, or to quantify information content of other discrete signaling events such as synaptic inputs; both of which are difficult to study using electrophysiological methods.

In vivo imaging of functional indicators has emerged as an important tool, largely because it possesses these capabilities. For example, using fluorescent calcium indicators, the functional properties of hundreds to tens of thousands of neurons can be simultaneously recorded in rodents (Dombeck et al., 2007;

Radvansky and Dombeck, 2018; Sheffield et al., 2017; Stirman et al., 2016; Stringer et al., 2019; Ziv et
al., 2013) or hundreds of thousands of neurons in zebrafish (Ahrens et al., 2013), as well as the whole
brain in invertebrates such as *C. elegans* (Nguyen et al., 2016) and *Drosophila* (Keller and Ahrens, 2015;
Mann et al., 2017). Furthermore, the genetic identity of the recorded neurons can be assured (Jing et al.,
2018a, 2018b, 2018c; Khoshkhoo et al., 2017; Sheffield et al., 2017) and subcellular structures are
accessible, allowing for functional recordings from synapses and dendrites using different functional
fluorescent indicators (e.g. Jing et al., 2018d; Marvin et al., 2019, 2018; Scholl et al., 2017; Sheffield et
al., 2017; Sheffield and Dombeck, 2015). However, functional fluorescence indicators generate signals
that are not a direct readout of the underlying quantal events. For example, somatic recordings from
neurons labeled with fluorescent calcium indicators reveal intensity variations that are correlated with
somatic AP firing rates but appear as a smoothed and varying amplitude version of the action potential
train. This transformation from action potential train to fluorescence trace is an active area of research
(Dana et al., 2018; Éltes et al., 2019; Greenberg et al., 2018), but it is often approximated by convolving
the action potential train with a kernel, which defines the indicator's response to a single action potential.
These kernels typically take the shape of a sharp increase in fluorescence followed by an exponential
decay to baseline (Chen et al., 2013; Dana et al., 2018; Pachitariu et al., 2018; Park et al., 2013; Yaksi and
Friedrich, 2006). Therefore, while functional fluorescence imaging and information theoretic
quantification may prove to be a powerful new combination of tools to study neural correlates of behavior,
it is critical to remember that functional fluorescence signals represent altered versions of the underlying
physiological events.

Caution is then needed when applying information metrics to continuous functional fluorescence traces,
yet the imaging community is already pushing forward and beginning to use information metrics,
particularly SMGM. This metric has already been applied to somatic calcium responses to compare the
information content of the same neurons across different behavioral epochs (Heys and Dombeck, 2018),
to quantify information across different populations of neurons in different brain regions (Hainmueller
and Bartos, 2018), to quantify information across different genetically identified neural populations
(Khoshkhoo et al., 2017), to examine encoding by subcellular structures (Rashid et al., 2020), and to
classify the significance of encoding particular parameters by individual neurons (Kinsky et al., 2018;
Mau et al., 2018; Rashid et al., 2020).

However, it is essential to recognize some of the assumptions underlying these information metrics are
violated by functional florescence recordings. All three metrics (SMGM, KSG and Binned Estimation)
assume stationarity in the neural response, which is violated by the elongated time responses and relatively
slow fluctuations of the fluorescence intensity of the reporters. When applied to spiking data, there is also
a change in units: rather than action potential counts, functional fluorescence traces are typically plotted
in units of florescence change with respect to the baseline ($\Delta F/F$). One possible solution to these issues
would be to deconvolve calcium traces to recover APs; however, deconvolution is an active area of
research, and the accuracy of these methods has recently been questioned (Evans et al., 2019). Ideally, the
calcium traces could be used directly to measure spiking information, without the need for such an in
between, potentially error introducing, step.

Quantifying the effects of the above violations on measurements of information using functional
fluorescence recordings using an analytical solution is particularly challenging with behaviorally
modulated neural recording data. However, a more tractable means of quantifying the effects would be to
use a simulation study to measure the induced biases and changes in measurement quality (Morris et al.,
2019). This strategy makes use of pseudo-randomly generated action potential traces and has the

advantage that the ground truth parameters of the simulations are known, while variability due to behavior
and other features can be incorporated. For example, this technique was previously used to examine
measures of theta phase precession, rhythmicity, neural correlation, and cellular dynamics (Climer et al.,
2015, 2013; Cohen and Kohn, 2011; Østergaard et al., 2018).

To provide the field with guidelines for the use of information metrics applied to functional fluorescence
recording data, we used computational simulation methods to create a library of ten thousand mock
neurons whose spiking output carry an exact, known (ground-truth) amount of information about the
animal's spatial location in its local environment. We used real behavioral data of spatial position over
time from mice navigating in virtual linear tracks and then simulated the spatial firing patterns of the mock
neurons using an inhomogeneous Poisson process framework (Brown et al., 2003; Climer et al., 2013;
Paninski, 2004). We then simulated fluorescent calcium responses for each neuron in each session by
convolving the action potential trains with calcium kernels for different indicators, primarily GCamp6f
(Chen et al., 2013), and then we added noise. MI metrics (between spatial location and the neural signals)
were then applied to the spiking or fluorescence traces to quantify the performance of the metrics for
estimating information. We focused on testing the performance of the SMGM method, and then compared
its performance to the Binned Estimation and KSG methods. We also applied a deconvolution algorithm
to test its performance. We found that the SMGM method applied directly to the mean $\Delta F/F$ intensity map
could best recover the ground truth information. We provide guidelines for the use of the SMGM metric
when applied to functional fluorescence recordings and demonstrate the appropriate application of these
guidelines to GCaMP6f population recordings from CA1 hippocampal pyramidal neurons in mice
navigating virtual linear tracks. From this CA1 analysis, we quantified the spatial information content of
large populations of pyramidal neurons and then performed Bayesian decoding of mouse position from
different information containing subsets of this large population. Interestingly, we found that the
population quantile with the lowest information values were still able to decode mouse position to the
closest quarter of the track. Thus, we provide new findings about the neural code for space that were made
possible by the information metrics and guidelines that we introduce here.

Results

The SMGM information metrics
Here we review the derivation of the SMGM information metrics and the underlying assumptions. For
illustrative purposes throughout this manuscript, we use the example of spatial encoding in which the
firing pattern of neurons carry information about the animal's location along a linear track; however, the
derivations, equations and conclusions generalize to encoded variables over other domains and
dimensionalities.

Consider a random variable X representing the positions an animal might take, with x being its value
measured at one time sample. The positions are subdivided into N spatial bins, such that x can take on the
values $\{1, 2, ..., N\}$. For our analyses, N=60. Consider a random variable Y representing the number of
action potentials a neuron might fire, where y is the count measured within a time sample. y can take on
the values of $\{0, 1, ..., +\infty\}$. X and Y are both discrete. If X and Y both obey the assumption that each time
sample is independent (i.e. they are stationary), then the mutual information (I, in bits per sample) between
X and Y is expressed as follows:

$$I(X;Y) = \sum_{i=1}^{N} \sum_{y=0}^{+\infty} p_{X,Y}(x_i, y) \log_2 \frac{p_{X,Y}(x_i, y)}{p_X(x_i)p_Y(y)} \qquad 1$$

172   Where $p_X(x_i)$ is the (marginal) probability that the animal is in the i^{th} spatial bin during a time sample,
173      $p_Y(y)$ is the probability that the neuron fires y action potentials in the time sample, and $p_{X,Y}(x_i, y)$ is the
174   joint probability that the neuron fires y action potentials and is in the i^{th} bin. Recall that $p_{X,Y}(x_i, y) =$
175      $p_{Y|X}(y|x_i)p_X(x_i)$, where $p_{Y|X}(y|x_i)$ is the conditional probability that the neuron fires y action potentials
176   given that the animal is in the i^{th} spatial bin. We can thus rewrite Equation 1 as follows:

$$I(X;Y) = \sum_{i=1}^{N} \sum_{y=0}^{+\infty} p_{Y|X}(y|x_i)p_X(x_i) \log_2 \frac{p_{Y|X}(y|x_i)}{p_Y(y)} \qquad 2$$

177   With the further assumption that the firing of the neuron follows Poisson statistics, we can then estimate
178   the mutual information as follows: let the action potential rate (action potential/s or Hz) in a single bin be
179   λ_i, and the average across the session be $\bar{\lambda}$. For an arbitrarily small time window Δt, the probability that
180   an action potential occurs in that window is $\Pr(Y = 1|x = i) = \lambda_i \Delta t$, with the probability that an action
181   potential occurs regardless of position as $\Pr(Y = 1) = \bar{\lambda}\Delta t$. We can thus rewrite Equation 2 as:

$$I(X;Y) = \sum_{i=1}^{N} \lambda_i \Delta t p_X(x_i) \log_2 \frac{\lambda_i}{\bar{\lambda}} \qquad 3$$

182   By integrating over one second ($\int_0^1 I(X;Y)d\Delta t$) we obtain the first key SMGM metric for spatial
183   information as measured by action potential firing, which is in units of bits per second:

$$\widehat{I_s^E} = \sum_{i=1}^{N} \lambda_i p_X(x_i) \log_2 \frac{\lambda_i}{\bar{\lambda}} \qquad 4$$

For notation, we will use a carrot (^) to indicate an information value that is measured from experiment,
the superscript (E in this case) to show the source of the data, and a subscript to show the units/formula
used (bits per second in this case). Thus, $\widehat{I_s^E}$ is the information measured via electrophysiology in bits per second.
This metric is linearly dependent on the average firing rate of the neuron, and this dependence is often
removed through normalization by the average firing rate to obtain the second key metric of spatial
information as measured by action potential firing, which is in units of bits/(second*Hz), or more
commonly, bits per action potential:

$$\widehat{I_{AP}^E} = \frac{1}{\bar{\lambda}} \sum_{i=1}^{N} \lambda_i p_X(x_i) \log_2 \frac{\lambda_i}{\bar{\lambda}} \qquad 5$$

Therefore, these two key metrics of spatial information are defined completely by quantities that can be
experimentally measured: the mean firing rate ($\bar{\lambda}$) from the action potential counts over the duration of
the recording, the action potential firing rate in the i^{th} bin from the average rate map (λ_i), and the

probability that the animal is in the i^{th} spatial bin from the normalized occupancy map ($p_X(x_i)$). The
quantity of and noise in these measurements effects the quality of the metric: in particular, undersampling
due to low firing rates or low trial counts induces a substantial positive bias (Treves and Panzeri, 1995).

In the derivation of these metrics, there are two key assumptions that are violated by functional
fluorescence recordings. First, the recordings do not follow Poisson statistics: instead of discrete counts
of action potentials (y), the functional fluorescence traces consists of a continuous relative change in
fluorescence ($\Delta F/F$), and instead of a firing rate map (λ_i) measured in Hz, average intensity maps in units
of $\Delta F/F$ are generated. The stationarity assumption is also violated: due to the slow decay, a time sample
of the fluorescence traces depend on the previous samples. The violation of these assumptions by
functional fluorescence recording will affect the precision and induce biases in the SMGM information
metrics. Since these effects have not previously been addressed or quantified, we measured these biases
here using a simulation study.

Figure 1. Procedures for generating a library of 10,000 neurons with known amounts of information. (A)
Five splines with a gradient of ground truth information (I_{AP}^E) representing the steps in generating a
continuous rate map ($\lambda(x)$) matching the desired target information, in this case 2 bits/action potential.
Red X's indicate control nodes that were moved to change the shape of the spline and minimize the squared
error to the target information. (B) Cross section of the error surface around the solution point as a

function of the position of node 3, and the trajectory taken by the solver to minimize the error and arrive
at the target. (C) Histograms of ground truth information resulting from repeating the procedure in A-B
10,000 times to target a range of ground truth information values in bits per second (I_S^E) and bits per
action potential (I_{AP}^E).(D) Splines representing $\lambda(x)$ at the solution point for a low (I_{AP}^E=0.04 bits per
action potential, left) and high (I_{AP}^E, 2 bits per action potential, right) information neuron. (E) Steps to
generate mock action potential and functional fluorescence data. (1) An example real behavior trace from
a mouse running on a linear track that was used to generate the simulated spiking. (2) The behavior in
combination with the rate maps generated in A-D were used to generate an instantaneous firing rate trace.
(3) The instantaneous rate was used to pseudorandomly generate action potentials, as shown in this mock
raster. (4) The action potential raster was convolved with the GCaMP6f kernel (red, inset) and noise was
added to generate a mock $\frac{\Delta F}{F}$ trace. (5) Large numbers of these traces were generated and used to assess
the effects of many simulation parameters on the estimators. (F-L) Spiking and fluorescence activity
patterns generated from the example simulated neurons shown in D and using a mean firing rate of 1 Hz.
(F) Behavioral trace in blue with action potential raster shown in red. (G) Lap-by-lap raster of the
neurons' firing vs mouse track position. (H) Lap by lap binned, firing rates vs mouse track position for
the neurons (I) Action potential raster (red) and mock calcium traces for the same behavioral period
shown in F. (J) Lap by lap mean binned fluorescence vs mouse position for the neurons. (K) Binned
average firing rate (λ_i, black) and fluorescence intensity (f_i, green) maps for the two neurons. These maps
were used for information analyses.

<u>*Building a ground truth library of 10,000 neurons with known values of information*</u>

To create a neuron with a known, ground truth information value, it was necessary to generate a continuous
(i.e. infinitesimally small bins) rate map ($\lambda(x)$) matching the desired information. To do this, we first
normalized the track length to 1 and assumed the animal's occupancy map to be spatially uniform
$\left(p_X(x_i) = \frac{1}{N}\right)$. We then created an exponentiated cubic spline with 5 randomly positioned nodes (Figure
1A) to build a starting continuous map of the normalized instantaneous firing rate, $\frac{\lambda}{\bar{\lambda}}(x)$, with the integral
normalized to 1. We calculated the ground truth amount of information in bits per action potential as
follows:

$$I_{AP}^E = \int_0^1 \frac{\lambda}{\bar{\lambda}}(x) \log_2\left(\frac{\lambda}{\bar{\lambda}}(x)\right) dx \qquad\qquad 6$$

The locations of the 5 nodes were then systematically varied [see Methods] to minimize the squared error
between the value calculated in Equation 6 and a target amount of information (Figure 1A,B), in the end
resulting in a mean error of 5.1*10^-9 bits/action potential and a mean absolute error of 1.5*10^-7
bits/action potential. The rate map at this convergence point was used for further analysis. This procedure
was repeated to generate 10,000 mock neurons with a range of (known and ground truth) information
values. Note that the value in Equation 5 cannot be higher than when all the action potentials arrive in one
spatial bin; the rate in that bin is $N\bar{\lambda}$. If we assume uniform occupancy $\left(p_X(x_i) = \frac{1}{N}\right)$, then the maximum
measureable information is $\log_2 N$, in our case, 5.9 bits/action potential with N=60 bins. Thus, the
information values considered here range between 0 and 6 bits/action potential (Figure 1C). We chose a

mean firing rate ($\bar{\lambda}$) for the neurons between 0.1 and 4 Hz, a range observed for a variety of different
cortical and hippocampal neurons during behavior (Buzsáki and Mizuseki, 2014; DeWeese et al., 2008;
O'Connor et al., 2010; Roxin et al., 2011; Shafi et al., 2007). From Equations 4-5, the ground truth
information in bits per second is $I_S^E = \bar{\lambda} I_{AP}^E$. I_S^E for these choices resulted in ground truth information
values between $I_S^E = 0$ and $I_S^E = 24$ bits/sec (Figure 1C). Example low ($I_{AP}^E = 0.04$ bits/action potential) and
mid ($I_{AP}^E = 2$ bits/action potential) rate maps are shown in Figure 1D.

These rate maps provided a basis for generating mock action potential firing data (and functional
fluorescence data, see below). Under real experimental conditions, recording duration and bin sizes are
finite and animal occupancy maps ($p_X(x_i)$) are not spatially uniform. These experimental limitations add
error to the estimate of a neuron's ground truth information value. Therefore, in order to accurately re-
create these limitations in our simulation study, we used real behavior datasets from head-restrained mice
running along a 3 m virtual linear track for water rewards (acquired as in Sheffield et al., 2017; Sheffield
and Dombeck, 2015). Unless otherwise indicated, all values reported will be the mean±standard
deviation. We selected at random from a library of 574 behavior sessions from mice navigating along
familiar tracks, and concatenated and truncated these sessions to create behavior sessions uniformly
sampled up to 60 minutes in duration (average 30.2±17.1 minutes), resulting in an average 132 ± 71.2
laps per session and an average running speed of 19.3 + 3.87 cm/sec (Figure 1E.1). This behavior, the
average firing rate ($\bar{\lambda}$), and the normalized rate map ($\frac{\lambda}{\bar{\lambda}}(x)$) from the mock neurons were used to create an
instantaneous firing rate trace (Figure 1E.2), sampled at 1 kHz, from which action potential times were
generated assuming Poisson firing statistics (Figure 1E.3). Example mock spiking in response to behavior
for low (0.04 bits/action potential) and mid (2 bits/action potential) information neurons can be seen in
Figure 1F-H. From these spiking responses, we then generated mock fluorescence traces by convolving
the raster with a double-exponential kernel matching the rise and fall times for GCaMP6f (Chen et al.,
2013, Figure 1E.4) and adding random noise to model shot noise. Mock fluorescence traces for the two
example neurons in Figure 1F-H can be seen in Figure 1I-J. The mock action potential and fluorescence
traces were used to create session mean spatial maps – of binned firing rate (λ_i in Hz) and change in
fluorescence (f_i in $\Delta F/F$), for information analyses (Figure 1K). By repeating this process, we built a
large dataset of spiking and fluorescence traces, generated from our library of mock neurons with known
amounts of information and using real animal spatial behavior. With tens of thousands of these mock
neuron recordings, we could then assess the effects of many simulation parameters on the information
values determined from the metrics - including firing rate, session duration, fluorescence kernel shape,
and ground truth information value.

Figure 2. Quantification of the precision of the SMGM bits per second metric using action potentials or functional fluorescence recordings. (A) Three representative mock neurons spanning the range of ground truth information values in bits per second (I_S^E). From top to bottom for each: mouse track position vs time, action potential raster, fluorescence calcium trace (green), and firing rate map (λ_i, black) and change in fluorescence map (f_i, green). (B-D) The ground truth bits per second values are well recovered when measured from action potential traces. (B) Information measured from action potential data using the SMGM bits per second metric ($\widehat{I_S^E}$) vs ground truth information (I_S^E). Each dot is a single mock neuron, the gray dashed line is the unity line (perfect measurement), the pink line is the line of best fit. Red circles show the examples in A. (C) Percentage error for the information measurements shown in B. (D) Heat

map of percentage error measurements shown in C. Black lines are 2 standard deviations, the white line
is the mean. (E-G) Effects of applying the SMGM bits per second metric to fluorescence traces. (E)
Information measured from mock GCaMP6f traces using the SMGM bits per second metric ($\widehat{I_s^F}$) vs ground
truth information (I_s^E). (F) Percentage error for the information measurements shown in E. (G) Heat map
of percentage error measurements shown in F. (H) Representative mock kernels mimicking responses
from different indicators. (I-K) The effect of kernel height on estimating ground truth information (I_s^E)
using the SMGM bits per second metric ($\widehat{I_s^F}$). Kernel height for the kernels shown in H are indicated by
colored triangles (I) Percentage error as a function of kernel height (J) Heat map of percentage error
measurements shown in I with mean (white) and 2 standard deviations (black). (K) The average
percentage error as a function of kernel height and ground truth information in SMGM bits per second
(I_s^E). (L-N) The effect of kernel width on estimating ground truth information (I_s^E) using the SMGM bits
per second metric ($\widehat{I_s^F}$). Kernel widths for the kernels shown in H are indicated by colored triangles. (L)
Percentage error as a function of kernel width (M) Heat map of percentage error measurements shown
in L with mean (white) and 2 standard deviations (black). (N) The average percentage error as a function
of kernel width.

Quantification of the accuracy and precision of the SMGM bits per second metric using functional
fluorescence recordings
We first applied the SMGM bits per second metric ($\widehat{I_s^E}$) to our mock action potential recording traces to
verify that they can recover our ground-truth information values given finite recording durations and bin
sizes, and non-uniform animal occupancy maps ($p_X(x_i)$). Figure 2A shows three mock neurons with
ground truth information values of I_s^E =3, 15 and 23 bits/sec. When the SMGM bits per second metric
($\widehat{I_s^E}$) was applied to the action potential traces from these example neurons, the information was well
recovered, with $\widehat{I_s^E}$=2.8, 15 and 24 bits/sec, respectively. The results from these examples also held across
the full 10,000 mock neuron library (Figures 2B-D), as a linear fit (y-intercept = 0.093$\pm$0.040, intercept
p=4.6*10^{-6} bits per second and slope = 0.97 $\pm$0.0030, slope p<<0.01) explained nearly all the variance
(R^2 = 0.97), the average error was 0.22$\pm$1.25 bits per second (1.0$\pm$0.69% error) and the absolute error
was 0.64$\pm$1.05 bits per second (8.4$\pm$0.69% error). There is a substantial positive bias for the lowest firing
rates and smallest number of trials (Supplemental Figure 1A-B) which has been previously well
characterized (Treves and Panzeri, 1995). Thus, the SMGM bits-per-second metric ($\widehat{I_s^E}$) recovers the
ground-truth information well using action potential recordings, with the only error coming from finite
recording time and variable animal behavior.

We next discuss the changes to the SMGM bits per second metric ($\widehat{I_s^E}$) commonly used for application to
functional fluorescence traces (Hainmueller and Bartos, 2018; Heys and Dombeck, 2018), and explore the
implications of these changes. Most simply, the mean firing rate ($\bar{\lambda}$) and the mean firing rate in a spatial
bin (λ_i) are replaced by the mean change in fluorescence ($\bar{f}$) and the mean change in fluorescence in a
bin (f_i). Making these substitutions in Equation 4 results in the information as measured by functional
fluorescence:

$$\widehat{I_s^F} = \sum_{i=1}^{N} f_i p_X(x_i) \log_2 \frac{f_i}{\bar{f}} \qquad\qquad 7$$

The fluorescence map f_i differs from the firing rate map λ_i in two ways. First, the fluorescence map is
approximated by the firing rate map scaled by a factor c, dependent on the height and width of the kernel

and measured in units of $\frac{\Delta F/F}{Hz}$, and second it is smoothed by the kernel (Figure 1E4). If we discount the
latter for a moment and focus on the scaling, $f \approx c\lambda$, we can see that substituting λ with $c\lambda$ in Equation 4
results in $\widehat{I_s^F} = c\widehat{I_s^E}$. The units for $\widehat{I_s^F}$ are no longer in bits per second, as it has previously been reported
(Hainmueller and Bartos, 2018), but are instead in units of $\frac{bits\ \Delta F/F}{sec\ Hz}$ or $\frac{bits\ \Delta F/F}{AP}$, which are difficult to
interpret (see *Guidelines for application of information theoretic metrics to functional fluorescence*
imaging data section for further discussion). The effect of smoothing is difficult to analytically quantify
since it both alters c by changing the average intensity and distorts the firing rate map. Therefore, to fully
quantify the impact of convolving an action potential recording with a functional fluorescence kernel on
recovering ground truth information, we used our mock fluorescence traces.

We applied a GCaMP6f modeled kernel to the 10,000 mock action potential traces to generate 10,000
mock fluorescence calcium traces. Figure 2A shows the fluorescence traces generated from three mock
neurons with ground truth information values of I_s^E =3, 15 and 23 bits/sec. The effects of the convolution
can be seen in the differences in scaling and shape between the fluorescence maps f_i and the firing rate
maps λ_i. When the fluorescence metric ($\widehat{I_s^F}$) was applied to the fluorescence traces from these example
neurons, the information recovered was $\widehat{I_s^F}$ = 3.8, 14 and 34 $\frac{bits\ \Delta F/F}{AP}$, respectively, indicating some
deviations from the ground truth information values assuming the units are comparable. The results from
these examples also held across the full 10,000 mock neuron library (Figures 2E-G), as there was a clear
scaling of the ground truth information and a consistent overestimation with a mean error of 2.0±3.28 AU
(19±30% error) and mean absolute error of 2.4±2.97 AU (22±27% error). The best-fit line of the
measured information ($\widehat{I_s^F}$) versus the ground truth information (I_s^E) had an intercept near 0 (0.088±0.048
$\frac{bits\ \Delta F/F}{AP}$, p=0.07). The slope of this fit was 1.17±0.0036 $\frac{\Delta F/F}{Hz}$ (p<<0.01), which provides a measure of the
scaling factor (c). This error was not corrected for with denser sampling: it remained consistent even at
high firing rates and many trials (Supplemental Figure 1C-D). In addition to this scaling effect caused by
c, smoothing of the rate map could induce nonlinearity in the relationship between $\widehat{I_s^F}$ and I_s^E. To test for
such an effect, we fit the measured information in Figure 2E with a saturating exponential and compared
the fits using a likelihood ratio test: the exponential did not significantly improve the fit (χ_1^2=2.8, p=0.095),
which indicates that smoothing by the kernel does not induce significant nonlinearities. Since c is
dependent on the height and width of the kernel and was measured here as 1.17±0.0036 $\frac{\Delta F/F}{Hz}$, then
considering the 0.19 $\frac{\Delta F}{F}$ height of the GCaMP6f kernel used here, we estimated the effect of the kernel
width on c as 6.1 s^{-1}. The consistent, positive bias observed in estimating information with $\widehat{I_s^F}$ (Figure 2E)
would be easy to correct for assuming the c factor, and therefore the kernel, were similar across all
measured neurons. This point is considered further in the *Guidelines for application of information metrics*
to functional fluorescence imaging data section below. We conclude that ground truth information, as
measured by the fluorescence SMGM bits per second metric ($\widehat{I_s^F}$), is transformed into different units and
is linearly scaled by a factor c dependent on the height and width of the kernel.

The amplitude (height) of the change in fluorescence can vary across indicators. The height of the kernel,
given a constant kernel width, should linearly scale c and the error in estimating information with $\widehat{I_s^F}$. To
explicitly test this prediction, we simulated an additional 5,000 fluorescence traces with kernels of varying
height (0-3 $\Delta F/F$, Figure 2I-K), but maintaining the same shape and width (from the GCaMP6f kernel),

and then measured the percent error in estimating information with $\widehat{I_s^F}$. As observed above for the
GCaMP6f example (Figure 2 E-G), the percent error in estimating information with $\widehat{I_s^F}$ shows little
dependence on ground truth information (Figure 2I-K). As a function of the height of the kernel, however,
the percent error (averaged over all ground truth information values) in estimating information with $\widehat{I_s^F}$ is
fit well with an increasing linear function (intercept = -92.5$\pm$11.7%, intercept p=3.6*10-^{15}, slope 619
$\pm$4.1%/$\frac{\Delta F}{F}$, slope p<<0.01, R^2= 0.81; Figure 2I-J). Over the wide array of available functional fluorescent
indicators in use today (Figure 2H), this leads to differences in error due to differences in transient height
alone of the indicator used. For the indicators shown in Figure 2J, there is an average height of 0.603$\pm$0.10
SD: the error spans from 25% for the kernel height reported for gCamp6f (0.19 $\Delta F/F$) to 254% for
gCamp7f (0.56 $\Delta F/F$). It should be noted that fluorescence ($\Delta F/F$) is always reported here as a fractional
change, not as a percentage (% $\Delta F/F$); if a kernel height of 19 % $\Delta F/F$ is used, the units would again
change. Thus, as expected, the percent error in estimating information with the SMGM bits per second
estimator ($\widehat{I_s^F}$) scales linearly with the height of the kernel.

The width of the kernel can vary widely across fluorescent indicators (Figure 2H), with "faster" indicators
boasting shorter rise and fall times. The combined effect of a longer rise and fall time is to smooth and
delay the action potential train; in other words, it acts as a causal low-pass filter. The cutoff period of this
low pass filter provides a measurement of the effective width of the kernel [see Methods]. The effect of
such differences in kernel shape on the error in estimating information with $\widehat{I_s^F}$ is difficult to measure
analytically. We therefore simulated an additional 5,000 fluorescence traces with kernels of different
kernel widths (but constant height of the GCaMP6f kernel), resulting in a range of kernel durations (rise
times: 1 ms to 1 second, fall times longer than the rise time up to 2 seconds), and then measured the percent
error in estimating information with $\widehat{I_s^F}$. Similarly, as observed above for the GCaMP6f and varying kernel
height examples (Figure 2 E-K), the percent error in estimating information with $\widehat{I_s^F}$ shows little
dependence on ground truth information (Figure 2L-N). Interestingly, however, the percent error
(averaged over all ground truth information values) in estimating information with $\widehat{I_s^F}$ shows a complex
nonlinear response as a function of the width of the kernel (Figure 2L-M). The error increases up to a
kernel width of ~2.5 seconds, at which point it saturates at ~250% error. This arises between a complex
interaction between changing the average value of the original action potential trace and flattening the
average fluorescence map (f_i). Over the wide array of available functional fluorescent indicators in use
today, this leads to large differences in error due to differences in width alone of the indicator used. For
example, an average error of -10.6$\pm$1.8% and absolute error of 20.0$\pm$1.0% were observed for iGluSnfR
the shortest indicator considered here at 0.52 seconds. For gCamp6s, the slowest indicator examined (2.54
seconds), the average was 235$\pm$18.9% and absolute error was 235$\pm$18.9%. To estimate the percent errors
for these five indicators considering differences in *both* height and duration, we used these 5 kernels to
generate mock fluorescence traces from the 10,000 neurons in Figure 2B-G. The resulting distributions,
estimated c values, and mean and absolute errors can be seen in Supplemental Figure 2. In summary, we
conclude that information, as measured by the fluorescence SMGM bits per second metric ($\widehat{I_s^F}$), is
transformed into different units and is linearly scaled by a factor (c) dependent on the height and width of
the kernel, with c linearly dependent on height and nonlinearly dependent on width. The error induced by
these transformations changes substantially over the range of kernel values of the different functional
indicators widely used today, and therefore are important factors to consider when designing and
interpreting functional imaging experiments (see *Guidelines for application of information metrics to*
functional fluorescence imaging data for further discussion).

Figure 3. Quantification of the precision of the SMGM bits per action potential metric using action potentials or functional fluorescence recordings. (A) Three representative mock neurons spanning the range of ground truth information values in bits per action potential (I_{AP}^E). From top to bottom for each: mouse track position vs time, action potential raster, fluorescence calcium trace (green), and firing rate map (λ_i, black) and change in fluorescence map (f_i, green). (B-D) The ground truth bits per action potential values are well recovered when measured from action potential traces. (B) Information measured from action potential data using the SMGM bits per action potential metric ($\widehat{I_{AP}^E}$) vs ground truth information (I_{AP}^E). Each dot is a single mock neuron, the gray dashed line is the unity line (perfect measurement). Red circles show the examples in A. (C) Percentage error for the information measurements shown in B. (D) Heat map of percentage error measurements shown in C. Black lines are 2 standard deviations, the white line is the mean. (E-G) Effects of applying the SMGM bits per action potential metric to fluorescence traces. (E) Information measured from mock GCaMP6f traces using the

SMGM bits per action potential metric ($\widehat{I^F_{AP}}$) vs ground truth information (I^E_{AP}). (F) Percentage error for
the information measurements shown in E. (G) Heat map of percentage error measurements shown in F.
(H-J) The effect of kernel height on estimating ground truth information (I^E_{AP}) using the SMGM bits per
second metric ($\widehat{I^F_{AP}}$). Kernel height for the kernels shown in Figure2H are indicated by colored triangles
(H) Percentage error as a function of kernel height (I) Heat map of percentage error measurements shown
in H with mean (white) and 2 standard deviations (black). (J) The average percentage error as a function
of kernel height and ground truth information in bits per action potential (I^E_{AP}). (K-M) The effect of kernel
width on estimating ground truth information (I^E_{AP}) using the SMGM bits per action potential metric ($\widehat{I^F_{AP}}$).
Kernel widths for the kernels shown in Figure 2H are indicated by colored triangles. (L) Percentage error
as a function of kernel width (M) Heat map of percentage error measurements shown in L with mean
(white) and 2 standard deviations (black). (N) The average percentage error as a function of kernel width.

Quantification of the accuracy and precision of the SMGM bits per action potential metric using
functional fluorescence recordings

We then applied the SMGM bits per action potential metric ($\widehat{I^E_{AP}}$) to our mock action potential recording
traces to verify that they can recover our ground-truth information values. Figure 3A shows three mock
neurons with ground truth information values I^E_{AP} =0.05, 1.8 and 4.2 bits/action potential. When the
SMGM bits per action potential metric ($\widehat{I^E_{AP}}$) was applied to the action potential traces from these example
neurons, the information was well recovered, with $\widehat{I^E_{AP}}$=0.06, 1.8 and 4.2 bits/action potential respectively.
The results from these examples also held across the full 10,000 mock neuron library (Figures 3B-D), as
a linear fit (y-intercept = 0.087±0.029, intercept p=2.8e-184 bits per second and slope = 0.93 ±0.0010,
slope p<<0.01) explained nearly all the variance (R^2 = 0.99), the average error was -0.071±0.23 bits per
AP (3.2±5.9% error) and the absolute error was 0.13±0.21 bits per second (8.1±9% error). However, the
data was even better fit with a saturating exponential (χ^2_1=1.6e3, p<<0.01) converging to 5.8 bits/action
potential as it approached the limit due to the finite bin count. There is a substantial positive bias for the
lowest firing rates and smallest number of trials (Supplemental Figure 1E-F) which has been previously
well characterized (Treves and Panzeri, 1995). Thus, the SMGM bits per-AP metric ($\widehat{I^E_{AP}}$) recovers the
ground-truth information well using action potential recordings (except at the largest ground truth
information values), with the primary error coming from finite recording time and variable animal
behavior.

We next discuss the changes needed to apply the SMGM bits per action potential metric ($\widehat{I^E_{AP}}$) to functional
fluorescence traces and explore the implications of these changes. Most simply, the mean firing rate ($\bar{\lambda}$)
and the mean firing rate in a spatial bin (λ_i) are replaced by the mean change in fluorescence ($\bar{f}$) and the
mean change in fluorescence in a bin (f_i). Making these substitutions in Equation 5 results in the
information as measured by functional fluorescence:

$$\widehat{I^F_{AP}} = \frac{1}{\bar{f}}\sum_{i=1}^{N} f_i p_X(x_i) \log_2 \frac{f_i}{\bar{f}} \qquad\qquad 8$$

As discussed above, the fluorescence map (f_i) can be approximated as a scaled version of the rate, that is,
$f = c\lambda$ and $\bar{f} = c\bar{\lambda}$. Thus, under this approximation, the c factors in Equation 8 cancel, leading to $\widehat{I^F_{AP}}$
equivalent to $\widehat{I^E_{AP}}$, with the same units of bits/action potential. This, of course, ignores the fact that the
kernel smooths the rate map, leading to a bias in the metric that is difficult to quantify analytically.

We then applied the fluorescence SMGM bits per action potential metric ($\widehat{I_{AP}^F}$) to our 10,000 mock
GCaMP6f traces. Figure 3A shows the fluorescence traces generated from three mock neurons with
ground truth information of I_{AP}^E=0.05, 1.8 and 4.2 bits/action potential. When the fluorescence metric ($\widehat{I_{AP}^F}$)
was applied to the fluorescence traces in these examples, the information recovered was 0.04, 1.8 and 3.5
bits/action potential, indicating some deviations – especially for the highest information neuron. These
results held for the 10,000 mock neuron library (Figure3E-G). At low information values, there was little
bias, but at higher information values the information recovered was substantially lower than the ground
truth information. The mean resulting error was -0.38$\pm$0.58 bits/action potential (-9.7$\pm$27.8%) and
absolute error of 0.39 (12.9$\pm$26.4%). This error was better fit with a saturating exponential than a linear
fit (χ_1^2=1.6e3, p<<0.01), with the average error less than 5% up to ground truth information of 1.8
bits/action potential and less than 10% up to 3.0 bits/action potential. At ground truth information values
higher than 3 bits/action potential, the average error was -1.06$\pm$0.595 (-22.5$\pm$9.44%) and absolute error
was 1.07$\pm$0.589 bits/action potential (22.6$\pm$9.21%). This error persisted even with denser sampling: it
remained consistent even at high firing rates and many trials (Supplemental Figure 1E-F). Thus, the
indicator induces relatively little error at lower information values (<3 bits/action potential), but the
smoothing effect of the kernel induces a nonlinear, negative bias to the estimator, particularly at ground
truth information values over 3 bits/action potential.

Although the height of the kernel can vary between different functional fluorescence indicators (Figure
2H), these height variations linearly scale the fluorescence map. Thus, since $\widehat{I_{AP}^F}$ involves normalization
by the mean change in fluorescence ($\bar{f}$), $\widehat{I_{AP}^F}$ should not depend on kernel height. To explicitly test this
prediction, we used the 5,000 fluorescence traces described in the previous section (Quantification of the
accuracy and precision of the SMGM bits per second metric using functional fluorescence recordings),
with kernels of varying height (0-3 ΔF/F), but maintaining the same shape and width (from the GCaMP6f
kernel), and then measured the percent error in estimating information with $\widehat{I_{AP}^F}$ (Figure 3H-J). Unlike for
the SMGM bits per second metric, the percent error (averaged over all ground truth information values)
in estimating information with $\widehat{I_{AP}^F}$ shows little or no dependence on the height of the kernel (p=0.43), but
a nonlinear dependence on ground truth information as in Figure 3E-G, with no significant difference in
the parameters of the saturating exponential fit (χ_2^2=1.67, p=0.43). Thus, as expected, the percent error in
estimating information with the SMGM bits per action potential metric ($\widehat{I_{AP}^F}$) does not vary with the height
of the kernel.

With little effect of kernel height on $\widehat{I_{AP}^F}$, the width of the kernel likely drives biases in the metric. We thus
used the 5,000 fluorescence traces generated from a range of different kernel durations (rise times: 1 ms
to 1 second, fall times longer than the rise time up to 2 seconds; but constant height of the GCaMP6f
kernel) from the previous section (*Quantification of the accuracy and precision of the SMGM bits per*
second metric using functional fluorescence recordings), and then measured the percent error in
estimating information with $\widehat{I_{AP}^F}$ (Figure 3 K-M). Similarly, as observed above for GCaMP6f and the
varying kernel height examples (Figure 3E-J), the percent error in estimating information with $\widehat{I_S^F}$ shows
a nonlinear dependence on ground truth information (Figure 3E-G). The percent error (averaged over all
ground truth information values) showed a nonlinear response as a function of the width of the kernel
(Figure 3K-L), with a steep increase in error for kernel widths >~1 second. Even for kernel widths <~1
second, the percent error was strongly dependent on the ground truth information value, with steep
increases in error for values >~2.5-3 bits/action potential (Figure 3M). Thus, as the kernels gets wider,

there is more negative bias at lower and lower information measured. The resulting errors are thus larger
for wider kernel indicators: for example, with a kernel width the same as gCaMP6s (2.54s), the error
exceeds -17% even at low (<0.25 bits/action potential) information, with average errors of -0.86 ± 1.0
bits/action potential ($-31\pm19\%$ error) and absolute errors of 0.87 ± 1.0 bits/action potential ($32.6\pm16\%$
error). In contrast, with a kernel width the same as iGluSnfR (0.52 seconds), the average error exceeded
5% at 3 bits per action potential and 10% at 3.7 bits per action potential with a mean error of -0.57 ± 1.00
($-8.0\pm14\%$) bits/action potential and absolute error of 0.41 ± 0.56 bits/action potential ($11\pm11\%$). To
estimate the percent errors for the five indicators shown in Figure 2H, taking into account differences in
both height and duration, we used the 5 kernels to generate mock fluorescence traces from the 10k neurons
in Figure3B-G. The resulting distributions, mean and absolute errors, and error thresholds can be seen in
Supplemental Figure 3.

In summary, we conclude that ground truth information, as measured by the fluorescence SMGM bits per
action potential metric ($\widehat{I_{AP}^{F}}$), retains the units and insensitivity to height scaling of the electrophysiological
metric ($\widehat{I_{AP}^{E}}$), but is nonlinearly biased by the smoothing of the fluorescence map dictated by the width of
the kernel. The estimation errors strongly depended on both the width of the kernel and the information
value being measured. Since these parameters change substantially over the different functional indicators
and different neuron types and behaviors that are commonly used today, they are important factors to
consider when designing and interpreting functional imaging experiments (see below for further
discussion).

Deconvolution may not be sufficient to eliminate biases
The framework presented here for comparing ground truth information with information measured with
the SMGM metrics can be extended to test the efficacy of other strategies for extracting mutual
information. In particular, a perfect AP inference method would alleviate the problems associated with
applying the SMGM metrics to functional fluorescence recordings. To test the utility of such a strategy in
measuring information, we applied a popular deconvolution algorithm, FOOPSI (Vogelstein et al., 2010,
see Methods), to the same 10,000 mock GCaMP6f time-series traces described above. Importantly, this
deconvolution algorithm (and other available algorithms) does not recover traces of relative spike
probability or exact spikes times, but instead produces sparse traces with arbitrary units, that have non-
zero values estimating the relative 'intensity' of spike production over time (d). This signal can be thought
of as a scaled estimate of the number of spikes per time bin, and thus the average intensity map will have
some similar properties to the florescence intensity maps – that is, we would expect the intensity maps
from deconvolution to approximate the relative firing rate scaled by some factor c, which has arbitrary
units.

We then measured information in these deconvolved d-traces using the SMGM metrics ($\widehat{I_{s}^{d}}$ and $\widehat{I_{AP}^{d}}$),
which are identical to ($\widehat{I_{s}^{F}}$ and $\widehat{I_{AP}^{F}}$), except SMGM is applied to d-traces instead of functional fluorescence
traces. When using the SMGM bits per second measure ($\widehat{I_{s}^{d}}$, Figure 4A), we found a clear scaling of the
ground truth information. The scaling factor was very small ($c=0.0346\pm5.5e\text{-}4$ A.U.), resulting in low
predicted information (mean error -11.0 ± 6.78 AU, mean absolute error 11.0 ± 6.76 AU, mean % error -
95.0 ± 11.5 %, mean absolute % error $95.0\pm6.87\%$). This error was significantly larger than when we
measured information directly from the florescence traces using $\widehat{I_{s}^{F}}$ (Figure 2E; 22.0% absolute error,
Ranksum $p\ll0.01$; c=1.17).

Assuming that the intensity map of the deconvolved d-traces are a scaled version of the true rate maps,
we could measure information using the SMGM bits per AP metric $\widehat{I_{AP}^d}$ without changing units (Figure
4B). Compared to the SMGM bits per AP metric applied to florescence (I_{AP}^F), on average there was some
reduction in the nonlinearity at higher ground truth information (I_{AP}^E) values when using $\widehat{I_{AP}^d}$, resulting in
linear fits closer to the unitary line ($\widehat{I_{AP}^d}$ slope $= 1.02 \pm 0.0027, R^2 = 0.76$ versus $\widehat{I_{AP}^F}$ slope $=$
0.78 ± 0.0011, $R^2 = 0.93$). However, information measured with $\widehat{I_{AP}^d}$ was still better fit with a saturating
exponential ($\chi_1^2 = 2.3e3$, p~0) converging to a saturation value of 5.51 bits per AP (compared to 5.78 for
$\widehat{I_{AP}^F}$). This resulted in a positive bias at lower levels of ground truth information. For ground truth
information values below 3 bits per AP, the average error for $\widehat{I_{AP}^d}$ was 0.556 ± 0.50 bits per AP ($68.9 \pm 104\%$)
compared to -0.060 ± 0.13 bits per AP ($2.75 \pm 0.36\%$) for $\widehat{I_{AP}^F}$. For ground truth information values above 3
bits per AP, the average error for $\widehat{I_{AP}^d}$ was -0.127 ± 0.76 bits per AP ($-1.0 \pm 16.8\%$) as compared to -1.04 ± 0.59
bits per AP ($-0.22 \pm 9.4\%$) for $\widehat{I_{AP}^F}$. Overall, there was more error when the SMGM bits per AP metric was
applied to deconvolved data compared to when applied directly to fluorescence traces ($\widehat{I_{AP}^d}$ mean absolute
error 0.60 ± 0.47 bits/AP ($52.7 \pm 89.1\%$) vs $\widehat{I_{AP}^F}$ was 0.13 ± 0.21 bits per second ($8.1 \pm 9\%$ error), Ranksum
p<<0.01). Thus, when comparing the recovery of ground truth information from functional fluorescence
traces using either direct application of the SMGM metrics ($\widehat{I_s^F}$ and $\widehat{I_{AP}^F}$) or the application of the SMGM
metrics to deconvolved z-traces ($\widehat{I_s^d}$ and $\widehat{I_{AP}^d}$), we found better recovery using the direct application
approach ($\widehat{I_s^F}$ and $\widehat{I_{AP}^F}$).

The KSG and Binned Estimators are poor estimators of MI in functional florescence data
In addition to SMGM, the KSG and Binned Estimation metrics have been developed for estimating mutual
information between variables. These other two metrics produce information measured in bits per second,
so they are only comparable to the SMGM bits per second estimator ($\widehat{I_s^F}$). The KSG metric uses the kth
nearest neighbor distances between points in the neural response and behavioral variable space to estimate
information (Kraskov et al., 2004). The Binned Estimation metric uses discrete bins to estimate the full
multidimensional joint probability distribution ($p_{(X,Y)}$ in Equation 1) to estimate mutual information
(Timme and Lapish, 2018). These two metrics estimate information across time samples and therefore are
dependent on firing rate like the SMGM bits/sec metric considered above. Further, the Binned Estimator
is sensitive to the precise method used for data binning, and thus we have used two commonly applied
binning methods: uniform and occupancy based bins.

We applied the KSG, Binned Estimator (Uniform bins), and Binned Estimator (Occupancy binned) (see
methods) to the same 10,000 mock GCaMP6f time-series traces and behavioral data used to assess the
SMGM approach. These methods all behaved similarly when applied to our simulations (Figure 4C-E),
so they will be discussed together here. The information values measured by these techniques correlated
with ground truth information in bits per second (I_s^E). Interestingly, however, unlike the SMGM bits per
second metric ($\widehat{I_s^F}$, Figure 2), the KSG and Binned Estimator results were better fit with a saturating
exponential than with a linear fit ($\chi_1^2 = 1.70e4$, 3.12e4, and 3.14e4 respectively, p~0). The KSG and Binned
Estimator methods overestimated the information at lower ground truth (I_s^E) values and saturated quickly
at higher values. For ground truth information (I_s^E) values below 10 bits per second, the mean absolute
errors were 10.3 ± 7.82 bits per second ($465 \pm 4,390\%$), 13.9 ± 8.33 bits per second ($539 \pm 2,324\%$), and

15.3$\pm$9.33 bits per second (600$\pm$2,721%) for the KSG, Binned Estimator (Uniform bins), and Binned

Estimator (Occupancy binned) respectively. This is in comparison to the 0.35$\pm$0.59 $\frac{bits^{\Delta F}/_F}{AP}$ (14$\pm$103%)

mean absolute error found using the SMGM bits per second metric ($\widehat{I_s^F}$) for ground truth information

values less than 10 bits per second. For ground truth information (I_s^E) values greater than 10 bits per

second, the mean absolute errors were 13.2$\pm$8.66 bits per second (85.0$\pm$64.4%), 14.3$\pm$9.46 bits per

second (93.3$\pm$72.0%) and 15.8$\pm$10.93 bits per second (103$\pm$80.2%) for the KSG, Binned Estimator

(Uniform bins), and Binned Estimator (Occupancy binned) respectively. This is in comparison to the

0.97$\pm$1.15 bits per second (5.85$\pm$6.73%) mean absolute errors found using the SMGM bits per second

metric ($\widehat{I_s^F}$) for ground truth information values greater than 10 bits per second. Over the full range of

ground truth values, mean absolute errors of 11.9$\pm$8.42 bits per second (254.8%), 14.1$\pm$8.94 bits per

second (292.5%) and 15.6$\pm$10.2 bits per second (325.5%) were found for the KSG, Binned Estimator

(Uniform bins), and Binned Estimator (Occupancy binned) respectively – an order magnitude larger than

the 2.4$\pm$2.97 AU (22$\pm$27% error) error seen using the SMGM bits per second metric ($\widehat{I_s^F}$). Therefore,

when comparing the recovery of ground truth information from functional fluorescence traces using either

the SMGM metric $\widehat{I_s^F}$ or the KSG and Binned Estimator metrics, we found better recovery using the

SMGM approach ($\widehat{I_s^F}$).

Method	Units	Source	Mean error	Mean abs error	Mean % error	Mean abs % err	C	p exponential	saturation level	90% saturation point
SMGM bits/sec	bits ΔF/F/AP	Skaggs et al, 1993	2.0	2.4	19.0	22.0	1.17	0.095	91	155.2
SMGM bits/AP	bits/AP	Skaggs et al, 1993	-0.38	0.4	-9.7	12.9	0.78	<<0.01	5.78	11.5
foopsi+SMGM bits/sec	AU	Vogelstein et al, 2010	-11.0	11.0	-95.0	95.5	0.0346	3.24E-08	0.84	31.93
Foopsi+SMGM bits/ap	bits/AP	Vogelstein et al, 2010	0.3	0.6	46.3	52.7	1.02	<<0.01	5.51	6.6
KSG	bits/sec	Kraskov et al, 2004	9.9	11.9	37.9	254.8	1.64	<<0.01	29.2	13.9
Binned MI (uniform bins)	bits/sec	Timm and Lapish, 2018	12.8	14.1	285.7	292.5	1.77	<<0.01	29	8.47
Binned MI (occupancy bins)	bits/sec	Timm and Lapish, 2018	15.0	15.6	322.5	325.5	1.94	<<0.01	32.2	9.26

Figure 4. Alternative techniques for measuring mutual information from functional fluorescence traces. (A-E) (Top) Information measured from mock GCaMP6f traces vs ground truth information. The gray line is the unity line, the pink line is the best fit saturating exponential. (Middle) Percentage error for the information measurements shown on top. (Bottom) Heat map of percentage error measurements shown in middle. (A) FOOPSI deconvolved traces using the SMGM bits per second metric ($\widehat{I_s^d}$). (B) FOOPSI deconvolved traces using the SMGM bits per AP metric ($\widehat{I_{AP}^d}$). (C) The KSG measure applied to GCaMP6f traces. (D) The Binned Estimator applied to GCaMP6f traces using uniform bins. (E) The Binned Estimator applied to GCaMP6f traces using equal occupancy bins. (F) Table of summary

Guidelines for application of information metrics to functional fluorescence imaging data
Taken together, the above results suggest that across the information metrics applied directly to functional
fluorescence traces, the SMGM metrics provide the most reliable and interpretable information
measurements. We thus suggest the following guidelines for use and interpretation of the SMGM metrics
as applied to fluorescence mutual information metrics ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$) defined in Equations 7 and 8.

The SMGM bits per second metric ($\widehat{I_S^F}$) is likely attractive to imaging researchers because the units suggest
that precise knowledge of action potential numbers and times are not required for its use. However, there
are several challenges when applying the SMGM bits per second metric to functional fluorescence
imaging data. First, the substitution of the change in fluorescence map (f) for the action potential firing
rate map (λ) introduces a change in units, from bits per second to $\frac{\Delta F}{F} \frac{bits}{AP}$, which is difficult to interpret and
relate back to bits per second. Second, the transformation of action potential firing rate to change in
fluorescence can be approximated by a c scaling factor ($f = c\lambda$), which is measured in $\frac{\Delta F/F}{Hz}$, a quantity
that is unknown *a priori*. If c is not consistent between the neurons of a population of interest, then the
information values will be scaled differently and cannot be directly compared (Figure 2). Since c is
dependent on the width and height of the indicator response to a single action potential (the kernel), it can
vary from neuron to neuron based on difference in indicator expression level, intracellular calcium
buffering, and many other factors (Aponte et al., 2008; Greenberg et al., 2018; Helmchen and Tank, 2015;
Park and Dunlap, 1998). More research will be needed to measure these parameters (e.g. Chen et al., 2012,
2013a), and thus c, across neurons. Some results suggest that there may be non-trivial amounts of
variability within a populations of neurons (Éltes et al., 2019; Greenberg et al., 2018). With the impact of
c on the SMGM bits per second metric, and the possible variability of c across a population of neurons,
how can researchers properly extract useful measurements of information using $\widehat{I_S^F}$? First, we note that *if*
experimental measurements reveal small and acceptable variations in c across the neurons of interest, then
the information values derived from $\widehat{I_S^F}$ can be normalized by this factor to recover information values in
units of SMGM bits per second (independent of $\Delta F/F$) that can be compared across neurons. Under the
assumption of a consistent kernel, approximations for c for common indicators can be found in
Supplemental Figure 2. Further, given small variations in c across the neurons of interest, the ratio of $\widehat{I_S^F}$
between neurons in the population provides a meaningful metric for comparisons. For example, such ratios
could be used to divide a population of neurons accurately into groups based on their information values
(e.g. three quantiles of information) or compare the information values between different functional
subtypes of neurons (e.g. between place and non-place cells). However, the metric can still be useful even
if experimental measurements reveal large and unacceptable variations in c across the neurons of interest,
or if experimental measurements of c do not exist. In such cases, since it is reasonable to assume that c is
consistent in the same neuron over time, comparisons across the same neuron can provide meaningful
insights by using a ratio of $\widehat{I_S^F}$ measured (from the same neuron) across different conditions. For example,
quantifying the neuron by neuron ratio of $\widehat{I_S^F}$ between different behavioral states or conditions of an animal
or task, such as between goal-directed vs non-goal-directed running down a linear track, would allow
researchers to make conclusions such the following: "The population of neurons in region Z carries X+/-

Y times more information during goal-directed than non-goal-directed running." Therefore, we conclude
that with careful consideration of the (known or unknown) variability of the fluorescence response kernel
(c), $\widehat{I_S^F}$ can be used to extract useful measurements of information, either direct measurements of
information across a population of neurons (with known and similar c), ratios of information between
different neurons of a population (with known and similar c) or differences across different conditions
within the same neuron (with c unknown or different across neurons).

The SMGM fluorescence bits per action potential metric ($\widehat{I_{AP}^F}$) results in the same units as the action
potential based metric $\widehat{I_{AP}^E}$, and therefore may provide imaging researchers with information values that
are relatively easy to interpret. However, this similarity in units is somewhat misleading since the number
and timing of action potentials are not directly measured with functional fluorescence traces and the
asymmetric and relatively slow dynamics of fluorescence indicators leads to shifting and smoothing of
the action potential rate map (λ). This issue can have the effect of inducing a significant negative bias in
information measurements, especially at high information values and with functional indicators with wider
kernels (Figure 3). This is the most important factor to consider when determining how researchers can
properly extract useful measurements of information using $\widehat{I_{AP}^F}$. The shifting and smoothing of the action
potential rate map by fluorescence effectively leads to crosstalk between adjacent spatial bins. Therefore,
it is critical to consider the size of the spatial bins in relation to the spatial shift and smoothing induced by
the indicator (effectively the kernel plotted in space, rather than time, using the animal's average running
velocity to transform from time to space). It is reasonable then to counteract the spatial shift and smoothing
effect by using larger bin sizes, but this only works up to a point since larger bins limit the maximum
amount of information possible to measure and may negatively bias the information values near this upper
limit, even for action potential based recordings (Figure 3B-D). In practice, using gCaMP6f and the rodent
spatial behavior and spatial bin sizes (5cm) used here, our analysis suggests that $\widehat{I_{AP}^F}$ provides reasonable
measurements of information for neurons with values up to 3 bits per action potential (Figure 3E-G) since
this is the point where the absolute error exceeds 10% (comparable to the mean absolute error when
measuring information from action potential data (8.4%)). Equivalent thresholds for other common
indicators are shown in Supplemental Figure 3. The error is exacerbated by slower indicators and thus
more accurate measurements of information will result from using the fastest, narrowest kernel indicators
available, assuming signal to noise and detection efficiency are comparable across the different width
indicators. We conclude that with careful consideration of the size of the spatial bins in relation to the
spatial shift and smoothing induced by the indicator, $\widehat{I_{AP}^F}$ can be used to extract useful measurements of
information, most accurately for neurons with < 3 bits/action potential under recording conditions similar
to those considered here. Previous research quantifying information in bits per action potential using $\widehat{I_{AP}^E}$
have found that the majority of neurons carry information in this range (< 3 bits/action potential)
(Bourboulou et al., 2019; Knierim et al., 1995; Lee et al., 2006; Markus et al., 1995; Poucet and Sargolini,
2013), with a few exceptions (Ji and Wilson, 2007). Although these levels are dependent on the number
of bins and bin dwell time, $\widehat{I_{AP}^F}$ should be widely applicable to quantifying information throughout the
brain during behavior.

Figure 5. Application of SMGM information metrics to functional fluorescence imaging data from hippocampus during spatial behavior. A.) Example field of hippocampal pyramidal neurons expressing GCaMP6f and imaged during linear track navigation. Active cell ROIs shown in yellow; traces for green cells shown in B. B.) Fluorescence DF/F traces (green) from two neurons in the field shown in A and the track position during the recording (blue). C.). Distribution of information values using the fluorescence SMGM bits per second metric ($\widehat{I_S^F}$,(top) and the fluorescence SMGM bits per action potential metric ($\widehat{I_{AP}^F}$, bottom). The gray line indicates the recommended cutoff for reliability using GCaMP6f. D.) Plot of $\widehat{I_S^F}$ vs $\widehat{I_{AP}^F}$ for each neuron. Place cells indicated in red and nonplace cells in blue. E.) Example non-place cells spanning the information ranges shown in C. Spatial fluorescence map (f_i) shown on left, and average change in fluorescence per track traversal on right. F.) Same as E, but for place cells. G.) Bayesian decoding of mouse's track position using different subpopulations of neurons for one example session. From top to bottom: All active neurons, all nonplace cells, and place cells, the first through third quantiles of the SMGM bits per second formulation ($\widehat{I_S^F}$), and the first through third quantiles of the SMGM bits per action potential formation ($\widehat{I_{AP}^F}$). The white dashed line indicates the ground truth position of the animal, the color map indicates the decoded position probability (peak-normalized posterior distribution). H.) Decoding accuracy (absolute position decoding error in units of % of track) pooled over all sessions for

each neuron group indicated in G. Black bars indicate significant differences by Holm-Bonferroni
corrected rank sum tests (α=0.05).

Example: application of information metrics to functional fluorescence imaging data from hippocampus
during spatial behavior

In this section we demonstrate use of the above guidelines for proper application and interpretation of the
SMGM fluorescence mutual information metrics ($\widehat{I_s^F}$ and $\widehat{I_{AP}^F}$) defined in Equations 7 and 8. We applied
these metrics to functional fluorescence recordings (gCaMP6f*)* from pyramidal neurons in CA1 of the
hippocampus acquired during mouse spatial behavior.

CA1 neurons expressing gCaMP6f (viral transfection, CaMKII promoter) were imaged with two-photon
microscopy though a chronic imaging window during mouse navigation along a familiar 1D virtual linear
track, as described previously (Figure 5A-B, Dombeck et al., 2010; Radvansky and Dombeck, 2018;
Sheffield et al., 2017). 8 fields of view from 4 mice were recorded in 8 total sessions (recording duration
8.8+/-1.3 minutes, number of traversals/session: 29+/-2.5, 3.6+/-0.3 laps/min, 3 m long track). From these
8 sessions, 1,500 neurons were identified from our segmentation algorithm (see Methods), and analysis
was restricted to the 964 neurons that displayed at least one calcium transient on at least 1/3 of the
traversals during the session. Among these 964 neurons, 304 (31.5%) had significant place fields and were
thus identified as place cells (see Methods), while the remaining 660 (68.5%) did not pass a place field
test and were thus identified as non-place cells.

By applying Equation 7 (using 5 cm sized spatial bins), we found a continuum of spatial information
values measured by the fluorescence SMGM bits per second metric ($\widehat{I_s^F}$) across the 964 CA1 neurons,
with an average value of $\widehat{I_s^F}$=4.31$\pm$0.12 $\frac{\text{bits}\,^{\Delta F}/_F}{\text{Hz}}$ (Figure 5C). The units for $\widehat{I_s^F}$ make direct use and
interpretation of these values difficult, however, because these recordings were all from pyramidal neurons
in a single area, here for illustrative purposes, we will presume that variations in c (discussed above) across
the 964 neurons of interest are small and acceptable, with the absolute value of c unknown. This allows
for comparisons of information ratios across different subsets of the population. For example, place cells
had 2.8$\pm$0.20 times more information than non-place cells using the SMGM bits per second metric
($\widehat{I_s^F}$=5.3$\pm$0.14 for place vs. 1.9$\pm$0.087 $\frac{\text{bits}\,^{\Delta F}/_F}{\text{Hz}}$ for non-place, Rank sum p=1.7e-63, Figure 5D-F), although
there was substantial overlap in information between the populations (see distributions in Figure 5D and
individual examples in Figure 5E,F). This also allows for accurate division of the 964 neurons into 3
quantiles based on information values, which we use below for spatial location decoding.

By applying Equation 8 (using 5 cm sized spatial bins), we found a continuum of spatial information
values measured by the fluorescence SMGM bits per action potential metric ($\widehat{I_{AP}^F}$) across the 964 CA1
neurons, with an average value of $\widehat{I_{AP}^F}$=1.65$\pm$0.023 bits/action potential (Figure 5C). The units for $\widehat{I_{AP}^F}$
allow for direct use and interpretation of these values, and notably, because most (97%) of the neurons
had values less than 3 bits/action potential, a mean absolute error of <10% can be assumed across the
distribution of SMGM bits per action potential values. When applied to the place and non-place cell
populations, we found that place cells had higher information than nonplace cells using the fluorescence
SMGM bits per action potential metric ($\widehat{I_{AP}^F}$=1.8$\pm$0.026 and 1.35$\pm$0.042 bits/action potential, Rank sum
p=4.6e-21, Figure 5D-F).

As a demonstration of the usefulness of using information metrics to analyze large functional fluorescence
population recordings, we explored the accuracy of decoding the animal's track position using different
subsets of neurons. We divided the 964 neurons into 9 groups: All neurons, place cells, non-place cells,
three quantiles based on the fluorescence SMGM bits per second metric, and three quantiles based on the
fluorescence SMGM bits per action potential metric. We then used a Bayesian decoder of the animals'
position (see Methods) separately for each of the 9 neuron groups in each of the 8 sessions (Figure 5G,H).
An individual session decoding example can be seen in Figure 5G. We quantified decoding accuracy using
the absolute position decoding error (% of track), and pooled this measure across sessions for each neuron
group (Figure 5H). The means and standard errors for each group are: All neurons (7.33 +/-2.5 %), place
cells (6.97 +/-1.9 %), non-place cells (20.9 +/-1.8 %), SMGM bits per second Q1 (21.9 +/-1.5 %), SMGM
bits per second Q2 (13.2 +/-2.4 %), SMGM bits per second Q3 (8.97 +/-2.4 %), SMGM bits per action
potential Q1 (17.6 +/-2.7%), SMGM bits per action potential Q2 (17.8 +/-3.1 %), SMGM bits per action
potential Q3 (10.4 +/-3.0 %). Interestingly, even the lowest quantile information groups still could be used
to determine animal track location to within ~1/5 of the track. This supports the idea that the hippocampal
code for space is carried by a large population of active neurons (Meshulam et al., 2016), and not just by
a select subpopulation with the highest information or most well-defined tuning curves. As could be
expected, place cells encoded the position of the animal better than nonplace cells and better than the
lowest quantile information groups (Holm-Bonferroni corrected Rank sum, α=0.05) and neurons in the
higher quantiles provided more accurate decoding. Thus, the fluorescence information metrics provide a
means to compare the relative contribution of hippocampal neurons with different information values to
decoding animal position.

Discussion

Here, we performed an in-depth simulation study to examine the application of the SMGM bits per second
and SMGM bits per action potential metrics of mutual information to functional fluorescence recordings.
Since these metrics were designed for action potential recordings and since functional fluorescence
recordings violate some of the assumptions that these metrics are based on, it was unclear if and how the
metrics could be used for functional fluorescence recordings. We created a library of ten thousand mock
neurons whose action potential output carried ground-truth amounts of information about the animal's
spatial location, and by using real behavioral recording data from mice navigating in virtual linear tracks,
we simulated the spatial firing patterns of the mock neurons. We then simulated fluorescent calcium
responses for each neuron in each session by convolving the action potential trains with calcium kernels
for different indicators, primarily GCamp6f (though see Supplemental Figures 2 and 3 for results from
other indicators), and then added noise.

We then derived fluorescence versions of the SMGM bits per second ($\widehat{I_S^F}$) and SMGM bits per action
potential metrics ($\widehat{I_{AP}^F}$) (Equations 7 and 8) and applied them to the fluorescence traces in order to quantify
the performance of the metrics for estimating information. We found that ground truth information, as
measured by the fluorescence SMGM bits per second metric ($\widehat{I_S^F}$), was transformed into different units and
was linearly scaled by a factor (c) dependent on the height and width of the kernel, with c linearly
dependent on height and nonlinearly dependent on width. The error induced by these transformations
changed substantially over the range of kernel values of the different functional indicators widely used
today, and therefore are important factors to consider when designing and interpreting functional imaging
experiments. We then found that ground truth information, as measured by the fluorescence SMGM bits
per action potential metric ($\widehat{I_{AP}^F}$), retains the units and insensitivity to height scaling of the

electrophysiological metric ($\widehat{I_{AP}^E}$), but is nonlinearly biased by the smoothing of the fluorescence map
dictated by the width of the kernel. The estimation errors strongly depended on both the width of the
kernel and the information value being measured. Since these parameters change substantially over the
different functional indicators and different neuron types and behaviors that are commonly used today,
they are important factors to consider when designing and interpreting functional imaging experiments.

Using our mock fluorescence traces, we also asked if an AP estimation method could relieve the biases in
the SMGM metrics. Applying the SMGM bits per second metric ($\widehat{I_s^d}$) to AP estimation traces from a
deconvolution algorithm (FOOPSI) resulted in a very low c value for recovered versus ground truth
information. When the SMGM bits per AP measure was applied ($\widehat{I_{AP}^d}$), the resulting measurements of
information were still nonlinear (compared to $\widehat{I_s^F}$), with a positive bias at lower values of ground truth
information. Overall, applying FOOPSI to fluorescence traces led to a poorer recovery of ground truth
information using SMGM compared to direct application of SMGM to the florescence traces ($\widehat{I_{AP}^F}$). This
finding is consistent with recent research that has questioned the accuracy of deconvolution methods
(Evans et al., 2019). We also tested other metrics to measure mutual information directly from the
fluorescence time traces (KSG, Binned Estimator (uniform bins), and Binned Estimator (occupancy
binned)) and found these alternatives produced highly variable, saturating measurements of recovered
versus ground truth information. This was in contrast to the SMGM bits per second measure ($\widehat{I_s^F}$) which
produced a linearly scaled bias with lower error.

Materials and Methods

Construction of action potential trains with known ground truth information
To construct mock neurons with ground truth information, we adapted the differential form of the action
potential information, in bits per action potential (Equation 6). To create a rate map, we first selected an
average firing rate and target ground truth information. The mean rate ($\bar{\lambda}$) was always between 0.1 and 30
Hz, the information in bits per action potential (I_{AP}^E) between 0 and 6 bits per AP, and the information in
bits per second (I_s^E) between 0 and 24. To more evenly sample each of these, we first randomly selected
the bits per second (I_s^E) or bits per action potential (I_{AP}^E) to target. If the information target was in bits per
action potential, both the information (I_{AP}^E) and mean firing rate ($\bar{\lambda}$) were chosen uniformly. Because the
information in bits per second $I_s^E = \bar{\lambda} I_{AP}^E$, the bits per second information was not uniformly sampled in
this case. If the target was to be in bits per second, both the bits per action potential (I_{AP}^E) and SMGM bits
per second (I_s^E) measures were first chosen uniformly. Because the rate $\bar{\lambda} = I_s^E / I_{AP}^E$, this was not chosen
uniformly. This procedure was repeated to maintain the bounds on $\bar{\lambda}$, resulting in a non-uniform sampling
of information. The final distribution (Figure 1C) was spread acceptably for further analysis.

The rate maps were constructed by spline interpolating across 5 control points with two anchored at each
end of the track, and taking the exponential for each point, and then normalizing by the numerically
calculated integral (Figure 1A, D). To create a map matching the target information, we began with a
random spline. The 'y' (relative rate) initial position of each node was chosen from a standard normal
distribution and the initial 'x' (track position) of the 3 center nodes was chosen uniformly. The nodes were
then systematically moved using the MATLAB built in optimizer 'fmincon' with constraints preventing
the crossing of the center nodes and keeping them on the track, and the 'OptimalityTolerance' option set
to 0 (Figure 1A).

We then randomly selected behavioral traces (see Methods "Behavior" section) and concatenated sessions
until a total time randomly chosen between 3 and 60 minutes was reached (Figure 1E, F; Figure 2A, Figure
3A). The track positions were normalized and used to build a conditional intensity function (CIF) from
the rate function above. The CIF was normalized to match an expected mean rate over the entire session,
and the MATLAB built-in 'poissrnd' function was used to generate action potential times, sampled at 1
kHz. Finally, the action potential times were binned according to the counts within mock imaging frames
sampled at 30 Hz.

Simulated $\frac{\Delta F}{F}$ traces

To construct the $\frac{\Delta F}{F}$ traces (Figure1E,J,K; Figure2A; Figure3A), we first created a single action potential
response kernel from the peak-normalized sum of two exponentials:

$$g(t) = \frac{e^{-a\,t} - e^{-b\,t}}{\left(\frac{a}{b}\right)^{\frac{a}{b-a}} - \left(\frac{a}{b}\right)^{\frac{b}{b-a}}} \qquad\qquad 9$$

Where t is the time since the action potential and a and b are chosen to minimize $\left(1 - g(\tau_{rise})\right)^2 +$
$\left(0.5 - g(\tau_{rise} + \tau_{fall})\right)^2$ where τ_{rise} is the rise time in seconds and τ_{fall} is the half-fall time in seconds.
Deviations in τ_{rise} and τ_{fall} from baseline were also measured. The kernel g(t) was then multiplied by the
indicator height.

The GCaMP6f, GCaMP6s, and jRGECO1a heights, rise and fall times were measured as responses to
single action potentials *in vivo* (Chen et al., 2013; Dana et al., 2019)*:* other kernels (Figure 2H,
Supplemental Figures 2 and 3) were approximated from other experiments presented in the references
(seen in Table 1).

Table 1. Properties of indicator kernels used

Kernel	Height $\Delta F/F$	Rise (s)	Fall (s)	Source
gCaMP6f	0.190	0.042	0.142	(Chen et al., 2013)
jRGECO1a	0.164	0.041	0.207	(Kalko et al., 2011)
gCaMP7f	0.560	0.063	0.276	(Dana et al., 2019)
gCaMP6s	0.230	0.179	0.550	(Chen et al., 2013)
iGluSnfR-A184S	0.300	0.022	0.106	(Marvin et al., 2018)

To define the width of the kernel (Figure2L-N, Figure3K-M), we considered the kernel as a low pass
filtered version of the action potentials. If we normalize the filter to mean 1, it has the Fourier transform
$\left(\frac{1}{a+2\pi f \hat{\jmath}} - \frac{1}{b+2\pi f \hat{\jmath}}\right)\left(\frac{ab}{b-a}\right)$. The kernel width was defined as the -3 dB (50%) cutoff period of this filter:

$f^{-1} = \frac{\sqrt{-a^2-b^2+\sqrt{a^4+14a^2b^2+b^4}}}{2\pi\sqrt{2}}$. For the simulations with different width kernels, a kernel width was

chosen between 0.01 and 10 seconds, a rise time between 0.001 and 1 second, and a fall time between the

rise time and 2 seconds. Then, a and b were chosen to minimize the squared error between these three

targets using the built in MATLAB optimizer 'fminsearch'.

White noise with a standard deviation of 0.15 $\frac{\Delta F}{F}$ was then added to the mock fluorescence traces.

Deconvolution
Deconvolution was performed using the previously described FOOPSI algorithm (Vogelstein et al.,
2010). λ was set at 0.02154, which maximized the correlation between the deconvolved trace and the
true spike train in a random sample of 500 simulated traces: all other parameters were optimized for
each trace.

KSG estimator
The previously described second KSG estimator (Kraskov et al., 2004) was used using the 5[th] nearest
neighbor distance.

Binned estimators
The binned mutual information estimators were used (Timme and Lapish, 2018). The activity trace was
divided into 10 bins, either evenly across the span of the activity (uniform binned) or variably so the
bins contained the same number of samples (occupancy binned). Position was similarly divided into 60
bins.

Bayesian decoding
The Bayesian decoder used here (Figure 5 G,H) was adapted from a previously described method (Zhang
et al., 1998). Decoding was performed on the likelihood that a significant transient occurred in a time
frame, trained on the first 80% of the session and tested on the last 20%. The session was divided into
Δt=0.1 second bins. The conditional likelihood that an animal is in position x_i given the number of active
frames during a time window (n) is $p(x_i|n) = p_X(x_i)\left(\prod_{j=1}^{M} f_{i,j}^{n_j}\right)e^{-\Delta t \sum_{j=1}^{M} f_{i,j}}$, Where $p_X(x_i)$ is the
(marginal) probability that the animal is in the i^{th} spatial bin during a time sample, $f_{i,j}$ is the average rate
of significant frames by the j^{th} neuron in the i^{th} spatial bin, n_j is the number of significant frames
observed during the time window in neuron j, and M is the total number of neurons. The decoded position
was selected as the one with maximum conditional likelihood.

Animals
10 to 12 week old male C57BL/6 mice (20-30g) were individually housed under a reverse 12 hr. light /
dark cycle, all experiments were conducted during the dark phase. All experiments were approved by the
Northwestern University Animal Care and Use committee.

Behavior
We used a previously described virtual reality set-up and task (Heys et al., 2014; Sheffield et al., 2017;
Sheffield and Dombeck, 2015); some of the behavior sessions used here has previously appeared in these
studies. Briefly, water scheduled, head fixed mice were trained to run on a cylindrical treadmill down a
3m virtual track to receive a water (4 µl) reward at the end of the track, and were subsequently teleported
to the beginning of the track after a 1.5 s delay. Behavioral sessions were included if the animal ran at

least 20 laps containing a continuous 40 cm run for which the velocity was over 7 cm/sec during a 5-30
minute session.

Mouse surgery and virus injected
We performed population calcium imaging of CA1 neurons as described previously(Sheffield et al., 2017;
Sheffield and Dombeck, 2015). Briefly, 30 nL of AAV1-SynFCaMP6f (University of Pennsylvania
Vector Core, 1.5*10^13 GC/ml) was injected through a small craniotomy over the right hippocampus (1.8
mm lateral, 2.3 mm caudal of Bregma; 1.25 mm below the surface of the brain) under Isoflurane (1-2%)
anesthesia. 7 days later, a hippocampal window and head plate was implanted as described previously
(Dombeck et al., 2010).

Two-photon imaging
Imaging was performed as previously described(Sheffield et al., 2017; Sheffield and Dombeck, 2015).
Scanimage 4 was used for microscope control and acquisition (Pologruto et al., 2003). Time series movies
(1024 or 512x256 pixels) were acquired at 50 Hz. A Digidata1440A (Molecular Devices) with Clampex
10.3 synchronized position on the linear track, reward timing, and the timing of image frames.

Image processing, ROI selection and calcium transient analysis
Images were processed as previously described (Sheffield et al., 2017; Sheffield and Dombeck, 2015),
with minor modifications. Briefly, rigid motion correction was performed using cross correlation as in
(Dombeck et al., 2010; Miri et al., 2011; Sheffield and Dombeck, 2015), but here using a Fast-Fourier
transform approximation on the full video. ROIs were defined as previously described (Mukamel et al.,
2009) (mu=0.6, 150 principal/independent components, s.d. threshold = 2.5, s.d. smoothing width=1, area
limits = 100-1200 pixels). $\frac{\Delta F}{F}$ traces were generated by normalizing around the 8th percentile of a 3 second
sliding window. Significant transients from both experimental and mock fluorescence traces were selected
by comparing the ratio of amplitudes and durations of positive to negative going transients with a false
positive rate <0.01% (Dombeck et al., 2010). Mock traces used the histograms generated from the mock
gCaMP6f traces (Figure 2-3) or from the specific matching indicator traces (Supplemental Figures 2,3):
experimental data histograms were built separately. All subsequent analyses were run using these
significant transients.

Behavior analysis
The mean virtual track velocity was defined as the total virtual track distance covered during the session
divided by the total duration of the session; slow and stop periods were included in this metric. All other
analyses were restricted to long running periods, where the animal exceeded a virtual track velocity of 4
cm/s and ran continuously for at least 40 cm.

Defining place fields
Place fields were defined by first creating the spatial fluorescence intensity map (f_i) with the 300 cm track
divided into 60, 5 cm bins. This map was smoothed via a 3 bin boxcar. Transients identified during run
periods were shuffled in order and to random intervals to create 1000 bootstrapped intensity maps.
Candidate fields were defined as regions of the original fluorescence map with values greater than 99%
of the bootstrapped maps. Fields were then retained if they were between 20 and 120 cm wide: significant
place cells retained at least one field that satisfied these criteria.

Funding

We acknowledge funding from the McKnight Foundation (DAD), the Chicago Biomedical Consortium with support from the Searle Funds at The Chicago Community Trust (DAD), and NIH (T32 AG20506 to JRC and 2R01MH101297 to DAD).

Supplemental Information

Figure S 1. Effect of recording density on information metrics. A. The mean error (top) and absolute error (bottom) between the ground truth information (I_S^E) and measured information in SMGM bits per second from mock action potentials ($\widehat{I_S^E}$) as a function of the mean firing rate ($\bar{\lambda}$). B. As A, but as a function of number of laps. C-D. As A-B, but for measurements from mock fluorescence traces. E-F. As A-B, but for the SMGM bits per action potential metric. G-H.) As C-D, but for the SMGM bits per action potential metric.

Indicator	Error (AU)	Absolute Error (AU)	% Error	Absolute % Error	$c\,(\frac{\Delta F/_F}{Hz})$
jRGECO1a	4.58±4.4	4.8±4.2	40.7±43.9	42.7±42.0	1.37
gCaMP7f	57.6± 36.0	57.6±36.0	488±170	488±170	5.75
gCaMP6s	35.7±25.1	35.7±25.1	305.9±146	306.09±146	3.93
iGluSnfR-A184S	4.54± 3.6	4.63±3.5	39.1±25.8	39.82±24.6	1.37

Figure S 2. Effects of applying the SMGM bits per second metric to fluorescence traces from different
common functional indicators. Top: Information measured from mock traces using the SMGM bits per
second metric ($\widehat{I_S^F}$) vs ground truth information (I_S^E). Each dot is a single mock neuron, the gray dashed
line is the unity line (perfect measurement), the pink line is the line of best fit. Middle: Percentage error
density plots. The white line is the mean, the black are ± one standard deviation. Bottom: Summary
statistics and estimates for the scaling factor c.

Indicator	Error (bits/AP)	Absolute Error (bits/AP)	% Error	Absolute % Error	5% cutoff (bits/AP)	10% Cutoff (bits/AP)
jRGECO1a	-0.56±0.76	0.61±0.73	-10.4±25.5	18.5±20.4	1.68	2.40
gCaMP7f	-0.79±0.87	0.80±0.86	-20.4±47.8	23.6±46.2	0.48	1.44
gCaMP6s	-1.2±1.1	1.2±1.1	-25.3±194.9	43.7±191.6	0.48	0.72
iGluSnfR-A184S	-0.32±0.53	0.36±0.5	-5.7±17.9	11.2±15.2	2.6	3.4

Figure S 3. Effects of applying the SMGM bits per action potential metric to fluorescence traces from different common functional indicators. Top: Information measured from mock traces using the SMGM bits per action potential metric ($\widehat{I_{AP}^{F}}$) vs ground truth information (I_{AP}^{E}). Each dot is a single mock neuron, the gray dashed line is the unity line (perfect measurement), the pink line is the exponential fit. Middle: Percentage error density plots. The white line is the mean, the black are ± one standard deviation. Bottom: Summary statistics and percent error cutoffs.

References

Ahrens MB, Orger MB, Robson DN, Li JM, Keller PJ. 2013. Whole-brain functional imaging at cellular resolution using light-sheet microscopy. *Nat Methods* **10**:413–420. doi:10.1038/nmeth.2434

Allen TA, Salz DM, McKenzie S, Fortin NJ. 2016. Nonspatial Sequence Coding in CA1 Neurons. *J Neurosci* **36**:1547–1563. doi:10.1523/JNEUROSCI.2874-15.2016

Aponte Y, Bischofberger J, Jonas P. 2008. Efficient Ca2+ buffering in fast-spiking basket cells of rat hippocampus. *J Physiol* **586**:2061–2075. doi:10.1113/jphysiol.2007.147298

Aronov D, Tank DW. 2014. Engagement of Neural Circuits Underlying 2D Spatial Navigation in a Rodent Virtual Reality System. *Neuron* **84**:442–456. doi:10.1016/j.neuron.2014.08.042

Belghazi MI, Baratin A, Rajeswar S, Ozair S, Bengio Y, Courville A, Hjelm RD. 2018. MINE: Mutual Information Neural Estimation. *35th Int Conf Mach Learn ICML 2018* **2**:864–873.

Belitski A, Gretton A, Magri C, Murayama Y, Montemurro MA, Logothetis NK, Panzeri S. 2008. Low-
frequency local field potentials and spikes in primary visual cortex convey independent visual
information. *J Neurosci* **28**:5696–5709. doi:10.1523/JNEUROSCI.0009-08.2008

Bourboulou R, Marti G, Michon FX, El Feghaly E, Nouguier M, Robbe D, Koenig J, Epsztein J. 2019.
Dynamic control of hippocampal spatial coding resolution by local visual cues. *Elife* **8**:1–30.
doi:10.7554/eLife.44487

Brandon MP, Bogaard AR, Libby CP, Connerney MA, Gupta K, Hasselmo ME. 2011. Reduction of
Theta Rhythm Dissociates Grid Cell Spatial Periodicity from Directional Tuning. *Science (80-)*
332:595–599. doi:10.1126/science.1201652

Brown EN, Barbieri R, Eden UT, Frank L. 2003. Likelihood Methods for Neural Spike Train Data
Analysis In: Feng J, editor. Computational Neuroscience: A Comprehensive Approach. CRC Pres.
pp. 252–281.

Buzsáki G, Mizuseki K. 2014. The log-dynamic brain: how skewed distributions affect network
operations. *Nat Rev Neurosci* **15**:264–78. doi:10.1038/nrn3687

Calton JL, Stackman RW, Goodridge JP, Archey WB, Dudchenko PA, Taube JS. 2003. Hippocampal
place cell instability after lesions of the head direction cell network. *J Neurosci* **23**:9719–9731.
doi:23/30/9719 [pii]

Chen Q, Cichon J, Wang W, Qiu L, Lee SJR, Campbell NR, DeStefino N, Goard MJ, Fu Z, Yasuda R,
Looger LL, Arenkiel BR, Gan WB, Feng G. 2012. Imaging Neural Activity Using Thy1-GCaMP
Transgenic Mice. *Neuron* **76**:297–308. doi:10.1016/j.neuron.2012.07.011

Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA, Orger MB,
Jayaraman V, Looger LL, Svoboda K, Kim DS. 2013. Ultrasensitive fluorescent proteins for
imaging neuronal activity. *Nature* **499**:295–300. doi:10.1038/nature12354

Chen Y, Huang L, He J, Zhao K, Cai R, Hao Z. 2019. HASS: High Accuracy Spike Sorting with
Wavelet Package Decomposition and Mutual InformationProceedings - 2018 IEEE International
Conference on Bioinformatics and Biomedicine, BIBM 2018. Institute of Electrical and Electronics
Engineers Inc. pp. 831–838. doi:10.1109/BIBM.2018.8621401

Climer JR, DiTullio R, Newman EL, Hasselmo ME, Eden UT. 2015. Examination of rhythmicity of
extracellularly recorded neurons in the entorhinal cortex. *Hippocampus* **25**:460–473.
doi:10.1002/hipo.22383

Climer JR, Newman EL, Hasselmo ME. 2013. Phase coding by grid cells in unconstrained
environments: two-dimensional phase precession. *Eur J Neurosci* **38**:2526–41.
doi:10.1111/ejn.12256

Cohen MR, Kohn A. 2011. Measuring and interpreting neuronal correlations. *Nat Neurosci* **14**:811–819.
doi:10.1038/nn.2842

Dana H, Sun Y, Mohar B, Hulse B, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R, Macklin JJ,

Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS, Hod Dana, Yi
Sun, Boaz Mohar, Brad Hulse, Jeremy P Hasseman, Getahun Tsegaye, Arthur Tsang, Allan Wong,
Ronak Patel, John J Macklin, Yang Chen, Arthur Konnerth, Vivek Jayaraman, Loren L Looger,
Eric R Schreiter, Karel Svoboda DSK. 2018. High-performance GFP-based calcium indicators for
imaging activity in neuronal populations and microcompartments. *bioRxiv* **16**:434589.
doi:10.1101/434589

Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R,
Macklin JJ, Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS.
2019. High-performance calcium sensors for imaging activity in neuronal populations and
microcompartments. *Nat Methods* **16**:649–657. doi:10.1038/s41592-019-0435-6

Dauwels J, Vialatte F, Musha T, Cichocki A. 2010. A comparative study of synchrony measures for the
early diagnosis of Alzheimer's disease based on EEG. *Neuroimage* **49**:668–693.
doi:10.1016/j.neuroimage.2009.06.056

DeWeese MR, Zador AM, Hromádka T. 2008. Sparse Representation of Sounds in the Unanesthetized
Auditory Cortex. *PLoS Biol* **6**:14. doi:10.1371/journal.pbio.Citation

Dombeck DA, Harvey CD, Tian L, Looger LL, Tank DW. 2010. Functional imaging of hippocampal
place cells at cellular resolution during virtual navigation. *Nat Neurosci* **13**. doi:10.1038/nn.2648

Dombeck DA, Khabbaz AN, Collman F, Adelman TL, Tank DW. 2007. Imaging large-scale neural
activity with cellular resolution in awake, mobile mice. *Neuron* **56**:43–57.
doi:10.1016/j.neuron.2007.08.003

Éltes T, Szoboszlay M, Kerti-Szigeti K, Nusser Z. 2019. Improved spike inference accuracy by
estimating the peak amplitude of unitary [Ca 2+] transients in weakly GCaMP6f-expressing
hippocampal pyramidal cells. *J Physiol* **597**:2925–2947. doi:10.1113/JP277681

Evans MH, Petersen RS, Humphries MD. 2019. On the use of calcium deconvolution algorithms in
practical contexts. *bioRxiv* 871137. doi:10.1101/871137

Frank LM, Brown EN, Wilson MA. 2001. A comparison of the firing properties of putative excitatory
and inhibitory neurons from CA1 and the entorhinal cortex. *J Neurophysiol* **86**:2029–2040.

Fu H, Rodriguez GA, Herman M, Emrani S, Nahmani E, Barrett G, Figueroa HY, Goldberg E, Hussaini
SA, Duff KE. 2017. Tau Pathology Induces Excitatory Neuron Loss, Grid Cell Dysfunction, and
Spatial Memory Deficits Reminiscent of Early Alzheimer's Disease. *Neuron* **93**:533-541.e5.
doi:10.1016/j.neuron.2016.12.023

Fyhn M, Molden S, Hollup S, Moser M-B, Moser EI. 2002. Hippocampal Neurons Responding to First-
Time Dislocation of a Target Object. *Neuron* **35**:555–566. doi:10.1016/S0896-6273(02)00784-5

Gao W, Kannan S, Oh S, Viswanath P. 2017. Estimating mutual information for discrete-continuous
mixturesAdvances in Neural Information Processing Systems. pp. 5987–5998.

García-Rosales F, Beetz MJ, Cabral-Calderin Y, Kössl M, Hechavarria JC. 2018. Neuronal coding of

multiscale temporal features in communication sequences within the bat auditory cortex. *Commun*
Biol **1**:1–14. doi:10.1038/s42003-018-0205-5

Gerrard JL, Burke SN, McNaughton BL, Barnes CA. 2008. Sequence Reactivation in the Hippocampus
Is Impaired in Aged Rats. *J Neurosci* **28**:7883–7890. doi:10.1523/JNEUROSCI.1265-08.2008

Gómez-Verdejo V, Martínez-Ramón M, Florensa-Vila J, Oliviero A. 2012. Analysis of fMRI time series
with mutual information. *Med Image Anal* **16**:451–458. doi:10.1016/j.media.2011.11.002

Greenberg DS, Wallace DJ, Voit K-M, Wuertenberger S, Czubayko U, Monsees A, Handa T,
Vogelstein JT, Seifert R, Groemping Y, Kerr JN. 2018. Accurate action potential inference from a
calcium sensor protein through biophysical modeling. *bioRxiv* 479055. doi:10.1101/479055

Grubb MS, Thompson ID. 2006. Quantitative Characterization of Visual Response Properties in the
Mouse Dorsal Lateral Geniculate Nucleus. *J Neurophysiol* **90**:3594–3607.
doi:10.1152/jn.00699.2003

Hainmueller T, Bartos M. 2018. Parallel emergence of stable and dynamic memory engrams in the
hippocampus. *Nature* **558**:292–296. doi:10.1038/s41586-018-0191-2

Harvey CD, Collman F, Dombeck DA, Tank DW. 2009. Intracellular dynamics of hippocampal place
cells during virtual navigation. *Nature* **461**:941–946. doi:nature08499 [pii]10.1038/nature08499

Hazama Y, Tamura R. 2019. Effects of self-locomotion on the activity of place cells in the hippocampus
of a freely behaving monkey. *Neurosci Lett* **701**:32–37. doi:10.1016/j.neulet.2019.02.009

Helmchen F, Tank DW. 2015. A single-compartment model of calcium dynamics in nerve terminals and
dendrites. *Cold Spring Harb Protoc* **2015**:155–167. doi:10.1101/pdb.top085910

Heys JG, Dombeck DA. 2018. Evidence for a subcircuit in medial entorhinal cortex representing
elapsed time during immobility. *Nat Neurosci* **21**:1574–1582. doi:10.1038/s41593-018-0252-8

Heys JG, Rangarajan K V., Dombeck DA. 2014. The functional micro-organization of grid cells
revealed by cellular-resolution imaging. *Neuron* **84**:1079–1090. doi:10.1016/j.neuron.2014.10.048

Hinman JR, Brandon MP, Climer JR, Chapman GW, Hasselmo ME. 2016. Multiple Running Speed
Signals in Medial Entorhinal Cortex. *Neuron* **91**:666–679. doi:10.1016/j.neuron.2016.06.027

Hok V, Chah E, Save E, Poucet B. 2013. Prefrontal Cortex Focally Modulates Hippocampal Place Cell
Firing Patterns. *J Neurosci* **33**:3443–3451. doi:10.1523/JNEUROSCI.3427-12.2013

Huang Y, Brandon MP, Griffin AL, Hasselmo ME, Eden UT. 2009. Decoding movement trajectories
through a T-maze using point process filters applied to place field data from rat hippocampal region
CA1. *Neural Comput* **21**:3305–3334. doi:10.1162/neco.2009.10-08-893

Hubel DH, Wiesel TN. 2009. Republication of The Journal of Physiology (1959) 148, 574-591:
Receptive fields of single neurones in the cat's striate cortex. 1959. *J Physiol* **587**:2721–32.
doi:10.1113/jphysiol.2009.174151

Jeong J, Gore JC, Peterson BS. 2001. Mutual information analysis of the EEG in patients with
Alzheimer's disease. *Clin Neurophysiol* **112**:827–835. doi:10.1016/S1388-2457(01)00513-2

Ji D, Wilson MA. 2007. Coordinated memory replay in the visual cortex and hippocampus during sleep.
Nat Neurosci **10**:100–107. doi:10.1038/nn1825

Jiaying Tang SS. 2015. Visual receptive field properties of neurons in the mouse LGN. *PLoS Comput*
Biol **18**:449–452. doi:10.5061/dryad.b2t22

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018a. Dendritic Inhibition in the Hippocampus Supports Fear Learning. *Nat Biotechnol*
36:726–737. doi:10.1126/science.1247485

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018b. State-Dependent Subnetworks of Parvalbumin-Expressing Interneurons in Neocortex.
Nat Biotechnol **36**:726–737. doi:10.1016/j.celrep.2019.02.005

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018c. Calcium Dynamics in Dendrites of Hippocampal CA1 Interneurons in Awake Mice.
Nat Biotechnol **36**:726–737. doi:10.3389/fncel.2019.00098

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018d. A genetically encoded fluorescent acetylcholine indicator for in vitro and in vivo
studies. *Nat Biotechnol* **36**:726–737. doi:10.1038/nbt.4184

Kalko EK V, Dukas R, Ratcliffe JM, Teeling EC, Haven N, Fattu JM, Bates ME, Simmons J a,
Riquimaroux H, Surlykke A, Bouffard FH, Lee DN, Dear SP, Horiuchi TK, Krishnaprasad PS,
Moss CF, Schuller G, Brudzynski SM, Syme D a, Hollingworth S, Lindstedt SL, Baylor SM, Mead
a F, Rome LC, Goller F, Spiers ILY, Leeuwen JL Van, Schachat F, Rossmanith GH, Hoh JFY,
Kelley DB, Vater M a, Zhao Y, Araki S, Wu J, Teramoto T, Chang Y-F, Nakano M, Abdelfattah
AS, Fujiwara M, Ishihara T, Nagai T, Campbell RE. 2011. An Expanded Palette of Genetically
Encoded Ca2+ Indicators. *Science (80-)* **333**:1888–1891. doi:10.1126/science.1208592

Keller PJ, Ahrens MB. 2015. Visualizing whole-brain activity and development at the single-cell level
using light-sheet microscopy. *Neuron* **85**:462–483. doi:10.1016/j.neuron.2014.12.039

Khoshkhoo S, Vogt D, Sohal VS. 2017. Dynamic, Cell-Type-Specific Roles for GABAergic
Interneurons in a Mouse Model of Optogenetically Inducible Seizures. *Neuron.*
doi:10.1016/j.neuron.2016.11.043

Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB. 2018. Hippocampal Place Fields
Maintain a Coherent and Flexible Map across Long Timescales. *Curr Biol* **28**:3578-3588.e6.
doi:10.1016/j.cub.2018.09.037

Knierim JJ, Kudrimoti HS, McNaughton BL. 1995. Place cells, head direction cells, and the learning of landmark stability. *J Neurosci* **15**:1648–1659.

Koenig J, Linder AN, Leutgeb JK, Leutgeb S. 2011. The Spatial Periodicity of Grid Cells Is Not Sustained During Reduced Theta Oscillations. *Science (80-)* **332**:592–595. doi:10.1126/science.1201685

Kraskov A, Stögbauer H, Grassberger P. 2004. Estimating mutual information. *Phys Rev E - Stat Physics, Plasmas, Fluids, Relat Interdiscip Top* **69**:16. doi:10.1103/PhysRevE.69.066138

Kraus BJ, Brandon MP, Robinson RJ, Connerney MA, Hasselmo ME, Eichenbaum HB. 2015. During Running in Place, Grid Cells Integrate Elapsed Time and Distance Run. *Neuron* **88**:578–589. doi:10.1016/j.neuron.2015.09.031

Kropff E, Carmichael JE, Moser M-B, Moser EI. 2015. Speed cells in the medial entorhinal cortex. *Nature* **523**:419–424. doi:10.1038/nature14622

Lee I, Griffin AL, Zilli EA, Eichenbaum HB, Hasselmo ME. 2006. Gradual translocation of spatial correlates of neuronal firing in the hippocampus toward prospective reward locations. *Neuron* **51**:639–650.

Leveteau J, MacLeod P. 1966. Olfactory discrimination in the rabbit olfactory glomerulus. *Science* **153**:175–6. doi:10.1126/science.153.3732.175

Linsenbardt DN, Timme NM, Lapish CC. 2019. Encoding of the Intent to Drink Alcohol by the Prefrontal Cortex Is Blunted in Rats with a Family History of Excessive Drinking. *eneuro* **6**:ENEURO.0489-18.2019. doi:10.1523/ENEURO.0489-18.2019

Liu P, Jerrard LE, Bilkey DK, Jarrard LE, Bilkey DK. 2004. Excitotoxic lesions of the pre- and parasubiculum disrupt the place fields of hippocampal pyramidal cells. *Hippocampus* **14**:107–116. doi:10.1002/hipo.10161

MacDonald CJ, Carrow S, Place R, Eichenbaum HB. 2013. Distinct hippocampal time cell sequences represent odor memories in immobilized rats. *J Neurosci* **33**:14607–16. doi:10.1523/JNEUROSCI.1537-13.2013

Mankin EA, Thurley K, Chenani A, Haas O V, Debs L, Henke J, Galinato M, Leutgeb JK, Leutgeb S, Leibold C. 2019. The hippocampal code for space in Mongolian gerbils. *Hippocampus* 1–15. doi:10.1002/hipo.23075

Mann K, Gallen CL, Clandinin TR. 2017. Whole-Brain Calcium Imaging Reveals an Intrinsic Functional Network in Drosophila. *Curr Biol* **27**:2389-2396.e4. doi:10.1016/j.cub.2017.06.076

Manyakov N V., Van Hulle MM. 2008. Synchronization in monkey visual cortex analyzed with an information-theoretic measure. *Chaos* **18**:037130. doi:10.1063/1.2949928

Markus EJ, Qin YL, Leonard B, Skaggs WEW, McNaughton BBL, Barnes CA. 1995. Interactions between location and task affect the spatial and directional firing of hippocampal neurons. *J*

Neurosci **15**:7079–7094. doi:20026608

Marvin JS, Scholl B, Wilson DE, Podgorski K, Kazemipour A, Müller JA, Schoch S, Quiroz FJU, Rebola N, Bao H, Little JP, Tkachuk AN, Cai E, Hantman AW, Wang SSH, DePiero VJ, Borghuis BG, Chapman ER, Dietrich D, DiGregorio DA, Fitzpatrick D, Looger LL. 2018. Stability, affinity, and chromatic variants of the glutamate sensor iGluSnFR. *Nat Methods* **15**:936–939. doi:10.1038/s41592-018-0171-3

Marvin JS, Shimoda Y, Magloire V, Leite M, Kawashima T, Jensen TP, Kolb I, Knott EL, Novak O, Podgorski K, Leidenheimer NJ, Rusakov DA, Ahrens MB, Kullmann DM, Looger LL, Malgoire V, Leite M, Kawashima T, Jensen TP, Knott EL, Novak O, Podgorski K, Leidenheimer NJ, Rusakov DA, Ahrens MB, Kullmann DM, Looger LL. 2019. A genetically encoded fluorescent sensor for in vivo imaging of GABA. *Nat Methods* **16**:763–770. doi:10.1038/s41592-019-0471-2

Mau W, Sullivan DW, Kinsky NR, Hasselmo ME, Howard MW, Eichenbaum H. 2018. The Same Hippocampal CA1 Population Simultaneously Codes Temporal Information over Multiple Timescales. *Curr Biol* **28**:1499-1508.e4. doi:10.1016/j.cub.2018.03.051

Meshulam L, Gauthier JL, Brody CD, Tank DW, Bialek W. 2016. Collective behavior of place and non-place neurons in the hippocampal network 1–17.

Min B-C. 2003. Analysis of Mutual Information Content for EEG Responses to Odor Stimulation for Subjects Classified by Occupation. *Chem Senses* **28**:741–749. doi:10.1093/chemse/bjg066

Miri A, Daie K, Burdine RD, Aksay E, Tank DW. 2011. Regression-Based Identification of Behavior-Encoding Neurons During Large-Scale Optical Imaging of Neural Activity at Cellular Resolution. *J Neurophysiol* **105**:964–980. doi:10.1152/jn.00702.2010

Morris TP, White IR, Crowther MJ. 2019. Using simulation studies to evaluate statistical methods. *Stat Med* **38**:2074–2102. doi:10.1002/sim.8086

Mukamel EA, Nimmerjahn A, Schnitzer MJ. 2009. Automated Analysis of Cellular Signals from Large-Scale Calcium Imaging Data. *Neuron* **63**:747–760. doi:10.1016/j.neuron.2009.08.009

Newman EL, Climer JR, Hasselmo ME. 2014. Grid cell spatial tuning reduced following systemic muscarinic receptor blockade. *Hippocampus* **24**:643–655. doi:10.1002/hipo.22253

Nguyen JP, Shipley FB, Linder AN, Plummer GS, Liu M, Setru SU, Shaevitz JW, Leifer AM. 2016. Whole-brain calcium imaging with cellular resolution in freely behaving Caenorhabditis elegans . *Proc Natl Acad Sci* **113**:E1074–E1081. doi:10.1073/pnas.1507110112

Nguyen MN, Hori E, Matsumoto J, Tran AH, Ono T, Nishijo H. 2013. Neuronal responses to face-like stimuli in the monkey pulvinar. *Eur J Neurosci* **37**:35–51. doi:10.1111/ejn.12020

Nguyen MN, Matsumoto J, Hori E, Maior RS, Tomaz C, Tran AH, Ono T, Nishijo H. 2014. Neuronal responses to face-like and facial stimuli in the monkey superior colliculus. *Front Behav Neurosci* **8**:1–18. doi:10.3389/fnbeh.2014.00085

Niell CM, Stryker MP. 2008. Highly Selective Receptive Fields in Mouse Visual Cortex. *J Neurosci* **28**:7520–7536. doi:10.1523/jneurosci.0623-08.2008

O'Connor DH, Peron SP, Huber D, Svoboda K. 2010. Neural activity in barrel cortex underlying vibrissa-based object localization in mice. *Neuron* **67**:1048–1061. doi:10.1016/j.neuron.2010.08.026

O'Keefe J. 1976. Place units in the hippocampus of the freely moving rat. *Exp Neurol* **51**:78–109.

Østergaard J, Kramer MA, Eden UT. 2018. Capturing Spike Variability in Noisy Izhikevich Neurons Using Point Process Generalized Linear Models. *Neural Comput* **30**:125–148. doi:10.1162/neco_a_01030

Pachitariu M, Stringer C, Harris KD. 2018. Robustness of Spike Deconvolution for Neuronal Calcium Imaging. *J Neurosci* **38**:7976–7985. doi:10.1523/JNEUROSCI.3339-17.2018

Paninski L. 2004. Maximum likelihood estimation of cascade point-process neural encoding models. *Netw Comput Neural Syst* **15**:243–262. doi:10.1088/0954-898X_15_4_002

Park D, Dunlap K. 1998. Dynamic regulation of calcium influx by G-proteins, action potential waveform, and neuronal firing frequency. *J Neurosci* **18**:6757–66.

Park E, Dvorak D, Fenton AA. 2011. Ensemble Place Codes in Hippocampus: CA1, CA3, and Dentate Gyrus Place Cells Have Multiple Place Fields in Large Environments. *PLoS One* **6**:e22349. doi:10.1371/journal.pone.0022349

Park IJ, Bobkov Y V., Ache BW, Principe JC. 2013. Quantifying bursting neuron activity from calcium signals using blind deconvolution. *J Neurosci Methods* **218**:196–205. doi:10.1016/j.jneumeth.2013.05.007

Pastalkova E, Itskov V, Amarasingham A, Buzsáki G. 2008. Internally generated cell assembly sequences in the rat hippocampus. *Science (80-)* **321**:1322–7. doi:10.1126/science.1159775

Pologruto TA, Sabatini BL, Svoboda K. 2003. ScanImage: Flexible software for operating laser scanning microscopes. *Biomed Eng Online* **2**:13. doi:10.1186/1475-925X-2-13

Poucet B, Sargolini F. 2013. A trace of your place. *Science (80-)* **340**:35–36. doi:10.1126/science.1237567

Radvansky BA, Dombeck DA. 2018. An olfactory virtual reality system for mice. *Nat Commun* **9**:839. doi:10.1038/s41467-018-03262-4

Rashid SK, Pedrosa V, Dufour MA, Moore JJ, Chavlis S, Delatorre RG, Poirazi P, Clopath C, Basu J. 2020. The dendritic spatial code: branch-specific place tuning and its experience-dependent decoupling. *bioRxiv* **10**:2020.01.24.916643. doi:10.1101/2020.01.24.916643

Robbe D, Buzsáki G. 2009. Alteration of theta timescale dynamics of hippocampal place cells by a cannabinoid is associated with memory impairment. *J Neurosci* **29**:12597–605.

doi:10.1523/JNEUROSCI.2407-09.2009

Robinson SE, Mandell AJ. 2015. Mutual information in a MEG complexity measure suggests regional
hyper-connectivity in schizophrenic probands. *Neuropsychopharmacology*.
doi:10.1038/npp.2014.217

Roxin A, Brunel N, Hansel D, Mongillo G, van Vreeswijk C. 2011. On the Distribution of Firing Rates
in Networks of Cortical Neurons. *J Neurosci* **31**:16217–16226. doi:10.1523/jneurosci.1677-
11.2011

Scholl B, Wilson DE, Fitzpatrick D. 2017. Local Order within Global Disorder: Synaptic Architecture of
Visual Space. *Neuron* **96**:1127-1138.e4. doi:10.1016/j.neuron.2017.10.017

Shafi M, Zhou Y, Quintana J, Chow C, Fuster J, Bodner M. 2007. Variability in neuronal activity in
primate cortex during working memory tasks. *Neuroscience* **146**:1082–1108.
doi:10.1016/j.neuroscience.2006.12.072

Sheffield MEJ, Adoff MD, Dombeck DA. 2017. Increased Prevalence of Calcium Transients across the
Dendritic Arbor during Place Field Formation. *Neuron* **96**:490-504.e5.
doi:10.1016/j.neuron.2017.09.029

Sheffield MEJ, Dombeck DA. 2015. Calcium transient prevalence across the dendritic arbour predicts
place field properties. *Nature* **517**:200–204. doi:10.1038/nature13871

Simonnet J, Brecht M. 2019. Burst Firing and Spatial Coding in Subicular Principal Cells. *J Neurosci*
39:3651–3662. doi:10.1523/jneurosci.1656-18.2019

Skaggs WE, McNaughton BL, Gothard KM. 1993. An Information-Theoretic Approach to Deciphering
the Hippocampal Code In: Hanson SJ, Cowan JD, Giles CL, editors. Advances in Neural
Information Processing Systems 5. Morgan-Kaufmann. pp. 1030–1037.

Soo FS, Schwartz GW, Sadeghi K, Berry MJ. 2011. Fine Spatial Information Represented in a
Population of Retinal Ganglion Cells. *J Neurosci* **31**:2145–2155. doi:10.1523/jneurosci.5129-
10.2011

Stackman RW, Taube JS. 1998. Firing properties of rat lateral mammillary single units: head direction,
head pitch, and angular head velocity. *J Neurosci* **18**:9020–9037.

Stirman JN, Smith IT, Kudenov MW, Smith SL. 2016. Wide field-of-view, multi-region, two-photon
imaging of neuronal activity in the mammalian brain. *Nat Biotechnol* **34**:857–862.
doi:10.1038/nbt.3594

Stringer C, Pachitariu M, Steinmetz N, Carandini M, Harris KD. 2019. High-dimensional geometry of
population responses in visual cortex. *Nature* **571**:361–365. doi:10.1038/s41586-019-1346-5

Sun C, Yang W, Martin J, Tonegawa S. 2020. Hippocampal neurons represent events as transferable
units of experience. *Nat Neurosci* **23**. doi:10.1038/s41593-020-0614-x

Timme NM, Lapish C. 2018. A Tutorial for Information Theory in Neuroscience. *eneuro*
5:ENEURO.0052-18.2018. doi:10.1523/ENEURO.0052-18.2018

Treves A, Panzeri S. 1995. The Upward Bias in Measures of Information Derived from Limited Data
Samples. *Neural Comput* **7**:399–407. doi:10.1162/neco.1995.7.2.399

Vogelstein JT, Packer AM, Machado TA, Sippy T, Babadi B, Yuste R, Paninski L. 2010. Fast
Nonnegative Deconvolution for Spike Train Inference From Population Calcium Imaging. *J*
Neurophysiol **104**:3691–3704. doi:10.1152/jn.01073.2009

Wachowiak M, Shipley MT. 2006. Coding and synaptic processing of sensory information in the
glomerular layer of the olfactory bulb. *Semin Cell Dev Biol* **17**:411–423.
doi:10.1016/j.semcdb.2006.04.007

Wang S, Borst A, Zaslavsky N, Tishby N, Segev I. 2017. Efficient encoding of motion is mediated by
gap junctions in the fly visual system. *PLoS Comput Biol* **13**:e1005846.
doi:10.1371/journal.pcbi.1005846

Wilent WB, Nitz DA. 2007. Discrete place fields of hippocampal formation interneurons. *J*
Neurophysiol **97**:4152–4161. doi:10.1152/jn.01200.2006

Yaksi E, Friedrich RW. 2006. Reconstruction of firing rate changes across neuronal populations by
temporally deconvolved Ca2+ imaging. *Nat Methods* **3**:377–383. doi:10.1038/nmeth874

Yartsev MM, Ulanovsky N. 2013. Representation of Three-Dimensional Space in the Hippocampus of
Flying Bats. *Science (80-)* **340**:367–372. doi:10.1126/science.1235338

Zhang K, Ginzburg I, McNaughton BL, Sejnowski TJ. 1998. Interpreting neuronal population activity
by reconstruction: unified framework with application to hippocampal place cells. *J Neurophysiol*
79:1017–1044.

Zhou J-L, Shatskikh TN, Liu X, Holmes GL. 2007. Impaired single cell firing and long-term
potentiation parallels memory impairment following recurrent seizures. *Eur J Neurosci* **25**:3667–
3677. doi:10.1111/j.1460-9568.2007.05598.x

Zinyuk L. 2000. Understanding hippocampal activity by using purposeful behavior: Place navigation
induces place cell discharge in both task-relevant and task-irrelevant spatial reference frames. *Proc*
Natl Acad Sci **97**:3771–3776. doi:10.1073/pnas.050576397

Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, Gamal A El, Schnitzer MJ. 2013. Long-
term dynamics of CA1 hippocampal place codes. *Nat Neurosci* **16**:264–266. doi:10.1038/nn.3329

1279