Information theoretic approaches to deciphering the neural code with functional fluorescence imaging

Jason R. Climer[1], PhD and Daniel A. Dombeck[1]*, Ph.D.

[1] Department of Neurobiology, Northwestern University, Evanston IL
*Lead Contact: d-dombeck@northwestern.edu

Abstract

Information theoretic metrics have proven useful in quantifying the relationship between behaviorally relevant parameters and neuronal activity with relatively few assumptions. However, these metrics are typically applied to action potential recordings and were not designed for the slow timescales and variable amplitudes typical of functional fluorescence recordings (e.g. calcium imaging). The lack of research guidelines on how to apply and interpret these metrics with fluorescence traces means the neuroscience community has yet to realize the power of information theoretic metrics. Here, we used computational methods to create mock action potential traces with known amounts of information. From these, we generated fluorescence traces and examined the ability of different information metrics to recover the known information values. We provide guidelines for how to use information metrics when applying them to functional fluorescence and demonstrate their appropriate application to GCaMP6f population recordings from mouse hippocampal neurons imaged during virtual navigation.

Introduction

Neurons encode parameters important for animal behavior, at least in part, through the rate of production of action potentials (APs). Evidence for this can be found from electrophysiological AP recordings of orientation tuning in the visual system (Hubel and Wiesel, 2009), chemical sensing in the olfactory system (Leveteau and MacLeod, 1966; Wachowiak and Shipley, 2006), and spatial encoding in the hippocampus (O'Keefe, 1976). Key to deciphering the neural code, therefore, is defining metric to quantify the relationship between behavioral parameter spaces and a neuron's spiking rate. There are many metrics used for quantification, and are often used to compare neural responses across conditions or in neurons with complex responses. The underlying assumptions of the different metrics then become important factors to consider when determining which one to use.

Information theory is growing in popularity in the neuroscience community, largely because it provides a means to quantify rate coding with relatively few assumptions. One useful information theoretic measure is mutual information (MI), which is typically measured in bits per unit time, and describes the increase in predictability of the neural response when behavioral parameters are known. Formally, mutual information is the information about one variable that can be extracted from another, such as the information about behavior that can be derived from observing neural activity. Mutual information can be applied to neurons with widely varying response properties because it:

1. Is a nonlinear metric, not requiring the linearity assumptions of correlation metrics (e.g. (Grubb and Thompson, 2006; Hinman et al., 2016; Jiaying Tang, 2015; Kropff et al., 2015),
2. Does not assume a response shape, as is typical with Gaussian field mapping metrics (e.g. Kraus et al., 2015; Soo et al., 2011) or metrics using exponential or polynomial curve fitting (Hinman et al., 2016; Jiaying Tang, 2015), and
3. Uses the full time trace or shape of the mean response profile, rather than defining receptive fields with thresholding (e.g. Harvey et al., 2009; Niell and Stryker, 2008; Pastalkova et al., 2008).

However, MI can be nontrivial to estimate from neural and behavioral recordings and its estimation is an ongoing area of research (Belghazi et al., 2018; Gao et al., 2017; Kraskov et al., 2004; Timme and Lapish, 2018).

Here we focus on the most widely used estimator of MI in neuroscience, the SMGM estimator developed by Skaggs, McNaughton, Gothard and Markus (Skaggs et al., 1993), though as a point of comparison, we also consider the Binned Estimator (Timme and Lapish, 2018) and a separate technique developed by Kraskov, Stogbauer and Grassberger (KSG, Kraskov et al., 2004). The Binned Estimator estimates the joint probability distribution using a 2D histogram of neural response vs. behavioral variable; this transforms continuous variables into discrete values (Timme and Lapish, 2018). KSG estimates mutual information by examining the distance between data-points in the neural activity-behavioral parameter space. The SMGM estimator, on the other hand, relies on the assumption that AP firing follows an inhomogeneous Poisson process. The SMGM estimator therefore requires binning of only the behavioral variable(s), in contrast to the Binned Estimator. The profile of firing rates vs. behavioral variable is then used to estimate the MI.

The relative simplicity of the SMGM estimator has added to its popularity and widespread use in neuroscience applications for estimating behavioral information contained in single unit AP recordings. This metric has proven useful in quantifying rate coding in place cells (Knierim et al., 1995; Lee et al., 2006; Markus et al., 1995; Poucet and Sargolini, 2013), complex spatial responses of hippocampal interneurons (Frank et al., 2001; Wilent and Nitz, 2007), odor sequence cells (Allen et al., 2016), time cells (MacDonald et al., 2013), head direction cells (Stackman and Taube, 1998), speed cells (Fyhn et al., 2002), and face differential neurons (Nguyen et al., 2014, 2013), and has been used across multiple different species (Hazama and Tamura, 2019; Mankin et al., 2019; Yartsev and Ulanovsky, 2013). Furthermore, as a single neuron metric it provides statistical power for comparisons. Thus, it has been used to quantify differences in rate coding across different brain regions (Simonnet and Brecht, 2019) and across experimental interventions such as lesions (Calton et al., 2003; Liu et al., 2004), inactivations (Brandon et al., 2011; Hok et al., 2013; Huang et al., 2009; Koenig et al., 2011), and applications of drugs (Newman et al., 2014; Robbe and Buzsáki, 2009). Further, it has been used to examine differences in encoding across different behaviors (Aronov and Tank, 2014; Park et al., 2011; Zinyuk, 2000), and disease states (Fu et al., 2017; Gerrard et al., 2008; Zhou et al., 2007). SMGM information is often normalized from measuring bits per unit time to instead measure bits per AP. This creates a measure sensitive only to the selectivity of a neuron, and not its average firing rate. Thus, SMGM is a powerful tool for measuring the neural code in electrophysiological recordings of APs.

The power of MI estimators has yet to be fully exploited by the neuroscience community. For example, the estimators have not yet been widely used to compare encoding properties of large numbers of genetically identified neurons, or to quantify information content of other discrete signaling events such as synaptic inputs; both of which are difficult to study using electrophysiological methods. *In vivo* imaging of functional indicators has emerged as an important tool, largely because it possesses these capabilities. For example, using fluorescent calcium indicators, the functional properties of large populations of neurons can be simultaneously recorded in rodents (Dombeck et al., 2007; Radvansky and Dombeck, 2018; Sheffield et al., 2017; Stirman et al., 2016; Stringer et al., 2019; Ziv et al., 2013) zebrafish (Ahrens et al., 2013), or invertebrates such as *C. elegans* (Nguyen et al., 2016) and *Drosophila* (Keller and Ahrens, 2015; Mann et al., 2017). Furthermore, *in vivo* imaging can assure the genetic identity of the recorded neurons (Jing et al., 2018a, 2018b, 2018c; Khoshkhoo et al., 2017; Sheffield et al., 2017) and can access subcellular structures, allowing for functional recordings from synapses and dendrites using different

functional fluorescent indicators (e.g. Jing et al., 2018d; Marvin et al., 2019, 2018; Scholl et al., 2017;
Sheffield et al., 2017; Sheffield and Dombeck, 2015).

However, these indicators generate signals that are different from the underlying quantal events. For
example, somatic calcium indicators reveal intensity variations that are correlated with somatic AP firing
rates but are a smoothed and varying amplitude version of the AP train. This transformation from AP train
to fluorescence trace is an active area of research (Dana et al., 2018; Éltes et al., 2019; Greenberg et al.,
2018), but it is often approximated by convolving the AP train with a kernel, which defines the indicator's
response to a single AP. The shape of the kernel is a function of the indicator expression level, intracellular
calcium buffering, amount of calcium influx, efflux rates, background fluorescence, resting calcium
concentration, and other factors. When measured in pyramidal neurons, average kernels typically take the
shape of a sharp increase in fluorescence followed by an exponential decay to baseline (Chen et al., 2013;
Dana et al., 2018; Pachitariu et al., 2018; Park et al., 2013; Yaksi and Friedrich, 2006). Therefore, while
functional fluorescence imaging and information theoretic quantification may prove to be a powerful new
combination of tools to study neural correlates of behavior, it is critical to remember that functional
fluorescence signals represent altered versions of the underlying physiological events.

Caution is then needed when applying information metrics to continuous functional fluorescence traces,
yet the imaging community is already beginning to use information metrics, particularly SMGM. This
metric has been applied to somatic calcium responses to compare the information content of the same
neurons across different behavioral epochs (Heys and Dombeck, 2018), across different populations of
neurons in different brain regions (Hainmueller and Bartos, 2018), across different genetically identified
neural populations (Khoshkhoo et al., 2017), or to examine encoding by subcellular structures (Rashid et
al., 2020), or to classify the significance of encoding particular parameters by individual neurons (Kinsky
et al., 2018; Mau et al., 2018; Rashid et al., 2020).

However, it is essential to recognize some of the assumptions underlying these information metrics are
violated by functional florescence recordings. All three metrics (SMGM, KSG and Binned Estimation)
assume stationarity in the neural response, which is violated by the elongated time responses and relatively
slow fluctuations of the fluorescence intensity of the reporters. When applied to spiking data, there is also
a change in units: rather than AP counts, functional fluorescence traces are typically plotted in units of
florescence change with respect to baseline (ΔF/F). One possible solution to these issues would be to
deconvolve calcium traces to recover APs; however, deconvolution is an active area of research, and the
accuracy of these methods has recently been questioned (Evans et al., 2019). Ideally, the calcium traces
could be used directly to measure spiking information, without the need for such an in between, potentially
error inducing, step.

Quantifying the effects of the above violations on measurements of information using functional
fluorescence recordings with an analytical solution is particularly challenging with behaviorally
modulated neural recording data. However, a more tractable means of quantifying the effects would be to
use a simulation study to measure the induced biases and changes in measurement quality (Morris et al.,
2019). This strategy makes use of pseudo-randomly generated AP traces and has the advantage that the
ground truth parameters of the simulations are known, while variability due to behavior and other features
can be incorporated (Climer et al., 2015, 2013; Cohen and Kohn, 2011; Østergaard et al., 2018).

To provide the field with guidelines for the use of information metrics applied to functional fluorescence
recording data, we used computational simulation methods to create a library of ten thousand mock

neurons whose spiking output carry an exact, known (ground-truth) amount of information about the
animal's spatial location in its environment. We used real behavioral data (available at
https://doi.org/10.7910/DVN/SCQYKR) of spatial position over time from mice navigating in virtual
linear tracks and then simulated the spatial firing patterns of the mock neurons using an inhomogeneous
Poisson process framework (Brown et al., 2003; Climer et al., 2013; Paninski, 2004). We then simulated
fluorescent calcium responses for each neuron in each session by convolving the AP trains with calcium
kernels for different indicators, primarily GCamp6f (Chen et al., 2013), and then we added noise. MI
metrics (between spatial location and the neural signals) were then applied to the spiking or fluorescence
traces to quantify the performance of the metrics for estimating information. We provide a user toolbox
(found at https://github.com/DombeckLab/infoTheory), which consists of Matlab functions to generate
libraries of model neurons with known amounts of information, to generate spiking or fluorescence time-
series from those model neurons, and to estimate neuron information from real or model spiking or
fluorescence time-series datasets using the three metrics considered here (SMGM, Binned Estimator,
KSG). We focused on testing the performance of the SMGM method, and then compared its performance
to the Binned Estimation and KSG methods, which do not have the underlying Poisson assumption
required for the SMGM approach. We also applied a deconvolution algorithm to test its performance. We
then applied this analysis to real datasets of hippocampal neuron populations from mice navigating in
virtual linear tracks. We quantified the spatial information content of the populations and then performed
Bayesian decoding of mouse position from different information containing subsets of this population.
Interestingly, we found that the population quantile with the lowest information values were still able to
decode mouse position to the closest quarter of the track. Thus, we provide new findings about the neural
code for space that were made possible by the information metrics and guidelines that we introduce here.

The SMGM method applied directly to the mean ΔF/F intensity map appeared to best recover the ground
truth information. We provide guidelines for the use of the SMGM metric when applied to functional
fluorescence recordings and demonstrate the appropriate application of these guidelines to GCaMP6f
population recordings from hippocampal neurons in mice navigating virtual linear tracks.

Results

The SMGM information metrics
Here we review the derivation of the SMGM information metrics and the underlying assumptions. For
illustrative purposes throughout this manuscript, we use the example of spatial encoding in which the
firing pattern of neurons carry information about the animal's location along a linear track; however, the
derivations, equations and conclusions generalize to encoded variables over other domains and
dimensionalities.

Consider a random variable X representing the positions an animal might take, with x being its value
measured at one time sample. The positions are subdivided into N spatial bins, such that x can take on the
values $\{1, 2, \dots, N\}$. For our analyses, $N=60$. Consider a random variable Y representing the number of APs
a neuron might fire, where y is the count measured within a time sample. y can take on the values of
$\{0, 1, \dots, +\infty\}$. X and Y are both discrete. If X and Y both obey the assumption that each time sample is
independent (i.e. they are stationary), then the mutual information (I, in bits per sample) between X and
Y is expressed as follows:

$$I(X;Y) = \sum_{i=1}^{N} \sum_{y=0}^{+\infty} p_{X,Y}(x_i, y) \log_2 \frac{p_{X,Y}(x_i, y)}{p_X(x_i)p_Y(y)} \qquad 1$$

169   Where $p_X(x_i)$ is the (marginal) probability that the animal is in the i^{th} spatial bin during a time sample,
170   $p_Y(y)$ is the probability that the neuron fires y APs in the time sample, and $p_{X,Y}(x_i,y)$ is the joint
171   probability that the neuron fires y APs and is in the i^{th} bin. Recall that $p_{X,Y}(x_i, y) = p_{Y|X}(y|x_i)p_X(x_i)$,
172   where $p_{Y|X}(y|x_i)$ is the conditional probability that the neuron fires y APs given that the animal is in the
173   i^{th} spatial bin. We can thus rewrite Equation 1 as follows:

$$I(X;Y) = \sum_{i=1}^{N} \sum_{y=0}^{+\infty} p_{Y|X}(y|x_i)p_X(x_i) \log_2 \frac{p_{Y|X}(y|x_i)}{p_Y(y)} \qquad 2$$

174   With the further assumption that the firing of the neuron follows Poisson statistics, we can then estimate
175   the mutual information as follows: let the AP rate (AP/s or Hz) in a single bin be λ_i, and the average across
176   the session be $\bar{\lambda}$. For an arbitrarily small time window Δt, the probability that an AP occurs in that window
177   is $\Pr(Y = 1|x = i) = \lambda_i \Delta t$, with the probability that an AP occurs regardless of position as $\Pr(Y = 1) =$
178   $\bar{\lambda}\Delta t$. We can thus rewrite Equation 2 as:

$$I(X;Y) = \sum_{i=1}^{N} \lambda_i \Delta t p_X(x_i) \log_2 \frac{\lambda_i}{\bar{\lambda}} \qquad 3$$

179   By integrating over one second ($\int_0^1 I(X;Y)d\Delta t$) we obtain the first key SMGM metric for spatial
180   information as measured by AP firing, which is in units of bits per second:

$$\widehat{I_s^E} = \sum_{i=1}^{N} \lambda_i p_X(x_i) \log_2 \frac{\lambda_i}{\bar{\lambda}} \qquad 4$$

For notation, we will use a carrot (^) to indicate an information value that is measured from experiment,
the superscript (E in this case) to show the source of the data, and a subscript to show the units/formula
used (bits per second in this case). Thus, $\widehat{I_s^E}$ is the information measured via electrophysiology in bits per second.
This metric is linearly dependent on the average firing rate of the neuron, and this dependence is often
removed through normalization by the average firing rate to obtain the second key metric of spatial
information as measured by AP firing, which is in units of bits/(second*Hz), or more commonly, bits per
AP:

$$\widehat{I_{AP}^E} = \frac{1}{\bar{\lambda}} \sum_{i=1}^{N} \lambda_i p_X(x_i) \log_2 \frac{\lambda_i}{\bar{\lambda}} \qquad 5$$

Therefore, these two key metrics of spatial information are defined completely by quantities that can be
experimentally measured: the mean firing rate ($\bar{\lambda}$) from the AP counts over the duration of the recording,

the AP firing rate in the i^{th} bin from the average rate map (λ_i), and the probability that the animal is in
the i^{th} spatial bin from the normalized occupancy map ($p_X(x_i)$). The quantity of and noise in these
measurements affects the quality of the metric: in particular, undersampling due to low firing rates or low
trial counts induces a substantial positive bias (Treves and Panzeri, 1995).

In the derivation of these metrics, there are two key assumptions that are violated by functional
fluorescence recordings. First, the recordings do not follow Poisson statistics: instead of discrete counts
of APs (y), the functional fluorescence traces consists of a continuous relative change in fluorescence
($\Delta F/F$), and instead of a firing rate map (λ_i) measured in Hz, average intensity maps in units of $\Delta F/F$ are
generated. The stationarity assumption is also violated: due to the slow decay, a time sample of the
fluorescence traces depend on the previous samples. The violation of these assumptions by functional
fluorescence recording will affect the precision and induce biases in the SMGM information metrics. Since
these effects have not previously been addressed or quantified, we measured these biases here using a
simulation study.

Building a ground truth library of 10,000 neurons with known values of information

To create a neuron with a known, ground truth information value, it was necessary to generate a continuous
(i.e. infinitesimally small bins) rate map ($\lambda(x)$) matching the desired information. To do this, we first
normalized the track length to 1 and assumed the animal's occupancy map to be spatially uniform
$\left(p_X(x_i) = \frac{1}{N}\right)$. We then created an exponentiated cubic spline with 5 randomly positioned nodes (Figure
1A) to build a starting continuous map of the normalized instantaneous firing rate, $\frac{\lambda}{\bar{\lambda}}(x)$, with the integral
normalized to 1. We calculated the ground truth amount of information in bits per AP as follows:

$$I_{AP}^E = \int_0^1 \frac{\lambda}{\bar{\lambda}}(x) \log_2 \left(\frac{\lambda}{\bar{\lambda}}(x)\right) dx \qquad 6$$

The locations of the 5 nodes were then systematically varied [see Methods] to minimize the squared error
between the value calculated in Equation 6 and a target amount of information (Figure 1A,B), in the end
resulting in a mean error of 5.1*10^-9 bits/AP and a mean absolute error of 1.5*10^-7 bits/AP. The rate
map at this convergence point was used for further analysis. This procedure was repeated to generate
10,000 mock neurons with a range of (known and ground truth) information values. Note that the value in
Equation 5 cannot be higher than when all the APs arrive in one spatial bin; the rate in that bin is $N\bar{\lambda}$. If
we assume uniform occupancy $\left(p_X(x_i) = \frac{1}{N}\right)$, then the maximum measureable information is $\log_2 N$, in
our case, 5.9 bits/AP with N=60 bins. Thus, the information values considered here range between 0 and
6 bits/AP (Figure 1C). We chose a mean firing rate ($\bar{\lambda}$) for the neurons between 0.1 and 30 Hz, a range
observed for a variety of different cortical and hippocampal neurons during behavior (Buzsáki and
Mizuseki, 2014; DeWeese et al., 2008; O'Connor et al., 2010; Roxin et al., 2011; Shafi et al., 2007). From
Equations 4-5, the ground truth information in bits per second is $I_S^E = \bar{\lambda} I_{AP}^E$. I_S^E for these choices resulted
in ground truth information values between I_S^E =0 and I_S^E =24 bits/sec (Figure 1C). Example low
(I_{AP}^E=0.04 bits/AP) and mid (I_{AP}^E=2 bits/AP) rate maps are shown in Figure 1D.

These rate maps provided a basis for generating mock AP firing data (and functional fluorescence data,
see below). Under real experimental conditions, recording duration and bin sizes are finite and animal

occupancy maps ($p_X(x_i)$) are not spatially uniform. These experimental limitations add error to the
estimate of a neuron's ground truth information value. Therefore, in order to accurately re-create these
limitations in our simulation study, we used real behavior datasets from head-restrained mice running
along a 3 m virtual linear track for water rewards (acquired as in Sheffield et al., 2017; Sheffield and
Dombeck, 2015). Unless otherwise indicated, all values reported will be the mean±standard deviation.
We selected at random from a library of 574 behavior sessions from mice navigating along familiar tracks
and concatenated and truncated these sessions to create behavior sessions uniformly sampled up to 60
minutes in duration (average 30.2±17.1 minutes), resulting in an average 132 ± 71.2 laps per session and
an average running speed of 19.3 + 3.87 cm/sec (Figure 1E.1). This behavior, the average firing rate ($\bar{\lambda}$),
and the normalized rate map ($\frac{\lambda}{\bar{\lambda}}(x)$) from the mock neurons were used to create an instantaneous firing
rate trace (Figure 1E.2), sampled at 1 kHz, from which AP times were generated assuming Poisson firing
statistics (Figure 1E.3). An example mock of spiking in response to behavior for low (0.04 bits/AP) and
mid (2 bits/AP) information neurons can be seen in Figure 1F-H. From these spiking responses, we then
generated mock fluorescence traces by convolving the raster with a double-exponential kernel matching
the rise and fall times for GCaMP6f (Chen et al., 2013, Figure 1E.4) and adding random Gaussian noise
to model shot noise. Mock fluorescence traces for the two example neurons in Figure 1F-H can be seen in
Figure 1I-J. The mock AP and fluorescence traces were used to create session mean spatial maps – of
binned firing rate (λ_i in Hz) and change in fluorescence (f_i in $\Delta F/F$), for information analyses (Figure
1K). By repeating this process, we built a large dataset of spiking and fluorescence traces, generated from
our library of mock neurons with known amounts of information and using real animal spatial behavior.
With tens of thousands of these mock neuron recordings, we could then assess the effects of many
simulation parameters on the information values determined from the metrics including firing rate, session
duration, fluorescence kernel shape, and ground truth information value.

Quantification of the accuracy and precision of the SMGM bits per second metric using functional
fluorescence recordings
We first applied the SMGM bits per second metric ($\widehat{I_s^E}$) to our mock AP recording traces to verify that
they can recover our ground-truth information values given finite recording durations and bin sizes, and
non-uniform animal occupancy maps ($p_X(x_i)$). Figure 2A shows three mock neurons with ground truth
information values of I_s^E =3, 15 and 23 bits/sec. When the SMGM bits per second metric ($\widehat{I_s^E}$) was applied
to the AP traces from these example neurons, the information was well recovered, with $\widehat{I_s^E}$=2.8, 15 and 24
bits/sec, respectively. The results from these examples also held across the full 10,000 mock neuron library
(Figures 2B-D), as a linear fit (y-intercept = 0.093±0.040, intercept p=4.6*10^{-6} bits per second and slope
= 0.97 ±0.0030, slope p<<0.01) explained nearly all the variance (R^2 = 0.97), the average error was
0.22±1.25 bits per second (1.0±0.69% error) and the absolute error was 0.64±1.05 bits per second
(8.4±0.69% error). There is a substantial positive bias for the lowest firing rates and smallest number of
trials (Figure 2-figure supplement 1A-B) which has been previously well characterized (Treves and
Panzeri, 1995), with average errors exceeding +10% for less than 6 minutes of recording, mean rate under
0.6 Hz, and under 11 trials. Thus, the SMGM bits-per-second metric ($\widehat{I_s^E}$) recovers the ground-truth
information well using AP recordings, with the only error coming from finite recording time and variable
animal behavior.

We next discuss the changes to the SMGM bits per second metric ($\widehat{I_s^E}$) commonly used for application to
functional fluorescence traces (Hainmueller and Bartos, 2018; Heys and Dombeck, 2018), and explore the
implications of these changes. Most simply, the mean firing rate ($\bar{\lambda}$) and the mean firing rate in a spatial

bin (λ_i) are replaced by the mean change in fluorescence ($\bar{f}$) and the mean change in fluorescence in a
bin (f_i). Making these substitutions in Equation 4 results in the information as measured by functional
fluorescence:

$$\widehat{I_s^F} = \sum_{i=1}^{N} f_i p_X(x_i) \log_2 \frac{f_i}{\bar{f}} \qquad\qquad 7$$

The fluorescence map f_i differs from the firing rate map λ_i in two ways. First, the fluorescence map is
approximated by the firing rate map scaled by a factor c, dependent on the height and width of the kernel
and measured in units of $\frac{\Delta F/F}{Hz}$, and second it is smoothed by the kernel (Figure 1E4). If we discount the
latter for a moment and focus on the scaling, $f \approx c\lambda$, we can see that substituting λ with $c\lambda$ in Equation 4
results in $\widehat{I_s^F} = c\widehat{I_s^E}$. The units for $\widehat{I_s^F}$ are no longer in bits per second, as it has previously been reported
(Hainmueller and Bartos, 2018), but are instead in units of $\frac{bits \, \Delta F/F}{sec \, Hz}$ or $\frac{bits \, \Delta F/F}{AP}$, which are difficult to
interpret (see *Guidelines for application of information theoretic metrics to functional fluorescence*
imaging data section for further discussion). The effect of smoothing is difficult to analytically quantify
since it both alters c by changing the average intensity and distorts the firing rate map. Therefore, to fully
quantify the impact of convolving an AP recording with a functional fluorescence kernel on recovering
ground truth information, we used our mock fluorescence traces.

We applied a GCaMP6f modeled kernel to the 10,000 mock AP traces to generate 10,000 mock
fluorescence calcium traces. Figure 2A shows the fluorescence traces generated from three mock neurons
with ground truth information values of $I_s^E = 3$, 15 and 23 bits/sec. The effects of the convolution can be
seen in the differences in scaling and shape between the fluorescence maps f_i and the firing rate maps λ_i.
When the fluorescence metric ($\widehat{I_s^F}$) was applied to the fluorescence traces from these example neurons, the
information recovered was $\widehat{I_s^F} = 0.13$, 0.47 and 1.1 $\frac{bits \, \Delta F/F}{AP}$, respectively, indicating deviation from the
ground truth information values assuming the units are comparable. The results from these examples also
held across the full 10,000 mock neuron library (Figures 2E-G), as there was a clear scaling of the ground
truth information and a consistent overestimation with a mean error of -11.1$\pm$6.7 AU (-96.0$\pm$1.3% error).
The best-fit line of the measured information ($\widehat{I_s^F}$) versus the ground truth information (I_s^E) had an
intercept near 0 (0.0029$\pm$0.0016 $\frac{bits \, \Delta F/F}{AP}$, p=0.07). The slope of this fit was 0.039$\pm$12e-4 $\frac{\Delta F/F}{Hz}$ (p<<0.01),
which provides a measure of the scaling factor (c). This error was not corrected for with denser sampling:
it remained consistent even at high firing rates and many trials (Figure 2-figure supplement 1C-D). In
addition to this scaling effect caused by c, smoothing of the rate map could induce nonlinearity in the
relationship between $\widehat{I_s^F}$ and I_s^E. To test for such an effect, we fit the measured information in Figure 2E
with a saturating exponential and compared the fits using a likelihood ratio test: the exponential did not
significantly improve the fit (χ_1^2=0.093, p=0.76), which indicates that smoothing by the kernel does not
induce significant nonlinearities. c is dependent on the height and width (the integral) of the kernel and
was measured here as 0.039$\pm$12e-4 $\frac{\Delta F/F}{Hz}$. The consistent, negative bias observed in estimating information
with $\widehat{I_s^F}$ (Figure 2E) would be easy to correct for assuming the c factor, and therefore the kernel, were
similar across all measured neurons. This point is considered further in the *Guidelines for application of*
information metrics to functional fluorescence imaging data section below. We conclude that ground truth

information, as measured by the fluorescence SMGM bits per second metric ($\widehat{I_s^F}$), is transformed into
different units and is linearly scaled by a factor c dependent on the height and width of the kernel.

The amplitude (height) of the change in fluorescence can vary across indicators and conditions. The height
of the kernel, given a constant kernel width, should linearly scale c and the error in estimating information
with $\widehat{I_s^F}$. To explicitly test this prediction, we simulated an additional 5,000 fluorescence traces with
kernels of varying height (0-3 $\Delta F/F$, Figure 2I-K), but that maintain the same shape and width (from the
GCaMP6f kernel), and then measured the percent error in estimating information with $\widehat{I_s^F}$. As observed
above for the GCaMP6f example (Figure 2 E-G), the percent error in estimating information with $\widehat{I_s^F}$ shows
little dependence on ground truth information (Figure 2I-K). However, as a function of the height of the
kernel, the percent error (averaged over all ground truth information values) in estimating information
with $\widehat{I_s^F}$ is fit well with an increasing linear function (intercept = -99.8±0.42%, intercept p<<0.01, slope
20.7 ±0.14%/$\frac{\Delta F}{F}$, slope p<<0.01, R^2= 0.80; Figure 2I-J). Over the wide array of available functional
fluorescent indicators in use today (Figure 2H), this leads to differences in error due to differences in
transient height of the indicator used alone. For the indicators shown in Figure 2H, there is an average
height of 0.603±0.10 SD $\Delta F/F$: the error spans from -95.8% for the kernel height reported for gCamp6f
(0.19 $\Delta F/F$) to -88.2% for gCamp7f (0.56 $\Delta F/F$). It should be noted that fluorescence ($\Delta F/F$) is always
reported here as a fractional change, not as a percentage (% $\Delta F/F$); if a kernel height of 19 % $\Delta F/F$ is
used, the units would again change. Thus, as expected, the percent error in estimating information with
the SMGM bits per second estimator ($\widehat{I_s^F}$) scales linearly with the height of the kernel.

The width of the kernel can vary widely across fluorescent indicators (Figure 2H), with "faster" indicators
boasting shorter rise and fall times. The combined effect of a longer rise and fall time is to smooth and
delay the AP train; in other words, it acts as a causal low-pass filter. The cutoff period of this low pass
filter provides a measurement of the effective width of the kernel [see Methods]. The effect of such
differences in kernel shape on the error in estimating information with $\widehat{I_s^F}$ is difficult to measure
analytically. We therefore simulated an additional 5,000 fluorescence traces with kernels of different
kernel widths (but constant height of the GCaMP6f kernel), resulting in a range of kernel durations (rise
times: 1 ms to 1 second, fall times longer than the rise time up to 2 seconds), and then we measured the
percent error in estimating information with $\widehat{I_s^F}$. Similarly, as observed above for the GCaMP6f and
varying kernel height examples (Figure 2 E-K), the percent error in estimating information with $\widehat{I_s^F}$ shows
little dependence on ground truth information (Figure 2N). Interestingly, the percent error (averaged over
all ground truth information values) in estimating information with $\widehat{I_s^F}$ shows a complex nonlinear response
as a function of the width of the kernel (Figure 2L-M). The error increases up to a kernel width of ~2.5
seconds, at which point it saturates at ~-85% error. This arises from an interaction between changing the
average value of the original AP trace and flattening the average fluorescence map (f_i). Over the wide
array of available functional fluorescent indicators in use today, this leads to differences in error due to
differences in width of the indicator used alone. For example, an average error of -97.1±0.63% were
observed for iGluSnfR, the shortest indicator considered here at 0.52 seconds. For gCamp6s, the slowest
indicator examined (2.54 seconds), the average was -89.6±4.6%. To estimate the percent errors for these
five indicators considering differences in *both* height and duration, we used these 5 kernels to generate
mock fluorescence traces from the 10,000 neurons in Figure 2B-G. The resulting distributions, estimated
c values, and mean and absolute errors can be seen in Figure 2-figure supplement 2. In summary, we
conclude that information, as measured by the fluorescence SMGM bits per second metric ($\widehat{I_s^F}$), is

transformed into different units and is linearly scaled by a factor (c) dependent on the height and width of
the kernel, with c linearly dependent on height and nonlinearly dependent on width. The error induced by
these transformations changes substantially over the range of kernel values of the different functional
indicators widely used today, and therefore these are important factors to consider when designing and
interpreting functional imaging experiments (see *Guidelines for application of information metrics to*
functional fluorescence imaging data for further discussion).

Quantification of the accuracy and precision of the SMGM bits per AP metric using functional
fluorescence recordings
The SMGM metric is commonly normalized by the mean rate to obtain a measurement in units of bits per
AP. We thus applied the SMGM bits per AP metric ($\widehat{I_{AP}^E}$) to our mock AP recording traces to verify that
they can recover our ground-truth information values. Figure 3A shows three mock neurons with ground
truth information values I_{AP}^E =0.05, 1.8 and 4.2 bits/AP. When the SMGM bits per AP metric ($\widehat{I_{AP}^E}$) was
applied to the AP traces from these example neurons, the information was well recovered, with $\widehat{I_{AP}^E}$=0.06,
1.8 and 4.2 bits/AP respectively. The results from these examples also held across the full 10,000 mock
neuron library (Figures 3B-D), as a linear fit (y-intercept = 0.087$\pm$0.029, intercept p=2.8e-184 bits per
second and slope = 0.93 $\pm$0.0010, slope p<<0.01) explained nearly all the variance ($R^{2=}$ 0.99), the average
error was -0.071$\pm$0.23 bits per AP (3.2$\pm$5.9% error) and the absolute error was 0.13$\pm$0.21 bits per second
(8.1$\pm$9% error). However, the data was better fit with a saturating exponential (χ_1^2=1.6e3, p<<0.01)
converging to 5.8 bits/AP as it approached the limit due to the finite bin count. There is a substantial
positive bias for the lowest firing rates and smallest number of trials (Figure 2-figure supplement 1E-F)
which has been previously well characterized (Treves and Panzeri, 1995). Thus, the SMGM bits per-AP
metric ($\widehat{I_{AP}^E}$) recovers the ground-truth information well using AP recordings (except at the largest ground
truth information values), with the primary error coming from finite recording time and variable animal
behavior.

We next discuss the changes needed to apply the SMGM bits per AP metric ($\widehat{I_{AP}^E}$) to functional
fluorescence traces and explore the implications of these changes. Most simply, the mean firing rate ($\bar{\lambda}$)
and the mean firing rate in a spatial bin (λ_i) are replaced by the mean change in fluorescence ($\bar{f}$) and the
mean change in fluorescence in a bin (f_i). Making these substitutions in Equation 5 results in the
information as measured by functional fluorescence:

$$\widehat{I_{AP}^F} = \frac{1}{\bar{f}} \sum_{i=1}^N f_i p_X(x_i) \log_2 \frac{f_i}{\bar{f}} \qquad\qquad 8$$

As discussed above, the fluorescence map (f_i) can be approximated as a scaled version of the rate, that is,
$f = c\lambda$ and $\bar{f} = c\bar{\lambda}$. Thus, under this approximation, the c factors in Equation 8 cancel, leading to $\widehat{I_{AP}^F}$
equivalent to $\widehat{I_{AP}^E}$, with the same units of bits/AP. This, of course, ignores the fact that the kernel smooths
the rate map, leading to a bias in the metric that is difficult to quantify analytically.

We then applied the fluorescence SMGM bits per AP metric ($\widehat{I_{AP}^F}$) to our 10,000 mock GCaMP6f traces.
Figure 3A shows the fluorescence traces generated from three mock neurons with ground truth
information of I_{AP}^E=0.05, 1.8 and 4.2 bits/AP. When the fluorescence metric ($\widehat{I_{AP}^F}$) was applied to the
fluorescence traces in these examples, the information recovered was 0.04, 1.8 and 3.5 bits/AP, indicating

some deviations – especially for the highest information neuron. These results held for the 10,000 mock
neuron library (Figure3E-G). At low information values, there was little bias, but at higher information
values the information recovered was substantially lower than the ground truth information. The mean
resulting error was -0.38±0.58 bits/AP (-9.7±27.8%) and absolute error of 0.39 (12.9±26.4%). This error
was better fit with a saturating exponential than a linear fit (χ_1^2=1.6e3, p<<0.01), with the average error
less than 5% up to ground truth information of 1.8 bits/AP and less than 10% up to 3.0 bits/AP. At ground
truth information values higher than 3 bits/AP, the average error was -1.06±0.595 (-22.5±9.44%) and
absolute error was 1.07±0.589 bits/AP (22.6±9.21%). This error persisted even with denser sampling: it
remained consistent even at high firing rates and many trials (Figure 2-figure supplement 1E-F). Thus, the
indicator induces relatively little error at lower information values (<3 bits/AP), but the smoothing effect
of the kernel induces a nonlinear, negative bias to the estimator, particularly at ground truth information
values over 3 bits/AP.

Although the height of the kernel can vary between different functional fluorescence indicators (Figure
2H), these height variations linearly scale the fluorescence map. Thus, since $\widehat{I_{AP}^F}$ involves normalization
by the mean change in fluorescence ($\bar{f}$), $\widehat{I_{AP}^F}$ should not depend on kernel height. To explicitly test this
prediction, we used the 5,000 fluorescence traces described in the previous section (Quantification of the
accuracy and precision of the SMGM bits per second metric using functional fluorescence recordings),
with kernels of varying height (0-3 ΔF/F), but that maintain the same shape and width (from the GCaMP6f
kernel). Then, we measured the percent error in estimating information with $\widehat{I_{AP}^F}$ (Figure 3H-J). Unlike for
the SMGM bits per second metric, the percent error (averaged over all ground truth information values)
in estimating information with $\widehat{I_{AP}^F}$ shows little or no dependence on the height of the kernel (p=0.43), but
a nonlinear dependence on ground truth information as in Figure 3E-G, with no significant difference in
the parameters of the saturating exponential fit (χ_2^2=1.67, p=0.43). Thus, as expected, the percent error in
estimating information with the SMGM bits per AP metric ($\widehat{I_{AP}^F}$) does not vary with the height of the
kernel.

With little effect of kernel height on $\widehat{I_{AP}^F}$, the width of the kernel likely drives biases in the metric. We thus
used the 5,000 fluorescence traces generated from a range of different kernel durations (rise times: 1 ms
to 1 second, fall times longer than the rise time up to 2 seconds; but constant height of the GCaMP6f
kernel) from the previous section (*Quantification of the accuracy and precision of the SMGM bits per*
second metric using functional fluorescence recordings), and then we measured the percent error in
estimating information with $\widehat{I_{AP}^F}$ (Figure 3 K-M). Similarly, as observed above for GCaMP6f and the
varying kernel height examples (Figure 3E-J), the percent error in estimating information with $\widehat{I_S^F}$ shows
a nonlinear dependence on ground truth information (Figure 3E-G). The percent error (averaged over all
ground truth information values) showed a nonlinear response as a function of the width of the kernel
(Figure 3K-L), with a steep increase in error for kernel widths >~1 second. Even for kernel widths <~1
second, the percent error was strongly dependent on the ground truth information value, with steep
increases in error for values >~2.5-3 bits/AP (Figure 3M). Thus, as the kernels gets wider, there is more
negative bias at lower and lower information measured. The resulting errors are thus larger for wider
kernel indicators: for example, with a kernel width the same as gCaMP6s (2.54s), the error exceeds -17%
even at low (<0.25 bits/AP) information, with average errors of -0.86±1.0 bits/AP (-31±19% error) and
absolute errors of 0.87±1.0 bits/AP (32.6±16% error). In contrast, with a kernel width the same as
iGluSnfR (0.52 seconds), the average error exceeded 5% at 3 bits per AP and 10% at 3.7 bits per AP with
a mean error of -0.57±1.00 (-8.0±14%) bits/AP and absolute error of 0.41± 0.56 bits/AP (11±11%). To

estimate the percent errors for the five indicators shown in Figure 2H, taking into account differences in
both height and duration, we used the 5 kernels to generate mock fluorescence traces from the 10k neurons
in Figure3B-G. The resulting distributions, mean and absolute errors, and error thresholds can be seen in
Figure 3-figure supplement 1.

Since the known information values in our library of 10,000 mock neurons were determined using the
SMGM metric, which includes the assumption that neuron firing follows an inhomogeneous Poisson
process, we next investigated whether the biases observed between AP and fluorescence metrics ($\widehat{I_S^E}$ versus
$\widehat{I_S^F}$ and $\widehat{I_{AP}^E}$ versus $\widehat{I_{AP}^F}$; Figures 2 and 3) in our mock neuron datasets were also observed in real neuron
recordings (i.e. real spiking that could deviate from Poisson firing).We therefore measured information in
a real spiking dataset from hippocampal neurons in rats running on a behavioral track (Chen et al., 2016;
Grosmark and Buzsaki, 2016; Grossmark et al., 2016). We generated mock fluorescence traces as we did
with simulated AP trains from our mock neurons, compared the information measured from APs versus
fluorescence ($\widehat{I_S^E}$ versus $\widehat{I_S^F}$ and $\widehat{I_{AP}^E}$ versus $\widehat{I_{AP}^F}$) in the real neuron recordings and found that the biases were
largely consistent with the simulated mock neuron datasets (Figure 5-figure supplement 1).

In summary, we conclude that ground truth information, as measured by the fluorescence SMGM bits per
AP metric ($\widehat{I_{AP}^F}$), retains the units and insensitivity to height scaling of the electrophysiological metric
($\widehat{I_{AP}^E}$), but is nonlinearly biased by the smoothing of the fluorescence map dictated by the width of the
kernel. The estimation errors strongly depended on both the width of the kernel and the information value
being measured. Since these parameters change substantially over the different functional indicators and
different neuron types and behaviors that are commonly used today, they are important factors to consider
when designing and interpreting functional imaging experiments (see below for further discussion).

Nonlinearity introduces further biases
The results presented in the previous two sections rely on the approximation that $\Delta F/F$ scales linearly
with the firing rate, which is not strictly true in practice (Dana et al., 2018; Éltes et al., 2019; Greenberg
et al., 2018). Calcium imaging can be more responsive to bursts of APs rather than isolated spikes, and
saturates at high firing rates. As an example for how to examine how nonlinearities between $\Delta F/F$ and
firing rate could affect the fluorescence SMGM metrics ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$), we applied a log-sigmoid
nonlinearity (Figure 3-figure supplement 3A) to the 10,000 mock GCaMP6f time-series traces described
above, based on the real behavior of GCaMP6f in cultured neurons (Dana et al., 2019, see Methods).
While the resulting measurements (Figure 3-figure supplement 3) of ground truth information, as
measured by the fluorescence SMGM metrics, are largely consistent with the results observed when using
the linear assumption (Figures 2 and 3), some quantitative difference can be seen. Thus, even a relatively
simple nonlinearity between $\Delta F/F$ and firing rate can add distortions to the amount of information
measured using the fluorescence SMGM approach.

Deconvolution may not be sufficient to eliminate biases
The framework presented here for comparing ground truth information with information measured with
the SMGM metrics can be extended to test the efficacy of other strategies for extracting mutual
information. In particular, a perfect AP inference method would alleviate the problems associated with
applying the SMGM metrics to functional fluorescence recordings. To test the utility of such a strategy in
measuring information, we applied a popular deconvolution algorithm, FOOPSI (Vogelstein et al., 2010,
see Methods), to the same 10,000 mock GCaMP6f time-series traces described above. Importantly, this

deconvolution algorithm (and other available algorithms) does not recover traces of relative spike probability or exact spikes times, but instead produces sparse traces with arbitrary units, that have non-zero values estimating the relative 'intensity' of spike production over time (d). This signal can be thought of as a scaled estimate of the number of spikes per time bin, and thus the average intensity map will have some similar properties to the florescence intensity maps – that is, we would expect the intensity maps from deconvolution to approximate the relative firing rate scaled by some factor c, which has arbitrary units.

We then measured information in these deconvolved d-traces using the SMGM metrics ($\widehat{I_S^d}$ and $\widehat{I_{AP}^d}$), which are identical to ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$), except SMGM is applied to d-traces instead of functional fluorescence traces. When using the SMGM bits per second measure ($\widehat{I_S^d}$, Figure 4A), we found a clear scaling of the ground truth information. The scaling factor was very small (c=1.15e-3±1.8e-5 A.U.), resulting in low predicted information (mean error -11.4±6.92 AU, mean % error -99.8±0.38%). This error was significantly larger than when we measured information directly from the florescence traces using $\widehat{I_S^F}$ (Figure 2E; 96.0% absolute error, Ranksum p<<0.01; c=0.0390). It is worth noting that the deconvolved trace d can be arbitrarily scaled, so in a sense this error is arbitrary. However, these are the results from the scaling chosen by a widely used deconvolution algorithm and the large error emphasize that the scale of d can have a large effect on the bits per second measure ($\widehat{I_S^d}$).

Assuming that the intensity map of the deconvolved d-traces are a scaled version of the true rate maps, we could measure information using the SMGM bits per AP metric $\widehat{I_{AP}^d}$ without changing units (Figure 4B). Compared to the SMGM bits per AP metric applied to florescence (I_{AP}^F), on average there was some reduction in the nonlinearity at higher ground truth information (I_{AP}^E) values when using $\widehat{I_{AP}^d}$, resulting in linear fits closer to the unitary line ($\widehat{I_{AP}^d}$ slope $= 1.02 \pm 0.0027, \mathrm{R}^2 = 0.76$ versus $\widehat{I_{AP}^F}$ slope $= 0.78\pm0.0011$, R^2=0.93). However, information measured with $\widehat{I_{AP}^d}$ was still better fit with a saturating exponential (χ_1^2=2.3e3, p~0) converging to a saturation value of 5.51 bits per AP (compared to 5.78 for $\widehat{I_{AP}^F}$), as expected since the algorithm is not expected to resolve spikes at orders of magnitude shorter timescales than the kernel. This resulted in a positive bias at lower levels of ground truth information. For ground truth information values below 3 bits per AP, the average error for $\widehat{I_{AP}^d}$ was 0.556±0.50 bits per AP (68.9±104%) compared to -0.060±0.13 bits per AP (2.75±0.36%) for $\widehat{I_{AP}^F}$. For ground truth information values above 3 bits per AP, the average error for $\widehat{I_{AP}^d}$ was -0.127±0.76 bits per AP (-1.0±16.8%) as compared to -1.04±0.59 bits per AP (-0.22±9.4%) for $\widehat{I_{AP}^F}$. Overall, there was more error when the SMGM bits per AP metric was applied to deconvolved data compared to when applied directly to fluorescence traces ($\widehat{I_{AP}^d}$ mean absolute error 0.60±0.47 bits/AP (52.7±89.1%) vs $\widehat{I_{AP}^F}$ was 0.13±0.21 bits per second (8.1±9% error), Ranksum p<<0.01). Thus, when comparing the recovery of ground truth information from functional fluorescence traces using either direct application of the SMGM metrics ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$) or the application of the SMGM metrics to deconvolved z-traces ($\widehat{I_S^d}$ and $\widehat{I_{AP}^d}$), we found better recovery using the direct application approach ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$).

The KSG and Binned Estimators are poor estimators of MI in functional florescence data
In addition to SMGM, the KSG and Binned Estimation metrics have been developed for estimating mutual information between variables. These other two metrics produce information measured in bits per second,

so they are only comparable to the SMGM bits per second estimator ($\widehat{I_s^F}$). The KSG metric uses the kth
nearest neighbor distances between points in the neural response and behavioral variable space to estimate
information (Kraskov et al., 2004). The Binned Estimation metric uses discrete bins to estimate the full
multidimensional joint probability distribution ($p_{(X,Y)}$ in Equation 1) to estimate mutual information
(Timme and Lapish, 2018). These two metrics estimate information across time samples and therefore are
dependent on firing rate like the SMGM bits/sec metric considered above. Further, the Binned Estimator
is sensitive to the precise method used for data binning, and thus we have used two commonly applied
binning methods: uniform and occupancy based bins.

We applied the KSG, Binned Estimator (Uniform bins), and Binned Estimator (Occupancy binned) (see
Methods) to the same 10,000 mock GCaMP6f time-series traces and behavioral data used to assess the
SMGM approach. These methods all behaved similarly when applied to our simulations (Figure 4C-E),
so they will be discussed together here. The information values measured by these techniques correlated
with ground truth information in bits per second (I_s^E). Interestingly, unlike the SMGM bits per second
metric ($\widehat{I_s^F}$, Figure 2), the KSG and Binned Estimator results were better fit with a saturating exponential
than with a linear fit (χ_1^2=1.70e4, 3.12e4, and 3.14e4 respectively, p~0). The KSG and Binned Estimator
methods overestimated the information at lower ground truth (I_s^E) values and saturated quickly at higher
values. For ground truth information (I_s^E) values below 10 bits per second, the mean absolute errors were
10.3$\pm$7.82 bits per second (465$\pm$4,390%), 22.2$\pm$12.6 bits per second (821.7$\pm$3,353%), and 24.2$\pm$14.0
bits per second (911$\pm$3,926%) for the KSG, Binned Estimator (Uniform bins), and Binned Estimator
(Occupancy binned) respectively (Figure 4F). This is in comparison to the $0.35\pm0.59\frac{bits\,^{\Delta F}/_F}{AP}$ (14$\pm$103%)
mean absolute error found using the SMGM bits per second metric ($\widehat{I_s^F}$) for ground truth information
values less than 10 bits per second. For ground truth information (I_s^E) values greater than 10 bits per
second, the mean absolute errors were 13.2$\pm$8.66 bits per second (85.0$\pm$64.4%), 25.4$\pm$16.2 bits per
second (165.2$\pm$118.3%) and 28.9$\pm$17.34 bits per second (186.3$\pm$124.9%) for the KSG, Binned Estimator
(Uniform bins), and Binned Estimator (Occupancy binned) respectively. This is in comparison to the
0.97$\pm$1.15 bits per second (5.85$\pm$6.73%) mean absolute errors found using the SMGM bits per second
metric ($\widehat{I_s^F}$) for ground truth information values greater than 10 bits per second. Over the full range of
ground truth values, mean absolute errors of 11.9$\pm$8.42 bits per second (254.8%), 23.5$\pm$14.8 bits per
second (456.7%) and 26.7$\pm$16.1 bits per second (509.0%) were found for the KSG, Binned Estimator
(Uniform bins), and Binned Estimator (Occupancy binned) respectively – an order magnitude larger than
the 2.4$\pm$2.97 AU (22$\pm$27% error) error seen using the SMGM bits per second metric ($\widehat{I_s^F}$). As a control,
we applied the Binned Estimators to AP traces and compared the estimated information to the ground
truth information to verify that the large errors observed (Figure 4 D,E) were caused when the estimators
were applied to fluorescence data (rather than simply a difference between the Binned Estimator, which
do not rely on a Poisson firing assumption, and the ground truth information established using SMGM,
which does rely on a Poisson firing assumption). We found the errors when applying the Binned
Estimators to AP traces were relatively small (Figure 4-figure supplement 2, mean absolute error
2.72$\pm$3.38 bits per second (41.1%) and 2.70$\pm$3.33 bits per second (41.0%) for the uniform and occupancy
based binning respectively). Therefore, when comparing the recovery of ground truth information from
functional fluorescence traces using either the SMGM metric $\widehat{I_s^F}$ or the KSG and Binned Estimator metrics,
we found better recovery using the SMGM approach ($\widehat{I_s^F}$).

Guidelines for application of information metrics to functional fluorescence imaging data
Taken together, the above results suggest that across the information metrics applied directly to functional
fluorescence traces, the SMGM metrics provide the most reliable and interpretable information
measurements. We thus suggest the following guidelines for use and interpretation of the SMGM metrics
as applied to fluorescence mutual information metrics ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$) defined in Equations 7 and 8.

The SMGM bits per second metric ($\widehat{I_S^F}$) is likely attractive to imaging researchers because the units suggest
that precise knowledge of AP numbers and times are not required for its use. However, there are several
challenges when applying the SMGM bits per second metric to functional fluorescence imaging data.
First, the substitution of the change in fluorescence map (f) for the AP firing rate map (λ) introduces a
change in units, from bits per second to $\frac{\Delta F}{F} \frac{bits}{AP}$, which is difficult to interpret and relate back to bits per
second. Second, the transformation of AP firing rate to change in fluorescence can be approximated by a
c scaling factor ($f = c\lambda$), which is measured in $\frac{\Delta F/F}{Hz}$, a quantity that is unknown *a priori*. If c is not
consistent between the neurons of a population of interest, then the information values will be scaled
differently and cannot be directly compared (Figure 2). Since c is dependent on the width and height of
the indicator response to a single AP (the kernel), it can vary from neuron to neuron based on difference
in indicator expression level, intracellular calcium buffering, and many other factors (Aponte et al., 2008;
Greenberg et al., 2018; Helmchen and Tank, 2015; Park and Dunlap, 1998). More research will be needed
to measure these parameters (e.g. Chen et al., 2012, 2013a), and thus c, across neurons. Some results
suggest that there may be non-trivial amounts of variability within a populations of neurons (Éltes et al.,
2019; Greenberg et al., 2018). With the impact of c on the SMGM bits per second metric, and the possible
variability of c across a population of neurons, how can researchers properly extract useful measurements
of information using $\widehat{I_S^F}$?

Guideline 1: First, we note that *if* experimental measurements reveal small and acceptable variations in c
across the neurons of interest, then the information values derived from $\widehat{I_S^F}$ can be normalized by this factor
to recover information values in units of SMGM bits per second (independent of $\Delta F/F$) that can be
compared across neurons.

Under the assumption of a consistent kernel, approximations for c for common indicators can be found in
Figure 2-figure supplement 2.

Guideline 2: Further, given small variations in c across the neurons of interest, the ratio of $\widehat{I_S^F}$ between
neurons in the population provides a meaningful metric for comparisons. For example, such ratios could
be used to divide a population of neurons accurately into groups based on their information values (e.g.
three quantiles of information) or compare the information values between different functional subtypes
of neurons (e.g. between place and non-place cells).

Guideline 3: The metric can still be useful even if experimental measurements reveal large and
unacceptable variations in c across the neurons of interest, or if experimental measurements of c do not
exist. In such cases, since it is reasonable to assume that c is consistent in the same neuron over time,
comparisons across the same neuron can provide meaningful insights by using a ratio of $\widehat{I_S^F}$ measured
(from the same neuron) across different conditions. For example, quantifying the neuron by neuron ratio
of $\widehat{I_S^F}$ between different behavioral states or conditions of an animal or task, such as between goal-directed

vs non-goal-directed running down a linear track, would allow researchers to make conclusions such as
the following: "The population of neurons in region Z carries X+/-Y times more information during goal-
directed than non-goal-directed running."

Therefore, we conclude that with careful consideration of the (known or unknown) variability of the
fluorescence response kernel (c), $\widehat{I_S^F}$ can be used to extract useful measurements of information, either
direct measurements of information across a population of neurons (with known and similar c), ratios of
information between different neurons of a population (with known and similar c) or differences across
different conditions within the same neuron (with c unknown or different across neurons).

The SMGM fluorescence bits per AP metric ($\widehat{I_{AP}^F}$) results in the same units as the AP based metric $\widehat{I_{AP}^E}$,
and therefore may provide imaging researchers with information values that are relatively easy to interpret.
However, this similarity in units is somewhat misleading since the number and timing of APs are not
directly measured with functional fluorescence traces and the asymmetric and relatively slow dynamics
of fluorescence indicators leads to shifting and smoothing of the AP rate map (λ). This issue can have the
effect of inducing a significant negative bias in information measurements, especially at high information
values and with functional indicators with wider kernels (Figure 3). This is the most important factor to
consider when determining how researchers can properly extract useful measurements of information
using $\widehat{I_{AP}^F}$. The shifting and smoothing of the AP rate map by fluorescence effectively leads to crosstalk
between adjacent spatial bins. Therefore, it is critical to consider the size of the spatial bins in relation to
the spatial shift and smoothing induced by the indicator (effectively the kernel plotted in space, rather than
time, using the animal's average running velocity to transform from time to space). It is reasonable then
to counteract the spatial shift and smoothing effect by using larger bin sizes, but this only works up to a
point since larger bins limit the maximum amount of information possible to measure and may negatively
bias the information values near this upper limit, even for AP based recordings (Figure 3B-D). Researchers
could potentially optimize the recovery of ground truth information by appropriately selecting bin size for
a particular indicator (see Figure 3-figure supplement 2 A-B).

In practice, using gCaMP6f and the rodent spatial behavior and spatial bin sizes (5cm) used here, our
analysis suggests that $\widehat{I_{AP}^F}$ provides reasonable measurements of information for neurons with values up to
3 bits per AP (Figure 3E-G) since this is the point where the absolute error exceeds 10% (comparable to
the mean absolute error when measuring information from AP data (8.4%)). Equivalent thresholds for
other common indicators are shown in Figure 3-figure supplement 1. The error is exacerbated by slower
indicators and thus more accurate measurements of information will result from using the fastest,
narrowest kernel indicators available, assuming signal to noise and detection efficiency are comparable
across the different width indicators.

Guideline 4: We conclude that with careful consideration of the size of the spatial bins in relation to the
spatial shift and smoothing induced by the indicator, $\widehat{I_{AP}^F}$ can be used to extract useful measurements of
information, most accurately for neurons with < 3 bits/AP under recording conditions similar to those
considered here.

Previous research quantifying information in bits per AP using $\widehat{I_{AP}^E}$ have found that the majority of neurons
carry information in this range (< 3 bits/AP) (Bourboulou et al., 2019; Knierim et al., 1995; Lee et al.,
2006; Markus et al., 1995; Poucet and Sargolini, 2013), with a few exceptions (Ji and Wilson, 2007).

Although these levels are dependent on the number of bins and bin dwell time, $\widehat{I_{AP}^F}$ should be widely
applicable to quantifying information throughout the brain during behavior.

Example: application of information metrics to functional fluorescence imaging data from hippocampus
during spatial behavior

In this section we demonstrate use of the above guidelines for proper application and interpretation of the
SMGM fluorescence mutual information metrics ($\widehat{I_S^F}$ and $\widehat{I_{AP}^F}$) defined in Equations 7 and 8. We applied
these metrics to functional fluorescence recordings (gCaMP6f) from pyramidal neurons in CA1 of the
hippocampus acquired during mouse spatial behavior.

CA1 neurons expressing gCaMP6f (viral transfection, *Camk2a* promoter) were imaged with two-photon
microscopy through a chronic imaging window during mouse navigation along a familiar 1D virtual linear
track, as described previously (Figure 5A-B, Dombeck et al., 2010; Radvansky and Dombeck, 2018;
Sheffield et al., 2017). 8 fields of view from 4 mice were recorded in 8 total sessions (recording duration
8.8+/-1.3 minutes, number of traversals/session: 29+/-2.5, 3.6+/-0.3 laps/min, 3 m long track). From these
8 sessions, 1,500 neurons were identified from our segmentation algorithm (see Methods), and analysis
was restricted to the 964 neurons that displayed at least one calcium transient on at least 1/3 of the
traversals during the session. Among these 964 neurons, 304 (31.5%) had significant place fields and were
thus identified as place cells (see Methods), while the remaining 660 (68.5%) did not pass a place field
test and were thus identified as non-place cells.

By applying Equation 7 (using 5 cm sized spatial bins), we found a continuum of spatial information
values measured by the fluorescence SMGM bits per second metric ($\widehat{I_S^F}$) across the 964 CA1 neurons,
with an average value of $\widehat{I_S^F}=0.14\pm0.0040\ \frac{\text{bits}\ \Delta F/F}{\text{Hz}}$ (Figure 5C). The units for $\widehat{I_S^F}$ make direct use and
interpretation of these values difficult, however, because these recordings were all from pyramidal neurons
in a single area, here for illustrative purposes, we will presume that variations in c (discussed above) across
the 964 neurons of interest are small and acceptable, with the absolute value of c unknown. This allows
for comparisons of information ratios across different subsets of the population. For example, place cells
had 2.8 ± 0.20 times more information than non-place cells using the SMGM bits per second metric
($\widehat{I_S^F}=0.18\pm0.0047$ for place vs. $0.063\pm0.0029\ \frac{\text{bits}\ \Delta F/F}{\text{Hz}}$ for non-place, Rank sum p=1.7e-63, Figure 5D-F),
although there was substantial overlap in information between the populations (see distributions in Figure
5D and individual examples in Figure 5E,F). This also allows for accurate division of the 964 neurons
into 3 quantiles based on information values, which we use below for spatial location decoding.

By applying Equation 8 (using 5 cm sized spatial bins), we found a continuum of spatial information
values measured by the fluorescence SMGM bits per AP metric ($\widehat{I_{AP}^F}$) across the 964 CA1 neurons, with
an average value of $\widehat{I_{AP}^F}=1.65\pm0.023$ bits/AP (Figure 5C). The units for $\widehat{I_{AP}^F}$ allow for direct use and
interpretation of these values, and notably, because most (97%) of the neurons had values less than 3
bits/AP, a mean absolute error of <10% can be assumed across the distribution of SMGM bits per AP
values. When applied to the place and non-place cell populations, we found that place cells had higher
information than nonplace cells using the fluorescence SMGM bits per AP metric ($\widehat{I_{AP}^F}=1.8\pm0.026$ and
1.35 ± 0.042 bits/AP, Rank sum p=4.6e-21, Figure 5D-F). This is consistent with mock fluorescence traces
generated from real neuron AP datasets (Figure 5-figure supplement 1B).

As a demonstration of the usefulness of using information metrics to analyze large functional fluorescence
population recordings, we explored the accuracy of decoding the animal's track position using different
subsets of neurons. We divided the 964 neurons into 9 groups: All neurons, place cells, non-place cells,
three quantiles based on the fluorescence SMGM bits per second metric, and three quantiles based on the
fluorescence SMGM bits per AP metric. We then used a Bayesian decoder of the animals' position (see
Methods) separately for each of the 9 neuron groups in each of the 8 sessions (Figure 5G,H). An individual
session decoding example can be seen in Figure 5G. We quantified decoding accuracy using the absolute
position decoding error (% of track), and pooled this measure across sessions for each neuron group
(Figure 5H). The means and standard errors for each group are: All neurons (7.33 +/-2.5 %), place cells
(6.97 +/-1.9 %), non-place cells (20.9 +/-1.8 %), SMGM bits per second Q1 (21.9 +/-1.5 %), SMGM bits
per second Q2 (13.2 +/-2.4 %), SMGM bits per second Q3 (8.97 +/-2.4 %), SMGM bits per AP Q1 (17.6
+/-2.7%), SMGM bits per AP Q2 (17.8 +/-3.1 %), SMGM bits per AP Q3 (10.4 +/-3.0 %). Interestingly,
even the lowest quantile information groups still could be used to determine animal track location to within
~1/5 of the track. This supports the idea that the hippocampal code for space is carried by a large
population of active neurons (Meshulam et al., 2016), and not just by a select subpopulation with the
highest information or most well-defined tuning curves. As could be expected, place cells encoded the
position of the animal better than nonplace cells and better than the lowest quantile information groups
(Holm-Bonferroni corrected Rank sum, α=0.05) and neurons in the higher quantiles provided more
accurate decoding. Thus, the fluorescence information metrics provide a means to compare the relative
contribution of hippocampal neurons with different information values to decoding animal position.

Discussion

Here, we performed an in-depth simulation study to examine the application of the SMGM bits per second
and SMGM bits per AP metrics of mutual information to functional fluorescence recordings. Since these
metrics were designed for AP recordings and since functional fluorescence recordings violate some of the
assumptions that these metrics are based on, it was unclear if and how the metrics could be used for
functional fluorescence recordings. We created a library of ten thousand mock neurons whose AP output
carried ground-truth amounts of information about the animal's spatial location, and by using real
behavioral recording data from mice navigating in virtual linear tracks, we simulated the spatial firing
patterns of the mock neurons. We then simulated fluorescent calcium responses for each neuron in each
session by convolving the AP trains with calcium kernels for different indicators, primarily GCamp6f
(though see Supplemental Figures 2 and 3 for results from other indicators), and then added noise.

We then derived fluorescence versions of the SMGM bits per second ($\widehat{I_s^F}$) and SMGM bits per AP metrics
($\widehat{I_{AP}^F}$) (Equations 7 and 8) and applied them to the fluorescence traces in order to quantify the performance
of the metrics for estimating information. We found that ground truth information, as measured by the
fluorescence SMGM bits per second metric ($\widehat{I_s^F}$), was transformed into different units and was linearly
scaled by a factor (c) dependent on the height and width of the kernel, with c linearly dependent on height
and nonlinearly dependent on width. The error induced by these transformations changed substantially
over the range of kernel values of the different functional indicators widely used today, and therefore are
important factors to consider when designing and interpreting functional imaging experiments. We then
found that ground truth information, as measured by the fluorescence SMGM bits per AP metric ($\widehat{I_{AP}^F}$),
retains the units and insensitivity to height scaling of the electrophysiological metric ($\widehat{I_{AP}^E}$), but is
nonlinearly biased by the smoothing of the fluorescence map dictated by the width of the kernel. The
estimation errors strongly depended on both the width of the kernel and the information value being

measured. Importantly, since these parameters change substantially over the different functional indicators
and different neuron types and behaviors that are commonly used today, they are important factors to
consider when designing and interpreting functional imaging experiments. For example, even for the same
indicator, the shape of the kernel is a function of intracellular calcium buffering, indicator concentration,
the amount of calcium influx, the efflux rates, background fluorescence and resting calcium concentration,
which can all vary across different cells. Additionally, the results presented here rely on the approximation
that $\Delta F/F$ scales linearly with the firing rate, which is not strictly true in practice. We show in Figure 3-
figure supplement 3 that even a relatively simple nonlinearity between $\Delta F/F$ and firing rate can add
distortions to the amount of information measured using the fluorescence SMGM approach. This
relationship between $\Delta F/F$ and firing rate can vary across different indicators and, since the Toolbox can
be used to vary this relationship, users can further explore this source of bias.

Some of the general features of the relationship between ground truth information and fluorescence
SMGM metrics can also be seen using an analytical approximation. For example, if we approximate the
rate map as a Gaussian firing field with mean rate $\bar{\lambda}$, simplify the kernel approximation to a single
exponential with falloff τ, assume constant, normalized movement speed v, and assume that the
convolution between the kernel and the mean rate map is nearly Gaussian, we can approximate the
relationship between the bits per second ground truth information I_S^E and measured fluorescence

information I_S^F as $I_S^F \approx -\dfrac{Av\tau\bar{\lambda}\log\left(4^{-\frac{I_S^E}{\bar{\lambda}}}+2\pi ev^2\tau^2\right)}{\log(4)}$ and between the bits per AP ground truth information

I_{AP}^E and measured fluorescence information I_{AP}^F as $I_{AP}^F \approx -\dfrac{\log\left(4^{-I_{AP}^E}+2e\pi v^2\tau^2\right)}{\log(4)}$. Similar to the numerical
simulations, these equations predict that the fluorescence bits per second metric is dominated by a
prefactor ($A\tau v\bar{\lambda}$ in the analytical case), and that the fluorescence bits per action potential metric
saturates at larger information values. Using 1000 values for I_{AP}^E and λ from the same distribution as our
simulations, we can approximate a c value for I_S^F of 0.011 $\frac{\Delta F/F}{Hz}$ (data not shown). Of course, our
numerical solutions provide more accurate approximations for the magnitude of these effects (and for
the magnitude of information values themselves) given that they include the more accurate double
exponential kernel (leading to a much larger kernel integral compared to a single exponential), signal
noise, and the realistic nonstationary speed, position and fluorescence signals.

In our approach, the known information values in our library of 10,000 mock neurons were determined
using the SMGM metric, which includes the assumption that neuron firing follows an inhomogenous
Poisson process. It is important to remember that the SMGM metric, which has been applied to spiking
data extensively over the past few decades, *requires* the use of a Poisson estimate of spiking probability—
i.e. the Poisson assumption is built into the original metric. In practice, even spiking data violates this and
other assumptions of the SMGM metric since real neurons do not strictly follow Poisson statistics (for
example they can display neural hysteresis) and animal behavior is non-stationary. Here we are building
from this existing framework and adding and testing whether it is possible to apply the metric to functional
fluorescence datasets. Even still, the Poisson assumption could have contributed to some of the biases
found when evaluating the fluorescence SMGM metrics with respect to ground truth information. We
explored this potential source of bias further using two different analyses. First, in Figure 4-figure
supplement 2, we applied the Binned Estimators (which do not rely on a Poisson firing assumption) to AP
traces and compared the estimated information to the ground truth information (which was established

using the SMGM metric that does rely on a Poisson firing assumption). We found the errors to be relatively
small, particularly in comparison to the errors induced by the Binned Estimators when applied to
fluorescence traces (Figure 4 D,E). Second, in Figure 5-figure supplement 1, we used a real spiking dataset
from hippocampal neurons in mice running on a behavioral track (i.e. real spiking neurons that can deviate
from Poisson firing) and generated mock fluorescence traces from the AP traces. When we compared the
information measured from the AP traces to the fluorescence traces, we found biases that were largely
consistent with those observed in Figure 2 and 3 from our simulated mock neuron datasets. Taken together,
these analyses indicate that any biases resulting from the Poisson assumption in the simulation procedure
appear to be small, particularly with respect to the biases introduced when AP traces are transformed into
functional fluorescence traces. Finally, in the Toolbox, we also include code to generate mock neurons
using a binned distribution, avoiding the Poisson assumption of SMGM. Thus users can further explore
sources of bias using a different ground truth dataset.
Using our mock fluorescence traces, we also asked if an AP estimation method could relieve the biases in
the SMGM metrics. Applying the SMGM bits per second metric ($\widehat{I_s^d}$) to AP estimation traces from a
deconvolution algorithm (FOOPSI) resulted in a low c value for recovered versus ground truth
information. When the SMGM bits per AP measure was applied ($\widehat{I_{AP}^d}$), the resulting measurements of
information were still nonlinear (compared to $\widehat{I_s^F}$), with a positive bias at lower values of ground truth
information. Overall, applying FOOPSI to fluorescence traces led to a poorer recovery of ground truth
information using SMGM compared to direct application of SMGM to the florescence traces ($\widehat{I_{AP}^F}$).
Importantly, this result from deconvolution is only specific to GCaMP6f, and conclusions should not be
drawn about other indicators or situations; users will be able to use the Toolbox to explore this area further.
We also tested other metrics to measure mutual information directly from the fluorescence time traces
(KSG, Binned Estimator (uniform bins), and Binned Estimator (occupancy binned)) and found these
alternatives produced highly variable, saturating measurements of recovered versus ground truth
information. This was in contrast to the SMGM bits per second measure ($\widehat{I_s^F}$) which produced a linearly
scaled bias with lower error.

Taken together, we find that the SMGM bits per AP metric can well recover the mutual information
between spiking and behavior. The SMGM bits per second metric is scaled such that comparisons should
be limited to within populations of well characterized neurons or for within neuron comparisons, e.g.
ratios of information across conditions. In general, researchers should use caution when applying measures
developed for AP data in fluorescence recordings: there's no guarantee that the assumptions that support
the measures hold for fluorescence data, and this can lead to difficult to interpret and biased results.

Materials and Methods

Toolbox and Data Availability
We provide a user toolbox (freely available at https://github.com/DombeckLab/infoTheory), which
consists of Matlab functions to generate libraries of model neurons with known amounts of information,
to generate spiking or fluorescence time-series from those model neurons, and to estimate neuron
information from real or model spiking or fluorescence time-series datasets using the three metrics
considered here (SMGM, Binned Estimator, KSG). This toolbox also contains tools to generate mock
neurons using a binned distribution, avoiding the Poisson assumption of SMGM. Behavioral data used to
generate the random traces is freely available at https://doi.org/10.7910/DVN/SCQYKR.

 Construction of AP trains with known ground truth information
To construct mock neurons with ground truth information, we adapted the differential form of the AP
information, in bits per AP (Equation 6). To create a rate map, we first selected an average firing rate and
target ground truth information. The mean rate ($\bar{\lambda}$) was always between 0.1 and 30 Hz, the information in
bits per AP (I_{AP}^E) between 0 and 6 bits per AP, and the information in bits per second (I_S^E) between 0 and
24. To more evenly sample each of these, we first randomly selected the bits per second (I_S^E) or bits per
AP (I_{AP}^E) to target. If the information target was in bits per AP, both the information (I_{AP}^E) and mean firing
rate ($\bar{\lambda}$) were chosen uniformly. Because the information in bits per second $I_S^E = \bar{\lambda}I_{AP}^E$, the bits per second
information was not uniformly sampled in this case. If the target was to be in bits per second, both the bits
per AP (I_{AP}^E) and SMGM bits per second (I_S^E) measures were first chosen uniformly. Because the rate $\bar{\lambda} =$
I_S^E/I_{AP}^E, this was not chosen uniformly. This procedure was repeated to maintain the bounds on $\bar{\lambda}$, resulting
in a non-uniform sampling of information. The final distribution (Figure 1C) was spread acceptably for
further analysis.

The rate maps were constructed by spline interpolating across 5 control points with two anchored at each
end of the track, and taking the exponential for each point, and then normalizing by the numerically
calculated integral (Figure 1A, D). To create a map matching the target information, we began with a
random spline. The 'y' (relative rate) initial position of each node was chosen from a standard normal
distribution and the initial 'x' (track position) of the 3 center nodes was chosen uniformly. The nodes were
then systematically moved using the MATLAB built in optimizer 'fmincon' with constraints preventing
the crossing of the center nodes and keeping them on the track, and the 'OptimalityTolerance' option set
to 0 (Figure 1A). This was accomplished using the 'genExpSpline' function' in the toolbox.

We then randomly selected behavioral traces (see Methods *Behavior* section) and concatenated sessions
until a total time randomly chosen between 3 and 60 minutes was reached (Figure 1E, F; Figure 2A, Figure
3A). This was accomplished using the 'loadBehaviorT' function in the toolbox. The track positions were
normalized and used to build a conditional intensity function (CIF) from the rate function above. The CIF
was normalized to match an expected mean rate over the entire session, and the MATLAB built-in
'poissrnd' function was used to generate AP times, sampled at 1 kHz. The was accomplished using the
'genSpikeTrain' function in the toolbox. Finally, the AP times were binned according to the counts within
mock imaging frames sampled at 30 Hz.

Simulated $\frac{\Delta F}{F}$ traces

To construct the $\frac{\Delta F}{F}$ traces (Figure1E,J,K; Figure2A; Figure3A), we first created a single AP response
kernel from the peak-normalized sum of two exponentials:

$$g(t) = \frac{e^{-a\,t} - e^{-b\,t}}{\left(\frac{a}{b}\right)^{\frac{a}{b-a}} - \left(\frac{a}{b}\right)^{\frac{b}{b-a}}} \qquad\qquad 9$$

Where t is the time since the AP and a and b are chosen to minimize $\left(1 - g(\tau_{rise})\right)^2 +$
$\left(0.5 - g(\tau_{rise} + \tau_{fall})\right)^2$ where τ_{rise} is the rise time in seconds and τ_{fall} is the half-fall time in seconds.

Deviations in τ_{rise} and τ_{fall} from baseline were also measured. The kernel g(t) was then multiplied by the
indicator height. The kernel parameters were generated using the 'fluorescenceKernel' function, and
evaluated using the 'doubleExp' function in the toolbox.

The GCaMP6f, GCaMP6s, and jRGECO1a heights, rise and fall times were measured as responses to
single APs *in vivo* (Chen et al., 2013; Dana et al., 2019)*:* other kernels (Figure 2H, Supplemental Figures
2 and 3) were approximated from other experiments presented in the references (seen in Table 1).

Table 1. Properties of indicator kernels used

Kernel	Height $\Delta F/F$	Rise (s)	Fall (s)	Source
gCaMP6f	0.190	0.042	0.142	(Chen et al., 2013)
jRGECO1a	0.164	0.041	0.207	(Kalko et al., 2011)
gCaMP7f	0.560	0.063	0.276	(Dana et al., 2019)
gCaMP6s	0.230	0.179	0.550	(Chen et al., 2013)
iGluSnfR-A184S	0.300	0.022	0.106	(Marvin et al., 2018)

To define the width of the kernel (Figure2L-N, Figure3K-M), we considered the kernel as a low pass
filtered version of the APs. If we normalize the filter to mean 1, it has the Fourier transform
$\left(\frac{1}{a+2\pi f\hat{\jmath}} - \frac{1}{b+2\pi f\hat{\jmath}}\right)\left(\frac{ab}{b-a}\right)$. The kernel width was defined as the -3 dB (50%) cutoff period of this filter:
$f^{-1} = \frac{\sqrt{-a^2-b^2+\sqrt{a^4+14a^2b^2+b^4}}}{2\pi\sqrt{2}}$. For the simulations with different width kernels, a kernel width was
chosen between 0.01 and 10 seconds, a rise time between 0.001 and 1 second, and a fall time between the
rise time and 2 seconds. Then, a and b were chosen to minimize the squared error between these three
targets using the built in MATLAB optimizer 'fminsearch'.

White noise with a standard deviation of $0.15\frac{\Delta F}{F}$ was then added to the mock fluorescence traces.

Nonlinearity
In our linear simulations used throughout this work, the fluorescence kernels associated with a fast
sequence of action potentials were approximated to sum linearly. In real cultured neurons, a summation
nonlinearity has been observed such that sequences of action potentials do not generate a linear summation
in ΔF/F (Dana et al., 2019). To simulate this nonlinearity, the $\frac{\Delta F}{F}$ trace was then further transformed as:

$$\frac{\Delta F'}{F} = sign\left(\frac{\Delta F}{F}\right) * \frac{6.264}{1 + e^{-3.251\,Re\left(\log_{10}\left(\frac{\Delta F}{F}\right)\right)}}$$

This equation was arrived at by fitting the measured responses in Dana et al, 2019, Figure 2C, which can
be compared to the nonlinearity used here (Figure 3-figure supplement 3A).

Deconvolution
Deconvolution was performed using the previously described FOOPSI algorithm (Friedrich et al., 2017;
Vogelstein et al., 2010). The regularization coefficient was set at 0.02154, which maximized the
correlation between the deconvolved trace and the true spike train in a random sample of 500 simulated
traces: all other parameters were optimized for each trace. Because the example regularization
coefficient provided by Friedrich et al, 2017 was 2.4, we also measured information values at 100
different values for the regularization coefficient between 0 and 3; this had little effect on the measured
information (Figure 4-figure supplement 1).

KSG estimator
The previously described second KSG estimator (Kraskov et al., 2004) was used using the 5^{th} nearest
neighbor distance.

Binned estimators
The binned mutual information estimators were used (Timme and Lapish, 2018). The activity trace was
divided into 10 bins, either evenly across the span of the activity (uniform binned) or variably so the
bins contained the same number of samples (occupancy binned). Position was similarly divided into 60
bins.

Bayesian decoding
The Bayesian decoder used here (Figure 5 G,H) was adapted from a previously described method (Zhang
et al., 1998). Decoding was performed on the likelihood that a significant transient occurred in a time
frame, trained on the first 80% of the session and tested on the last 20%. The session was divided into
Δt=0.1 second bins. The conditional likelihood that an animal is in position x_i given the number of active
frames during a time window (n) is $p(x_i|n) = p_X(x_i) \left(\prod_{j=1}^{M} f_{i,j}^{n_j} \right) e^{-\Delta t \sum_{j=1}^{M} f_{i,j}}$, Where $p_X(x_i)$ is the
(marginal) probability that the animal is in the i^{th} spatial bin during a time sample, $f_{i,j}$ is the average rate
of significant frames by the j^{th} neuron in the i^{th} spatial bin, n_j is the number of significant frames
observed during the time window in neuron j, and M is the total number of neurons. The decoded position
was selected as the one with maximum conditional likelihood.

Animals
10 to 12 week old male C57BL/6 mice (20-30g) were individually housed under a reverse 12 hr. light /
dark cycle, all experiments were conducted during the dark phase. All experiments were approved by the
Northwestern University Animal Care and Use committee.

Behavior
We used a previously described virtual reality set-up and task (Heys et al., 2014; Sheffield et al., 2017;
Sheffield and Dombeck, 2015); some of the behavior sessions used here has previously appeared in these
studies. Briefly, water scheduled, head fixed mice were trained to run on a cylindrical treadmill down a
3m virtual track to receive a water (4 ul) reward at the end of the track, and were subsequently teleported
to the beginning of the track after a 1.5 s delay. Behavioral sessions were included if the animal ran at
least 20 laps containing a continuous 40 cm run for which the velocity was over 7 cm/sec during a 5-30
minute session.

Mouse surgery and virus injected
We performed population calcium imaging of CA1 neurons as described previously(Sheffield et al., 2017;
Sheffield and Dombeck, 2015). Briefly, 30 nL of AAV1-SynFCaMP6f (University of Pennsylvania
Vector Core, 1.5*10^13 GC/ml) was injected through a small craniotomy over the right hippocampus (1.8
mm lateral, 2.3 mm caudal of Bregma; 1.25 mm below the surface of the brain) under Isoflurane (1-2%)
anesthesia. 7 days later, a hippocampal window and head plate was implanted as described previously
(Dombeck et al., 2010).

Two-photon imaging
Imaging was performed as previously described(Sheffield et al., 2017; Sheffield and Dombeck, 2015).
Scanimage 4 was used for microscope control and acquisition (Pologruto et al., 2003). Time series movies
(1024 or 512x256 pixels) were acquired at 50 Hz. A Digidata1440A (Molecular Devices) with Clampex
10.3 synchronized position on the linear track, reward timing, and the timing of image frames.

Image processing, ROI selection and calcium transient analysis
Images were processed as previously described (Sheffield et al., 2017; Sheffield and Dombeck, 2015),
with minor modifications. Briefly, rigid motion correction was performed using cross correlation as in
(Dombeck et al., 2010; Miri et al., 2011; Sheffield and Dombeck, 2015), but here using a Fast-Fourier
transform approximation on the full video. ROIs were defined as previously described (Mukamel et al.,
2009) (mu=0.6, 150 principal/independent components, s.d. threshold = 2.5, s.d. smoothing width=1, area
limits = 100-1200 pixels). $\frac{\Delta F}{F}$ traces were generated by normalizing around the 8th percentile of a 3 second
sliding window. Significant transients from both experimental and mock fluorescence traces were selected
by comparing the ratio of amplitudes and durations of positive to negative going transients with a false
positive rate <0.01% (Dombeck et al., 2010). Mock traces used the histograms generated from the mock
gCaMP6f traces (Figure 2-3) or from the specific matching indicator traces (Supplemental Figures 2,3):
experimental data histograms were built separately. All subsequent analyses were run using these
significant transients.

Behavior analysis
The mean virtual track velocity was defined as the total virtual track distance covered during the session
divided by the total duration of the session; slow and stop periods were included in this metric. All other
analyses were restricted to long running periods, where the animal exceeded a virtual track velocity of 4
cm/s and ran continuously for at least 40 cm.

Defining place fields
Place fields were defined by first creating the spatial fluorescence intensity map (f_i) with the 300 cm track
divided into 60, 5 cm bins. This map was smoothed via a 3 bin boxcar. Transients identified during run
periods were shuffled in order and to random intervals to create 1000 bootstrapped intensity maps.
Candidate fields were defined as regions of the original fluorescence map with values greater than 99%
of the bootstrapped maps. Fields were then retained if they were between 20 and 120 cm wide: significant
place cells retained at least one field that satisfied these criteria.

Figures

Figure 1. Procedures for generating a library of 10,000 neurons with known amounts of information. (A) Five splines with a gradient of ground truth information (I_{AP}^E) representing the steps in generating a continuous rate map ($\lambda(x)$) matching the desired target information, in this case 2 bits/AP. Red X's indicate control nodes that were moved to change the shape of the spline and minimize the squared error to the target information. (B) Cross section of the error surface around the solution point as a function of the position of node 3, and the trajectory taken by the solver to minimize the error and arrive at the target. (C) Histograms of ground truth information resulting from repeating the procedure in A-B 10,000 times to target a range of ground truth information values in bits per second (I_S^E) and bits per AP (I_{AP}^E).(D) Splines representing $\lambda(x)$ at the solution point for a low (I_{AP}^E=0.04 bits per AP, left) and high (I_{AP}^E, 2 bits per AP, right) information neuron. (E) Steps to generate mock AP and functional fluorescence data. (1) An example real behavior trace from a mouse running on a linear track that was used to generate the simulated spiking. (2) The behavior in combination with the rate maps generated in A-D were used to generate an instantaneous firing rate trace. (3) The instantaneous rate was used to pseudorandomly generate APs, as shown in this mock raster. (4) The AP raster was convolved with the GCaMP6f kernel (red, inset) and noise was added to generate a mock $\frac{\Delta F}{F}$ trace. (5) Large numbers of these traces were generated and used to assess the effects of many simulation parameters on the estimators. (F-L) Spiking and fluorescence activity patterns generated from the example simulated neurons shown in D and using a

mean firing rate of 1 Hz. (F) Behavioral trace in blue with AP raster shown in red. (G) Lap-by-lap raster of the neurons' firing vs mouse track position. (H) Lap by lap binned, firing rates vs mouse track position for the neurons (I) AP raster (red) and mock calcium traces for the same behavioral period shown in F. (J) Lap by lap mean binned fluorescence vs mouse position for the neurons. (K) Binned average firing rate (λ_i, black) and fluorescence intensity (f_i, green) maps for the two neurons. These maps were used for information analyses.

Figure 2. Quantification of the precision of the SMGM bits per second metric using APs or functional fluorescence recordings. (A) Three representative mock neurons spanning the range of ground truth information values in bits per second (I_S^E). From top to bottom for each: mouse track position vs time, AP

raster, fluorescence calcium trace (green), and firing rate map (λ_i, black) and change in fluorescence map
(f_i, green). (B-D) The ground truth bits per second values are well recovered when measured from AP
traces. (B) Information measured from AP data using the SMGM bits per second metric ($\widehat{I_s^E}$) vs ground
truth information (I_s^E). Each dot is a single mock neuron, the gray dashed line is the unity line (perfect
measurement), the pink line is the line of best fit. Red circles show the examples in A. (C) Percentage
error for the information measurements shown in B. (D) Heat map of percentage error measurements
shown in C. Black lines are 2 standard deviations, the white line is the mean. (E-G) Effects of applying
the SMGM bits per second metric to fluorescence traces. (E) Information measured from mock GCaMP6f
traces using the SMGM bits per second metric ($\widehat{I_s^F}$) vs ground truth information (I_s^E). (F) Percentage error
for the information measurements shown in E. (G) Heat map of percentage error measurements shown in
F. (H) Representative mock kernels mimicking responses from different indicators. (I-K) The effect of
kernel height on estimating ground truth information (I_s^E) using the SMGM bits per second metric ($\widehat{I_s^F}$).
Kernel height for the kernels shown in H are indicated by colored triangles (I) Percentage error as a
function of kernel height (J) Heat map of percentage error measurements shown in I with mean (white)
and 2 standard deviations (black). (K) The average percentage error as a function of kernel height and
ground truth information in SMGM bits per second (I_s^E). (L-N) The effect of kernel width on estimating
ground truth information (I_s^E) using the SMGM bits per second metric ($\widehat{I_s^F}$). Kernel widths for the kernels
shown in H are indicated by colored triangles. (L) Percentage error as a function of kernel width (M) Heat
map of percentage error measurements shown in L with mean (white) and 2 standard deviations (black).
(N) The average percentage error as a function of kernel width.

<u>Figure 2-figure supplement 1.</u> Effect of recording density on information metrics. A. The mean error (top)
and absolute error (bottom) between the ground truth information (I_S^E) and measured information in
SMGM bits per second from mock APs ($\widehat{I_S^E}$) as a function of the mean firing rate ($\bar{\lambda}$). B. As A, but as a
function of number of laps. C-D. As A-B, but for measurements from mock fluorescence traces. E-F. As
A-B, but for the SMGM bits per AP metric. G-H.) As C-D, but for the SMGM bits per AP metric.

Indicator	Error (AU)	Absolute Error (AU)	% Error	Absolute % Error	$c\left(\frac{\Delta F/F}{Hz}\right)$
jRGECO1a	-10.9±6.7	10.9±6.6	-94.8±8.5	95.0±6.1	0.0482
gCaMP7f	-9.12±5.9	9.17±5.8	-77.6±36.6	80.0±31.0	0.200
gCaMP6s	-9.81±6.2	9.84±6.1	-84.5±24.8	85.8±19.8	0.140
iGluSnfR-A184S	-10.9±6.6	10.9±6.6	-94.7±8.9	94.9±6.5	0.0490

<u>Figure 2-figure supplement 2.</u> Effects of applying the SMGM bits per second metric to fluorescence traces
from different common functional indicators. Top: Information measured from mock traces using the
SMGM bits per second metric ($\widehat{I_S^F}$) vs ground truth information (I_S^E). Each dot is a single mock neuron,
the gray dashed line is the unity line (perfect measurement), the pink line is the line of best fit. Middle:
Percentage error density plots. The white line is the mean, the black are $\pm$ one standard deviation. Bottom:
Summary statistics and estimates for the scaling factor c.

Figure 3. Quantification of the precision of the SMGM bits per AP metric using APs or functional fluorescence recordings. (A) Three representative mock neurons spanning the range of ground truth information values in bits per AP (I_{AP}^E). From top to bottom for each: mouse track position vs time, AP raster, fluorescence calcium trace (green), and firing rate map (λ_i, black) and change in fluorescence map (f_i, green). (B-D) The ground truth bits per AP values are well recovered when measured from AP traces. (B) Information measured from AP data using the SMGM bits per AP metric ($\widehat{I_{AP}^E}$) vs ground truth information (I_{AP}^E). Each dot is a single mock neuron, the gray dashed line is the unity line (perfect measurement). Red circles show the examples in A. (C) Percentage error for the information measurements shown in B. (D) Heat map of percentage error measurements shown in C. Black lines are 2 standard deviations, the white line is the mean. (E-G) Effects of applying the SMGM bits per AP metric to fluorescence traces. (E) Information measured from mock GCaMP6f traces using the SMGM bits per AP metric ($\widehat{I_{AP}^F}$) vs ground truth information (I_{AP}^E). (F) Percentage error for the information measurements

shown in E. (G) Heat map of percentage error measurements shown in F. (H-J) The effect of kernel height
on estimating ground truth information (I_{AP}^E) using the SMGM bits per second metric ($\widehat{I_{AP}^F}$). Kernel height
for the kernels shown in Figure2H are indicated by colored triangles (H) Percentage error as a function of
kernel height (I) Heat map of percentage error measurements shown in H with mean (white) and 2 standard
deviations (black). (J) The average percentage error as a function of kernel height and ground truth
information in bits per AP (I_{AP}^E). (K-M) The effect of kernel width on estimating ground truth information
(I_{AP}^E) using the SMGM bits per AP metric ($\widehat{I_{AP}^F}$). Kernel widths for the kernels shown in Figure 2H are
indicated by colored triangles. (L) Percentage error as a function of kernel width (M) Heat map of
percentage error measurements shown in L with mean (white) and 2 standard deviations (black). (N) The
average percentage error as a function of kernel width.

Indicator	Error (bits/AP)	Absolute Error (bits/AP)	% Error	Absolute % Error	5% cutoff (bits/AP)	10% Cutoff (bits/AP)
jRGECO1a	-0.56±0.76	0.61±0.73	-10.4±25.5	18.5±20.4	1.68	2.40
gCaMP7f	-0.79±0.87	0.80±0.86	-20.4±47.8	23.6±46.2	0.48	1.44
gCaMP6s	-1.2±1.1	1.2±1.1	-25.3±194.9	43.7±191.6	0.48	0.72
iGluSnfR-A184S	-0.32±0.53	0.36±0.5	-5.7±17.9	11.2±15.2	2.6	3.4

1009

Figure 3-figure supplement 1. Effects of applying the SMGM bits per AP metric to fluorescence traces
from different common functional indicators. Top: Information measured from mock traces using the
SMGM bits per AP metric ($\widehat{I_{AP}^F}$) vs ground truth information (I_{AP}^E). Each dot is a single mock neuron, the
gray dashed line is the unity line (perfect measurement), the pink line is the exponential fit. Middle:
Percentage error density plots. The white line is the mean, the black are ± one standard deviation. Bottom:
Summary statistics and percent error cutoffs.

1016

<u>Figure 3-figure supplement 2.</u> Effect of number of bins on the SMGM metrics. A. The mean percentage
error (top) and the standard deviation (bottom) for the bits per second measure ($\widehat{I_S^F}$) applied to 10,000
mock gCamp6f traces, with ground truth information on the X-axis and 2-60 bins (3m track) on the Y-
axis. B. As A, for the bits per AP measure ($\widehat{I_{AP}^F}$). C. The mean percentage error (top) and the standard
deviation (bottom) for the bits per second measure ($\widehat{I_S^F}$) applied to 20,000 mock fluorescence traces with
differing kernel width. Kernel width is on the X-axis, number of bins is on the Y-axis. D. As C, but for
the bits per AP measure ($\widehat{I_{AP}^F}$).

1024

Figure 3-figure supplement 3. Effects of a sigmoid nonlinearity between ΔF/F and firing rate. Here we applied a log-sigmoid nonlinearity to the 10,000 mock GCaMP6f time-series traces and then measured information using the fluorescence SMGM metrics. (A) Nonlinearity applied to AP-to-florescence trace transformation (B) Information measured from AP data using the SMGM bits per second ($\widehat{I_S^F}$) vs ground truth information (I_S^E). Each dot is a single mock neuron, the gray dashed line is the unity line (perfect measurement). (C) Percentage error for the information measurements shown in B. (D) Heat map of percentage error measurements shown in C. Black lines are 2 standard deviations, the white line is the mean. (E-G) As B-D, but for the bits per AP measured ($\widehat{I_{AP}^F}$) versus the ground truth information in bits per AP (I_{AP}^E).

Method	Units	Source	Mean error	Mean abs error	Mean % error	Mean abs % err	C	p exponential	saturation level	90% saturation point
SMGM bits/sec	bits ΔF/F /AP	Skaggs et al, 1993	-11.1	11.1	-96.0	96.0	0.0390	0.760	39.7	2.3e3
SMGM bits/AP	bits/AP	Skaggs et al, 1993	-0.38	0.4	-9.7	12.9	0.78	<<0.01	5.78	11.5
foopsi+SMGM bits/sec	AU	Vogelstein et al, 2010	-11.4	11.4	-99.8	99.8	1.15e-3	0.129	0.02	32.0
Foopsi+SMGM bits/ap	bits/AP	Vogelstein et al, 2010	0.3	0.6	46.3	52.7	1.02	<<0.01	5.51	6.6
KSG	bits/sec	Kraskov et al, 2004	9.9	11.9	37.9	254.8	1.64	<<0.01	29.2	13.9
Binned MI (uniform bins)	bits/sec	Timm and Lapish, 2018	12.8	14.1	285.7	292.5	1.77	<<0.01	29	8.47
Binned MI (occupancy bins)	bits/sec	Timm and Lapish, 2018	15.0	15.6	322.5	325.5	1.94	<<0.01	32.2	9.26

1034

Figure 4. Alternative techniques for measuring mutual information from functional fluorescence traces. (A-E) (Top) Information measured from mock GCaMP6f traces vs ground truth information. The gray line is the unity line, the pink line is the best fit saturating exponential. (Middle) Percentage error for the information measurements shown on top. (Bottom) Heat map of percentage error measurements shown in middle. (A) FOOPSI deconvolved traces using the SMGM bits per second metric ($\widehat{I_s^d}$). (B) FOOPSI

1040   deconvolved traces using the SMGM bits per AP metric ($\widehat{I^d_{AP}}$). (C) The KSG measure applied to GCaMP6f
1041   traces. (D) The Binned Estimator applied to GCaMP6f traces using uniform bins. (E) The Binned
1042   Estimator applied to GCaMP6f traces using equal occupancy bins. (F) Table of summary statistics for
1043   each measure. P exponential is the p-value from the chi^2 test used to determine if a saturating exponential
1044   fit is better than a linear fit for the measured vs ground truth information plots.

1045

1046   Figure 4-figure supplement 1. Effect of changing the regularization coefficient in deconvolution (Friedrich
1047   et al., 2017; Vogelstein et al., 2010) on the measured information. (A) The measured information using
1048   the fluorescence bits per second metric applied to 10,000 mock gCamP6f traces using regularization
1049   coefficients between 0 and 3. (B) As A, but for the fluorescence bits per AP metric.

Figure 4-figure supplement 2. The binned estimator applied to AP traces and then compared to ground truth information. (Top) Information measured from mock AP traces vs ground truth information. The gray line is the unity line, the pink line is the best fit saturating exponential. (Middle) Percentage error for the information measurements shown on top (same scale as shown in Figure 4). (Bottom) Heat map of percentage error measurements shown in middle. (A) The Binned Estimator applied to AP traces using uniform bins. (B) The Binned Estimator applied to AP traces using equal occupancy bins.

1057

Figure 5. Application of SMGM information metrics to functional fluorescence imaging data from hippocampus during spatial behavior. A.) Example field of hippocampal pyramidal neurons expressing GCaMP6f and imaged during linear track navigation. Active cell ROIs shown in yellow; traces for green cells shown in B. B.) Fluorescence DF/F traces (green) from two neurons in the field shown in A and the track position during the recording (blue). C.). Distribution of information values using the fluorescence SMGM bits per second metric ($\widehat{I_S^F}$,(top) and the fluorescence SMGM bits per AP metric ($\widehat{I_{AP}^F}$, bottom). The gray line indicates the recommended cutoff for reliability using GCaMP6f. D.) Plot of $\widehat{I_S^F}$ vs $\widehat{I_{AP}^F}$ for each neuron. Place cells indicated in red and nonplace cells in blue. E.) Example non-place cells spanning the information ranges shown in C. Spatial fluorescence map (f_i) shown on left, and average change in fluorescence per track traversal on right. F.) Same as E, but for place cells. G.) Bayesian decoding of mouse's track position using different subpopulations of neurons for one example session. From top to bottom: All active neurons, all nonplace cells, and place cells, the first through third quantiles of the

SMGM bits per second formulation ($\widehat{I_S^F}$), and the first through third quantiles of the SMGM bits per AP formation ($\widehat{I_{AP}^F}$). The white dashed line indicates the ground truth position of the animal, the color map indicates the decoded position probability (peak-normalized posterior distribution). H.) Decoding accuracy (absolute position decoding error in units of % of track) pooled over all sessions for each neuron group indicated in G. Black bars indicate significant differences by Holm-Bonferroni corrected rank sum tests (α=0.05).

Figure 5-figure supplement 1. The SMGM estimators as applied to real AP data from a real spiking dataset from hippocampal neurons in mice running on a behavioral track (Chen et al., 2016; Grosmark and Buzsaki, 2016; Grossmark et al., 2016). (A) Example real place cell. From top to bottom: rat track position vs time, real AP raster, mock fluorescence calcium trace generated from real AP trace by convolving APs with GCaMP6f kernel and adding noise (green), and firing rate map (λ_i, black) and change in mock fluorescence map (f_i, green). (B)Plot of $\widehat{I_S^F}$ vs $\widehat{I_{AP}^F}$ for each neuron. Place cells indicated in red and nonplace cells in blue. (C) The SMGM bits per second metric applied to the real AP traces ($\widehat{I_S^E}$) versus the mock fluorescence traces ($\widehat{I_S^F}$) (generated from the real AP traces). (D) The percentage difference between the SMGM bits per second metric applied to the real AP traces ($\widehat{I_S^E}$) and the mock fluorescence traces ($\widehat{I_S^F}$). The mean and standard deviation are indicated in black. In magenta, the mean and standard deviation of 5,000 of the mock neuron traces seen in Figures 2 and 3, sampled to have the same firing rates as the real neurons and shortened to the same mean session duration. (E) Density plot for the data shown in D. (F-H) As C-E, but for the bits per AP metric ($\widehat{I_{AP}^E}$ versus $\widehat{I_{AP}^F}$).

References

Ahrens MB, Orger MB, Robson DN, Li JM, Keller PJ. 2013. Whole-brain functional imaging at cellular resolution using light-sheet microscopy. *Nat Methods* **10**:413–420. doi:10.1038/nmeth.2434

Allen TA, Salz DM, McKenzie S, Fortin NJ. 2016. Nonspatial Sequence Coding in CA1 Neurons. *J Neurosci* **36**:1547–1563. doi:10.1523/JNEUROSCI.2874-15.2016

Aponte Y, Bischofberger J, Jonas P. 2008. Efficient Ca2+ buffering in fast-spiking basket cells of rat hippocampus. *J Physiol* **586**:2061–2075. doi:10.1113/jphysiol.2007.147298

Aronov D, Tank DW. 2014. Engagement of Neural Circuits Underlying 2D Spatial Navigation in a Rodent Virtual Reality System. *Neuron* **84**:442–456. doi:10.1016/j.neuron.2014.08.042

Belghazi MI, Baratin A, Rajeswar S, Ozair S, Bengio Y, Courville A, Hjelm RD. 2018. MINE: Mutual Information Neural Estimation. *35th Int Conf Mach Learn ICML 2018* **2**:864–873.

Bourboulou R, Marti G, Michon FX, El Feghaly E, Nouguier M, Robbe D, Koenig J, Epsztein J. 2019. Dynamic control of hippocampal spatial coding resolution by local visual cues. *Elife* **8**:1–30. doi:10.7554/eLife.44487

Brandon MP, Bogaard AR, Libby CP, Connerney MA, Gupta K, Hasselmo ME. 2011. Reduction of Theta Rhythm Dissociates Grid Cell Spatial Periodicity from Directional Tuning. *Science (80-)* **332**:595–599. doi:10.1126/science.1201652

Brown EN, Barbieri R, Eden UT, Frank L. 2003. Likelihood Methods for Neural Spike Train Data Analysis In: Feng J, editor. Computational Neuroscience: A Comprehensive Approach. CRC Pres. pp. 252–281.

Buzsáki G, Mizuseki K. 2014. The log-dynamic brain: how skewed distributions affect network operations. *Nat Rev Neurosci* **15**:264–78. doi:10.1038/nrn3687

Calton JL, Stackman RW, Goodridge JP, Archey WB, Dudchenko PA, Taube JS. 2003. Hippocampal place cell instability after lesions of the head direction cell network. *J Neurosci* **23**:9719–9731. doi:23/30/9719 [pii]

Chen Q, Cichon J, Wang W, Qiu L, Lee SJR, Campbell NR, DeStefino N, Goard MJ, Fu Z, Yasuda R, Looger LL, Arenkiel BR, Gan WB, Feng G. 2012. Imaging Neural Activity Using Thy1-GCaMP Transgenic Mice. *Neuron* **76**:297–308. doi:10.1016/j.neuron.2012.07.011

Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS. 2013. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**:295–300. doi:10.1038/nature12354

Chen Z, Grosmark AD, Penagos H, Wilson MA. 2016. Uncovering representations of sleep-associated hippocampal ensemble spike activity. *Sci Rep* **6**:32193. doi:10.1038/srep32193

Climer JR, DiTullio R, Newman EL, Hasselmo ME, Eden UT. 2015. Examination of rhythmicity of extracellularly recorded neurons in the entorhinal cortex. *Hippocampus* **25**:460–473.

doi:10.1002/hipo.22383

Climer JR, Newman EL, Hasselmo ME. 2013. Phase coding by grid cells in unconstrained
environments: two-dimensional phase precession. *Eur J Neurosci* **38**:2526–41.
doi:10.1111/ejn.12256

Cohen MR, Kohn A. 2011. Measuring and interpreting neuronal correlations. *Nat Neurosci* **14**:811–819.
doi:10.1038/nn.2842

Dana H, Sun Y, Mohar B, Hulse B, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R, Macklin JJ,
Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS, Hod Dana, Yi
Sun, Boaz Mohar, Brad Hulse, Jeremy P Hasseman, Getahun Tsegaye, Arthur Tsang, Allan Wong,
Ronak Patel, John J Macklin, Yang Chen, Arthur Konnerth, Vivek Jayaraman, Loren L Looger,
Eric R Schreiter, Karel Svoboda DSK. 2018. High-performance GFP-based calcium indicators for
imaging activity in neuronal populations and microcompartments. *bioRxiv* **16**:434589.
doi:10.1101/434589

Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R,
Macklin JJ, Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS.
2019. High-performance calcium sensors for imaging activity in neuronal populations and
microcompartments. *Nat Methods* **16**:649–657. doi:10.1038/s41592-019-0435-6

DeWeese MR, Zador AM, Hromádka T. 2008. Sparse Representation of Sounds in the Unanesthetized
Auditory Cortex. *PLoS Biol* **6**:14. doi:10.1371/journal.pbio.Citation

Dombeck DA, Harvey CD, Tian L, Looger LL, Tank DW. 2010. Functional imaging of hippocampal
place cells at cellular resolution during virtual navigation. *Nat Neurosci* **13**. doi:10.1038/nn.2648

Dombeck DA, Khabbaz AN, Collman F, Adelman TL, Tank DW. 2007. Imaging large-scale neural
activity with cellular resolution in awake, mobile mice. *Neuron* **56**:43–57.
doi:10.1016/j.neuron.2007.08.003

Éltes T, Szoboszlay M, Kerti-Szigeti K, Nusser Z. 2019. Improved spike inference accuracy by
estimating the peak amplitude of unitary [Ca 2+] transients in weakly GCaMP6f-expressing
hippocampal pyramidal cells. *J Physiol* **597**:2925–2947. doi:10.1113/JP277681

Evans MH, Petersen RS, Humphries MD. 2019. On the use of calcium deconvolution algorithms in
practical contexts. *bioRxiv* 871137. doi:10.1101/871137

Frank LM, Brown EN, Wilson MA. 2001. A comparison of the firing properties of putative excitatory
and inhibitory neurons from CA1 and the entorhinal cortex. *J Neurophysiol* **86**:2029–2040.

Friedrich J, Zhou P, Paninski L. 2017. Fast online deconvolution of calcium imaging data. *PLOS*
Comput Biol **13**:e1005423. doi:10.1371/journal.pcbi.1005423

Fu H, Rodriguez GA, Herman M, Emrani S, Nahmani E, Barrett G, Figueroa HY, Goldberg E, Hussaini
SA, Duff KE. 2017. Tau Pathology Induces Excitatory Neuron Loss, Grid Cell Dysfunction, and
Spatial Memory Deficits Reminiscent of Early Alzheimer's Disease. *Neuron* **93**:533-541.e5.

doi:10.1016/j.neuron.2016.12.023

Fyhn M, Molden S, Hollup S, Moser M-B, Moser EI. 2002. Hippocampal Neurons Responding to First-
Time Dislocation of a Target Object. *Neuron* **35**:555–566. doi:10.1016/S0896-6273(02)00784-5

Gao W, Kannan S, Oh S, Viswanath P. 2017. Estimating mutual information for discrete-continuous
mixturesAdvances in Neural Information Processing Systems. pp. 5987–5998.

Gerrard JL, Burke SN, McNaughton BL, Barnes CA. 2008. Sequence Reactivation in the Hippocampus
Is Impaired in Aged Rats. *J Neurosci* **28**:7883–7890. doi:10.1523/JNEUROSCI.1265-08.2008

Greenberg DS, Wallace DJ, Voit K-M, Wuertenberger S, Czubayko U, Monsees A, Handa T,
Vogelstein JT, Seifert R, Groemping Y, Kerr JN. 2018. Accurate action potential inference from a
calcium sensor protein through biophysical modeling. *bioRxiv* 479055. doi:10.1101/479055

Grosmark AD, Buzsaki G. 2016. Diversity in neural firing dynamics supports both rigid and learned
hippocampal sequences. *Science (80-)* **351**:1440–1443. doi:10.1126/science.aad1935

Grossmark AD, Long J, Buzsáki G. 2016. Recordings from hippocampal area CA1, PRE, during and
POST novel spatial learning. doi:dx.doi.org/10.6080/K0862DC5

Grubb MS, Thompson ID. 2006. Quantitative Characterization of Visual Response Properties in the
Mouse Dorsal Lateral Geniculate Nucleus. *J Neurophysiol* **90**:3594–3607.
doi:10.1152/jn.00699.2003

Hainmueller T, Bartos M. 2018. Parallel emergence of stable and dynamic memory engrams in the
hippocampus. *Nature* **558**:292–296. doi:10.1038/s41586-018-0191-2

Harvey CD, Collman F, Dombeck DA, Tank DW. 2009. Intracellular dynamics of hippocampal place
cells during virtual navigation. *Nature* **461**:941–946. doi:nature08499 [pii]10.1038/nature08499

Hazama Y, Tamura R. 2019. Effects of self-locomotion on the activity of place cells in the hippocampus
of a freely behaving monkey. *Neurosci Lett* **701**:32–37. doi:10.1016/j.neulet.2019.02.009

Helmchen F, Tank DW. 2015. A single-compartment model of calcium dynamics in nerve terminals and
dendrites. *Cold Spring Harb Protoc* **2015**:155–167. doi:10.1101/pdb.top085910

Heys JG, Dombeck DA. 2018. Evidence for a subcircuit in medial entorhinal cortex representing
elapsed time during immobility. *Nat Neurosci* **21**:1574–1582. doi:10.1038/s41593-018-0252-8

Heys JG, Rangarajan K V., Dombeck DA. 2014. The functional micro-organization of grid cells
revealed by cellular-resolution imaging. *Neuron* **84**:1079–1090. doi:10.1016/j.neuron.2014.10.048

Hinman JR, Brandon MP, Climer JR, Chapman GW, Hasselmo ME. 2016. Multiple Running Speed
Signals in Medial Entorhinal Cortex. *Neuron* **91**:666–679. doi:10.1016/j.neuron.2016.06.027

Hok V, Chah E, Save E, Poucet B. 2013. Prefrontal Cortex Focally Modulates Hippocampal Place Cell
Firing Patterns. *J Neurosci* **33**:3443–3451. doi:10.1523/JNEUROSCI.3427-12.2013

Huang Y, Brandon MP, Griffin AL, Hasselmo ME, Eden UT. 2009. Decoding movement trajectories
through a T-maze using point process filters applied to place field data from rat hippocampal region
CA1. *Neural Comput* **21**:3305–3334. doi:10.1162/neco.2009.10-08-893

Hubel DH, Wiesel TN. 2009. Republication of The Journal of Physiology (1959) 148, 574-591:
Receptive fields of single neurones in the cat's striate cortex. 1959. *J Physiol* **587**:2721–32.
doi:10.1113/jphysiol.2009.174151

Ji D, Wilson MA. 2007. Coordinated memory replay in the visual cortex and hippocampus during sleep.
Nat Neurosci **10**:100–107. doi:10.1038/nn1825

Jiaying Tang SS. 2015. Visual receptive field properties of neurons in the mouse LGN. *PLoS Comput*
Biol **18**:449–452. doi:10.5061/dryad.b2t22

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018a. Dendritic Inhibition in the Hippocampus Supports Fear Learning. *Nat Biotechnol*
36:726–737. doi:10.1126/science.1247485

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018b. State-Dependent Subnetworks of Parvalbumin-Expressing Interneurons in Neocortex.
Nat Biotechnol **36**:726–737. doi:10.1016/j.celrep.2019.02.005

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018c. Calcium Dynamics in Dendrites of Hippocampal CA1 Interneurons in Awake Mice.
Nat Biotechnol **36**:726–737. doi:10.3389/fncel.2019.00098

Jing M, Zhang P, Wang G, Feng J, Mesik L, Zeng J, Jiang H, Wang S, Looby JC, Guagliardo NA,
Langma LW, Lu J, Zuo Y, Talmage DA, Role LW, Barrett PQ, Zhang LI, Luo M, Song Y, Zhu JJ,
Li Y. 2018d. A genetically encoded fluorescent acetylcholine indicator for in vitro and in vivo
studies. *Nat Biotechnol* **36**:726–737. doi:10.1038/nbt.4184

Kalko EK V, Dukas R, Ratcliffe JM, Teeling EC, Haven N, Fattu JM, Bates ME, Simmons J a,
Riquimaroux H, Surlykke A, Bouffard FH, Lee DN, Dear SP, Horiuchi TK, Krishnaprasad PS,
Moss CF, Schuller G, Brudzynski SM, Syme D a, Hollingworth S, Lindstedt SL, Baylor SM, Mead
a F, Rome LC, Goller F, Spiers ILY, Leeuwen JL Van, Schachat F, Rossmanith GH, Hoh JFY,
Kelley DB, Vater M a, Zhao Y, Araki S, Wu J, Teramoto T, Chang Y-F, Nakano M, Abdelfattah
AS, Fujiwara M, Ishihara T, Nagai T, Campbell RE. 2011. An Expanded Palette of Genetically
Encoded Ca2+ Indicators. *Science (80-)* **333**:1888–1891. doi:10.1126/science.1208592

Keller PJ, Ahrens MB. 2015. Visualizing whole-brain activity and development at the single-cell level
using light-sheet microscopy. *Neuron* **85**:462–483. doi:10.1016/j.neuron.2014.12.039

Khoshkhoo S, Vogt D, Sohal VS. 2017. Dynamic, Cell-Type-Specific Roles for GABAergic
Interneurons in a Mouse Model of Optogenetically Inducible Seizures. *Neuron*.
doi:10.1016/j.neuron.2016.11.043

Kinsky NR, Sullivan DW, Mau W, Hasselmo ME, Eichenbaum HB. 2018. Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales. *Curr Biol* **28**:3578-3588.e6. doi:10.1016/j.cub.2018.09.037

Knierim JJ, Kudrimoti HS, McNaughton BL. 1995. Place cells, head direction cells, and the learning of landmark stability. *J Neurosci* **15**:1648–1659.

Koenig J, Linder AN, Leutgeb JK, Leutgeb S. 2011. The Spatial Periodicity of Grid Cells Is Not Sustained During Reduced Theta Oscillations. *Science (80-)* **332**:592–595. doi:10.1126/science.1201685

Kraskov A, Stögbauer H, Grassberger P. 2004. Estimating mutual information. *Phys Rev E - Stat Physics, Plasmas, Fluids, Relat Interdiscip Top* **69**:16. doi:10.1103/PhysRevE.69.066138

Kraus BJ, Brandon MP, Robinson RJ, Connerney MA, Hasselmo ME, Eichenbaum HB. 2015. During Running in Place, Grid Cells Integrate Elapsed Time and Distance Run. *Neuron* **88**:578–589. doi:10.1016/j.neuron.2015.09.031

Kropff E, Carmichael JE, Moser M-B, Moser EI. 2015. Speed cells in the medial entorhinal cortex. *Nature* **523**:419–424. doi:10.1038/nature14622

Lee I, Griffin AL, Zilli EA, Eichenbaum HB, Hasselmo ME. 2006. Gradual translocation of spatial correlates of neuronal firing in the hippocampus toward prospective reward locations. *Neuron* **51**:639–650.

Leveteau J, MacLeod P. 1966. Olfactory discrimination in the rabbit olfactory glomerulus. *Science* **153**:175–6. doi:10.1126/science.153.3732.175

Liu P, Jerrard LE, Bilkey DK, Jarrard LE, Bilkey DK. 2004. Excitotoxic lesions of the pre- and parasubiculum disrupt the place fields of hippocampal pyramidal cells. *Hippocampus* **14**:107–116. doi:10.1002/hipo.10161

MacDonald CJ, Carrow S, Place R, Eichenbaum HB. 2013. Distinct hippocampal time cell sequences represent odor memories in immobilized rats. *J Neurosci* **33**:14607–16. doi:10.1523/JNEUROSCI.1537-13.2013

Mankin EA, Thurley K, Chenani A, Haas O V, Debs L, Henke J, Galinato M, Leutgeb JK, Leutgeb S, Leibold C. 2019. The hippocampal code for space in Mongolian gerbils. *Hippocampus* 1–15. doi:10.1002/hipo.23075

Mann K, Gallen CL, Clandinin TR. 2017. Whole-Brain Calcium Imaging Reveals an Intrinsic Functional Network in Drosophila. *Curr Biol* **27**:2389-2396.e4. doi:10.1016/j.cub.2017.06.076

Markus EJ, Qin YL, Leonard B, Skaggs WEW, McNaughton BBL, Barnes CA. 1995. Interactions between location and task affect the spatial and directional firing of hippocampal neurons. *J Neurosci* **15**:7079–7094. doi:20026608

Marvin JS, Scholl B, Wilson DE, Podgorski K, Kazemipour A, Müller JA, Schoch S, Quiroz FJU,

Rebola N, Bao H, Little JP, Tkachuk AN, Cai E, Hantman AW, Wang SSH, DePiero VJ, Borghuis
BG, Chapman ER, Dietrich D, DiGregorio DA, Fitzpatrick D, Looger LL. 2018. Stability, affinity,
and chromatic variants of the glutamate sensor iGluSnFR. *Nat Methods* **15**:936–939.
doi:10.1038/s41592-018-0171-3

Marvin JS, Shimoda Y, Magloire V, Leite M, Kawashima T, Jensen TP, Kolb I, Knott EL, Novak O,
Podgorski K, Leidenheimer NJ, Rusakov DA, Ahrens MB, Kullmann DM, Looger LL, Malgoire V,
Leite M, Kawashima T, Jensen TP, Knott EL, Novak O, Podgorski K, Leidenheimer NJ, Rusakov
DA, Ahrens MB, Kullmann DM, Looger LL. 2019. A genetically encoded fluorescent sensor for in
vivo imaging of GABA. *Nat Methods* **16**:763–770. doi:10.1038/s41592-019-0471-2

Mau W, Sullivan DW, Kinsky NR, Hasselmo ME, Howard MW, Eichenbaum H. 2018. The Same
Hippocampal CA1 Population Simultaneously Codes Temporal Information over Multiple
Timescales. *Curr Biol* **28**:1499-1508.e4. doi:10.1016/j.cub.2018.03.051

Meshulam L, Gauthier JL, Brody CD, Tank DW, Bialek W. 2016. Collective behavior of place and non-
place neurons in the hippocampal network 1–17.

Miri A, Daie K, Burdine RD, Aksay E, Tank DW. 2011. Regression-Based Identification of Behavior-
Encoding Neurons During Large-Scale Optical Imaging of Neural Activity at Cellular Resolution. *J*
Neurophysiol **105**:964–980. doi:10.1152/jn.00702.2010

Morris TP, White IR, Crowther MJ. 2019. Using simulation studies to evaluate statistical methods. *Stat*
Med **38**:2074–2102. doi:10.1002/sim.8086

Mukamel EA, Nimmerjahn A, Schnitzer MJ. 2009. Automated Analysis of Cellular Signals from Large-
Scale Calcium Imaging Data. *Neuron* **63**:747–760. doi:10.1016/j.neuron.2009.08.009

Newman EL, Climer JR, Hasselmo ME. 2014. Grid cell spatial tuning reduced following systemic
muscarinic receptor blockade. *Hippocampus* **24**:643–655. doi:10.1002/hipo.22253

Nguyen JP, Shipley FB, Linder AN, Plummer GS, Liu M, Setru SU, Shaevitz JW, Leifer AM. 2016.
Whole-brain calcium imaging with cellular resolution in freely behaving Caenorhabditis elegans .
Proc Natl Acad Sci **113**:E1074–E1081. doi:10.1073/pnas.1507110112

Nguyen MN, Hori E, Matsumoto J, Tran AH, Ono T, Nishijo H. 2013. Neuronal responses to face-like
stimuli in the monkey pulvinar. *Eur J Neurosci* **37**:35–51. doi:10.1111/ejn.12020

Nguyen MN, Matsumoto J, Hori E, Maior RS, Tomaz C, Tran AH, Ono T, Nishijo H. 2014. Neuronal
responses to face-like and facial stimuli in the monkey superior colliculus. *Front Behav Neurosci*
8:1–18. doi:10.3389/fnbeh.2014.00085

Niell CM, Stryker MP. 2008. Highly Selective Receptive Fields in Mouse Visual Cortex. *J Neurosci*
28:7520–7536. doi:10.1523/jneurosci.0623-08.2008

O'Connor DH, Peron SP, Huber D, Svoboda K. 2010. Neural activity in barrel cortex underlying
vibrissa-based object localization in mice. *Neuron* **67**:1048–1061.
doi:10.1016/j.neuron.2010.08.026

O'Keefe J. 1976. Place units in the hippocampus of the freely moving rat. *Exp Neurol* **51**:78–109.

Østergaard J, Kramer MA, Eden UT. 2018. Capturing Spike Variability in Noisy Izhikevich Neurons
Using Point Process Generalized Linear Models. *Neural Comput* **30**:125–148.
doi:10.1162/neco_a_01030

Pachitariu M, Stringer C, Harris KD. 2018. Robustness of Spike Deconvolution for Neuronal Calcium
Imaging. *J Neurosci* **38**:7976–7985. doi:10.1523/JNEUROSCI.3339-17.2018

Paninski L. 2004. Maximum likelihood estimation of cascade point-process neural encoding models.
Netw Comput Neural Syst **15**:243–262. doi:10.1088/0954-898X_15_4_002

Park D, Dunlap K. 1998. Dynamic regulation of calcium influx by G-proteins, action potential
waveform, and neuronal firing frequency. *J Neurosci* **18**:6757–66.

Park E, Dvorak D, Fenton AA. 2011. Ensemble Place Codes in Hippocampus: CA1, CA3, and Dentate
Gyrus Place Cells Have Multiple Place Fields in Large Environments. *PLoS One* **6**:e22349.
doi:10.1371/journal.pone.0022349

Park IJ, Bobkov Y V., Ache BW, Principe JC. 2013. Quantifying bursting neuron activity from calcium
signals using blind deconvolution. *J Neurosci Methods* **218**:196–205.
doi:10.1016/j.jneumeth.2013.05.007

Pastalkova E, Itskov V, Amarasingham A, Buzsáki G. 2008. Internally generated cell assembly
sequences in the rat hippocampus. *Science (80-)* **321**:1322–7. doi:10.1126/science.1159775

Pologruto TA, Sabatini BL, Svoboda K. 2003. ScanImage: Flexible software for operating laser
scanning microscopes. *Biomed Eng Online* **2**:13. doi:10.1186/1475-925X-2-13

Poucet B, Sargolini F. 2013. A trace of your place. *Science (80-)* **340**:35–36.
doi:10.1126/science.1237567

Radvansky BA, Dombeck DA. 2018. An olfactory virtual reality system for mice. *Nat Commun* **9**:839.
doi:10.1038/s41467-018-03262-4

Rashid SK, Pedrosa V, Dufour MA, Moore JJ, Chavlis S, Delatorre RG, Poirazi P, Clopath C, Basu J.
2020. The dendritic spatial code: branch-specific place tuning and its experience-dependent
decoupling. *bioRxiv* **10**:2020.01.24.916643. doi:10.1101/2020.01.24.916643

Robbe D, Buzsáki G. 2009. Alteration of theta timescale dynamics of hippocampal place cells by a
cannabinoid is associated with memory impairment. *J Neurosci* **29**:12597–605.
doi:10.1523/JNEUROSCI.2407-09.2009

Roxin A, Brunel N, Hansel D, Mongillo G, van Vreeswijk C. 2011. On the Distribution of Firing Rates
in Networks of Cortical Neurons. *J Neurosci* **31**:16217–16226. doi:10.1523/jneurosci.1677-
11.2011

Scholl B, Wilson DE, Fitzpatrick D. 2017. Local Order within Global Disorder: Synaptic Architecture of

Visual Space. *Neuron* **96**:1127-1138.e4. doi:10.1016/j.neuron.2017.10.017

Shafi M, Zhou Y, Quintana J, Chow C, Fuster J, Bodner M. 2007. Variability in neuronal activity in primate cortex during working memory tasks. *Neuroscience* **146**:1082–1108. doi:10.1016/j.neuroscience.2006.12.072

Sheffield MEJ, Adoff MD, Dombeck DA. 2017. Increased Prevalence of Calcium Transients across the Dendritic Arbor during Place Field Formation. *Neuron* **96**:490-504.e5. doi:10.1016/j.neuron.2017.09.029

Sheffield MEJ, Dombeck DA. 2015. Calcium transient prevalence across the dendritic arbour predicts place field properties. *Nature* **517**:200–204. doi:10.1038/nature13871

Simonnet J, Brecht M. 2019. Burst Firing and Spatial Coding in Subicular Principal Cells. *J Neurosci* **39**:3651–3662. doi:10.1523/jneurosci.1656-18.2019

Skaggs WE, McNaughton BL, Gothard KM. 1993. An Information-Theoretic Approach to Deciphering the Hippocampal Code In: Hanson SJ, Cowan JD, Giles CL, editors. Advances in Neural Information Processing Systems 5. Morgan-Kaufmann. pp. 1030–1037.

Soo FS, Schwartz GW, Sadeghi K, Berry MJ. 2011. Fine Spatial Information Represented in a Population of Retinal Ganglion Cells. *J Neurosci* **31**:2145–2155. doi:10.1523/jneurosci.5129-10.2011

Stackman RW, Taube JS. 1998. Firing properties of rat lateral mammillary single units: head direction, head pitch, and angular head velocity. *J Neurosci* **18**:9020–9037.

Stirman JN, Smith IT, Kudenov MW, Smith SL. 2016. Wide field-of-view, multi-region, two-photon imaging of neuronal activity in the mammalian brain. *Nat Biotechnol* **34**:857–862. doi:10.1038/nbt.3594

Stringer C, Pachitariu M, Steinmetz N, Carandini M, Harris KD. 2019. High-dimensional geometry of population responses in visual cortex. *Nature* **571**:361–365. doi:10.1038/s41586-019-1346-5

Timme NM, Lapish C. 2018. A Tutorial for Information Theory in Neuroscience. *eneuro* **5**:ENEURO.0052-18.2018. doi:10.1523/ENEURO.0052-18.2018

Treves A, Panzeri S. 1995. The Upward Bias in Measures of Information Derived from Limited Data Samples. *Neural Comput* **7**:399–407. doi:10.1162/neco.1995.7.2.399

Vogelstein JT, Packer AM, Machado TA, Sippy T, Babadi B, Yuste R, Paninski L. 2010. Fast Nonnegative Deconvolution for Spike Train Inference From Population Calcium Imaging. *J Neurophysiol* **104**:3691–3704. doi:10.1152/jn.01073.2009

Wachowiak M, Shipley MT. 2006. Coding and synaptic processing of sensory information in the glomerular layer of the olfactory bulb. *Semin Cell Dev Biol* **17**:411–423. doi:10.1016/j.semcdb.2006.04.007

Wilent WB, Nitz DA. 2007. Discrete place fields of hippocampal formation interneurons. *J*
Neurophysiol **97**:4152–4161. doi:10.1152/jn.01200.2006

Yaksi E, Friedrich RW. 2006. Reconstruction of firing rate changes across neuronal populations by
temporally deconvolved Ca2+ imaging. *Nat Methods* **3**:377–383. doi:10.1038/nmeth874

Yartsev MM, Ulanovsky N. 2013. Representation of Three-Dimensional Space in the Hippocampus of
Flying Bats. *Science (80-)* **340**:367–372. doi:10.1126/science.1235338

Zhang K, Ginzburg I, McNaughton BL, Sejnowski TJ. 1998. Interpreting neuronal population activity
by reconstruction: unified framework with application to hippocampal place cells. *J Neurophysiol*
79:1017–1044.

Zhou J-L, Shatskikh TN, Liu X, Holmes GL. 2007. Impaired single cell firing and long-term
potentiation parallels memory impairment following recurrent seizures. *Eur J Neurosci* **25**:3667–
3677. doi:10.1111/j.1460-9568.2007.05598.x

Zinyuk L. 2000. Understanding hippocampal activity by using purposeful behavior: Place navigation
induces place cell discharge in both task-relevant and task-irrelevant spatial reference frames. *Proc*
Natl Acad Sci **97**:3771–3776. doi:10.1073/pnas.050576397

Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, Gamal A El, Schnitzer MJ. 2013. Long-
term dynamics of CA1 hippocampal place codes. *Nat Neurosci* **16**:264–266. doi:10.1038/nn.3329

1390