Brief sensory deprivation triggers plasticity of neurotransmitter-synthesising enzyme

expression in genetically labelled olfactory bulb dopaminergic neurons

Darren J Byrne, Marcela Lipovsek, Matthew S Grubb

Centre for Developmental Neurobiology, Institute of Psychiatry, Psychology and Neuroscience

(IoPPN), King's College London, UK

Correspondence: matthew.grubb@kcl.ac.uk

Abstract

In the glomerular layer of the olfactory bulb, local dopaminergic interneurons play a key role in

regulating the flow of sensory information from nose to cortex. These dual dopamine- and GABA-

releasing cells are capable of marked experience-dependent changes in the expression of

neurotransmitter-synthesising enzymes, including tyrosine hydroxylase (TH). However, such

plasticity has most commonly been studied in cell populations identified by their expression of the

enzyme being studied, and after long periods of sensory deprivation. Here, instead, we used brief 1-

or 3-day manipulations of olfactory experience in juvenile mice, coupled with a conditional genetic

approach that labelled neurons contingent upon their expression of the dopamine transporter (DAT-

tdTomato). This enabled us to evaluate the potential for faster changes in neurotransmitter-

synthesising enzyme expression in an independently identified population of neurons. Our labelling

strategy showed good specificity for olfactory bulb dopaminergic neurons, whilst also revealing a

minority sub-population of non-dopaminergic DAT-tdTomato cells that expressed the calcium-

binding protein calretinin. Crucially, the proportions of these neuronal subtypes were not affected

by brief alterations in sensory experience. Short-term olfactory manipulations also produced no

significant changes in immunofluorescence for the GABA-synthesising enzyme GAD67. However, in

bulbar DAT-tdTomato neurons brief sensory deprivation was accompanied by a transient drop in

immunofluorescence for the dopamine-synthesising enzyme dopa decarboxylase (DDC), and a

sustained decrease in TH expression. Careful characterisation of an independently identified,

genetically labelled neuronal population therefore enabled us to uncover experience-dependent

changes in neurotransmitter-synthesising enzyme expression that are more rapid than previously

appreciated.

Introduction

Dopaminergic neurons in the mammalian olfactory bulb form a neurochemically-distinct population of glomerular layer interneurons (Hökfelt *et al.*, 1975; Kosaka & Kosaka, 2007, 2011). They are dual-transmitter cells, releasing both dopamine and GABA to locally modulate information being sent from nose to cortex (Borisovska *et al.*, 2013; Liu *et al.*, 2013, 2016; Banerjee *et al.*, 2015; Pignatelli & Belluzzi, 2017; Vaaga *et al.*, 2017). Production of these transmitters is undertaken by specialised enzymatic pathways, which include tyrosine hydroxylase (TH) as the rate-limiting step in dopamine synthesis, and DOPA decarboxylase (DDC) as its immediate downstream partner (Halász *et al.*, 1977). Glutamate decarboxylase (GAD)-67, on the other hand, is the GABA-synthesising enzyme that is ubiquitously expressed in this neuronal class (Kiyokage *et al.*, 2010; Kosaka *et al.*, 2019).

Expression levels of these neurotransmitter-synthesising enzymes are a well characterised target of activity-dependent plasticity in olfactory bulb dopaminergic neurons. TH, especially, is exquisitely sensitive to changes in sensory input. Since the landmark descriptions of decreased TH enzymatic activity and immunofluorescence in the bulb following damage to the peripheral olfactory system (Nadi *et al.*, 1981; Baker *et al.*, 1983), a wealth of evidence has shown that both *Th* mRNA and TH protein levels in bulbar cells track alterations in olfactory sensory neuron activity (Baker *et al.*, 1984, 1993, 1999; Kosaka *et al.*, 1987; Baker, 1990; Stone *et al.*, 1990; Cho *et al.*, 1996; Philpot *et al.*, 1998; Saino-Saito *et al.*, 2004; Bastien-Dionne *et al.*, 2010; Cave *et al.*, 2010; Weiss *et al.*, 2011; Grier *et al.*, 2016). Indeed, these TH changes are so reliable and sensitive that they are often used as confirmation that particular olfactory system manipulations have been effective (Wilson & Sullivan, 1995; Mandairon *et al.*, 2006; Kass *et al.*, 2013). Olfactory bulb GAD67 has received rather less attention, but nevertheless has also been shown to undergo experience-dependent changes in both mRNA (*Gad1*) and protein levels (Parrish-Aungst *et al.*, 2011; Lau & Murthy, 2012; Banerjee *et al.*, 2013; Wang *et al.*, 2017).

The vast majority of these activity-dependent enzyme expression changes have been detected following relatively prolonged manipulations of sensory input, usually on the order of weeks. However, there are indications that transmitter-synthesising enzyme expression can change more rapidly. A significant decrease in whole-bulb *Th* mRNA has been observed after only 2 d of unilateral naris occlusion (Cho *et al.*, 1996). *In vitro*, increases in TH immunofluorescence or TH-GFP transgene levels have been observed after just 24 h of sustained depolarisation in dissociated and slice cultures, respectively (Akiba *et al.*, 2007; Chand *et al.*, 2015). We also recently found that a 24 h period of *in vivo* sensory deprivation was associated with a decrease in relative immunofluorescence

in olfactory bulb TH-positive neurons, without any accompanying change in their density within the

glomerular layer (Galliano *et al.*, 2020).

A significant caveat is that, to date, these changes in neurotransmitter-synthesising enzyme

expression have all been detected either by employing tissue-level analyses that encompass the full

range of bulbar cell types, or by using the measured variable itself (e.g. TH immunofluorescence) to

identify individual cells to be studied. Ideally, gaining cell-type-specific information regarding

plasticity in enzyme expression would instead use an independent marker of cell identity. This

would enable experience-dependent changes to be identified amongst a consistently identified

group of neurons, without the potential distortion that can result when a plastic variable determines

which cells to measure. Here, we therefore took advantage of a conditional mouse transgenic line,

DAT^{IREScre} (Bäckman *et al.*, 2006), which when crossed with an appropriate reporter line generates

selective fluorescent label in cells that express the dopamine transporter (DAT-tdTomato). We

quantitatively characterised the selectivity of this line and found that it labels a small subset of

calretinin-expressing cells in addition to a majority of olfactory bulb dopaminergic neurons.

Crucially, this subtype specificity was unaffected by manipulations used to induce brief olfactory

deprivation. We found that such short-term, naturally relevant sensory manipulations do not affect

relative levels of GAD67 immunofluorescence in olfactory bulb DAT-tdTomato cells. In contrast,

deprivation does lead to transient and sustained decreases in the relative expression of DDC and TH,

respectively, within this independently identified population of neurons.

Methods

Animals

Mice of either gender were used and housed under a 12 h light:dark cycle in an environmentally

controlled room with access to water and food ad libitum. Day of birth was designated as P0.

DAT^{IREScre} mice (B6.SJL-*Slc6a3*$^{tm1.1(cre)Bkmn/J}$, Jax stock 006660; Bäckman *et al.*, 2006) were crossed with

Rosa26-floxed stop tdTomato reporter mice (Gt(ROSA)26Sor$^{tm9(CAG-tdTomato)Hze}$, Jax stock 007909) to

generate DAT-tdTomato mice. Wild type C57/Bl6 mice (Charles River) were used to back cross each

generation of transgenic mice. All experiments were performed at King's College London under the

auspices of UK Home Office personal and project licences held by the authors.

Unilateral naris occlusion

Olfactory sensory deprivation was induced by unilateral naris occlusion, using a custom made plug

(Cummings *et al.*, 1997). Each plug was constructed by knotting suture (Ethicon Ethilon

Monofilament Polyamide 6) around a piece of unscented dental floss pulled through the lumen of

PTFE tubing with an outer diameter of 0.6 mm and inner diameter of 0.3 mm (VWR International,

cat#: S1810 04). Mice were briefly anaesthetised with isoflurane and the plug trimmed, coated in

petroleum jelly and inserted into the right naris. Mice were returned to their home cage and left for

1 or 3 days before tissue collection. Presence of the nose plug was confirmed by examining the nasal

cavity after olfactory bulb dissection *ex vivo*. Sham controls, used for qPCR experiments only, had a

nose plug inserted into the right naris and immediately removed. Anaesthetised controls, used for all

other experiments, were handled and anaesthetised briefly with isoflurane but had no plug inserted

into either naris.

Transcardial perfusion and histology

P28 and P30 DAT-tdTomato mice were anaesthetised with an intraperitoneal overdose of

pentobarbital, then transcardially perfused with 20 mL phosphate buffered saline (PBS) with

20 U/mL heparin, followed by 20 mL of 4 % paraformaldehyde (PFA) in PIPES buffer (3 % sucrose,

60 mM PIPES, 25 mM HEPES, 5 mM EGTA, and 1 mM MgCl$_2$). PFA was prepared in batches, frozen at

-20 °C and defrosted shortly before use. On completion of perfusion, the mouse was decapitated,

and the brain removed from the skull and post-fixed in 4% PFA overnight. The brain was then

washed with PBS (3x5 minutes) and stored in PBS with 0.05% sodium azide for up to 6 months. For
sectioning, brains were embedded in 6% agarose and sliced coronally at 50 µm using a vibratome
(Leica VT1000s; Wetzlar, Germany). Slices were stored in PBS and 0.05% sodium azide at 4 °C before
staining.

Immunohistochemistry

Free-floating slices were incubated at room temperature (RT; 18-24 °C) for 2 hours in a blocking and
permeabilising solution (10% normal goat serum in PBS containing 0.25% Triton-X and 0.02% sodium
azide). They were then incubated in primary antibody diluted in the blocking solution at 4 °C for
between 18-24 hours (see Table 1 for antibodies used and their working dilution; endogenous
tdTomato fluorescence was always enhanced with immunolabel for red fluorescent protein). For
immunofluorescence intensity analysis, slices were incubated for approximately 72 hours in primary
antibody solution. Slices were then washed with PBS (3x5 minutes) and incubated in the species-
appropriate secondary antibody (Alexa Fluor® Life Technologies) diluted 1: 1,000 in the blocking
solution at RT for 2 hours. After 3x5 minutes washes in PBS, slices were mounted on glass slides with
MOWIOL® (Millipore).

Antigen	Host Species	Working Dilution	Supplier (product code)
Calbindin	Rabbit	1: 5,000	Swant (CB38a)
Calretinin	Mouse	1: 5,000	Millipore (MAB1568)
	Rabbit	1: 5,000	Swant (7699/3H)
cFos	Mouse	1: 1,000	Santa Cruz (sc166940)
Dopa decarboxylase (DDC)	Mouse	1: 1,000	Santa Cruz (sc-293287)
Glutamate decarboxylase 67 (Gad67)	Mouse	1: 500	Millipore (MAB351R)
Red fluorescent protein (tdTomato)	Rabbit	1: 1,000	Clontech (632496)
	Rat	1: 1,000	Chromotek (5F8)
Tyrosine hydroxylase (TH)	Mouse	1: 1,000	Millipore (MAB318)
	Rabbit	1: 1,000	Millipore (AB152)

Table 1: Details of primary antibodies

Image acquisition and experimental design

All images were acquired with a laser scanning confocal microscope (Zeiss LSM 710 Axio Examiner

Z1) using appropriate excitation and emission filters, a pinhole of 1 AU and a 40x oil immersion

objective. Laser power, gain and offset were manually set to prevent fluorescence saturation and

ensure optimum signal-to-noise.

For analyses of relative immunofluorescence intensity, we aimed to minimise and control for

between-section experimental variability. To reduce the effect of the homecage environment, each

experimental 'set' of four mice belonged to the same litter and was exposed to the same external

odour environment until unilateral naris occlusion or control procedure at P27. All mice were

returned to the homecage and left for 1 or 3 days. Perfusions and post-fixing were performed using

the same batch of PFA for each set, and mice within each set were perfused within 30 minutes of

each other. Brains from the same set were trimmed to the same size and co-embedded in 6%

agarose to ensure all brains would be sliced together. The agarose block was cut with landmarks to

ensure each mouse could be reliably identified. Co-embedded slices were processed for

immunohistochemistry as described above, with primary antibodies applied for 72 hours. Therefore,

all control and occluded sections on each slice were exposed to identical reagents at identical

concentrations for identical periods of time. The olfactory bulb sections in a co-embedded slice were

then imaged on the same day using identical laser and detection settings that were optimised for

that co-embedded slice. Z-stacks were taken from different parts of an olfactory bulb slice, setting

the start and end Z positions using the DAT-tdTomato fluorescence. Images were acquired at a

resolution of 512 x 512 pixels (0.277 µm /pixel) and a Z step of 1 µm.

Images for analysing colocalisation of DAT-tdTomato with other markers were acquired at a

resolution of 512 x 512 pixels (0.277 µm /pixel) and a Z step of 1 µm, with multiple Z stacks taken

around the entire olfactory bulb section.

All immunohistochemistry, imaging and image analysis of tissue from mice that had been

manipulated was performed blind to experimental group.

Image analysis

All image analysis was performed using ImageJ software (Fiji). Cells were counted manually using the

cell counter plugin and only if their soma was entirely inside the image area. Soma size and

fluorescence intensity were measured at the neuron's widest area in the Z-stack, identified when the

soma was larger than in the slice above and below. For measuring immunofluorescence intensity,

neurons were manually identified for analysis solely by the expression of DAT-tdTomato. TH and

GAD67 immunolabel were located primarily outside the nucleus, therefore, average

immunofluorescence intensity (mean grey value; MGV) was measured in the somatic cytoplasm

only, using a hand-drawn polygonal region of interest (ROI). For DDC and cFos, MGV was measured

across the entire soma, again using a hand-drawn polygonal ROI. All tdTomato cells that had the

entire cell body within the image area and for which the widest soma area could be confirmed were

measured in each stack.

For immunofluorescence intensity analysis, background MGV values were taken from unlabelled

bulbar regions of each Z-stack and were subtracted from the MGV of each cell in that Z-stack. MGV

values were then normalised to the control average (1 and 3 day) in each slice.

Electrophysiology

For acute slice preparation, mice were deeply anaesthetised with isoflurane and decapitated. The

olfactory bulb and frontal cortices were rapidly removed and transferred into ice-cold sucrose slicing

medium (containing in mM: 240 sucrose, 5 KCl, 1.25 Na_2HPO_4, 2 $MgSO_4$, 1 $CaCl_2$, 26 $NaHCO_3$ and 10

D-glucose, bubbled with 95% O_2 °and 5% CO_2). 300 µm horizontal sections of the olfactory bulb were

cut using a Leica vibratome (VT1000S) (Wetzlar, Germany) and were transferred to standard artificial

CSF (ACSF; containing in mM: 124 NaCl, 5 KCl, 1.25 Na_2HPO_4, 2 $MgSO_4$, 2 $CaCl_2$, 26 $NaHCO_3$ and 20 D-

glucose; with 10 µM NBQX, 50 µM APV, 10 µM SR-95531 to block major ionotropic synaptic

receptors, continuously bubbled with 95% O_2 and 5% CO_2, pH 7.4, ~310 mOsm). Slices were left to

recover at RT for at least 45 minutes before experiments began.

For whole-cell recordings, individual slices were secured in a chamber mounted on a Nikon FN1

microscope. DAT-tdTomato neurons were visualised with a 40x water immersion objective and

tdTomato fluorescence was revealed by under LED illumination (CoolLED pE-100) with appropriate

excitation and emission filters (ET575/50m, CAIRN Research, UK). Slices were continuously

superfused with ACSF heated to physiologically relevant temperature (34 ± 1 C) using an in-line

solution heater (TC-344B, Warner Instruments). Recordings were carried out using a Heka EPC10/2

amplifier coupled to PatchMaster acquisition software. Signals were digitised and sampled at 20 kHz.

Recording electrodes were pulled from borosilicate glass [outer diameter (OD), 1.5 mm; inner

diameter (ID), 1.17 mm] (GT100T-10, Harvard Apparatus) with a vertical puller (PC-10, Narishige) and

the tip fire-polished with a microforge (MF-830, Narishige). Pipettes with 3-5 MΩ resistance were

filled with K-gluconate-based intracellular solution, containing, in mM: 124 K-gluconate, 9 KCl, 10

KOH, 4 NaCl, 10 HEPES, 28.5 Sucrose, 4 Na$_2$ATP, 0.4 Na$_3$GTP, plus 0.15 % biocytin; pH ~7.3, 280-

290 mOsm. To study multiple action potential firing, 500 ms somatic current injections of Δ+5 pA

were applied from a holding potential of -60 mV in current-clamp mode with 100 % bridge balance,

until the point where the neuron entered depolarisation block.

After completion of electrophysiology experiments, the pipette was gradually removed with a small

amount of positive pressure applied to try and dislodge the patched neuron from the pipette. Once

the pipette was separated from the cell, the acute slice was transferred to 4% PFA. The slice was

fixed overnight at 4° C, washed with PBS (3x5 minutes) and stored in PBS and 0.05 % sodium azide at

4° C, for up to 3 months. Standard immunohistochemistry protocols were used (see above).

Streptavidin 488 (Alexa Fluor® Life Technologies) diluted 1: 1,000 was included with the secondary

antibodies to label biocytin in the patched neuron.

Quantitative PCR

All experimental steps for qPCR were performed in compliance with the MIQE guidelines (Bustin *et*

al., 2009). One or 3 days after naris occlusion or control sham surgery, wild-type mice were

sacrificed by cervical dislocation followed by decapitation. The brain was quickly removed and

placed in cold PBS. The right olfactory bulb was dissected and placed in RNAlater (Ambion) overnight

at 4° C and stored at -20° C until further processing. Total RNA extraction was performed using the

RNAqueous® Micro Kit (Life Technologies). RNA was eluted in 20 µl of elusion buffer and quantified

using Nanodrop (Thermo Fisher Scientific) and Qubit (RNA HS Assay Kit, Life Technologies). 300-

500 ng of total RNA were used for cDNA synthesis with SuperScript™ II Reverse Transcriptase (Life

Technologies), 0.5 mM dNTPs (Life Technologies), 5 µM OligodT23VN (IDT), 10 µM DTT (Life

Technologies) and 0.2 Units RNase inhibitor (SUPERase•In™, Invitrogen). RNA was denatured for

5 min at 65° C, before adding the reaction mix. Retrotranscription was performed at 42° C for

90 min, followed by 20 min at 70° C for enzyme inactivation. Standard RT-PCRs were run for all

samples, including non-template controls, to corroborate amplification of single bands of the

appropriate size. Cycling conditions were: 95° C for 3 min, (95 C, 15 s; 65-55° C (-0.5° C per cycle),

20 s; 68° C, 25 s), 20 cycles, (95° C, 15 s; 55° C, 20 s; 68° C, 25 s), 20 cycles, 68° C, 5 min, 4° C, hold.

Reactions were run on a SimpliAmp™ Thermal Cycler (Applied Biosystems). Primers used were:

ActB_For, cctctatgccaacacagtgc; ActB_Rev, cttctgcatcctgtcagcaa; Th_For, tgcctcctcacctatgcact;

Th_Rev, gtcagccaacatgggtacg. Primers were designed using the Universal Probe Library Assay Design

Centre (Roche), compatible with UPLs 157 and 3 for ActB and Th primer pairs, respectively.

qPCR reactions were run using the FastStart Essential DNA Probes Master mix (Roche) on the

LightCycler ® 96 Real-Time PCR System (Roche). Final primer concentration was 0.5 µM and probe

concentration was 0.1 µM. Cycling conditions were: 95° C for 10 min, (95° C, 10 s; 60° C, 30 s),

55 cycles, 4° C, hold. Technical duplicates were performed for all samples and reactions.

Amplification efficiency for all primer pairs was determined by obtaining a standard curve. Briefly,

the cDNAs from each cDNA synthesis reaction were pooled and serially diluted. qPCR reactions were

performed as above. Ct amplification curve values were plotted as a function of dilution factor and

the amplification efficiency determined from the slope of the linear regression as in $E = 10(-1/slope)-$

1. Amplification efficiency for ActB primers was 1.07 ± 0.02 and for Th primers, 1.01 ± 0.03. Fold

changes in Th expression levels were obtained using the $2^{(-ddCt)}$ method (Livak & Schmittgen, 2001;

Taylor *et al.*, 2019), first normalising Th Ct levels for each sample to ActB Ct levels and subsequently

normalising to the average of the control (sham) conditions.

Statistical analysis

Statistical analysis was carried out using Prism (GraphPad), Matlab (Mathworks) or SPSS (IBM). All

datasets were described as mean ± SEM, unless otherwise stated, n values reported in the text refer

to number of cells, and N values refer to number of mice. Details of all individual statistical tests are

reported alongside the appropriate results. α values were set to 0.05 (except where a Bonferroni

correction was carried out for multiple comparisons) and all comparisons were two tailed. For

multiple group comparisons with ANOVA followed by Bonferroni post-tests, all groups either passed

the D'Agostino and Pearson omnibus test for normality (GraphPad Prism), and/or could be fitted by

a Gaussian distribution that explained ≥ 90 % of their variance (custom Matlab script written by

MSG). Fisher's exact test was performed when analysing proportions. Due to the design of this test,

data had to be analysed twice, by combining groups based on manipulation and then duration of

manipulation. A Bonferroni correction was therefore carried out for multiple comparisons in this

instance (α = 0.05/2 = 0.025). qPCR normalised TH expression data were compared with separate

unpaired t-tests for 1 day and 3 day samples, which also required Bonferroni correction for multiple

comparisons (again, α = 0.05/2 = 0.025).

Results

Using DAT-tdTomato mice to label olfactory bulb dopaminergic neurons

We used a conditional genetic strategy to label dopaminergic neurons in the olfactory bulb by crossing DAT[IREScre] mice (Bäckman *et al.*, 2006) with a floxed tdTomato reporter line. This resulted in strong tdTomato fluorescence in small cells surrounding glomerular neuropil in the glomerular layer, along with the odd scattered tdTomato-positive cell in the superficial external plexiform layer (EPL), in keeping with previous reports of dopaminergic cell localisation within the olfactory bulb (Fig. 1A; (e.g. Hökfelt *et al.*, 1975; Banerjee *et al.*, 2013; Kosaka *et al.*, 2019). To confirm the identity of these DAT-tdTomato neurons, we co-stained fixed olfactory bulb slices from juvenile postnatal day (P)28 mice for the rate-limiting enzyme in the synthesis of dopamine, tyrosine hydroxylase (TH). Although necessary for both dopamine and noradrenaline production, the absence of local noradrenergic cells in the olfactory bulb (Hökfelt *et al.*, 1975; Rosser *et al.*, 1986) makes TH a selective marker for dopaminergic neurons in this region. In control animals we found a high incidence of co-labelled neurons, with ~75% of DAT-tdTomato cells staining positively for TH (Fig. 1B,C; % of tdTomato-positive neurons ± SEM: TH-positive, 74 ± 4 %; TH-negative, 26 ± 4 %; n = 598 cells, N = 3; Vaaga *et al.*, 2017). However, by no means all genetically labelled cells in the DAT-tdTomato mouse were TH-positive. We reasoned that the ~25 % of non-TH-expressing DAT-tdTomato cells might be 'true' dopaminergic neurons that have low, undetectable levels of TH expression due to long-term activity-dependent down-regulation (e.g. Baker *et al.*, 1983). However, co-staining for another dopamine-synthesising enzyme dopa decarboxylase (DDC), the expression of which is known to be resistant to long-term changes in ongoing activity (Stone *et al.*, 1990; Baker *et al.*, 1993; Cave *et al.*, 2010), showed that this is not the case. We found very good agreement between TH- and DDC-positive cells in the olfactory bulb glomerular layer (Fig. 1B; 99 ± 0.5 % of TH-positive neurons were also DDC-positive; 89 ± 2 % of DDC-positive neurons were also TH-positive; n = 598; N = 3), but we found only 3 % of DAT-tdTomato cells were positive for DDC without expressing TH (Fig. 1B,C; % of tdTomato-positive neurons ± SEM: TH- and DDC-positive, 74 ± 4 %; DDC-positive but TH-negative, 3 ± 7 %; TH-positive but DDC-negative, 0.5 ± 0.3 %; TH- and DDC-negative, 23 ± 2 %; n = 598; N = 3).

If ~20-25 % of DAT-tdTomato cells in the olfactory bulb are therefore non-dopaminergic, what type of cell are they? Red fluorescent neurons never had the large, pear-shaped soma characteristic of local excitatory neurons, the external tufted cells, in the glomerular layer (Hayar *et al.*, 2004). This made them most likely to be GABAergic juxtaglomerular interneurons, which in mouse are divided into three mutually exclusive subgroups expressing either TH or the calcium-binding proteins

calbindin and calretinin (Kohwi *et al.*, 2007; Kosaka & Kosaka, 2007; Panzanelli *et al.*, 2007; Parrish-

Aungst *et al.*, 2007). We never observed any DAT-tdTomato neurons that co-labelled for calbindin

(Fig. 1D), but we did find DAT-tdTomato cells that were positive for calretinin (Fig. 1E; Ninkovic *et al.*,

2010). Indeed, the tdT-positive/TH-negative subgroup was almost entirely accounted for by cells

expressing calretinin (Fig. 1E, F; % of tdTomato-positive neurons ± SEM: TH-positive only,

74.1 ± 2.24 %; calretinin-positive only, 21.71 ± 0.9 %; n = 1404; N = 3). Importantly, although we

observed some rare tdTomato-positive cells that were neither TH- nor calretinin-positive (Fig. 1F;

tdTomato-positive only, 4.2 ± 1.4%, consistent with the 3 % of tdTomato cells found above to be

DDC-positive but TH-negative), we never saw any tdTomato-positive cells co-labelled for both TH

and calretinin (n = 0 / 1404 cells from N = 3 mice). This suggests that the calretinin-positive DAT-

tdTomato cells are non-dopaminergic calretinin neurons that have at least transiently activated DAT

expression at some point in their history, rather than being dopaminergic neurons that also co-

express calretinin.

To check these cells' distinct properties, we obtained whole-cell patch-clamp recordings of DAT-

tdTomato neurons in acute olfactory bulb slices. Although these cells are structurally fragile and

extremely difficult to recover for post-recording histology (Galliano *et al.*, 2018), we were able to

find very rare example neurons that had been successfully filled with biocytin during our recordings

and then co-labelled for glomerular layer markers. While a confirmed TH-positive DAT-tdTomato

neuron had characteristic multiple spike production in response to a prolonged 500 ms current

injection in current-clamp, the one calretinin-positive tdTomato cell we managed to recover

displayed only a single, small spike followed by a sustained depolarised plateau potential, as

previously described to be typical of this bulbar cell type (Fig. 1G; Fogli Iseppe *et al.*, 2016; Benito *et*

al., 2018). The calretinin-positive cells labelled in DAT-tdTomato mice are therefore likely to share

their major features with the general population of calretinin-expressing juxtaglomerular neurons.

Figure 1

Figure 1. Cell-type identity of olfactory bulb DAT-tdTomato neurons. **A)** tdTomato label in DAT-tdTomato olfactory bulb; **i)** aerial view diagram of mouse brain, showing location of olfactory bulb coronal section in **ii**; **ii)** coronal section of olfactory bulb from a DAT-tdTomato mouse; box shows location of detail in **iii**; **iii)** tdTomato-positive neurons are located predominantly in the olfactory bulb's glomerular layer (GL); ONL, olfactory nerve layer; EPL, external plexiform layer. **B)** Example maximum intensity projection images from the glomerular layer in the olfactory bulb of a control DAT-tdTomato mouse, showing co-label for TH and DDC. Yellow star indicates location of a neuron showing expression of all three markers. Arrowheads indicate location of neurons expressing two markers: red = DAT-tdTomato and DDC; white = TH and DDC. Arrows indicate location of neurons expressing a single marker: red = DAT-tdTomato; white = DDC. **C)** Mean ± SEM percentage of tdTomato+ neurons positive for TH and/or DDC. Grey circles indicate percentages from individual mice. **D)** Example maximum intensity projection images showing immunofluorescence for calbindin in the olfactory bulb glomerular layer of a control DAT-tdTomato mouse. **E)** Example maximum intensity projection images showing immunofluorescence for TH and calretinin in the glomerular layer of the olfactory bulb in a control DAT-tdTomato mouse. Red arrowheads indicate example neurons expressing DAT-tdTomato and calretinin, but not TH. **F)** Mean ± SEM percentage of tdTomato+ neurons expressing TH and calretinin. Grey circles indicate percentages from individual mice. **G)** Action potential firing patterns in response to 500 ms somatic current injection (left) and immunohistochemical characterisation (right) in example biocytin-filled DAT-tdTomato neurons that were co-labelled for either TH (top) or calretinin (bottom).

Sensory deprivation via brief unilateral naris occlusion reduces cFos expression in olfactory bulb DAT-tdTomato neurons

Having shown that the DAT-tdTomato mouse line labels olfactory bulb neurons that are predominantly dopaminergic but include a distinct subset of calretinin-positive cells, we went on to use this line for naturally-relevant manipulations of sensory experience (Fokkens *et al.*, 2012; Galliano *et al.*, 2020). In juvenile mice from postnatal day (P)27, we induced brief 1- or 3-day periods of sensory deprivation without damaging the olfactory epithelium, via unilateral naris occlusion with a plastic plug (Fig.2A; Cummings *et al.*, 1997; Kikuta *et al.*, 2015; Cheetham *et al.*, 2016; Galliano *et al.*, 2020). Because of interpretational issues caused by abnormal airflow through the contralateral naris in such manipulations (Coppola, 2012; Kass *et al.*, 2013), we did not compare deprived versus spared olfactory bulbs in the same animals, and instead made all comparisons between the right bulbs of individual occluded or control mice (see Methods).

To validate the effectiveness of our sensory deprivation approach for reducing activity in olfactory bulb DAT-tdTomato neurons *in vivo*, we examined immunofluorescence for the activity-dependent immediate early gene cFos (Liu *et al.*, 1999). Co-embedding olfactory bulbs from all four treatment groups (Fig. 2A; 1 day control, 1 day occluded; 3 day control; 3 day occluded) prior to sectioning allowed the best possible consistency in immunostaining protocols (see Methods; Vlug *et al.*, 2005; Galliano *et al.*, 2020). It also enabled us to normalise labelling intensity within co-embedded sets in order to combine analyses across multiple individuals. With this approach we found that DAT-tdTomato cells from occluded mice had significantly lower cFos immunofluorescence after both 1 and 3 days of occlusion (Fig. 2B-D; normalised intensity (%) mean ± SEM: 1 day control, 102.5 ± 3.83%, n = 316 cells, N = 2 mice; 1 day occluded, 56.85 ± 2.1%, n = 358, N = 2; 3 day control, 97.53 ± 3.3%, n = 317, N = 2; 3 day occluded, 62.87 ± 2.63%, n = 309, N = 2. Multi-level ANOVA controlling for set variance, effect of manipulation, $F_{1,1298} = 60.73$, $p < 0.0001$; effect of time, $F_{1,1298} = 1.4$, $p = 0.24$; effect of manipulation x time interaction, $F_{1,1299} = 3.4$, $p = 0.07$. Bonferroni post- test comparing control versus occluded ($\alpha = 0.025$): 1 day, $F_{1,673} = 115.76$, $p < 0.00005$; 3 day, $F_{1,626} = 68.56$, $p < 0.00005$). Brief sensory deprivation therefore reliably lowers activity levels in olfactory bulb DAT-tdTomato neurons, and does so in a sustained manner over 1 to 3 days.

Figure 2

Figure 2. Sensory deprivation via brief unilateral naris occlusion reduces cFos expression in olfactory
bulb DAT-tdTomato neurons. **A)** Schematic of naris occlusion protocol and tissue co-embedding for
comparison of relative immunofluorescence. OE, Olfactory epithelium; OB, Olfactory bulb; P,
Postnatal day; a, 1 day control; b, 1 day occluded; c, 3 day control; d, 3 day occluded. **B)** Example
maximum intensity projection images showing immunofluorescent label for cFos in the glomerular
layer of the olfactory bulb in 1 and 3 day control and occluded DAT-tdTomato mice. All images are
from the same co-embedded preparation, acquired and presented with identical settings. **C)**
Cumulative distribution of normalised cFos intensity (%) in DAT-tdTomato neurons in each treatment
group. **D)** Line plots showing overall mean ± SEM normalised cFos intensity (%) in each treatment
group; unfilled circles show mean normalised cFos intensity (%) for individual mice. Multi-level
ANOVA controlling for set variance; effect of manipulation, **** $p < 0.0001$; effect of time, ns: non-
significant. Bonferroni post-test ($\alpha = 0.025$); 1 day, effect of manipulation, **** $p < 0.00005$; 3 day,
effect of manipulation, **** $p < 0.00005$.

Brief naris occlusion does not alter the sub-type composition of olfactory bulb DAT-tdTomato neurons

Given the mixed dopaminergic and calretinin-expressing nature of the population labelled in the DAT-tdTomato mouse olfactory bulb (Fig. 1), it was important to check whether our effective short-term sensory manipulations (Fig. 2) altered this sub-type composition in any way. Crucially, we found no change in the proportion of DAT-tdTomato cells that were calretinin-positive after 1 or 3 days of sensory deprivation (Fig. 3A,B; % of calretinin-positive DAT-tdTomato neurons; 1 day control, 20.95 %, n = 506, N = 3; 1 day occluded, 24.5 %, n = 502, N = 3; 3 day control, 23.91 %, n = 343, N = 2; 3 day occluded, 25 %, n = 424, N = 3. Fisher's exact test with α = 0.025, effect of manipulation, p = 0.22; effect of time, p = 0.40).

Olfactory bulb dopaminergic neurons are not a unitary, homogeneous cell type (Galliano *et al.*, 2018; Kosaka *et al.*, 2019), so it was also important to assess the representation of dopaminergic subtypes amongst DAT-tdTomato neurons following sensory deprivation. Work from our laboratory and others has identified a key distinction between, on the one hand, small-soma olfactory bulb dopaminergic cells that lack an axon and can be generated throughout life and, on the other, large-soma olfactory bulb dopaminergic cells that possess an axon and are exclusively generated in early embryonic development (Galliano *et al.*, 2018; Kosaka *et al.*, 2019). The latter large-soma sub-population, although present, is significantly under-represented by DAT-tdTomato label, which is dominated by small-soma anaxonic cells (Galliano *et al.*, 2018). Using a previously established soma area cut-off value of 70 μm^2 to define olfactory bulb dopaminergic cells as 'small' (Galliano *et al.*, 2018), we confirmed here that this subtype accounts for > 90 % of DAT-tdTomato neurons (Fig.3C,D). Crucially, the high proportion of such small cells represented in the DAT-tdTomato line – which will also include the small-soma calretinin-positive subgroup (Parrish-Aungst *et al.*, 2007) – was not significantly affected by 1 or 3 day naris occlusion (Fig.3C,D; % of 'small' cells; 1 day control, 95.22 %, n = 460, N = 3; 1 day occluded, 91.18 %, n = 408, N = 3; 3 day control, 93.62 %, n = 298, N = 2; 3 day occluded, 96.75 %, n = 461, N = 3. Fisher's exact test with α = 0.025, effect of manipulation, p = 0.75; effect of time, p = 0.067). Brief periods of sensory deprivation therefore do not significantly alter the cellular sub-composition of olfactory bulb glomerular layer neurons labelled in DAT-tdTomato mice.

     Figure 3

Figure 3. Brief naris occlusion does not alter the sub-type composition of olfactory bulb DAT-
tdTomato neurons. **A)** Example maximum intensity projection images showing immunofluorescence
for tdTomato and calretinin in the glomerular layer of the olfactory bulb in 1 and 3 day control and
occluded DAT-tdTomato mice. Red arrowheads indicate example neurons co-expressing
DAT-tdTomato and calretinin. **B)** Line plots showing, for each treatment group, overall % of
tdTomato+ neurons that also express calretinin. Open circles show % values for individual mice.
Fisher's exact test with Bonferroni correction for multiple comparisons (α = 0.025); ns, non-
significant. **C)** Histogram of soma sizes for all tdTomato+ neurons in each treatment condition.
Dotted line shows 70 μm^2 threshold for cells classed as 'small'. **D)** Small cell % data; all conventions
as in **B**.

Brief sensory deprivation does not affect GAD67 expression in olfactory bulb DAT-tdTomato neurons

Having confirmed that brief unilateral naris occlusion effectively lowers activity levels in the

olfactory bulb DAT-tdTomato population without affecting its subtype composition, we used this

mouse line to investigate potential experience-dependent plasticity in the expression of

neurotransmitter-synthesising enzymes. We focused first on the GABA-synthesising enzyme isoform

GAD67, which is ubiquitously expressed in dual GABA- and dopamine-releasing olfactory bulb

neurons (Kosaka & Kosaka, 2007; Kiyokage *et al.*, 2010; Kosaka *et al.*, 2019), and has been previously

reported to undergo experience-dependent changes in both mRNA expression and protein levels

after prolonged manipulations (Parrish-Aungst *et al.*, 2011; Lau & Murthy, 2012; Banerjee *et al.*,

2013; Wang *et al.*, 2017). However, here we found no changes in relative GAD67

immunofluorescence intensity in olfactory bulb DAT-tdTomato neurons after 1 or 3 days of unilateral

naris occlusion (Fig.4; normalised immunofluorescence intensity (%) mean ± SEM: 1 day control,

97.43 ± 3.1 %, n = 261, N = 3; 1 day occluded, 92.93 ± 2.9 %, n = 314, N = 3; 3 day control, 103.5 ±

3.7 %, n = 192, N = 2; 3 day occluded, 94.03 ± 2.88 %, n = 304, N = 3. Multi-level ANOVA controlling

for set variance, effect of manipulation, $F_{1,1071} = 0.09$, p = 0.77; effect of time, $F_{1,1071} = 1.6$, p = 0.21;

effect of manipulation x time interaction, $F_{1,1071} = 0.62$, p = 0.43).

Figure 4

Figure 4. Brief sensory deprivation does not affect GAD67 expression in olfactory bulb DAT-
tdTomato neurons. **A)** Example maximum intensity projection images showing immunofluorescence
for GAD67 in the glomerular layer of the olfactory bulb in 1 and 3 day control and occluded DAT-
tdTomato mice. All images are from the same co-embedded preparation, acquired and presented
with identical settings. **B)** Cumulative distribution of normalised GAD67 intensity (%) in DAT-
tdTomato neurons in each treatment group. **C)** Line plots showing overall mean ± SEM of
normalised GAD67 intensity (%) in each treatment group; unfilled circles show mean normalised
GAD67 intensity (%) for individual mice. Multi-level ANOVA controlling for set variance; effect of
manipulation and effect of time, ns: non-significant.

DDC expression in olfactory bulb DAT-tdTomato neurons is only slightly and transiently reduced by

unilateral naris occlusion

The enzyme DDC catalyses the direct synthesis of dopamine from L-DOPA, but unlike TH has

received relatively little attention regarding its potential for activity-dependent expression. Indeed,

two previous studies found no effect of odour deprivation on *Ddc* mRNA levels in the olfactory bulb

following prolonged, >1 month-long naris closure (Stone *et al.*, 1990; Cave *et al.*, 2010). Here, we

found that brief sensory deprivation produced a small but significant decrease in the relative

immunofluorescence intensity of DDC in olfactory bulb DAT-tdTomato neurons (Fig. 5; normalised

immunofluorescence intensity (%) mean ± SEM: 1 day control, 97.99 ± 2.61%, n = 417, N = 2; 1 day

occluded, 82.88 ± 2.47%, n = 425, N = 2; 3 day control, 104.2 ± 3.45%, n = 198, N = 1; 3 day occluded,

96.03 ± 2.61%, n = 412, N = 2. Multi-level ANOVA controlling for set variance, effect of manipulation,

$F_{1,1452}$ = 10.19, p = 0.001; effect of time, $F_{1,1452}$ = 1.95, p = 0.16; effect of manipulation x time

interaction, $F_{1,1452}$ = 1.44, p = 0.22). Post-hoc tests demonstrated that this effect was driven largely

by a reduction in DDC immunofluorescence only after 1 day of occlusion, with expression of DDC in 3

day occluded DAT-tdTomato neurons returning back close to control levels (Fig. 5C; Bonferroni post-

test comparing control versus occluded with α = 0.025: 1 day, $F_{1,842}$ = 17.70, p < 0.0001; 3 day, $F_{1,610}$ =

3.39, p = 0.066). Small and transient changes in relative DDC levels are therefore possible in

olfactory bulb dopaminergic cells experiencing sustained decreases in activity.

Figure 5

Figure 5. DDC expression in olfactory bulb DAT-tdTomato neurons is slightly reduced only after 1 day of unilateral naris occlusion. **A)** Example maximum intensity projection images showing immunofluorescence for DDC in the glomerular layer of the olfactory bulb in 1 and 3 day control and occluded DAT-tdTomato mice. All images are from the same co-embedded preparation, acquired and presented with identical settings. **B)** Cumulative distribution of normalised DDC intensity (%) in DAT-tdTomato neurons in each treatment group. **C)** Line plots showing overall mean ± SEM of normalised DDC intensity (%) in each treatment group; unfilled circles show mean normalised DDC intensity (%) for individual mice. Multi-level ANOVA controlling for set variance; effect of manipulation, ** $p < 0.01$; effect of time, ns: non-significant. Bonferroni post-test ($\alpha = 0.025$); 1 day, effect of manipulation, ** $p < 0.005$; 3 day, effect of manipulation, ns: non-significant.

Brief sensory deprivation reduces TH expression in olfactory bulb DAT-tdTomato neurons

Olfactory bulb dopaminergic neurons have a well-described ability to regulate TH expression in an activity-dependent manner, including after unilateral naris occlusion by plug insertion (Nadi *et al.*, 1981; Baker *et al.*, 1983; Kosaka *et al.*, 1987; Cummings *et al.*, 1997). The most rapid previously-published example of this found that 2 days of odour deprivation induced a small, but significant, decrease in *Th* mRNA in the occluded olfactory bulb compared to open-side control (Cho *et al.*, 1996). In cultured bulbar neurons or slices, however, just one day of elevated activity was sufficient to increase TH immunofluorescence intensity or TH-GFP transgene expression (Akiba *et al.*, 2007; Chand *et al.*, 2015). We also recently found that 24 h naris occlusion was accompanied by a significant drop in immunofluorescence for TH in the ipsilateral olfactory bulb, without producing a change in TH-positive cell density (Galliano *et al.*, 2020). However, the above approaches had the drawback that they used TH expression itself as a means to identify samples for TH quantification. We therefore used our conditional genetic labelling strategy to assess rapid deprivation-induced changes in TH expression in an independently identified neuronal population.

Brief sensory deprivation was indeed associated with a decrease in the relative intensity of TH immunofluorescence in olfactory bulb DAT tdTomato neurons (Fig.6A-C; normalised immunofluorescence intensity (%) mean ± SEM: 1 day control, 95.27 ± 3.35%, n = 460, N = 3; 1 day occluded, 70.96 ± 3.16%, n = 408, N = 3; 3 day control, 107.3 ± 4.64%, n = 298, N = 2; 3 day occluded, 57 ± 2.32%, n = 461, N = 3. Multi-level ANOVA controlling for set variance, effect of manipulation, $F_{1,1546.} = 2.9$, $p = 0.09$; effect of time, $F_{1,1626} = 3.1$, $p = 0.08$). This effect was highly significant after just 1 day (Fig. 6C; Bonferroni post-test comparing control versus occluded with $\alpha = 0.025$: 1 day, $F_{1,867} = 27.41$, $p < 0.00005$), but was even stronger after 3 days (Fig. 6C; post-test: 3 day, $F_{1,639} = 76.36$, $p < 0.00005$; effect of manipulation x time interaction in main ANOVA, $F_{1,1613} = 11.67$, $p = 0.001$). Continuous naris occlusion therefore produces a rapid downregulation in TH immunofluorescence in independently identified olfactory bulb dopaminergic neurons, and this effect becomes stronger over time.

Finally, we also observed rapid occlusion-induced downregulation of *Th* expression at the mRNA level, as assessed via qPCR on whole-OB samples from wild-type mice. Compared to sham-treated bulbs, OBs on the occluded side had significantly lower levels of Th expression after both 1 day and 3 days of sensory deprivation (Fig.6D; 1 day mean ± SEM fold change: sham, 1.02 ± 0.07; occluded, 0.66 ± 0.04; t-test with $\alpha = 0.025$, $t_{18} = 4.69$, $p = 0.0002$, N = 20; 3 day: sham, 1.04 ± 0.13; occluded, 0.53 ± 0.03; Welch's t-test with $\alpha = 0.025$, $t_{7.73} = 4.00$, $p = 0.0043$; N = 16).

Figure 6

Figure 6. Brief sensory deprivation reduces TH expression in olfactory bulb DAT-tdTomato neurons.
A) Example maximum intensity projection images showing immunofluorescence for TH in the
glomerular layer of the olfactory bulb in 1 and 3 day control and occluded DAT-tdTomato mice. All
images are from the same co-embedded preparation, acquired and presented with identical
settings. **B)** Cumulative distribution of normalised TH intensity (%) in DAT-tdTomato neurons in each
treatment group. **C)** Line plots showing overall mean ± SEM of normalised TH intensity (%) in each
treatment group; unfilled circles show mean normalised TH intensity (%) for individual mice. Multi-
level ANOVA controlling for set variance; effect of manipulation x time interaction, **** p < 0.0001;
effect of manipulation and effect of time, ns: non-significant. Bonferroni post-test (α = 0.025); 1 day,
effect of manipulation, **** p < 0.00005; 3 day, effect of manipulation, **** p < 0.00005. **D)** *Th*
mRNA expression by whole-bulb qPCR. Circles show fold change for individual bulbs vs relevant
sham control mean; lines show mean ± SEM fold change for each group. t-test (α = 0.025); 1 day,
***, p = 0.0002; 3 day, **, p = 0.0043.

Discussion

Here we used a conditional genetic strategy to identify olfactory bulb neurons expressing the dopamine transporter. DAT-tdTomato neurons were predominantly dopaminergic, but also included a subset of non-dopaminergic, calretinin-positive cells. Importantly, while the activity of DAT-tdTomato cells could be readily manipulated with brief periods of sensory deprivation, these manipulations did not alter the labelled neurons' subtype composition. We found that 1-3 days of unilateral naris occlusion was insufficient to alter relative levels of immunolabel for the GABA-synthesising enzyme GAD67 in these cells. In contrast, similarly brief sensory manipulations were associated with significant activity-dependent changes in the levels of enzymes required for the synthesis of dopamine.

Identifying olfactory bulb dopaminergic neurons using DAT^{IREScre} mice

Using transgenic approaches to label specific cell types in the brain requires thorough characterisation of their specificity. To study dopaminergic neurons in the olfactory bulb, several previous studies have utilised conditional or knock-in strategies based on expression driven from the *Th* promoter (e.g. Saino-Saito *et al.*, 2004; Pignatelli *et al.*, 2005; Kiyokage *et al.*, 2010; Adam & Mizrahi, 2011; Liu *et al.*, 2013; Wachowiak *et al.*, 2013). However, *Th* mRNA expression is more widespread than TH protein expression in the bulb, with the former being observed amongst presumptive immature migrating neurons, as well as amongst TH-immunopositive neurons concentrated in the glomerular layer (Saino-Saito *et al.*, 2004; Pignatelli *et al.*, 2009; Banerjee *et al.*, 2013). In other brain regions, the specificity of TH-Cre mice for labelling dopaminergic neurons also appears to be rather low (Lammel *et al.*, 2015). We therefore opted for a conditional genetic labelling strategy based on Cre recombinase expression under the control of the DAT promoter, which has been reported to be more specific to TH-positive neurons in general (Lammel *et al.*, 2015), and which has already been used successfully to target olfactory bulb dopaminergic neurons. In those bulb studies, which either employed the DAT^{IREScre} line used here (Vaaga *et al.*, 2017; Short & Wachowiak, 2019; Sato *et al.*, 2020) a DAT-Cre knock-in model (Ninkovic *et al.*, 2010; Banerjee *et al.*, 2015), or a BAC transgenic DAT-CreER^{T2} inducible line (Bywalez *et al.*, 2016), approximately 85% of genetically-labelled neurons were found to be TH-positive by immunohistochemistry (Banerjee *et al.*, 2015; Vaaga *et al.*, 2017). This is broadly in line with the ~75% co-label we found here, with cross-study variability possibly arising from differences in the Cre-dependent reporter lines utilised

and/or differences in label detection criteria. Nevertheless, the general approach of using DAT-Cre

mice to label dopaminergic neurons in the olfactory bulb clearly produces reasonable specificity.

However, we found that not all DAT-tdTomato olfactory bulb cells were TH-positive, and that the

large majority of non-dopaminergic genetically labelled neurons (~20 %) expressed the calcium-

binding protein calretinin. Calretinin expression in DAT-Cre-positive bulbar neurons has been

previously reported, but not quantified, in a separate knock-in line (Ninkovic *et al.*, 2010). This

suggests that off-target label of calretinin cells is not a peculiarity of one particular genetic

manipulation, but instead reflects a biological link between these neurons and their dopaminergic

neighbours. Indeed, fate mapping neuronal precursors that generate glomerular layer GABAergic

interneurons has demonstrated that the decision to differentiate into a dopaminergic neuron over a

calretinin neuron relies on just a few transcription factors. Loss of Pax6, Dlx2 or Meis2 reduces the

proportion of postnatal dopaminergic neurons that are generated, while increasing the number of

calretinin neurons (Brill *et al.*, 2008; Agoston *et al.*, 2014). Overexpression of Dlx2 increases the

proportion of dopaminergic neurons and decreases the calretinin population (Brill *et al.*, 2008). The

transcription factor Er81 is also expressed in both calretinin and dopaminergic neurons in the

glomerular layer and is important in mice for regulating the expression of *Th* (Cave *et al.*, 2010).

Therefore, the molecular determinants of dopaminergic versus calretinin identities appear to be

closely interdependent. It remains unclear whether tdTomato expression in some calretinin neurons

is due to transient expression of DAT during their development, or whether this subset of cells

continues to express DAT postnatally. In the former case, obtaining transgenic label via inducible

activation of DAT-Cre in mature mice (Bywalez *et al.*, 2016) might result in greater specificity for

bulbar dopaminergic neurons. Low levels of bulbar DAT expression (Cockerham *et al.*, 2016)

unfortunately precluded us from addressing this question using immunohistochemistry, but targeted

interrogation of single-cell sequencing datasets (e.g. Zeisel *et al.*, 2018) might provide an answer in

the near future. Regardless, it is clear that this minority calretinin-expressing population needs to be

considered when interpreting the results of experiments performed on 'dopaminergic' neurons

targeted via DAT-Cre mouse lines (Wachowiak *et al.*, 2013; Banerjee *et al.*, 2015; Bywalez *et al.*,

2016; Vaaga *et al.*, 2017; Galliano *et al.*, 2018, 2020; Sato *et al.*, 2020). Thankfully, though, the

distinct and unusually immature-like properties of glomerular layer calretinin-positive cells, including

their extremely low excitability and apparent lack of GABA release (Fogli Iseppe *et al.*, 2016; Benito

et al., 2018; Tavakoli *et al.*, 2018), mean that their presence as a minority of DAT-Cre-targeted cells is

unlikely to have significantly biased the conclusions of those previous studies.

Our main goal was to achieve reasonable specificity in genetically labelling olfactory bulb

dopaminergic neurons, but we note that our approach was by no means comprehensive in labelling

all olfactory bulb dopaminergic neurons. As is the case for many Cre lines (Heffner *et al.*, 2012), we

saw mosaicism in the fact that many immunohistochemically-defined TH-positive neurons did not

express the tdTomato marker (see Fig. 1B,E; % of TH-positive neurons that were tdTomato-negative,

36 ± 0.2 %, n = 692 cells, N = 3 mice; Galliano *et al.*, 2018). The DAT^{IREScre} line is also known to

particularly under-represent one subtype of bulbar dopaminergic neuron – the large-soma, axon-

bearing, embryonically-generated subclass (Galliano *et al.*, 2018) – and we found again here that

small-soma, presumed anaxonic neurons dominate the tdT-labelled population. However, large-

soma, axon-positive cells are already rare amongst the olfactory bulb dopaminergic population, and

they can certainly still be labelled in DAT-tdTomato mice (Galliano *et al.*, 2018). So, although the

DAT^{IREScre} line produces genetic label which is heavily biased towards one particular dopaminergic

subpopulation in the olfactory bulb, it is far from being entirely subtype specific. Overall, the

strategy of using DAT-based transgenic lines to target dopaminergic neurons in the olfactory bulb

can produce good levels of specificity, but care needs to be taken to fully characterise the resulting

label before such experiments can be safely interpreted.

Experience-dependent plasticity of GAD67

With brief periods of sensory deprivation, we saw no evidence here for changes in relative GAD67

immunofluorescence amongst olfactory bulb DAT-tdTomato-expressing neurons. Given the mixed

calretinin- and TH-positive composition of our tdTomato-labelled population (Fig. 1), this result could

theoretically have been produced if changes in GAD67 levels in dopaminergic neurons were

perfectly offset by inverse, and stronger, changes in the smaller calretinin-positive tdTomato

subgroup. We did not co-label for calretinin and GAD67 in this study, while the literature contains

conflicting reports as to whether calretinin-positive neurons in the olfactory bulb express GAD67 at

all (Panzanelli *et al.*, 2007; Parrish-Aungst *et al.*, 2007; opposing results are likely due to the different

transgenic versus immunohistochemical labelling methods employed in these studies). Even if

calretinin-positive tdTomato cells do express GAD67, however, the most parsimonious

interpretation of our data remains that dopaminergic bulbar neurons do not rapidly alter their

GAD67 levels in response to sensory deprivation. This fits with previous work measuring general

GAD (i.e. GAD65+67) expression in the bulb, which found no significant alterations following more

long-term sensory deprivation (Kosaka *et al.*, 1987; Stone *et al.*, 1991; Baker *et al.*, 1993). In contrast,

reduced expression of GAD67, but not GAD65, has been observed after 7 days of unilateral naris

occlusion and deafferentation via measures of glomerular layer immunofluorescence and whole-

bulb protein expression (Parrish-Aungst *et al.*, 2011). Decreased expression of *Gad1* mRNA and

GAD67 protein have also been described following chemical ablation of olfactory sensory neurons

(Lau & Murthy, 2012) or unilateral naris occlusion (Banerjee *et al.*, 2013; Wang *et al.*, 2017).

However, these studies used longer-duration manipulations and did not distinguish between

different bulbar neuronal populations. Therefore, either 1-3 days of unilateral naris occlusion is

insufficient to induce activity dependent changes in GAD67, or measures based on

immunofluorescence intensity may not be sensitive enough to always detect changes in GAD67, or

these changes are driven in neurons not labelled in the DAT-tdTomato mouse line.

Experience-dependent plasticity of dopamine-synthesising enzymes

We report here the first description of activity-dependent changes in olfactory bulb DDC levels. This

effect was small and transient, with relative DDC immunofluorescence in DAT-tdTomato neurons

returning back to baseline after just 3 days of sensory deprivation. Such temporary downregulation

might explain why previous studies – which used much longer periods of sensory manipulation - did

not find similar effects (Stone *et al.*, 1990; Cave *et al.*, 2010), although expression of DDC in the

midbrain has been reported to be affected within a few hours of activation or blockade of dopamine

receptors (Hadjiconstantinou *et al.*, 1993; Zhu *et al.*, 1993). Further work is needed to confirm

whether downregulation of bulbar DDC after 1 day of occlusion is a consistent activity-dependent

modification. If so, it might reflect an initial strategy whereby dopaminergic neurons downregulate

multiple nodes in the dopamine synthesis pathway, before sufficiently decreased TH activity can

alone – as the rate-limiting step – control levels of transmitter production.

Our data showing decreased TH expression after olfactory sensory deprivation are entirely

consistent with a large body of prior work describing this effect (Nadi *et al.*, 1981; Baker *et al.*, 1983,

1984, 1993, 1999; Kosaka *et al.*, 1987; Baker, 1990; Stone *et al.*, 1990; Cho *et al.*, 1996; Philpot *et al.*,

1998; Saino-Saito *et al.*, 2004; Bastien-Dionne *et al.*, 2010; Cave *et al.*, 2010; Weiss *et al.*, 2011; Grier

et al., 2016), including some indications that it can occur after relatively brief manipulations (Cho *et

al.*, 1996; Akiba *et al.*, 2007; Chand *et al.*, 2015; Galliano *et al.*, 2020). It is novel, however, that our

methodological approach demonstrated activity-dependent TH alterations in cells identified in an

entirely TH-independent manner. As discussed above, this approach was not completely selective,

labelling a minority of calretinin-positive neurons in addition to dopaminergic cells. However, since

we never observed a DAT-tdTomato neuron that was both calretinin-positive and dopaminergic, and

since the proportions of calretinin-positive cells did not change significantly with occlusion (Fig. 3),

this off-target labelled sub-population cannot account for the relative changes in TH and DDC

immunofluorescence we observed. Indeed, if anything, the presence of a significant proportion of

essentially TH- or DDC-negative cells amongst the DAT-tdTomato samples we analysed may have
made our ability to detect relative differences in enzyme levels less sensitive.

We also note the caveat that, of course, changes in immunofluorescence intensity cannot be used to
infer activity-dependent plasticity in *absolute* levels of protein expression. With this approach,
results can be easily confounded by variability in the duration of immunohistochemistry steps, the
composition of reagents, and/or individual image acquisition settings. We did take every possible
measure to mitigate against such factors, comparing tissue from mice that were fixed on the same
day with the same fixative solution, comparing sections that were physically co-embedded prior to
all stages of histological processing, and comparing images that were taken on the same day with
the same acquisition settings. We also ensured that we only ever made *relative* comparisons of
immunofluorescence. So, while we are confident that, for example, TH levels decreased in DAT-
tdTomato cells after 1 day of sensory deprivation, we do not claim any insight into the magnitude of
this decrease. We do, however, have corroborative, fully quantitative qPCR data to show that – at
least at the whole-bulb level – brief sensory deprivation is associated with an approximate halving of
TH mRNA levels.

Mechanisms

It remains unclear precisely how activity-dependent changes in TH expression occur, though the
immediate depolarisation-activated pathways appear to require signalling through L-type CaV1
Channels and cAMP (Cigola *et al.*, 1998), and *Th* gene expression is at least partly controlled by
changes in secondary DNA structure influencing the binding of heterogeneous nuclear
ribonucleoproteins (hnRNPs) to its proximal promoter (Banerjee *et al.*, 2014; Wang *et al.*, 2017). Our
findings suggest that these processes can work over reasonably quick timescales. However, given
the similarity in promoter-based changes in secondary structure and hnRNP binding between the *Th*
and *Gad1* genes (Wang et al., 2017), it will be interesting to consider how these processes can work
more rapidly, more specifically, and/or be coupled with targeted changes in mRNA and protein
turnover, to produce selective downregulation of TH without any changes in GAD67 after a few days
of naris occlusion.

Implications for sensory processing

Plasticity of TH expression is believed to be a way in which the olfactory bulb can tune its levels of

dopaminergic signalling according to changes in incoming sensory-evoked activity. Given some of

the known effects of dopamine in glomerular networks, this could potentially have the effect of

counteracting input activity perturbation. For example, the best-described circuit-level effect of

dopamine in the bulb is its inhibitory action on the release of glutamate from olfactory sensory

neuron axon terminals, which it achieves via D2 receptor activation (Hsia *et al.*, 1999; Berkowicz &

Trombley, 2000; Ennis *et al.*, 2001; Maher & Westbrook, 2008; McGann, 2013; Vaaga *et al.*, 2017). If

sensory input levels are chronically decreased, it might be adaptive to reduce the strength of this

inhibition, perhaps by decreasing expression of the rate-limiting enzyme for neurotransmitter

synthesis. This might put the system into a state of higher-gain readiness for the low levels of input

that will initially arrive if and when sensory stimulation recommences. Indeed, the volume-

transmission nature of sensory neuron presynaptic inhibition (Pinching & Powell, 1971; McGann,

2013), coupled with the high threshold and slow dynamics of bulbar dopamine release (Borisovska *et*

al., 2013), might make for a slow modulatory process that is well suited to days-scale modification

via plasticity of enzyme expression. However, given the strong presynaptic inhibition also mediated

by GABA$_B$ receptor activation on OSN terminals (McGann, 2013), it is unclear why dopamine- but not

GABA-synthesising enzymes are targets for rapid activity-dependent regulation in this circuit. It may

be that GABAergic inhibition is modulated by distinct plastic mechanisms that do not target

transmitter synthesis. Alternatively, distinct mechanisms of GABAergic and dopaminergic inhibition

that can occur within the same presynaptic terminals (Burke *et al.*, 2018) might be individually tuned

according to sensory experience.

In contrast, the potential functional implications of TH plasticity on the less-understood postsynaptic

effects of bulbar dopamine transmission are harder to predict. Dopamine can act via D1-type

receptors to promote rebound firing after interglomerular GABAergic inhibition of external tufted

cells (Liu *et al.*, 2013), so perhaps decreased TH levels might lead to more prolonged lateral

inhibition within the glomerular layer. The implications of such an effect for information processing

in sensory-deprived bulbar circuits might be best addressed via computational network modelling.

Most important, though, is the fact that, although experience-dependent plasticity in olfactory bulb

TH expression has been described for decades, it is still not known how (or whether) changes in TH

mRNA and protein levels give rise to alterations in the amount of dopamine actually released.

Future experiments should be able to control the activity of dopaminergic neurons whilst monitoring

the resulting effects of dopamine release (e.g. Vaaga *et al.*, 2017), in order to quantitatively compare

the outcome in control versus sensory-deprived tissue.

Acknowledgements

This work was supported by a Medical Research Council 4-year PhD studentship to DJB, and a

Leverhulme Trust Research Grant (RPG-2016-095) and ERC Consolidator Grant (725729;

FUNCOPLAN) to MSG. We wish to thank Beatriz Rico for access to equipment, Laura Andreae and

Elisa Galliano for comments on the manuscript, and Elisa Galliano, Andres Crespo Vieira, Christiane

Hahn and other past and present members of the Grubb laboratory for technical assistance and

stimulating discussion.

References

Adam, Y. & Mizrahi, A. (2011) Long-term imaging reveals dynamic changes in the neuronal
composition of the glomerular layer. *J. Neurosci.*, **31**, 7967–7973.
Agoston, Z., Heine, P., Brill, M.S., Grebbin, B.M., Hau, A.-C., Kallenborn-Gerhardt, W., Schramm, J.,
Götz, M., & Schulte, D. (2014) Meis2 is a Pax6 co-factor in neurogenesis and dopaminergic
periglomerular fate specification in the adult olfactory bulb. *Development*, **141**, 28–38.
Akiba, Y., Sasaki, H., Saino-Saito, S., & Baker, H. (2007) Temporal and spatial disparity in cFOS
expression and dopamine phenotypic differentiation in the neonatal mouse olfactory bulb.
Neurochem. Res., **32**, 625–634.
Bäckman, C.M., Malik, N., Zhang, Y., Shan, L., Grinberg, A., Hoffer, B.J., Westphal, H., & Tomac, A.C.
(2006) Characterization of a mouse strain expressing Cre recombinase from the 3'
untranslated region of the dopamine transporter locus. *Genesis*, **44**, 383–390.
Baker, H. (1990) Unilateral, neonatal olfactory deprivation alters tyrosine hydroxylase expression but
not aromatic amino acid decarboxylase or GABA immunoreactivity. *Neuroscience*, **36**, 761–
771.
Baker, H., Cummings, D.M., Munger, S.D., Margolis, J.W., Franzen, L., Reed, R.R., & Margolis, F.L.
(1999) Targeted deletion of a cyclic nucleotide-gated channel subunit (OCNC1): biochemical
and morphological consequences in adult mice. *J. Neurosci.*, **19**, 9313–9321.
Baker, H., Kawano, T., Albert, V., Joh, T.H., Reis, D.J., & Margolis, F.L. (1984) Olfactory bulb dopamine
neurons survive deafferentation-induced loss of tyrosine hydroxylase. *Neuroscience*, **11**,
605–615.
Baker, H., Kawano, T., Margolis, F.L., & Joh, T.H. (1983) Transneuronal regulation of tyrosine
hydroxylase expression in olfactory bulb of mouse and rat. *J. Neurosci.*, **3**, 69–78.
Baker, H., Morel, K., Stone, D.M., & Maruniak, J.A. (1993) Adult naris closure profoundly reduces
tyrosine hydroxylase expression in mouse olfactory bulb. *Brain Res.*, **614**, 109–116.
Banerjee, A., Marbach, F., Anselmi, F., Koh, M.S., Davis, M.B., Garcia da Silva, P., Delevich, K., Oyibo,
H.K., Gupta, P., Li, B., & Albeanu, D.F. (2015) An Interglomerular Circuit Gates Glomerular
Output and Implements Gain Control in the Mouse Olfactory Bulb. *Neuron*, **87**, 193–207.
Banerjee, K., Akiba, Y., Baker, H., & Cave, J.W. (2013) Epigenetic control of neurotransmitter
expression in olfactory bulb interneurons. *Int. J. Dev. Neurosci.*, **31**, 415–423.
Banerjee, K., Wang, M., Cai, E., Fujiwara, N., Baker, H., & Cave, J.W. (2014) Regulation of tyrosine
hydroxylase transcription by hnRNP K and DNA secondary structure. *Nat Commun*, **5**, 5769.
Bastien-Dionne, P.-O., David, L.S., Parent, A., & Saghatelyan, A. (2010) Role of sensory activity on
chemospecific populations of interneurons in the adult olfactory bulb. *J. Comp. Neurol.*, **518**,
1847–1861.
Benito, N., Gaborieau, E., Sanz Diez, A., Kosar, S., Foucault, L., Raineteau, O., & De Saint Jan, D.
(2018) A Pool of Postnatally Generated Interneurons Persists in an Immature Stage in the
Olfactory Bulb. *J. Neurosci.*, **38**, 9870–9882.
Berkowicz, D.A. & Trombley, P.Q. (2000) Dopaminergic modulation at the olfactory nerve synapse.
Brain Res., **855**, 90–99.
Borisovska, M., Bensen, A.L., Chong, G., & Westbrook, G.L. (2013) Distinct modes of dopamine and
GABA release in a dual transmitter neuron. *J. Neurosci.*, **33**, 1790–1796.
Brill, M.S., Snapyan, M., Wohlfrom, H., Ninkovic, J., Jawerka, M., Mastick, G.S., Ashery-Padan, R.,
Saghatelyan, A., Berninger, B., & Götz, M. (2008) A dlx2- and pax6-dependent transcriptional
code for periglomerular neuron specification in the adult olfactory bulb. *J. Neurosci.*, **28**,
6439–6452.
Burke, K.J., Keeshen, C.M., & Bender, K.J. (2018) Two Forms of Synaptic Depression Produced by
Differential Neuromodulation of Presynaptic Calcium Channels. *Neuron*, **99**, 969-984.e7.
Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M., Mueller, R., Nolan, T.,
Pfaffl, M.W., Shipley, G.L., Vandesompele, J., & Wittwer, C.T. (2009) The MIQE guidelines:

minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.*, **55**, 611–622.

Bywalez, W.G., Ona-Jodar, T., Lukas, M., Ninkovic, J., & Egger, V. (2016) Dendritic Arborization Patterns of Small Juxtaglomerular Cell Subtypes within the Rodent Olfactory Bulb. *Front Neuroanat*, **10**, 127.

Cave, J.W., Akiba, Y., Banerjee, K., Bhosle, S., Berlin, R., & Baker, H. (2010) Differential regulation of dopaminergic gene expression by Er81. *J. Neurosci.*, **30**, 4717–4724.

Chand, A.N., Galliano, E., Chesters, R.A., & Grubb, M.S. (2015) A distinct subtype of dopaminergic interneuron displays inverted structural plasticity at the axon initial segment. *J. Neurosci.*, **35**, 1573–1590.

Cheetham, C.E.J., Park, U., & Belluscio, L. (2016) Rapid and continuous activity-dependent plasticity of olfactory sensory input. *Nat Commun*, **7**, 10729.

Cho, J.Y., Min, N., Franzen, L., & Baker, H. (1996) Rapid down-regulation of tyrosine hydroxylase expression in the olfactory bulb of naris-occluded adult rats. *J. Comp. Neurol.*, **369**, 264–276.

Cigola, E., Volpe, B.T., Lee, J.W., Franzen, L., & Baker, H. (1998) Tyrosine hydroxylase expression in primary cultures of olfactory bulb: role of L-type calcium channels. *J. Neurosci.*, **18**, 7638–7649.

Cockerham, R., Liu, S., Cachope, R., Kiyokage, E., Cheer, J.F., Shipley, M.T., & Puche, A.C. (2016) Subsecond Regulation of Synaptically Released Dopamine by COMT in the Olfactory Bulb. *J. Neurosci.*, **36**, 7779–7785.

Coppola, D.M. (2012) Studies of olfactory system neural plasticity: the contribution of the unilateral naris occlusion technique. *Neural Plast.*, **2012**, 351752.

Cummings, D.M., Henning, H.E., & Brunjes, P.C. (1997) Olfactory bulb recovery after early sensory deprivation. *J. Neurosci.*, **17**, 7433–7440.

Ennis, M., Zhou, F.M., Ciombor, K.J., Aroniadou-Anderjaska, V., Hayar, A., Borrelli, E., Zimmer, L.A., Margolis, F., & Shipley, M.T. (2001) Dopamine D2 receptor-mediated presynaptic inhibition of olfactory nerve terminals. *J. Neurophysiol.*, **86**, 2986–2997.

Fogli Iseppe, A., Pignatelli, A., & Belluzzi, O. (2016) Calretinin-Periglomerular Interneurons in Mice Olfactory Bulb: Cells of Few Words. *Front Cell Neurosci*, **10**, 231.

Fokkens, W.J., Lund, V.J., Mullol, J., Bachert, C., Alobid, I., Baroody, F., Cohen, N., Cervin, A., Douglas, R., Gevaert, P., Georgalas, C., Goossens, H., Harvey, R., Hellings, P., Hopkins, C., Jones, N., Joos, G., Kalogjera, L., Kern, B., Kowalski, M., Price, D., Riechelmann, H., Schlosser, R., Senior, B., Thomas, M., Toskala, E., Voegels, R., Wang, D.Y., & Wormald, P.J. (2012) EPOS 2012: European position paper on rhinosinusitis and nasal polyps 2012. A summary for otorhinolaryngologists. *Rhinology*, **50**, 1–12.

Galliano, E., Franzoni, E., Breton, M., Chand, A.N., Byrne, D.J., Murthy, V.N., & Grubb, M.S. (2018) Embryonic and postnatal neurogenesis produce functionally distinct subclasses of dopaminergic neuron. *Elife*, **7**.

Galliano, E., Hahn, C., Browne, L., Villamayor, P.R., & Grubb, M.S. (2020) Brief sensory deprivation triggers cell type-specific structural and functional plasticity in olfactory bulb neurons. *bioRxiv*, 2020.05.10.086926.

Grier, B.D., Belluscio, L., & Cheetham, C.E.J. (2016) Olfactory Sensory Activity Modulates Microglial-Neuronal Interactions during Dopaminergic Cell Loss in the Olfactory Bulb. *Front Cell Neurosci*, **10**, 178.

Hadjiconstantinou, M., Wemlinger, T.A., Sylvia, C.P., Hubble, J.P., & Neff, N.H. (1993) Aromatic L-amino acid decarboxylase activity of mouse striatum is modulated via dopamine receptors. *J. Neurochem.*, **60**, 2175–2180.

Halász, N., Ljungdahl, A., Hökfelt, T., Johansson, O., Goldstein, M., Park, D., & Biberfeld, P. (1977) Transmitter histochemistry of the rat olfactory bulb. I. Immunohistochemical localization of monoamine synthesizing enzymes. Support for intrabulbar, periglomerular dopamine neurons. *Brain Res.*, **126**, 455–474.

Hayar, A., Karnup, S., Shipley, M.T., & Ennis, M. (2004) Olfactory Bulb Glomeruli: External Tufted
Cells Intrinsically Burst at Theta Frequency and Are Entrained by Patterned Olfactory Input. *J.*
Neurosci., **24**, 1190–1199.
Heffner, C.S., Herbert Pratt, C., Babiuk, R.P., Sharma, Y., Rockwood, S.F., Donahue, L.R., Eppig, J.T., &
Murray, S.A. (2012) Supporting conditional mouse mutagenesis with a comprehensive cre
characterization resource. *Nat Commun*, **3**, 1218.
Hökfelt, T., Halasz, N., Ljungdahl, A., Johansson, O., Goldstein, M., & Park, D. (1975) Histochemical
support for a dopaminergic mechanism in the dendrites of certain periglomerular cells in the
rat olfactory bulb. *Neurosci. Lett.*, **1**, 85–90.
Hsia, A.Y., Vincent, J.D., & Lledo, P.M. (1999) Dopamine depresses synaptic inputs into the olfactory
bulb. *J. Neurophysiol.*, **82**, 1082–1085.
Kass, M.D., Pottackal, J., Turkel, D.J., & McGann, J.P. (2013) Changes in the neural representation of
odorants after olfactory deprivation in the adult mouse olfactory bulb. *Chem. Senses*, **38**,
77–89.
Kikuta, S., Sakamoto, T., Nagayama, S., Kanaya, K., Kinoshita, M., Kondo, K., Tsunoda, K., Mori, K., &
Yamasoba, T. (2015) Sensory deprivation disrupts homeostatic regeneration of newly
generated olfactory sensory neurons after injury in adult mice. *J. Neurosci.*, **35**, 2657–2673.
Kiyokage, E., Pan, Y.-Z., Shao, Z., Kobayashi, K., Szabo, G., Yanagawa, Y., Obata, K., Okano, H., Toida,
K., Puche, A.C., & Shipley, M.T. (2010) Molecular identity of periglomerular and short axon
cells. *J. Neurosci.*, **30**, 1185–1196.
Kohwi, M., Petryniak, M.A., Long, J.E., Ekker, M., Obata, K., Yanagawa, Y., Rubenstein, J.L.R., &
Alvarez-Buylla, A. (2007) A subpopulation of olfactory bulb GABAergic interneurons is
derived from Emx1- and Dlx5/6-expressing progenitors. *J. Neurosci.*, **27**, 6878–6891.
Kosaka, K. & Kosaka, T. (2007) Chemical properties of type 1 and type 2 periglomerular cells in the
mouse olfactory bulb are different from those in the rat olfactory bulb. *Brain Res.*, **1167**, 42–
55.
Kosaka, T. & Kosaka, K. (2011) "Interneurons" in the olfactory bulb revisited. *Neurosci. Res.*, **69**, 93–
99.
Kosaka, T., Kosaka, K., Hama, K., Wu, J.Y., & Nagatsu, I. (1987) Differential effect of functional
olfactory deprivation on the GABAergic and catecholaminergic traits in the rat main olfactory
bulb. *Brain Res.*, **413**, 197–203.
Kosaka, T., Pignatelli, A., & Kosaka, K. (2019) Heterogeneity of tyrosine hydroxylase expressing
neurons in the main olfactory bulb of the mouse. *Neurosci. Res.*,.
Lammel, S., Steinberg, E.E., Földy, C., Wall, N.R., Beier, K., Luo, L., & Malenka, R.C. (2015) Diversity of
transgenic mouse models for selective targeting of midbrain dopamine neurons. *Neuron*, **85**,
429–438.
Lau, C.G. & Murthy, V.N. (2012) Activity-dependent regulation of inhibition via GAD67. *J. Neurosci.*,
32, 8521–8531.
Liu, N., Cigola, E., Tinti, C., Jin, B.K., Conti, B., Volpe, B.T., & Baker, H. (1999) Unique regulation of
immediate early gene and tyrosine hydroxylase expression in the odor-deprived mouse
olfactory bulb. *J. Biol. Chem.*, **274**, 3042–3047.
Liu, S., Plachez, C., Shao, Z., Puche, A., & Shipley, M.T. (2013) Olfactory bulb short axon cell release of
GABA and dopamine produces a temporally biphasic inhibition-excitation response in
external tufted cells. *J. Neurosci.*, **33**, 2916–2926.
Liu, S., Puche, A.C., & Shipley, M.T. (2016) The Interglomerular Circuit Potently Inhibits Olfactory
Bulb Output Neurons by Both Direct and Indirect Pathways. *J. Neurosci.*, **36**, 9604–9617.
Livak, K.J. & Schmittgen, T.D. (2001) Analysis of Relative Gene Expression Data Using Real-Time
Quantitative PCR and the 2−ΔΔCT Method. *Methods*, **25**, 402–408.
Maher, B.J. & Westbrook, G.L. (2008) Co-transmission of dopamine and GABA in periglomerular
cells. *J. Neurophysiol.*, **99**, 1559–1564.

Mandairon, N., Sacquet, J., Jourdan, F., & Didier, A. (2006) Long-term fate and distribution of
newborn cells in the adult mouse olfactory bulb: Influences of olfactory deprivation.
Neuroscience, **141**, 443–451.

McGann, J.P. (2013) Presynaptic inhibition of olfactory sensory neurons: new mechanisms and
potential functions. *Chem. Senses*, **38**, 459–474.

Nadi, N.S., Head, R., Grillo, M., Hempstead, J., Grannot-Reisfeld, N., & Margolis, F.L. (1981) Chemical
deafferentation of the olfactory bulb: plasticity of the levels of tyrosine hydroxylase,
dopamine and norepinephrine. *Brain Res.*, **213**, 365–377.

Ninkovic, J., Pinto, L., Petricca, S., Lepier, A., Sun, J., Rieger, M.A., Schroeder, T., Cvekl, A., Favor, J., &
Götz, M. (2010) The transcription factor Pax6 regulates survival of dopaminergic olfactory
bulb neurons via crystallin αA. *Neuron*, **68**, 682–694.

Panzanelli, P., Fritschy, J.M., Yanagawa, Y., Obata, K., & Sassoè-Pognetto, M. (2007) GABAergic
phenotype of periglomerular cells in the rodent olfactory bulb. *Journal of Comparative
Neurology*, **502**, 990–1002.

Parrish-Aungst, S., Kiyokage, E., Szabo, G., Yanagawa, Y., Shipley, M.T., & Puche, A.C. (2011) Sensory
experience selectively regulates transmitter synthesis enzymes in interglomerular circuits.
Brain Res., **1382**, 70–76.

Parrish-Aungst, S., Shipley, M.T., Erdelyi, F., Szabo, G., & Puche, A.C. (2007) Quantitative analysis of
neuronal diversity in the mouse olfactory bulb. *J. Comp. Neurol.*, **501**, 825–836.

Philpot, B.D., Men, D., McCarty, R., & Brunjes, P.C. (1998) Activity-dependent regulation of
dopamine content in the olfactory bulbs of naris-occluded rats. *Neuroscience*, **85**, 969–977.

Pignatelli, A., Ackman, J.B., Vigetti, D., Beltrami, A.P., Zucchini, S., & Belluzzi, O. (2009) A potential
reservoir of immature dopaminergic replacement neurons in the adult mammalian olfactory
bulb. *Pflugers Arch.*, **457**, 899–915.

Pignatelli, A. & Belluzzi, O. (2017) Dopaminergic Neurones in the Main Olfactory Bulb: An Overview
from an Electrophysiological Perspective. *Front Neuroanat*, **11**, 7.

Pignatelli, A., Kobayashi, K., Okano, H., & Belluzzi, O. (2005) Functional properties of dopaminergic
neurones in the mouse olfactory bulb. *J. Physiol. (Lond.)*, **564**, 501–514.

Pinching, A.J. & Powell, T.P. (1971) The neuropil of the periglomerular region of the olfactory bulb. *J.
Cell. Sci.*, **9**, 379–409.

Rosser, A.E., Hökfelt, T., & Goldstein, M. (1986) LHRH and catecholamine neuronal systems in the
olfactory bulb of the mouse. *Journal of Comparative Neurology*, **250**, 352–363.

Saino-Saito, S., Sasaki, H., Volpe, B.T., Kobayashi, K., Berlin, R., & Baker, H. (2004) Differentiation of
the dopaminergic phenotype in the olfactory system of neonatal and adult mice. *J. Comp.
Neurol.*, **479**, 389–398.

Sato, T., Homma, R., & Nagayama, S. (2020) Direct comparison of odor responses of homologous
glomeruli in the medial and lateral maps of the mouse olfactory bulb. *eNeuro*,.

Short, S.M. & Wachowiak, M. (2019) Temporal Dynamics of Inhalation-Linked Activity across Defined
Subpopulations of Mouse Olfactory Bulb Neurons Imaged In Vivo. *eNeuro*, **6**.

Stone, D.M., Grillo, M., Margolis, F.L., Joh, T.H., & Baker, H. (1991) Differential effect of functional
olfactory bulb deafferentation on tyrosine hydroxylase and glutamic acid decarboxylase
messenger RNA levels in rodent juxtaglomerular neurons. *J. Comp. Neurol.*, **311**, 223–233.

Stone, D.M., Wessel, T., Joh, T.H., & Baker, H. (1990) Decrease in tyrosine hydroxylase, but not
aromatic L-amino acid decarboxylase, messenger RNA in rat olfactory bulb following
neonatal, unilateral odor deprivation. *Brain Res. Mol. Brain Res.*, **8**, 291–300.

Tavakoli, A., Schmaltz, A., Schwarz, D., Margrie, T.W., Schaefer, A.T., & Kollo, M. (2018) Quantitative
Association of Anatomical and Functional Classes of Olfactory Bulb Neurons. *J. Neurosci.*, **38**,
7204–7220.

Taylor, S.C., Nadeau, K., Abbasi, M., Lachance, C., Nguyen, M., & Fenrich, J. (2019) The Ultimate qPCR
Experiment: Producing Publication Quality, Reproducible Data the First Time. *Trends in
Biotechnology*, **37**, 761–774.

Vaaga, C.E., Yorgason, J.T., Williams, J.T., & Westbrook, G.L. (2017) Presynaptic gain control by
endogenous cotransmission of dopamine and GABA in the olfactory bulb. *J. Neurophysiol.*,
117, 1163–1170.
Vlug, A.S., Teuling, E., Haasdijk, E.D., French, P., Hoogenraad, C.C., & Jaarsma, D. (2005) ATF3
expression precedes death of spinal motoneurons in amyotrophic lateral sclerosis-SOD1
transgenic mice and correlates with c-Jun phosphorylation, CHOP expression, somato-
dendritic ubiquitination and Golgi fragmentation. *Eur. J. Neurosci.*, **22**, 1881–1894.
Wachowiak, M., Economo, M.N., Díaz-Quesada, M., Brunert, D., Wesson, D.W., White, J.A., &
Rothermel, M. (2013) Optical dissection of odor information processing in vivo using
GCaMPs expressed in specified cell types of the olfactory bulb. *J. Neurosci.*, **33**, 5285–5300.
Wang, M., Cai, E., Fujiwara, N., Fones, L., Brown, E., Yanagawa, Y., & Cave, J.W. (2017) Odorant
Sensory Input Modulates DNA Secondary Structure Formation and Heterogeneous
Ribonucleoprotein Recruitment on the Tyrosine Hydroxylase and Glutamic Acid
Decarboxylase 1 Promoters in the Olfactory Bulb. *J. Neurosci.*, **37**, 4778–4789.
Weiss, J., Pyrski, M., Jacobi, E., Bufe, B., Willnecker, V., Schick, B., Zizzari, P., Gossage, S.J., Greer,
C.A., Leinders-Zufall, T., Woods, C.G., Wood, J.N., & Zufall, F. (2011) Loss-of-function
mutations in sodium channel Nav1.7 cause anosmia. *Nature*, **472**, 186–190.
Wilson, D.A. & Sullivan, R.M. (1995) The D2 antagonist spiperone mimics the effects of olfactory
deprivation on mitral/tufted cell odor response patterns. *J. Neurosci.*, **15**, 5574–5581.
Zeisel, A., Hochgerner, H., Lönnerberg, P., Johnsson, A., Memic, F., van der Zwan, J., Häring, M.,
Braun, E., Borm, L.E., La Manno, G., Codeluppi, S., Furlan, A., Lee, K., Skene, N., Harris, K.D.,
Hjerling-Leffler, J., Arenas, E., Ernfors, P., Marklund, U., & Linnarsson, S. (2018) Molecular
Architecture of the Mouse Nervous System. *Cell*, **174**, 999-1014.e22.
Zhu, M.Y., Juorio, A.V., Paterson, I.A., & Boulton, A.A. (1993) Regulation of striatal aromatic L-amino
acid decarboxylase: effects of blockade or activation of dopamine receptors. *Eur. J.*
Pharmacol., **238**, 157–164.
957