Massively parallel functional testing of *MSH2* missense variants conferring Lynch Syndrome risk

Xiaoyan Jia[1], Bala Bharathi Burugula[1], Victor Chen[1], Rosemary M. Lemons[1], Sajini Jayakody[1], Mariam Maksutova[1], Jacob O. Kitzman[1,2,*]

[1]Department of Human Genetics, University of Michigan, Ann Arbor, MI, USA
[2]Department of Computational Medicine & Bioinformatics, University of Michigan, Ann Arbor, MI, USA
*correspondence to kitzmanj@umich.edu

Introduction

The lack of functional evidence for the majority of missense variants limits their clinical interpretability, and poses a key barrier to the broad utility of carrier screening. In Lynch Syndrome (LS), one of the most highly prevalent cancer syndromes, nearly 90% of clinically observed missense variants are deemed 'variants of uncertain significance' (VUS). To systematically resolve their functional status, we performed a massively parallel screen in human cells to identify loss-of-function missense variants in the key DNA mismatch repair factor *MSH2*. The resulting sequence-function map is substantially complete, covering 94% of the 17,746 possible variants, and is highly concordant (≥96%) with existing functional data and expert clinicians' interpretations. The large majority (89%) of missense variants were functionally neutral, perhaps unexpectedly in view of its evolutionary conservation. These data provide ready-to-use functional evidence to resolve the ~1,300 extant missense VUSs in *MSH2*, and may facilitate the prospective classification of newly discovered variants in the clinic.

Main text

Lynch Syndrome (LS, OMIM:120435), the first discovered familial cancer syndrome[1], confers predisposition to colorectal and endometrial cancers due to the loss of DNA mismatch repair (MMR) and resulting mutagenic burden. Most affected individuals inherit a single loss-of-function variant in one of four MMR factors (*MSH2, MLH1, MSH6, PMS2*), followed by a somatic 'second hit' inactivating the remaining functional copy. Due to high population prevalence (≥1:300)[2,3], clear genetic etiology, and potential for prevention through intensified surveillance, MMR gene variants are considered clinically actionable[4].

Missense changes together comprise 20-30% of LS variants[5], and are particularly challenging to interpret: their functional impacts may range from minimal to profound, most are individually rare, and they exhibit incomplete penetrance[6]. Consequently, the overwhelming majority of LS gene missense variants listed in ClinVar (4,762/5,473, 87.0%) are deemed 'variants of uncertain significance', or VUS **(Supplementary Fig.1)**, and cannot be used to guide diagnosis.

To address these challenges, we performed a deep mutational scan[7] to systematically measure the functional impact of missense variants in the major Lynch Syndrome gene *MSH2.* We established a human cell system to model *MSH2* variant function using the mismatch repair proficient[8] cell line HAP1 **(Fig. 1a,c)**. First, to disrupt

Fig. 1 | Overview of *MSH2* functional screen. a, Generating and testing *MSH2* variant function in isogenic human cells. Cells with pathogenic loss-of-function variants (top) are resistant to 6-thioguanine (6-TG), while complementation by a functional variant (middle) restores 6-TG sensitivity. In a pooled screen (bottom), a mixed library of *MSH2* missense variants is transduced, and 6-TG treatment selects for loss-of-function variants. **b,** 6-TG resistance reflects MMR function in human cells. Clonogenic assay of HAP1 *MSH2* KO cells transduced with either functional (wild-type) *MSH2* cDNA (upper row) or loss-of-function missense variant A636P (bottom row), without (left) or with 6-TG (1 µM, right). **c,** Readout using deep sequencing to quantify abundances of each *MSH2* allele before and after 6-TG selection. The resulting loss-of-function (LOF) score is positive for deleterious variants and negative for functionally neutral ones.

MMR, we derived clonal *MSH2* knockout cells bearing a 19-bp frameshift deletion in *MSH2* exon 6, and
verified that it lacked detectable MSH2 protein expression **(Supplementary Fig. 2)**. To restore MMR, we
stably reintroduced into cells either wild-type *MSH2* cDNA (KO+WT) or a pathogenic founder allele
(KO+A636P) on inducible lentiviral constructs. After inducing expression with doxycycline, wild-type
MSH2 protein reached near-endogenous levels, while A636P was only barely detectable, consistent with
the known destabilizing effect of this variant[9] **(Supplementary Fig. 3)**.

As a readout for *MSH2* function, we leveraged selection with the purine analog 6-thioguanine (6-TG),
which has been used to assay MMR activity in diverse systems[10,11]. Incorporation of 6-TG is selectively
toxic to MMR-proficient cells, as it creates lesions that the MMR machinery recognizes but cannot repair,
culminating in replication arrest[12]. As expected, KO cells showed increased 6-TG resistance compared
to parental HAP1 cells, while 6-TG sensitivity was restored by reintroduction of wild-type *MSH2* but not
the pathogenic A636P mutant **(Fig. 1b** and **Supplementary Fig. 3)**. To mimic simultaneously testing
thousands of mutations, we prepared a mixed culture of KO+WT cells and KO+A636P cells, with the
respective *MSH2* expression constructs bearing distinct, identifying barcode libraries *in cis*
(Supplementary Fig. 4). We quantified barcode abundances by deep sequencing (**Methods**), and found
barcodes linked to the pathogenic variant A636P were strongly enriched, relative to WT barcodes, by 6-
TG treatment (median fold change: 179, *P*<2.2x10^{-16}, Kolmogorov-Smirnov test). This result confirms the
feasibility of highly multiplexed MMR functional tests in isogenic human cell mixtures.

To model known and novel *MSH2* variants, we applied single amino-acid saturation mutagenesis[13] to
generate libraries comprising every possible missense, synonymous and nonsense variant (**Methods**)
MSH2 cDNA (2,802 bp) was divided into 21 short tiles, to enable tracking by amplicon sequencing, and
to mitigate the impact of recombination during lentiviral replication[14]. The mutant cDNA library for each
tile was cloned into the inducible lentiviral vector mentioned above, followed by a C-terminal 2A linker
and a blasticidin resistance gene to permit selection for complete in-frame cDNA expression
(Supplementary Fig. 3). HAP1 *MSH2* KO cells were transduced at low multiplicity (<0.1) to yield a
population in which each cell expressed a single mutant *MSH2* variant. Sequencing of integrated libraries
from genomic DNA confirmed that nearly all single-codon mutations were present (97.3%) with highly
uniform representation **(Supplementary Fig. 5)**.

Each *MSH2* mutant cell pool was then selected *en masse* for MMR deficiency. To allow comparison
across pools, each was spiked with barcoded control cells (KO+WT: 10% of cells; KO+A636P: 0.5%).
Pools were then grown under selective (6-TG+blasticidin) or mock (blasticidin only) conditions for two
passages **(Supplementary Fig. 6)**. As expected, the spiked-in *MSH2* A636P cells were consistently
enriched over WT by 6-TG selection relative to mock treatment (median fold change: 105;
Supplementary Fig. 7).

To track the functional status of each *MSH2* variant, we performed amplicon sequencing of each tile
throughout the course of selection (**Supplementary Fig. 7-9**). Loss-of-function (LOF) scores for each
variant were calculated as the log$_2$-ratio of that variant's frequency after 6-TG treatment divided by its
frequency after mock treatment, such that deleterious variants should score positively and neutral
variants negatively **(Fig. 1c)**. LOF scores strongly correlated across replicate 6-TG selections (mean
pairwise Pearson's r=0.77, **Supplementary Fig. 10**). As expected, mock treatment had little effect upon
most variants' frequencies, except for truncating variants which terminate translation before the
downstream blasticidin resistance marker and were therefore depleted (**Supplementary Fig. 11**).

The resulting sequence-function map was substantially complete, covering 94.4% of possible *MSH2*
single-codon missense variants **(Fig. 2)**. Missense variant scores were bimodal, with the large majority
(89.4%) scoring negatively, suggesting a high degree of tolerance to single amino acid substitution (**Fig.
2g**). Indeed, for over half of MSH2 residues (510/934), substitution to any other amino acid was tolerated.
As expected, synonymous variants predominantly scored negatively (820/841, 97.5%), with exceptions
likely caused by second-site mutations outside the sequenced tile. Most of the scored missense variants
were represented by multiple equivalent codons (on average, 3.0), and summarizing these internal

replicates i nto a single, amino-acid
level LOF score res ulted in an overall
false discovery rate of 0.95%. To verify
these results were generalizable
across cell types, we repeated the
screen for two tiles (tile 14 and 15) in
another MMR-proficient human cell
line, HEK293, with largely concordant
results (**Supplementary Fig. 12**).

We sought to validate these pooled
measurements by comparison to
traditional, low-throughput functional
studies. We collated a set of 126
MSH2 missense variants previously
characterized as functionally
deleterious or neutral by individual
cell-based[10,11,15–19] and/or biochemical
assays[20,21] (**Supplementary Tables
1-2**). For the great majority (121/126,
96.0%; **Fig. 3a**), our function scores
agreed with earlier reports. Among 3 of
the 5 discordant variants, our function
scores were internally consistent
across multiple codons encoding the
same amino acid substitution, and with
similar substitutions at the same
residue (**Supplementary Fig. 13**).
These may reflect false negatives on
the part of the previous screens, or
differences between this human cell
model and previous yeast or *in vitro*
assays.

Bioinformatic tools are frequently used
to interpret clinical variants in the
absence of other evidence[22]. We
compared our LOF scores'
classification performance with that of
six computational predictors, taking
the previously reported functional
evidence as a truth set (**Fig. 3b**). Our
LOF scores classified these variants
with a Matthews Correlation
Coefficient of 0.93, higher than all six
tools' (range: 0.50-0.77). General-
purpose, pathway-agnostic classifiers
exhibited poor specificity, possibly
reflecting the challenge of
distinguishing functional effects
between variants in highly conserved
genes such as *MSH2.* Although the
bioinformatic predictions were
modestly correlated with our

Fig. 2 | Sequence function map of *MSH2* missense variants. a, MSH2 secondary structure (cyan: alpha helices, pink: beta sheets), and **b**, protein domains. **c**, Heatmap of loss-of-function (LOF) scores after mutating each of 934 residues in MSH2 (rows) to each of 19 other possible amino acids (columns). Negative scores (blue) indicate functionally neutral variants, while positive scores (red) indicate deleterious ones; Yellow denotes wild-type amino acid, gray: no data. **d**, Fractions of substitutions which are disruptive (LOF score>0) by position. **e**, Number of scored missense variants by position (max. possible: 19). **f**, Evolutionary conservation score (PSIC) by position. **g**, Histogram of scores by variant class.

experimental measurements for
variants in the functional validation
set, they were markedly less
concordant for other variants at those
same residues, or throughout *MSH2*
at large (**Supplementary Fig. 14**). As
variant effect predictors are often
trained on the limited number of
known variants with available
classifications, their divergence with
our experimental measurements may
reflect overfitting.

To assess the suitability of these
scores for classifying variant
pathogenicity, we intersected them
with clinical variant databases. We
retrieved all *MSH2* missense variants
with consensus annotations from
InSight[5] and/or expert review in
ClinVar, totaling 75 variants. After
excluding 9 variants previously known
or predicted to impact splicing[23]
(SpliceAI score>0.2), our LOF scores
agreed with clinical annotations for 63
of the 66 remaining variants (**Fig. 3c**).
These data provide evidence to
facilitate interpretation of the 1,374
variants without classification (i.e.,
VUS) or with conflicting reports. Our
scores predicted that 112/1,374
(8.2%) of these unresolved variants
are functionally deleterious, similar to
the overall rate of LOF among all
missense variants reachable by a
single base mutation (7.7%; *P*=0.51,
two-sided binomial test). This result

Fig. 3 | Concordance with existing functional and clinical classifications. a, Measured LOF scores (y-axis) for published *MSH2* missense variants (n=209 reports covering 126 distinct variants), previously characterized by individual cellular[10,11,15–19] or biochemical assays[20,21]. **b**, Precision-recall curves show LOF scores outperform all tested bioinformatic callers in predicting functional status of variants in (**a**), with area under the curve indicated for each. **c**, LOF scores for patient missense variants from ClinVar and InSight databases, by variant classification. **d**, LOF scores versus allele frequency for missense variants in the gnomAD database[24]; size of circle denotes number of heterozygous variants; red squares mark missense variants observed in homozygous state in ≥1 individual. Gray region denotes singletons/doubletons (gnomAD allele count ≤2 individuals).

highlights the pervasiveness of benign missense variation in clinical databases and identifies a subset of
variants likely to be causal for LS.

We next examined how these function scores varied in apparently healthy populations (**Fig. 3d**). The
gnomAD database[24] lists 744 of the scored *MSH2* missense variants, nearly all rare (742/744 with minor
allele frequency < 0.5%). Among these, only 18 scored as deleterious in our assay (all singletons or
doubletons except the founder allele A636P), a significant depletion (*P*=6.9x10⁻⁷, two-sided binomial test)
relative to randomly drawn, size-matched sets of single-base missense variants. All 14 *MSH2* missense
variants observed in the homozygous state in gnomAD scored as functionally neutral, as expected given
that inherited biallelic MMR loss causes pediatric-onset cancer syndromes. These results reflect
survivorship bias which depletes adult cohorts of deleterious variants in genes underlying disorders such
as LS, as previously noted for *MSH2* deletions[25].

Deep mutational scans can inform protein structure-function relationships[26]. On aggregate, chemically
conservative substitutions across MSH2 were rarely deleterious (5.63%), compared to substitutions
replacing hydrophobic with charged residues (28.0%; **Supplementary Fig. 15**). Mapping function scores
onto the crystal structure[27] (**Fig. 4a**), we noted several regions of mutational intolerance corresponding

Fig. 4 | Structural insights from deep mutational scanning data. a, MutSα crystal structure (PDB:2O8E), with MSH6 surface in light gray, and MSH2 shaded by the number of deleterious missense variants at each residue (length of bar in legend proportional to count of residues within denoted range of LOF mutations). **b**, Extreme intolerance to substitution in MSH2 ATPase domain (upper), and connector domain (lower); selected residues or beta sheets shown with spheres or arrows, respectively. **c**. LOF score heatmap (as in **Fig. 2c**) showing mutational intolerance at Walker A motif of the ATPase domain (upper) and selected connector domain residues (lower) corresponding to (**b**).

to structural features. The phosphate-binding Walker A motif (residues 674-676) is required for the
essential ATPase activity of MSH2[28] and was almost completely intolerant to mutation (50/55 mutations
LOF, **Fig. 4b**). We also noticed strong constraint at glycine 126, possibly reflecting the requirement for
flexibility to accommodate conformational shifts between the mismatch binding and connector domains
(**Fig. 4c**)[27]. The connector domain's hydrophobic core, comprising three beta sheets and inward-facing
residues of the surrounding helices, was also largely functionally intolerant to substitutions. Conversely,
despite its name, the N-terminal mismatch binding domain (residues 1-124) was largely devoid of
missense LOF variants, consistent with its dispensability in yeast[29], and observations that *MSH2* start
codon loss is not strongly pathogenic in humans[30]. By contrast, prokaryotic mutS homodimerizes, with
both N termini making extensive DNA contacts essential for mismatch recognition[31]. Several positively
charged clamp domain residues (K512, R524, K546), which interact with the DNA backbone by hydrogen
bonding[27], were strongly intolerant to substitution, especially to negatively charged residues. Although
computational predictions of protein stability[32] showed modest overall correlation with LOF scores
(Pearson's r=0.39, **Supplementary Fig. 14**), we noted a few exceptions including the ATPase Walker
motifs, which showed strong constraint in our data but were not predicted to be destabilizing
(**Supplementary Fig. 15**), suggesting mechanisms other than protein instability.

We also examined the scores of variants accepted by evolution. Given the strong pressure to maintain
genome integrity, *MSH2* variants rising to fixation in extant lineages are expected to be functionally
neutral, except for potential epistatic effects of other mutations *in cis* or trans[33]. We aligned the human
MSH2 protein sequence with seven orthologs distributed across eukaryota, identifying 1707 amino acid
substitutions relative to the human sequence. Of those scored, nearly all (1,627/1,639, 99.3%) had
negative LOF scores (**Supplementary Fig. 16**), indicating these substitutions are functionally neutral in
the context of the human MSH2 sequence, and supporting these scores' specificity.

We have described a nearly complete sequence-function map for the key Lynch Syndrome gene *MSH2*,
and showed its predictive value in separating functionally intact variants from deleterious ones. This
screen exploited the direct link between molecular dysfunction – the inability to correct replicative errors
– and pathogenesis in Lynch Syndrome, and provides a blueprint for comprehensive functional
annotation of the three other major LS genes. A limitation of our cDNA-based measurements is that they
cannot model effects upon splicing, although among known LS variants and in other saturation screens[34],

splice disruption accounts for only a small fraction of pathogenic variants. Other genes with more varied
functions, or unknown pathogenic mechanisms, may require multiple assays for different aspects of
function, or may benefit from more generalized functional readouts such as protein stability[35],
localization[36], or target expression[37]. These functional scores, when rationally combined with other lines
of clinical evidence[22], will enable more accurate *MSH2* variant classification, and provide carriers of
deleterious variants the opportunity to learn their status and seek medical interventions[38].
254

Acknowledgements

We thank M. Meisler, A. Antonellis, E. Stoffel, S. Hemker, and A. Glazer for helpful discussions, J. Moran for providing sequencer access, and J. Moldovan for assistance with clonogenic assays. This work was supported by the National Institute of General Medical Sciences (5R01GM129123 to J.O.K.) and a Precision Health Fellowship from the University of Michigan (to X.J.).

Author contributions

J.O.K. conceived of and supervised the project. J.O.K. and X.J. designed the experiments. X.J. and B.B. executed the experiments with assistance from V.C., R.M.L., S.J., and M.M. X.J. and J.O.K. performed the data analysis. X.J. and J.O.K. prepared the manuscript with input from all other authors.

Competing interests

The authors declare no competing interests.

267
268

References

1. Lynch, H. T. & Krush, A. J. Cancer family 'G' revisited: 1895-1970. *Cancer* **27**, 1505–1511 (1971).
2. Win, A. K. *et al.* Prevalence and Penetrance of Major Genes and Polygenes for Colorectal Cancer. *Cancer Epidemiol. Biomarkers Prev.* **26**, 404–412 (2017).
3. Haraldsdottir, S. *et al.* Comprehensive population-wide analysis of Lynch syndrome in Iceland reveals founder mutations in MSH6 and PMS2. *Nat. Commun.* **8**, 14755 (2017).
4. Kalia, S. S. *et al.* Recommendations for reporting of secondary findings in clinical exome and genome sequencing, 2016 update (ACMG SF v2. 0): a policy statement of the American College of Medical Genetics and Genomics. *Genet. Med.* **19**, 249 (2017).
5. Thompson, B. A. *et al.* Application of a 5-tiered scheme for standardized classification of 2,360 unique mismatch repair gene variants in the InSiGHT locus-specific database. *Nat. Genet.* **46**, 107–115 (2014).
6. Hampel, H. & de la Chapelle, A. How do we approach the goal of identifying everybody with Lynch Syndrome? *Fam. Cancer* (2013).
7. Fowler, D. M., Stephany, J. J. & Fields, S. Measuring the activity of protein variants on a large scale using deep mutational scanning. *Nat. Protoc.* **9**, 2267–2284 (2014).
8. Harmsen, T., Klaasen, S., van de Vrugt, H. & Te Riele, H. DNA mismatch repair and oligonucleotide end-protection promote base-pair substitution distal from a CRISPR/Cas9-induced DNA break. *Nucleic Acids Res.* **46**, 2945–2955 (2018).
9. Foulkes, W. D. *et al.* The Founder Mutation MSH2*1906G→C Is an Important Cause of Hereditary Nonpolyposis Colorectal Cancer in the Ashkenazi Jewish Population. *Am. J. Hum. Genet.* **71**, 1395–1412 (2002).
10. Gammie, A. E. *et al.* Functional characterization of pathogenic human MSH2 missense mutations in Saccharomyces cerevisiae. *Genetics* **177**, 707–721 (2007).
11. Houlleberghs, H. *et al.* Oligonucleotide-directed mutagenesis screen to identify pathogenic Lynch syndrome-associated MSH2 DNA mismatch repair gene variants. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 4128–4133 (2016).
12. Yan, T., Berry, S. E., Desai, A. B. & Kinsella, T. J. DNA mismatch repair (MMR) mediates 6-thioguanine genotoxicity by introducing single-strand breaks to signal a G2-M arrest in MMR-proficient RKO cells. *Clin. Cancer Res.* **9**, 2327–2334 (2003).
13. Kitzman, J. O., Starita, L. M., Lo, R. S., Fields, S. & Shendure, J. Massively parallel single-amino-acid mutagenesis. *Nat. Methods* **12**, 203–6, 4 p following 206 (2015).
14. Sack, L. M., Davoli, T., Xu, Q., Li, M. Z. & Elledge, S. J. Sources of Error in Mammalian Genetic Screens. *G3* **6**, 2781–2790 (2016).
15. Rath, A. *et al.* Functional interrogation of Lynch syndrome-associated MSH2 missense variants via CRISPR-Cas9 gene editing in human embryonic stem cells. *Hum. Mutat.* **40**, 2044–2056 (2019).
16. Martinez, S. L. & Kolodner, R. D. Functional analysis of human mismatch repair gene mutations identifies weak alleles and polymorphisms capable of polygenic interactions. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 5070–5075 (2010).
17. Mastrocola, A. S. & Heinen, C. D. Lynch syndrome-associated mutations in MSH2 alter DNA repair and checkpoint response functions in vivo. *Hum. Mutat.* **31**, E1699–E1708 (2010).
18. Wielders, E. A. L. *et al.* Functional analysis of MSH2 unclassified variants found in suspected Lynch syndrome patients reveals pathogenicity due to attenuated mismatch repair. *J. Med. Genet.* **51**, 245–253 (2014).
19. Brown, G. D. Biochemical and Cellular Studies of Human MSH2 ATPase Domain Mutants Support the Molecular Switch Model of Mismatch Repair In Vivo. (University of Connecticut, 2014).
20. Drost, M. *et al.* A rapid and cell-free assay to test the activity of lynch syndrome-associated MSH2 and MSH6 missense variants. *Human Mutation* vol. 33 488–494 (2012).
21. Drost, M. *et al.* A functional assay–based procedure to classify mismatch repair gene variants in Lynch syndrome. *Genet. Med.* **21**, 1486–1496 (2019).
22. Brnich, S. E. *et al.* Recommendations for application of the functional evidence PS3/BS3 criterion using the ACMG/AMP sequence variant interpretation framework. *Genome Med.* **12**, 3 (2019).
23. Jaganathan, K. *et al.* Predicting Splicing from Primary Sequence with Deep Learning. *Cell* **176**,

535–548.e24 (2019).

24. Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature* **581**, 434–443 (2020).

25. Aguirre, M., Rivas, M. A. & Priest, J. Phenome-wide Burden of Copy-Number Variation in the UK Biobank. *Am. J. Hum. Genet.* **105**, 373–383 (2019).

26. Gray, V. E., Hause, R. J. & Fowler, D. M. Analysis of Large-Scale Mutagenesis Data To Assess the Impact of Single Amino Acid Substitutions. *Genetics* **207**, 53–61 (2017).

27. Warren, J. J. *et al.* Structure of the Human MutSα DNA Lesion Recognition Complex. *Mol. Cell* **26**, 579–592 (2007).

28. Graham, W. J., Putnam, C. D. & Kolodner, R. D. The properties of Msh2–Msh6 ATP binding mutants suggest a signal amplification mechanism in DNA mismatch repair. *J. Biol. Chem.* (2018).

29. Kumar, C. *et al.* Multiple Factors Insulate Msh2–Msh6 Mismatch Repair Activity from Defects in Msh2 Domain I. *J. Mol. Biol.* **411**, 765–780 (2011).

30. Cyr, J. L., Brown, G. D., Stroop, J. & Heinen, C. D. The predicted truncation from a cancer-associated variant of the MSH2 initiation codon alters activity of the MSH2-MSH6 mismatch repair complex. *Molecular Carcinogenesis* vol. 51 647–658 (2012).

31. Obmolova, G., Ban, C., Hsieh, P. & Yang, W. Crystal structures of mismatch repair protein MutS and its complex with a substrate DNA. *Nature* vol. 407 703–710 (2000).

32. Nielsen, S. V. *et al.* Predicting the impact of Lynch syndrome-causing missense mutations from structural calculations. *PLoS Genet.* **13**, e1006739 (2017).

33. Jordan, D. M. *et al.* Identification of cis-suppression of human disease mutations by comparative genomics. *Nature* **524**, 225–229 (2015).

34. Findlay, G. M. *et al.* Accurate classification of BRCA1 variants with saturation genome editing. *Nature* **562**, 217–222 (2018).

35. Matreyek, K. A. *et al.* Multiplex assessment of protein variant abundance by massively parallel sequencing. *Nat. Genet.* **50**, 874–882 (2018).

36. Penn, W. D. *et al.* Probing biophysical sequence constraints within the transmembrane domains of rhodopsin by deep mutational scanning. *Sci Adv* **6**, eaay7505 (2020).

37. Majithia, A. R. *et al.* Prospective functional classification of all possible missense variants in PPARG. *Nat. Genet.* **48**, 1570–1575 (2016).

38. Hampel, H. & de la Chapelle, A. The search for unaffected individuals with Lynch syndrome: do the ends justify the means? *Cancer Prev. Res.* **4**, 1–5 (2011).

Methods

General

Except as noted, cell culture media and supplements were obtained from Gibco, PCR reactions used Kapa HiFi Polymerase kit (Kapa Systems), nucleic acid cleanup kits were from Zymo Research, chemicals were from Sigma, cloning enzymes were from NEB, and oligonucleotides were synthesized by IDT or Eurofins. Primer oligonucleotide sequences are listed in Supplementary Table 3 and PCR cycling programs are listed in Supplementary Table 4.

Cell culture

Human HAP1 cells (Horizon Discovery) were cultured in IMDM; HEK293 and 293T/17 cells (ATCC) were cultured in DMEM, each supplemented with 10% fetal bovine serum, 100 U/ml penicillin and 100 g/ml streptomycin. During routine passaging, cells were washed with PBS pH 7.4 and dissociated with TrypLE Express. Cell lines were tested monthly for mycoplasma; all were negative.

MSH2 knockout cell lines

Clonal knockout lines were derived by transient transfection with plasmids expressing SpCas9 and guide RNAs targeting *MSH2*. For HAP1, a single guide sequence (primers JBW0001/2) targeting *MSH2* exon 6 was cloned into pSpCas9(BB)-2A-GFP (from Feng Zheng lab, Addgene #48138) as described[39]. For HEK293, four guides (primers VC0019/20, VC0021/22, VC0023/24, VC0027/28), two flanking each side of the *MSH2* locus, were each likewise cloned into pSpCas9(BB)-2A-PuroV2 (Feng Zheng lab, Addgene #62988) and used in combination. Wild-type HAP1 or HEK293 cells were transfected with the respective plasmid(s) using Lipofectamine 3000 (Thermo). Clonal HAP1 *MSH2* knockout cells were isolated by sorting single GFP positive cells using a FACS Aria II instrument (BD), and colonies were screened for *MSH2* exon 6 frameshift mutations. Clonal HEK293 *MSH2* knockout cells were isolated by dilution, expanded, and screened by PCR for the presence of a full-gene knockout deletion junction(s), and the absence of an internal target in exon 12 (Supplementary Fig. 12).

MSH2 cloning

Wild-type *MSH2* cDNA was PCR amplified from a lentiORF plasmid (Dharmacon OHS5897), and fused in-frame by HiFi Assembly to a blasticidin resistance marker amplified from pLentiCas9Blast (Feng Zheng lab, AddGene #52962), separated by a P2A self-cleaving peptide linker. The p.Ala636Pro mutation was introduced by site-directed mutagenesis. Each resulting construct was subcloned the doxycycline-inducible lentiviral expression plasmid pCW57.1 (from David Root lab, Addgene #41393) by digestion with SalI and NheI and ligation with T4 DNA ligase.

Western blotting

Cells were seeded on 6-well plates, grown to confluence, and lysed by the addition of RIPA buffer and protease inhibitor cocktail (Sigma-Aldrich). Protein extracts were run for 30 min at 200 V on denaturing Bolt 4-12% Bis-Tris Plus Gels in 1× Bolt MOPS SDS buffer (Invitrogen), transferred to a nitrocellulose membrane for 1.5 h at 10 V, and blocked at 4˚C overnight in 1X TBST (20 mM Tris base, 0.15 M NaCl, 0.1% Tween-20, 2% milk). Blots were probed with a 1:2000 dilution of anti-MSH2 (FE11, Thermo) and as loading control, 1:5000 dilution of anti-beta actin (BA3R, Invitrogen), followed by a 1:1000 dilution of HRP-conjugated goat anti-mouse IgG (H+L) secondary antibody (#62-6520, Invitrogen). Detection was performed with SuperSignal West Pico Chemiluminescent substrate (Thermo) and ChemiDoc XRS system (BioRad).

Library mutagenesis

MSH2 cDNA was divided into 21 tiles (Supplementary Table 5), and each was subjected to programmed allelic series mutagenesis[13] modified as follows. Mutagenic PCR reactions used PrimeStar GXL polymerase (Takara), and 45mer mutagenesis primers (Supplementary Table 3) were individually synthesized. Each codon was randomized in an individual PCR (program MUT_1), with a common

forward primer (JOK_0184) upstream of *MSH2,* and a codon-specific degenerate (NNN) reverse primer.
Resulting partial-length mutant cDNA was subjected to 10 cycles of strand extension (program MUT_2)
with limiting template (250 pg) of wild-type cDNA plasmid. Finally, each full-length mutant cDNA was
PCR amplified with flanking outer primers (program MUT_3, primers JOK_0184 and JOK_0185). Per-
codon products were pooled within each tile and purified by SPRI bead cleanup[40]. PCR products digested
with DpnI, to deplete starting plasmid, and NheI-HF and SalI-HF, for cloning into the lentiviral vector as
described above. Ligation products were transformed into Endura electrocompetent *E. coli* (Lucigen). A
small aliquot of the transformant culture was plated to monitor library complexity, while the remainder
was expanded for 12-16 hours at 32ºC in Luria Broth with 100 ug/ml ampicillin (LB+amp). We required
total colony forming units $\geq 4 \times 10^4$, for mean $\geq$14-fold coverage per variant. Lentiviral tile plasmid libraries
were isolated using ZymoPURE II Plasmid kit, and converted to shotgun libraries for deep sequencing
by Tn5 tagmentation[41], to assess mutational representation and uniformity (Supplementary Fig. 9).
Absent or under-represented codons were individually retrieved from the earlier mutagenesis steps,
pooled, cloned as above, and spiked into each tile library as needed to equalize coverage. Finally, these
mutant tile libraries, as well as control clones (WT and A636P), were modified to add trackable barcodes.
Each clone or library was linearized with SalI-HF (NEB) downstream of the *MSH2*-2A-bl^R ORF, and a
degenerate 20-mer (primer jklab0172) was inserted by HiFi Assembly (NEB). Resulting barcoded clone
pools were transformed, expanded, and isolated as described above.
339
Lentiviral preparation and transduction
Barcoded *MSH2*-2A-bl^R cDNA libraries were packaged into lentivirus by co-transfecting HEK293T/17
cells (ATCC) with the transfer plasmid pool plus envelope and packaging vectors pMD2.G and psPAX2
(from Didier Trono's lab, Addgene #12259 and #12260). For each pool, 4.4×10^6 cells were plated in a
100 mm dish, then transfected with 17 ug total plasmid (2.0:1.0:1.3 transfer:envelope:packaging ratio),
using Lipofectamine 3000 (Thermo). Media was replaced at 6 h post transfection, and viral supernatants
were collected at 24 h, passed through an 0.45 micron filter (Millipore), and used immediately or stored
at -80ºC until use. Viral titer was estimated by transduction with a dilution series followed by puromycin
selection and CellTiterGlo (Promega) cell proliferation assay[42], and verified by counting unique barcodes
in the transduced population (barc-seq, described below). For each tile, *MSH2* knockout HAP1 or
HEK293 cells were transduced with mutant library at low multiplicity of infection (< 0.1), by applying 1.5
ml viral supernatant (with 8 µg mL^{-1} polybrene) to each of four 100mm dishes (Corning) containing
$\sim 7.5 \times 10^6$ cells. After 48 h, transduced cells were selected by addition of 1 µg mL^{-1} puromycin, which was
included in all subsequent steps.
354
Mismatch repair functional selection
Cells expressing loss-of-function *MSH2* variants were enriched by adding 6-thioguanine (6-TG) to the
culture media. To find an optimal concentration, parental and *MSH2* KO cell lines were plated in
quadruplicate wells of a 96-well plate and treated with a range of 6-TG concentrations. Cell growth after
4 days was measured using the CellTiterGlo 2 kit (Promega) and a GloMax luminometer (Promega), and
survival was normalized to untreated wells (Supplementary Fig. 2).
361
For library selections, each mutant tile cell population was spiked with barcoded knock-in control cells
(*MSH2* WT: to 10% of the population, A636P: to 0.5%), then split and expanded in triplicate 10 cm dishes
(Supplementary Fig. 6) in media supplemented with 1 µg mL^{-1} doxycycline to induce *MSH2* expression.
From each baseline passage ('P0') culture, 10^7 cells were collected for gDNA extraction, and cells were
split into 6-TG treated and mock-treated cultures, with media containing 1 µg mL^{-1} puromycin, 1 µg mL^{-1}
doxycycline, 5 µg mL^{-1} blasticidin (to enforce *MSH2* expression), with and without 1 µM 6-TG,
respectively. Cells were grown to confluence for two rounds, at each round harvesting $\geq 10^7$ cells for
gDNA extraction and seeding with 3×10^6 cells to avoid population bottlenecks.
370
Amplification and sequencing of integrated *MSH2*
After selections, genomic DNA was isolated from $1-3 \times 10^7$ cells using the Quick-DNA Midiprep Plus kit.
To provide sufficiently redundant sampling of integrated *MSH2* libraries, for each sample, 16 replicate
PCR reactions were assembled, each with 0.5 µg of gDNA as template (template copy number of 2.4×10^6

genome equivalents per tile). Integrated inserts (~3.7 kbp), including *MSH2* cDNA and associated
barcode, were PCR amplified with PrimeSTAR GXL (program MSH2gDNA). Each set of 16 replicate
amplicons was pooled and purified with SPRI beads. Next, tileseq libraries were prepared from each pool
of amplified integrated *MSH2* constructs. Tile-specific primers (Supplementary Table 3) were used to
further amplify the tile mutagenized in each sample in a PCR reaction (program TILESEQ) with 10 ng
full-length amplicon ($\geq 2.6 \times 10^9$ copies) as template. Dual-indexed illumina adapters (sequences available
upon request) were then added by a subsequent round of PCR. The resulting short amplicon libraries
thus provide overlapping read coverage for most of each tile. Tile-seq libraries were pooled for paired
end 150bp Illumina sequencing ($\geq 1 \times 10^6$ paired reads per sample, n=189 libraries). Library complexity
(the number of distinct transduced clones) was assessed by barc-seq. Barcodes were amplified from the
full-length *MSH2* amplicon and converted to illumina amplicon libraries, as for tileseq, but using primers
jklab0077 and jklab0078 (program BCSEQ) . Each tile's starting population ('P0') had ≥20,000 barcodes,
indicating at least that number of unique initial transduction events (Supplementary Fig. 7). Catalogs of
barcodes associated with each of the two *MSH2* control clones (WT, A636P), and for each tile library,
were established by barc-seq (as described above) from their respective gDNAs at P0 (after
transduction). Finally, tile-seq libraries were prepared from a fully wild-type *MSH2* plasmid DNA, to
measure and correct by position-specific error rates, primarily single base substitutions which varied
considerably by position and type (Supplementary Fig. 8)
393
Sequence data processing
Shotgun sequencing reads were aligned to a set of *MSH2* cDNA sequences with each codon sequentially
replaced with 'NNN' using bbmap (https://jgi.doe.gov/data-and-tools/bbtools/) in semi-perfect mode, to
identify reads containing at most one mutated codon with between one and three edits. Custom python
scripts converted the resulting alignments to a per-sample table of counts for every possible codon
mutation. For tile-seq libraries, paired reads were overlapped and error-corrected with PEAR v0.9.6 (ref.
[43]); only perfectly matching pairs were used. For each tile, all possible single-codon mutation sequences
were enumerated, and the count of reads exactly matching each mutant sequence was tabulated. Reads
with additional mutations or sequencing errors were discarded. Barc-seq reads were clustered with
starcode v1.3 (ref. [44]) to define and count unique barcodes. To determine the number of distinct barcodes
present, robust to sequencing or amplification artifacts, we ranked clustered barcode groups by read
count and determined the minimum number of barcodes for which all equally or more abundant barcodes
cumulatively accounted for ≥90% of the total read counts.
407
408
Variant enrichment scoring
The raw frequency of each per-codon mutation *i* in each sample *s* was calculated, with a pseudocount of
0.9: $f'_{i,s} = \frac{(c_{i,s}+0.9)}{\sum_k (c_{k,s}+0.9)}$. Position specific error rates from wild-type *MSH2* plasmid tile-seq were subtracted
and normalized to yield corrected frequencies: $f_{i,s} = \frac{max\,(f'_{i,s}-f'_{i,WT},0)}{\sum_k max\,(f'_{k,s}-f'_{k,WT},0)}$. Variants with frequency
≤1/200,000 in the initial transduced cell population (P0) were removed from further analysis. Next,
variants substantially enriched or depleted after mock treatment (absolute $\log_2$ ratio > 2, after two
passages blasticidin vs P0) were likewise removed; these variants may be *in cis* with a frameshift or
premature truncating mutation beyond the sequenced tile. A loss-of-function (LOF) score was taken as
the $\log_2$ratio of the frequency after 6-TG treatment over that after mock treatment. For amino acid
substitutions represented by more than one equivalent codon substitution, the median was taken across
those codons' LOF scores to yield an amino acid-level LOF score. Within each replicate, variants were
removed when represented by exactly two LOF scores with opposite signs. Finally, the median score
was taken for each amino acid LOF score across the three replicates; when only two replicate scores
remained and had discordant signs, they were removed. False discovery rate was estimated by random
sampling of synonymous variants' read counts, with replacement and keeping triplicates together. To
mimic the number of codons corresponding to a given mutant amino acid, synonymous variants were
sampled in groups of size one to six. Counts were processed to LOF scores and filtered as described
above, combining first within each simulated missense amino acid and then across replicates; final LOF

scores greater than zero were taken as false discoveries. Overall FDR was estimated as the mean FDR
for amino acids with each number of different variant codons (1, 2, 3, 4, or 6 codons), weighted by the
fraction of variant amino acids with that number of codons.
430
Clinical and population variants
MSH2 missense variants and classifications were obtained from ClinVar
(https://www.ncbi.nlm.nih.gov/clinvar/) on April 14, 2020. Redundant variants were removed, and likely
splice-disruptive variants with a SpliceAI[23] score of ≥0.2 were filtered out, along with those where the
ClinVar record supported a splicing disruptive mechanism (Supplementary Table 2). Only variants with
expert panel reviews were included in comparisons to LOF scores. *MSH2* missense variants and
classifications were accessed from InSiGHT (https://www.insight-group.org/variants/databases/). *MSH2*
missense population variation was obtained from the gnomAD database[24], combining whole-genome and
exome calls from versions 2.1.1 and 3.0. The expected number of LOF missense mutations was modeled
with a binomial distribution parameterized by n=744 (the number of scored missense variants in gnomAD)
and p taken as the sum of LOF missense point variants' relative likelihoods of occurring *de novo*[45],
equivalent to drawing sets of size 744 from the universe of possible missense SNVs with probabilities
scaled by their individual relative likelihoods.
444
445   **Bioinformatic classification**
446   All *MSH2* missense variants were scored by the following bioinformatic tools: CADD[46], PolyPhen-2[47],
447   REVEL[48], PON-MMR[49], MAPP-MMR[50], and FoldX (scores obtained from ref.[32]). For CADD and REVEL,
448   scores were available only for missense variants reachable by single-base variants (SNVs); for amino-
449   acid substitutions that could arise from more than one SNV, the mean of those SNVs' scores was taken.
450
MSH2 structure and conservation
The MSH2 crystal structure[27] was obtained from the Protein Databank (accession 2O8E) and rendered
with PyMOL. Amino acid secondary structure assignment was extracted from pre-calculated DSSP
files[51]. Surface accessibility was calculated using an online tool (http://cib.cf.ocha.ac.jp/bitool/ASA/).
MSH2 ortholog protein sequences were downloaded from Ensembl (https://useast.ensembl.org/) and
aligned with MUSCLE[52].
457
458   **Data availability**
459   Processed count and final LOF score data are available from NCBI Gene Expression Omnibus (GEO
460   accession XXXXX), and corresponding raw sequencing data are posted to SRA (accession XXXXX).
461

References

39.   Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**, 2281–2308 (2013).
40.   Rohland, N. & Reich, D. Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Res.* **22**, 939–946 (2012).
41.   Picelli, S. *et al.* Tn5 transposase and tagmentation procedures for massively scaled sequencing projects. *Genome Res.* **24**, 2033–2040 (2014).
42.   Joung, J. *et al.* Genome-scale CRISPR-Cas9 knockout and transcriptional activation screening. *Nat. Protoc.* **12**, 828–863 (2017).
43.   Zhang, J., Kobert, K., Flouri, T. & Stamatakis, A. PEAR: a fast and accurate Illumina Paired-End reAd mergeR. *Bioinformatics* **30**, 614–620 (2014).
44.   Zorita, E., Cuscó, P. & Filion, G. J. Starcode: sequence clustering based on all-pairs search. *Bioinformatics* **31**, 1913–1919 (2015).
45.   Carlson, J. *et al.* Extremely rare variants reveal patterns of germline mutation rate heterogeneity in humans. *Nat. Commun.* **9**, 3753 (2018).
46.   Rentzsch, P., Witten, D., Cooper, G. M., Shendure, J. & Kircher, M. CADD: predicting the deleteriousness of variants throughout the human genome. *Nucleic Acids Res.* **47**, D886–D894

(2019).

47. Adzhubei, I. A. *et al.* A method and server for predicting damaging missense mutations. *Nat. Methods* **7**, 248–249 (2010).

48. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am. J. Hum. Genet.* **99**, 877–885 (2016).

49. Ali, H., Olatubosun, A. & Vihinen, M. Classification of mismatch repair gene missense variants with PON-MMR. *Hum. Mutat.* **33**, 642–650 (2012).

50. Chao, E. *et al.* Accurate classification of missense variants in MLH1/MSH2 with MAPP-MMR. *Cancer Res.* **68**, LB–14–LB–14 (2008).

51. Touw, W. G. *et al.* A series of PDB-related databanks for everyday needs. *Nucleic Acids Research* vol. 43 D364–D368 (2015).

52. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).