Non-dietary Fructose Metabolism Contributes to the Warburg effect in Cancers

Bing Han[1]*, Lu Wang[1]*, Meilin Wei[1], Cynthia Rajani[1], Runming Wei[1], Jingye Wang[1], Haining Yang[1], Michele Carbone[1], Guoxiang Xie[1,2], Wei Jia[1,3,4]

Abstract:

Fructose metabolism is increasingly recognized as a preferred energy source for cancer cell proliferation. However, dietary fructose rarely enters the bloodstream. Therefore, it remains unclear how cancer cells acquire a sufficient amount of fructose to supplement their energy needs. Here we report that the cancer cells can convert glucose into fructose through intra- and extracellular polyol pathways. The fructose metabolism bypasses normal aerobic respiration's self-control to supply excessive metabolites to glycolysis and causes the Warburg effect. Inhibition of fructose production drastically suppressed glycolysis and ATP production in cancers. Furthermore, we determined that a glucose transporter, SLC2A8/GLUT8, exports intracellular fructose to other cells in the tumor microenvironment. Taken together, our study identified overlooked fructose resources for cancer cells as an essential part of their metabolic reprogramming and caused the Warburg effect.

Declaration of Interests
The authors declare no competing interests.

---

[1] Cancer Biology Program, University of Hawaii Cancer Center, Honolulu, HI, 96813, USA.
[2] Human Metabolomics Institute, Inc., Shenzhen, Guangdong, 518109, China.
[3] Shanghai Key Laboratory of Diabetes Mellitus and Center for Translational Medicine, Shanghai Jiao Tong University Affiliated Sixth People's Hospital, Shanghai, 200233, China.
[4] School of Chinese Medicine, Hong Kong Baptist University, Kowloon Tong, Hong Kong.
*Equal contributions.
Correspondence: WJia@cc.hawaii.edu.

Statement of Significance:

Our findings in this study suggest that the Warburg effect is actually achieved by means of fructose metabolism, instead of glucose metabolism alone. Fructose metabolism results in accelerated glycolysis and an increased amount of ATP and key intermediates for anabolic metabolism.

Introduction

Cancer cells consume excessive amounts of glucose that can be obtained from cancer-mediated metabolic reprogramming to sustain their unrestricted proliferation [1]. We previously reported that cancer cells such as acute myeloid leukemia cells exhibited enhanced fructose utilization under low glucose conditions and that pharmacologic blockade of fructose utilization suppressed AML cell proliferation and enhanced the therapeutic efficacy of Ara-C [2]. Our recent findings, along with reports from other groups [1-3], suggest that fructose is a preferred carbon source for cancer cells. We found that high expression of the fructose transporter SLC2A5/GLUT5 was found in multiple types of cancers, enabling the increased uptake of fructose from the environment and enhancement of cancer growth and development. However, dietary fructose is mostly converted to glucose, glycogen, and organic acids in the intestine and liver [4]. Therefore, the fasting blood levels of fructose are about 1,000-fold lower (~0.005 mM) than glucose (~5.5 mM) under normal physiological conditions. Therefore, we hypothesized that this unique metabolic feature in cancer cells must arise from additional fructose sources and increased advantages of utilizing fructose over glucose by cancer cells. The polyol pathway is commonly present in somatic cells. It converts about 3% of intracellular glucose to fructose under normal physiological conditions in a two-step process [5]. The first step involves reducing glucose to sorbitol utilizing the reducing equivalents from NADPH and the aldose reductase enzyme, AKR1B1. The second step is the oxidation of sorbitol to fructose via sorbitol dehydrogenase (SORD) and the conversion of NAD+ to NADH.

Dr. Otto Heinrich Warburg first observed the Warburg effect, which described a phenomenon that the cancer cells tend to ferment glucose to lactic acid even in aerobic conditions, so-called "aerobic glycolysis." The current understanding of this phenomenon is that glucose fermentation promotes biomass synthesis in cancers to maintain cell proliferation [6]. And

this phenomenon is also found in non-cancer high-proliferating cells, such as T lymphocytes [7]. But the metabolic mechanism of the Warburg effect is still unknown. This study revealed that non-dietary sources of fructose from both intra- and extracellular polyol pathways supported cancer metabolism and contributed significantly to the Warburg effect of cancers.

Results

Non-dietary sources are significant suppliers of fructose to cancers

This study revealed overlooked sources of fructose produced by both intra- and extracellular polyol pathways, which were upregulated due to cancer-mediated metabolic reprogramming. First, we identified significant levels of ^{13}C-fructose in 11 out of 12 cancer cell lines that were incubated with ^{13}C-glucose (Fig. 1A). Basal levels of fructose were also detected in the two control cell lines, the pancreatic epithelial cell, HPDE6C7, and the mammary epithelial cell line, MCF-10A. We then tested the fructose-conversion rate in the A549 cell line. The results showed that A549 cells converted glucose to fructose rapidly and that their intracellular ^{13}C-fructose level reached a saturation point after 2 hours of treatment with 25 mM ^{13}C-glucose. We also found that the intracellular ^{13}C-fructose level was equal to intracellular ^{13}C-glucose after 2 hours (Fig.1B). Therefore, we estimated that about 50% of the glucose taken up by A549 cells was converted to fructose.

In addition to intracellular production of fructose, we unexpectedly found that in two common commercial brands of FBS, there was approximately double the amount of fructose relative to glucose (Fig. 1C). This phenomenon could be due to either fetal bovine serum naturally having higher fructose levels, or it could be due conversion of FBS glucose into fructose slowly after harvest. To identify the reason for high FBS fructose, we selected regular,

heat-inactivated (HI), active charcoal striped, and dialyzed FBSs purchased from two major commercial vendors (VWR and Gibson) for this study. We adjusted every FBS sample to 25mM of ^{13}C-glucose and cultured them at 37°C 5% CO2 for 0, 8, 24, and 48 hours. The ^{13}C-fructose levels increased with culture time in all the FBS samples and different conversion rates were observed depending on the type of FBS sample (Fig. 1D). This result revealed a novel source of fructose in blood: a mechanism that naturally converts glucose to fructose.

The polyol pathway is the only known physiological reaction to convert glucose to fructose intracellularly. We analyzed two enzymes involved in the polyol pathway, AKR1B1 and SORD, at both gene (Fig. S1A) and protein (Fig. S1B) levels and found that all cancer and control cell lines expressed varying levels of these two genes. We then examined the human TCGA pan-cancer database (Fig. S1C), and confirmed that all human cancers express AKR1B1 and SORD, which suggested that the polyol pathway was active as an essential metabolic feature in all types of cancer. To identify whether the polyol pathway also processes extracellular glucose-fructose conversion, we then assayed the AKR1B1 and SORD proteins in FBS by Western Blot (Fig. S1D). Both of these two components were found in all tested samples with varying amounts. We then blocked the intracellular and extracellular polyol pathway by AKR1B1 gene knocked out (KO) (Fig. S1E) or epalrestat [8], an AKR1B1 inhibitor, respectively, to verify the role of the polyol pathway in glucose-fructose conversion. Inhibition of of AKR1B1 by either method effectively decreased the intracellular (Fig. 1E) and extracellular (Fig. 1F) fructose concentrations. These results provide evidence for the activity of the polyol pathway in cancer cells and serum.

As a result of AKR1B1 gene KO in cancer cells, the fructose specific metabolite, fructose-1-phosphate (F-1-P), was significantly decreased as shown in Fig. 1G and Fig.S1F. Inhibition of

the polyol pathway did not have the same effect on the production of the two glucose-specific metabolites, glucose-6-phosphate (G-6-P) and fructose-6-phosphate (F-6-P), in the four cancer cell lines (Fig. 1G and Fig. S1G). Therefore, AKR1B1 KO blocked [13]C-fructose production from the polyol pathway but did not alter glucose metabolism. Thus, it appeared that once fructose is produced in cancer cells, its subsequent metabolism is independent of glucose metabolism. Fructose-specific metabolism is different from glucose metabolism in that it has fewer steps, and the pathway is not reversible [9]. Cancer cells commonly overexpress glucose transporters, such as SLC2A1/GLUT1 and SLC2A3/GLUT3, to take in a large amount of extracellular glucose to support their high rates of proliferation [10]. Glucose is known to be the preferred carbon source for the production of structural components necessary for cell growth and division. When cancer cells metabolize glucose to synthesize biomass, a massive amount of NADPH is generated as well [6]. Hence high glucose and high NAPDH levels are commonly found in cancer cells. Interestingly, in this study, we found that the accumulation of intracellular glucose and NADPH in cancer cells facilitated the activation of the polyol pathway (Fig. 1G).

To determine whether the glucose level affects the activation of the polyol pathway, we compared the expressions of AKR1B1 and SORD in cancer cells cultured under different glucose levels. We found that glucose availability regulated the polyol pathway. We selected four different types of cancer cells with varying fructose production capabilities for this study. Under high glucose (25 mM) concentrations, the cancer cells significantly increased their AKR1B1 expression levels compared to those grown in low glucose (2.5 mM) (Fig.1H). The response of SORD expression to different glucose concentrations varied in the four cancer cell lines (Fig.S1H). This observation may also provide a mechanistic explanation for the much-increased cancer risk in the diabetic patient population [11]. Increased blood glucose activates the

polyol pathway, not only causing tissue damage due to increased production of sorbitol but also providing damaged cells with additional nutrient support to facilitate their malignant transformation.

Fig.1. Cancer cells produce high concentrations of fructose

(**A**) HPDE6C7 and MCF-10A cells are considered normal cells controls. Along with these controls, we tested a wide array of different cancer cell lines using ^{13}C labeled metabolic flux analysis. From these data, it was seen that although most of the cells affected the conversion of glucose to fructose, there was variability in how much fructose was produced. Notably, A549 cells produced the most fructose. (**B**) Based on the results in Fig. 1A, we tested the rate of fructose production in A549 cells. The red line represents intracellular ^{13}C-fructose production, and the blue line represents the intracellular ^{13}C-glucose level. Data are represented as mean ± SEM. (**C**) The concentration of glucose and fructose in two major commercial FBSs. (**D**) The ^{13}C-fructose production rate in 6 different FBSs. (**E**) AKR1B1 KO blocked fructose production in tested cancer cells. (**F**) AKR1B1 inhibitor, epalrestat, suppressed the ^{13}C-fructose transformation in FBS. (**G**) Sketch of the polyol pathway and fructose metabolism. In A549 cells, fructose specific metabolite, fructose-1-phosphate (F-1-P), highly depended on the polyol pathway. Only a trace amount of F-1-P can be detected in AKR1B1 KO cancer cells, while the compromised polyol pathway did not significantly affect glucose metabolism. The amounts of glucose specific metabolites, glucose-6-phosphate and fructose-6-phosphate (G-6-P and F-6-P), had no significant changes in A549 AKR1B1 KO cancer cells. (**H**) We tested the regulation of expressions of AKR1B1 according to glucose level in different cancer cells. The AKR1B1 expression was promoted significantly by high glucose level (25mM) in all tested cell lines. Data are represented as mean ± SEM. *: t-test $p < 0.05$; **: t-test $p < 0.01$.

Fructose metabolism enhances glycolysis in cancer cells

To investigate the role of fructose metabolism in cancer cells, we compared the AKR1B1 KO (the polyol pathway deficient) A549 cells with their wild-type counterparts (control) in the Seahorse Cell Glycolysis Stress test and Mito Stress test assays [12]. The assays showed drastically compromised glycolysis in the AKR1B1 KO cells (Fig. 2A) but no significant difference in mitochondrial functions (Fig. 2B) between the two cell lines. We then examined metabolites involved in glycolysis and mitochondrial respiration. Targeted metabolomics profiling showed that the glycolytic metabolites, such as serine [13], pyruvate, and lactate, were significantly decreased in AKR1B1 KO cells (Fig. 2C-E and Fig S2A). Meanwhile, metabolites related to aerobic respiration, such as succinate, fumarate, malate, and aspartate [14] were slightly decreased or not significantly altered (Fig. 2F-I and Fig.S2B). These results indicated that fructose

metabolism promoted glycolysis, leading to enhanced amino-acid (Fig. 2C, F and Fig. S2A, C) and pyruvate (Fig. 2D) production. Pyruvate was first provided to the TCA cycle in mitochondria (Fig. 2G-I), and after TCA metabolism reached the saturation point, the excess amount of pyruvate was converted to lactate (Fig. 2E). We observed similar metabolic phenotypes in AKR1B1 silenced CY790 cells (Fig S3A-F), showing high fructose production close to that of A549 cells. Metabolic flux analysis showed that wild-type cells had a significantly faster pyruvate-producing rate than the polyol pathway-deficient cells (Fig. 2J). The malate production curve showed a similar trend to pyruvate production (Fig. 2K), suggesting that the polyol pathway accelerated pyruvate production and rapidly fueled the TCA cycle. The lactate production increased at a much slower rate than those of pyruvate and malate in the wild-type cells (Fig. 2L), suggesting that after the TCA-cycle reached its saturation point, additional pyruvate derived from fructose metabolism was then converted to lactate. Such metabolic outcome is the phenomenon that Dr. Otto Warburg observed about 100 years ago.

To evaluate the contribution of fructose metabolism to the Warburg effect, we knocked out or silencing the Hexokinase-1 (HK1) and Hexokinase-2 (HK2) genes from A549 cells (Fig. S3J and K). The hexokinases phosphorylate glucose to G-6-P (Fig. 1G) and start the glycolysis. Both the hexokinases HK1 and HK2 are expressed in cancers. Knock out of HK1 and HK2 effectively blocked glucose metabolism in A549 cells[15]. As described above, the A549 cells could process about 50% of glucose to fructose, i.e., the amounts of glucose and fructose are almost equal in this cell. Blockage of glucose and fructose metabolisms in this cell line should thus be comparable. Interestingly, the amount of ATP was drastically depleted in both glucose metabolism deficient and fructose metabolism deficient cells at similar levels (Fig. 2M and Fig. S3L). We then compared the extracellular acidification rates (ECAR), a symbolic index of the Warburg effect, in the AKR1B1 KO and HK KO or silencing cells. The data showed that glucose

or fructose metabolism blockage resulted in a similar level of acidification decrease in the A549 cancer cells (Fig. 2N and Fig. S3M).

These results implied both glucose and fructose were equally capable at fueling glycolysis. As we know, glucose metabolism is a self-limited process with negative regulation of PFK and HK by glycolysis products such as ATP and G-6-P[16]. To verify whether metabolites from glucose metabolism affected fructose conversion, we further tested the AKR1B1 and SORD expression in glucose metabolism deficient cells. The results showed that compared to control cells, AKR1B1 and SORD expressions did not significantly change in HK KO cells (Fig.2O and P, and Fig. S3N and O). These implied that fructose metabolism was relatively independent of glucose metabolism.

As a tightly controlled process, glucose metabolism will reach a dynamic balance to supply limited materials and energy for regular cell metabolism. But in highly proliferating cancer cells, it is not sufficient. Under such circumstances, the fructose metabolism will bypass these control elements of classic glycolysis and keep providing Glyceraldehyde 3-phosphate (GA3P) into the later glycolysis stage (Fig. 2Q). The additional GA3P from fructose metabolism will not significantly fuel the TCA cycle, which is almost saturated by classic glycolysis. Instead, the enhanced glycolysis by fructose metabolism will tend to increase the biomass synthesis and production of lactic acid.

Fig.2. Fructose metabolism enhances glycolysis in cancer cells
(**A**) The Seahorse Cell Glycolysis Stress test showed compromised glycolysis in AKR1B1 KO A549 cells. (**B**) The Mito Stress test showed no significant change in mitochondrial function between A549 wild-type control and AKR1B1 KO cells. The direct metabolites from glycolysis, serine (**C**), pyruvate (**D**), and lactate (**E**) were significantly decreased in AKR1B1 KO cells. In comparison, TCA cycle-related metabolites, such as succinate (**I**), fumarate (**H**), malate (**G**), and aspartate (**F**) were only slightly reduced in the KO cells. We also compared the flux of pyruvate (**J**), malate (**K**), and lactate (**L**) between wild-type control and AKR1B1 KO A549 cells when feeding with glucose after starving. The KO cells had slower rates in both of these metabolites. These changes would be expected to lead to diminished ATP production, as we observed (**M**) in AKR1B1 KO cells as well as HK KO cells. We then compared glucose and fructose metabolisms effects on glycolysis using the Seahorse glycolysis stress test (**N**). To evaluate whether glucose metabolism affects the polyol pathway, AKR1B1 and SORD expressions were tested in glucose metabolism compromised cells (**O** and **P**). (**Q**) Sketch of fructose metabolism contributions to the Warburg effect. Data are represented as mean ± SEM. *: t-test $p<0.05$; **: t-test $p<0.01$.

Cancer cells export endogenous fructose through the sugar transporter SLC2A8/GLUT8.

Cancer cells absorb extracellular fructose predominantly through the fructose transporter, SLC2A5/GLUT5. Interestingly, we detected [13]C-fructose in culture supernatants of all cell lines incubated with [13]C-glucose, except CY790, a mesothelioma cell line (Fig. 3A). This finding suggests that fructose produced in cancer cells can be transported to the extracellular environment. We compared the two mesothelioma cell lines, CY790 and CY289, one with a trace amount of fructose and the other with high concentration fructose (7 μM vs. 894 μM) found in the supernatant, respectively. The SLC2A5/GLUT5 and SLC2A8/GLUT8 expression levels were distinct between the two cell lines (Fig. 3B). We detected SLC2A8 expression in CY289 but not in CY790 cells, suggesting that the difference in SLC2A8 expression between the two cell lines may be responsible for the different fructose concentrations in supernatants. We then knocked out the expression of SLC2A8 in A549 cells using a CRISPR/Cas9 technique (Fig. 3C) as this cell line demonstrated strong fructose-exporting ability with the highest concentration of fructose in the culture supernatant among 14 cell lines (Fig. 3A). The A549 SLC2A8 knockout

cells showed significantly higher intracellular fructose levels and lower extracellular fructose levels compared with the A549 wild-type (Fig. 3D), indicating that fructose export was effectively shut down. SLC2A8 is widely expressed in almost all cancer cell lines, as shown in Fig. 3E, similar to the expression of the two genes encoding AKR1B1 and SORD in the polyol pathway. This finding was also validated by the TCGA human cancer database (Fig. 3F), which reported that SLC2A8 was expressed across the spectrum of all types of human cancer.

This novel finding led us to hypothesize that fructose-producing cancer cells can make this additional energy source available to other cells in the tumor microenvironment. Generally, the nutrient supply is compromised inside a fast-growing tumor mass. However, with the polyol pathway and SLC2A8-mediated fructose exchange, it may be possible that the cancer cells in the outermost layer of the tumor can access circulating glucose and convert it to fructose and then export it into the tumor microenvironment for utilization by other cancer cells in the inner layer of the tumor cell mass (Fig. 3G).

Fig. 3. Cancer cells export endogenous fructose through the sugar transporter SLC2A8/GLUT8

(A) We found that cancer cells that produced fructose can export it to the extracellular environment, and that fructose distribution is varied in different cancer cells. Some cells tended to keep fructose in the cytoplasm, and others exported more fructose to the culture medium. This may be due to differences in fructose transporter expression in different cancer cells. **(B)** We tested potential fructose transporter[17-19] expressions in two mesothelioma cell lines with different fructose distribution patterns and found that the expression of SLC2A8 was different between the

two cell lines. **(C)** We generated SLC2A8 KO A549 cells by CRISPR/Cas9 technology and compared them with the control sgRNA group. **(D)** In SLC2A8 KO A549 cells, fructose export was highly suppressed, and the extracellular/intracellular fructose ratio was significantly reduced compared with the control group, while the intracellular fructose/glucose ratio was increased considerably. **(E)** SLC2A8 was also found to be widely expressed in cancer cells and **(F)** in human cancers according to the TCGA database, including: A. acute myeloid leukemia; B. thymoma; C. mesothelioma; D. thyroid carcinoma; E. cholangiocarcinoma; F. pancreatic adenocarcinoma; G. lung adenocarcinoma; H. breast invasive carcinoma; I. esophageal carcinoma; J. kidney clear cell carcinoma; K. head & neck squamous cell carcinoma; L. ovarian serous cystadenocarcinoma; M. lung squamous cell carcinoma; N. cervical & endocervical cancer; O. skin cutaneous melanoma; P. kidney papillary cell carcinoma; Q. uterine carcinosarcoma; R. pheochromocytoma & paraganglioma; S. stomach adenocarcinoma; T. brain lower grade glioma; U. glioblastoma multiforme; V. sarcoma; W. uterine corpus endometrioid carcinoma; X. diffuse large B−cell lymphoma; Y. testicular germ cell tumor; Z. prostate adenocarcinoma; a. bladder urothelial carcinoma; b. uveal melanoma; c. colon adenocarcinoma; d. rectum adenocarcinoma; e. liver hepatocellular carcinoma; f. kidney chromophobe; g. adrenocortical cancer. **(G)** Sketch of the fructose communication between cancer cells. Data are represented as mean ± SEM. **: *t*-test p<0.01.

Fructose enhances cancer cell proliferation and malignancy

To determine how fructose metabolism benefits cancer cells, we measured the cell responses to fructose in AKR1B1 KO cells and also cells overexpressed with AKR1B1. The AKR1B1 KO cells showed a decrease in malignant phenotypes *in vitro* and *in vivo*, while cells overexpressed in AKR1B1, on the other hand, demonstrated significantly increased rates of growth and tumor development in a mouse model (Fig. 4, Fig. S4-5). To study the role of fructose metabolism via the polyol pathway in cancer cell growth, we generated AKR1B1 KO (Fig. 1E and S1E) and AKR1B1 overexpressed cancer cell lines (Fig. S4A, B) using CRISPR/Cas9 knockout and CRISPRa/Cas9-VPR overexpression technology. In the original AKR1B1 high expression cancer cells, such as A549 and CY790 cells, the CRISPRa/Cas9-VPR had a negligible effect on the AKR1B1 expression. In U87 and BxPC3 cells with moderate expression levels of AKR1B1, however, fructose production was increased by 3-5 fold with overexpression of AKR1B1 (Fig. S4B).

We tested the migration and invasion ability of AKR1B1 KO A549 cells or AKR1B1 silenced CY790 cells. The knockout or silencing of AKR1B1 resulted in compromised cell motion in both the wound healing assay (Fig. 4A and Fig. S5A) and transwell assay (Fig.4B and Fig. S5B), compared with wild-type control cells. *In vitro* cell proliferation tests showed that the growth rate of AKR1B1 KO or silenced cancer cells was highly suppressed (Figs. 4C, Fig. S4C, and Fig. S5C). On the other hand, overexpression of AKR1B1 not only increased fructose production but also significantly increased tumor growth rate (right panels of Figs. 4C and Fig. S4C). We then investigated the effect of the polyol pathway on cancer proliferation *in vivo*. We injected 0.1 x10^6 wild-type or gene manipulated (AKR1B1 KO, overexpression, or AKR1B1 silenced) cancer cells, which constitutively expressed luciferase, in nude mice at day 0, respectively. Luciferase activity of each tumor increased with tumor mass. Every three days, we measured the luminescence intensity of tumors by D-luciferin injection. Compared with wild-type cells, AKR1B1 compromised (KO or silenced) cancer cells showed significantly reduced tumor growth rates (Fig. 4D, Fig. S4D, S5D, and S5E) while AKR1B1 overexpression promoted cancer cell growth (Fig. 4E, and Fig. S4E). These results confirmed that fructose metabolism enhanced cancer malignancy.

Fig. 4. Fructose enhances cancer cell proliferation and malignancy (A) The A549 AKR1B1 KO cells showed suppressed cell migration in the wound healing assay and (B) transwell assay. (C) The AKR1B1 KO cells slowed down cancer cell growths, and overexpression of AKR1B1 caused increased growth rates of U87 cells *in vitro*. (D) An *in vivo* xenograft model also showed growth suppression when AKR1B1 KO A549 cells were used, (E) there was increased cancer cell proliferation when U87 cells were overexpressed in AKR1B1. Data are represented as

mean ± SEM. *: *t*-test p<0.05; **: *t*-test p<0.01.

Discussion

Our study identified the polyol pathway to be a crucial fructose-producing mechanism that is important for cancer-mediated metabolic reprogramming leading to enhanced glycolysis, cell proliferation, and malignancy. Compared with glucose, fructose metabolism results in accelerated glycolysis and an increased amount of ATP, demonstrating a typical glycolytic flux phenotype. We suspect that cancer metabolism involving the intra-/extracellular polyol pathway-derived fructose may constitute an essential part of the Warburg effect. Under the aerobic condition, the abundant ATP from the TCA cycle limits the glycolytic rate. Almost all of the glycolysis production will be exhausted by the TCA cycle with nothing remaining to be converted to significant amounts of lactate. In cancers, a large amount of glucose taken in by overexpressed GLUT1 and/or GLUT3 activates the polyol pathway and fructose metabolism, bypassing glycolysis limitations and producing excess glycolytic metabolites for biomass and lactate production. It results in the acidification of cancer cells under aerobic conditions. Additionally, there is a selective advantage for cancer cells to utilize fructose to support their rapid proliferation because fructose lacks cellular metabolic homeostasis machinery, unlike glucose, which is strictly regulated by either insulin or glucagon. As the polyol pathway is not cancer-specific (comparable levels of AKR1B1 and SORD were found in normal epithelial cell lines, Fig S1A and B), we hypothesize that it profoundly impacts all fast-growing cells, including activated immune cells, germinal cells, and stem cells.

We also found in this study that high glucose levels stimulated the expression of AKR1B1. Therefore, it is not difficult to understand that the polyol pathway is activated to lower the blood glucose levels in diabetic patients and, thus, implicated in diabetic complications. The intermediate molecule in the polyol pathway, sorbitol, cannot cross cell membranes and, when it

accumulates from increased cellular production, causes microvascular damage to the retina, kidney, and nerves [5]. Given the recent findings that increased fructose level promotes cancer malignancy, the polyol pathway's activation may be responsible for the increased risk of cancer development in the diabetic population.

As the polyol pathway is not essential to energy metabolism in normal cells via the glycolysis pathway, inhibition of AKR1B1 activity may represent a novel, effective, and safe strategy for cancer therapy and preventive cancer treatments for diabetic patients. Currently, several AKR1B1 inhibitors are undergoing clinical trials to treat diabetic complications [8,20]. These compounds may have the potential to be useful as direct treatments for cancer and preventive cancer treatments for diabetic patients.

To summarize, the cancer cells utilize the polyol pathway to produce fructose and gain more energy and biomass from fructose metabolism than the equivalent amount of glucose. Fructose metabolism results in increased lactate production, demonstrating a typical glycolytic phenotype. Cancer metabolism involving polyol pathway-derived fructose may contribute to the Warburg effect.

Methods

Cell Culture

A549 (lung carcinoma), EKVX (lung adenocarcinoma), SW-620 (colorectal adenocarcinoma), HCT116 (colorectal carcinoma), U251 (malignant glioblastoma), MCF7 (breast adenocarcinoma), and MDA-MB-231 (breast adenocarcinoma) cells were obtained from NCI-60 cell storage of the University of Hawaii Cancer Center. BxPC3 (pancreas adenocarcinoma), PSN1 (pancreas adenocarcinoma), and U87 MG (malignant glioblastoma) cells were ordered from Sigma-Aldrich. HPDE6c7 (pancreatic duct epithelial cell), and MCF-10A (breast

fibrocystic disease) cells were ordered from American Type Culture Collection (ATCC). CY289 and CY790 (mesothelioma) cells were gifts from Dr. Carbrone and Dr. Yang's lab. All cell lines were adapted to culture and maintained in DMEM (Thermo Fisher Scientific) with 10% FBS under 37°C and 5% CO_2.

Gene knock out and active by CRISPR/Cas9

We used Dharmacon Edit-R All-in-one Lentiviral and Edit-R CRISPRa Lentiviral systems to create gene knock out (KO) and gene overexpression cell lines. The sgRNA sequences are as follows: CGACCTGAAGCTGGACTACC (GSGH11935-247543511, for Human AKR1B1 KO, Dharmacon), GCCGCGGCGGCCTTCCCCAA (GSGH11887-247076986, for Human AKR1B1 overexpression, Dharmacon), GACAACTATACCTGCCAGGT (GSGH11935-247493945, for Human SLC2A8 KO, Dharmacon), GTTCAAGATGGCGCCCAGTG (GSGH11935-247518767, for Human HK1, Dharmacon), TAAGCGGTTCCGCAAGGAGA (CS-HCP001744-LvSG03-1-B, for Human HK2, GeneCopoeia). The shRNA sequences are as follows: GGTGGAGATGATCTTAAACAA (HSH005861-LVRU6GP, for Human AKR1B1, GeneCopoeia), CAATGCCTGCTACATGGAGGA (CS-HSH1530L-LVRU6GP-01, for Human HKs, GeneCopoeia). Lentivirus was packed using the Dharmacon Trans-Lentiviral ORF Packaging Kit with Calcium Phosphate (TLP5916), and cells were infected to create stable cell lines according to the manufacturer's instruction.

Western Blotting

Immunoblotting was performed as previously described [2]. Briefly, the transfer-ready membrane was blocked 1 hour in TBS-T containing 5% nonfat milk at room temperature, followed by incubation with primary antibody at 4°C overnight. The anti-AKR1B1 antibody (Abcam, ab62795), anti-SORD antibody (Abcam, ab185705), anti-SLC2A8 antibody (LifeSpan

BioSciences, LS-C757596), and anti-Vinculin antibody (Cell Signaling Technology, E1E9V) were used at 1:1000 dilution respectively. The secondary antibody was horseradish peroxidase-conjugated anti-rabbit antibody used at a 1:5,000 dilution. Vinculin was used as a protein loading control.

RT PCR

RNA extraction and reverse transcription were performed by TRIzol (Thermo Fisher Scientific) and iScript™ cDNA Synthesis Kit (Bio-Rad Laboratories) according to the manufacturer's instructions. PCR primer sequences were as follows. Primers for AKR1B1 PCR: TAC TCA GCT ACA ACA GGA ACT G, AGG CAA GAA ACA CAG GTA TAG G, probe: TTG TTG AGC TGT ACC TCC CAC AAG G; Primers for SORD PCR: ACT CCA GAG CCA AAA GAG C , CAT CCT CAG CAA GAC CTC AT, probe: AGG ATA GTT CTC CAG GCG CAA GTC; Primers for SLC2A8 PCR: GCC AAG TTC AAG GAC AGC A, ATG ACC ACA CCT GAC AAG AC, probe: ATG ATG AGA GCC GCC ACA GCT; Primers for ACTB PCR: GCG AGA AGA TGA CCC AGA T, CCA GTG GTA CGG CCA GA, probe: CCA TGT ACG TTG CTA TCC AGG CTG T; Primers for SLC2A5 PCR: CAA GCA TGG AGC AAC AGG AT, GAA GGA TGA CCC AAA GGC A, probe: AGC ATG AAG GAA GGG AGG CTG AC; Primers for SLC2A2 PCR: CTG GAA GAA GCA TAT CAG GAC T, GCT GAT GAA AAG TGC CAA GTG, probe: CTG AGA GCG GTT GGA GCA ATT TCA C; Primers for SLC2A7 PCR: CCG TCT CCA TGT TTC CTC TG, CCA CTT TGC TGA CTC CCA TC, probe: CCC ACG AGC AAT GAC CCC AAC A; Primers for SLC2A11 PCR: GTC AGC AGC AAT CCT GTT TG, GTA CAT GGG CTG GAT GTT CA, probe: AGC AGT CTT CCC AGC ATG ATC ATC TC. We performed real-time PCR with the iTaq™ Universal Probes Supermix (Bio-Rad Laboratories) by Lightcycle 480 (Roche Molecular Systems).

Seahorse Extracellular Flux Assays

Gene-modified cancer cells were maintained in serum-free HuMEC Ready Medium before the assay. The Seahorse XF Glycolysis Stress Test Kit and Cell Mito Stress Test Kit (Agilent) were used to measuring cell glycolysis and mitochondrial function, respectively, using the Agilent Seahorse XF96 analyzer [12] following the manufacturer's instructions.

Cell Proliferation Assay

All gene-modified cancer cells were first adapted to HuMEC Ready Medium (Thermo Fisher Scientific) and then plated at 1000 cells/well in HuMEC Ready Medium followed by the culture at 37°C under 5% CO_2. A cell proliferation assay was performed by using the Cell Counting Kit-8 (CCK-8) (Dojindo Molecular Technologies) after the indicated time point of culture. 10 μl of the CCK8 dye was added to each plate well and then incubated for 1 hour at 37°C under 5%CO_2. The optical density (OD) was then read at 450 nm directly using a microplate reader.

***In vitro* Cell Migration and Invasion Assays**

The ibidi Culture-Insert 4 Well in μ-Dish(ibidi, 80466) was used for the wound healing assay, and the Corning BioCoat Matrigel Invasion Chamber (Corning, 354480) was used for the transwell assay according to the manufacturer's instructions. Gene-modified cancer cells were seeded in HuMEC Ready Medium and cultured at 37°C under 5% CO_2. The images of wound healing data were analyzed by NIH ImageJ.

Xenograft Formation

Nude mice (4–6 weeks old) from Charles River Laboratory were injected right and left sides respectively with 200 μl Matrigel (Corning) slurry (prepared at a 1:1 ratio with 1X PBS) containing 0.1 million control or gene manipulated cancer cells that constitutively expressed luciferase (AMS Biotechnology Limited). Luciferase activity of each tumor increased with tumor mass by the time. Every three days, we measured the luminance intensity of tumors by intraperitoneal injection of 50 mg/mouse D-luciferin (Thermo Fisher Scientific) into anesthetized mice, followed by the detection of live images using the Xenogen IVIS (PerkinElmer) at 30 min post-injection.

Chemicals and reagents

Diisopropylethylamine (DIPEA), 3-nitrophenylhydrazine (3-NPH)·HCl, 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) HCl, and pyridine (HPLC grade), glucose, fructose, glucose 6-phosphate, fructose 6-phosphate, glucose 1-phosphate, fructose 1-phosphate, fructose 1,6-bisphosphate, glyceraldehyde 3-phosphate, dihydroxyacetone phosphate, 2-phosphoglycerate, pyruvate, 6-phosphogluconate, ribulose 5-phosphate, ribose 5-phosphate, sedoheptulose 7-phosphate, erythrose 4-phosphate, citrate, isocitrate, succinate, fumarate, malate, lactate, α-ketoglutarate, oxaloacetate, serine, glutamate, glutamine, aspartate, hydropyruvate were purchased from Sigma-Aldrich (St Louis, MO). Optima LC-MS grade water, methanol (MeOH), isopropanol, formic acid and acetonitrile (ACN) were purchased from Thermo Fisher Scientific (Waltham, MA). [13C6]-glucose and [13C6]-fructose D-Glucose U-13C6 and D-Fructose U-13C6 were purchased from Cambridge Isotope Laboratories (Tewksbury, MA). FBS: Premium Grade Fetal Bovine Serum (VWR, 97068-085), Performance Plus Fe tal Bovine Serum (gibco, 16000-036), Charcoal Stripped Fetal Bovine Serum (gibco, A33821-01), Dialyzed Fetal Bovine Serum(gbico, a33820-01). Heat Inactive was performed

under 56 °C for 30 min. DMEM (gibco, 11966-025). Epalrestat (Cayman Chemical, 15214). 13C-Glucose (Cambridge Isotope Laboratories, CLM-1396). Amino Acids: L-Arginine, Hydrochloride (MilliporeSigma, 181003), Histidine hydrochloride (MilliporeSigma, H0755000), L-Lysine (MilliporeSigma, 62840), L-Phenylalanine (MilliporeSigma, P5482), L-Valine (MilliporeSigma, V0513). PBS (gibco, 14190-144).

Standard solutions

Each standard compound was accurately weighed and dissolved in water to prepare stock solutions with a final concentration of 20 mM for each standard. The final working standard solution mix contained 28 metabolites (detailed in Table 1) with a final concentration of 400 µM/metabolite. For quantification, a 7-point calibration curve was constructed by using a series of 1:2 serial dilutions from the highest concentration of calibration mixture with water. The stock solutions were kept at -20 °C.

Stable isotope tracer analysis and metabolite quantification

Cell culture, labeling, and sample collection: All tested cells were cultured in DMEM containing 10% FBS under 37°C/ 5% CO_2 before treatment. After washing with PBS, all tested cells were treated with DMEM containing 25mM D-Glucose U-13C6（Cambridge Isotope Laboratories (Tewksbury, MA)）and 2% BSA under 37°C/5% CO_2 for 24 hours or indicated time points for flux rate tests. [13]C labeled metabolites in medium and cell pellets were identified and quantified by using UPLC- QTOF- MS.

UPLC-QTOF-MS analysis: Cells were washed twice with phosphate-buffered saline (Thermo Fisher Scientific) (pH 7.4) and then carefully scraped into 1.5 mL safe lock centrifuge tubes with 10 mg of beads (0.5 mm). Metabolites were extracted by adding 0.5 mL of 50% MeOH (-20°C), followed by homogenization for 3 min using a Bullet Blender Tissue Homogenizer (Next Advance, Inc., Averill Park, NY) and centrifugation at 13, 500 g for 10 min at 4°C.

The resulting 50 μL of supernatant and also cell culture medium samples were transferred to 1.5 mL tube and mixed with 10 μL of 200 mM 3-NPH solution and 10 μL of the mixed 96 mM of EDC/pyridine methanolic solution. Derivatization was conducted by incubation at 30°C for 1 h before evaporation to dryness under nitrogen. 400 μL of 50% aqueous methanol was used to re-suspend the samples. The supernatants were used for UPLC-QTOFMS analysis according to previous reports with minor modifications [21,22].

Cell Metabolomics Analysis

Cell culture and sample collection: Cells were adapted to HuMEC Ready Medium and fed by fresh medium 24h before harvest for metabolomics assay. To perform the metabolite flux test, cells were starved for 16 hours in DMEM containing 2% BSA without glucose before feeding them with the HuMEC Ready Medium. Cells were then harvested cells at indicated time points.

Cell Metabolite extraction and derivatization: Cells were washed twice with phosphate-buffered saline (Mediatech, Manassas, VA) (1 mM pH 7.4) and then carefully scraped into 1.5 mL safe lock centrifuge tubes with containing 10 mg of beads (0.5 mm). Metabolites were extracted by adding 0.5 mL of 50% MeOH (-20°C), followed by homogenization for 3 min using a Bullet Blender Tissue Homogenizer (Next Advance, Inc., Averill Park, NY) and centrifugation at 13, 500 g for 10 min at 4°C. Derivatization was performed according to the method of Han et al.[23] with minor modifications. 50 μL of supernatant or standard was transferred to a 1.5 mL tube and

mixed with 10 μL of 200 mM 3-NPH solution, and 10 μL of the mixed 96 mM of EDC/pyridine methanolic solution. Derivatization was conducted by incubation at 30°C for 1 h before evaporation to dryness under nitrogen. 400 μL of 50% aqueous methanol was used to re-suspend the samples. The derivatives were analyzed via UPLC-QTOFMS.

Cell extracts analysis by ultra-performance liquid chromatography/xevo G2 quadrupole time-of-flight tandem mass spectrometry: The metabolites in culture medium and cell pellets were analyzed using the ACQUITY UPLC I-Class System coupled with a Xevo G2-S QTOF (ACQUITY UPLC- Xevo G2 QTOF, Waters Corp., Milford, MA). Chromatographic separation of phosphate and carboxylate was performed on the ACQUITY UPLC with a conditioned autosampler at 10°C, using an Acquity BEH C18 column (100 mm × 2.1 mm i.d.,1.7 μm particle size) (Waters, Milford, MA, USA). The column temperature was maintained at 40°C. The mobile phase consisting of water with 5 mM DIPEA in water (solvent A) and acetonitrile/isopropanol=7/3 (solvent B) was pumped at a flow rate of 0.3 mL/ min. The gradient elution program was as follows: 0–8 min, 1-20% B; 8–16 min, 20–98% B; 16.0–16.1 min, 98–1% B; 16.1–18.0 min, 1% B for equilibration of the column. The injection volume was 5 μL. The Xevo G2 QTOF mass spectrometer was used in negative ESI mode for data acquisition using UPLC/MS. Typical source conditions were as follows: capillary voltage, 3.0 kV; sample cone, 40 V; source temperature, 120 °C; desolvation temperature 450 °C; cone gas flow rate 50 L/ h; desolvation gas (N2) flow rate 900 L/h. All analyses were performed using the lockspray, which ensured accuracy and reproducibility. Leucine–enkephalin (5 ng /mL) was used as the lockmass generating a reference ion in positive mode at m/z 554.2615 and was introduced by a lockspray at 10 μL /min for accurate mass acquisition. Data acquisition was achieved using MS. The injection volume was 5 μL.

As shown in Fig. S6, a total of 28 phosphate and carboxylate metabolites were detected by UPLC-QTOF-MS. Table S1 showed the identification of metabolites in cells using UPLC-QTOF mass spectrometry.

References

Goncalves, M. D. *et al.* High-fructose corn syrup enhances intestinal tumor growth in mice. *Science* **363**, 1345-1349, doi:10.1126/science.aat8515 (2019).

Chen, W. L. *et al.* Enhanced Fructose Utilization Mediated by SLC2A5 Is a Unique Metabolic Feature of Acute Myeloid Leukemia with Therapeutic Potential. *Cancer Cell* **30**, 779-791, doi:10.1016/j.ccell.2016.09.006 (2016).

Chen, W. L. *et al.* GLUT5-mediated fructose utilization drives lung cancer growth by stimulating fatty acid synthesis and AMPK/mTORC1 signaling. *JCI Insight* **5**, doi:10.1172/jci.insight.131596 (2020).

Jang, C. *et al.* The Small Intestine Converts Dietary Fructose into Glucose and Organic Acids. *Cell Metab* **27**, 351-361 e353, doi:10.1016/j.cmet.2017.12.016 (2018).

Yabe-Nishimura, C. Aldose reductase in glucose toxicity: a potential target for the prevention of diabetic complications. *Pharmacol Rev* **50**, 21-33 (1998).

Vander Heiden, M. G., Cantley, L. C. & Thompson, C. B. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* **324**, 1029-1033, doi:10.1126/science.1160809 (2009).

Abdel-Haleem, A. M. *et al.* The Emerging Facets of Non-Cancerous Warburg Effect. *Front Endocrinol (Lausanne)* **8**, 279, doi:10.3389/fendo.2017.00279 (2017).

Goto, Y., Hotta, N., Shigeta, Y., Sakamoto, N. & Kikkawa, R. Effects of an aldose reductase inhibitor, epalrestat, on diabetic neuropathy. Clinical benefit and indication for the drug assessed from the results of a placebo-controlled double-blind study. *Biomed Pharmacother* **49**, 269-277, doi:10.1016/0753-3322(96)82642-4 (1995).

Sun, S. Z. & Empie, M. W. Fructose metabolism in humans - what isotopic tracer studies tell us. *Nutr Metab (Lond)* **9**, 89, doi:10.1186/1743-7075-9-89 (2012).

Krzeslak, A. *et al.* Expression of GLUT1 and GLUT3 glucose transporters in endometrial and breast cancers. *Pathol Oncol Res* **18**, 721-728, doi:10.1007/s12253-012-9500-5 (2012).

Ohkuma, T., Peters, S. A. E. & Woodward, M. Sex differences in the association between diabetes and cancer: a systematic review and meta-analysis of 121 cohorts including 20 million individuals and one million events. *Diabetologia* **61**, 2140-2154, doi:10.1007/s00125-018-4664-5 (2018).

Bononi, A. *et al.* BAP1 regulates IP3R3-mediated Ca(2+) flux to mitochondria suppressing cell transformation. *Nature* **546**, 549-553, doi:10.1038/nature22798 (2017).

Amelio, I., Cutruzzola, F., Antonov, A., Agostini, M. & Melino, G. Serine and glycine metabolism in cancer. *Trends Biochem Sci* **39**, 191-198, doi:10.1016/j.tibs.2014.02.004 (2014).

Sullivan, L. B. *et al.* Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in Proliferating Cells. *Cell* **162**, 552-563, doi:10.1016/j.cell.2015.07.017 (2015).

Xu, S. *et al.* Hexokinase 2 is targetable for HK1 negative, HK2 positive tumors from a wide variety of tissues of origin. *J Nucl Med*, doi:10.2967/jnumed.118.212365 (2018).

Jeremy M Berg, J. L. T., and Lubert Stryer. in *Biochemistry   The Glycolytic Pathway Is Tightly Controlled.* Ch. 16.2, (W H Freeman, 2002).

Debosch, B. J., Chen, Z., Saben, J. L., Finck, B. N. & Moley, K. H. Glucose transporter 8 (GLUT8) mediates fructose-induced de novo lipogenesis and macrosteatosis. *J Biol Chem* **289**, 10989-10998, doi:10.1074/jbc.M113.527002 (2014).

Manolescu, A., Salas-Burgos, A. M., Fischbarg, J. & Cheeseman, C. I. Identification of a hydrophobic residue as a key determinant of fructose transport by the facilitative hexose transporter SLC2A7 (GLUT7). *J Biol Chem* **280**, 42978-42983, doi:10.1074/jbc.M508678200 (2005).

Scheepers, A. *et al.* Characterization of the human SLC2A11 (GLUT11) gene: alternative promoter usage, function, expression, and subcellular distribution of three isoforms, and lack of mouse orthologue. *Mol Membr Biol* **22**, 339-351, doi:10.1080/09687860500166143 (2005).

Quattrini, L. & La Motta, C. Aldose reductase inhibitors: 2013-present. *Expert Opin Ther Pat* **29**, 199-213, doi:10.1080/13543776.2019.1582646 (2019).

Bononi, A. *et al.* Germline BAP1 mutations induce a Warburg effect. *Cell Death Differ* **24**, 1694-1704, doi:10.1038/cdd.2017.95 (2017).

Paglia, G. *et al.* Monitoring metabolites consumption and secretion in cultured cells using ultra-performance liquid chromatography quadrupole-time of flight mass spectrometry (UPLC-Q-ToF-MS). *Anal Bioanal Chem* **402**, 1183-1198, doi:10.1007/s00216-011-5556-4 (2012).

Han, J., Lin, K., Sequria, C., Yang, J. & Borchers, C. H. Quantitation of low molecular weight sugars by chemical derivatization-liquid chromatography/multiple reaction monitoring/mass spectrometry. *Electrophoresis* **37**, 1851-1860, doi:10.1002/elps.201600150 (2016).

SI Fig.1.

Fig.S2.

Fig.S3.

Fig.S4.

Fig.S5.

Fig.S6.

Table S1　Identification of metabolites in cells using UPLC-QTOF mass spectrometry.

		Retention Time (min)	Formula	Calculated mass M-H	Measured value	RE (ppm)
glycolysis	D-Glucose	7.74	C12H17N3O7	314.0994	314.0984	-3.2
	D-Fructose	8.6	C12H17N3O7	314.0994	314.0989	-1.6
	Glucose 1-phosphate	2.23	C12H18N3O10P	394.0657	394.0648	-2.3
	Fructose 1-phosphate	8.11	C12H18N3O10P	529.1090	529.1091	0.2
	Fructose 6-phosphate	8.4	C12H18N3O10P	529.1090	529.1090	0.0
	Glucose 6-phosphate	7.96	C18H22N6O11P	529.1090	529.1089	-0.2
	Fructose 1,6-bisphosphate	6.01	C18H24N6O14P2	609.0747	609.0744	-0.5
	D-Glyceraldehyde 3-phosphate	9.13	C15H17N6O8P	439.0767	439.0734	-7.5
	3-phosphoglyceric acid	7.38	C15H17N6O9P	455.0722	455.0688	-7.5
	Dihydroxyacetone phosphate	9.18	C15H17N6O8P	439.0767	439.0734	-7.5
	Lactate	7.91	C9H10N3O4	224.0677	224.0662	-6.7
	Pyruvate	10.91	C9H9N3O4	357.0930	357.0930	0.0
Pentose phosphate pathway	6-Phosphogluconic acid	6.97	C18H23N6O12P	545.1033	545.1035	0.4
	D-Sedoheptulose 7-phosphate	8.26	C19H25N6O12P	559.119	559.1197	1.3
	D-Erythrose 4-phosphate	8.79	C16H19N6O9P	469.0873	469.085	-4.9
	D-Ribose 5-phosphate	8.51	C17H20N6O10P	499.0979	499.0966	-2.6
	D-Ribulose 5-phosphate	8.66	C17H21N6O10P	499.0984	499.0968	-3.2
Citric acid cycle	Succinic acid	10.57	C16H15N6O6	387.1059	387.1028	-8.0
	Isocitric acid	7.22	C18H17N6O9	461.1057	461.1042	-3.3
	Citric acid	7.73	C18H17N6O9	461.1062	461.1029	-7.2
	Oxoglutaric acid	11.17	C23H20N9O8	550.1440	550.1436	-0.7
	L-Malic acid	10.29	C16H16N6O7	403.1002	403.0966	-8.9
	Oxalacetic acid	11.1	C22H19N9O8	536.1278	536.1284	1.1
Others	L-Serine	6.04	C9H11N4O4	239.0786	239.0754	-13.4
	Glutamic acid	7.84	C11H11N4O4	263.078	263.0742	-14.4
	L-Glutamine	7.04	C11H14N5O4	280.1046	280.1003	-15.4
	Aspartic Acid	10.23	C16H16N7O6	402.1162	402.113	-8.0
	Hydroxypyruvate	10.72	C15H13N6O6	373.0897	373.0858	-10.5
	Fumaric acid	9.43	C16H13N6O6	385.0897	385.0856	-10.6