Aster repulsion drives local ordering in an active system

Jorge de-Carvalho[1], Sham Tlili[2,#], Lars Hufnagel[3], Timothy E. Saunders[2,4,5,*], Ivo A. Telley[1,*]

[1] Instituto Gulbenkian de Ciência, Fundação Calouste Gulbenkian, Oeiras, Portugal

[2] Mechanobiology Institute, National University of Singapore, Singapore

[3] European Molecular Biology Laboratory, Heidelberg, Germany

[4] Department of Biological Sciences, National University of Singapore, Singapore

[5] Institute of Molecular and Cellular Biology, A*Star, Proteos, Singapore

[*] For correspondence: dbsste@nus.edu and itelley@igc.gulbenkian.pt

[#] Present address: Aix-Marseille Université | AMU · Institut de Biologie du Développement de

Marseille-Luminy (UMR 7288 IBDML)

Abstract:

Biological systems are a form of active matter, that often undergo rapid changes in their material

state, *e.g.* liquid to solid transitions. Yet, such systems often also display remarkably ordered

structures. It remains an open question as to how local ordering occurs within active systems.

Here, we utilise the rapid early development of *Drosophila melanogaster* embryos to uncover

the mechanisms driving short-ranged order. During syncytial stage, nuclei synchronously divide

(within a single cell defined by the ellipsoidal eggshell) for nine cycles after which most of the

nuclei reach the cell cortex. Despite the rapid nuclear division and repositioning, the spatial

pattern of nuclei at the cortex is highly regular. Such precision is important for subsequent

cellularisation and morphological transformations. We utilise *ex vivo* explants and mutant

embryos to reveal that microtubule asters ensure the regular distribution and maintenance of

nuclear positions in the embryo. For large networks of nuclei, such as in the embryo, we predict

– and experimentally verify – the formation of force chains. The *ex vivo* extracts enabled us to

deduce the force potential between single asters. We use this to predict how the nuclear division

axis orientation in small *ex vivo* systems depend on aster number. Finally, we demonstrate that,

upon nucleus removal from the cortex, microtubule force potentials can reorient subsequent

nuclear divisions to minimise the size of pattern defects. Overall, we show that short-ranged

microtubule-mediated repulsive interactions between asters can drive ordering within an active

system.

Keywords

Centrosome, cytoskeleton, force chains, packing, ex vivo, microtubules, syncytium

Introduction

Proliferation of the genome is a cornerstone of early development in all animals, generally achieved by cell division. Almost all insects first segregate genome copies into hundreds of nuclei (syncytium) and only at a specific nuclear density transform the single cell into a tissue[1]. The distribution and separation of nuclei in syncytia typically display surprising positional uniformity[2–4]. This uniformity is likely required for precise gene patterning and fate determination[5]. How the embryo controls internuclear distance so robustly has been a decades-long topic of debate[6–12].

In *Drosophila melanogaster*, early nuclear divisions are meta-synchronous, whereby nuclei gradually fill the inner cellular space until, nine division cycles or ~80 minutes post-fertilisation, 300–400 of them arrive at the cell cortex[13]. The nuclei are subsequently embedded within a two-dimensional topology near the cortex of the ellipsoidal embryo, where they undergo four more rounds of division to generate ~6000 nuclei[14] prior to cellularisation. Some nuclei do not reach the cell cortex, resulting in the final number after 13 divisions at the cortex being less than 8192 (2^{13}). The spatiotemporal synchronisation of nuclear divisions is governed by a reaction-diffusion process emerging from nuclei[4]. Furthermore, global nuclear positioning in the early embryo is crucial for synchronisation[9].

Microtubule dynamics is critical for nuclear migration to the cortex at nuclear cycle (n.c.) 9 and for their regular distribution upon arrival[15–17]. Nuclei are embedded in a regular matrix that either pulls or pushes them apart, leading to precise internuclear distances. In interphase of the cell cycle, microtubules are organised in radial arrays called 'asters', which are nucleated and organised by the centrosome[18–20]. The centrosome acts as the main microtubule organising centre (MTOC) and promotes polymerisation and focusing of microtubules[21]. Two asters are linked to each nucleus, being functional elements of the bipolar spindle during mitosis. Asters are required for nuclear separation following chromosome segregation in the embryo[22]. Embryos lacking core centrosomal components do not regularly distribute nuclei and abort development after a few division cycles[23–25]. Conversely, embryos that do not form actin caps and membrane furrowing, which are the precursor of uninuclear cell formation and are thought to help separate nuclei at the cortex, still show regular nuclear distribution at n.c. 10 and 11 [10]. The causality and the mode of mechanical separation during and after nuclear division remains an open problem, primarily due to limited visualisation in living samples and the growing mechanical complexity during development.

Here, by exploiting embryonic explants[26], which reduces complexity, and a cell cycle regulation

mutant[27,28] to uncouple microtubule organisation from nuclear division, we determine how

microtubule interactions spatially organise nuclei. In this system, we uncover the physical

principles of separation for simple nuclear arrays, and derive the microtubule-driven mechanical

separation potential. We find that nuclei behave as cargo associated to self-organising

microtubule asters which have repulsive properties. These results contrast with uninuclear model

systems, where cortical pulling and hydrodynamic forces appear to drive aster movement and

centring[19,29,30]. Our work reveals the underlying local biophysical interactions that pack and

order nuclei within a rapidly changing active system.

Results

Synchronous nuclear duplication within a constant surface area poses a geometrical challenge

(Fig. 1a, Supplementary Video 1). Spindle elongation during nuclear division[22] should cause

transiently smaller distances between spindles (and their asters) unless some leave the surface

after division. Alternatively, spindles may reorient their division axis to optimise the spacing

between nuclei. We hypothesised that spatial patterns are related to mechanical interactions

between spindle asters. Thus, we quantified neighbour distances (Fig. 1b), focusing on the

distance d between centrosomes belonging to nearest neighbour nuclei ('non-sister').

Importantly, despite organelle duplication, synchronous spindle expansion (defined by s, Fig

1b), and apparent collective nuclear movement in a finite space, the distance distribution between

neighbouring centrosomes within n.c. 10 did not exhibit noticeable decrease as a compensation

(Fig. 1c, black dots, see also Methods). Similarly, or possibly as a result, the nuclear separation

distance D only mildly decreased (Fig. 1d, black dots) during anaphase and telophase when

nuclear duplication and mitotic separation occurs (Fig. 1d, blue dots). In subsequent division

cycles, the mean distance between centrosomes gradually decreased while nuclear density

doubled. Still, we observed no abrupt decrease or oscillation in the nuclear separation during

duplication (Fig. 1e, Supplementary Fig. 1). These measurements suggest that a rigid mechanical

connection exists between centrosomes. Thus, we hypothesised the presence of a repulsive

mechanism between neighbouring nuclei or asters in the highly viscous, and effectively over-

damped cytoplasm.

There appears to be no long-range order in spindle orientation across the embryo[12], but we asked

whether there are local patterns. Defects may result in local ordering, akin to how subtle

variations in sand corn size and shape induce fractures in sand piles with a range of lengths[31,32].

In the embryo, we investigated whether there were chains of aligned spindles (Fig. 1f and

Methods). Calculating the probability of a given chain length, L, in different cycles (Fig. 1g-h,

Supplementary Fig. 2) we find $P(L) \sim L^{-\alpha}$, with $\alpha \approx 2.0 \pm 0.3$ in n.c. 13 (compared to $\alpha \approx 3.6 \pm$

0.5 with randomised spindle orientation). The value of α did not appear to decrease with

increasing nuclear density (Fig. 1h), though it was dependent on the thresholds for defining

chains (Methods). The value for α is similar to the exponent in cluster size variability in a range

of other physical models[33–35]. Our results support the presence of short-range interactions driving

spindle alignment.

We posited that repulsive interactions between astral microtubules underlie the forces

determining the magnitude and spatial extent of the interactions determining spindle position

and orientation. We consider a simple model of aster-aster interaction (Fig. 2a): (i) asters have a

radial microtubule structure nucleated from a centrosome; (ii) asters are self-repulsive due to

microtubule interactions, generating a "dumbbell"-like potential for each nucleus; and (iii)

nearest neighbour interactions dominate over longer-ranged interactions. Each spindle has a

single rotational degree of freedom (Fig. 2a, bottom). From our model (Methods), we predict

(Fig. 2b): (i) two nearby isolated spindles will align in parallel and orthogonal to their connecting

line (i.e., $\phi \sim 90°$); (ii) three equidistant spindles align at $\phi \sim 60°$ to each other; (iii) four and more

spindles align randomly. The system is geometrically frustrated for n>3 resulting in a broad

range of possible states with the same energy minimum[36].

In embryos, with hundreds of spindles, these predictions for small systems cannot be tested

experimentally. Thus, we took advantage of our embryo explant assay, which enables us to study

a small number of spindles in quasi-2D spaces[26] (Fig. 2c-d, Supplementary Video 2). We

measured spindle axis orientation ϕ relative to the separation axis (Fig. 1f). When a single

spindle in an explant divides (Fig. 2d(i)), the two subsequent spindles align in parallel (Fig.

2d(ii)). Further divisions resulted in random spindle orientation, even when the spindles were

uniformly distributed (Fig. 2d(iii), Supplementary Video 2). For three spindles, the alignment

was biased away from random (p=0.04, Supplementary Fig. 3b-c), with some arrangements

showing 60° alignment (Fig. 2d(iv), Supplementary Fig. 3a). However, we also observed cases

where two out of three spindles aligned at 90°, likely due to the spindles not being positioned

equidistant. The experimental observations were in agreement with our model predictions when

stochasticity was introduced (Fig. 2e-f, Supplementary Fig. 3d-e and Methods). To conclude, the

orientation and alignment of two-, three-, and four- syncytial mitotic spindles can be described

by a model of mechanical dumbbells with nearest-neighbour repulsive interactions.

Centrosomes are essential for nuclear organisation in the embryo[23]. Are centrosomes, rather than

nuclei, the active positioning structures in the early embryo? We propose that the aster repulsion

is proportional to microtubule probability density and likely decays rapidly with distance δ from

the MTOC (Fig. 2g). To test these ideas, we utilised *giant nuclei (gnu)* mutant embryos, which

undergo DNA endoreplication without mitosis; chromosome segregation is inhibited, leading to

one or few overgrowing (polyploid) nuclei, while centrosomes continue to duplicate and

split[27,28,37] (Supplementary Video 3). We produced embryo explants from *gnu* mutant embryos

and studied the positioning properties of a small number of microtubule asters in quasi-2D

spaces. Asters consistently moved towards the centre of the explant, even when initially located

near the boundary after cytosol deposition (Supplementary Video 4). We measured the shortest

distance of the centrosome from the boundary at steady state, which varied between R/2 and the

maximum distance R (Fig. 3a). Deviation from precise centring may be due to lipid droplets

(green circles in Supplementary Video 4) forming exclusion zones. Typically, individual asters

appear to self-centre within a restricted space, consistent with a radial force potential. Such short-

ranged centring forces could be generated by microtubule polymerisation acting against the

boundary of the water–oil interface[38,39]. Alternatively, hydrodynamic drag caused by

microtubule motor transport together with radial asymmetry of asters can generate a net pulling

force towards the centre[19]. In either scenario, once radial symmetry is restored the net force drops

to zero.

How far do asters separate? We investigated the steady-state distribution of two-aster

configurations in explants. Inter-aster interaction, if existing, must balance with the force

involved in moving asters away from the boundary. Two asters reached a steady-state separation

distance that scaled with explant size and boundary distances, typically up to 45 μm, but did not

scale further in larger explants. Consequently, the shortest boundary distances were not always

diametral (Fig. 3b, inset). For most explants the inter-aster distance and the shortest boundary

distances were similar (Supplementary Fig. 4a-c). Interestingly, two asters did not separate

according to equal force but approximately partitioned the available space ($d = b_1 = b_2 = $

$2R/3$). Some care is needed here though as (i) not all experiments covered the entire time course

of aster separation; and (ii) some asters were likely initially positioned farther apart by the

extraction procedure (red dots). Combined, these observations further support the presence of

short-ranged repulsive interactions between asters and between aster and boundary. The two

asters may mechanically interact via overlapping microtubules[40,41], while astral microtubules

may simply hit against the boundary interface, which acts as an immovable hard wall.

In explants, as the number of asters further increases, the shortest distance between the aster and
boundary decreases (Fig. 3c-d). Interestingly, their steady-state position often assumed highly
ordered, almost crystalline configurations (Fig. 3c). Explants with higher numbers of asters had
reduced aster separation distances, likely due to increased internal compression of larger 2D aster
networks (Fig. 3e). Finally, in *gnu* mutant embryos, asters organise into a regularly spaced
network and reach a dynamic equilibrium once the cortex becomes fully occupied
(Supplementary Video 5). Quantifying inter-aster distance revealed a surprisingly stable and
reproducible pattern along time and between embryos (Fig. 3f, Supplementary Fig 4d-e).
Notably, the inter-aster distance d_a in *gnu* mutants at steady state and in multi-aster explants are
comparable, matching the non-sibling distance d observed in wildtype embryos (Fig. 1c),
strongly indicating that positional patterning of centrosomes in the syncytial embryo can happen
independently of embryo cortical factors, nuclei, spindle assembly and mitotic regulation.

Aster dynamics provide information about the underlying mechanical interactions. Single asters
located near the boundary after cytosol deposition stayed for up to 10 min, but they always
eventually migrated (Fig. 4a, Supplementary Fig. 5b & Video 4). Single asters moved rapidly
after separation from the boundary, with a maximum velocity of 0.05±0.02 µm/s, at around 20%
of its final distance from the explant boundary (Fig. 4b, Supplementary Fig. 5, Methods), before
linearly decelerating and stopping between 15–35 µm from the boundary (Supplementary Fig.
5b-d). There was only weak correlation between the final aster position and explant size, perhaps
due to steric effects from lipid droplets (Supplementary Fig. 5a-b). We noticed fewer
microtubules oriented outwards when the aster was near the boundary (Fig. 4c), suggesting a
lack of space for microtubule elongation. Indeed, in some samples we observed splay of
microtubules near the explant boundary (Supplementary Fig. 5e). Finally, we analysed the
dynamics of the lipid droplets that are also present in the explant, to infer about passive behaviour
from hydrodynamic effects. There was a droplet exclusion zone of ~10 µm around each aster.
As the aster moved away from the boundary, the lipid droplets streamed around the aster,
maintaining their exclusion (Supplementary Fig. 5f-g). We quantified the motion of lipid
droplets with and without an aster present (Supplementary Fig. 5h). In the absence of an aster,
the lipid droplets appeared to move randomly ($x_{rms} \sim t^{1/2}$). In the presence of an aster, lipid
droplets moved faster, and appeared to move in a more directed manner ($x_{rms} \sim t^{2/3}$). Such
behaviour is consistent with the aster having a repulsive force potential that can act on the
surrounding lipid droplets and other boundary constraints.

Next, we tracked aster pairs during separation (Fig. 4d-e, Supplementary Fig. 6 & Video 6). The

final distance between asters correlated well with explant size (Supplementary Fig. 6a).

Interestingly, the peak separation velocity was always near half the final aster separation distance

(Fig. 4e), independent of final separation distance. This contrasts with the single aster scenario

(Fig. 4b), suggesting that the overall effective forces are different in the two cases, consistent

with our observations in Fig. 3c-e. The aster separation could be driven by overlap and

crosslinking of microtubules[17], or by mutual contact leading to repulsion by microtubules of both

asters. We quantified the microtubule intensity between the separating asters (Fig. 4f, left) and

generated kymographs of the microtubule fluorescence intensity at half the separation distance

(Fig. 4f, right). The intensity exponentially decayed (Supplementary Fig. 6e), consistent with

models of dynamic microtubule length distribution[42,43]. When aster separation ceased there was

almost no measurable microtubule signal between the asters.

For a viscous material, the velocity, v, of an object is dependent on the applied force F: $v \approx \gamma F$,

where γ is the effective viscous drag coefficient. Naïvely interpreting the microtubule

distribution as the resulting force profile does not match with the observed separation velocity

profile. However, multiplying the microtubule distribution by an effective *slipping* term, $f_{slip} =$

$f_0 \frac{x^2}{x_0^2 + x^2}$ ($x_0 \approx 15 \ \mu m$), results in an excellent fit to our observed aster separation velocities (Fig.

4g). We attribute this effective slipping to molecular friction between microtubules[44]. Finally,

we noticed that the microtubule density between the asters was often not maximal along the

shortest distance between the asters (Supplementary Fig. 6f-g), suggesting that the contact

interfaces between asters is more complex than assumed above.

To explore the nature of the microtubule interaction, we performed a series of inhibitory

treatments. We generated explants with two asters in the course of separation and pulse-injected

a defined volume of 200 μM colchicine, which causes acute depolymerisation of microtubules.

Upon injection the asters stopped separating and sometimes inverted their direction of motion

(Fig. 4h, Supplementary Video 7). We then tested whether active molecular machinery was

required for aster repulsion by inhibiting ATP consumption. We injected a series of

concentrations of sodium azide into explants that contained a separating pair of asters

(Supplementary Video 7). Adding sodium azide decreased the initial recoil velocity (dashed lines

in Fig. 4h) and also resulted in a considerable reduction in aster separation. However, even at

very high concentrations of sodium azide, we still observed residual motion, suggesting that both

actively driven microtubule-mediated separation and passive physical contact driven separation

occur.

From our observations, we conclude that aster separation is caused by a net repulsive force
between asters and between aster and boundary. However, either pushing or pulling forces can
cause the separation dynamics and centring[29,45,46]. We performed targeted UV photo-ablation
experiments in explants and generated ellipse-shaped ablations positioned asymmetrically
around one aster, affecting microtubules on the left side more than on the right side of the aster
(Fig. 5a, Supplementary Video 8). If hydrodynamic effects from vesicle transport along
microtubules drives aster motion[19], we expect a displacement to the right (positive) after
ablation. Conversely, if the net force applied on microtubules favours pushing on MTOC, we
expect a displacement to the left (negative). Indeed, asters consistently moved to the left,
supporting a dominating effect of microtubule-driven pushing (Fig. 5b). As a control, we
performed the same perturbation in explants that were injected with the microtubule inhibitor
colchicine (Fig. 5c). Under this condition, asters moved very slowly to the right (positive), which
is consistent with a weak hydrodynamic effect from other contractile sources (*e.g.* actomyosin[47]).

To further test these conclusions, we performed two types of ablation in explants containing two
asters (Fig. 5d, Supplementary Video 9): (1) light pulses emitted along an ellipse around both
asters to destroy microtubules in the periphery; (2) light pulses emitted along a line between the
two asters to destroy microtubules between asters. If forces are attractive, then ablation type 1
will stop separation while ablation type 2 will lead to an acceleration. If forces are repulsive, we
predict the opposite response. We found a slight acceleration for peripheral ablation and a strong
deceleration with recovery for central ablation (Fig. 5e). Separation recovered likely because of
fast regrowth of microtubules after ablation (in the range of μm/min[48]). In summary, the dynamic
behaviour of asters in our explants is consistent with a model of radially symmetric microtubule
repulsion.

But, what is the relevance of our findings *in vivo*? Specifically, do the aster interactions enable
the embryo to pack the nuclei in a more regular manner? Heterogeneities in nuclear density are
a common phenomenon in early embryos resulting from aberrant cortical migration or nuclear
internalisation due to mitotic failure (Fig. 5f, Supplementary Fig. 7a). We predicted that nuclei
neighbouring a low-density region will orientate their division axis towards that region, where
repulsion is weakest. We identified such low-density regions and quantified the subsequent
division orientation of the surrounding nuclei (Methods), which confirmed our prediction (Fig.
5g, Supplementary Fig. 7b, Supplementary Video 10). Repeating the same analysis on regions
of uniform nuclear density showed no correlation in the division angle (Supplementary Fig. 7c).
We also generated acute density reductions by UV ablation. Using low laser damage, the target

nuclei failed to divide and detached from the cortex lowering local nuclear density

(Supplementary Fig. 7d-e). Subsequently, the surrounding nuclei adjusted their division axis to

orientate into the perturbed region (Supplementary Fig. 7f). Combining our results from

spontaneous low-density regions and ablated embryos (including larger ablations of 3–5 nuclei),

we see that the microtubule repulsion mechanism is efficient in adjusting the angle of division

to compensate for heterogeneities in nuclear packing (Fig. 5h).

Discussion

Robust embryonic development critically depends on homogenous delivery of nuclei to the cell

cortex and subsequent maintenance of a regular nuclear distribution despite further division

cycles[15,16]. Recent work has shown how nuclear divisions are synchronised and, as a

consequence, how nuclei are globally distributed around the embryo cortex[4,9]. However, these

results assumed that nuclei are positioned regularly after each round of duplication[3]. Yet, what

ensures such local order of nuclei? To answer this requires deepening our understanding of the

biophysical principles defining the orientation of the spindle axis and how nuclei separate and

reposition during division cycles. Our explant experiments demonstrate that, in the absence of

perturbation by neighbour interactions, the ground-state orientation of a spindle is orthogonal to

the previous division axis (Fig. 2f). This can be explained by the stereotypical migration of the

two centrosomes from their common origin, each along one quadrant of the nucleus, until they

form the poles of the bipolar spindle[49]. Hence, orthogonality of spindle axes likely emerges from

the geometric nature of bipolar structures and symmetry considerations. Surprisingly though, a

system of four or more spindles in a two-dimensional space evolves towards random

orientations, arguing against active spindle orientation control by the cell. Our analysis suggests

that force balancing and energetic minimisation in a noisy two-dimensional environment

dynamically determine where nuclei are positioned and in which orientation they propagate upon

division. Interestingly, in larger networks, small positional irregularities result in division axis

orientation towards the low-density region, enabling the nuclear distribution to homogenise

quickly and act as a self-repair mechanism. Finally, it would be interesting to compare aster force

driving spindle alignment in the early embryo with similar microtubule-driven processes across

cells, such as mitosis in polarised tissue growth[50].

Why is a high spatial regularity of nuclei important for the embryo? After n.c. 13, the embryo

transforms into a multicellular embryo by engulfing each nucleus with plasma membrane[1]. The

nearest neighbour internuclear distance defines cell size. Therefore, a narrow distance

distribution leads to a uniform size of cells that subsequently assume distinct function during

body part definition. Analysis of information decoding in the *Drosophila* embryo has showed how each individual cell unambiguously reads its current position, which defines a specific function later in development[5]. But these results are dependent on the interpreting units (*i.e.* the nuclei) being uniformly distributed around the embryo. Therefore, we can conclude that a robust mechanism defining cell size and position is crucial as size irregularity would effectively decrease positional precision.

Acknowledgements

We thank members of the IAT and TES labs for fruitful discussions, and Virgile Viasnoff, Gianluca Grenci, Tetsuya Hiraiwa, Jacques Prost and Thomas Surrey for constructive comments and feedback. We thank the staff of the Fly Facility, the Advanced Imaging Facility (AIF) and the Technical Support Service at the Instituto Gulbenkian de Ciência (IGC). Transgenic fly stocks were obtained from Bloomington Drosophila Stock Center (NIH 741P40OD018537). We acknowledge financial support from: Human Frontiers Science Program (HFSP), awarded to IAT and TES and supporting JdC and ST (RGY0083/2016); Fundação Calouste Gulbenkian (FCG), the Fundação para Ciência e a Tecnologia (FCT) supporting IAT (Investigador FCT1072IF/00082/2013); EU FP7-PEOPLE-2013-CIG (Nº 818743) awarded to IAT and supporting JdC; LISBOA-01-0145-FEDER-007654 supporting IGC's core operation; LISBOA-01-7460145-FEDER-022170 (Congento) supporting the Fly Facility; PPBI-POCI-01-0145-FEDER-022122 supporting the AIF, all co-financed by FCT (Portugal) and Lisboa Regional Operational Program (Lisboa2020) under the PORTUGAL2020 Partnership Agreement (European Regional Development Fund).

Figures

(Figures 1–5)

Fig. 1 – Spindle separation and orientation in the syncytial blastoderm of *Drosophila melanogaster* reveals robust distance maintenance and local orientation patterns. (a) Time lapse (maximum intensity Z-projections) (Supplementary Video 1) from an embryo expressing H2Av::RFP (magenta) and Spd2::GFP (green) during n.c. 10 (scale bar, 20 µm left, 10 µm right). Each panel corresponds to the indicated mitotic phase. **(b)** Morphological identification of mitotic phases and hierarchical classification of distances; extended figure legend in Supplementary Fig. 1. **(c-d)** Inter-aster (**c**) and nuclear (**d**) distance during n.c. 10 (n=15 spindles, N=5 embryos). **(e)** Schematic of average inter-aster distances during blastoderm n.c. 10–13 (also Supplementary Fig. 1). **(f)** Maximum intensity Z-projection of an embryo in n.c. 11, expressing Jupiter::mCherry (scale bar, 20 µm). Schematic illustrates how neighbouring spindles belonging to an alignment ('force') chain form angles θ and ϕ relative to each other (details in Supplementary Fig. 2a-b). Resulting chains are shown on the right; lines denote connections that satisfy two (thick) or only one (thin) of the conditions defining a chain (Methods). **(g)** Cumulative probability function of chain length for different cycles (n=7 embryos each for n.c. 10, 11, 12, 13). P-value calculated from Kolmogorov-Smirnov test. **(h)** Scaling of chain length probability with chain length for n.c. 10–13. Lines represent fit to $\beta L^{-\alpha}$, where L is the chain length and α is the scaling exponent (Methods).

Figure 2 – Division axis orientation for 2–4 spindles in a cytosolic explant are consistent
with a simple dumbbell model of aster repulsion. (a) Schematic of an aster, polymerised and
organised by a centrosome (green), which translates into a concentric repulsion potential, here
represented by dashed circles. When two asters are coupled to a nucleus (magenta) the repulsive
potentials translate into a dumbbell potential with a rotational degree of freedom. (b) Scheme of
dumbbells representing spindles in consecutive divisions. The graphs below show the expected
probability of division angles ϕ between dumbbell axes from free energy considerations
(Methods). Right panel shows predicted alignment for a three-spindle arrangement. (c)
Schematic of cytosol extraction from a *Drosophila* syncytial embryo and explant formation. (d)
Maximum intensity Z-projections of explants from embryos expressing Jupiter::GFP (grey) and
H2Av::RFP (not shown) containing different numbers of spindles (10 μm scale bar). (e) Angle
between the division axes of two spindles (left, n=32 explants) and between nearest neighbour
spindles in a four-spindle scenario (right, n=40 explants) as measured in experiments (blue) or
obtained from simulation (magenta, Methods). (f) Cumulative probability of the angle between
division axes for two-, three- (n=5 explants) and four-spindle scenarios. Dashed lines are model
predictions. (g) Proposed repulsion force as function of distance from the centrosome (green in

352 **a**) with peak at ~15 μm and negligible for >45 μm. The dashed line represents the short-distance

353 interaction regime that is below the diffraction limit of optical resolution.

Figure 3 – Free asters separate and achieve steady-state distance regularity. (a) Explant with single aster: scatter plot of the shortest distance to the boundary (*b*, see inset) as a function of the explant radius (*R*) (n=78). Magenta line: linear regression with zero intercept. **(b)** Explant with two asters: scatter plot of inter-aster distance (d_a, see inset) as a function of the maximum

projected separation M, calculated from the explant diameter (2R) and the boundary distance of

each of the two asters (n=54). The blue dashed line represents the estimated upper limit of the

interaction distance between two asters (~45 µm). Red dots represent cases where the two asters

were likely positioned far apart during explant generation. The yellow dots are cases of small

explants where projection leads to overestimation of b (Methods) **(c)** Maximum intensity Z-

projections of explants containing multiple asters extracted from *gnu* mutant embryos expressing

RFP::β-Tubulin (magenta) and Spd2::GFP (green) (scale bar, 10 µm). Dashed yellow circle

represents the explant boundary, and white dashed lines highlight the symmetry in the aster

distribution. **(d)** Distribution plot of shortest boundary distance (b, top scheme) and the ratio

b/R from explants containing 3 (n=19), 4 (n=11), 5 (n=8), 6 (n=5) and 7 (n=3) asters. Magenta

bars represent mean value. **(e)** Distribution plot of inter-aster distance (d_a, see inset) of single

explants containing nine or more asters. **f)** Scatter plots of inter-aster distance from five *gnu*

mutant embryos in order of increasing aster density.

Figure 4 – Aster dynamics in explants depends on microtubule distribution and interactions. (a) Maximum intensity Z-projections of a single aster moving away from the boundary of an explant produced from *gnu* mutant embryos expressing RFP::β-Tubulin

(magenta) and Spd2::GFP (green). Scale bar, 20 μm **(b)** Average migration velocity of single

asters away from the explant boundary (n=7). Distance normalised by the final, steady-state

distance for each aster (also Supplementary Fig. 5a). **(c)** Average microtubule density (black

line, inferred from RFP::β-Tubulin signal) along shortest distance to explant boundary,

normalised by the maximum intensity within each experiment. Grey traces are individual

experiments (n=7). **(d)** As in (**a**) but for an explant containing two separating asters. **(e)** Aster

separation velocity as a function of normalised separation distance (n=9). For each experiment,

distance is normalised by the final, steady-state separation distance (also Supplementary Fig. 6).

(f) (left) Colourmap of normalised microtubule density (as in **c**) between two separating asters.

(right) Kymograph of microtubule intensity between the asters during separation. Scale bars, 2

min (horizontal) 5 μm (vertical). **(g)** Fitting to average separation velocity (circles) considering

microtubule intensity and a microtubule slipping term. Microtubule density was either fitted

beforehand (solid line, Supplementary Fig. 6e) or directly included (dashed line). **(h)** Aster

separation dynamics upon injection of buffer (control, n=3), 0.5 mM (n=3), 10 mM (n=4),

100mM (n=3) sodium azide, or 0.2 mM (n=3) colchicine. [*] denotes $p<0.05$. Grey or coloured

areas around average curves in **b**, **c**, **e** and **h** denote ± 1 s.d.

a single aster asymmetric ablation

$t = 0$

$t = \Delta t$

b – colchicine

c + colchicine

d (1) ablation around two separating asters   (2) ablation between two asters

$t = 0$

$t = \Delta t$

e type 1 type 2 ablation

f 0:00   8:00   24:00

g cycle 10 cycle 11 cycle 12 cycle 13

neighbor nucleus

φ position of internalization

h

Figure 5 – Microtubule-dependent repulsion provides a mechanism for spindle alignment towards lower density. (a) Schematic of single aster eccentric circular UV laser ablation (magenta dashed line); t=0 min denotes ablation time. **(b,c)** Aster displacement before and after ablation in explants unperturbed (**b**) or treated (**c**) with colchicine. Arrows represent average displacement magnitude and direction, and vertical and horizontal grey bars denote ± 1 s.d. of displacement in x and y, respectively. **(d)** Two-aster ablation procedure: (1) ellipse ablation around two asters during separation; (2) linear ablation between two asters. **(e)** Change of inter-aster distance upon laser ablation (at t = 0 min) as described in (**d**). **(f)** Maximum intensity Z-projections from an embryo expressing H2Av::mCherry (magenta) and Spd2::GFP (green) in n.c. 12–13. Yellow arrow (top panel) denotes internalisation of a nucleus. The centrosomes remain at the embryo cortex (yellow circle, middle panel). Division axes of neighbouring spindles (yellow lines) orientate towards the location of internalisation in n.c. 13 (bottom panel). Scale bar, 20 μm; time in min:sec. **(g)** Probability density function of the division angle orientation φ of neighbouring nuclei to regions of low nuclear density in n.c. 10–13 (n=67, 116, 96, 73 angles from N=10, 15, 15, 15 embryos in n.c. 10, 11, 12, 13 respectively). **(h)** Cumulative distribution function of division axis angle at the end of n.c. 13 towards artificially generated holes generated by single-pulse UV laser ablation (n=108 angles from 15 embryos). The dashed black line represents random division orientation (also Supplementary Fig. 7).

Materials and Methods

Fly strains

Flies with genotypes w[1118]; +; endo>Jupiter::GFP (stock no. 6836, Bloomington) and w*; +; endo>H2Av::RFP (stock no. 23650; Bloomington) were crossed to generate recombinant progeny. Similarly, flies expressing fluorescent reporters recombined on the 2[nd] chromosome were produced by crossing the following stocks: w*; endo>H2Av::RFP; + (stock no. 23651; Bloomington); w*; pUbq> β-Tubulin::EGFP; + (stock no. 109603, Kyoto) ; w*; pUbq>RFP::β2-Tubulin; + (see Reference[51]) and w[1118]; pUbq>Spd2::GFP; + (all provided by Mónica Bettencourt Dias, Instituto Gulbenkian de Ciência, Portugal). All resulting recombinant fly lines are homozygous viable. w[1118]; +; Jupiter::mCherry flies were generated by and obtained from Nick Lowe in Daniel St Johnston's lab (The Gurdon Institute, United Kingdom). H2Av::mCherry flies were generated as described elsewhere[13]. Two different mutants of giant nucleus (*gnu*), namely w*; +; gnu[305]/TM3 (discontinued stock no. 3321; Bloomington) and w*; +; gnu[Z3-3770A]/TM3 (discontinued stock no. 38440; Bloomington), were each balanced with w[1118]; CyO/Sco; MKRS/TM6B (stock no. 3703, Bloomington). Above described recombined

lines on the 2nd chromosome were individually crossed with gnu mutants and kept as balanced
stocks. Finally, trans-heterozygous were generated for gnu^{305}/gnu$^{Z3-3770A}$ mutants, whereby only
flies homozygous for the fluorescent reporters on the 2nd chromosome were selected for
increased signal collection during live microscopy. These trans-heterozygotes laid fertilized eggs
which undergo several embryonic rounds of chromatin replication and centrosome duplication,
allowing for the study and quantification of asters at the embryo cortex.

Embryo collection and sample preparation
We followed established procedures[52] of fly husbandry, keeping flies at 25°C under 50-60%
humidity. For embryo collections, young adult flies were transferred to a cage coupled to an
apple juice agar plate. After 2–3 rounds of egg laying synchronisation, developing embryos were
collected every 30–60 minutes. In the case of *gnu* mutants, embryos were collected at different
time intervals, ranging from 30 min up to 4h. Embryos were dechorionated by short immersion
in 7% sodium hypochlorite solution (VWR). After extensive rinsing with water, embryos were
aligned and immobilised in a thin strip of heptane glue placed on 22x22mm coverslips, and
covered with halocarbon oil (Voltalef 10S, Arkema).

Microscopy
Time-lapse acquisitions were conducted on a Nikon Eclipse Ti-E microscope equipped with a
Yokogawa CSU-W Spinning Disk confocal scanner and a piezoelectric stage (737.2SL, Physik
Instrumente). For embryo imaging, 15 μm (31 planes) Z-series stacks were acquired every 15s,
using a Plan Fluor 40x 1.3NA oil immersion objective, the 488nm and 561nm laser lines, and an
Andor Zyla 4.2 sCMOS camera to acquire images. For explants up to 100μm in diameter, we
used a Plan Apo VC 60x 1.2NA water immersion objective with 2x post-magnification and an
Andor iXon3 888 EMCCD camera. When needed, the Andor Zyla 4.2 sCMOS camera was
selected to acquire a 2x wider field of view with the same spatial resolution or, alternatively, the
Apo λ S LWD 40x 1.15NA water immersion objective.

Single embryo explant assay
Embryo extractions were performed as previously described[26,53]. Briefly, cytosol from wild-type
embryos between telophase and subsequent interphase of cycle 8 was extracted by puncturing
the vitelline membrane with a sharp glass micropipette and flow control by operating a bi-
directional syringe pump. Small explants of cytosol (in the picolitre range) were deposited on
poly-L-lysine coated glass surface under halocarbon oil. Time-lapse acquisitions typically
started in late interphase or prophase. In the case of *gnu* mutant embryos, most extractions were
performed when few centrosomes (between 5 and 40) were visible at the anterior-lateral cortex.

During extractions, shear stress was avoided to prevent structural damages and undesirable

molecular dissociations that induce premature mitotic failures or aberrant microtubule structures.

In *gnu* mutant embryos, repeated use of the same extraction micropipette is not recommended.

Explants from wildtype embryos initially containing a single nucleus were selected for time-

lapse imaging of subsequent mitotic divisions. Explants from *gnu* mutants initially containing a

single free aster near oil interface or two free asters in close proximity were selected for time-

lapse imaging of aster separation. All experiments were conducted at 25±1 °C.

Pharmacological perturbation of embryo explants

Pharmacological perturbations were performed by adding different drugs (colchicine at 0.2 mM,

sodium azide at 0.5, 10, or 100 mM) diluted in cytoplasm-compatible buffer (50 mM HEPES,

pH 7.8, 100 mM KCl, 1 mM MgCl$_2$). Solutions were directly administrated to the explants using

a fine pipette (pulled using a Narishige PC-100 Puller with a: 2-step (69% + 55%) heating

protocol and with 4 mm drop length) connected to an Eppendorf FemtoJet® 4i pump. The final

drug dilution in the explants was of approximately 1:10 (solution:cytosol). Buffer injections were

conducted as control.

Laser ablation system

The laser ablation systems used for experiments with intact embryos (at EMBL Heidelberg,

described elsewhere[22]) and embryo extracts (at Instituto Gulbenkian de Ciência, implemented

by IA Telley on the microscope described above) were conceptually identical. A Crylas FTSS-

355-Q pulsed laser emitting 355 nm, 1.1 ns pulses, 15µJ pulse energy at 1 KHz was aligned with

a beam expander (16x), a scan head (SCANcube 7, Scanlab, Germany) coupled to an f-theta lens

(f=56 mm, anti-reflection coating for 340–370 nm, SCANLAB AG, Germany). The focus point

of the f-theta lens was aligned to be parfocal to the focal plane of the objective, using a tube lens

(f=200 mm, Ø=30 mm, 355 nm AR coated, OWIS, Germany) and a dichroic mirror (T387

DCLP, Chroma) in the upper stage filter wheel. Any scattered light was blocked at the emission

side with a RazorEdge LP 355 dichroic mirror OD6 @ 355nm (Chroma). The system was

controlled with homemade journals for Metamorph software (Molecular Devices Inc.). The

optimal laser power was set to ensure microtubule ablation while avoiding thermal expansion of

cytoplasm, with post-ablation microtubule signal recovery matching known polymerisation

dynamics. This combination of conditions proved to be efficient at ablating target structures

beyond fluorophore bleaching. In explants containing a single aster, astral microtubules were

asymmetrically ablated by positioning an ellipsoid off-centre (21.7 by 10.8 µm, 4 times, 15s

interval, 0.54 µm step, laser power: 25%) (Fig. 5a). In explants containing two asters, astral

microtubules were ablated using an ellipsoid (21.7 by 10.8 µm, 3 times, 15s interval, 0.54 µm

step, laser power: 10–15%) roughly centred at the mid-point between the two asters, while

interpolar microtubules were ablated using linear ablations (21.7 µm, 3 times, 15s interval 0.54

µm step, laser power: 10–15%) perpendicular to the axis connecting the asters (Fig. 5d).

Distance analysis in embryos

Automated positional detection of the signals from centrosomes and nuclei (or chromatin) was

performed by applying a Gaussian blur filter (radius: 1–2 pixels) and using the plugin TrackMate

v3.5.1 in FIJI ImageJ[54,55]. The coordinates of detected spots were imported into MATLAB® for

assignment and distance calculation. The connection between poles belonging to a spindle

structure was assigned in a custom-made script requiring user input, on an area containing 15–

40 spindles for each mitotic phase and per embryo. For each spindle–assigned coordinate

positions, the nearest neighbour positions were determined using the Delaunay triangulation

functions in Matlab® yielding a connectivity list. Thereby, a spindle structure is defined as a

combination of n centrosome and m chromatin positions (n,m) with the following numbers for

mitotic phases: late interphase (2,1); anaphase A (2,1); anaphase B (2,2); telophase (4,2), early

interphase (4,2). With this assignment, the duplicated organelles dissociate at the transition from

telophase to early interphase, so that two related nuclei become independent neighbours. Next,

the 3D Euclidean distances between relevant positions were calculated from position coordinates

with a computer-assisted manual heritage classification. The distance between separating

chromosomes $\bar{D}$ was calculated from the two chromatin entities within a spindle. Spindle length

s was calculated from the distance between two centrosomes belonging to each spindle. In phases

with four centrosomes per spindle, two at each pole, spindle length was defined as the smallest

distance between opposite centrosomes (four possible combinations). The sister centrosome

distance $\bar{s}$ was calculated from centrosome pairs at each spindle pole. Inter-aster distance d

(corresponding to the distance between non-sister centrosomes) was calculated between

different, neighbouring spindles by selecting all centrosomes not associated with the same

spindle from the nearest neighbour connectivity list. Inter-nuclear distance D was calculated

between nearest neighbour nuclei or chromosomes not belonging to the same spindle. Finally,

the arithmetic mean and standard deviation of the distance distributions within a single embryo

were calculated and overlaid for division cycles 10–13 (Supplementary Fig. 1). In *gnu* mutant

embryos, inter-aster distance d_a was calculated from a selected region containing 15–20

centrosomes, for each embryo at different time points using triangulation and neighbourhood

connectivity list. Centrosomes located at the anterior hemisphere were excluded to avoid the

influence of the giant polyploid nucleus. The *gnu* mutants present variable centrosome densities

depending on age and other unknown factors. We analysed the variation of distance distribution

for five different embryos with similar densities during intervals of 10 min. All data plots were

generated in MATLAB®.

Spindle alignment analysis in embryos

For each embryo, we took the time point two minutes prior to the first onset of nuclear division.

The angle of the centrosome pair was used to define an orientation angle for the spindle.

In calculating the force chains, we did the following for each cycle:

1. For each spindle i, identify nearby spindles (taken as 1.3 times {average nearest neighbour

distance});

2. Calculate the orientation angle difference between pairs of nearest spindles. This value can

range between 0° and 90° (as there is no direction to the spindle orientation). We define this angle

as θ_{ij} for each pair of spindles denoted by i and j respectively.

3. Find the vector between each spindle and its nearest neighbours $\mathbf{x}_{ij}$. Find the angle, ϕ, between

the orientation of the spindles i and j relative to $\mathbf{x}_{ij}$. As the orientation does not have a specific

direction, for angles >90° we define $\Phi = 180 - \phi$.

4. We tested different thresholds on θ and ϕ (annotated by $\Delta\theta$ and $\Delta\phi$) to find the chain length

frequency (Supplementary Fig. 2c). For a spindle to belong to a chain, at least the angle θ had

to meet the condition $0 \leq \theta \leq \Delta\theta$. In addition, a second condition was defined: $\Delta\phi \leq \phi \leq 90°$.

Neighbouring spindles that meet both conditions belong to a chain that exhibits alignment both

along the chain axis and between spindles. In the below table, we give the fitted value of α (as

defined in main text) for different constraints on θ and ϕ.

Cycle	$\theta = \phi = 45^\circ$	$\theta = 45^\circ, \phi = 30^\circ$	$\theta = 30^\circ, \phi = 45^\circ$	$\theta = \phi = 30^\circ$
10	1.9	2.7	2.7	3.4
11	1.9	3.3	2.9	4.0
12	2.1	3.1	3.4	3.8
13	2.0	3.4	3.0	4.6

549 Table 1A: α for different conditions on chain alignments for experimental data

Cycle	$\theta = \phi = 45^\circ$	$\theta = 45^\circ, \phi = 30^\circ$	$\theta = 30^\circ, \phi = 45^\circ$	$\theta = \phi = 30^\circ$
10	4.0	5.1	4.7	4.6
11	4.1	5.0	4.8	6.0
12	3.8	4.7	4.7	5.5
13	3.6	4.8	4.4	5.6

Table 1B: α for different conditions on chain alignments for randomised data

Spindle alignment in explants

Extracts initially containing 2, 3 and 4 dividing nuclei were analysed in terms of spindle axis

orientation by analysis of microtubule reporters at the onset of anaphase B. Using MATLAB®

home-made scripts, the two minor orthogonal angles (ϕ_1 and ϕ_2) were determined by manual

clicking at spindle poles. These angles can range between 0º (parallel orientation) and 90º

(perpendicular orientated).

Dumbbell model of nuclear alignment in explants

For the model, we consider a simple phenomenological model: $H = J \sum_{<i,j>} (\vec{S_i} \cdot \vec{r}_{i,j})^2$, where $\vec{S_i}$

represents the orientation of aster i (with $|S_i| = 1$) and $\vec{r}_{i,j}$ is the unit vector between aster i and

its nearest neighbours j. The model is quadratic as there is no preferential direction for S. In this

case, energy is minimised if nuclei align perpendicular to the vector of separation, $\vec{r}_{ij}$, between

nuclei. In the case of two nuclear spindles, it is clear this results in parallel aligned nuclear

spindles, both of which are perpendicular to the vector between the nuclei. For three spindles,

positioned at (0,0), (1,0) and $(\frac{1}{2}, \frac{\sqrt{3}}{2})$, the energy is minimised for $S_1 = \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), S_2 = (\frac{1}{2}, \frac{\sqrt{3}}{2})$

and $S_3 = (1,0)$ (direction of the vectors can also be inverted). We use a Metropolis algorithm to

simulate the alignment of asters at different effective temperatures T. There is only one

parameter, defined by $J/k_B T$, which represents the competition between alignment force (J) and

random fluctuations ($k_B T$). In Supplementary Fig. 3d-e, we show simulation outputs for $\frac{J}{k_B T} =$

10^{-3} and 10^0. Result presented in Fig. 2 are for $\frac{J}{k_B T} = 10^{-1}$.

Of course, we can consider more complex models, such as $H = J \sum_{<i,j>} (1 -$

$\overrightarrow{(\vec{S_i} \cdot \vec{S_j})^2}) (\vec{S_i} \cdot \vec{r}_{i,j})^2$ which incorporate both terms involving neighbouring aster alignment and

their alignment relative to $\vec{r}_{ij}$. There is also similarity to models of nematic ordering in liquid

crystals[33], which have recently been applied to other biological systems[56]. Studies of self-

propelled particles with repulsive interactions are also relevant, where longer ranged interactions

are also considered[57]. Our aim here is to simply show how simple dumbbell-like repulsion

(which results in one rotational degree of freedom) can lead to different behaviours depending

on the system topology, not to build a precise model for how such potentials interact.

Analysis of free asters in explants – distance distributions

Distance between asters and from aster to the boundary were obtained in explants at steady state,

i.e. where asters did not move anymore (usually 30–45 min after explant deposition). The inter-

aster distance was determined as Euclidean distance in 3D. We defined the *boundary distance*

(b, b_1, b_2) as the shortest distance from the aster to the interface between glass, oil and cytosol,

determined manually using the Fiji measurement tools (at a precision of ±0.5 µm). To determine

the explant boundary on the glass (approximated with a circle of radius R), maximum intensity

projections of both fluorescence emission channels was assessed to trace the interface between

the glass, oil and cytosol. For larger explants with high aspect ratio – a quasi-2D situation – the

definition of *boundary distance* served as good approximation for a boundary in two dimensions.

However, in small explants where the aspect ratio is not as high, two asters sometimes aligned

considerably in the third dimension. In these cases, the definition for *boundary distance* led to

an underestimation of the maximum projected inter-aster distance $M = 2R - b_1 - b_2$; it

becomes a geometric problem in 3D and the longest dimension is not necessarily in the plane of

the glass-explant interface. This is evident for some data points in small explants (yellow dots in

Fig. 3b).

Analysis of free asters in explants – dynamics

The coordinates of free asters were obtained by applying a Gaussian blur filter (radius: 1–2

pixels) and using the plugin TrackMate v3.5.1 of Fiji ImageJ[54,55]. The coordinates of detected

spots were imported into MATLAB® for assignment and distance calculation similarly as

mentioned above.

The instant relative velocity was calculated using the formula: $v_i = \frac{d_{i+1} - d_{i-1}}{t_{i+1} - t_{i-1}}$, where d is the 3D

Euclidian distance and t is time in the flanking time points of the measure point.

For unperturbed experiments, data was normalised to the maximum distance achieved in the

separated phase in order to correct for scaling effect during splitting dynamics (Fig. 4). This data

was fitted to the phenomenological equation (Supplementary Fig. 6b):

604
$$u = a + b \cdot \tanh\left(\frac{t}{\tau}\right)$$

To analyse the lipid droplets, we performed a similar analysis using Fiji TrackMate. Seven

extracts were analysed with an aster present, with over 100 individual tracks of lipid droplets.

RMS distance was then extracted across the entire time course of imaging. Similar analysis was

performed in extracts without an aster. Curves (Supplementary Fig. 5h) were fitted using the 'fit'

function in MATLAB®, with $r^2 = 0.98$ ($b \cdot t^{\frac{1}{2}}$ for no-aster data), $r^2 = 0.96$ ($a_1 \cdot t^{\frac{1}{2}}$ for 1-aster data),

and $r^2 = 0.99$ ($a_2 \cdot t^{\frac{2}{3}}$ for 1-aster data). Fitting the general function $a \cdot t^c$ gives a best fit for $c =$

0.66 for the one aster data.

Microtubule profile quantification

For single asters (Fig. 4a-c), we quantified the microtubule intensity using the intensity of the
RFP::β-Tubulin signal. Taking the point when asters were either 5 μm or 20 μm from the explant
boundary, we used FIJI to measure the microtubule intensity along a 10 μm straight line from the
edge and through the aster. The line had a width of 2 μm. For each experiment, we normalised
the total intensity by the maximum measured value and then binned the data in 0.2 μm bins.
Hence, the recorded intensity does not reach one, and the mean intensity only reaches a
maximum around 0.8 as the maximum value does not occur at the same position.
Similar analysis was performed for the scenario with two asters (Fig. 4d-g). In this case, the
centroids of the asters were used to define a straight line along which the microtubule intensity
was measured throughout the process of aster separation. From this straight line between the
asters, we also generated the kymograph shown in Fig. 4f right.

Analysis of free asters in explants – perturbations with drugs and UV ablation

For comparison between control and perturbation experiments, data was time-aligned to the
perturbation time-point (t=0) and plotted as average ± s.d. from at least three replicates for each
condition. The change of inter-aster distance during the first 3 s after drug injection was
estimated by linear regression, and the confidence interval of the slope served as significance
test. Differences in steady-state inter-aster distance were tested by comparing the pools of
distances from the last 3 s, using Wilcoxon signed-rank test. A significance level of 0.05 was
defined prior to testing. In the case of UV ablations, the position of the aster five frames before
ablation was defined as coordinate origin. The two main axes of the ellipsoid, along which the
pulsed ablation was performed, defined the cartesian coordinate system. A displacement vector
of the current aster position relative to the origin was calculated for each time point. The mean
and standard deviation of axial (Δx) and lateral (Δy) displacement was plotted in time (Fig. 5b-
c).

Analysis of Nuclei Internalisation in Embryos – angle probability distributions from cuts

Embryos expressing H2Av::mCherry were segmented using level sets and watershed algorithms
in MATLAB®. Regions of low nuclear density were identified as pixels that were positioned
greater than 20% of the average nucleus separation from the nearest nucleus. The centre of mass
of the low-density region was identified. The division angle orientation φ of the neighbouring
nuclei was measured relative to the centre of mass. Therefore, a nucleus dividing directly into
the region of low density would be assigned an angle of 0º, and a nucleus dividing perpendicular
to the region would be assigned an angle 90º. As the division does not have a preferred direction,
the angle range is between 0º and 90º. A similar analysis was performed for the laser ablations,

where the centre of the low-density region (artificially generated by ablating nuclei) was used to

determine the relative angle of the division axis for the neighbouring nuclei.

References

1. Lecuit, T. & Wieschaus, E. Polarized Insertion of New Membrane from a Cytoplasmic
Reservoir during Cleavage of the Drosophila Embryo. *J Cell Biol* **150**, 849–860 (2000).

2. Foe, V. E., Field, C. M. & Odell, G. M. Microtubules and mitotic cycle phase modulate
spatiotemporal distributions of F-actin and myosin II in Drosophila syncytial blastoderm
embryos. *Development* **127**, 1767–87 (2000).

3. Kanesaki, T., Edwards, C. M., Schwarz, U. S. & Grosshans, J. Dynamic ordering of nuclei
in syncytial embryos: a quantitative analysis of the role of cytoskeletal networks. *Integr Biol* **3**,
1112–1119 (2011).

4. Deneke, V. E., Melbinger, A., Vergassola, M. & Di Talia, S. Waves of Cdk1 Activity in S
Phase Synchronize the Cell Cycle in Drosophila Embryos. *Dev Cell* **38**, 399–412 (2016).

5. Petkova, M. D., Tkačik, G., Bialek, W., Wieschaus, E. F. & Gregor, T. Optimal Decoding of
Cellular Identities in a Genetic Network. *Cell* **176**, 844-855.e15 (2019).

6. Dassow, G. von & Schubiger, G. How an actin network might cause fountain streaming and
nuclear migration in the syncytial Drosophila embryo. *J Cell Biol* **127**, 1637–1653 (1994).

7. Valdés-pérez, R. E. & Minden, J. S. Drosophila melanogaster syncytial nuclear divisions are
patterned: time-lapse images, hypothesis and computational evidence. *J Theor Biol* **175**, 525–
532 (1995).

8. Kaiser, F. *et al.* Mechanical Model of Nuclei Ordering in Drosophila Embryos Reveals
Dilution of Stochastic Forces. *Biophys J* **114**, 1730–1740 (2018).

9. Deneke, V. E. *et al.* Self-Organized Nuclear Positioning Synchronizes the Cell Cycle in
Drosophila Embryos. *Cell* **177**, 925-941.e17 (2019).

10. Postner, M. A., Miller, K. G. & Wieschaus, E. F. Maternal effect mutations of the sponge
locus affect actin cytoskeletal rearrangements in Drosophila melanogaster embryos. *J Cell Biol*
119, 1205–1218 (1992).

11. Dutta, S., Djabrayan, N. J.-V., Torquato, S., Shvartsman, S. Y. & Krajnc, M. Self-similar
dynamics of nuclear packing in the early Drosophila embryo. *Biophys J* **117**, 743–750 (2019).

12. Lv, Z. *et al.* The Emergent Yo-yo Movement of Nuclei Driven by Cytoskeletal Remodeling
in Pseudo-synchronous Mitotic Cycles. *Curr Biol* **30**, 1–10 (2020).

13. Foe, V. E. & Alberts, B. M. Studies of nuclear and cytoplasmic behaviour during the five
mitotic cycles that precede gastrulation in Drosophila embryogenesis. *J Cell Sci* **61**, 31–70
(1983).

14. Krzic, U., Gunther, S., Saunders, T. E., Streichan, S. J. & Hufnagel, L. Multiview light-
sheet microscope for rapid in toto imaging. *Nat Methods* **9**, 730–3 (2012).

15. Zalokar, M. & Erk, I. Division and migration of nuclei during early embryogenesis of
Drosophila melanogaster. *J Microscopie Biol Cell* **25**, 97–106 (1976).

16. Hatanaka, K. & Okada, M. Retarded nuclear migration in Drosophila embryos with
aberrant F-actin reorganization caused by maternal mutations and by cytochalasin treatment.
Development **111**, 909–20 (1991).

17. Baker, J., Theurkauf, W. & Schubiger, G. Dynamic changes in microtubule configuration
correlate with nuclear migration in the preblastoderm Drosophila embryo. *J Cell Biol* **122**,
113–121 (1993).

18. Callaini, G. & Riparbelli, M. G. Centriole and centrosome cycle in the early Drosophila
embryo. *J Cell Sci* **97**, 539–43 (1990).

19. Tanimoto, H., Sallé, J., Dodin, L. & Minc, N. Physical forces determining the persistency
and centring precision of microtubule asters. *Nat Phys* **14**, 848–854 (2018).

20. Mitchison, T. *et al.* Growth, interaction, and positioning of microtubule asters in extremely
large vertebrate embryo cells. *Cytoskeleton* **69**, 738–750 (2012).

21. Paz, J. & Lüders, J. Microtubule-Organizing Centers: Towards a Minimal Parts List.
Trends Cell Biol **28**, 176–187 (2018).

22. Telley, I. A., Gáspár, I., Ephrussi, A. & Surrey, T. Aster migration determines the length
scale of nuclear separation in the Drosophila syncytial embryo. *J Cell Biol* **197**, 887–895
(2012).

23. Basto, R. *et al.* Flies without Centrioles. *Cell* **125**, 1375–1386 (2006).

24. Vaizel-Ohayon, D. & Schejter, E. D. Mutations in centrosomin reveal requirements for
centrosomal function during early Drosophila embryogenesis. *Curr Biol* **9**, 889–898 (1999).

25. Megraw, T. L., Li, K., Kao, L. R. & Kaufman, T. C. The centrosomin protein is required
for centrosome assembly and function during cleavage in Drosophila. *Development* **126**, 2829–
39 (1999).

26. Telley, I. A., Gáspár, I., Ephrussi, A. & Surrey, T. A single Drosophila embryo extract for
the study of mitosis ex vivo. *Nat Protoc* **8**, 310–324 (2013).

27. Freeman, M., Nüsslein-Volhard, C. & Glover, D. M. The dissociation of nuclear and
centrosomal division in gnu, a mutation causing giant nuclei in Drosophila. *Cell* **46**, 457–468
(1986).

28. Lee, L. A., Hoewyk, D. V. & Orr-Weaver, T. L. The Drosophila cell cycle kinase PAN GU
forms an active complex with PLUTONIUM and GNU to regulate embryonic divisions. *Gene*
Dev **17**, 2979–2991 (2003).

29. Grill, S. W., Howard, J., Schäffer, E., Stelzer, E. H. K. & Hyman, A. A. The Distribution of
Active Force Generators Controls Mitotic Spindle Position. *Science* **301**, 518–521 (2003).

30. Wühr, M., Dumont, S., Groen, A. C., Needleman, D. J. & Mitchison, T. J. How does a
millimeter-sized cell find its center? *Cell Cycle* **8**, 1115–1121 (2009).

31. Bak, P., Tang, C. & Wiesenfeld, K. Self-organized criticality: An explanation of the 1/ f
noise. *Phys Rev Lett* **59**, 381–384 (1987).

32. Hughes, D. & Paczuski, M. Large Scale Structures, Symmetry, and Universality in
Sandpiles. *Phys Rev Lett* **88**, 054302 (2001).

33. Gennes, P. G. de, Prost, J. & Pelcovits, R. The Physics of Liquid Crystals. *Phys Today* **48**,
70–71 (1995).

34. Chakraborty, A. & Manna, S. S. Space-filling Percolation. *Phys Rev E* **89**, 032103 (2014).

35. Jo, W. S., Yi, S. D., Baek, S. K. & Kim, B. J. Cluster-size heterogeneity in the two-
dimensional Ising model. *Phys Rev E* **86**, 032103 (2012).

36. Moessner, R. & Chalker, J. T. Properties of a Classical Spin Liquid: The Heisenberg
Pyrochlore Antiferromagnet. *Phys Rev Lett* **80**, 2929–2932 (1998).

37. Freeman, M. & Glover, D. M. The gnu mutation of Drosophila causes inappropriate DNA
synthesis in unfertilized and fertilized eggs. *Gene Dev* **1**, 924–930 (1987).

38. Holy, T. E., Dogterom, M., Yurke, B. & Leibler, S. Assembly and positioning of
microtubule asters in microfabricated chambers. *PNAS* **94**, 6228–6231 (1997).

39. Dogterom, M. & Yurke, B. Measurement of the Force-Velocity Relation for Growing
Microtubules. *Science* **278**, 856–860 (1997).

40. Bieling, P., Telley, I. A. & Surrey, T. A Minimal Midzone Protein Module Controls
Formation and Length of Antiparallel Microtubule Overlaps. *Cell* **142**, 420–432 (2010).

41. Subramanian, R., Ti, S.-C., Tan, L., Darst, S. A. & Kapoor, T. M. Marking and Measuring
Single Microtubules by PRC1 and Kinesin-4. *Cell* **154**, 377–390 (2013).

42. Jeune-Smith, Y. & Hess, H. Engineering the length distribution of microtubules
polymerized in vitro. *Soft Matter* **6**, 1778–1784 (2010).

43. Howard, J. *Mechanics of motor proteins and cytoskeleton*. (Sinauer Associates), 2001,
ISBN 978-0878933334.

44. Forth, S., Hsia, K.-C., Shimamoto, Y. & Kapoor, T. M. Asymmetric Friction of Nonmotor
MAPs Can Lead to Their Directional Motion in Active Microtubule Networks. *Cell* **157**, 420–
432 (2014).

45. Laan, L. *et al.* Cortical dynein controls microtubule dynamics to generate pulling forces
that position microtubule asters. *Cell* **148**, 502–14 (2012).

46. Zhu, J., Burakov, A., Rodionov, V. & Mogilner, A. Finding the cell center by a balance of
dynein and myosin pulling and microtubule pushing: a computational study. *Mol Biol Cell* **21**,
4418–27 (2010).

47. Kinneret, K. Non Equilibrium Steady State Dynamics of Contractile Actin Networks.
Biophys J **110**, 174a (2016).

48. Rogers, S. L., Rogers, G. C., Sharp, D. J. & Vale, R. D. Drosophila EB1 is important for
proper assembly, dynamics, and positioning of the mitotic spindle. *J Cell Biol* **158**, 873–884
(2002).

49. Robinson, J. T., Wojcik, E. J., Sanders, M. A., McGrail, M. & Hays, T. S. Cytoplasmic
Dynein Is Required for the Nuclear Attachment and Migration of Centrosomes during Mitosis
in Drosophila. *J Cell Biol* **146**, 597–608 (1999).

50. Nestor-Bergmann, A., Goddard, G. & Woolner, S. Force and the spindle: mechanical cues
in mitotic spindle orientation. *Semin Cell Dev Biol* **34**, 133–9 (2014).

51. Inoue, Y. H. *et al.* Mutations in orbit/mast reveal that the central spindle is comprised of
two microtubule populations, those that initiate cleavage and those that propagate furrow
ingression. *J Cell Biol* **166**, 49–60 (2004).

52. Schubiger, G. & Edgar, B. Using inhibitors to study embryogenesis. *Methods Cell Biol* **44**,
697–713 (1994).

53. de-Carvalho, J., Deshpande, O., Nabais, C. & Telley, I. A. A cell-free system of Drosophila
egg explants supporting native mitotic cycles. *Methods Cell Biol* **144**, 233–257 (2018).

54. Tinevez, J.-Y. *et al.* TrackMate: An open and extensible platform for single-particle
tracking. *Methods* **115**, 80–90 (2016).

55. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat
Methods* **9**, 676–682 (2012).

56. Saw, T. B. *et al.* Topological defects in epithelia govern cell death and extrusion. *Nature*
544, 212–216 (2017).

57. Menzel, A. M. & Ohta, T. Soft deformable self-propelled particles. *EPL* **99**, 58001 (2012).

777
778

Supplementary Figures

Supplementary Fig. 1 – Timeline of average inter-nuclear and inter-aster distances. Averages from five blastoderm embryos during cycles 10, 11, 12 and 13. Each cycle is characterised by late interphase (late IP), anaphase A (APA), anaphase B (APB), telophase (TP) and early interphase (early IP). Morphological criteria for identification of mitotic phases and hierarchical classification of distances; *Nuclear-based*: D – non-sister nuclei identified as nearest neighbours. $\bar{D}$ – sister chromatids/nuclei. *Aster-based*: s – sister asters ~ spindle length. $\bar{s}$ – sisters in the following mitotic cycle, i.e, s becomes $\bar{s}$ after early interphase of the ensuing mitotic cycle. d – non-sister asters identified as nearest neighbours (distribution data from cycle 10 is presented in Fig. 1, details on the measurements in Methods).

Supplementary Fig. 2 – Scaling of chain length probability with chain length for different
inclusion criteria. (a) Schematic of geometric considerations in 2D of neighbour spindle
alignment. Spindle axes are in cyan. Left shows almost parallel division axes (θ small),
orientated perpendicular to the vector between the two spindles (ϕ almost 90º). Right shows
almost colinear alignment (θ small, ϕ small). The spindle does not have a specific vector
direction, so angles are between 0º and 90º. A chain is defined by spindles fulfilling one or two
of the following conditions; (i) $0 \leq \theta \leq \Delta\theta$ (ii) $\Delta\phi \leq \phi \leq 90°$ (Methods). **(b)** Representative
embryo in n.c. 10, 11, 12 and 13 showing alignment chains. Solid purple lines denote nearest
neighbours that satisfy the two chain conditions with angles $\Delta\phi = 45°$ and $\Delta\theta = 45°$. Thin
dashed red lines denote nearest neighbours that satisfy only chain condition (i). **(c)** Logarithmic
plots of probability as a function of chain length for different chain conditions. Filled symbols
denote experimental data and open symbols are from simulations of randomised orientations.
Lines represent a fit to $\beta C^{-\alpha}$, where C is the chain length and α is the scaling exponent (fitting
performed with *fit* in MATLAB®); solid lines are fits to experimental data and dashed lines are
fits to simulated data of randomised orientations. We generated 1000 random orientations for
each embryo at each cycle.

Supplementary Fig. 3 – Division orientation of three spindles in a cytosolic explant, and

influence of effective temperature. (a) 3D plot of the shortest distance (black solid line)

between three spindle centres (open circles). The spindle orientations (dashed lines) and

positions of associated asters (filled circles) illustrate the adjustment of orientation and position

in this simple system. **(b)** Histogram of the angle ϕ between spindle axis and the connecting line

(inset) for the three-spindle scenario from experimental (blue, n=7) and *in silico* (magenta) data

($J/k_BT = 10^{-1}$). **(c)** Cumulative probability of relative division axis for the 3-spindle scenario and

randomised orientation. p-value was determined from Kolmogorov–Smirnov test. **(d, e)**

Cumulative probabilities of spindle axes orientation for two-, three- and four-spindle scenarios

at different effective simulation temperature. Dashed lines are model predictions with either low

(d) or high **(e)** effective temperature (thermal noise).

Explant

a Steady-state distance between the two asters

b Boundary distance of each asters in a two-aster scenario

c Boundary distances in the two-aster scenario

Embryo

d Aster positioning in the *gnu* mutant embryo

e Steady state inter-aster distance in the *gnu* embryo

Supplementary Figure 4 – Distance analysis of asters in explants and embryos in the absence of nuclei. (a) Scatter plot of shortest distance to explant boundary (b_1 and b_2) as a function of the radius R in explants containing two asters (n=54; details in inset of Fig. 3d). The magenta line represents the linear regression. Black dashed lines represent half and full radius distance. **(b)** Scattered plot of inter-aster distance (d_a) as a function of the radius (R) of explants containing two asters (n=54). Most measured data points fall between the dashed lines denoting the explant radius ($Y = R$) and half of the radius ($Y = \frac{1}{2}R$). The magenta line represents the linear regression. **(c)** Correlation plot of the boundary distances b_1 and b_2 referred to in panel (**a**) with Pearson's correlation coefficient r. **(d)** Maximum intensity Z-projection of a *gnu* mutant embryo expressing RFP::β-Tubulin (grey) and Spd2::GFP (green) (left, scale bar 20 µm), with magnification of three asters (right, scale bar 10 µm). **(e)** Schematic of the measurement of inter-aster distance d between nearest neighbour asters (left), and scatter plots of d_a during consecutive intervals of 10 min for the same embryo.

Supplementary Figure 5 – Analysis of migration dynamics in explants containing a single aster. (a) Correlation between explant radius R and final aster distance to the explant boundary after reaching a steady state (n=7 for **a-d**). The solid line represents the linear regression to experimental data. **(b)** Trajectories of normalised aster distance to the explant boundary from independent experiments. **(c)** Migration velocity as a function of boundary distance. **(d)** Aster separation velocity as a function of time, where t=0 is defined as the time when the aster lies midway between the explant edge and the final position of the aster. Solid line represents average over all measurements. **(e)** Z-projection of a 3D image stack of a small explant containing one aster that exemplifies microtubule buckling and splay near the explant boundary (scale bar 5 μm). **(f)** Velocity field of yolk droplets around the single aster as it escapes from the explant boundary. Image rotation and frame matching were performed to overlay all experiments such that the aster, at each time point is positioned at (0,0) and moves in the direction (0,-1). Track colour coding denotes displacement angle, where turquoise corresponds to 0° and red to 90°. **(g)** Velocity profile of asters (black) and 35 lipid droplet (magenta) tracks within a distance of 20 μm from the aster, where t=0 is the time point at which the aster starts moving away from the explant boundary. **(h)** Root-mean-square (RMS) displacement of lipid droplets in the explants. Droplet movement analysed with (black) and without (magenta, n=6) an aster. Error bars s.e.m. and lines represent fits to models shown in legend. Measurements in explants with aster fit best to a model exhibiting some directionality, see Methods for further discussion.

a final inter-aster distance [μm] vs droplet radius R [μm]

$r^2 = 0.84$

b normalised aster separation vs time [s]

$0.52 + 0.44 \tanh{(t/193)}$

c separation velocity [μm/s] vs aster separation (μm)

d separation velocity [μm/s] vs time [s]

e normalised microtubule intensity vs aster separation [μm]

$0.16 + 0.56 e^{-x/9.8}$

f RFP::βTub
Spd2::GFP

g Normalised microtubule intensity vs perpendicular distance from aster midpoint [μm]

Supplementary Figure 6 – Analysis of separation dynamics in explants containing two
asters. (a) Correlation between explant radius R and the final distance between asters after
reaching a steady state (n=9). The solid line represents the linear regression to experimental data.
(b) Normalised aster separation distance versus time, where t=0 is defined as the time when the
aster lies midway between the explant edge and the final position of the aster. Solid line
represents the fit to a hyperbolic tangent function (n=9). **(c)** Aster separation velocity as a
function of aster separation, as defined above. Each colour corresponds to a different experiment
(n=8). **(d)** Aster separation velocity versus time, where t=0 is defined as above. Solid line denotes
the fitting to the time derivative of the *tanh* function shown in **(b)** (n=9). **(e)** Normalised
microtubule intensity at the midpoint perpendicular axis between asters in function of aster
separation distance. Open markers denote average values and error bars the standard deviation.
Solid line represents the fitting to exponential decay (n=7), fitting performed using MATLAB
fit function. **(f)** Two colour maximum intensity Z-projection of two separating asters in an
explant with fluorescent reporters for Spd2::GFP (green) and RFP::β-Tubulin (magenta). A void
of microtubules in the centre between two separating asters is observed during the initial
acceleration phase. Scale bar, 10 μm. **(g)** Normalised microtubule intensity along the
perpendicular midpoint axis of separating aster during the initial acceleration phase. Solid line
denotes average normalised (for each experiment, as discussed in Methods) microtubule density
and dashed lines ±1s.d. (n=6). There was significant variation between samples and the
microtubule signal was often weak, making a detailed analysis challenging.

Supplementary Figure 7 – Analysis of nuclei division orientation *in vivo*

(a) (i) Embryo expressing H2Av::mCherry just prior to anaphase B of n.c. 13 . (ii) Areas of low nuclear density were identified by finding pixels with no nuclei within 20% of the average nuclear separation and the centroid of these areas found (red diamonds). (iii) The nuclei (red squares) neighbouring the low density region were then identified. **(b)** Examples of nuclear division near a region of low nuclear density. The division axis orientation of the neighbouring nuclei to a region of low density was measured (red bars) and the subsequent position of the

nuclei after division identified (turquoise squares). Change in area of low nuclear density region from before (blue) to after (red) area is shown. **(c)** Analysis performed as in Fig. 5g, but for arbitrary locations selected within the embryo. **(d)** Artificial generation of a region of low nuclear density by ablating a nucleus (Methods). Black arrow in top left panel identifies ablated nucleus; ablation at time t=0 min. Images from maximum intensity projection of embryo expressing H2Av::mCherry. Red arrows at 8 min denote division axis orientation of neighbouring nuclei to the ablated nucleus. **(e)** Similar to **(d)**, except three nuclei are ablated to generate a larger region of low nuclear density. **(f)** Rose plots of division axis orientation for nuclei adjacent to ablated nuclei, for single (left column) and multi-nuclei (right column) ablations.

Supplementary Video Legends

Supp. Video 1: Maximum intensity Z-projection from a 3D time-lapse movie of an embryo expressing H2Av::RFP (magenta) and Spd2::GFP (green) from interphase of n.c. 10 until the end of n.c. 13 (time in min:sec, scale bar 20 μm). In support of Fig. 1.

Supp. Video 2: Maximum intensity Z-projection from a 3D time-lapse movies of three distinct cycling explants starting with a single spindle extracted from embryos expressing Jupiter::GFP (grey) and H2Av::RFP (magenta) (time in min:sec, scale bar 10 μm). In support of Fig. 2.

Supp. Video 3: Maximum intensity Z-projection from a 3D time-lapse movie of a *gnu* mutant embryo, approximately 1h old embryo expressing β-Tubulin::EGFP (grey) (time in hr:min:sec, scale bar 10 μm). In support of Fig. 3.

Supp. Video 4: Maximum intensity Z-projection from a 3D time-lapse movie of an explant generated from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green), containing a single aster moving away from the explant boundary (time in min:sec, scale bar 10 μm). In support of Fig. 3 and 4.

Supp. Video 5: Maximum intensity Z-projection from a 3D time-lapse movie of a *gnu* mutant embryo, approximately 4h old embryo expressing β-Tubulin::GFP (grey) (time in min:sec, scale bar 10 μm). In support of Fig. 3.

Supp. Video 6: Maximum intensity Z-projection from a 3D time-lapse movie of an explant generated from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green), containing two separating asters (time in min:sec, scale bar 10 μm). In support of Fig. 4.

Supp. Video 7: Maximum intensity Z-projection from a 3D time-lapse movie of explants generated from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green), containing two separating asters, after pulse injection of solutions: control with buffer (left), 10 mM sodium azide (centre) and 0.2 mM of colchicine (time in min:sec, scale bar 10 μm). In support of Fig. 4.

Supp. Video 8: Maximum intensity Z-projection from a 3D time-lapse movie of explants generated from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green) containing a single aster (time in min:sec, scale bar 10 μm). The aster was allowed to equilibrate followed by an asymmetric elliptic ablation (yellow line at times 00:15 to 01:00) performed in control explants (no injection) and in explants supplemented with 0.2mM of colchicine. In support of Fig. 5.

Supp. Video 9: Maximum intensity Z-projection from a 3D time-lapse movie of an explant containing two separating asters from a *gnu* mutant embryo expressing RFP::β-Tubulin (magenta) and Spd2::GFP (green) (time in min:sec, scale bar 10 μm). The elliptic ablation (yellow line from 00:15 to 00:45) was performed when asters were ~7 μm apart. In support of Fig. 5.

Supp. Video 10: Maximum intensity Z-projection from a 3D time-lapse movie of a wildtype embryo expressing H2Av::mCherry (magenta) and Spd2::GFP (green) during n.c. 12 and 13, in response to spontaneous nuclear internalisation (time in min:sec, scale bar 20 μm). In support of Fig. 5.