# Acquisition of ionic copper by a bacterial outer membrane protein

Satya Prathyusha Bhamidimarri[1], Tessa R. Young[2], Muralidharan Shanmugam[3], Sandra Soderholm[4], Bastien Belzunces[5], Arnaud Baslé[1], Chris Skylaris[5], Dirk Bumann[4], Syma Khalid[5], Bert van den Berg[1#]

[1]Biosciences Institute, The Medical School, Newcastle University, Newcastle upon Tyne NE2 4HH, UK

[2]Department of Biosciences, Durham University, DH1 3LE Durham, UK.

[3]Manchester Institute of Biotechnology, University of Manchester, 131 Princess Street, Manchester M1 7DN, UK

[4] Focal Area Infection Biology, University of Basel, CH-4056 Basel, Switzerland

[5]School of Chemistry, University of Southampton, Southampton SO17 1BJ, UK

[#] Corresponding author: bert.van-den-berg@ncl.ac.uk

**Abstract**

Copper, while toxic in excess, is an essential micronutrient in all kingdoms of life due to its essential role in the structure and function of many proteins. Proteins mediating ionic copper import are known in eukaryotes, but have not yet been described in prokaryotes. Here we show that *Pseudomonas aeruginosa* OprC is a TonB-dependent transporter that mediates acquisition of ionic copper. Crystal structures of wild type and mutant OprC variants with silver and copper, as well as ICP-MS and electron paramagnetic resonance (EPR), suggest that binding of Cu(I) occurs via a surface-exposed methionine track leading towards an unprecedented CxxxM-HxM metal binding site that binds Cu(I) directly and can facilitate reduction of Cu(II) via the cysteine thiol. Together with quantitative proteomics and growth assays, our data identify OprC as an abundant component of bacterial copper biology that enables copper acquisition and potentially copper storage under a wide range of environmental conditions.

**Introduction**

Metals fulfil cellular functions that cannot be met by organic molecules and are indispensable for the biochemistry of life in all organisms. Copper is the third-most abundant transition metal in biological systems after iron and zinc. It has key roles as structural component of proteins or catalytic cofactor for enzymes[1], most notably associated with the biology of oxygen and in electron transfer. On the other hand, an excess of copper can be deleterious due to its ability to catalyse production of hydroxyl radicals[2, 3]. Excessive copper may also disrupt protein structure by interaction with the polypeptide backbone, or via replacement of native metal cofactors from proteins, thus abolishing enzymatic activities via mismetallation[4, 5]. Thus, cellular copper levels and availability must be tightly controlled. Bacterial copper homeostasis systems are well characterised[6]. Specific protein machineries are involved in fine-tuning the balance of intracellular copper trafficking, storage and efflux according to cellular requirement, in such a way that copper is always bound inside the cell. This control is executed by periplasmic and cytosolic metalloregulators, which activate transcription of periplasmic multi-copper oxidases, metallochaperones, copper-sequestering proteins[7, 8] and transporters[9-11]. To date, relatively few families of integral membrane proteins have been validated as copper transporters, and these have different structures and transport mechanisms[12]. The $P_{1B}$-type ATPases such as CopA are responsible for Cu(I) efflux from the cytosol via several metal binding domains, using energy released from ATP hydrolysis[13-15]. A second class of copper export proteins are RND-type tripartite pumps such as CusABC, which efflux Cu(I) by utilising the proton-motive force[16-18]. Relatively few copper influx proteins have been identified. The inner membrane copper importer CcoA is a major facilitator superfamily (MFS)-type

transporter involved in fine-tuning the trafficking of copper into the cytosol and required for
cytochrome c oxidase maturation[19, 20]. The Ctr family of copper transporters is responsible for
Cu(I) translocation into the cell without requiring external sources of energy[21]. However, Ctr
homologs are found only in eukaryotes, and the molecular mechanisms by which copper ions
enter bacteria is still a matter of debate. The exception is copper import via metallophores like
methanobactin, a small Cu-chelating molecule that is secreted by methanotropic bacteria and
most likely taken up via TonB-dependent transporters, analogous to iron-siderophores[22].
80
*Pseudomonas aeruginosa* is a versatile and ubiquitous Gram-negative bacterium that
promotes infections in several hosts and multiple sites. It is a notorious opportunistic pathogen
in humans, and is mainly responsible for the development of chronic lung infection in cystic
fibrosis patients[23, 24]. *P. aeruginosa* has a number of TonB-dependent transporters (TBDTs) in
the outer membrane (OM) dedicated to the acquisition of different iron-siderophore complexes
such as pyochelin and pyoverdin[25]. In addition, *P. aeruginosa* contains another TBDT, termed
OprC (*PA3790*), whose function has remained enigmatic. Nakae *et al*. suggested that OprC
binds Cu(II) with micromolar affinities[26]. Transcription of OprC was recently found to be
repressed in the presence of Cu(II) in the external medium under aerobic conditions[27],
suggesting a role for OprC in copper acquisition. Very recently, the blue copper protein azurin
was reported to be secreted by a *P. aeruginosa* Type VI secretion system and to interact with
OprC, suggesting a role of the latter in Cu(II) uptake[27].
93
To clarify the role of OprC in copper biology, we have determined X-ray crystal structures of
wild type and mutant OprC proteins in the absence and presence of copper and silver, and
characterised metal binding via ICP-MS, EPR and quantum chemistry calculations. OprC
indeed has the typical structure of a TBDT, and differences between the Cu-loaded and Cu-
free protein demonstrate changes in tertiary structure that likely lead to TonB interaction.
Strikingly, metal binding experiments show that while OprC can bind to both Cu (I) and Cu (II)
*in vitro*, only Cu (I) is present at the principal CxxxM-HxM metal binding site, suggesting that
Cu(I) is the imported redox state and that reduction of Cu(II) may be mediated by OprC.

**Figure 1. OprC is a TonB-dependent transporter.** Cartoon representation of (**a,c**) Cu-loaded OprC and (**b,d**) Cu-free OprC (OprC$_{AA}$). The N-terminal plug domain is shown separately for both forms (**e,f**). Structures are shown in rainbow from N-terminus (blue) to C-terminus (red); copper is represented as a magenta sphere. The arrow in (**f**) highlights the visibility of the Ton box in apo-OprC.

## Results

### OprC is a TonB-dependent transporter that binds ionic copper

The structure of OprC crystallised under aerobic conditions in the presence of 2 mM CuCl$_2$ was solved using single wavelength anomalous dispersion (SAD), using data to 2.0 Å resolution (Methods; Supplementary Table 1; Supplementary Fig. 1). As indicated by the successful structure solution, OprC contains bound copper and shows the typical fold of a TBDT, with a large 22-stranded β-barrel occluded by an N-terminal ~15 kDa plug domain (Fig. 1). The copper binding site comprises residues Cys143 and Met147 in the plug domain and His323 and Met325 in the barrel wall. The CxxxM-HxM configuration, which coordinates the copper in a tetrahedral manner (Fig. 2a, b), is highly unusual and has, to our knowledge, not been observed before in copper homeostasis proteins. The most similar site is present for one of the copper ions of the valence-delocalised Cu$_A$ dimer in cytochrome c oxidase, where the

copper ion is coordinated by 2Cys+1Met+1His[28, 29]. The other relevant sites are class I Type I
copper proteins like pseudoazurin and plastocyanin, where copper is coordinated by
2His+1Cys+1Met[30]. Interestingly, and unlike class I Type I copper proteins, concentrated
solutions and OprC crystals are colourless in the presence of Cu(II). Another notable feature
of the OprC structure becomes apparent when analysing the positions of the methionine
residues. As shown in Fig. 2d, out of the 15 visible methionines in OprC, 10 are organised in
such a way that they form a distinct "track" leading from the extracellular surface towards the
copper binding site. An additional two methionines are not visible due to loop disorder, but
given their positions (Met448, Met558) they will be a part of the methionine track. We propose
that the methionine track serves to bind copper with low affinity and may guide the metal
towards the principal binding site. It should be noted that the anomalous difference maps of
OprC crystallised in the presence of Cu(II) do not show any evidence for weaker, secondary
copper sites (Supplementary Fig. 1).

135
**Figure 2. OprC has an unusual CxxxM-HxM binding site and a methionine binding track.**
**a,** Stick models of copper-coordinating residues Cys143, Met147, Met325 and His323.
Electron density in gray mesh (2Fo-Fc map contoured at 2.0σ, carve = 2.0) is shown for the
binding site residues C/M-H/M and the copper atom (anomalous difference map shown in

magenta, contoured at 3.0σ, carve = 2.25). **b,** Distances between coordinating residues and
metal show that copper is coordinated via 1 thiolate (from Cys), two thioethers (from Met), and
one imidazole nitrogen from His. **c,** Mutation of binding site residues Cys143 and Met147 to
alanines abolishes copper binding (2Fo-Fc map contoured at 2.0σ, carve = 2.0). **d,e,** OM plane
**(d)** and extracellular views **(e)** showing the thioether atoms of all methionine residues present
in OprC as yellow spheres.

**Copper binding by OprC is highly specific and near-irreversible.**

Following structure determination of copper-bound OprC, several attempts were made to
produce a structure of copper-free OprC. First, the protein was purified and crystallised without
added copper; this gave a structure that was identical to the one already obtained and
contained bound copper that presumably originated from the LB medium. As expression in
rich media always yielded OprC with ~1 equivalent copper as judged by ICP-MS, various
attempts to lower the copper content were made. Removal of bound copper from purified
protein with combinations of denaturants (up to 4.0 M urea) and EDTA were not successful.
Expression in chelex-treated minimal medium reduced, in the best cases, the metal content
of the wild type to ~45 % equivalency as assessed by ICP-MS (Fig. 3a). Subsequent
incubation of OprC for 30 min in the presence of either 3 or 10 equivalents Cu(II) followed by
size exclusion chromatography (SEC) and ICP-MS analysis of the peak fractions leads to co-
elution and occupancy of 1 equivalent copper (Fig. 3a). However, co-incubation with 1 mM
EDTA (~100-fold excess) does not result in copper loading, suggesting that EDTA effectively
withholds copper from OprC (Supplementary Fig. 2). As-purified OprC does not contain zinc,
a common contaminant in metal-binding proteins, indicating that OprC is highly specific.
Indeed, incubation of purified OprC in the presence of 3 or 10 equivalents Zn does not result
in zinc co-elution (Supplementary Fig. 2). To obtain copper-free OprC, we constructed a
variant in which the binding site residues Cys143 and Met147 were both mutated to alanines,
generating copper-free protein (OprC$_{AA}$) after purification from rich media (Fig. 3a; AA). Even
after equilibration of OprC$_{AA}$ for 30 min with 3 or 10 equivalents Cu(II) no co-elution with metal
is observed (Fig. 3a), indicating that high-affinity copper binding is completely abolished. It
should be noted that OprC$_{AA}$ contains a His-tag, demonstrating that Cu(II) does not bind to the
His-tag of OprC with an affinity high enough to survive SEC, possibly due to the presence of
0.5 mM EDTA in the SEC buffer.

The fact that it is not possible to obtain copper-free wild type protein, even after taking
extensive precautions, suggests that copper binds to OprC with very high affinity. To explore
this further, we performed a copper extraction assay with a large excess of
bathocuproinedisulfonic acid (BCS) under reducing conditions (Methods). BCS is a high-
affinity Cu(I) chelator (logβ$_2$ 20.8) and forms a 2:1 complex with Cu(I), namely [Cu(BCS)$_2$]$^{-3}$,
with a molar extinction coefficient of 13,300 cm$^{-1}$ M$^{-1}$ at 483nm, enabling quantitation of Cu(I).

For copper-loaded OprC$_{WT}$, only 20% copper was removed after 24 hrs at room temperature,
and the temperature had to be increased to 60 °C to obtain near-quantitative extraction of
copper (~90% after 24 hours) (Fig. 3b). For reasons that are unclear, the orange-coloured
$[Cu(BCS)_2]^{-3}$ complex was hard to separate from OprC, and BCS-treated OprC did not bind
copper anymore, suggesting an irreversible change in the protein due to the harsh incubation
conditions. These results demonstrate that copper is kinetically trapped inside OprC and is,
for all intents and purposes, irreversibly bound. This is compatible with the consensus
transport mechanism of TBDTs, in which the interaction with TonB is thought to result in the
disruption of the binding site and subsequent release of the substrate[31].
188

**Figure 3. OprC binds to up 1 equivalent copper near-irreversibly. a,** Copper content of
wild type OprC and binding site mutant proteins before (blue) and after aerobic incubation with
either 3 (red) or 10 equivalents Cu(II) (green) for 30 min followed by analytical SEC and quan-
titation of the copper content of the OprC peak by ICP-MS. "As purified" metal contents were
obtained from MM-grown *E. coli* cells for WT OprC and from LB-grown cells for all mutant
proteins. All proteins except where stated contain a C-terminal hexa-histidine tag. The in-
crease in copper content in "as purified" tagless WT protein is most likely due to the use of a
second IMAC column after tag cleavage. Reported values are averages ± s.d. (n = 3) **b,** Cop-
per is kinetically trapped in OprC. Time course of copper extraction experiments showing %
bound copper for OprC WT (blue), OprC$_{AA}$ (green) and OprC M147A (red), at room tempera-
ture (open symbols) and 60 °C (filled symbols). The inset shows % bound copper in the first
few minutes after starting the experiment. OprC$_{AA}$ served as a control. Reported values are
averages ± s.d. (n = 3)
203

**Conformational changes upon copper binding.**

The OprC$_{AA}$ structure was solved by molecular replacement using Cu-bound OprC as the
search model (Fig.1b, d, f; Fig. 2c). The binding site residues of both structures occupy very
similar positions, indicating that the introduced mutations abolish copper binding without
generating gross changes in the binding site. Superposition of the structures (Figs. 4 a,b)
shows that for the remainder of the transporter, structural changes upon copper binding are
confined to the vicinity of the copper binding site, with parts far removed virtually unchanged

(overall Cα r.m.s.d ~ 1.0 Å). The largest change is observed for loop L11, which undergoes an

inward-directed motion of ~ 8.0 Å upon copper binding (Figs. 4 d,e and Supplementary Fig.

3). A similar inward-directed but smaller change occurs for loop L8. Some loop tips (*e.g.* L4,

L5, L6) in OprC$_{AA}$ lack electron density for a limited number of residues, suggesting increased

mobility. However, given that the proteins were crystallised in different space groups (OprC,

C222$_1$; OprC$_{AA}$ P22$_1$2$_1$), such differences could also be the result of different crystal packing.

Overall, the conformational changes of the loops decrease the accessibility of the copper

binding site. However, the main reason why the bound copper is inaccessible to solvent is

because the binding site residues Met147 and Met325, together with Asn145, effectively form

a lid on the copper ion. In the double mutant, copper becomes solvent accessible due to the

absence of the Met147 side chain (Fig. 4d and Supplementary Fig. 3).

**Figure 4. OprC structural changes upon copper binding. a,b** Cartoon superposition from
the OM plane (**a**) and extracellular environment (**b**) of OprC (coloured green) and OprC$_{AA}$
(blue), indicating locations of loops L5, L8 and L11; copper is represented as a magenta
sphere. The plug domains of OprC and OprC$_{AA}$ are coloured light green and light blue,
respectively. **c,** N-terminal plug domains indicating the location of the TonB box, which is
invisible in Cu-bound OprC. Arrows indicate the missing density for Glu88-Pro94 in OprC. **d,**
Surface slab representations from the OM plane, showing the presence of a solvent pocket in

OprC$_{AA}$ that is generated by the absence of Met147 (arrow). For orientation purposes, the
location of OprC-bound copper is shown in both structures. **e,** Side surface views showing the
conformational changes of L8 and L11 (coloured red) as a result of copper binding. As in **(d)**,
the bound copper of OprC is shown in both proteins.

The current consensus mechanism for TonB-dependent transport postulates that ligand
binding on the extracellular side generates conformational changes that are propagated to the
periplasmic side of the plug and increase the periplasmic accessibility of the Ton box for
subsequent interaction with TonB[32]. In OprC$_{AA}$, N-terminal density is visible up to Leu66 (*i.e.*,
the first 10 residues of the mature protein are disordered) including the Ton box
([68]PSVVTGV[75]), which is tucked away against the plug domain and the barrel wall. In Cu-
bound OprC, the density between Glu88 and Pro94 is hard to interpret and, more importantly,
no density is observed before Pro79, including the Ton box (Figs. 1e, f and 4c). Thus, the
structures of OprC in the absence and presence of ligand conform to the consensus transport
mechanism, in the sense that the TonB box is accessible for potential interaction with
periplasmic TonB only in the substrate-loaded protein. This would prevent non-productive
interactions with transporters that do not have substrate bound[32].

**The OprC methionine track binds Cu(I) but not Cu(II).**
Which redox state of the copper ion is present in OprC? Since it is challenging to maintain
copper in its +1 state during crystallisation, we used silver (Ag$^+$) as a proxy for Cu(I) and
determined the co-crystal structure of WT OprC in the presence of 2 mM AgNO$_3$ (Methods).
Data was collected at 8000 eV, at which energy the anomalous signal of copper is very small
(0.6 e$^-$, compared to 4.2 e$^-$ for Ag). Strikingly, and in sharp contrast to copper (Supplementary
Fig. 1), the anomalous map of OprC WT crystallised in the presence of silver shows not one
but three anomalous peaks. The first, strong peak (Ag1; 23$\sigma$) is located at the same site as in
OprC crystallised with copper, and is coordinated by the same residues (Cys143, Met147,
His323 and Met325; Fig. 5). The other two sites have lower occupancy (Ag2, ~10$\sigma$ and Ag3,
~10$\sigma$) and are each coordinated by two methionines in the methionine track (Met286 and
Met339 for Ag2; Met341 and Met343 for Ag3). This structure therefore suggests that the
methionine track provides several low-affinity binding sites for Cu(I) ions, and that the high
affinity CxxxM-HxM site binds Cu(I).

To obtain more information on the individual residue contribution to copper binding, we next
generated the complete set of single alanine mutants of the principal binding site residues
(C143A, M147A, H323A and M325A), and determined copper binding via analytical SEC and
ICP-MS. For all single mutants, the copper content after purification from LB was below 10%,
except for M147A (~20 %) (Fig. 3a). Upon incubation with 3 or 10 equivalents Cu(II), various

occupancies were obtained. C143A has no bound copper even after incubation with 10
equivalently Cu (II), suggesting this residue has a crucial role. H323A (~30%) and in particular

**Figure 5 OprC binds Cu(I). a-e**, Anomalous difference maps of (**a**) OprC wild type, (**b,d**) C143A and (**c,e**) H323A variants crystallised in the presence of (**a,d,e**) Ag or (**b,c**) Cu(II), and collected at different energies. The inset to (**a**) shows a close-up of the anomalous difference peaks (magenta) near the principal binding site in wild type OprC, with binding residues labelled and represented as stick models. Sulphurs are coloured yellow. For clarity, the metal used in co-crystallisation and the energy used for data collection are shown underneath each panel. The OprC plug domain is coloured blue.

M147A (~60%) have relatively high occupancies after copper incubation and SEC, indicating

that these residues contribute less towards binding. Of the four ligands, the M147 thioether is

the furthest away from the copper in the crystal structure (Fig. 2b), which may explain why it

contributes the least to ligand binding. Interestingly, removal of bound copper is much faster

in the M147A mutant compared to OprC$_{WT}$ (Fig. 3b), suggesting that solvent exclusion by the

M147 side chain (Fig. 4d) is the main reason why copper is kinetically trapped in the wild type

protein.

**Cu(II) is reduced by the principal binding site cysteine.**

The fact that wild type OprC and most single alanine mutants can be loaded via Cu(II)

incubation (Fig. 3a) suggests that Cu(II) is also a substrate for the transporter. However, the

complete lack of copper in the C143A mutant is intriguing (Fig. 3a) and could be explained in

two ways. If one assumes that OprC can bind Cu(II), removal of the cysteine could lead to

much lower affinity for Cu(II), so that after SEC nothing remains bound. Alternatively, if OprC

only binds Cu(I), the lack of the cysteine could prevent reduction of added Cu(II). To distinguish

between these two possibilities, we determined the crystal structures of the OprC C143A and

H323A mutants in the presence of either Cu(II) or Ag(I). For each crystal, two datasets were

collected at different energies. The first energy (8800 eV) is below the copper K-edge (8979

eV), while the second one is above the edge (9175 eV), giving small and large anomalous

signals for copper (0.5 and 3.7 e$^-$ respectively). For silver, substantial anomalous signal is

present at both energies (~3.6 e$^-$). Thus, bound copper is expected to give a strong anomalous

peak only at 9175 eV, while silver will give comparable peaks at both energies. For C143A co-

crystallised with copper, no anomalous peaks are visible at both energies (Fig. 5b), showing

a lack of bound copper. Crucially, in the presence of silver, the same three anomalous peaks

are visible as for wild type OprC (Figs. 5 a, d), strongly suggesting that the C143A mutant does

still bind Cu(I). The H323A mutant, on the other hand, still binds copper supplied as Cu(II),

and also binds silver (Figs. 5 c, e). Together, these data therefore suggest that (i) Cys143

reduces Cu(II) to Cu(I) and (ii) OprC does not bind Cu(II) specifically. To support the notion

that the principal OprC binding site is a reducing environment, we performed quantum

chemistry calculations for a 70-atom system consisting of the OprC binding site, containing

either Cu(I) or Cu(II) (Methods). The absolute energies show that the system with Cu(I) is 221

and 133 kcal/mol more stable compared to Cu(II) for the hydrogenated and dehydrogenated

forms of Cys respectively (Supplementary Table 4). Energy Decomposition Analysis (EDA) of

the Cu(I) interaction with Cys143, Met147, Met325 and His323, revealed the total interaction

energy of the four residues with Cu(I) to be 96 kcal/mol more stable when cysteine is

dehydrogenated (oxidised). The metal protein stabilisation interactions are dominated by

electrostatic interactions, with polarization and charge-transfer playing only a minor role.
Additional details are provided in the Supplementary Information.

To shed additional light on the redox state of the bound copper, continuous wave EPR (cw-
EPR) spectra were recorded on OprC$_{WT}$ and single/double mutants. As-purified OprC WT con-
taining 0.6 equivalents copper was EPR silent (Supplementary Fig. 4; WT-1), confirming the
bound species is Cu(I). All other analysed variants were EPR silent as well. Subsequently,
proteins were incubated with 0.25 equivalents of CuSO$_4$ and EPR spectra were recorded. The
EPR spectra of all the OprC variants showed the appearance of the typical EPR signals as-
sociated with a Type(II) copper centre. These EPR signals are significantly different from the
standard CuSO$_{4,}$ Cu(II) EPR signals, as inferred from the visual inspection and also from the
spectral simulations. The EPR spectra of the OprC variants show nicely resolved $^{63,65}$Cu(II)
hyperfine coupling along the parallel region, due to the interaction of an unpaired electron spin
($S$= ½) of Cu(II) with the nuclear spin of ($I$ = 3/2) of $^{63,65}$Cu nuclei, as indicated by the magenta
goal-post in Supplementary Fig. 4. The intense EPR signal observed at ~3300 G shows super-
hyperfine splittings for the OprC$_{AA}$ double mutant and OprC-M147A, arising plausibly from
$^{1}$H/$^{14}$N nuclei in close proximity of the Cu(II) ion. The intensities of the EPR signals increased
upon addition of further Cu(II) equivalents (Supplementary Fig. 4 and Supplementary Table
5). The additional Cu(II) loading caused changes in EPR line widths, which results in broad-
ening and/or disappearance of the super-hyperfine splittings for OprC M147A. The analysis of
the EPR spectra of OprC$_{WT}$ and OprC-M147A shows the presence of two type(II) Cu(II) EPR
active species ($S$ = ½ species), which are not clearly visible by the naked eye in the case of
the M147A variant. By contrast, OprC$_{AA}$ and M147H yielded EPR signals arising from single
contributing EPR active species. The spectra suggest that Cu(II) does bind to OprC, but not
at the principal binding site, as judged from the facts that (i) the spectra resembled that of a
Type II copper center [33][34, 35] and (ii) Cu(II) also binds to OprC$_{AA}$. The EPR spectra were suc-
cessfully simulated (Supplementary Fig. 5) using the spin-Hamiltonian parameters provided
in Supplementary Table 5. Interestingly, the EPR signals for both wild type OprC and the
M147A variant decrease slowly upon prolonged incubation (Supplementary Fig. 4). Collec-
tively, the EPR data suggest that Cu(II) binds to OprC at one or more sites distinct from the
principal binding site, and, via unknown routes, eventually ends up in the principal binding site,
where it is reduced to Cu(I). It is possible, and perhaps even likely, that the Cu(II) binding sites
as inferred by EPR are adventitious due to the fact that the protein is present in detergent and
has much more surface (*e.g.* the entire periplasmic face) available for binding compared to
the physiological situation, in which OprC is embedded in the OM and accepts copper from
the extracellular environment. The addition of 1-10 equivalents of hydrogen peroxide oxidant
directly to the 'as purified' OprC$_{WT}$ has no significant effect on the electronic structure of the

bound Cu(I) ion as reflected by the lack of changes in the observed EPR spectra. This is
consistent with the solvent inaccessibility of the principal copper binding site in OprC$_{WT}$.
393
To confirm that OprC binds the Cu(II) redox state *in vitro*, M147A and OprC$_{AA}$ samples were
analysed by recording intrinsic tryptophan fluorescence upon Cu(II) addition. OprC$_{AA}$ was used
as a control for which high-affinity copper binding is abolished. After excitation at 300 nm, an
emission peak at 340nm was observed in both variants. Subsequent titrations with Cu(II) so-
lution showed very similar reductions in the 340 nm emission peak in both OprC M147A and
OprC$_{AA}$ (Supplementary Fig. 6). In both cases, saturation is not reached at 3 equivalents cop-
per, suggesting that Cu(II) binds non-specifically to OprC, consistent with the position of this
metal at the top of the Irving-Williams series. Indeed, at high concentrations (~0.1 mM), OprC
precipitates in the presence of millimolar Cu(II). Since OprC has 15 tryptophan residues it is
not possible to define the locations of these binding sites.
404

**OprC is abundant in *P. aeruginosa* during infection.**

To determine the abundance of OprC in *Pseudomonas aeruginosa* UCBPP-PA14 or
*Acinetobacter baumannii* ATCC 19606 in infected lung tissues, we employed a sensitive
targeted proteomics approach with parallel reaction monitoring, on a high resolution and
accurate mass instrument with absolute quantification using heavy-isotope labeled reference
peptides (Methods). The results showed that in both mouse and rat pneumonia models, OprC
was present at 1,000 to 10,000 molecules per *P. aeruginosa* cell, making it one of the five
most abundant TonB-dependent transporters. As a comparison, the most abundant TonB-
dependent transporter, FpvA, had 8,000 to 33,000 molecules per cell. We also assessed OprC
abundance in *A. baumannii*, due to the facts that both proteins are highly similar (50%
sequence identity, Supplementary Fig. 8) and that both bacteria are important human
pathogens with a similar, low-permeability OM. In *A. baumannii*, OprC was less abundant in
mouse and rat pneumonia models (40 to 400 molecules per cell), while the most abundant
TonB-dependent transporters BfnH and BauA were present at 500 to 3'000 molecules per cell.
419

**OprC does not confer copper toxicity in *P. aeruginosa in vitro*.**

To demonstrate that OprC imports copper, we initially performed growth experiments in an *E.*
*coli ΔcopA* strain that should be more susceptible to copper stress. Cells were grown at vari-
ous external Cu(II) concentrations (0-6 mM) while expressing plasmid-based wild type OprC
or OprC$_{AA}$. Toxicity was observed in all three strains above 1 mM CuSO$_4$, but no differences
were observed between the strains. However, the presence of abundant porins in *E. coli*, to-
gether with the high copper concentrations, make porin-mediated copper influx likely, and this
would mask any contribution from OprC. In addition, it is also not clear whether *E. coli* TonB

couples to *P. aeruginosa* OprC. Given that OprC is highly abundant and is expressed during

anaerobiosis[26, 36, 37], we next investigated whether OprC causes *in vitro* copper or silver toxicity

in *P. aeruginosa* under anaerobic conditions. In rich media, excessive copper concentrations

(mM) are required to cause toxicity, and no differences are observed for PA14 wild type and

Δ*oprC* strains (Supplementary Fig. 7), in line with recent data showing that *oprC* is repressed

in the presence of mM external copper under aerobic conditions[27]. We next assessed minimal

media, in which toxicity was observed at μM concentrations of copper (Fig. 6). However, PA14

wild type and Δ*oprC* strains still had similar phenotypes, and only occasionally did we observe

better growth for the Δ*oprC* strain over a very narrow range of copper concentrations (Supple-

mentary Fig. 7c, d). Silver causes toxicity at 10-fold lower concentrations compared to copper,

but again, no clear differences in growth were observed between wild type PA14 and the

Δ*oprC* strain (Fig. 6). One reason for the lack of robust OprC-mediated toxicity is that *oprC*

might also be repressed by external copper under anaerobic conditions.

**Figure 6. Lack of OprC does not alleviate copper toxicity.** *P. aeruginosa* PA14 growth under anaerobic conditions was monitored during copper and silver stress in minimal media. Growth curves of (**a,c**) WT and (**b,d**) Δ*oprC* in LR media supplemented with 100 mM sodium nitrate at increasing concentrations of added Cu(II) (**a,b**) or Ag (**c,d**).

## Discussion

Our combined data define OprC as a specific importer of Cu(I), which is coordinated by a very unusual CxxxM-HxM binding motif in which the cysteine provides a reducing environment. The presence of a methionine binding track in OprC and the involvement of two methionine residues in the principal binding site are consistent with the preference of Cu(I) for S-containing ligands. Methionines also coordinate Cu(I) in other known copper transporters, such as the bacterial CusABC and CopA exporters and the eukaryotic Ctr1 copper importer[16, 18, 38-40]. These transporters have been shown to also transport silver, but structures with silver have been determined only for the Cus system[16, 38]. Our structural data strongly suggest that OprC can also import silver. With regards to Cu(II), the picture that emerges from our *in vitro* data with detergent-solubilised protein, is that Cu(II) binds non-specifically to OprC and subsequently migrates to the principal binding site via an unknown mechanism, where it is reduced to Cu(I). The Cu(II) affinity of 2.6 μM reported by an early study[26] most likely resulted from non-specific binding, since the rich media used[41] to culture *P. aeruginosa* would have generated OprC with 100% copper occupancy in the principal binding site. Whether Cu(II) is a substrate for OprC *in vivo* remains an open question. A recent study in *P. aeruginosa* proposed a novel copper uptake mechanism in copper-limited conditions which involves secretion of the copper binding protein azurin by a CueR-regulated Type VI secretion system. The secreted azurin would scavenge Cu(II) from the environment and load it onto OprC via a direct interaction, conferring a competitive advantage under copper-limiting conditions[27]. While this model requires further corroboration, our data show that the imported species will be Cu(I).

The copper site most similar to that in OprC occurs in class I Type I copper proteins like cytochrome c oxidase, pseudoazurin and plastocyanin, electron transfer proteins that coordinate Cu(I) or Cu(II) via a 2Cys+1Met+1His site/2His+1Cys+1Met site. Interestingly, substitution of either of the two OprC binding site methionines with histidine generates M147H and M325H transporters that bind copper with similar efficiency as wild type OprC and, at least in the case of M325H, has a pseudoazurin-like active site with tetrahedral geometry (Supplementary Fig. 8). However, both proteins are still colourless and the spectrum of the Cu-loaded M147H variant is EPR-silent (Supplementary Fig. 4), suggesting the principal binding sites in these variants also comprise a reducing environment and bind Cu(I).

Given that copper, and in particular the more toxic Cu(I), is a known antimicrobial, the question
arises why bacteria have proteins dedicated to Cu(I) acquisition from the environment. Indeed,
in contrast to *e.g.* iron that is withheld from a pathogen by the host during infection, host-
derived copper is a "nutritional immunity" antimicrobial response. Thus, bacterial virulence is
generally attenuated by mutations, particularly in transporters, that cause copper sensitivity[42].
No data have yet been reported on the role of OprC in *P. aeruginosa* virulence, but a recent
study reported decreased virulence of an *A. baumannii* OprC knockout[43], which is the opposite
of what one might expect from the copper nutritional immunity model[42]. In this context, our
proteomics data showing that OprC is very abundant in a *P. aeruginosa* mouse infection model
are particularly intriguing. Considering the near-irreversible copper binding, we speculate that
OprC could also play a role as a copper reservoir on the cell surface. As a TBDT, Cu-loaded
OprC depends on TonB for transport, and it is possible that (hypothetical) regulation of the
TonB-OprC interaction could "titrate" copper to the periplasmic space as required, when ex-
ternal copper is low. Conversely, in the presence of high concentrations of external copper,
OprC would be loaded with copper, but this would not lead to import in the absence of an
interaction with TonB. This model could provide an alternative explanation for the lack of a
robust OprC-dependent growth phenotype in *P. aeruginosa* under copper-replete conditions
(Fig. 6). Metallo-proteomes in bacteria are not well characterized, making it challenging to
estimate copper requirements[44]. Total cell quantitation suggests a copper concentration of
~40 µM when cells are grown in metal-restricted medium[45]. Assuming a cellular volume of ~1
$\mu m^3$ ($10^{-15}$ l), this would correspond to ~25,000 molecules per cell. Thus, a pool of 5,000-
10,000 Cu-loaded OprC molecules could provide a substantial part of the total copper content
of the cell, most of which will be present as an essential cofactor in enzymes and bound to
chaperones and storage proteins. Given this need for copper on the one hand and its toxicity
on the other, it is clear that regulation of any copper import protein is crucial, possibly both at
the gene and protein level. Unfortunately, and in contrast to the many copper stress genes
that, as part of the CopR or CueR regulons, are upregulated under aerobic conditions during
copper stress[10, 11, 46-49], nothing is known about how OprC is downregulated during such stress.
Intriguingly, *oprC* (*PA3790*) is in an operon with *PA3789*, which encodes for an uncharacter-
ised inner membrane protein containing PepSY domains, hinting at a peptidase function. An-
other protein strongly downregulated during copper stress is PA5030, which is an MFS trans-
porter with a large number of His+Met residues (26 out of 438 residues), suggesting it could
mediate copper delivery to the cytoplasm, possibly in concert with OprC and an as yet uniden-
tified periplasmic protein[49].
514

Bacterial copper import via metallophores is well established. For example, methanotrophic bacteria produce methanobactins that bind copper with micromolar affinity, and are imported by MbnT[22]. In *E. coli,* yersiniabactin (Ybt) binds Cu(II), which is taken up by the FyuA receptor[50]. Thus, OprC is the first example of a TBDT that mediates copper import without a metallophore. The TBDT with the closest substrate specificity to OprC is the ionic zinc transporter ZnuD from *Neisseria meningitidis*, the structure of which has been solved[51]. Large structural differences between OprC and ZnuD exist for the extracellular loops (overall C$\alpha$ r.m.s.d. ~5.9 Å). ZnuD has several discrete low-affinity binding sites that may guide the metal towards the high-affinity binding site[51]. In OprC, a distinctive "methionine track" provides low-affinity binding sites to guide copper to the high affinity site. Interestingly, while the extracellular loops between OprC and ZnuD are very different and the overall sequence identity is only 28%, the metal binding sites are located at very similar positions and only 2.8 Å apart (Supplementary Fig. 9), suggesting that the transport channel formed via TonB interaction may be similar. Inspection of the ZnuD structure shows that the zinc binding site is excluded from solvent, and we propose that the zinc ion in ZnuD is kinetically trapped, analogous to copper in OprC.

OprC shares ~60 % identity to NosA from *Pseudomonas stutzeri*, for which no structure is available. Like OprC, NosA is expressed under anaerobic conditions and repressed in the presence of $\mu$M concentrations Cu(II)[52-54]. *P. stutzeri* NosA antibodies did not react with *P. aeruginosa*[52], but our structure identifies NosA as an OprC ortholog, since the CxxxM-HxM copper binding motif and some of the methionine track residues are conserved (Supplementary Fig. 10). NosA is important during denitrification in *P. stutzeri* JM300 and was proposed to load copper either directly or indirectly to the periplasmic $N_2O$ reductase NosZ[52-54]. However, a more recent report for a different *P. stutzeri* strain found no difference between NosZ activity and copper content for a *nosA* knockout[55]. In addition, OprC/NosA also occurs in a number of non-denitrifying *Proteobacteria* such as *Salmonella enterica, Klebsiella pneumoniae* and *Acinetobacter baumannii* (Supplementary Fig. 10), showing that NosZ maturation is not a general function of OprC. The occurrence of OprC in some (*e.g. S. enterica*) but not in other (*e.g. E. coli*) Enterobacteria is intriguing, given that the OM of these bacteria is relatively permeable to small molecules due to abundant general porins such as OmpC and OmpF.

The ability of OprC to take up Cu(I) could be important for biofilms, which are anaerobic to various degrees depending on the location inside the biofilm[56, 57]. Oxygen tension also reduces as lung chronic disease mediated by *P. aeruginosa* progresses, turning airway mucus into an anaerobic environment in cystic fibrosis patients that will favour the availability of Cu(I)[58]. *P.*

551 *aeruginosa* is capable of anaerobic respiration by using nitrate, nitrite or nitrous oxide as
552 terminal electron acceptor[59], and OprC has been shown to be induced under anaerobic
553 conditions[36]. Together, our data suggest that OprC is a novel type of copper transporter for the
554 efficient scavenging of copper under a wide range of physiological conditions, including the
555 anaerobiosis that is a hallmark of disease caused by *P. aeruginosa* in humans.
556

**Methods**
**Recombinant production of *Pseudomonas aeruginosa* OprC.** The mature version of the
gene coding for *oprC* of *P. aeruginosa* PAO1 (UniProt ID; PA3790)[60], starting with His56 as
determined by Nakae et al.[26], was synthesized to include a 7 x His tag at the N-terminus
(Eurofins, UK), cloned into the pB22 arabinose-inducible expression vector[61] and transformed
into chemically competent *Escherichia coli* DH5$\alpha$ cells. After expression and processing by
signal peptidase, the N-terminal sequence of this construct is
NVRLQHHHHHHHLEAEEHSQHQ-. A second version of this construct was constructed in a
pB22 version containing a tobacco etch virus (TEV) site after the His7-tag. Correct sequences
were confirmed by DNA sequencing (Eurofins, UK) using both forward and reverse plasmid-
specific primers. The OprC$_{AA}$ mutant was produced by changing the key amino acids Cys143
and Met147 to alanine residues using the KLD Quickchange site-directed mutagenesis kit
(New England Biolabs, UK) and specific primers containing both mutation sites (forward: 5`-
tcgcgcggatgcaccaaccagctatattagc-3`; reverse: 5`-ttcggggcggcgccaagcatcatgc-3`). The single
mutants C143A, M147A, H323A, M325A, M147H and M325H were made in similar ways.
572

OprC recombinant protein production and purification was performed as follows: *E. coli* C43
Δ*cyo* was electroporated with expression vector, recovered for 60 minutes in LB (Sigma, UK)
at 37 °C, and plated on LB agar (Sigma, UK) containing 100 µg mL$^{-1}$ ampicillin (Melford, UK).
Transformants were cultured in LB medium or in LeMasters-Richards (LR) minimal medium
with glycerol (2-3 g/l) as carbon source. All media contained 100 µg mL$^{-1}$ ampicillin. For rich
media, cells were grown (37 °C, 180rpm) until OD600 ~0.6, when protein expression was
induced with 0.1% arabinose for 4-5 h at 30 °C or overnight at 16 °C (150 rpm). For LR media,
a small overnight pre-culture in LB was used at 1/100 v/v to inoculate an LR-medium pre-
culture early in the morning (typically 1 ml preculture for 100 ml of cells was used), which was
grown during the day at 37 °C. After late afternoon inoculation, large-scale cultures (typically
6-8 l) were grown overnight at 30 °C until OD 0.4-0.7, followed by induction with 0.1%
arabinose at 30 °C for 6-8 hours. Cells were harvested by centrifugation (5,000 rpm, 20
minutes, 4 °C), and pellets homogenized in 20 mM Tris (Sigma), 300 mM NaCl (Fisher) pH
8.00 (TBS buffer), in the presence of 10 mM ethylene diamine tetraacetic acid (EDTA, Sigma).

Cells were broken by one pass through a cell disruptor (Constant Systems 0.75 kW operated

at 23 kpsi), centrifuged at 42,000 rpm for 45 minutes at 4 °C (45Ti rotor; Beckman), and the

resulting total membrane fraction was homogenized in TBS buffer containing 1.5% Lauryl-

dimethylamine oxide (LDAO) (Sigma, UK). Membrane proteins were extracted by stirring (60

minutes, 4 °C), centrifuged (42,000 rpm in 45Ti rotor, 30 minutes, 4°C), and the membrane

extract was loaded on a Chelating Sepharose Fast Flow bed resin (~10 ml; GE Healthcare,

UK) previously activated with 200 mM $NiCl_2$ (Sigma) and equilibrated in TBS containing 0.15 %

n-dodecyl-beta-D-maltopyranoside (DDM). After washing with 15 column volumes buffer with

30 mM imidazole, protein was eluted with 0.25 M imidazole buffer (Fisher), incubated with 20

mM EDTA (30 minutes, 4 °C), and loaded on a Superdex 200 16/600 size exclusion column

equilibrated with 10 mM Hepes, 100 mM NaCl, 0.05% DDM, 10 mM EDTA, pH 7.5. Peak

fractions were pooled and concentrated using a 50 MWCO Amicon filter (Millipore, UK),

analyzed on SDS-PAGE, flash-frozen in liquid nitrogen and stored at -80C. Typical yields of

purified wild type and most mutant OprC proteins ranged between 2-5 mg per l media grown

at 16 °C. All media and buffer components were made in fresh milli-Q water.

Protein preparations intended for crystal trials were pooled and buffer-exchanged to 10 mM

Hepes 100 mM NaCl, 0.4% tetraethylene glycol mono-octyl ether ($C_8E_4$) (Anatrace, US), pH

7.5. $NaNO_3$ was substituted for NaCl for protein preparations intended for crystal trials with

silver. Protein preparations to be used for metal analysis after removal of the His-tag

underwent a slightly different protocol. The elution fraction from immobilized metal affinity

chromatography (IMAC) was buffer-exchanged to 50 mM Tris, 0.5 mM EDTA, 0.2 mM TCEP,

100 mM NaCl, 0.05% DDM (Anatrace, US), pH 7.50, and submitted to TEV protease digestion

(ratio 1 mg TEV: 10 mg protein, 4 °C, overnight). Samples were submitted to a second IMAC

column, where flow-through and wash fractions were combined for the subsequent SEC step

in 10 mM HEPES 100 mM NaCl 0.05 % DDM 0.5 mM EDTA pH 7.5. Protein concentration

was determined by BCA assay (Thermo Scientific, UK) and by UV/Vis absorbance at 280nm

(considering OprC $E_{0.1\%}$ = 1.6 as determined by ProtParam).

**In vitro metal binding assays and Inductively Coupled Plasma Mass Spectrometry (ICP-**

**MS).** OprC samples intended for metal binding assays were exchanged into respective chelex-

treated buffers without EDTA and were equilibrated with different equivalents of Cu(II) or Zn,

for 30 min at room temperature (n=3). Protein concentrations used were in the range of 10 –

20 μM. Samples were loaded on an analytical Superdex 200 Increase 10/300GL (GE

Healthcare) column, equilibrated in 10 mM Hepes, 100 mM NaCl, 0.05% DDM, 0.5 mM EDTA

pH 7.5.  Size exclusion peaks were pooled, concentrated and quantified for protein by UV

absorbance at 280 nm. Protein samples were diluted 10-fold in 2.5% $HNO_3$. Analytical metal

standards of 0- 500 ppb were prepared by serial dilution from individual metal stocks (VWR, UK) and were matrix-matched to protein samples. Samples and standard curves were analysed by ICP-MS using a Thermo X-series instrument (Thermo Fisher Scientific), and samples were typically screened for the presence of $^{63}$Cu and $^{65}$Zn.

**Copper extraction (demetallation) experiments.** OprC$_{WT}$, OprC$_{AA}$ and M147A samples were incubated with 3 equivalents copper for 30 min and then loaded onto a Superdex S-200 Increase 10/300GL column equilibrated with 10 mM HEPES 100 mM NaCl 0.05 % DDM 0.5 mM EDTA pH 7.5. Peak fractions were pooled, concentrated and quantified by UV absorbance at 280 nm. Samples were exchanged into respective chelex-treated buffer without EDTA. For demetallation experiments, 20 µM of copper-bound proteins were taken in duplicates and incubated with 100-fold excess of the copper chelator bathocuproine disulfonate (BCS) (Sigma) and 100-fold excess of the reducing agent hydroxyl amine (NH$_2$OH) (Sigma) at 60 °C and room temperature. The amount of extracted Cu(I) was quantified by measuring the formation of coloured copper chelate complex, using the complex molar extinction coefficient $\varepsilon_{483nm}$ = 13,300 M$^{-1}$ cm$^{-1}$ [62].

**Fluorescence spectroscopy**

OprC samples in chelex-treated buffers without EDTA were diluted in 10 mM HEPES, 100 mM NaCl, 0.05% DDM (pH 7.5) to a volume of 250 µL and a concentration of 20 µM, and samples were titrated with consecutive additions of Cu(II). Fluorescence emission spectra were recorded on a Varian Cary Eclipse spectrophotometer (Agilent). The excitation wavelength was 300 nm and emission spectra were recorded between 300 – 500 nm, with excitation slit of 5 nm, at a scan rate of 600 nm/min. Data was plotted and analysed for peak changes in the region of 300-400 nm due to Cu(II) binding[63]. Experiments were performed in triplicates under aerobic conditions, and reactions were equilibrated for 5 minutes at room temperature prior to recording each fluorescence spectrum.

**Protein crystallization, data collection and structure determination.** Sitting-drop crystallization trials were set up using a Mosquito crystallization robot (TTP Labtech) with commercial screens (MemGold1 and MemGold2, Molecular Dimensions) at 20 °C. To obtain the initial structure of Cu-bound OprC, the protein (~12 mg/ml) was incubated with 3 mM CuCl$_2$ for 1 hr at room temperature, followed by setting up crystallisation trials. A number of initial hits were obtained and were subsequently optimised by manual hanging drop vapour diffusion using larger drops (typically 1-1.5 ul protein + 1 ul reservoir). Well-diffracting crystals (~3 Å resolution at a home source) were obtained in 0.1 M NaCl/0.15 M NH$_4$SO$_4$/0.1 M MES pH 6.5/18-22% PEG1000. Crystals were cryoprotected with mother liquor lacking CuCl$_2$

containing 10% PEG400 for ~5-10 s and flash-frozen in liquid nitrogen. Diffraction data were collected at Diamond Light Source (Didcot, UK) at beamline i02. For the best crystal, belonging to space group $C222_1$, 720 degrees of data were collected at an energy of 8994 eV, corresponding to the K-edge of copper (Supplementary Table 1). Data were auto processed by xia2. The structure was solved via single anomalous dispersion (SAD) via AUTOSOL in Phenix[64]. Two copper sites were found, one for each OprC molecule in the asymmetric unit (Supplementary Fig. 1). The phases were of sufficient quality to allow automated model building via Phenix AUTOBUILD, generating ~60% of the structure and using data to 2.0 Å. The remainder of the structure was built manually, via iterative cycles of refinement in Phenix and model building in COOT[65]. Metal coordination was analysed by the Check-my-metal server[66]. The final refinement statistics are listed in Supplementary Table 1. Subsequently, crystals were also obtained without any copper supplementation of the protein. These were isomorphous to those described above and obtained under identical conditions. Molecular replacement indicated the presence of copper and an identical structure to that obtained above (data not shown). $OprC_{AA}$ crystals (~10 mg/ml protein) were obtained and optimized by hanging drop vapor diffusion as described above, and diffraction-quality crystals were obtained in the same conditions as for Cu-OprC, *i.e.* 0.1 M sodium chloride/0.15 M ammonium sulfate/0.01 M MES sodium pH 6.5/19% (w/v) PEG1000. Interestingly however, the $OprC_{AA}$ crystals belong to a different space group ($P22_12_1$), most likely as a result of structural differences between both OprC variants. Diffraction data were collected at Diamond Light Source (Didcot, UK) at beamline I24. Diffraction data were processed in XDS[67]. The structure was solved by molecular replacement (MR) using Phaser, with wildtype OprC as the search model. Model building was done in COOT and refinement in Phenix. As for Cu-OprC, the data collection and refinement statistics are shown in Supplementary Table 1. C143A and H323A proteins (~10 – 12 mg/ml protein) were incubated with 2 mM $CuSO_4$ at room temperature for 1 hr, followed by co-crystallisation. Diffracting crystals for both C143A and H323A in the presence of copper were obtained in 0.34 M Ammonium sulfate/0.1 M Sodium citrate pH 5.5/12 -16 % w/v PEG 4000 and were cryo-protected using mother liquor lacking $CuSO_4$ and with 25% ethylene glycol for ~10 s and flash-frozen in liquid nitrogen. M147H and M325H crystals were obtained in the same condition as those for wild type OprC. For co-crystallisation with silver, OprC proteins were incubated with 2 mM AgNO3 for 1 hour at room temperature, followed by co-crystallisation. Well-diffracting $OprC_{WT}$ crystals with silver were obtained under the same conditions as in the presence of copper. For the best $OprC_{WT}$ crystal, belonging to space group $C222_1$, 999 degrees of data were collected at an energy of 8000 eV to obtain anomalous signals for Ag. C143A and H323A crystals (~10-12 mg/ml protein) with Ag were obtained from 0.2 M Choline chloride/0.1 M Tris pH 7.5/12-16 % w/v PEG 2000 MME and 0.5 M Potassium chloride/0.05 M HEPES pH 6.5/12-16 % v/v PEG 400, respectively. Crystals

were cryoprotected for 5-20 s with mother liquor lacking $AgNO_3$ but containing 25% ethylene glycol for C143A and 20 % PEG 400 for H323A. For C143A and M325A crystallised in the presence of copper or silver, datasets of 360 degrees each were collected at energies of 8800 and 9175 eV, using different parts of the same crystal (Supplementary Tables 2 and 3).

**Electron Paramagnetic Resonance Spectroscopy.**

Electron Paramagnetic Resonance (EPR) measurements were carried out using a Bruker ELEXSYS-E500 X-band EPR spectrometer operating in continuous wave mode, equipped with an Oxford variable-temperature unit and ESR900 cryostat with Super High-Q resonator. All EPR samples were prepared in quartz capillary tubes (outer diameter; 4.0 mm, inner diameter 3.0 mm) and frozen immediately in liquid $N_2$ until further analysis. The experimental setup and conditions were similar to reported earlier[68]. The low temperature EPR spectra were acquired using the following conditions: sweep time of 84 s, microwave power of 0.2 mW, time constant of 81 ms, average microwave frequency of 9.44 GHz and modulation amplitude of 5 G, T = 20 K. The concentration of $OprC_{WT}$ and mutants varied from 210-260 $\mu$M in 10 mM HEPES, 100 mM NaCl, 0.03 % DDM (n-dodecyl-D-maltoside), pH 7.5 and 10 mM HEPES, 100 mM $NaNO_3$, 0.4 % C8E4 (tetraethylene glycol monooctyl ether) buffer pH 8, respectively. Analysis and simulations of the cw-EPR spectra were performed using EasySpin toolbox (5.2.28) for the Matlab (R2017a) program package[69, 70].

**Quantum chemistry calculations.** A truncated version of the crystal structure of OprC (PDB ID 6FOK) consisting of only the copper ion, Cys143, Met147, His323 and Met2325 was used as input for single point energy calculations. The link atoms between these four amino acids and the rest of the protein were capped with hydrogen atoms, giving a total of 70 atoms. The calculations were performed on the system (i) without copper, (ii) with Cu(I) or (iii) with Cu(II). In addition, two possibilities were considered for the state of the cysteine: hydrogenated (reduced) or dehydrogenated (oxidised). The calculations were performed in linear-scaling Density Functional Theory (DFT) with the PBE exchange-correlation functional in ONETEP[71]. Norm-conserving pseudopotentials, an energy cutoff of 800 eV and a kernel cutoff of 1000 Bohr in open-boundary conditions were used in a cubic box with sides of 30 Å in length. Quantum chemistry calculations produce huge energies because they are the energies of all the electrons (including core electrons) interacting with the nuclei and themselves. However, the total energies are meaningless and what is chemically important are differences between different conformations and different bonding situations, which are typically much smaller as they concern re-arrangements of valence electrons. For this reason, it is crucial to converge the huge total energies very tightly in order to capture the chemically interesting energy differences.

### *In vivo* metal toxicity assays

For metal toxicity assays in *E. coli,* JW0473-3 (*ΔcopA*)[72] were transformed with plasmids containing OprC$_{WT}$, OprC$_{AA}$ or empty cloning vector (pB22). The copper exporter *copA* mutant was obtained from the Keio collection; Growth assays were set up in flat-bottomed 96-well plates (Thermo Scientific) containing LB-ampicillin supplemented with 0.01% arabinose. Different concentrations (0-6mM) of Cu(II) (CuSO$_4$) were tested in order to assess growth phenotypes under metal excess conditions. Cultures in triplicates were inoculated with 1:100 dilution of overnight pre-inoculum from isolated colonies. Assays were performed in a plate reader (Tecan Infinite M200), with 37C incubation and regular orbital shaking of 30 seconds at every 5 minutes, for a total of 25 hours. Time points were collected with 30 mins intervals. For metal toxicity assays in *P. aeruginosa*, PA14 WT and PA14 *Δoprc* strains were used and assays were performed in anaerobic conditions. Pre- cultures were grown under anaerobiosis in rich LB or LR minimal media supplemented with 100 mM sodium nitrate. The Cu(II) (CuSO$_4$) range tested varied from 0 – 7 mM in rich media to 0 – 8 µM in LR media.  Cultures in triplicates were inoculated with precultures grown in respective media under anaerobic conditions without any excess metal added. Plates were set up and sealed inside an anaerobic chamber (Don Whitley Scientific). For silver toxicity, 0 – 1 µM Ag (AgNO$_3$) was used and assays were performed as described for Cu (II) in LR minimal media supplemented with 100 mM NaNO$_3$. Experiments were performed in triplicates for all tested conditions.

### Animal infection models.

*Intra-tracheal instillation model*: specific pathogen free (SPF) immunocompetent male Sprague-Dawley rats weighing 100 - 120 g or male CD-1 mice weighing 20 - 25 g were infected by depositing an agar bead containing around $10^7$ colony-forming units *Acinetobacter baumannii* ATCC 19606 and *Pseudomonas aeruginosa* UCBPP-PA14, deep into the lung via nonsurgical intra-tracheal intubation[73]. In brief, animals were anesthetized with isoflurane (5%) and oxygen (1.5 L/min) utilizing an anesthesia machine. Depth of anesthesia was evaluated by absence of gag reflex; if the reflex was present, the animal was placed back under anesthesia until the reflex disappeared. No animals were utilized until they were fully anesthetized.  Animals were infected via intra-bronchial instillation of molten agar suspension (rats- 100 µl) (mice- 20 µl) via intra-tracheal intubation, and then allowed to recover. Animals were returned in their home cages and observed until recovered from anesthesia. At 24 h post infection, animals were sacrificed and lung was homogenized in sterile saline using a lab blender. All procedures are in accordance with protocols approved by the GSK Institutional Animal Care and Use Committee (IACUC), and meet or exceed the standards of the American

771 Association for the Accreditation of Laboratory Animal Care (AAALAC), the United States
772 Department of Health and Human Services and all local and federal animal welfare laws.
773

**Sample workup for proteomics**
The sample workup protocol was optimized to deplete host material while maintaining *A.*
*baumannii* and *P. aeruginosa* viability until lysis. All buffers and equipment were used at 0 to
4 °C to minimize proteome changes during sample workup. The sample volume (maximum of
1 ml) was estimated and an equal volume of 1% Tergitol in PBS was added followed by
vigorous vortexing for 30 s. After centrifugation at 500 x g for 5 min, the supernatant was
transferred to a fresh tube, and the pellet was extracted again with 2 ml 0.5% Tergitol in PBS.
The supernatant was combined with the first supernatant and centrifuged at 18'000 x g for 5
min. The pellet was washed with 2 ml and again centrifuged at 18'000 x g for 5 min. The
supernatant was removed, and the pellet was resuspended in 100 µL 5% sodium
deoxycholate, 5 mM Tris (2-carboxyethyl) phosphine hydrochloride, 100 mM $NH_4HCO_3$. The
sample was incubated at 90°C for 1 min. and then stored at -80 °C. Samples were thawed
and sonicated for 2 x 20 s (1 s interval, 100% power). Proteins were alkylated with 10 mM
iodoacetamide for 30 min in the dark at room temperature. Samples were diluted with 0.1M
ammonium bicarbonate solution to a final concentration of 1% sodium deoxycholate before
digestion with trypsin (Promega) at 37°C overnight (protein to trypsin ratio: 50:1). After
digestion, the samples were supplemented with TFA to a final concentration of 0.5% and HCl
to a final concentration of 50 mM. Precipitated sodium deoxycholate was removed by
centrifugation at 4°C and 14'000 rpm for 15 min. Peptides in the supernatant were desalted
on C18 reversed phase spin columns according to the manufacturer's instructions (Macrospin,
Harvard Apparatus), dried under vacuum, and stored at −80°C until further processing.
795

**Parallel reaction monitoring**
Heavy proteotypic peptides (JPT Peptide Technologies GmbH) were chemically synthesized
for *A. baumannii* and *P. aeruginosa* outer membrane proteins. Peptides were chosen
dependent on their highest detection probability and their length ranged between 7 and 20
amino acids. Heavy proteotypic peptides were spiked into each sample as reference peptides
at a concentration of 20 fmol of heavy reference peptides per 1 µg of total endogenous protein
mass. For spectrum library generation, we generated parallel reaction-monitoring (PRM)[74]
assays from a mixture containing 500 fmol of each reference peptide. The setup of the µRPLC-
MS system was as described previously[75]. Chromatographic separation of peptides was
carried out using an EASY nano-LC 1000 system (Thermo Fisher Scientific) equipped with a
heated RP-HPLC column (75 µm x 37 cm) packed in-house with 1.9 µm C18 resin (Reprosil-
AQ Pur, Dr. Maisch). Peptides were separated using a linear gradient ranging from 97%

solvent A (0.15% formic acid, 2% acetonitrile) and 3% solvent B (98% acetonitrile, 2% water,

0.15% formic acid) to 30% solvent B over 60 minutes at a flow rate of 200 nl/min. Mass

spectrometry analysis was performed on Q-Exactive HF mass spectrometer equipped with a

nanoelectrospray ion source (both Thermo Fisher Scientific). Each MS1 scan was followed by

high-collision-dissociation (HCD) of the 10 most abundant precursor ions with dynamic

exclusion for 20 seconds. Total cycle time was approximately 1 s. For MS1, 3e6 ions were

accumulated in the Orbitrap cell over a maximum time of 100 ms and scanned at a resolution

of 120,000 FWHM (at 200 m/z). MS2 scans were acquired at a target setting of 1e5 ions,

accumulation time of 50 ms and a resolution of 30,000 FWHM (at 200 m/z). Singly charged

ions and ions with unassigned charge state were excluded from triggering MS2 events. The

normalized collision energy was set to 35%, the mass isolation window was set to 1.1 m/z and

one microscan was acquired for each spectrum.

The acquired raw-files were converted to the mascot generic file (mgf) format using the

msconvert tool (part of ProteoWizard, version 3.0.4624 (2013-6-3)). Converted files (mgf

format) were searched by MASCOT (Matrix Sciences) against normal and reverse sequences

(target decoy strategy) of the UniProt database of *Acinetobacter baumannii* strains ATCC

19606 and ATCC 17978, and Pseudomonas aeruginosa UCBPP-PA14, as well as commonly

observed contaminants. The precursor ion tolerance was set to 20 ppm and fragment ion

tolerance was set to 0.02 Da. Full tryptic specificity was required (cleavage after lysine or

arginine residues unless followed by proline), three missed cleavages were allowed,

carbamidomethylation of cysteins (+57 Da) was set as fixed modification and arginine (+10

Da), lysine (+8 Da) and oxidation of methionine (+16 Da) were set as variable modifications.

For quantitative PRM experiments the resolution of the orbitrap was set to 30,000 FWHM (at

200 m/z) and the fill time was set to 50 ms to reach a target value of 1e6 ions. Ion isolation

window was set to 0.7 Th (isolation width) and the first mass was fixed to 100 Th. Each

condition was analyzed in biological triplicates. All raw-files were imported into Spectrodive

(Biognosys AG) for protein and peptide quantification.

**Author contributions**

BvdB designed the study. SPB and BvdB expressed, purified, and crystallized proteins. AB

collected the diffraction data. SPB and BvdB analysed the diffraction data and refined the

structures. SPB performed metal binding and *in vivo* growth experiments. TRY performed ICP-

MS analyses. MS performed cw-EPR measurements, simulations and interpreted the data.

BB performed quantum energy calculations, supervised by SK. SH performed proteomics ex-

periments, supervised by DB. BvdB and SPB wrote the paper.

**Acknowledgements**

We would like to thank Scott Sucoloski, Jennifer Hoover, Josh West (Glaxo Smith Kline) for providing proteomics samples. We would also like to thank Kevin Waldron (Newcastle University) for useful discussions and for carrying out initial ICP-MS analyses. We also thank Deenah Osman and Nigel Robinson (Durham University) for elemental analyses by ICP-MS and helpful discussions, supported by awards BB/L009226/1 and BB/R002118/1 from the Biotechnology and Biological Sciences Research Council, BBSRC, UK. We are indebted to the Diamond Light Source for beam time (proposals mx9948, mx13587 and mx18598) and beamline assistance. MS acknowledges the EPSRC National (UK) EPR Research Facility and Service for use of the EPR spectrometers. MS and BvdB thank Luisa Ciano for the useful discussions at an early stage. The research leading to these results was in part conducted as part of the Translocation consortium ([www.translocation.eu](www.translocation.eu)) and has received support from the Innovative Medicines Initiatives Joint Undertaking under Grant Agreement No. 115525, resources that are composed of financial contributions from the European Union's seventh framework programme (FP7/2007–2013) and European Federation of Pharmaceutical Industries and Associations companies in-kind contribution. BvdB would also like to acknowledge the Royal Society for salary support.

**Accession codes**

Coordinates and structure factors have been deposited in the Protein Data Bank (http://www.ebi.ac.uk/pdbe/) with accession codes 6FOK (OprC$_{WT}$), 6FOM (OprC$_{AA}$), 6Z8Q (OprC$_{WT}$ Ag 8000 ev), 6Z9I (OprC$_{C143A}$ Ag 8800 eV), 6Z99 (OprC$_{C143A}$ Ag 9175 eV), 6Z8Y (OprC$_{C143A}$ Cu 8800 eV), 6Z8Z (OprC$_{C143A}$ Cu 9175 eV), 6Z8T (OprC$_{H323A}$ Ag 8800 eV), 6Z8U (OprC$_{H323A}$ Ag 9175 eV), 6Z8R (OprC$_{M147H}$), 6Z8S (OprC$_{M325H}$), 6Z9N (OprC$_{H323A}$ Cu 9175 eV), 6Z9Y (OprC$_{H323A}$ Cu 8800 eV).

**References**

1.    Hodgkinson, V. & Petris, M.J. Copper homeostasis at the host-pathogen interface. *The Journal of biological chemistry* **287**, 13549-13555 (2012).

2.    Harrison, J.J., Ceri, H. & Turner, R.J. Multimetal resistance and tolerance in microbial biofilms. *Nature reviews. Microbiology* **5**, 928-938 (2007).

3.    Lemire, J.A., Harrison, J.J. & Turner, R.J. Antimicrobial activity of metals: mechanisms, molecular targets and applications. *Nature reviews. Microbiology* **11**, 371-384 (2013).

4.    Braymer, J.J. & Giedroc, D.P. Recent developments in copper and zinc homeostasis in bacterial pathogens. *Current opinion in chemical biology* **19**, 59-66 (2014).

5.    Robinson, N.J. & Winge, D.R. Copper metallochaperones. *Annual review of*
*biochemistry* **79**, 537-562 (2010).

6.    Hernandez-Montes, G., Arguello, J.M. & Valderrama, B. Evolution and diversity of
periplasmic proteins involved in copper homeostasis in gamma proteobacteria. *BMC*
*microbiology* **12**, 249 (2012).

7.    Zimmermann, M. et al. PcoE--a metal sponge expressed to the periplasm of copper
resistance *Escherichia coli*. Implication of its function role in copper resistance. *Journal*
*of inorganic biochemistry* **115**, 186-197 (2012).

8.    Vita, N. et al. A four-helix bundle stores copper for methane oxidation. *Nature* **525**,
140-143 (2015).

9.    Arguello, J.M., Raimunda, D. & Padilla-Benavides, T. Mechanisms of copper
homeostasis in bacteria. *Frontiers in cellular and infection microbiology* **3**, 73 (2013).

10.    Zhang, X.X. & Rainey, P.B. Regulation of copper homeostasis in *Pseudomonas*
*fluorescens SBW25*. *Environmental microbiology* **10**, 3284-3294 (2008).

11.    Hu, Y.H., Wang, H.L., Zhang, M. & Sun, L. Molecular analysis of the copper-responsive
CopRSCD of a pathogenic *Pseudomonas fluorescens* strain. *Journal of microbiology*
**47**, 277-286 (2009).

12.    Rubino, J.T. & Franz, K.J. Coordination chemistry of copper proteins: how nature
handles a toxic cargo for essential function. *Journal of inorganic biochemistry* **107**,
129-143 (2012).

13.    Boal, A.K. & Rosenzweig, A.C. Structural biology of copper trafficking. *Chemical*
*reviews* **109**, 4760-4779 (2009).

14.    Ma, Z., Jacobsen, F.E. & Giedroc, D.P. Coordination chemistry of bacterial metal
transport and sensing. *Chemical reviews* **109**, 4644-4681 (2009).

15.    Gonzalez-Guerrero, M. & Arguello, J.M. Mechanism of Cu+-transporting ATPases:
soluble Cu+ chaperones directly transfer Cu+ to transmembrane transport sites.
*Proceedings of the National Academy of Sciences of the United States of America* **105**,
5992-5997 (2008).

16.    Su, C.C. et al. Crystal structure of the CusBA heavy-metal efflux complex of
*Escherichia coli*. *Nature* **470**, 558-562 (2011).

17.    Outten, F.W., Huffman, D.L., Hale, J.A. & O'Halloran, T.V. The independent cue and
cus systems confer copper tolerance during aerobic and anaerobic growth in
*Escherichia coli*. *The Journal of biological chemistry* **276**, 30670-30677 (2001).

18.    Kulathila, R., Kulathila, R., Indic, M. & van den Berg, B. Crystal structure of *Escherichia*
*coli* CusC, the outer membrane component of a heavy metal efflux pump. *PloS one* **6**,
e15610 (2011).

19.    Khalfaoui-Hassani, B. et al. Widespread Distribution and Functional Specificity of the
918           Copper Importer CcoA: Distinct Cu Uptake Routes for Bacterial Cytochrome c
919           Oxidases. *mBio* **9** (2018).

920    20.    Ekici, S., Yang, H., Koch, H.G. & Daldal, F. Novel transporter required for biogenesis
921           of cbb3-type cytochrome c oxidase in *Rhodobacter capsulatus. mBio* **3** (2012).

922    21.    De Feo, C.J., Aller, S.G., Siluvai, G.S., Blackburn, N.J. & Unger, V.M. Three-
923           dimensional structure of the human copper transporter hCTR1. *Proceedings of the*
924           *National Academy of Sciences of the United States of America* **106**, 4237-4242 (2009).

925    22.    Dassama, L.M., Kenney, G.E., Ro, S.Y., Zielazinski, E.L. & Rosenzweig, A.C.
926           Methanobactin transport machinery. *Proceedings of the National Academy of*
927           *Sciences of the United States of America* **113**, 13027-13032 (2016).

928    23.    Govan, J.R. & Deretic, V. Microbial pathogenesis in cystic fibrosis: mucoid
929           *Pseudomonas aeruginosa* and *Burkholderia cepacia. Microbiological reviews* **60**, 539-
930           574 (1996).

931    24.    Lyczak, J.B., Cannon, C.L. & Pier, G.B. Establishment of *Pseudomonas aeruginosa*
932           infection: lessons from a versatile opportunist. *Microbes and infection* **2**, 1051-1060
933           (2000).

934    25.    Chevalier, S. et al. Structure, function and regulation of *Pseudomonas aeruginosa*
935           porins. *FEMS microbiology reviews* **41**, 698-722 (2017).

936    26.    Yoneyama, H. & Nakae, T. Protein C (OprC) of the outer membrane of *Pseudomonas*
937           *aeruginosa* is a copper-regulated channel protein. *Microbiology* **142 ( Pt 8)**, 2137-2144
938           (1996).

939    27.    Han, Y. et al. A *Pseudomonas aeruginosa* type VI secretion system regulated by CueR
940           facilitates copper acquisition. *PLoS pathogens* **15**, e1008198 (2019).

941    28.    Tsukihara, T. et al. The whole structure of the 13-subunit oxidized cytochrome c
942           oxidase at 2.8 A. *Science* **272**, 1136-1144 (1996).

943    29.    Robinson, H. et al. Structural basis of electron transfer modulation in the purple CuA
944           center. *Biochemistry* **38**, 5677-5683 (1999).

945    30.    Shibata, N. et al. Novel insight into the copper-ligand geometry in the crystal structure
946           of *Ulva pertusa* plastocyanin at 1.6-A resolution. Structural basis for regulation of the
947           copper site by residue 88. *The Journal of biological chemistry* **274**, 4225-4230 (1999).

948    31.    Celia, H. et al. Structural insight into the role of the Ton complex in energy transduction.
949           *Nature* **538**, 60-65 (2016).

950    32.    Noinaj, N., Guillier, M., Barnard, T.J. & Buchanan, S.K. TonB-dependent transporters:
951           regulation, structure, and function. *Annual review of microbiology* **64**, 43-60 (2010).

952    33.    Frandsen, K.E. et al. The molecular basis of polysaccharide cleavage by lytic
953           polysaccharide monooxygenases. *Nature chemical biology* **12**, 298-303 (2016).

34. Hedison, T.M., Heyes, D.J., Shanmugam, M., Iorgu, A.I. & Scrutton, N.S. Solvent-
slaved protein motions accompany proton coupled electron transfer reactions
catalysed by copper nitrite reductase. *Chemical communications* **55**, 5863-5866
(2019).

35. Simmons, T.J. et al. Structural and electronic determinants of lytic polysaccharide
monooxygenase reactivity on polysaccharide substrates. *Nature communications* **8**,
1064 (2017).

36. Filiatrault, M.J. et al. Effect of anaerobiosis and nitrate on gene expression in
*Pseudomonas aeruginosa*. *Infection and immunity* **73**, 3764-3772 (2005).

37. Yamano, Y., Nishikawa, T. & Komatsu, Y. Cloning and nucleotide sequence of
anaerobically induced porin protein E1 (OprE) of *Pseudomonas aeruginosa PAO1*.
*Molecular microbiology* **8**, 993-1004 (1993).

38. Xue, Y. et al. Cu(I) recognition via cation-pi and methionine interactions in CusF.
*Nature chemical biology* **4**, 107-109 (2008).

39. Ren, F. et al. X-ray structures of the high-affinity copper transporter Ctr1. *Nature*
*communications* **10**, 1386 (2019).

40. Stoyanov, J.V., Magnani, D. & Solioz, M. Measurement of cytoplasmic copper, silver,
and gold with a lux biosensor shows copper and silver, but not gold, efflux by the CopA
ATPase of *Escherichia coli*. *FEBS letters* **546**, 391-394 (2003).

41. Yoshihara, E. & Nakae, T. Identification of porins in the outer membrane of
*Pseudomonas aeruginosa* that form small diffusion pores. *The Journal of biological*
*chemistry* **264**, 6297-6301 (1989).

42. Ladomersky, E. & Petris, M.J. Copper tolerance and virulence in bacteria. *Metallomics*
*: integrated biometal science* **7**, 957-964 (2015).

43. Abdollahi, S., Rasooli, I. & Mousavi Gargari, S.L. The role of TonB-dependent copper
receptor in virulence of *Acinetobacter baumannii*. *Infection, genetics and evolution :*
*journal of molecular epidemiology and evolutionary genetics in infectious diseases* **60**,
181-190 (2018).

44. Cvetkovic, A. et al. Microbial metalloproteomes are largely uncharacterized. *Nature*
**466**, 779-782 (2010).

45. Schalk, I.J. & Cunrath, O. An overview of the biological metal uptake pathways in
*Pseudomonas aeruginosa*. *Environmental microbiology* **18**, 3227-3246 (2016).

46. Thaden, J.T., Lory, S. & Gardner, T.S. Quorum-sensing regulation of a copper toxicity
system in *Pseudomonas aeruginosa*. *Journal of bacteriology* **192**, 2557-2568 (2010).

47. Teitzel, G.M. et al. Survival and growth in the presence of elevated copper:
transcriptional profiling of copper-stressed *Pseudomonas aeruginosa*. *Journal of*
*bacteriology* **188**, 7242-7256 (2006).

48.  Frangipani, E., Slaveykova, V.I., Reimmann, C. & Haas, D. Adaptation of aerobically
growing *Pseudomonas aeruginosa* to copper starvation. *Journal of bacteriology* **190**,
6706-6717 (2008).

49.  Quintana, J., Novoa-Aponte, L. & Arguello, J.M. Copper homeostasis networks in the
bacterium *Pseudomonas aeruginosa*. *The Journal of biological chemistry* **292**, 15691-
15704 (2017).

50.  Koh, E.I., Robinson, A.E., Bandara, N., Rogers, B.E. & Henderson, J.P. Copper import
in *Escherichia coli* by the yersiniabactin metallophore system. *Nature chemical biology*
**13**, 1016-1021 (2017).

51.  Calmettes, C. et al. The molecular mechanism of Zinc acquisition by the neisserial
outer-membrane transporter ZnuD. *Nature communications* **6**, 7996 (2015).

52.  Lee, H.S., Hancock, R.E. & Ingraham, J.L. Properties of a *Pseudomonas stutzeri* outer
membrane channel-forming protein (NosA) required for production of copper-
containing N2O reductase. *Journal of bacteriology* **171**, 2096-2100 (1989).

53.  Mokhele, K., Tang, Y.J., Clark, M.A. & Ingraham, J.L. A *Pseudomonas stutzeri* outer
membrane protein inserts copper into N2O reductase. *Journal of bacteriology* **169**,
5721-5726 (1987).

54.  Lee, H.S., Abdelal, A.H., Clark, M.A. & Ingraham, J.L. Molecular characterization of
nosA, a *Pseudomonas stutzeri* gene encoding an outer membrane protein required to
make copper-containing N2O reductase. *Journal of bacteriology* **173**, 5406-5413
(1991).

55.  Wunsch, P., Herb, M., Wieland, H., Schiek, U.M. & Zumft, W.G. Requirements for
Cu(A) and Cu-S center assembly of nitrous oxide reductase deduced from complete
periplasmic enzyme maturation in the nondenitrifier *Pseudomonas putida*. *Journal of*
*bacteriology* **185**, 887-896 (2003).

56.  Flemming, H.C. & Wingender, J. The biofilm matrix. *Nature reviews. Microbiology* **8**,
623-633 (2010).

57.  Kragh, K.N. et al. Role of Multicellular Aggregates in Biofilm Formation. *mBio* **7**,
e00237 (2016).

58.  Alvarez-Ortega, C. & Harwood, C.S. Responses of *Pseudomonas aeruginosa* to low
oxygen indicate that growth in the cystic fibrosis lung is by aerobic respiration.
*Molecular microbiology* **65**, 153-165 (2007).

59.  Hassett, D.J. et al. *Pseudomonas aeruginosa* hypoxic or anaerobic biofilm infections
within cystic fibrosis airways. *Trends in microbiology* **17**, 130-138 (2009).

60.  Stover, C.K. et al. Complete genome sequence of *Pseudomonas aeruginosa PAO1*,
an opportunistic pathogen. *Nature* **406**, 959-964 (2000).

61.    Guzman, L.M., Belin, D., Carson, M.J. & Beckwith, J. Tight regulation, modulation, and
1028           high-level expression by vectors containing the arabinose PBAD promoter. *Journal of*
1029           *bacteriology* **177**, 4121-4130 (1995).

1030    62.    Bagchi, P., Morgan, M.T., Bacsa, J. & Fahrni, C.J. Robust affinity standards for Cu(I)
1031           biochemistry. *Journal of the American Chemical Society* **135**, 18549-18559 (2013).

1032    63.    Wijekoon, C.J., Young, T.R., Wedd, A.G. & Xiao, Z. CopC protein from *Pseudomonas*
1033           *fluorescens SBW25* features a conserved novel high-affinity Cu(II) binding site.
1034           *Inorganic chemistry* **54**, 2950-2959 (2015).

1035    64.    Adams, P.D. et al. PHENIX: a comprehensive Python-based system for
1036           macromolecular structure solution. *Acta crystallographica. Section D, Biological*
1037           *crystallography* **66**, 213-221 (2010).

1038    65.    Emsley, P., Lohkamp, B., Scott, W.G. & Cowtan, K. Features and development of
1039           Coot. *Acta crystallographica. Section D, Biological crystallography* **66**, 486-501 (2010).

1040    66.    Zheng, H. et al. Validation of metal-binding sites in macromolecular structures with the
1041           CheckMyMetal web server. *Nature protocols* **9**, 156-170 (2014).

1042    67.    Kabsch, W. Xds. *Acta crystallographica. Section D, Biological crystallography* **66**, 125-
1043           132 (2010).

1044    68.    Karimov, R.R. & Hartwig, J.F. Transition-Metal-Catalyzed Selective Functionalization
1045           of C(sp(3) )-H Bonds in Natural Products. *Angewandte Chemie* **57**, 4234-4241 (2018).

1046    69.    Stoll, S. & Schweiger, A. EasySpin, a comprehensive software package for spectral
1047           simulation and analysis in EPR. *Journal of magnetic resonance* **178**, 42-55 (2006).

1048    70.    Stoll, S. & Schweiger, A. in ESR spectroscopy in membrane biophysics, Vol. 27. (eds.
1049           M.A. Hemminga & L.J. Berliner) 299-321 (Springer, Boston, MA, 2007).

1050    71.    Prentice, J.C.A. et al. The ONETEP linear-scaling density functional theory program.
1051           *The Journal of chemical physics* **152**, 174111 (2020).

1052    72.    Baba, T. et al. Construction of *Escherichia coli K-12* in-frame, single-gene knockout
1053           mutants: the Keio collection. *Molecular systems biology* **2**, 2006 0008 (2006).

1054    73.    Hoover, J.L. et al. A Robust Pneumonia Model in Immunocompetent Rodents to
1055           Evaluate Antibacterial Efficacy against *S. pneumoniae, H. influenzae*, *K. pneumoniae*,
1056           *P. aeruginosa* or *A. baumannii. Journal of visualized experiments : JoVE* (2017).

1057    74.    Peterson, A.C., Russell, J.D., Bailey, D.J., Westphall, M.S. & Coon, J.J. Parallel
1058           reaction monitoring for high resolution and high mass accuracy quantitative, targeted
1059           proteomics. *Molecular & cellular proteomics : MCP* **11**, 1475-1488 (2012).

1060    75.    Ahrne, E. et al. Evaluation and Improvement of Quantification Accuracy in Isobaric
1061           Mass Tag-Based Protein Quantification Experiments. *Journal of proteome research*
1062           **15**, 2537-2547 (2016).

1063