Locus coeruleus spiking differently correlates with somatosensory cortex activity and pupil diameter

Hongdian Yang[1][*], Bilal A. Bari[2], Jeremiah Y. Cohen[2], Daniel H. O'Connor[2][*]

[1] Department of Molecular, Cell and Systems Biology, University of California, Riverside, CA 92521, USA
[2] Department of Neuroscience, Brain Science Institute, and Kavli Neuroscience Discovery Institute, Johns Hopkins School of Medicine, Baltimore, MD, 21205, USA
[*] Correspondence: hongdian@ucr.edu, dan.oconnor@jhmi.edu

ABSTRACT

We examined the relationships between activity in the locus coeruleus (LC), activity in the primary somatosensory cortex (S1), and pupil diameter in mice performing a tactile detection task. While LC spiking consistently preceded S1 membrane potential depolarization and pupil dilation, the correlation between S1 and pupil was more heterogeneous. Furthermore, the relationships between LC, S1 and pupil varied on timescales of sub-seconds to seconds within trials. Our data suggest that pupil diameter can be dissociated from LC spiking and cannot be used as a stationary index of LC activity.

INTRO

Multiple lines of evidence implicate the locus coeruleus/norepinephrine (LC/NE) system in perceptual task performance. First, LC activity modulates feedforward processing of sensory stimuli[1–3], and impacts sensory cortex states[4,5]. Second, LC activity correlates with task performance[6,7] and pupil diameter[7–9]. Finally, pupil diameter is thought to index arousal and has been found to be correlated with neuronal and behavioral detection or discrimination sensitivity[10–15]. Since sensory cortex activity impacts perceptual reports[16,17], these observations suggest the hypothesis that LC/NE modulates sensory cortex activity and affects perceptual task performance, and that this effect can be monitored noninvasively via the easy-to-measure pupil diameter. Testing this hypothesis requires simultaneous measurement of (1) LC activity, (2) cortical activity, ideally subthreshold membrane potential, and (3) pupil diameter, all during perceptual task performance. Here, we recorded spiking activity of optogenetically-tagged LC units together with pupil diameter in mice performing a tactile detection task[18]. In a subset of experiments, we also performed simultaneous whole-cell current clamp recordings in S1 (Fig. 1).

RESULTS

First, we report the analysis of LC and pupil recordings during behavior (e.g., Fig. 2a). Consistent with prior reports[8,9], cross-correlogram analysis revealed that LC activity and pupil diameter were correlated across entire sessions, with pupil dilation following LC spikes (peak correlation coefficient: 0.15 ± 0.02; time lags: 2.61 ± 0.39 s, n = 39, Fig. 2b). Mean LC spiking activity aligned to trial onsets showed prominent responses to a tone delivered at the beginning of each trial, as well as in trials where mice made Go responses (Hit and False Alarm trials, Fig. 2a, c). LC spiking activity to the tone was comparable to Go responses (P = 0.24, Fig. 2d, Methods). On Hit trials, where mice successfully licked to the whisker stimulus, pre-stimulus LC activity (measured in a

0.5-s window prior to stimulus onset) was slightly but significantly lower than Miss trials, where
mice failed to lick to the whisker stimulus (Fig. 2e). We note that on Miss trials LC responded
weakly to whisker stimulus alone (< 0.5 sp/s above baseline, Fig. S1). LC activity measured in a
short window (0.2-s) after stimulus onset was larger on Hits compared with Misses (Fig. 2e; the
same trend holds for 0.1-s window, data not shown). Ideal-observer analysis showed that both
pre- and post-stimulus LC activity significantly predicted perceptual reports of the mice on a trial-
by-trial basis, with choice probabilities[18] of 0.47 ± 0.014 (P = 0.032, n = 43) for pre-stimulus and
0.59 ± 0.017 (P = 4.6e-6, n = 43) for post-stimulus LC activity, respectively (Fig. 2e). LC activity
aligned to the time of licking showed that spiking responses began ~200 ms prior to licking (Fig.
2f).
In striking contrast, pupil diameter minimally increased in response to the tone. Instead,
pupil strongly dilated on Hit and False Alarm trials, in which mice made Go responses (Fig. 2a, c,
d; tone vs. Go: P = 6.4e-5, n = 36, Methods)[14]. Interestingly, pupil response to the tone was larger
on Misses compared to Hits, and significantly predicted perceptual choices of the mice (Fig. S2).
Pupil diameter changes (ΔPupil) aligned to the time of licking showed that pupil responses
occurred after licking (Fig. 2f).
Together, these data show that LC and pupil responses were positively correlated. Both
LC activity and pupil diameter increased during Go responses, but LC also strongly responded to
the tone, a salient sensory cue that alerted mice of trial onsets. Thus, LC activity and pupil
diameter appear to reflect different sets of task events during this behavior.
Next, we analyzed recordings where we simultaneously measured membrane potential
(V_m) of S1 neurons (mostly from layer 2/3, Fig. S3) along with LC spiking and/or pupil diameter
during the detection task. Our goal was to determine how LC spiking related to cortical activity
and to pupil diameter during task performance. We used spike-triggered averages (STAs) to
quantify how individual spikes from single LC units correlated with changes in V_m and pupil
diameter. LC spike-triggered V_m analyses revealed that LC spikes were associated with a small
depolarization in cortical neurons (1.39 ± 0.35 mV, n = 12, Fig. 3a-c). On average, V_m
depolarization associated with an LC spike peaked after the spike, with short time lags from an
LC spike to peak depolarization in S1 (0.17 ± 0.06 s, Fig. 3a-c, Fig. S4).
Consistent with the previous cross-correlogram analysis based on a larger set of LC-pupil
recordings (Fig. 2b), here STA analysis showed that pupil diameter increased in association with
individual spikes from LC single units (0.03 ± 0.01 mm, n = 7), with peak dilation occurring roughly
ten-fold slower than peak V_m depolarization (time lags from an LC spike to peak pupil dilation:
1.89 ± 0.25 s, Fig. 3d-f).
Given that pupil diameter and LC activity are positively correlated, and that pupil diameter
has been often considered to index LC activity[15,19], we next tested whether the pupil-S1
relationship resembled the LC-S1 relationship. Cross-correlogram analyses revealed
heterogeneous correlations between pupil diameter and S1 V_m, with both positive and negative
correlations as well as positive and negative time lags (peak correlation coefficient: 0.05 ± 0.04;
time lags: - 0.22 ± 1.01 s, n = 19, Fig. 3g-i). We further examined how well LC spiking and pupil
diameter can predict cortical V_m fluctuations at different timescales. We found that LC activity was
superior in predicting (correlating with) cortical dynamics faster than ~200-300 ms (exponential
decay time constant: 1.02 ± 0.09 s vs. 6.59 ± 0.60 s, P = 1.3e-4, n = 19, Fig. 3j).
Together, these data show that LC spikes preceded S1 depolarizations and pupil dilations.
LC spiking correlated with both V_m and pupil diameter changes, but on vastly different timescales
(~0.2 s vs. ~2 s). Our data also show that pupil diameter changes are heterogeneously correlated
with S1 V_m fluctuations (in terms of their temporal relationship and correlation strength), and can
only track slow V_m fluctuations.
Individual trials in our detection task contained distinct events, including the tone that
alerted mice of the trial start ("Tone"), the whisker stimulus on Go trials ("Stimulus"), and licks
("Lick"), as well as other periods in which mice did not receive stimuli or make lick responses

("Quiet"). For a more granular perspective on how LC spiking correlated with changes in V_m and
pupil diameter, we computed LC spike-triggered averages separately in these different event
windows (task epochs, Methods).
While single LC spikes were associated with prominent changes in both cortical V_m and
pupil diameter, we found that these associations strikingly depended on task epochs: V_m
depolarization associated with an LC spike had the biggest response to tone/licking and almost
no response during the quiet periods (Fig. 3k). In contrast, pupil dilation associated with an LC
spike had the biggest response to licking and almost no response to the tone (Fig. 3l). In addition,
peak pupil dilation and peak V_m depolarization appeared to have different dependencies on LC
spike counts, with a roughly monotonic relationship between pupil and LC, and a much weaker
dependence between V_m and LC (Fig. S5). Thus, the correlations between LC spiking and V_m,
and between LC spiking and pupil diameter, are non-stationary, even on the timescale of a few
seconds. Importantly, these epoch-dependencies were different for V_m and pupil - with the biggest
response occurring to the tone for V_m, and the smallest response occurring to the tone for pupil -
suggesting that the correlations between LC activity and V_m and pupil each reflect distinct
unmeasured factors.
118
119 DISCUSSION
120
We found that pre-stimulus baseline LC spiking predicted behavioral responses. Thus,
fluctuations in LC/NE activity may in part underlie perceptual task performance. However, the
effect was weak, possibly due to the use of an auditory cue that puts the mice in a more
homogeneous arousal state. As a result, factors other than the fluctuations of arousal also likely
contribute to cortical choice probabilities observed in prior work with this task[18]. In other tasks
without such alerting cues, task performance may have a stronger dependence on arousal and
pre-stimulus LC activity.
LC responded strongly to an auditory cue (tone) meant to alert the mice to the beginning
of a trial. While this tone carried no information about the presence of a tactile stimulus or reward
on any given trial, and therefore was not associated with a particular movement response, it did
inform the mice about the time when a tactile stimulus could occur (in our task the duration
between the tone and stimulus onset was fixed). The robust LC spiking responses to this cue are
therefore consistent with LC's role in promoting alertness or preparedness to detect a weak
stimulus. We also found that LC responded to operant licking responses, which is consistent with
earlier work showing that LC encoded overt decision execution[20].
Our data show that while LC spiking and pupil diameter correlate well at long timescales,
and both can predict changes in cortical dynamics, LC does so an order of magnitude faster.
Moreover, the correlation between pupil and V_m is much more heterogeneous than between LC
and V_m. Importantly, the relationships between LC activity, S1 V_m and pupil depended on task
epoch. Because these epochs changed on the timescale of a few seconds, our data imply that
pupil diameter can be dissociated from LC spiking and cannot be used as a stationary index of
LC activity. However, comparing across repeats of similar epochs should yield a more accurate
prediction of LC spiking by pupil diameter. That is, in attempting to use pupil diameter as a proxy
for LC spiking, our data suggest it would be useful to separately normalize distinct task epochs.
Future work should examine the LC-pupil relationship using fine-scale analyses that consider the
behavioral states at a granular level specific to individual tasks.
147
148
149 AUTHOR CONTRIBUTIONS

H.Y. performed all experiments with help from B.A.B. H.Y., B.A.B. and D.H.O. analyzed data. H.Y., J.Y.C. and D.H.O. planned the project. H.Y. and D.H.O. wrote the paper with input from B.A.B. and J.Y.C.

ACKNOWLEDGMENTS
We thank E. Zagha for comments on the manuscript; Dwight E. Bergles for DBH-Cre mice. This work was supported by UCR startup (H.Y.), Klingenstein-Simons Fellowship Awards in Neuroscience (H.Y.), NIH grants F30MH110084 (B.A.B.), 1R01NS107355 (H.Y.), 1R01NS112200 (H.Y.), R01NS089652 (D.H.O.), 1R01NS104834-01 (D.H.O., J.Y.C), and P30NS050274.

REFERENCES

1. Rodenkirch, C., Liu, Y., Schriver, B. J. & Wang, Q. Locus coeruleus activation enhances thalamic feature selectivity via norepinephrine regulation of intrathalamic circuit dynamics. *Nat. Neurosci.* **22**, 120–133 (2019).
2. Hirata, A., Aguilar, J. & Castro-Alamancos, M. A. Noradrenergic activation amplifies bottom-up and top-down signal-to-noise ratios in sensory thalamus. *J. Neurosci.* **26**, 4426–36 (2006).
3. Devilbiss, D. M., Page, M. E. & Waterhouse, B. D. Locus ceruleus regulates sensory encoding by neurons and networks in waking animals. *J. Neurosci.* **26**, 9860–72 (2006).
4. Polack, P.-O., Friedman, J. & Golshani, P. Cellular mechanisms of brain state–dependent gain modulation in visual cortex. *Nat. Neurosci.* **16**, 1331–1339 (2013).
5. Constantinople, C. M. & Bruno, R. M. Effects and mechanisms of wakefulness on local cortical networks. *Neuron* **69**, 1061–8 (2011).
6. Usher, M., Cohen, J. D., Servan-Schreiber, D., Rajkowski, J. & Aston-Jones, G. The role of locus coeruleus in the regulation of cognitive performance. *Science* **283**, 549–54 (1999).
7. Rajkowski, J., Kubiak, P. & Aston-Jones, G. Locus coeruleus activity in monkey: Phasic and tonic changes are associated with altered vigilance. *Brain Res. Bull.* **35**, 607–616 (1994).
8. Joshi, S., Li, Y., Kalwani, R. M. & Gold, J. I. Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron* **89**, 221–234 (2016).
9. Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B. & Wang, Q. Dynamic Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from Sympathetic, Not Parasympathetic, Contributions. *Cell Rep.* **20**, 3099–3112 (2017).
10. McGinley, M. J., David, S. V. & McCormick, D. A. Cortical Membrane Potential Signature of Optimal States for Sensory Signal Detection. *Neuron* **87**, 179–192 (2015).
11. Reimer, J. *et al.* Pupil Fluctuations Track Fast Switching of Cortical States during Quiet Wakefulness. *Neuron* **84**, 355–362 (2014).
12. Vinck, M., Batista-Brito, R., Knoblich, U. & Cardin, J. A. Arousal and Locomotion Make Distinct Contributions to Cortical Activity Patterns and Visual Encoding. *Neuron* **86**, 740–754 (2015).
13. Schriver, B. J., Bagdasarov, S. & Wang, Q. Pupil-linked arousal modulates behavior in rats performing a whisker deflection direction discrimination task. *J. Neurophysiol.* **120**, 1655–1670 (2018).
14. Lee, C. R. & Margolis, D. J. Pupil Dynamics Reflect Behavioral Choice and Learning in a Go/NoGo Tactile Decision-Making Task in Mice. *Front. Behav. Neurosci.* **10**, 1–14 (2016).
15. McGinley, M. J. *et al.* Waking State: Rapid Variations Modulate Neural and Behavioral

Responses. *Neuron* **87**, 1143–1161 (2015).

16. Sachidhanandam, S., Sreenivasan, V., Kyriakatos, A., Kremer, Y. & Petersen, C. C. H.
Membrane potential correlates of sensory perception in mouse barrel cortex. *Nat.*
Neurosci. **16**, 1671–1677 (2013).

17. Miyashita, T. & Feldman, D. E. Behavioral detection of passive whisker stimuli requires
somatosensory cortex. *Cereb. Cortex* **23**, 1655–1662 (2013).

18. Yang, H., Kwon, S. E., Severson, K. S. & O'Connor, D. H. Origins of choice-related
activity in mouse somatosensory cortex. *Nat. Neurosci.* **19**, 127–134 (2016).

19. Aston-Jones, G. & Cohen, J. D. An integrative theory of locus coeruleus-norepinephrine
function: adaptive gain and optimal performance. *Annu. Rev. Neurosci.* **28**, 403–50
(2005).

20. Kalwani, R. M., Joshi, S. & Gold, J. I. Phasic Activation of Individual Neurons in the Locus
Ceruleus/Subceruleus Complex of Monkeys Reflects Rewarded Decisions to Go But Not
Stop. *J. Neurosci.* **34**, 13656–13669 (2014).

METHODS:

Mice were DBH-Cre (B6.FVB(Cg)-Tg(Dbh-cre) KH212Gsat/Mmucd, 036778-UCD, MMRRC);
Ai32 (RCL-ChR2(H134R)/EYFP, 012569, JAX), singly housed in a vivarium with reverse light-
dark cycle (12 hr each phase). Male and female mice of 6-12 weeks were implanted with
titanium head posts as described previously[18]. After recovery, mice were trained to perform a
Go/NoGo single whisker detection task as described previously[18].

Custom microdrives with eight tetrodes and an optic fiber[21] (0.39 NA, 200 um core) were built to
make extracellular recordings from LC neurons. Each tetrode comprised four nichrome wires
(100-300 KΩ). A ~1 mm diameter craniotomy was made (centered at -5.2 mm caudal and 0.85
mm lateral relative to bregma) for implanting the tetrodes to a depth of 2.7 mm relative to the
brain surface. The microdrive was advanced in steps of ~100 um each day until reaching LC,
identified by optogenetic tagging of DBH+ neurons expressing ChR2, tail pinch response, wide
extracellular spike waveforms and post-hoc electrolytic lesions. Broadband voltage traces were
acquired at 30 kHz (Intan Technologies), and filtered between 0.1 and 10 kHz. Signals were
then bandpass filtered between 300 and 6000 Hz, and spikes were detected using a threshold
of 4-6 standard deviations. The timestamp of the peak of each detected spike, as well as a 1-ms
waveform centered at the peak were extracted from each channel for offline spike sorting using
MClust[22]. At the conclusion of the experiments, brains were perfused with PBS followed by 4%
PFA, post-fixed overnight, then cut into 100 μm coronal sections and stained with anti-Tyrosine
Hydroxylase (TH) antibody (Millipore AB152).

Pupil video was acquired at 50 Hz using a PhotonFocus camera and StreamPix 5 software.
Light from a 940 LED was passed through a condenser lens and directed to the right eye,
reflected off a mirror, and directed into a 0.25X telecentric lens. WaveSurfer
(https://www.janelia.org/open-science/wavesurfer) triggered individual camera frames
synchronized with electrophysiological recordings.

In a subset of animals, we performed simultaneous intracellular current clamp (whole-cell)
recordings in conjunction with LC recording and/or pupil tracking during behavior. A craniotomy
over the C2 barrel was made based on intrinsic signal imaging[18]. In some cases, we also made
craniotomies over nearby barrels based on the known somatotopy of S1[23,24] to increase yield.

Whole-cell recording procedures, quality control and data processing were performed as
described previously[18].
252
For Fig. 2d, LC responses to the tone were calculated using a 300-ms window starting at tone
onset, and LC responses to Go were calculated using a 300-ms window starting 200 ms after
stimulus onset to capture peak responses. These estimates were based on LC response
profiles in Fig. 2c. Pupil responses to the tone were calculated using a 1-s window starting 1 s
after tone onset. This estimate was primarily based on pupil response profile during CR trials
(e.g., Fig. 2a, c, indicated by the grey bar), where there was no whisker stimulus or licking
response. Pupil responses to Go were calculated using a 1-s window starting 1.5 s after
stimulus onset (e.g., FA trials in Fig. 2a, c, indicated by the black bar). Based on the temporal
profiles of pupil diameter in different trial types shown in Fig. 2a, c, and that the whisker stimulus
started 1 s after tone onset, pupil responses to tone and Go can be segregated. These
estimates were consistent with the results showing that pupil dilated 1-2 s after LC spikes (Fig.
2b, and Fig. 3d-f).
265
266   For Fig. 2e, pre-stimulus LC baseline activity was calculated using a 500-ms window ending 50
267   ms before stimulus onset. Post-stimulus activity was calculated using a 200-ms window starting
268   20 ms after stimulus onset, before licking responses[18]. Choice probabilities were computed as
269   described previously[18].
270
271   To compute lick-aligned changes in LC spiking and pupil diameter, we only used licks that
272   occurred at least 0.5 s after the previous lick. To compute LC spike triggered S1 V_m and pupil,
273   we only used LC spikes that occurred at least 0.5 s after the previous spike. For STA analysis,
274   peak ΔV_m or ΔPupil was defined as the largest positive or negative value within the observed
275   window (± 1 s or ± 10 s, respectively).
276
For cross-correlogram analysis, each LC spike train was convolved with a 400-ms wide
Gaussian kernel. Peak correlation coefficients were defined as the largest positive or negative
value within the observed window (± 1 s or ± 10 s). To examine how well LC spiking and pupil
diameter could predict cortical V_m fluctuations at different timescales (Fig. 3j), V_m was high-pass
filtered at 0, 0.2, 0.4, 0.6, 0.8, 1, 2, 3, 4 and 5 Hz separately. Cross-correlogram analysis
between the filtered V_m and LC (pupil) activity was then performed as described above, and
absolute values of peak correlation coefficients were used.
284
285   Task epochs were defined as: "Tone" epochs: -0.2 s to 0.3 s with respect to tone onset;
286   "Stimulus" epochs: -0.2 s to 0.3 s with respect to stimulus onset (i.e., only on Go trials); "Licking"
287   epochs: -0.2 s to 0.3 s with respect to licks that occurred at least 0.5 s after the previous lick;
288   "Quiet" epochs: non-overlapping 0.5 s segments excluding the three types of epoch defined
289   previously during the entire session.
290
Thirty-nine LC-pupil pairs were included in Fig. 2b, including single- and multi-units, with and
without S1 recordings. For the rest of Fig. 2, LC analysis included forty-three recordings, each
with at least 4 Hit and 4 Miss trials. Among those, thirty-six were with pupil recordings, and were
used for pupil analysis. Twelve pairs of S1 whole-cell and LC single-unit recordings were
included in Fig. 3a-c, k, seven of which were with pupil recordings and included in Fig. 3d-f.
Nineteen S1- -pupil recordings were included in Fig. 3g-j. Twenty pairs of LC SU and pupil
recordings were included in Fig. 3l, with and without S1 recordings.
298

299 Data were reported as mean ± s.e.m. unless otherwise noted. Statistical tests were by two-tailed
300 Wilcoxon signed rank unless otherwise noted.
301

21. Cohen, J. Y., Haesler, S., Vong, L., Lowell, B. B. & Uchida, N. Neuron-type-specific
signals for reward and punishment in the ventral tegmental area. *Nature* **482**, 85–8
(2012).
22. Redish, A. D. MClust Spike sorting toolbox Documentation for version 4.4. (2014).
Available at: http://redishlab.neuroscience.umn.edu/MClust/MClust-4.4.pdf.
23. Welker, C. & Woolsey, T. A. Structure of layer IV in the somatosensory neocortex of the
rat: Description and comparison with the mouse. *J. Comp. Neurol.* **158**, 437–453 (1974).
24. Wilson, M. a, Johnston, M. V, Goldstein, G. W. & Blue, M. E. Neonatal lead exposure
impairs development of rodent barrel field cortex. *Proc. Natl. Acad. Sci. U. S. A.* **97**,
5540–5545 (2000).

FIGURE 1: Cortical membrane potential, LC spike rate, and pupil recorded during a tactile detection task.

(a) Task schematic, trial structure and all trial types of the single-whisker detection task[18]. (b) Schematic of tetrode recording in LC, whole-cell recording in S,1 and pupil tracking during the task.

(c) Expression of ChR2 in a Dbh;Ai32 mouse. (ChR2-EYFP: green; Tyrosine Hydroxylase, TH: red).

(d) Left: Responses of a ChR2-expressing LC unit to opto-tagging (lightning bolts: blue light pulses) and tail pinch. Middle: LC unit responses to 12 blue light pulses (200-ms) aligned to individual pulse onset. Ticks represent spikes. PSTH is shown at the bottom. Right: Typical wide

waveforms of LC units and an electrolytic lesion (arrow: lesion site) in the LC (white) showing the recording location.

(e) Example simultaneously recorded LC activity, S1 V_m, and pupil.

FIGURE 2: LC and pupil responses during behavior.
(a) Example LC recording with pupil tracking. Left: LC spike raster separated by trial types. Right: Mean pupil diameter (± s.e.m.) separated by trial types. Grey and black arrows indicate tone and stimulus onsets, respectively. Grey and black bars indicate the time windows during which pupil responses to tone and to Go (behavioral responses) were quantified, respectively. We note that based on the temporal profiles of pupil diameter in different trial types (i.e., in the presence or absence of tactile stimulus or licking), and that tactile stimulus starts 1 s after tone onset, pupil responses to tone and Go can be segregated (Methods).
(b) Top: Cross-correlogram between LC spike train and pupil diameter. Individual LC spikes were convolved with a 400-ms wide Gaussian kernel. Bottom: Histogram of peak correlation coefficient (left), and time lags (right) between LC spike train and pupil diameter for each paired recording (magenta dot: mean). Both distributions are significantly larger than 0 (peak correlation coefficient: 0.15 ± 0.02, P = 8.3e-7; time lags: 2.61 ± 0.39 s, P = 7.8e-7, n = 39).
(c) Trial-aligned LC spike rate (top), and pupil diameter (bottom) averaged by different trial types. Grey and black arrows indicate tone and stimulus onsets, respectively.
(d) Left: LC responses to tone (T) and Go responses (G) during Hit trials with median indicated. Tone vs. Go: 4.79 (3.70 – 6.66) sp/s vs. 4.68 (3.33 – 7.26) sp/s, median (IQR), P = 0.24, n = 43. Right: Pupil responses to tone and Go responses during Hit trials with median indicated. Tone vs. Go: 0.003 (-0.015 – 0.015) mm vs. 0.027 (-0.010 – 0.063) mm, median (IQR), P = 6.4e-5, n = 36. Grey lines indicate individual recordings.
(e) Top: Pre-stimulus (baseline) and post-stimulus (evoked) LC spike rate for Hit and Miss trials with median indicated (Baseline: Hit vs. Miss, 0.66 (0.30 – 3.51) sp/s vs. 1.55 (0.68 – 3.00) sp/s, median (IQR), P = 0.0083; Evoked: Hit vs. Miss, 3.24 (1.78 – 5.49) sp/s vs. 1.82 (0.95 – 3.45) sp/s, median (IQR), P = 5.5e-7, n = 43). Grey lines indicate individual recordings. Bottom: Histogram of choice probability for Hit vs. Miss trials based on baseline and evoked LC activity (magenta dots: mean). Choice probabilities are significantly deviated from 0.5. Baseline: 0.47 ± 0.014, P = 0.032; Evoked: 0.59 ± 0.017, P = 4.6e-6, n = 43.
(f) Lick-aligned LC spike rate (top) and pupil diameter (ΔPupil, bottom) averaged by trial types: Hit (blue), FA (green).

FIGURE 3: Correlations between LC spikes, S1 V_m and pupil diameter depend on task epoch.

(a) Two examples of LC spike-triggered average ΔV_m.

(b) Group mean of LC spike-triggered average ΔV_m (± s.e.m., n = 12)

(c) Histograms of peak ΔV_m and peak lags (showing all LC-S1 pairs) with means indicated (magenta dots). Both distributions are significantly larger than 0 (Peak ΔV_m: 1.39 ± 0.35 mV, P = 4.9e-4; Peak lags: 0.17 ± 0.06 s, P = 4.9e-4, n = 12).

(d) Two examples of LC spike-triggered average ΔPupil.

(e) LC spike-triggered average ΔPupil group mean (± s.e.m., n = 7).

(f) Histograms of peak ΔPupil and peak lags (showing all LC-Pupil pairs) with means indicated (magenta dots). Both distributions are significantly larger than 0 (Peak ΔPupil: 0.03 ± 0.01 mm, P = 0.016; Peak lags: 1.89 ± 0.25 s, P = 0.016, n = 7).

(g) Two examples of Pupil-V_m cross-correlograms.

(h) Group mean of Pupil-V_m cross-correlograms (± s.e.m., n = 19).

(i) Histograms of peak Pupil-V_m correlation coefficient and peak lags (showing all S1-Pupil pairs) with means indicated (magenta dots). Both distributions are not significantly deviated from 0 (Peak correlation coefficient: 0.05 ± 0.04, P = 0.33; Peak lags: - 0.22 ± 1.01 s, P = 0.87, n = 19).

(j) Top: Peak correlation coefficient for LC-V_m and Pupil-V_m pairs after progressive high-pass filtering of S1 V_m. Bottom: Exponential curve fitted time constants for Pupil-V_m are larger than LC-V_m (1.02 ± 0.09 s vs. 6.59 ± 0.60 s, P = 1.3e-4, n = 19).

(k) Left: LC spike-triggered ΔV_m separated by task epoch: tone, stimulus, lick and quiet. Right: Bar graphs of peak ΔV_m for each epoch. Dots indicate individual paired recordings. Repeated-measure ANOVA, P = 1.4e-4, n = 12. Post-hoc Tukey-Kramer tests revealed that peak ΔV_m in lick, stimulus and tone epochs were not different from each other. Lick vs. Stim, P = 1.00; Lick vs. Tone, P = 0.76; Stim vs. Tone, P = 0.94. Peak ΔV_m in quiet epochs was lower. Quiet vs. Lick, P = 0.0059; Quiet vs. Stim, P = 0.0038; Quiet vs. Tone, P = 0.0041.

(l) Left: LC spike-triggered ΔPupil separated by task epoch. Right: Bar graphs of peak ΔPupil for each epoch. Dots indicate individual paired recordings. Repeated-measure ANOVA, P = 1.3e-9, n = 20. Post-hoc Tukey-Kramer tests revealed that peak ΔPupil in lick and stimulus epochs were larger than in tone and quiet epochs. Lick vs. Stim, P = 0.10; Tone vs. Quiet, P = 0.76; Lick vs. Tone, P = 3.7e-7; Lick vs. Quiet, P = 6.2e-4; Stim vs. Tone, P = 1.1e-4; Stim vs. Quiet, P = 0.0027.

Figure S1. LC responded minimally to whisker stimulation when mice did not make a licking response (Miss trials, Baseline vs. Evoked: 1.55 (0.68-3.00) sp/s vs. 1.82 (0.95-3.45) sp/s, median (IQR), P = 0.02, n = 43).

Figure S2. (a) Pupil responses to the tone for Hit and Miss trials with median indicated. Hit vs. Miss, 0.003 (-0.015 – 0.015) mm vs. 0.0083 (-0.0005 – 0.029) mm, median (IQR), P = 0.0062, n = 36. Grey lines indicate individual recordings. (b) Histogram of choice probability for Hit vs. Miss trials based on pupil responses to the tone (magenta dot: mean). Choice probability is significantly deviated from 0.5 (0.44 ± 0.021, P = 0.0036, n = 36).

Figure S3. Histograms of the depth of S1 whole-cell recordings. Red: 12 S1 recordings included in the LC-S1 pairs in Fig. 3a-c, 3k. Grey: 19 S1 recordings included in the Pupil-S1 pairs in Fig. 3g-j.

Figure S4. (a) Cross-correlogram between LC spike train and S1 V_m. Individual LC spikes were convolved with a 400-ms wide Gaussian kernel.
(b) Histogram of peak correlation coefficient (left), and time lag (right) between LC spike train and S1 V_m for each paired recording (magenta dot: mean).

Figure S5. (a) Left: Peak ΔV_m vs. LC spike counts by epochs. Right: Cumulative histograms showing numbers of trials that go into the plots when broken down by LC spike counts. (b) Left: Peak ΔPupil vs. LC spike counts by epochs. Right: Cumulative histograms showing numbers of trials that go into the plots when broken down by LC spike counts.