Multiple spatial behaviours govern social network positions in a wild ungulate

Gregory F Albery[1,2*], Alison Morris[2], Sean Morris[2], Josephine M Pemberton[2], Tim H. Clutton-Brock[2,3], Daniel H Nussey[2], Josh A Firth[4,5]

1. Department of Biology, Georgetown University, Washington, DC
2. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, UK
3. Department of Zoology, University of Cambridge, Cambridge, UK
4. Department of Zoology, University of Oxford, Oxford, UK
5. Merton College, University of Oxford, Oxford, UK

*gfalbery@gmail.com

Abstract

The structure of wild animal social systems depends on a complex combination of intrinsic and extrinsic drivers. Population structuring and spatial behaviour are key determinants of individuals' observed social behaviour, but quantifying these spatial components alongside multiple other drivers remains difficult due to data scarcity and analytical complexity. We used a 43-year dataset detailing a wild red deer population to investigate how individuals' spatial behaviours drive social network positioning, while simultaneously assessing other potential contributing factors. Using Integrated Nested Laplace Approximation (INLA) multi-matrix animal models, we demonstrate that social network positions are shaped by two-dimensional landscape locations, pairwise space sharing, individual range size, and spatial and temporal variation in population density, alongside smaller but detectable impacts of a selection of individual-level phenotypic traits. These results indicate strong, multifaceted spatiotemporal structuring in this society, emphasising the importance of considering multiple spatial components when investigating the causes and consequences of sociality.

Word Count: **Abstract**: 143 words; **Main text**: 4720 words
Number of figures: 4; Number of tables: 0; Number of references: 53

Authorship Statement
GFA conceived the study, analysed the data, and wrote the manuscript, advised by JAF. AM and SM collected the data. JAF, JMP, THCB, and DN commented on the manuscript.

Data Accessibility Statement
The code used here is available at https://github.com/gfalbery/Spocial_Deer. On acceptance, the data will be uploaded to the same repo, which will be archived on Zenodo.

Introduction

Social behaviour is an integral component of an animal's phenotype, driving processes including disease transmission, mating, learning, and selection (Croft *et al.* 2008; VanderWaal *et al.* 2014; Krause *et al.* 2015; Firth *et al.* 2018; Sah *et al.* 2018; Silk *et al.* 2019; Firth 2020). Contemporary studies of animal behaviour often use social networks to derive individual-level social network positions, under the notion that between-individual variation in network positioning is indicative of between-individual variation in social behaviour (Franks *et al.* 2010; Krause *et al.* 2015; Sosa *et al.* 2020). However, an animal's position in its social network is also dependent on its own spatial behaviour (Webber & Vander Wal 2018; Albery *et al.* 2020a), and on a range of extrinsic factors: demography determines local population density and structuring (Shizuka & Johnson 2019)**,** while the environment shapes resource distributions, movement corridors, and emergent patterns of space use, all of which will influence the architecture of the social system (Firth & Sheldon 2016; Webber & Vander Wal 2018; Farine & Sheldon 2019; He *et al.* 2019). As such, it is important to consider spatial behaviour and environmental context when assessing the causes and consequences of individual-level social network positioning (Webber & Vander Wal 2018; He *et al.* 2019; Albery *et al.* 2020a), yet doing so remains difficult in most systems due to the complexity of spatial-social analyses that incorporate these processes.

The spatial drivers of social network structure are poorly understood because they are highly multivariate and (therefore) difficult to analyse. On the one hand, there is strong support for simpler "first-order" associations between spatial and social behaviour. For example, spatial proximity and social connections are often correlated, because individuals that share more space are more likely to associate or interact. This finding holds for diverse taxa including elk (Vander Wal *et al.* 2014), raccoons (Robert *et al.* 2012), birds (Firth & Sheldon 2016), and myriad other systems. Similarly, spatial and social network centrality are occasionally found to correlate (Mourier *et al.* 2019), as are temporal variation in population density and social contact rates (Sanchez & Hudgens 2015), and the social environment can drive spatial behaviour (Firth & Sheldon 2016; Spiegel *et al.* 2016). Spatial behaviours can be summarised using a wide range of metrics, including individuals' spatial activity levels (e.g. home range area), pairwise space sharing (e.g. distances or home range overlaps), demographic structure (e.g. temporal population size or local conspecific density), and point location on the two-dimensional landscape. For example, are more social individuals simply wider-ranging, leading them to make more contacts? Do they most often inhabit areas of high population density or well-used movement corridors? These variable spatial components take a combination of different data structures, and are therefore difficult to include in the same models, particularly in large numbers and alongside a range of other

individual-level phenotypes. It is therefore unclear to what extent individuals' social network

positions emerge from 1) their own social behaviour; 2) their own spatial behaviour; 3) their

situation within the population and broader social network; 4) other aspects of their biotic and

abiotic environment such as landscape structure; and 5) intrinsic phenotypic traits that

researchers are commonly interested in investigating.

Several frameworks have been proposed to facilitate the untangling of spatial and social

processes in wild animals (Jacoby & Freeman 2016; Silk *et al.* 2018, 2019; Webber &

Vander Wal 2018; Mourier *et al.* 2019; Albery *et al.* 2020a). To date, statistical methodology

focusses on incorporating spatial behaviours into the node-and-edge structure of network

data, using e.g. null network permutations (Firth & Sheldon 2016), spatially embedded

networks (Daraganova *et al.* 2012), and nested "networks of networks" composed of

movement trajectories (Mourier *et al.* 2019). Many such analyses involve reducing

movement patterns into some form of spatial network based on home range overlap or

spatial proximity between dyads (Mourier *et al.* 2019). For example, statistical models

named "animal models" can examine spatial variation by fitting such matrices as variance

components, potentially alongside other dyadic similarity matrices (i.e., pairwise measures of

similarity), to quantify genetic and non-genetic contributions to individuals' phenotypes

(Kruuk 2004; Stopher *et al.* 2012b; Regan *et al.* 2016; Thomson *et al.* 2018; Webber &

Vander Wal 2018). As yet, the focus on controlling for spatial autocorrelation using space

sharing and network permutations has contributed to a lack of clarity concerning the role that

spatial behaviour and environmental context play in driving social network positioning

(Albery *et al.* 2020a).

Studies across ecological disciplines increasingly use Integrated Nested Laplace

Approximation (INLA) models to control for spatial autocorrelation in a multitude of contexts

(Lindgren *et al.* 2011; Lindgren & Rue 2015; Zuur *et al.* 2017). As well as including fixed and

random effects to quantify individual-level drivers, these models can incorporate dyadic

space sharing components (Holand *et al.* 2013) and stochastic partial differentiation

equation (SPDE) effects to model 2-dimensional spatial patterns in the response variable,

thereby controlling for and estimating spatiotemporal variation associated with fine-scale

positioning within the landscape (Albery *et al.* 2019). As such, these models offer an exciting

opportunity to test and compare the roles of a range of spatial behaviours and

autocorrelation structures, alongside phenotypic drivers, in determining social network

positioning.

We address this question using the long-term study in the Isle of Rum red deer (*Cervus*

elaphus). These study animals comprise an unmanaged wild population with a contiguous

fission-fusion social system (Clutton-Brock *et al.* 1982). They experience strong

environmental gradients and exhibit spatial autocorrelation in a number of important

phenotypes: individuals with greater home range overlap have more similar behavioural and

life history traits (Stopher *et al.* 2012b), and those in closer proximity have more similar

parasite burdens (Albery *et al.* 2019); further, as with other matrilineal mammalian systems,

closely related individuals frequently associate (Clutton-Brock *et al.* 1982) and live closer

together (Stopher *et al.* 2012b). Individuals have highly repeatable home ranges (Stopher *et*

al. 2012b) that decline in size over their lifetimes, predicting declining survival probability

(Froy *et al.* 2018). The study area has a strong spatial gradient in resource availability, with

high-quality grazing heavily concentrated in the far north of the system, and with most

individuals aggregating around this area, such that population density decreases outwards

towards the edge of the study population (Clutton-Brock *et al.* 1982). As such, the deer

comprise an ideal system for assessing spatial-social relationships in the wild.

To assess how individuals' spatial behaviours translate to social network positions, we

constructed fine-scale social networks from 43 years of censuses of the study population.

We derived 8 different individual-level network positioning measures of varying complexity

that are important to different social processes (Krause *et al.* 2015; Sosa *et al.* 2020). Using

multi-matrix animal models in INLA, we examined whether spatial locations, space sharing,

home range area, and local population density explained variation in network position

metrics, alongside a range of individual-, temporal-, and population-level factors. Specifically,

we aimed to test two hypotheses: that the structure of the social network would be highly

dependent on the distribution of population density in space; and that individuals' social

network centrality would be largely explained by their ranging behaviour, where wide-ranging

individuals were more likely to be socially well-connected. We further expected that space

sharing and point locations would uncover substantial spatial autocorrelation in social

network positioning, and that different social network metrics would exhibit different spatial

patterns and vary drivers. This not only comprises a large-scale empirical examination of the

factors shaping social network positions in this extensively monitored wild mammal, but also

provides a methodological advancement in developing powerful, flexible new methods

(INLA-based multi-matrix animal models) with broad potential for examining spatial-social

processes in this and other systems.

Methods

Study system and censusing

The study was carried out on an unpredated long-term study population of red deer on the

Isle of Rum, Scotland (57°N,6°20'W). The natural history of this matrilineal mammalian

system has been studied extensively (Clutton-Brock et al. 1982), and we focussed on
females aged 3+ years, as these individuals have the most complete associated census
data, and few males live in the study area except during the mating period. Individuals are
monitored from birth, providing substantial life history and behavioural data, and >90% of
calves are caught and tagged, with tissue samples taken (Clutton-Brock *et al.* 1982). The
population thus has comprehensive genomic data, allowing high-powered quantitative
genetic analyses: most individuals born since 1982 have been genotyped at >37,000 SNPs,
distributed throughout the genome (e.g. Huisman, Kruuk, Ellis, Clutton-Brock, & Pemberton,
2016). Census data were collected for the years 1974-2017, totalling 423,070 census
observations. Deer were censused by field workers five times a month, for eight months of
the year, along one of two alternating routes (Clutton-Brock *et al.* 1982). Individuals'
identities, locations (to the nearest 100M), and group membership were recorded. Grouping
events were estimated by seasoned field workers according to a variant of the "chain rule"
(e.g. Castles et al., 2014), where individuals grazing in a contiguous group within close
proximity of each other (under ~10 metres) were deemed to be associating, with mean 130.4
groups observed per individual across their lifetime (range 6-943). The mortality period falls
between Jan-March, when there is least available food, and minimal mortality occurs outside
this period. We only used census records in each May-December period, from which we
derived annual social network position measures as response variables (Figure 1-2). We
elected to investigate this seasonal period because it stretches from the spring calving
period until the beginning of the mortality period, simplifying network construction and
avoiding complications arising from mortality events. Our dataset totalled 3356 annual
observations among 532 grown females (Figure 1).

In this system, female reproduction imposes substantial costs for immunity and parasitism
(Albery *et al.* 2020c), and for subsequent survival and reproduction (Clutton-Brock, Albon, &
Guinness, 1989; Froy, Walling, Pemberton, Clutton-Brock, & Kruuk, 2016). If a female
reproduces, she produces 1 calf per year in the spring, generally beginning in May; the "deer
year" begins on May 1 for this reason. Here, reproductive status was classified into the
following four categories using behavioural observations: True Yeld (did not give birth);
Summer Yeld (the female's calf died in the summer, before 1[st] October); Winter Yeld (the
female's calf died in the winter, after 1[st] October); and Milk (calf survived to 1[st] May the
following calendar year).

Generating spatial and social matrices

All code is available online at https://github.com/gfalbery/Spacial_Deer. We constructed the
home range overlap (HRO) matrix using the R package AdeHabitatHR (Calenge 2011),
following previous methodology (Stopher *et al.* 2012b; Regan *et al.* 2016; Froy *et al.* 2018).

First, using a kernel density estimation method, we derived lifetime home ranges for each

individual with more than five census observations. Previous analysis has shown that this

system is robust to the number of observations used to generate home ranges (Froy *et al.*

2018). We used lifetime home ranges to fit one value per individual in the animal models;

individual ranges (and range sizes) correlate strongly from year to year (Stopher *et al.*

2012b; Froy *et al.* 2018). We derived proportional HRO of each dyad using Bhattacharya

Affinity (following Stopher *et al.* 2012b), producing values between 0-1 (i.e. no overlap to

complete overlap).

To control for individuals' two-dimensional point locations, we used a Stochastic Partial

Differentiation Equation (SPDE) effect in INLA. This effect models the distance between

points to calculate spatial autocorrelation, using Matern covariance (Lindgren *et al.* 2011).

This random effect used individuals' annual centroids (mean easting and northing in a given

year) or lifetime centroids (mean easting and northing across all observations) as point

locations to approximate spatial variation in the response variable (Lindgren *et al.* 2011;

Albery *et al.* 2019).

We used a genomic relatedness matrix (GRM) using homozygosity at 37,000 Single

Nucleotide Polymorphisms, scaled at the population level (Yang *et al.* 2011; for a population-

specific summary, see Huisman *et al.* 2016). This matrix is well-correlated with pedigree-

derived relatedness metrics (Huisman *et al.* 2016). HRO was well-correlated with distance

between lifetime centroids (i.e., closer individuals tended to share more range), and both

were weakly but significantly correlated with genetic relatedness (Supplementary Figure 1).

To test whether social network positions could be explained by population density, we

derived the local density of individuals again using AdeHabitatHR (Calenge 2011). We

generated density kernels of observations, and then assigned individual deer their local

population density based on their location on this kernel, following previous methodology

developed in badgers (Albery *et al.* 2020b). This local density value was then fitted as a fixed

explanatory variable. We used four different density metrics, each examining the density of a

different observation type: lifetime centroids ("lifetime density"); annual centroids ("annual

density"); all observations across the study period ("sighting density"); and all observations in

the focal year ("annual sighting density"). Only one such density metric was fitted at once.

We also calculated annual home range areas (HRA) by taking the 70% isopleth of each

individual's annual space use distribution, following previous methodology (Froy *et al.* 2018).

This HRA variable was fitted as a fixed effect in the same way as local density.

We constructed a series of 43 annual social networks using "gambit of the group," where

individuals in the same grouping event (as described above) were taken to be associating

(Franks *et al.* 2010). Dyadic associations were calculated using the 'simple ratio index'

(Cairns & Schwager 1987) derived as a proportion of total sightings (grouping events) in

which the focal individuals were seen together: $Sightings_{A,B}/(Sightings_A+Sightings_B-$

$Sightings_{A,B})$, or $Intersect_{A,B}/Union_{A,B}$. In this dyadic matrix, 0=never seen together and

1=never seen apart.

Statistical Analysis

Metrics. Using the annual social networks, we derived eight individual-level network metrics

which are commonly used across animal social networks and have been considered in

detail(Whitehead 2008; Brent 2015; Krause *et al.* 2015; Firth *et al.* 2017). We set each of

these network metrics for use as response variables in separate INLA Generalised Linear

Mixed Models (GLMMs) with a Gaussian family specification. In increasing order of

complexity, our measures included four direct sociality metrics, which only take into account

an individual's connections with other individuals: 1) Group Size – the average number of

individuals a deer associated with per sighting; 2) Degree – the number of unique individuals

she was observed with; 3) Strength – sum of all their weighted social associations to others;

4) Mean Strength – the average association strength to each of the unique individuals she

was observed with (equivalent to strength divided by degree). We also included four more

complex "indirect" metrics (all using algorithms as specified from (Csardi & Nepusz 2006)),

which also take into account an individual's connections' connections: 5) Eigenvector

centrality – which considers the sum of their own connections and the sum of their

associates' connections; 6) Weighted Eigenvector – which is akin to eigenvector centrality

but also accounts for the weights of theirs, and their associates, connections; 7)

Betweenness – the number of shortest paths that pass through the focal individual to

traverse the whole network; 8) Clustering (local) – the tendency for an individual's contacts

to be connected to one another, forming triads. The raw, untransformed correlations were

assessed for all metrics, and R lay between -0.5 and 0.879 across metrics (Supplementary

Figure 2). When modelling them as response variables, to approximate normality, all social

metrics were square root-transformed apart from eigenvector centralities (which were left

untransformed), group size (which was cube root-transformed), and betweenness (which

was log(X+1)-transformed). Each social network metric was fitted as a response variable in

a separate model set (as outlined conceptually in Figure 1).

Base model structure. We ensured that all models followed the same base structure.

Random effects included individual identity and year (categorical random intercepts), as well

as the genetic relatedness matrix. Fixed effects included Age (continuous, in years),

Reproductive Status (four categories: True Yeld; Summer Yeld; Winter Yeld; and Milk), and

Number of observations (continuous, log-transformed to approximate normality), as well as

year-level continuous factors including Year (continuous) and that year's study Population

Size (log-transformed). All continuous response and explanatory variables were

standardised to have a mean of zero and a standard deviation of 1. Fixed effect estimates

were provided by the mean and 95% credibility intervals of the posterior estimate

distribution.

Adding spatial components. To investigate the divergent value of different spatial

behaviours, we iteratively added spatial effects to the base model, investigating which

behaviours best fit the data. These spatial behaviours corresponded to four broad

components in Figure 1: space sharing (HRO matrix); home range area (HRA); point

locations (SPDE effect); and local population density (density fixed effect). For space

sharing, we only used one metric: lifetime HRO (see above). For point locations, we selected

between 1) lifetime centroids; 2) annual centroids; and 3) annual centroids with a

spatiotemporally varying annual field. For density, we used the four metrics outlined above

("lifetime", "annual", "sighting", and "annual sighting" density). To distinguish between

competitive models we used Deviance Information Criterion (DIC). In each round, we added

each spatial behaviour individually and then kept the best-fitting one, until all four had been

added or their addition did not improve the model, using a cutoff of 2 DIC.

Comparing all spatial and non-spatial drivers. To compare the relative importance of all

fixed and random effects, we examined the model's predicted values and their correlations

with the observed values, representing the proportion of the variance that was explained by

the model (i.e., R^2). We used the model to predict each social behaviour metric, and

iteratively held each explanatory variable's predictions at the mean, one at a time. We then

assessed the squared correlations of these values with the observed values, relative to

those of the full model. Variables with greater effects in the model produced less accurate

predicted values when held constant.

Results

Spatial behaviours were important in determining all eight individual-level social network

position variables. The non-spatial model was far the poorest-fitting for all eight metrics, and

the DIC changes associated with adding spatial components were substantial (Figure 3A).

Generally, wide-ranging individuals and those living in areas of greater population density

tended to be more central, and space sharing and point location effects both revealed

substantial spatial autocorrelation (Figure 3).

As expected, home range area and population density had substantial effects on social

network centrality (Figure 3). Population density was positively associated with all centrality

measures except betweenness and clustering (Figure 3A), and the best-fitting density metric

was annual density. Individuals with larger home ranges likewise tended to be more social,

except in the case of clustering (no effect) and mean strength, which were negatively

associated with HRA (Figure 3A).

Notably, point location-based SPDE effects tended to improve model fit over these fixed

effects, and had a greater effect than on model fit space sharing HRO effects, even when

conceptualised at the same timescale (i.e., across the individual's lifetime). Investigating the

R^2 components of the models containing only HRO (i.e., without SPDE effects) revealed that

in general spatial overlap accounted for more variation than the genetic matrix

(Supplementary Figure 3), but comparing these with the other models revealed that the point

location effects contributed more than either of these matrices (Figure 3B). Annually varying

centroids further improved model fit, and allowing the spatial field to vary between years in

our spatiotemporal models improved models even more (Figure 3A). Although the space

sharing and genomic relatedness matrices had similar sized impacts on the full models

(Figure 3B), removing the SPDE effect resulted in a substantial increase in the HRO effect,

but with very little impact on the GRM's R^2 (Supplementary Figure 3). These findings were

relatively consistent across all metrics (Figure 3A-B), although the SPDE effect was notably

smaller for betweenness (Figure 3B). Taken together, these results reveal that lifetime space

sharing was good at accounting for variation in social behaviour, but that its effect was

surpassed by increasingly complex temporal formulations of point location effects.

We compared the importance of all fixed and random effects by predicting selectively from

the model, revealing overwhelmingly strong effects of spatiotemporal drivers (Figure 3B).

Our models fit well and explained a substantial amount of variation in social network

centrality (>70%); the majority of the fit was lent by a combination of the INLA SPDE effect,

fixed effects of local population density, and random effects of year (Figure 3B). Space

sharing (HRO) and home range area (HRA) had comparatively small effects.

Observations also had a notable impact for Degree, Betweenness, and Clustering (Figure

3B). Fixed effects for year and observation numbers were generally strong and significantly

positive across metrics, except in the case of clustering, for which observation number's

effect was significantly negative (Figure 3B). There were also small positive effects of

population size on betweenness and degree centrality (Figure 3B).

Although individual-level drivers (reproduction, age, and individual identity) had a negligible

impact on all variables' R^2 (Figure 3B), many had a significant effect (i.e., their 95%

credibility intervals did not overlap with zero; Figure 3C). Individuals whose calves lived to

the winter and then either died before the 1st May ("Winter Yeld") or survived ("Milk") were

generally less central than those that did not give birth ("True Yeld") or whose calf died

before 1st October ("Summer Yeld"). Similarly, there were minor age-related decreases in

network centrality for the direct metrics (Group Size, Degree, and Strength; Figure 3C).

To investigate spatial patterns of sociality when accounting for our fixed and random effects,

we projected the annual SPDE random effect in two-dimensional space (Figure 4;

Supplementary Figures 5-12). The spatial distributions of network centrality metrics were

highly variable, but direct metrics generally peaked in the centre of the study system and

decreased outwards (Figure 4). Mean Strength was an exception, being lowest in the centre

and increasing outward (Figure 4D); Clustering was patchily distributed, such that no clear

pattern was evident (Figure 4H); and Betweenness was slightly offset, being highest in the

north-northeast of the study area rather than in the central north (Figure 4G). The range of

autocorrelation also varied among metrics; Betweenness and Clustering had notably shorter

ranges than the other metrics (Supplementary Figure 4). We also plotted the spatial fields

through time, revealing substantial variation in the spatial fields across the study period

(Supplementary Figures 5-12).

331 Discussion

332 The role of spatial behaviour in driving social network structure

The position individuals occupy within their social networks can affect many aspects of their

ecology and evolution (Krause *et al.* 2015; Firth *et al.* 2018; Sah *et al.* 2018), and our results

confirm the powerful role of fine-scale spatial context in shaping such traits (e.g. Farine &

Sheldon, 2019; Mourier et al., 2019; Webber & Vander Wal, 2018). Capitalising on our

models' ability to compare the influence of a wide range of spatial and non-spatial

components, we found that spatial behaviour and environmental context were the most

important determinants of social network centrality -- more so than a suite of individual-level

phenotypes and demographic factors. Individuals with larger ranges and inhabiting higher-

density areas were more central in the social network, revealing the important role of

individual spatial activity levels and location within the broader population structure. As

expected, models were further improved when we incorporated pairwise space sharing and

two-dimensional point locations, demonstrating that an individual's social network position is

not determined simply by the density of nearby individuals and by its own spatial activity, but

by other aspects of the fine-scale surrounding environment such as microclimate, resource

distribution, and landscape architecture (Spiegel *et al.* 2018; Webber & Vander Wal 2018;

He *et al.* 2019). Reciprocally, individuals may be altering their spatial behaviour, e.g. opting

to share more space or live closer together if they are more socially connected (Firth &

Sheldon 2016; Spiegel *et al.* 2016). As such, we propose that social network studies should

more regularly incorporate both space sharing and (temporally varying) point locations in
their statistical approaches to anticipate these effects, alongside specific spatial behaviours
thought to drive social network position. This practice will help to buffer for the fact that the
spatial environment not only correlates with social proximity, but can alter the fabric of the
network itself.

The landscape of sociality

One of the foremost advantages of our approach is the ability to flexibly investigate two-
dimensional spatial patterns of social network centrality. This allowed us to qualitatively
assess the spatial structure of the social network, while providing clues towards the causal
factors. Most notably, betweenness peaked in the north-northeast of the system, likely
because the far northeastern community is relatively isolated from the rest of the population
due to the landscape structure (Figure 2), so that many 'social paths' that traverse the
population (the criteria for betweenness centrality) go through individuals in this intermediate
(north-northeast) area. That is, individuals living in this area are more likely to be connected
to both the far eastern communities and the central and western ones.

As expected, direct centrality metrics (group size, degree, and strength) were affected by
local population density, which peaks in the central north study area due to the concentration
of high quality grazing (Clutton-Brock *et al.* 1982). Individuals' resource selection behaviours
increase local density in this area (Clutton-Brock *et al.* 1982), and will increase social
connectivity as a result (Ostfeld *et al.* 1986; Sanchez & Hudgens 2015; Webber & Vander
Wal 2018). This comprises strong evidence for density-related increases in social contact
frequency, and accentuates the vital importance of considering resource distribution, habitat
selection, and population structure when examining social network correlates (Spiegel *et al.*
2016; Webber & Vander Wal 2018; Farine & Sheldon 2019; He *et al.* 2019). However,
because density was accounted for as a fixed effect in the models, the spatial patterns of
location effects for the direct metrics did not strictly follow the spatial pattern of density.
Instead, these metrics peaked in the centre of the study population, demonstrating that
individuals living in this central region are more well-connected *when accounting for*
population density. Combining these spatial components allowed us to effectively
differentiate what we do know (that greater population density drives increased social
connectedness) from what we do not (the drivers of greater sociality for individuals in the
central area). Without using the SPDE effect (i.e., relying only on generalised pairwise space
sharing rather than accounting for specific two-dimensional spatial patterns), these insights
into these patterns may have been harder to detect. An alternative method could involve
splitting the population into subpopulations and analysing them separately or comparing
them, but this method has been shown to be less powerful in this population (Albery *et al.*

2019), and is ultimately based on arbitrary choices if a population is mixed. The causes of
the spatial distribution of clustering remain unresolved, but the pattern highlights areas
where individuals are connected together in triads or tight cliques, and appears to be
negatively correlated with betweenness (Figure 4). For traits such as this, it is unlikely that a
simpler explanatory variable could be formulated to quantify the spatial-social processes at
play.

Regardless of the causes of the spatial patterns, such fine-scale variation across the
landscape holds important ecological consequences, particularly for the more complex
network metrics. For instance, the areas of high clustering may act as 'incubator' areas
where cliques can develop new socially influenced behaviours (Centola 2018; Guilbeault *et*
al. 2018; Firth 2020) such as cooperative behaviours (Rand *et al.* 2011). The high contact
rates in the northern central areas might sustain high local burdens of directly transmitted
diseases (Cote & Poulin 1995), while individuals inhabiting the high-betweenness
intermediate areas may be important for transmitting novel diseases across the population
as a whole (VanderWaal *et al.* 2014).

Analytical benefits of INLA animal models

Analyses using multiple layers of different behaviours are well-suited to extricating space
and sociality in wild animal systems (Silk *et al.* 2018; Webber & Vander Wal 2018; Finn *et al.*
2019), and there is increasing conceptual and analytical overlap with the related field of
movement ecology (Jacoby & Freeman 2016; Mourier *et al.* 2019; Pasquaretta *et al.* 2020).
Notably, many spatial-social studies suffer from the necessity to reduce complex movement
patterns into simpler metrics, which risks losing important information in the process. As
such, recent studies have pushed for researchers to incorporate movement trajectories
themselves into complex network data structures (Mourier *et al.* 2019). Our approach allows
incorporation of multiple dyadic and non-dyadic behavioural measures, and with several
analytical timescales, offering an alternative workaround to this problem. Although other
methods can control for point locations (e.g. using autoregressive processes and
row/column effects; Stopher *et al.* 2012b), INLA models allow greater precision, fit quickly,
and allow incorporation of spatiotemporal structuring. Furthermore, plotting the SPDE effect
in two dimensions, as in Figure 4, gives an easily interpretable and intuitive portrayal of
network traits in space that can be hard to visualise using other methods. For these reasons,
we highly recommend further exploration of INLA animal models as a flexible method with
which to extricate individual, demographic, spatial, and temporal contributors to sociality
where sample sizes are sufficient (Thomson *et al.* 2018; Webber & Vander Wal 2018). In
addition to carrying out network-level manipulations (Daraganova *et al.* 2012; Davis *et al.*
2015; Firth & Sheldon 2016; Farine 2017), researchers concerned about spatial confounding

could implement relatively familiar linear models of social behaviour, but with additional

spatial components such as SPDE random effects and similarity matrix variance

components, with trustworthy and interpretable results (Albery *et al.* 2020a).

Although space accounted for an overwhelming amount of variation, many non-spatial

factors had substantial effects. The categorical random effect for interannual variation was

substantial, and there were detectable linear annual effects and population size effects, as

expected given the important roles of demography in structuring social networks (Shizuka &

Johnson 2019). Interestingly, there was a substantial positive association with study year

that was not attributable to the growth in population size over the same period. It is possible

that this represents a change in the deer's social phenotypes over time, although the

potential specific mechanisms now would benefit from further examination. Individual-level

factors had weaker contributions to model fit and smaller effect sizes: most notably, genetic

and individual random effects were negligible when spatial autocorrelation was accounted

for, confirming the importance of considering space when assessing heritability

independently of space in this population (Stopher *et al.* 2012a). Nevertheless, individual-

level effects were encouragingly still detectable and significant, particularly for simpler

"direct" metrics. It is possible that more complex social network positions are less

determined by individual social behaviours, particularly for animals with fission-fusion

societies such as the deer; this hypothesis could be tested using similar spatial-social

analyses in a number of other systems. This finding demonstrates that even when spatial

structuring plays a vital role in determining social network structure, controlling for this

structuring analytically can reveal important, conservative individual-level effects. Future

analyses within this population, and potentially other long-term studies, could take

advantage of this framework by including environmental drivers such as food availability and

climatic factors to explain patterns of social connectivity, while further unpicking the causes

of the individual-level trends that we observed.

451
Figure 1: Data processing and analysis pipeline, demonstrating how behavioural census
data were collected, used to derive social and spatial behavioural traits, and fitted in INLA
animal model GLMMs. Numbers in brackets represent sample sizes, and only include
females aged 3+ years. Blue arrows represent social behaviour; red arrows represent spatial
behaviours. See methods for the fixed and random effects. The text box displays the
definitions for the different spatial effects.
458

Figure 2: Spatial structuring of the 2016 social association network as a representative
example. A: the spatial locations (centroids) of individual deer, connected by their social
associations. Line opacity and width are weighted by connection strength. Ten axis units =
1KM. B: the same social network with the nodes positioned in a network spring-layout
(Csardi & Nepusz 2006) and then expanded into an even, circular grid according to their
nearest spatial positions in A. The points' (i.e. nodes') sizes and colours show individuals'
strength centrality (large and red=high strength; small and blue=low strength). Thickness of
the lines (i.e. edges) connecting them shows dyadic association strength between
individuals.
469

470
Figure 3: Model outputs demonstrating strong effects of spatial and non-spatial drivers on
social network positions. A: DIC changes associated with addition of different spatial
components, for all eight social network centrality measures. Variables are arranged in order
of mean contribution to model fit, which varied little among response variables. Different
colours correspond to different network centrality response variables, with the same colour
key as panel C. GRM = Genomic Relatedness Matrix. HRA = Home Range Area. HRO =
home range overlap. The SPDE models are differentiated into those using annual centroids
("SPDE") and the version with spatiotemporally varying annual spatial fields ("tSPDE"). B:
Variance accounted for by each variable for all eight network position measures, expressed
as contribution to R^2 in the annual model (squared correlation between observed and
predicted values). Different shades correspond to different variables. fYear = year as a
categorical random effect. HRA = Home Range Area. GRM = Genomic Relatedness Matrix.
HRO = home range overlap. Name = individual identity. NObs = number of observations
(i.e., sampling bias). PopN = population size. Status = reproductive status. SPDE = point
location effects estimated using the Stochastic Partial Differentiation Equation effect in the

INLA models. For all response variables, individual level effects (Age, Reproductive Status,
Name) had a negligible effect. C: Fixed effect estimates for the models. Fixed effects are
grouped into individual factors (age and three reproductive status effects), annual factors
(continuous time in years since study began, and annual population size), and sampling
factors (observation number). Reproductive status effects are separated into four levels: did
not reproduce (the intercept); calf died in the first few months of life ("Summer Yeld"); calf
died during the winter ("Winter Yeld"); and calf survived to May the following year ("Milk").
Different colours correspond to different network centrality response variables. Points
represent the posterior mean; error bars denote the 95% credibility intervals for the effects.
Asterisks denote significant variables (i.e., those whose estimates did not overlap with zero).
Significant variables are fully opaque, while non-significant ones are transparent.
497

499 Figure 4: Spatial fields for the SPDE random effect for each response variable, taken from
500 the INLA animal models and based on annual centroid point locations. Metrics can be
501 conceptualised as simpler "direct" metrics (top row) and more complex "indirect" metrics
502 (bottom row). Darker colours correspond to greater values. Each axis tick corresponds to
503 1km; for the values associated with the Easting and Northings, see Figure 1.

Acknowledgements

We thank Scottish Natural Heritage and its predecessors for permission to work on the Isle
of Rum NNR. The field project has been supported by grants mainly from the UK NERC with
some additional funding from BBSRC, the Royal Society and ERC. We thank all who have
contributed to the maintenance of the project over time, especially Loeske Kruuk. We thank
multiple dedicated field workers who have contributed to field data collection, especially
Fiona Guinness who collected the first 20 years of census data. GFA was funded by NSF
grant number 1414296, and by a Bruce McEwen Career Development Fellowship the Animal
Models for the Social Dimensions of Health and Aging Research Network (NIH/NIH R24
AG065172). JAF was supported by a fellowship from Merton College and BBSRC
(BB/S009752/1) and funding from NERC (NE/S010335/1). We thank Amy Sweeny and

Quinn Webber for comments on the manuscript, as well as Matt Silk, Orr Spiegel, and one
anonymous reviewer.

References

Albery, G.F., Becker, D.J., Kenyon, F., Nussey, D.H. & Pemberton, J.M. (2019). The fine-
scale landscape of immunity and parasitism in a wild ungulate population. *Integr.*
Comp. Biol., icz016, 1–11.

Albery, G.F., Kirkpatrick, L., Firth, J.A. & Bansal, S. (2020a). Unifying spatial and social
network analysis in disease ecology. *J. Anim. Ecol.*, 1–17.

Albery, G.F., Newman, C., Ross, J.G.B., Macdonald, D.W., Bansal, S. & Buesching, C.D.
(2020b). Negative density-dependent parasitism in a group-living carnivore.

Albery, G.F., Watt, K.A., Keith, R., Morris, S., Morris, A., Kenyon, F., *et al.* (2020c).
Reproduction has different costs for immunity and parasitism in a wild mammal. *Funct.*
Ecol., 34, 229–239.

Brent, L.J.N. (2015). Friends of friends: Are indirect connections in social networks important
to animal behaviour? *Anim. Behav.*, 103, 211–222.

Cairns, S.J. & Schwager, S.J. (1987). A comparison of association indices. *Anim. Behav.*,
35, 1454–1469.

Calenge, C. (2011). *Home range estimation in R: the adehabitatHR package*. Available at:
https://cran.r-project.org/web/packages/adehabitatHR/index.html. Last accessed 10
March 2020.

Castles, M., Heinsohn, R., Marshall, H.H., Lee, A.E.G., Cowlishaw, G. & Carter, A.J. (2014).
Social networks created with different techniques are not comparable. *Anim. Behav.*,
96, 59–67.

Centola, D. (2018). *How Behavior Spreads: The Science of Complex Contagions.* Princeton
Analytical Sociology Series.

Clutton-Brock, T.H., Albon, S.D. & Guinness, F.E. (1989). Fitness costs of gestation and
lactation in wild mammals. *Nature*, 337, 260–262.

Clutton-Brock, T.H., Guinness, F.E. & Albon, S.D. (1982). *Red Deer: Behavior and Ecology*
of Two Sexes. University of Chicago Press, Chicago, IL.

Cote, I.M. & Poulin, R. (1995). Parasitism and group size in social animals: A meta-analysis.
Behav. Ecol., 6, 159–165.

Croft, D.P., James, R. & Krause, J. (2008). *Exploring animal social networks*. Princeton
University Press.

Csardi, G. & Nepusz, T. (2006). The igraph software package for complex network research.
InterJournal Complex Syst., Complex Sy, 1695.

Daraganova, G., Pattison, P., Koskinen, J., Mitchell, B., Bill, A., Watts, M., *et al.* (2012).
Networks and geography: Modelling community network structures as the outcome of
both spatial and network processes. *Soc. Networks*, 34, 6–17.

Davis, S., Abbasi, B., Shah, S., Telfer, S. & Begon, M. (2015). Spatial analyses of wildlife
contact networks. *J. R. Soc. Interface*, 12.

Farine, D.R. (2017). A guide to null models for animal social network analysis. *Methods Ecol.*
Evol., 8, 1309–1320.

Farine, D.R. & Sheldon, B.C. (2019). Stable multi-level social structure is maintained by
habitat geometry in a wild bird population. *bioRxiv*, 1–30.

Finn, K.R., Silk, M.J., Porter, M.A. & Pinter-Wollman, N. (2019). The use of multilayer
network analysis across social scales in animal behaviour. *Anim. Behav.*, 149, 7–22.

Firth, J.A. (2020). Considering complexity: animal social networks and behavioural
contagions. *Trends Ecol. Evol.*, 35, 100–104.

Firth, J.A., Cole, E.F., Ioannou, C.C., Quinn, J.L., Aplin, L.M., Culina, A., *et al.* (2018).
Personality shapes pair bonding in a wild bird social system. *Nat. Ecol. Evol.*, 2, 1696–
1699.

Firth, J.A. & Sheldon, B.C. (2016). Social carry-over effects underpin trans-seasonally linked
structure in a wild bird population. *Ecol. Lett.*, 19, 1324–1332.

Firth, J.A., Voelkl, B., Crates, R.A., Aplin, L.M., Biro, D., Croft, D.P., *et al.* (2017). Wild birds
respond to flockmate loss by increasing their social network associations to others.
Proc. R. Soc. B Biol. Sci., 284.

Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with
the gambit of the group. *Behav. Ecol. Sociobiol.*, 64, 493–503.

Froy, H., Börger, L., Regan, C.E., Morris, A., Morris, S., Pilkington, J.G., *et al.* (2018).
Declining home range area predicts reduced late-life survival in two wild ungulate
populations. *Ecol. Lett.*, 21, 1001–1009.

Froy, H., Walling, C.A., Pemberton, J.M., Clutton-brock, T.H. & Kruuk, L.E.B. (2016).

Relative costs of offspring sex and offspring survival in a polygynous mammal. *Biol.*
Lett., 12, 20160417.

Guilbeault, D., Becker, J. & Centola, D. (2018). Complex Contagions: A Decade in Review.
pp. 3–25.

He, P., Maldonado-chaparro, A.A. & Farine, D.R. (2019). The role of habitat configuration in
shaping social structure: a gap in studies of animal social complexity. *Behav. Ecol.*
Sociobiol., 73.

Holand, A.M., Steinsland, I., Martino, S. & Jensen, H. (2013). Animal Models and Integrated
Nested Laplace Approximations. *Genes|Genomes|Genetics*, 3, 1241–1251.

Huisman, J., Kruuk, L.E.B., Ellis, P.A., Clutton-Brock, T.H. & Pemberton, J.M. (2016).
Inbreeding depression across the lifespan in a wild mammal population. *Proc. Natl.*
Acad. Sci., 113, 201518046.

Jacoby, D.M.P. & Freeman, R. (2016). Emerging Network-Based Tools in Movement
Ecology. *Trends Ecol. Evol.*, 31, 301–314.

Krause, J., James, R., Franks, D.W. & Croft, D.P. (2015). *Animal social networks*. Oxford
University Press, Oxford, UK.

Kruuk, L.E.B. (2004). Estimating genetic parameters in natural populations using the "animal
model." *Philos. Trans. R. Soc. B Biol. Sci.*, 359, 873–890.

Lindgren, F. & Rue, H. (2015). Bayesian Spatial Modelling with R-INLA. *J. Stat. Softw.*, 63,
1–25.

Lindgren, F., Rue, H. & Lindstrom, J. (2011). An explicit link between Gaussian fields and
Gaussian Markov random fields: the stochastic partial differential equation approach. *J.*
R. Stat. Soc. B, 73, 423–498.

Mourier, J., Lédée, E.J.I. & Jacoby, D.M.P. (2019). A multilayer perspective for inferring
spatial and social functioning in animal movement networks. *bioRxiv*.

Ostfeld, R.S., Lidicker, W.Z., Heske, E.J., Ostfeld, R.S., Lidicker, W.Z. & Heske, E.J. (1986).
The Relationship between Habitat Heterogeneity, Space Use, and Demography in a
Population of California Voles. *J. Mammal.*, 45, 433–442.

Pasquaretta, C., Dubois, T., Gomez-Moracho, T., Perilhon Delepoulle, V., Le Loc'h, G.,
Heeb, P., *et al.* (2020). Analysis of temporal patterns in animal movement networks.
Methods Ecol. Evol., 0–1.

Rand, D.G., Arbesman, S. & Christakis, N.A. (2011). Dynamic social networks promote
cooperation in experiments with humans. *Proc. Natl. Acad. Sci. U. S. A.*, 108, 19193–
19198.

Regan, C.E., Pilkington, J.G., Berenos, C., Pemberton, J.M., Smiseth, P.T. & Wilson, A.J.
(2016). Accounting for female space sharing in St. Kilda Soay sheep (Ovis aries)
results in little change in heritability estimates. *J. Evol. Biol.*, 30, 96–111.

Robert, K., Garant, D. & Pelletier, F. (2012). Keep in touch: Does spatial overlap correlate
with contact rate frequency? *J. Wildl. Manage.*, 76, 1670–1675.

Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network
structure: A synthesis across social systems. *J. Anim. Ecol.*, 87, 546–558.

Sanchez, J.N. & Hudgens, B.R. (2015). Interactions between density, home range
behaviors, and contact rates in the Channel Island fox (*Urocyon littoralis*). *Ecol. Evol.*,
5, 2466–2477.

Shizuka, D. & Johnson, A.E. (2019). How demographic processes shape animal social
networks. *Behav. Ecol.*, 1–11.

Silk, M.J., Finn, K.R., Porter, M.A. & Pinter-Wollman, N. (2018). Can Multilayer Networks
Advance Animal Behavior Research? *Trends Ecol. Evol.*, 33, 376–378.

Silk, M.J., Hodgson, D., Rozins, C., Croft, D., Delahay, R., Boots, M., *et al.* (2019).
Integrating social behaviour, demography and disease dynamics in network models:
applications to disease management in declining wildlife populations. *Philos. Trans. R.*
Soc. B, 374, 20180211.

Sosa, S., Sueur, C. & Puga-Gonzalez, I. (2020). Network measures in animal social network
analysis: their strengths, limits, interpretations and uses. *Methods Ecol. Evol.*

Spiegel, O., Leu, S.T., Sih, A. & Bull, C.M. (2016). Socially interacting or indifferent
neighbours□? Randomization of movement paths to tease apart social preference and
spatial constraints. *Methods Ecol. Evol.*, 971–979.

Spiegel, O., Sih, A., Leu, S.T. & Bull, C.M. (2018). Where should we meet? Mapping social
network interactions of sleepy lizards shows sex-dependent social network structure.
Anim. Behav., 136, 207–215.

Stopher, K. V., Nussey, D.H., Guinness, F., Morris, A., Pemberton, J.M., Clutton-Brock, T.H.,
et al. (2012a). Re-mating across years and intralineage polygyny are associated with
greater than expected levels of inbreeding in wild red deer. *J. Evol. Biol.*, 25, 2457–

2469.

Stopher, K. V, Walling, C. a, Morris, A., Guinness, F.E., Clutton-brock, T.H., Pemberton,
J.M., *et al.* (2012b). Shared spatial effects on quantitative genetic parameters:
accounting for spatial autocorrelation and home range overlap reduces estimates of
heritability in wild red deer. *Evolution*, 66, 2411–26.

Thomson, C.E., Winney, I.S., Salles, O.C. & Pujol, B. (2018). A guide to using a Multiple-
Matrix animal model to disentangle genetic and nongenetic causes of phenotypic
variance. *PLoS One*, 13, e0197720.

VanderWaal, K.L., Atwill, E.R., Isbell, L.A. & McCowan, B. (2014). Linking social and
pathogen transmission networks using microbial genetics in giraffe (Giraffa
camelopardalis). *J. Anim. Ecol.*, 83, 406–414.

Vander Wal, E., Laforge, M.P. & McLoughlin, P.D. (2014). Density dependence in social
behaviour: Home range overlap and density interacts to affect conspecific encounter
rates in a gregarious ungulate. *Behav. Ecol. Sociobiol.*, 68, 383–390.

Webber, Q.M.R. & Vander Wal, E. (2018). An evolutionary framework outlining the
integration of individual social and spatial ecology. *J. Anim. Ecol.*, 87, 113–127.

Whitehead, H. (2008). *Analyzing animal societies : quantitative methods for vertebrate*
social analysis. University of Chicago Press.

Yang, J., Lee, S.H., Goddard, M.E. & Visscher, P.M. (2011). GCTA: A Tool for Genome-wide
Complex Trait Analysis. *Am. J. Hum. Genet.*, 88, 76.

Zuur, A.F., Ieno, E.N. & Saveliev, A.A. (2017). *Beginner's guide to spatial, temporal, and*
spatial-temporal ecological data analysis with R-INLA. Highstat Ltd.

673   Supplementary Figure 1: Correlation among pairwise values in lifetime point location distances, home
674   range overlap matrix, and genomic relatedness matrix. The data have been fitted with a Generalised
675   Additive Model (GAM) smooth.

Supplementary Figure 2: Pairwise correlations among network position response variables. Values
were transformed, scaled to have a mean of 0 and a standard deviation of 1, with outliers removed,
before analysis. The data have been fitted with a Generalised Additive Model (GAM) smooth.

Supplementary Figure 3: Variance accounted for by each variable for all eight network position
measures, expressed as contribution to R^2 in the models with home range overlap. This output differs
from Figure 3B in the main text because the model did not include the SPDE random effect. Different
shades correspond to different variables. fYear = year as a categorical random effect. GRM =
Genomic Relatedness Matrix. HRO = home range overlap. Name = individual identity. NObs =
number of observations (i.e., sampling bias). PopN = population size. Status = reproductive status.
For all response variables, individual level variables (Age, Reproductive Status, Identity) had a
negligible effect.

Supplementary Figure 4: The INLA SPDE autocorrelation ranges for each response variable. Different
shades and line types correspond to different response variables. The X axis is in kilometres.

Supplementary Figure 5: Annual spatial fields for the SPDE random effect for group size, taken from
the INLA animal models and based on annual centroid point locations. Darker colours correspond to
greater values. Each axis tick corresponds to 1km distance; for the values associated with the Easting
and Northings, see Figure 1 in the main text.

Supplementary Figure 6: Annual spatial fields for the SPDE random effect for degree centrality, taken from the INLA animal models and based on annual centroid point locations. Darker colours correspond to greater values. Each axis tick corresponds to 1km distance; for the values associated with the Easting and Northings, see Figure 1 in the main text.

Supplementary Figure 7: Annual spatial fields for the SPDE random effect for strength centrality, taken from the INLA animal models and based on annual centroid point locations. Darker colours correspond to greater values. Each axis tick corresponds to 1km distance; for the values associated with the Easting and Northings, see Figure 1 in the main text.

Supplementary Figure 8: Annual spatial fields for the SPDE random effect for mean strength
centrality, taken from the INLA animal models and based on annual centroid point locations. Darker
colours correspond to greater values. Each axis tick corresponds to 1km distance; for the values
associated with the Easting and Northings, see Figure 1 in the main text.

Supplementary Figure 9: Annual spatial fields for the SPDE random effect for Eigenvector centrality,
taken from the INLA animal models and based on annual centroid point locations. Darker colours
correspond to greater values. Each axis tick corresponds to 1km distance; for the values associated
with the Easting and Northings, see Figure 1 in the main text.

Supplementary Figure 10: Annual spatial fields for the SPDE random effect for weighted Eigenvector centrality, taken from the INLA animal models and based on annual centroid point locations. Darker colours correspond to greater values. Each axis tick corresponds to 1km distance; for the values associated with the Easting and Northings, see Figure 1 in the main text.

Supplementary Figure 11: Annual spatial fields for the SPDE random effect for betweenness
centrality, taken from the INLA animal models and based on annual centroid point locations. Darker
colours correspond to greater values. Each axis tick corresponds to 1km distance; for the values
associated with the Easting and Northings, see Figure 1 in the main text.

Supplementary Figure 12: Annual spatial fields for the SPDE random effect for clustering, taken from
the INLA animal models and based on annual centroid point locations. Darker colours correspond to
greater values. Each axis tick corresponds to 1km distance; for the values associated with the Easting
and Northings, see Figure 1 in the main text.

738