Spatial point locations explain a range of social network positions in a wild ungulate

Gregory F Albery[1,2], Alison Morris[2], Sean Morris[2], Josephine M Pemberton[2], Tim H. Clutton-Brock[2,3], Daniel H Nussey[2], Josh A Firth[4,5]

1. Department of Biology, Georgetown University, Washington, DC
2. Institute of Evolutionary Biology, University of Edinburgh, Edinburgh, UK
3. Department of Zoology, University of Cambridge, Cambridge, UK
4. Department of Zoology, University of Oxford, Oxford, UK
5. Merton College, University of Oxford, Oxford, UK

Abstract

The structure of wild animal social systems governs many eco-evolutionary processes, and is determined by a complex combination of intrinsic and extrinsic drivers. Spatial structuring is a key determinant of sociality, but quantifying spatial components alongside multiple other drivers remains difficult due to data scarcity and analytical complexity. We used a 43-year dataset detailing a wild red deer population to investigate how individuals' spatial behaviours drive social network positioning, while simultaneously assessing other potential contributing factors. Using Integrated Nested Laplace Approximation (INLA) multi-matrix animal models, we demonstrate important roles of space sharing, two-dimensional point locations, and especially annually varying spatiotemporal locations, alongside reduced but detectable impacts of demography, genetics, and individual-level traits. Interestingly, spatial patterns and other drivers differed considerably among different social network metrics. These results indicate strong, multifaceted spatiotemporal structuring, emphasising the importance of considering multiple components of spatial behaviour when investigating the causes and consequences of sociality.

Introduction

Social behaviour is an integral component of an animal's phenotype, driving processes including disease transmission, mating, learning, and selection (Croft *et al.* 2008; Krause *et al.* 2015; Firth *et al.* 2018; Sah *et al.* 2018; Firth 2020). Due to its generality and flexibility, social network analysis has recently become an important method for studying wild animal social systems (Krause *et al.* 2015; Webber & Vander Wal 2019). Contemporary studies of animal behaviour often use social networks to derive individual-level traits (i.e. social network positions), under the notion that between-individual variation in network positioning is indicative of between-individual variation in social behaviour (Krause *et al.* 2015). However, social networks are also shaped by many extrinsic factors: demography determines population structure and the available individuals to interact with (Shizuka & Johnson 2019)**,** while the environment governs resource distributions, movement corridors, and emergent patterns of space use, all of which will influence the architecture of the social system (Firth & Sheldon 2016; Webber & Vander Wal 2018; Farine & Sheldon 2019; He *et al.* 2019). As such, when assessing individual-level social network positions and associated eco-evolutionary consequences, it is important to consider the population's environmental context and individual space use as factors that may affect association patterns. Such analyses can provide insights into the extent to which intrinsic *versus* extrinsic processes contribute to social network positioning (Lusseau *et al.* 2006), and can reveal how social preferences and space use interact (Firth & Sheldon 2016; Spiegel *et al.* 2016; Webber & Vander Wal 2018), indicating underlying forces governing behavioural and social processes.

Several frameworks have been proposed to facilitate the untangling of spatial and social processes in wild animals (Jacoby & Freeman 2016; Silk *et al.* 2018, 2019; Webber & Vander Wal 2018; Mourier *et al.* 2019). To date, associated statistical methodology focusses on incorporating spatial activity into the node-and-edge structure of network data, including e.g. null network permutations (Firth & Sheldon 2016), spatially embedded networks (Daraganova *et al.* 2012), and nested super-networks composed of movement trajectories (Mourier *et al.* 2019). Many such analyses involve reducing movement patterns into some form of spatial network based on home range overlap or spatial proximity between dyads (Mourier *et al.* 2019). For example, statistical models named "animal models" can examine spatial variation by fitting such matrices as variance components, potentially alongside other dyadic similarity matrices, to quantify genetic and non-genetic contributions to individuals' phenotypes (Kruuk 2004; Stopher *et al.* 2012b; Regan *et al.* 2016; Thomson *et al.* 2018; Webber & Vander Wal 2018). However, movement paths are highly multivariate, and dyadic home range overlaps represent just one aspect of spatial behaviour (Mourier *et al.* 2019;

Pasquaretta *et al.* 2020). As such, controlling for structuring using space sharing alone
potentially risks missing important variation.

On the one hand, there is now much evidence supporting the intuitive concept that spatial
proximity and social activity are aligned: that is, individuals that share more space are more
likely to associate or interact, for elk (Vander Wal *et al.* 2014), raccoons (Robert *et al.* 2012),
foxes (Sanchez & Hudgens 2015), great tits (Firth & Sheldon 2016), and myriad other
systems. However, while important, this concept is subtly distinct from fine-scale spatial
assessment of heterogeneity in social behaviour – i.e., whether different areas of the
landscape encourage different social network structures. For instance, if environmental
gradients across a study system alter individuals' behaviour, then this will alter local social
network structure, so that individuals will exhibit different social network traits depending on
their locations in space – independent of their pairwise space sharing. Most notably,
resource distributions are often extremely heterogeneous, altering habitat selection,
aggregation in space, and therefore social network structure (Webber & Vander Wal 2018).
Consequently, individuals living in high-resource areas may (for example) have many social
partners. This phenomenon can have far-reaching consequences, for example by enhancing
disease exposure in resource-supplemented populations (Becker *et al.* 2015). Notably, due
partly to the analytical challenges, few studies of spatial-social structuring have examined
the effect of point locations (i.e., where an animal is on a landscape) compared to space
sharing (i.e., how much a pair of animals overlap in an unspecified space), and/or compared
their influence to individual-level phenotypic traits.

Studies across ecological disciplines are increasingly using Integrated Nested Laplace
Approximation (INLA) models to control for spatial autocorrelation in a multitude of contexts
(Lindgren *et al.* 2011; Lindgren & Rue 2015; Zuur *et al.* 2017). This approach uses
individuals' point locations to model 2-dimensional spatial patterns in the response variable,
thereby controlling for and estimating spatiotemporal variation associated with fine-scale
positioning within the landscape (Albery *et al.* 2019). Neither animal models nor INLA have
yet been used to examine how spatial processes shape social network positions, despite
their enormous potential for doing so. Further, although animal models can be fitted using
INLA (Holand *et al.* 2013), there has been no exploration of how dyadic space sharing and
individual point locations can be fitted simultaneously within animal models, how these
factors compete with each other, and what insights this framework can provide.

As with spatial behaviour, sociality can be summarised using a range of different metrics
(Krause *et al.* 2015; Sosa *et al.* 2020). Social network measures can represent individuals'
own social connections, such as 'degree' (their number of social ties), or 'average strength'

(the mean strength of their associations); alternatively, more complex network metrics may
represent individuals' positions within the wider network, including 'betweenness' (the extent
to which they bridge different social groups), 'Eigenvector centrality' (the centrality of their
associates), and 'clustering' (the propensity for their associates to be connected to one
another). These diverse metrics can provide insights into different aspects of social
behaviour, while arising from different spatial movement syndromes: for example, individuals
that consistently inhabit densely populated locations may commonly reoccur with the same
individuals and thus experience high average strength, while individuals that move around
various locations or inhabit restricted movement corridors may connect otherwise
disconnected social groups together, thereby exhibiting high betweenness centrality (Firth *et*
al. 2017). Further, different metrics can also provide insights into various consequences of
sociality: for instance, while spreading a highly contagious disease may depend heavily on
an animal's degree (unique number of partners), their influence on the spread of a learned
behaviour may depend more on local clustering (Firth 2020). Because few studies use
multiple spatial behaviour metrics (as outlined above) and few include other potential drivers
in the same models, it is unclear how spatial factors compare to other individual-level and
demographic factors in shaping sociality, and whether different spatial behaviours are
important for determining different social network metrics.

The Isle of Rum red deer (*Cervus elaphus*) study population is an unmanaged wild
population with a fission-fusion social system (Clutton-Brock *et al.* 1982). They exhibit spatial
autocorrelation in a number of important phenotypes: individuals with greater home range
overlap have more similar behavioural and life history traits (Stopher *et al.* 2012b), and those
in closer proximity have more similar parasite burdens (Albery *et al.* 2019); further, as with
other matrilineal mammalian systems, closely related individuals frequently associate
(Clutton-Brock *et al.* 1982) and live closer together (Stopher *et al.* 2012b). Individuals have
highly repeatable home ranges (Stopher *et al.* 2012b) which decline in size over their
lifetimes, predicting declining survival probability (Froy *et al.* 2018). As such, the deer
comprise an ideal system for assessing spatial-social relationships in the wild.

To assess how individuals' spatial behaviours translate to social network positions, we
constructed fine-scale social networks from 43 years of censuses describing social
groupings across the study population. We derived 8 different individual-level network
positioning measures of varying complexity which are often proposed to be important to
different social processes (Krause *et al.* 2015; Sosa *et al.* 2020). Using multi-matrix animal
models in INLA (Thomson *et al.* 2018), we examined whether spatial locations and home
range overlap explained variation in network position metrics, alongside a range of
individual-, temporal-, and population-level factors. Additionally, we investigated how various

spatiotemporal autocorrelation structures compared in determining social behaviour. We expected that a) locations on the landscape would determine a substantial proportion of social network positioning, comparable or greater than space sharing and other individual-level drivers; b) that this influence would be temporally varying; and c) that different social network metrics would exhibit different spatial patterns and vary in their relationship to the other drivers. We furthermore predicted that the spatial gradient in population density would have a strong effect on social network structure. This not only comprises a large-scale empirical examination of the factors shaping social network positions in this extensively monitored wild mammal, but also provides a methodological advancement in developing powerful, flexible new methods (INLA-based multi-matrix animal models) with broad potential for examining spatial-social processes in this and other systems.

Methods

Study system and censusing

This study was carried out on a long-term study population of red deer on the Isle of Rum, Scotland (57°N,6°20'W). We focussed on females aged 3+ years, as these individuals have the most complete associated census data. Individuals are monitored from birth, providing substantial life history and behavioural data, and >90% of calves are caught and tagged, with tissue samples taken (Clutton-Brock *et al.* 1982). The population thus has comprehensive genomic data, allowing high-powered quantitative genetic analyses: most individuals born since 1982 have been genotyped at >37,000 SNPs, distributed throughout the genome (e.g. Huisman, Kruuk, Ellis, Clutton-Brock, & Pemberton, 2016). Census data were collected for the years 1974-2017, totalling 423,070 census observations. Deer were censused by field workers five times a month, for eight months of the year, along one of two alternating routes (Clutton-Brock *et al.* 1982). Individuals' identities, locations (to the nearest 100M), and group membership were recorded. Grouping events were estimated by seasoned field workers according to a variant of the "chain rule" (e.g. Castles et al., 2014), where individuals grazing in a contiguous group within close proximity of each other (under ~10 metres) were deemed to be associating. Our dataset totalled 3356 annual observations among 532 grown females, with mean 112 groups observed per individual (Figure 1).

In this system, female reproduction imposes substantial costs for immunity and parasitism (Albery *et al.* 2020), and for subsequent fitness (Clutton-Brock, Albon, & Guinness, 1989; Froy, Walling, Pemberton, Clutton-Brock, & Kruuk, 2016). If a female reproduces, she produces 1 calf per year in the spring, generally beginning in May; the "deer year" begins on May 1 for this reason. Here, reproductive status was classified into the following four categories using behavioural observations: True Yeld (did not give birth); Summer Yeld (the

169 female's calf died in the summer, before 1st October); Winter Yeld (the female's calf died in

170 the winter, after 1st October); and Milk (calf survived to 1st May the following calendar year).

Generating spatial and social matrices

All code is available online at https://github.com/gfalbery/INLA_N_Out. We constructed the

HRO matrix using the R package AdeHabitatHR, following previous methodology (Stopher *et*

al. 2012b; Regan *et al.* 2016; Froy *et al.* 2018). First, using a kernel density estimation

method, we derived lifetime home ranges for each individual with more than five census

observations. We used lifetime home ranges to fit one value per individual in the animal

models; individual ranges (and range sizes) correlate strongly from year to year (Stopher *et*

al. 2012b; Froy *et al.* 2018). We derived proportional home range overlap (HRO) of each

dyad using Bhattacharya Affinity (following Stopher *et al.* 2012b), producing values between

0-1 (i.e. no overlap to complete overlap).

To control for individuals' two-dimensional point locations we used a Stochastic Partial

Differentiation Equation (SPDE) effect, in which distance between points is used to calculate

spatial autocorrelation using Matern covariance (Lindgren *et al.* 2011). This random effect

used individuals' annual centroids (mean easting and northing in a given year) or lifetime

centroids (mean easting and northing across all observations) as point locations to

approximate spatial variation in the response variable (Lindgren *et al.* 2011; Albery *et al.*

2019). We used a genomic relatedness matrix (R_{GRM}) using homozygosity at 37,000 Single

Nucleotide Polymorphisms, scaled at the population level (Yang *et al.* 2011; for a population-

specific summary, see Huisman *et al.* 2016). This matrix is well-correlated with pedigree-

derived relatedness metrics (Huisman *et al.* 2016). Home range overlap was well-correlated

with distance between lifetime centroids (i.e., closer individuals tended to share more range),

and both were weakly but significantly correlated with genetic relatedness (Supplementary

Figure 1).

We constructed annual social networks using "gambit of the group," where individuals in the

same grouping event (as described above) were taken to be associating (Franks *et al.*

2010). Dyadic associations were calculated using the 'simple ratio index' (Cairns &

Schwager 1987) derived as a proportion of total sightings (grouping events) in which the

focal individuals were seen together: Sightings$_{A,B}$/(Sightings$_A$+Sightings$_B$-Sightings$_{A,B}$), or

Intersect$_{A,B}$/Union$_{A,B}$. In this dyadic matrix, 0=never seen together and 1=never seen apart.

We constructed a series of 43 annual networks constructed only from census records in

each May-December period, from which we derived annual social network position

measures as response variables (Figure 1-2). We elected to investigate this seasonal period

because it stretches from the spring calving period until the beginning of the mortality period,
simplifying network construction and avoiding complications arising from mortality events.

207
Figure 1: Data processing and analysis pipeline, demonstrating how behavioural census
data were collected, used to derive social and spatial behavioural traits, and fitted in INLA
animal model GLMMs. Numbers in brackets represent sample sizes, and only include
females aged 3+ years. Blue arrows represent social behaviour; red arrows represent spatial
behaviours. See methods for the fixed and random effects. The text box displays the
definitions for the different spatial effects.
214

Figure 2: Spatial structuring of the 2016 social association network as a representative
example. A: the spatial locations (centroids) of individual deer, connected by their social
associations. Line opacity and width are weighted by connection strength. Ten axis units =
1KM. B: the same social network expanded into an even, circular grid according to their
nearest spatial positions in A). The points' (i.e. nodes') sizes and colours show individuals'
strength centrality (large and red=high strength; small and blue=low strength). Thickness of
the lines (i.e. edges) connecting them shows dyadic association strength between
individuals.

Statistical Analysis

We derived eight individual-level network metrics from the annual social networks for use as
response variables in INLA Generalised Linear Mixed Models (GLMMs) with a Gaussian
family specification. In increasing order of complexity, our measures included four direct
metrics: 1) Group Size – the average number of individuals a deer associated with per
sighting; 2) Degree – the number of unique individuals she was observed with; 3) Strength –
sum of all their weighted social associations to others; 4) Mean Strength – the average
association strength to each of the unique individuals she was observed with (equivalent to
strength divided by degree). We also included four more complex "indirect" metrics: 5)
Eigenvector centrality – akin to the sum of her unique associates' degrees; 6) Weighted
Eigenvector – akin to the sum of her associates' strengths weighted by their association to
her; 7) Betweenness – the number of shortest paths that pass through the focal individual to
traverse the whole network; 8) Clustering (local) – the tendency for an individual's contacts
to be connected to one another, forming triads. The raw, untransformed correlations were
assessed for all metrics (Supplementary Figure 2); when modelling them as response

variables, to approximate normality, all social metrics were square root-transformed apart
from eigenvector centralities (which were left untransformed), group size (which was cube
root-transformed), and betweenness (which was log-transformed). Each social network
metric was fitted as a response variable in a separate model set (as outlined conceptually in
Figure 1). We ensured that all models followed the same base structure. Random effects
included individual identity and year (categorical), as well as the genetic relatedness matrix.
Fixed effects included Age (continuous, in years), Reproductive Status (four categories: True
Yeld; Summer Yeld; Winter Yeld; and Milk), and Number of observations (continuous, log-
transformed), as well as year-level continuous factors including Year (continuous) and that
year's study Population Size (log-transformed). All continuous response and explanatory
variables were standardised to have a mean of zero and a standard deviation of 1.

To investigate the divergent effects of different spatial behaviours, we iteratively added
different combinations of spatial random effects to the base model, in increasingly complex
formulations. First, we added the HRO (space sharing) matrix. Next, we added an INLA
Stochastic Partial Differentiation Equation (SPDE or point location) spatial effect based on
lifetime centroids, to investigate whether point locations and space sharing behaved similarly
when at the same timescale (i.e., across individual lifetimes). Next, to investigate whether
finer temporal scales improved our inference of spatial effects, we altered the SPDE effect to
use annual rather than lifetime centroids. Finally, we fitted the annual centroids in a
spatiotemporal model structure, allowing entirely different (uncorrelated) spatial fields for the
SPDE effect for each year. Only one of the SPDE random effects was fitted at once, and the
best-fitting model was identified using changes in Deviance Information Criterion (DIC). A
conservative change of 10ΔDIC was used to differentiate between competing models – i.e.,
any variable that decreased DIC by more than 10 was deemed significant.

To compare the variance accounted for by all fixed and random effects, we examined the
model's predicted values and their correlations with the observed values. We used the
model to predict each social behaviour metric, and iteratively held each explanatory
variable's predictions at the mean, one at a time. We then assessed the squared correlations
of these values with the observed values (i.e., R^2), relative to those of the full model.
Variables with greater effects in the model produced less accurate predicted values when
held constant. Animal models generally extract the variance components (random effects) of
genetic and non-genetic contributors to quantify heritability (Kruuk 2004). However, in INLA
models, the SPDE variance components are hard to estimate where the range parameter is
large; as such, predicting using the model is a preferred approach for our purposes (Finn
Lindgren, pers. comms.). Nevertheless, the variance components were found to largely
mirror our results when inspected.

Figure 3: Model outputs demonstrating strong effects of spatial and non-spatial drivers on social network positions. A: DIC changes associated with addition of different spatial components, for all eight social network centrality measures. Models are arranged in order of model fit, and the rank order varied little among response variables. Different colours correspond to different network centrality response variables, with the same colour key as panel C. The SPDE models are differentiated into those using individuals' lifetime centroids ("LifetimeSPDE") and annual centroids ("AnnualSPDE") and the spatiotemporally varying annual spatial fields ("SpatiotemporalSPDE"). B: Variance accounted for by each variable for all eight network position measures, expressed as contribution to R^2 in the annual model (squared correlation between observed and predicted values). Different shades correspond to different variables. fYear = year as a categorical random effect. GRM = Genomic

Relatedness Matrix. HRO = home range overlap. Name = individual identity. NObs = number
of observations (i.e., sampling bias). PopN = population size. Status = reproductive status.
SPDE = point location effects estimated using the Stochastic Partial Differentiation Equation
effect in the INLA models. For all response variables, individual level effects (Age,
Reproductive Status, Name) had a negligible effect. C: Fixed effect estimates for the models.
Fixed effects are grouped into individual factors (age and three reproductive status effects),
annual factors (continuous time in years since study began, and annual population size), and
sampling factors (observation number). Reproductive status effects are separated into four
levels: did not reproduce (the intercept); calf died in the first few months of life ("Summer
Yeld"); calf died during the winter ("Winter Yeld"); and calf survived to May the following year
("Milk"). Different colours correspond to different network centrality response variables.
Points represent the posterior mean; error bars denote the 95% credibility intervals for the
effects. Asterisks denote significant variables (i.e., those whose estimates did not overlap
with zero). Significant variables are fully opaque, while non-significant ones are transparent.
303

Results

Spatial behaviours were extremely important in determining all eight individual-level social

network position measures. The non-spatial model was by far the worst-fitting for all eight

response variables, and the DIC changes associated with adding spatial components were

substantial (Figure 3A). Notably, point location-based SPDE effects tended to improve model

fit more than space sharing HRO effects, even when conceptualised at the same timescale

(i.e., across the individual's lifetime; Figure 1A). Investigating the R^2 components of the

models containing only HRO (i.e., without SPDE effects) revealed that in general spatial

overlap accounted for more variation than the genetic matrix (Supplementary Figure 3), but

comparing these with the other models revealed that the point location effects contributed

more than either of these matrices (Figure 3B). Annually varying centroids further improved

model fit, and allowing the spatial field to vary between years in our spatiotemporal models

improved models even more (Figure 3A). Although the space sharing and genomic

relatedness matrices had similar sized impacts on the full models (Figure 3B), removing the

SPDE effect resulted in a substantial increase in the HRO effect, but with very little impact

on the GRM's R^2 (Supplementary Figure 3). These findings were relatively consistent across

all metrics (Figure 3A-B), although the SPDE effect was notably smaller for betweenness

(Figure 3B). Taken together, these results reveal that lifetime space sharing was good at

accounting for variation in social behaviour, but that its effect was surpassed by increasingly

complex temporal formulations of point location effects.

We compared the importance of all fixed and random effects by predicting selectively from

the model, revealing overwhelmingly strong effects of spatiotemporal factors (Figure 3B).

Our models fit well and explained a substantial amount of variation in social network

centrality (>70%), and the majority of the model's fit was lent by the INLA SPDE effect

(Figure 3B). Observations also had a notable impact for Degree, Betweenness, and

Clustering, and the categorical random effect for year had a substantial effect across all

response variables (Figure 3B). Fixed effects for year and observation numbers were

generally strong and significantly positive across metrics, except in the case of clustering,

which was significantly negative (Figure 3B). There were also small positive effects of

population size on betweenness and degree centrality (Figure 3B).

Although individual-level drivers (reproduction, age, and individual identity) had a negligible

impact on all variables' R2 (Figure 3B), many had a significant effect (i.e., their 95%

credibility intervals did not overlap with zero; Figure 3C). Individuals whose calves lived to

the winter and then either died before the 1st May ("Winter Yield") or survived ("Milk") were

generally less central than those that did not give birth ("True Yield") or whose calf died

before 1st October ("Summer Yeld"). Similarly, there were minor age-related decreases in

network centrality for the direct metrics (Group Size, Degree, and Strength; Figure 3C).

To investigate spatial patterns of sociality, we projected the annual SPDE random effect in

two-dimensional space (Figure 4; Supplementary Figures 5-12). As expected, the spatial

distributions of network centrality metrics were highly variable, but generally peaked in the

central north area of the study system and decreased outwards (Figure 4). Mean Strength

was an exception, being lowest in the high-density areas and increasing outward (Figure

4D); Clustering was patchily distributed, such that no clear pattern was evident (Figure 4H);

and Betweenness was slightly offset, being highest in the north-northeast of the study area

rather than in the central north (Figure 4G). The range of autocorrelation also varied among

metrics; Betweenness and Clustering had notably shorter ranges than the other metrics

(Supplementary Figure 4). We also plotted the spatial fields through time, revealing

substantial variation in the spatial fields across the study period (Supplementary Figures 5-

12).

Figure 4 (Previous page): Spatial fields for the SPDE random effect for each response
variable, taken from the INLA animal models and based on annual centroid point locations.
Metrics can be conceptualised as simpler dyadic "direct" metrics (top row) and more
complex "indirect" metrics (bottom row). Darker colours correspond to greater values. Each
axis tick corresponds to 1km; for the values associated with the Easting and Northings, see
Figure 1.

Discussion

The role of spatial behaviour in driving social network positions

The position individuals occupy within their social networks can affect many aspects of their

ecology and evolution (Krause *et al.* 2015; Firth *et al.* 2018; Sah *et al.* 2018), and our results

confirm the powerful role of fine-scale spatial activity in shaping such traits (e.g. Farine &

Sheldon, 2019; Mourier et al., 2019; Webber & Vander Wal, 2018). Although pairwise home

range overlap was important in determining social centrality measures, point locations were

substantially more important, and allowed us to more easily account for spatiotemporal

variation. Inter-annual variation in spatial effects proved especially influential in our models,

across increasingly sophisticated spatiotemporal model formulations. While point locations

were superior to home range overlaps even at the same coarse timescale (i.e., across

lifetimes), our models universally revealed benefits of incorporating temporally varying

spatial behaviours. Moreover, the autocorrelation range and the importance of different

behavioural components differed notably across centrality measures, suggesting that

different spatial processes play a role in determining different network positions. As such, we

propose that social network studies should more regularly incorporate both space sharing

and (temporally varying) point locations in their statistical approaches to anticipate these

effects. This practice will help to buffer for the fact that the spatial environment not only
correlates with social proximity, but can alter the fabric of the network itself.

The landscape of sociality

Spatial patterns were quite varied among metrics but were nevertheless amenable to
interpretation. Most notably, the spatial distributions of direct metrics (group size, degree,
and strength) were very similar and likely attributable to the concentration of resources in the
form of high quality grazing, which peaks in the central north study area (Clutton-Brock *et al.*
1982). Individuals' resource selection behaviours increase local density in this area (Clutton-
Brock *et al.* 1982), and will increase social connectivity as a result (Ostfeld *et al.* 1986;
Sanchez & Hudgens 2015; Webber & Vander Wal 2018). This comprises strong evidence for
density-related increases in social contact frequency, and accentuates the vital importance
of considering resource distribution, habitat selection, and population structure when
examining social network correlates (Spiegel *et al.* 2016; Webber & Vander Wal 2018;
Farine & Sheldon 2019; He *et al.* 2019). For these simpler, direct metrics, it is possible that a
measure of local spatial population density (e.g. Coulson *et al.* 1997) could be fitted to
control for and estimate spatial-social confounding. In contrast, betweenness peaked in the
north-northeast of the system, likely because the northeastern community is slightly isolated
from the rest of the population (Figure 2), so that many 'social paths' that traverse the
population (the criteria for betweenness centrality) go through individuals in this intermediate
area. The causes of the spatial distribution of clustering remain unresolved, but the pattern
highlights areas where individuals are connected together in triads or tight cliques, and
appears to be negatively correlated with betweenness (Figure 4). For these traits, it is
unlikely that a simpler explanatory variable could be formulated to quantify the spatial-social
processes at play.

Regardless of the causes of the spatial patterns, such fine-scale variation across the
landscape holds important eco-evolutionary consequences, particularly for the more
complex network metrics. For instance, the areas of high clustering may act as 'incubator'
areas where cliques can develop new socially influenced behaviours (Firth 2020) such as
cooperative behaviours (Rand *et al.* 2011). The high contact rates in the northern central
areas might sustain high local burdens of directly transmitted diseases (Cote & Poulin 1995),
while individuals inhabiting the high-betweenness intermediate areas may be important for
transmitting novel diseases across the population as a whole (VanderWaal *et al.* 2014).
Without using the SPDE effect (i.e., relying only on generalised pairwise space sharing
rather than accounting for specific two-dimensional spatial patterns), these insights into
these patterns would not have been possible.

Analytical benefits of INLA animal models

Analyses using multiple layers of different behaviours are well-suited to extricating space and sociality in wild animal systems (Silk *et al.* 2018; Webber & Vander Wal 2018; Finn *et al.* 2019), and there is increasing conceptual and analytical overlap between movement-based and network-based approaches (Jacoby & Freeman 2016; Mourier *et al.* 2019; Pasquaretta *et al.* 2020). Notably, many spatial-social studies suffer from the necessity to reduce complex movement patterns into simpler metrics, which risks losing important information in the process. As such, recent studies have pushed for researchers to incorporate movement trajectories themselves into complex network data structures (Mourier *et al.* 2019). Our approach allows incorporation of multiple dyadic and non-dyadic behavioural measures, and with several analytical timescales, offering an alternative workaround to this problem. Although other methods can control for point locations (e.g. using autoregressive processes and row/column effects; Stopher *et al.* 2012b), INLA models allow greater precision, fit quickly, and allow easy incorporation of spatiotemporal structuring. Furthermore, plotting the SPDE effect in two dimensions, as in Figure 4, gives an easily interpretable and intuitive portrayal of network traits in space that can be hard to visualise using other methods. For these reasons, we highly recommend further exploration of INLA animal models as a flexible method with which to extricate individual, demographic, spatial, and temporal contributors to sociality where sample sizes are sufficient (Thomson *et al.* 2018; Webber & Vander Wal 2018). In addition to carrying out network-level manipulations (Daraganova *et al.* 2012; Davis *et al.* 2015; Firth & Sheldon 2016; Farine 2017), researchers concerned about spatial confounding could implement relatively familiar linear models of social behaviour, but with additional spatial components such as SPDE random effects and similarity matrix variance components, with trustworthy and interpretable results (Albery et al., in review).

Our approach also allowed us to quantify the impacts of multiple non-spatial drivers of network centrality and compare them with spatial behaviour. Although space accounted for an overwhelming amount of variation, many other factors had substantial effects. The categorical random effect for interannual variation was substantial, and there were detectable linear annual effects and population size effects, as expected given the important roles of demography in structuring social networks (Shizuka & Johnson 2019). Individual-level factors had weaker contributions to model fit and smaller effect sizes: most notably, genetic and individual random effects were negligible when spatial autocorrelation was accounted for, confirming the importance of considering space when assessing heritability independently of space in this population (Stopher *et al.* 2012a). Notably, previous analyses in this system revealed that accounting for space substantially reduced heritability estimates for spatial behaviour (home range size), but less so for life history characteristics (Stopher *et*

al. 2012a). We impress that the finding of extreme spatial dependence in social behaviour

does not necessarily imply that other traits will be subject to a similar reduction in heritability,

although incorporating point locations may be similarly revealing about the non-spatial

drivers of such traits.

Nevertheless, individual-level effects were encouragingly still detectable and significant,

particularly for simpler "direct" metrics. It is possible that more complex social network

positions are less determined by individual social behaviours, particularly for animals with

relatively strong fission-fusion dynamics (i.e., heavy mixing) such as the deer; this

hypothesis could be tested using similar spatial-social analyses in a number of other

systems. This finding demonstrates that even when spatial structuring plays a vital role in

determining social network structure, controlling for this structuring analytically can reveal

important, conservative individual-level effects. Future analyses within this population, and

potentially other long-term studies, could take advantage of this framework by including

environmental drivers such as food availability and climatic factors to explain patterns of

social connectivity, while further unpicking the causes of the individual-level trends that we

observed.

Acknowledgements

We thank Scottish Natural Heritage and its predecessors for permission to work on the Isle

of Rum NNR. The field project has been supported by grants mainly from the UK NERC with

some additional funding from BBSRC, the Royal Society and ERC. We thank all who have

contributed to the maintenance of the project over time, especially Loeske Kruuk. We thank

multiple dedicated field workers who have contributed to field data collection, especially

Fiona Guinness who collected the first 20 years of census data. JAF was supported by a

fellowship from Merton College and BBSRC (BB/S009752/1) and funding from NERC

(NE/S010335/1). We thank Amy Sweeny and Quinn Webber for comments on the

manuscript.

References

Albery, G.F., Becker, D.J., Kenyon, F., Nussey, D.H. & Pemberton, J.M. (2019). The fine-

scale landscape of immunity and parasitism in a wild ungulate population. *Integr.*

Comp. Biol., icz016, 1–11.

Albery, G.F., Watt, K.A., Keith, R., Morris, S., Morris, A., Kenyon, F., *et al.* (2020).

Reproduction has different costs for immunity and parasitism in a wild mammal. *Funct.*

Ecol., 34.

Becker, D.J., Streicker, D.G. & Altizer, S. (2015). Linking anthropogenic resources to wildlife-
pathogen dynamics: a review and meta-analysis. *Ecol. Lett.*, n/a-n/a.

Cairns, S.J. & Schwager, S.J. (1987). A comparison of association indices. *Anim. Behav.*,
35, 1454–1469.

Castles, M., Heinsohn, R., Marshall, H.H., Lee, A.E.G., Cowlishaw, G. & Carter, A.J. (2014).
Social networks created with different techniques are not comparable. *Anim. Behav.*,
96, 59–67.

Clutton-Brock, T.H., Albon, S.D. & Guinness, F.E. (1989). Fitness costs of gestation and
lactation in wild mammals. *Nature*, 337, 260–262.

Clutton-Brock, T.H., Guinness, F.E. & Albon, S.D. (1982). *Red Deer: Behavior and Ecology*
of Two Sexes. University of Chicago Press, Chicago, IL.

Cote, I.M. & Poulin, R. (1995). Parasitism and group size in social animals: A meta-analysis.
Behav. Ecol., 6, 159–165.

Coulson, T., Albon, S., Guinness, F., Pemberton, J. & Clutton-Brock, T. (1997). Population
Substructure, Local Density, and Calf Winter Survival in Red Deer (Cervus elaphus).
Ecology, 78, 852.

Croft, D.P., James, R. & Krause, J. (2008). *Exploring animal social networks*. Princeton
University Press.

Daraganova, G., Pattison, P., Koskinen, J., Mitchell, B., Bill, A., Watts, M., *et al.* (2012).
Networks and geography: Modelling community network structures as the outcome of
both spatial and network processes. *Soc. Networks*, 34, 6–17.

Davis, S., Abbasi, B., Shah, S., Telfer, S. & Begon, M. (2015). Spatial analyses of wildlife
contact networks. *J. R. Soc. Interface*, 12.

Farine, D.R. (2017). A guide to null models for animal social network analysis. *Methods Ecol.*
Evol., 8, 1309–1320.

Farine, D.R. & Sheldon, B.C. (2019). Stable multi-level social structure is maintained by
habitat geometry in a wild bird population. *bioRxiv*, 1–30.

Finn, K.R., Silk, M.J., Porter, M.A. & Pinter-Wollman, N. (2019). The use of multilayer
network analysis across social scales in animal behaviour. *Anim. Behav.*, 149, 7–22.

Firth, J.A. (2020). Considering complexity: animal social networks and behavioural
contagions. *Trends Ecol. Evol.*, 35, 100–104.

Firth, J.A., Cole, E.F., Ioannou, C.C., Quinn, J.L., Aplin, L.M., Culina, A., *et al.* (2018).
Personality shapes pair bonding in a wild bird social system. *Nat. Ecol. Evol.*, 2, 1696–
1699.

Firth, J.A. & Sheldon, B.C. (2016). Social carry-over effects underpin trans-seasonally linked
structure in a wild bird population. *Ecol. Lett.*, 19, 1324–1332.

Firth, J.A., Voelkl, B., Crates, R.A., Aplin, L.M., Biro, D., Croft, D.P., *et al.* (2017). Wild birds
respond to flockmate loss by increasing their social network associations to others.
Proc. R. Soc. B Biol. Sci., 284.

Franks, D.W., Ruxton, G.D. & James, R. (2010). Sampling animal association networks with
the gambit of the group. *Behav. Ecol. Sociobiol.*, 64, 493–503.

Froy, H., Börger, L., Regan, C.E., Morris, A., Morris, S., Pilkington, J.G., *et al.* (2018).
Declining home range area predicts reduced late-life survival in two wild ungulate
populations. *Ecol. Lett.*, 21, 1001–1009.

Froy, H., Walling, C.A., Pemberton, J.M., Clutton-brock, T.H. & Kruuk, L.E.B. (2016).
Relative costs of offspring sex and offspring survival in a polygynous mammal. *Biol.*
Lett., 12, 20160417.

He, P., Maldonado-chaparro, A.A. & Farine, D.R. (2019). The role of habitat configuration in
shaping social structure: a gap in studies of animal social complexity. *Behav. Ecol.*
Sociobiol., 73.

Holand, A.M., Steinsland, I., Martino, S. & Jensen, H. (2013). Animal Models and Integrated
Nested Laplace Approximations. *Genes|Genomes|Genetics*, 3, 1241–1251.

Huisman, J., Kruuk, L.E.B., Ellis, P.A., Clutton-Brock, T.H. & Pemberton, J.M. (2016).
Inbreeding depression across the lifespan in a wild mammal population. *Proc. Natl.*
Acad. Sci., 113, 201518046.

Jacoby, D.M.P. & Freeman, R. (2016). Emerging Network-Based Tools in Movement
Ecology. *Trends Ecol. Evol.*, 31, 301–314.

Krause, J., James, R., Franks, D.W. & Croft, D.P. (2015). *Animal social networks*. Oxford
University Press, Oxford, UK.

Kruuk, L.E.B. (2004). Estimating genetic parameters in natural populations using the "animal
model." *Philos. Trans. R. Soc. B Biol. Sci.*, 359, 873–890.

Lindgren, F. & Rue, H. (2015). Bayesian Spatial Modelling with R-INLA. *J. Stat. Softw.*, 63,
1–25.

Lindgren, F., Rue, H. & Lindstrom, J. (2011). An explicit link between Gaussian fields and
Gaussian Markov random fields: the stochastic partial differential equation approach. *J.*
R. Stat. Soc. B, 73, 423–498.

Lusseau, D., Wilson, B., Hammond, P.S., Grellier, K., Durban, J.W., Parsons, K.M., *et al.*
(2006). Quantifying the influence of sociality on population structure in bottlenose
dolphins. *J. Anim. Ecol.*, 75, 14–24.

Mourier, J., Lédée, E.J.I. & Jacoby, D.M.P. (2019). A multilayer perspective for inferring
spatial and social functioning in animal movement networks. *bioRxiv.*

Ostfeld, R.S., Lidicker, W.Z., Heske, E.J., Ostfeld, R.S., Lidicker, W.Z. & Heske, E.J. (1986).
The Relationship between Habitat Heterogeneity, Space Use, and Demography in a
Population of California Voles. *J. Mammal.*, 45, 433–442.

Pasquaretta, C., Dubois, T., Gomez-Moracho, T., Perilhon Delepoulle, V., Le Loc'h, G.,
Heeb, P., *et al.* (2020). Analysis of temporal patterns in animal movement networks.
Methods Ecol. Evol., 0–1.

Rand, D.G., Arbesman, S. & Christakis, N.A. (2011). Dynamic social networks promote
cooperation in experiments with humans. *Proc. Natl. Acad. Sci. U. S. A.*, 108, 19193–
19198.

Regan, C.E., Pilkington, J.G., Berenos, C., Pemberton, J.M., Smiseth, P.T. & Wilson, A.J.
(2016). Accounting for female space sharing in St. Kilda Soay sheep (Ovis aries)
results in little change in heritability estimates. *J. Evol. Biol.*, 30, 96–111.

Robert, K., Garant, D. & Pelletier, F. (2012). Keep in touch: Does spatial overlap correlate
with contact rate frequency? *J. Wildl. Manage.*, 76, 1670–1675.

Sah, P., Mann, J. & Bansal, S. (2018). Disease implications of animal social network
structure: A synthesis across social systems. *J. Anim. Ecol.*, 87, 546–558.

Sanchez, J.N. & Hudgens, B.R. (2015). Interactions between density, home range
behaviors, and contact rates in the Channel Island fox (*Urocyon littoralis*). *Ecol. Evol.*,
5, 2466–2477.

Shizuka, D. & Johnson, A.E. (2019). How demographic processes shape animal social
networks. *Behav. Ecol.*, 1–11.

Silk, M.J., Finn, K.R., Porter, M.A. & Pinter-Wollman, N. (2018). Can Multilayer Networks
Advance Animal Behavior Research? *Trends Ecol. Evol.*, 33, 376–378.

Silk, M.J., Hodgson, D., Rozins, C., Croft, D., Delahay, R., Boots, M., *et al.* (2019).

Integrating social behaviour, demography and disease dynamics in network models:
applications to disease management in declining wildlife populations. *Philos. Trans. R.*
Soc. B, 374, 20180211.

Sosa, S., Sueur, C. & Puga-Gonzalez, I. (2020). Network measures in animal social network
analysis: their strengths, limits, interpretations and uses. *Methods Ecol. Evol.*

Spiegel, O., Leu, S.T., Sih, A. & Bull, C.M. (2016). Socially interacting or indifferent
neighbours□? Randomization of movement paths to tease apart social preference and
spatial constraints, 971–979.

Stopher, K. V., Nussey, D.H., Guinness, F., Morris, A., Pemberton, J.M., Clutton-Brock, T.H.,
et al. (2012a). Re-mating across years and intralineage polygyny are associated with
greater than expected levels of inbreeding in wild red deer. *J. Evol. Biol.*, 25, 2457–
2469.

Stopher, K. V, Walling, C. a, Morris, A., Guinness, F.E., Clutton-brock, T.H., Pemberton,
J.M., *et al.* (2012b). Shared spatial effects on quantitative genetic parameters:
accounting for spatial autocorrelation and home range overlap reduces estimates of
heritability in wild red deer. *Evolution*, 66, 2411–26.

Thomson, C.E., Winney, I.S., Salles, O.C. & Pujol, B. (2018). A guide to using a Multiple-
Matrix animal model to disentangle genetic and nongenetic causes of phenotypic
variance. *PLoS One*, 13, e0197720.

VanderWaal, K.L., Atwill, E.R., Isbell, L.A. & McCowan, B. (2014). Linking social and
pathogen transmission networks using microbial genetics in giraffe (Giraffa
camelopardalis). *J. Anim. Ecol.*, 83, 406–414.

Vander Wal, E., Laforge, M.P. & McLoughlin, P.D. (2014). Density dependence in social
behaviour: Home range overlap and density interacts to affect conspecific encounter
rates in a gregarious ungulate. *Behav. Ecol. Sociobiol.*, 68, 383–390.

Webber, Q.M.R. & Vander Wal, E. (2018). An evolutionary framework outlining the
integration of individual social and spatial ecology. *J. Anim. Ecol.*, 87, 113–127.

Webber, Q.M.R. & Vander Wal, E. (2019). Trends and perspectives on the use of animal
social network analysis in behavioural ecology: a bibliometric approach. *Anim. Behav.*,
149, 77–87.

Yang, J., Lee, S.H., Goddard, M.E. & Visscher, P.M. (2011). GCTA: A Tool for Genome-wide
Complex Trait Analysis. *Am. J. Hum. Genet.*, 88, 76.

Zuur, A.F., Ieno, E.N. & Saveliev, A.A. (2017). *Beginner's guide to spatial, temporal, and*

spatial-temporal ecological data analysis with R-INLA. Highstat Ltd.

612