Heat Shock Protein 27 versus Estrogen Therapy for Post-Menopausal Atherosclerosis:

Rethinking Mechanisms of Cholesterol Lowering

Running Title: *HSP27 vs. E2 Rx Post-Menopause*

Nadia Maarouf PhD,[1] Yong-Xiang Chen MD, PhD[1], Chunhua Shi PhD[1], Jingti Deng MD, PhD[1],

Catherine Diao MSc[1], Matthew Rosin MSc[1], Vipul Shrivastava MSc[1], Zarah Batulan PhD[1],

Jingwen Liu PhD[2], Edward R. O'Brien MD[1#]

[#]corresponding author

[1]Department of Cardiac Sciences, Libin Cardiovascular Institute of Alberta, University of

Calgary Cumming School of Medicine, Calgary, Alberta, Canada

[2]Department of Veterans Affairs Palo Alto Health Care System, Palo Alto, California

Corresponding Author:

Edward R. O'Brien, MD, FRCPC, FAHA, FACC, FESC

Professor, Division of Cardiology, Department of Cardiac Sciences,

Libin Cardiovascular Institute of Alberta

Health Research & Innovation Centre, Rm GC68

3280 Hospital Drive NW, Calgary, Alberta,

Canada T2N 4Z6

Tel. 403-220-4525

Fax 403-210-9739

e-mail: ermobrie@ucalgary.ca

Abbreviations

AAb – anti-HSP27 (or anti-HSP25) antibodies

ApoE$^{-/-}$ - apolipoprotein E null

a.u. – absorbance units

E2 - 17β estradiol

HFD – high fat diet

HPRT - hypoxanthine-phosphoribosyltransferase

HSP27 – Heat Shock Protein 27

HSP27$^{o/e}$ – HSP27 over-expressing

LDL-C – low density lipoprotein cholesterol

LDLR – low density lipoprotein receptor

MP - menopause

mRNA – messenger RNA

NF-κB - nuclear factor kappa light chain enhancer of activated B cells

ON - overnight

OVX - ovariectomy

PBS - phosphate buffered saline

PCSK9 - Proprotein Convertase Subtilisin/Kexin type 9

rC1 – recombinant C-terminal truncation of HSP27 spanning amino acids 90-205

rHSP27 – recombinant HSP27

SDS-PAGE - sodium dodecyl sulfate and polyacrylamide gel

1   **Abstract**

The estrogen-inducible protein, Heat Shock Protein 27 (HSP27) attenuates experimental

atherogenesis by reducing cholesterol levels in the blood and artery wall. Blood levels of HSP27

and natural antibodies to HSP27 are higher in healthy controls compared to patients with

cardiovascular disease. Recently, we showed that vaccination of *ApoE$^{-/-}$* mice with recombinant

HSP25 (rHSP25, the murine ortholog of human HSP27) boosts anti-HSP25 levels and attenuates

atherogenesis in male and female *ApoE$^{-/-}$* mice. As estrogens and ovarian function influence

HSP27 synthesis, cellular release and blood levels, we hypothesize that menopause will result in

loss of HSP27 atheroprotection. Hence, we now compare the efficacy of rHSP25 vaccination *vs.*

estradiol (E2) therapy in preventing the increase in atherogenesis that occurs with surgical

menopause. Ovariectomized (OVX) *ApoE$^{-/-}$* fed a high fat diet were randomized to receive 5-

weekly rHSP25 vaccinations, a subcutaneous E2 slow release pellet or a combination thereof.

Anti-HSP25 IgG antibodies were generated in mice vaccinated with rHSP25 or rC1, an inactive

truncated form of HSP27 that acted as a control. OVX increased atherosclerotic burden by up to

65% and total plasma cholesterol levels by up to 23%. Compared to rC1/OVX (control) mice

there were dramatic reductions in atherogenesis with the following treatments: -43% with

rHSP25 vaccination, -52% with E2 therapy, and -82% with combined rHSP25 + E2 treatment

(p<0.0001 for all). Relative to the rC1/OVX control mice, terminal total cholesterol levels were

reduced by -22% with rHSP25 vaccination, -31% with E2 therapy and -45% with combined

rHSP25 + E2 treatment (p<0.0001 for all). Of note, plasma PCSK9 levels were higher in

E2/OVX mice than all other groups (p<0.0001). In contrast, OVX mice vaccinated with rHSP25

had PCSK9 levels that were -41% lower than rC1/OVX control mice (p<0.0001). Similarly,

PCSK9 plasma protein and hepatic mRNA levels in the E2/OVX group were higher by 52% and

148% relative to rC1/OVX control mice; (p<0.0001 for both). Hepatic LDLR mRNA levels in

the E2 treated mice were unchanged, but markedly increased with rHSP25 vaccination. In human

HepG2 hepatocytes E2 increased PCSK9 promoter activity in a dose-dependent manner - an

effect that could be inhibited with pan-estrogen receptor blocker ICI 182780. In contrast, the

combination of [rHSP27 + PAb] decreased PCSK9 promoter activity by 64%. In conclusion, E2

therapy attenuates atherogenesis post-OVX but is accompanied by increased PCSK9 expression.

In contrast, rHSP25 vaccination attenuates post-OVX atherogenesis and increases LDLR without

altering PCSK9 expression. These data suggest that for the post-menopausal preventative

treatment of atherosclerosis perhaps rHSP27 immunotherapy on its own, or in combination with

E2 should be considered.

Keywords:

Atherosclerosis

Cholesterol

PCSK9

LDL Receptor

Estrogen

Menopause

Introduction

Our laboratory is interested in understanding why atherosclerosis is relatively silent in women

compared to men – at least until menopause (MP) when estrogen levels plummet. Noting the

relative abundant presence of estrogen receptors in the artery wall, we embarked on an objective

discovery project looking for estrogen receptor associated proteins that might help explain

important biological differences. These studies led to the discovery that Heat Shock Protein 27

(HSP27) is as an estrogen receptor beta associated protein. The relationship between HSP27 and

estrogen is complex, as HSP27 acts as a repressor of estrogen-mediated transcription *in vitro*, yet

its expression and extracellular release are also partially regulated by estrogens (e.g., there is an

estrogen response element found in the HSP27 promoter) [1,2]. Importantly, our group [1] and four

others (using objective proteomic approaches) [3-6] showed that HSP27 expression is inversely

corelated with the degree of coronary artery plaque burden.

Accordingly, we then demonstrated in a series of experiments that augmenting HSP27 levels

reduces atherogenesis [7-10]. Moreover, when $ApoE^{-/-}$ mice that over-express HSP27 are subjected

to ovariectomy (OVX) HSP27 levels are lower and athero-protection is lost, unless estrogenic

therapy is administered [2]. Clinically, elevated HSP27 blood levels are associated with a lower 5-

year risk of myocardial infarction, stroke or cardiovascular death [8]. In women who undergo

premature surgical MP because of an inherited risk of cancer, we note that HSP27 levels drop in

the first post-operative year [11].

Interestingly, natural IgG auto-antibodies to HSP27 (AAbs) are detectable in human blood, and

recently we demonstrated how HSP27 immune complexes (ICs) form, dock at the cell membrane

engaging with Toll-like receptor 4 (TLR4) and compete with LPS to reduce inflammatory

signaling [12]. Via NF-κB pathway activation HSP27 ICs increase the expression of the low

density lipoprotein receptor (LDLR) [13]. Unlike the effect of statins, the effect of HSP27 on

LDLR expression is divergent from the sterol regulatory element binding protein 2-mediated

dual regulation of LDLR and Proprotein Convertase Subtilisin/Kexin type 9 (PCKS9), the

negative regulator of cholesterol metabolism. Furthermore, vaccination of mice with rHSP25

(murine ortholog of human HSP27), augments anti-HSP25 antibodies, reduces atherogenesis by

markedly decreasing total plasma cholesterol levels and increases LDLR expression [13].

Interestingly, with rHSP25 vaccination PCSK9 levels fall, and while this may in part be due to

reduced expression, it is likely due to increased PCSK9 clearance by abundant LDLRs [13].

Therefore, in this study we sought to determine if rHSP25 vaccination can combat atherogenesis

post-OVX, employing a surgical model of MP in *ApoE$^{-/-}$* mice and using estradiol (E2) as a

comparator. *In vitro*, we also explore the effects of the HSP27 IC *vs*. E2 on the expression of

PCSK9. Our results indicate that rHSP25 vaccination and E2 therapy are both effective in

attenuating the early stages of atherogenesis. However, we were surprised to find that E2

increases PCSK9 transcription without altering that of LDLR, while HSP27 therapy lowers

cholesterol (and PCSK9) by increasing LDLR expression.

Materials and Methods

Recombinant protein preparation

Plasmids encoding HIS-tagged full-length HSP27 were used to express the following

recombinant proteins: i) the mouse ortholog rHSP25, or the C-terminal, biologically inactive

(rC1) fragment of HSP27 (AA93-205) (**Fig. 1A**). For *in vitro* studies involving human

hepatocytes, we also generated rHSP27, as previously described [14]. Using a pET-21a vector, with

the plasmids were transformed into an *Escherichia coli* expression strain (DE3). Recombinant

proteins were purified with a Ni-NTA resin and Q-Sepharose[TM] (GE Healthcare). Endotoxin was

removed by Pierce High-Capacity Endotoxin Removal Resin (ThermoFisher Scientific,

Waltham, MA, USA). The purity of the final recombinant proteins was determined to be more

than 99% by sodium dodecyl sulfate and polyacrylamide gel (SDS-PAGE) with an endotoxin

concentration lower than 2 units/mg protein measured by Limulus Amebocyte Lysate

PYROGENT[TM] 125 Plus (Lonza). In previous experiments we demonstrated that these

recombinant proteins were functionally devoid of significant endotoxin contamination, as the

addition of polymyxin B had no effect on the ability of these proteins to activate NF-κB [9,15].

Murine Surgical Model of Menopause and Atherogenesis

The following animal protocol was approved by the Animal Care and Use Committee at the

University of Calgary and is in compliance with the Canadian Council on Animal Care (**Fig.**

1B). Five-week-old *ApoE*[-/-] female mice were purchased from the Jackson Laboratories (JAX®,

Canada; C57BL/6 genetic background, Jackson Laboratory, Bar Harbor, ME, USA; stock

002052: B6.129P2-*ApoE*[tm1Unc]/J) and after one week of acclimatization were randomized to

bilateral OVX *vs.* a surgical sham operation (i.e., ovaries not removed). For all surgical

procedures, mice were given a mixture of isoflurane/oxygen (2.5%, 1L/min respectively).

Briefly, the dorsum of each mouse was incised and both ovaries were removed. Muscle and skin

layers were closed with sterile synthetic bio-absorbable sutures (Ethicon, Somerville, NJ).

Buprenorphine (0.05 mg/kg subcutaneously) was administered twice-daily for 3 days post-

surgery for analgesia. For the mice randomized to receive E2 supplementation, after recovering

for one-week post-OVX a pellet containing 0.1 mg of E2 (60-day release formulation; Innovative

Research America) was implanted subcutaneously between the ear and shoulder using a trocar.

This strength of E2 pellet was selected because in a pilot experiment using higher concentration

E2 pellets (0.25 mg/60-day) there was a 3-fold increase in the uterine / fallopian tube weight in

the E2/OVX *vs.* rC1/OVX mice. Mice were also randomized to rHSP25 or rC1 vaccination, as

well as the combination of rHSP25 vaccination plus an E2 pellet. Vaccination was performed

weekly with recombinant HSP25 (rHSP25; 100 µg, 75 µL) or 'rC1', a truncated HSP27 peptide

consisting of the C-terminal half of the peptide, (100 µg, 75 µL) mixed with an aluminum

hydroxide gel adjuvant (Alhydrogel® 2%, Al, 25 µL; Invivogen vac-alu-250, San Diego, CA).

One week after the commencement of E2 and/or vaccinations, the mice began a 5-week

atherogenic High Fat Diet (HFD) containing: 1.25% cholesterol and 15.8% fat, (ENVIGO

Harlan Teklad diets) until euthanasia. In addition, control experiments were conducted

comparing PBS *vs.* rC1 treatments in sham and OVX mice (**Supplemental Fig. 1A**).

Mice were housed (n=3 or 4 per cage) in a temperature controlled (20ºC) facility with a 12 h

light/ dark cycles and had free access to autoclaved food and water. A total of 64 mice were

randomized to 8 groups (8 mice/group). Plasma samples were collected weekly and animal

weight recorded prior to treatment administration. At euthanasia, mice were placed under

anesthesia with isoflurane/oxygen mixture (5%, 1L/min) and adequate sedation was verified by

toe and/or tail pinching to ensure absence of a withdrawal reflex response. The thoracic cavity

was opened by cutting the ribs lateral to the sternum to expose the heart, and whole blood was

collected by inserting a 25-gauge needle into the left ventricle. After blood collection, the right

atrium pierced to permit hemorrhage, and systemic perfusion fixation with phosphate-buffered

saline (PBS) thrice or until the liver turned pale in color, followed by 10% NBF. The heart and

aorta were dissected, removed and immersed in NBF. The uteri and fallopian tubes were also

harvested, dissected of all fat and connective tissue, photographed and weighed in order to assess

the end organ effect of E2 *vs.* rHSP25 therapies.

Murine anti-HSP25 IgG antibody levels

Levels of IgG AAb were measured in murine plasma using an ELISA developed in our

laboratory (**Fig. 1C**). Briefly, NUNC maxisorp plates (ThermoFisher) were coated with rHSP25

at a concentration of 500 ng/well in carbonate-bicarbonate buffer overnight (ON) at room

temperature. The wells were blocked with 1% BSA/PBST, washed in phosphate buffered saline

tween (PBST), and incubated with plasma at a final dilution of 1:2,000 in 1% BSA for two hrs

followed by 3 more washes in PBST. A horse radish peroxidase (HRP) labeled a goat anti-mouse

IgG (H&L) antibody (catalogue #115-035-062; Jackson Immunoresearch, West Grove, PA) was

used as a detection antibody at a dilution of 1:5,000 and incubated for one hr at room

temperature. Finally, substrate solution (3,3',3.5'-Tetramethylbenzidine Liquid Substrate, TMB;

Millipore Sigma; Oakville, ON) was added to each well and incubated for 10 mins avoiding

direct light. The reaction was stopped by 2N H_2SO_4 and the optical density quantified at 450 nm,

using Synergy Mx plate reader (BioTek). To establish an internal standard for the measurement

of HSP25 auto-antibodies, plasma from a healthy control subject was diluted 500 times in 1%

BSA/PBST and arbitrarily defined as 50 absorbance units (a.u.).

Evaluation of *en face* Atherosclerotic Lesion Area

The aorta was dissected from the ascending to the thoracic segments ending at the 4[th] bronchial

branches and the adventitia was removed. The primary incision followed the lesser curvature of

the arch opening up the aorta longitudinally. A second incision was made along the greater

curvature of the arch down to the level of the left subclavian artery. The aortic lipid content was

evaluated using oil red O staining (described previously) [10] and photographed using an Olympus

SZ-61 stereo microscope with an Olympus UC30 camera (aortic magnification ×12; **Fig. 2A,**

Supplemental Fig. 1B). The *en face* atherosclerotic aortic lesions were analyzed using Image-

Pro software (Media Cybernetics) to calculate the total and atherosclerotic lesion areas. The

extent of atherosclerosis was expressed as the percentage of surface area of the entire aorta

covered by atherosclerotic lesions [8].

Murine Plasma Cholesterol Profiles and Lipoprotein Fractions

Total plasma cholesterol levels were determined using an enzymatic colorimetric assay kit

(Wako Pure Chemical Industries, Ltd, 439-17501) (**Fig. 2C, Supplemental Fig. 1D, 2C, 2D**).

The lipoprotein cholesterol distribution profile (VLDL, LDL, HDL) was evaluated in pooled

plasma samples under fasting conditions after fractionation by fast protein liquid

chromatography (FPLC) gel filtration on a single Superose 6 column (**Fig. 2D**).

LDLR and PCSK9 Gene Expression in Murine Liver Tissue

As previously described, RNA was isolated from mouse liver tissue and used for Real-time PCR

analysis of LDLR and PCSK9 gene expression (**Fig. 3A, 4B**).[16] The specific murine primer

sequence pairs, including those for the control gene, hypoxanthine-phosphoribosyltransferase

(hHPRT1), are provided in **Supplemental Table 1**. RNA expression is presented as log fold

change.

LDLR Protein Expression in Murine Liver Tissue

Protein was harvested from the murine livers for SDS-PAGE. Multiple Western blots were

obtained for each protein assayed, and representative data are shown in **Fig. 3B, 3C**. Mouse liver

tissue (200-300 mg) was placed inside a Precellys® homogenizing tube with 4 glass beads and

500 µL of RIPA lysis buffer (50 mM Tris HCL pH 7.4, 1% Triton X-100, 0.5% Sodium

Deoxycholate, 150 mM NaCl, 2mM EDTA, protease inhibitor cocktail). Tissue was

homogenized during two 30 s cycles at 6,000 RPM. The lysate was transferred to 2 mL tubes and

further processed using a handheld sonicator at 50% amplitude for two 3 s pulses, centrifuged 45

mins at 16,000×g, and the supernatant layer below the fat and above the pellet was transferred to

a new tube. Samples were centrifuged again for 10 mins at 16,000×g and the supernatants were

transferred to new tubes. Protein lysates were quantified using a Bio-Rad DC Protein Assay, with

equal amounts of protein (40-80 µg per well) loaded onto 10% Tris-Glycine gels for SDS-PAGE

before being transferred to 0.2 µm nitrocellulose membranes ON at 4°C. Membranes were

blocked in 5% Milk in TBS-Tween for 1h, washed, incubated in fresh primary antibodies with

5% bovine serum albumin (BSA) ON at 4°C, washed, and incubated in secondary antibody

before ECL detection using an ImageQuant LAS4000 imager (GE Healthcare). The primary

antibodies used for Western Blots prepared using murine liver lysates were rabbit anti-LDLR

(1:5,000; Abcam ab52818) and mouse anti-β-actin (1:10,000; sc-47778, Santa Cruz

Biotechnology). The secondary antibodies used for WB were: anti-rabbit IgG HRP (Abcam

ab97051) and anti-mouse IgG HRP (Jackson Immunoresearch 115-035-062).

PCSK9 Measurement

Plasma levels of PCSK9 (**Fig. 4A**) were measured using an ELISA kit specific to murine PCSK9

(R&D, DY3985) as per manufacturer's instructions. Samples were diluted 1:100 in 1% BSA and

a PCSK9 standard curve was generated for each experiment.

PCSK9 Promoter Activity Assays

Cells from the human hepatic HepG2 cell line were grown in complete media (with 10% FBS

and antibiotics) and seeded ON at 4×10^4 cells/well in a 48-well culture plates, then: i) transiently

transfected with a PCSK9 promoter firefly luciferase reporter construct, incorporated in the

plasmids pGL3-PCSK9-D1 [17], and ii) co-transfected with a transfection control vector

(pBLSV40 plasmid constitutively expressing a *Renilla* luciferase; generously donated by Dr.

David Proud University of Calgary, Calgary, AB), so that the dual luciferase activities could be

measured. The stoichiometry was: 500 ng of the PCSK9-D1 promoter construct and 25 ng of

pBLSV40 vector, using 1 μL of Lipofectamine®2000 Transfection Reagent (ThermoFisher).

Cells were then treated with recombinant proteins (rHSP27, rC1), antibodies (unrelated IgG

control, PAb), and combinations thereof for 24 hrs. PCSK9 promoter-driven firefly luciferase

activity and control renilla luciferase activity were measured using the Firefly & Renilla

Luciferase Single Tube Assay Kit (#30081-1; Biotium, Fremont, CA) according to the

manufacturer's protocol, with some minor modifications. Briefly, cells were washed with PBS

then lysed in 110 µL of passive lysis buffer. Firefly luciferase activity was measured first by

combining 30 µL of cell lysate with 50 µL of firefly working solution in a 12×75mm round-

bottom tube, measuring luminescence, then adding 5 µL of *Renilla* working solution to the same

tube and again measuring luminescence. Luminescence was detected using a Monolight 3010

Luminometer (BD Biosciences). Absolute firefly luciferase activity was normalized against

Renilla luciferase activity to correct for transfection efficiency. Triplicate wells were assayed for

each transfection condition, and 3 independent transfection assays were performed for each

reporter construct. The data was log transformed in order to facilitate parametric statistical

analyses (**Fig. 4C, 4D**).

Statistical analysis

For all data, continuous variables are expressed as mean ± standard error of the mean (SEM).

Statistical significance was assessed by using a one-way ANOVA (Graph Pad Prism 8, La

Jolla,CA, USA) and Tukey's comparison of the groups. The direct comparison of unpaired

treatment groups was performed using a Student's t-test. Quantitative PCR gene expression data

were log-transformed and are reported as log-fold change in order to more normally distribute

the data and permit parametric statistical analyses. A p-value < 0.05 was considered significant.

Results

a) Overview of rHSP25 Vaccination *vs*. E2 Therapy Experiment in OVX mice

To test the hypothesis that rHSP25 vaccination reduces atherogenesis in OVX mice, we

subjected atherosclerosis-prone *ApoE*$^{-/-}$ mice to OVX, followed by 6 weekly subcutaneous

injections with rHSP25 with the adjuvant Alum (**Fig. 1A, 1B**). Another group of mice underwent

a sham surgery (the ovaries were not removed) followed by rHSP25 vaccination. For comparison

with the clinical scenario of hormonal therapy for MP, a group of OVX *ApoE*$^{-/-}$ mice underwent

subcutaneous implantation of a slow release E2 pellet, while another group received both E2 and

rHSP25 vaccination. The inactive truncated C-terminus of HSP27, rC1 (mixed with Alum), was

administered as a negative control to OVX or sham surgery mice. All mice (7-8 per group)

received a HFD for 5 weeks, beginning one week after the initiation of therapies.

b) **Validation of the Vaccination response**

The rise in IgG anti-HSP25 levels were similar for all mice vaccinated with rHSP25 whether it

was combined with E2 or not (**Fig. 1C**). The antibody response to vaccination with rC1 was

similar to that with rHSP25, thereby indicating that rC1 remained immunogenic. E2 or PBS

treated mice had very low (background) levels of AAbs.

c) **Validation of the Murine Model of Menopause and Physiological Estrogen Therapy**

We confirmed that this OVX model was appropriate for the study of MP atherogenesis by

showing that OVX increased atherosclerotic burden and cholesterol levels. *ApoE*$^{-/-}$ mice

subjected to OVX had more aortic atherosclerosis than sham operated mice [e.g., PBS sham *vs.*

PBS OVX: 45% increase (11.8 ± 1.2 vs. 17.1 ± 1.7; respectively, p=0.03); and rC1 OVX

compared to rC1 sham: 65% increase (18.2 ± 1.9 *vs.* 11.0 ± 0.7; respectively, p=0.006)]

(Supplemental Fig. 1B, 1C). Similarly, total plasma cholesterol levels 1-day pre-euthanasia

were higher in OVX mice compared to sham operated mice [e.g., PBS/sham compared to

PBS/OVX: 23% higher (1,119 ± 32 *vs.* 1,374 ± 34; respectively, p=0.0001), and rC1/sham

compared to rC1/OVX: 21% higher (1,136 ± 23 *vs.* 1,369 ± 55; respectively, p=0.002)]

(Supplemental Fig. 1D). As measurement of estrogen levels in mice is technically challenging,

we confirmed that the estrogenic therapy in the E2 pellet (0.1 mg/60 day) provided a

physiological level of hormone replacement by examining the uteri at the end of the experiment.

The uterine weights of the E2 treated mice were similar to those of the sham mice, thereby

confirming that the E2 dose was likely in a physiological range (**Fig. 1D, 1E**). In contrast, the

rHSP25/OVX uteri were similar to those of the rC1/OVX mice showing 4.5-fold atrophy *vs.*

rC1/sham mice uteri; thus, indicating that rHSP25 had no direct or indirect estrogenic properties.

d)   **Validation of rC1 as a Negative Control Treatment in OVX / Atherogenesis Model**

Four control groups (PBS/sham, rC1/sham, PBS/OVX and rC1/OVX) were used to assess the

validity of using rC1 as a negative control. rC1 is an excellent control treatment as it is prepared

in an identical manner as rHSP25 using recombinant technology in *E. coli*. Although these

recombinant proteins are treated to reduce endotoxin contaminants to very low levels, if there is

residual endotoxin present in one protein it is balanced by the same risk in the other protein. The

degree of atherosclerosis was similar for the sham operated mice that received PBS *vs.* rC1 (11.8

± 1.19 *vs.* 11.03 ± 0.65 respectively; p=0.58), as well as the OVX mice that received PBS *vs.* rC1

(17.1 ± 1.67 *vs.* 18.2 ± 1.93; respectively, p=0.68) (**Supplemental Fig. 2A, 2B**). Likewise,

cholesterol levels were similar amongst the PBS sham *vs.* rC1 sham mice (1,119 ± 31.64 *vs.*

1,136 ± 22.83; respectively, p=0.67) and the PBS OVX *vs*. rC1 OVX mice (1,374 ± 33.75 *vs*.

1,369 ± 54.88; respectively, p= 0.94) (**Supplemental Fig. 2C, 2D**). Therefore, for the purpose of

comparing the effect of the active treatment groups (rHSP25 or E2), the rC1/OVX group was

used as the reference group for the effect of OVX, as the degree of atherogenesis as well as the

cholesterol levels were similar to those found in the PBS/OVX cohort.

e) **rHSP25 Vaccination and E2 Therapy Post-OVX are Comparable for Attenuation of**

Atherogenesis and Lowering Cholesterol

Mice subjected to OVX alone (rC1/OVX) had 51% more aortic atherosclerosis compared to non-

OVX (rC1/sham) mice (16.7 ± 1.4 *vs.* 11.0 ± 0.6; **Fig. 2A, 2B**). The atherosclerotic burden was

similar in the rHSP25/OVX and E2/OVX mice, and markedly reduced compared to rC1/OVX

(p<0.0001). Interestingly, the combination of [rHSP25 + E2] in the OVX mice showed the

lowest atherosclerotic burden (-82% *vs*. rC1/OVX; p<0.0001). Finally, the atherosclerotic burden

was lower in the rHSP25/sham *vs.* rC1/sham mice (p=0.04).

While the baseline cholesterol levels in all mice fed a normal chow diet were similar (515 ± 82

mg/dL), by the second week of the HFD the levels peaked and then fell by varying degrees over

the ensuing 3 weeks of the HFD. Terminal plasma cholesterol levels (**Fig. 2C**) were measured 1-

day prior to euthanasia, with the highest levels found in the rC1/OVX group (1,393 ± 57 mg/dl),

which was 21% higher (p<0.0001) than the rC1/sham mice (1,148 ± 23 mg/dl). Both the

rHSP25/OVX and E2/OVX mice had similar cholesterol levels that were lower than the

rC1/OVX mice (1,068 ± 8 and 964 ± 27; both p<0.0001 *vs.* rC1/OVX; respectively), while the

lowest levels were found in the combined therapy [rHSP25 + E2/OVX] mice (736 ± 24;

p<0.0001 *vs.* all OVX groups). Cholesterol levels were lower in the rHSP25/sham compared to

rC1/sham mice (p<0.0001). As demonstrated by FPLC, the LDL-C fraction was reduced in both

the E2/OVX and rHSP25/OVX groups relative to the rC1/OVX mice (**Fig. 2D**).

f) **Unlike rHSP25 Vaccination, E2 Therapy Did not Increase LDLR Expression Post-OVX**

rHSP25 alone or in combination with E2 markedly increased hepatic LDLR mRNA expression

(+219% for the rHSP25/OVX *vs.* rC1/OVX and +165% for the [rHSP25 + E2] *vs.* E2/OVX

mice; p<0.0001 for both; **Fig. 3A**). LDLR protein expression also increased in the mice

vaccinated with rHSP25 (50% for the rHSP25/OVX *vs.* rC1/OVX, 51% for the [E2 +

rHSP25/OVX] *vs.* E2/OVX; p=0.04 and p=0.006, respectively; **Fig. 3B, 3C**). Surprisingly, E2

therapy in isolation had no effect on LDLR mRNA or protein expression.

g) **E2 but Not rHSP25 Vaccination Increases PCSK9 Gene Expression**

Plasma PCSK9 levels were much higher in E2/OVX (464 ± 13 ng/ml) compared to the other four

treatment groups (*p*<0.0001) – including a 52% increase relative to rC1/OVX (**Fig. 4A**).

Combining [rHSP25 + E2] therapy post-OVX resulted in a 23% reduction in PCSK9 levels (357

± 12 ng/ml; p=0.0001) relative to E2/OVX alone. Plasma PCSK9 levels were 26% lower in the

rHSP25/sham *vs.* rC1/sham mice (p=0.046). Hepatic PCSK9 mRNA expression was 148%

higher in the E2/OVX compared to rC1/OVX and all other murine treatment groups (p<0.0001)

(**Fig. 4B**).

The PCSK9 promoter is thought to contain two estrogen response elements (EREs), as predicted

in silico using the MatInspector analysis tool (optimized matrix threshold = 0.9 for both;

https://www.genomatix.de/online_help/help_matinspector/matinspector_help.html). Hence, we

assessed PCSK9 promoter activity in HepG2 cells treated for 24 hrs with escalating

concentrations of E2 (0.1, 10, 100nM) and noted a robust dose-dependent increase in activity

(+505% with 100 nM E2, p=0.0005) that was reduced by 46% (p=0.02) with the pan estrogen

receptor blocker, ICI 182,780 (**Fig. 4C**). Likewise, treating HepG2 cells for 24 hrs with rHSP27

alone (100 µg) had no effect, yet there was a 64% decrease in PCSK9 promoter activity (p=0.01)

when cells were treated with [rHSP27 + PAb] (**Fig. 4D**). Moreover, there was a remarkable

303% increase in PCSK9 promoter activity with E2 treatment (100 nM; p<0.0001) that shrunk

by 43% (p<0.0001) with [rHSP27 + PAb] treatment.

Discussion

Using a surgical model of MP in atherosclerosis-prone *ApoE*$^{-/-}$ mice fed a HFD we demonstrate

that rHSP25 vaccination, like E2 therapy, is atheroprotective. As well, we confirmed a recent

observation [13] that rHSP25 vaccination promotes the expression of the LDLR – an effect that is

not associated with E2 therapy. Unlike rHSP25 vaccination, E2 therapy is associated with

increased PCSK9 synthesis and plasma levels (**Fig. 4A, 4B**), and showed a dose dependent

increase in PCSK9 promoter activity that was partially abrogated with estrogen receptor

blockade or concomitant treatment with [rHSP27 + PAb] (**Fig. 4C, 4D**).

While it was surprising to find that E2 treatment did not increase LDLR expression, perhaps it

was predictable? The specific mechanisms that account for the relative protection against

atherosclerotic complications in pre-MP women are poorly understood, yet there is a common

belief that one of the main cardiovascular benefits of estrogens is the lowering of LDL-C due to

an increase in LDLR expression. However, on reviewing the pre-clinical and clinical literature,

the effects of estrogens on LDLR vary. For example, in rats LDLR activity and expression are

reduced with OVX but improve with estrogen therapy [18,19]. But, in non-OVX mice treated with

(even) supra-physiological E2 concentrations, LDLR expression does not change [20,21]. As the

ApoE[-/-] mice in our study did not show an increase in LDLR with E2, this does not seem to be

due to an inherent problem with LDLR expression that is unique to this strain of mice – as it is

clear that these same mice showed a robust increase in LDLR expression in response to rHSP25

vaccination. Finally, it is interesting to note that E2 treatment is even atheroprotective in *LDLR[-/-]*

mice – thereby pointing to beneficial effects of estrogens that may be at a distance from LDLR

modulation [22].

Similarly, the evidence that E2 modulates LDLR expression in human hepatocytes is far from

convincing [23,24]. For example, there is indirect evidence of increased LDLR binding activity in

liver biopsy tissue from two male patients receiving estrogen therapy for prostate cancer [25]. The

doses of estrogenic compounds administered to these men were not in a physiological range, and

their cholesterol levels were not lower than the normal. Given that >1,000 human hepatic genes

display a sex-bias in their expression, particularly those known to participate in or regulate lipid

metabolism, and that the LDLR gene is more highly expressed in the female liver, the value of

these experimental results in male subjects should be interpreted with caution [26].

PCSK9 was only discovered less than 2 decades ago, and there is a lot to learn about its

transcriptional regulation, including the potential role of two estrogen response elements in its

promoter. Nonetheless, the estrogen mediated upregulation of PCSK9 expression noted in the

current studies does fit with clinical observations showing that: i) PCSK9 levels are higher in

pre-MP women than men [27], and ii) PCSK9 levels are markedly elevated in pregnant compared

to non-pregnant age-matched women (493 *vs.* 290 ng/ml) [28]. This would fit with the fact that in

the third trimester of pregnancy E2 levels increase 25-50 fold and total cholesterol levels rise 50-

75% [29]. Hence, we postulate that E2 drives these increases in PCSK9 and cholesterol levels in

pregnancy. Whether the surges in E2 levels that occur during the normal menstrual cycle or with

exogenous estrogenic therapies have the same effects on PCSK9 levels remains to be studied.

One study that does not fit with our observations on the role of E2 in upregulating PCSK9 levels

is the Dallas Heart Study. Specifically, these investigators found that PCSK9 levels are higher in

post- *vs.* pre-MP women, and that there was no difference in PCSK9 levels between women

receiving estrogenic compounds or not [27]. However, it is not clear if the higher PCSK9 levels in

the older post-MP women are clearly related to loss of ovarian function, or that the estrogenic

therapy that they received was physiological. Indeed, a number of other studies suggest that

PCSK9 levels simply correlate with age, and this may be difficult to discount too [30-32].

Although we did not see major differences in atheroprotection with E2 *vs.* rHSP25 therapy in

vivo, we have uncovered some subtle mechanistic differences that require further consideration –

particularly the increase in PCSK9 with estrogens. To summarize the results of our studies, we

constructed a working hypothesis incorporating the interplay between estrogens and HSP27, as

well as PCSK9 and LDLR (**Fig. 5**). Briefly, E2 upregulates PCSK9 expression, yet it is also well

known that E2 increases HSP27 expression and cellular release [2,33,34]. HSP27 markedly increases

LDLR expression, and given the rapid recycling of the LDLR, this increase in LDLR synthesis

may overpower any change in PCSK9 levels by binding PCKS9 and subjecting it to intracellular

disposal. However, in the MP scenario estrogen levels are diminished, and while PCSK9

expression is not augmented because of this, there may also be inadequate HSP27 levels to

stimulate LDLR expression and the clearance of PCSK9. As a result, LDL-C and PCSK9 levels

are higher compared to pre-MP. Hence, what needs to be emphasized about these data are: i) the

dramatic HSP27-induced increase in LDLR (also reported elsewhere recently)[13] that not only

reduces LDL-C but also PCSK9 levels, and ii) the previously unrecognized E2-mediated

increase in PCSK9 expression. As long as E2 therapy is increasing HSP27 (and therefore LDLR)

levels to the same extent there will be clearance of PCSK9 and a balanced lipoprotein profile.

However, we do not know if this is uniformly the case for all estrogenic compounds (e.g., there

may estrogen receptor specific subtleties with regards to LDLR and PCSK9 expression that have

yet to be uncovered). Further study is required, particularly if PCSK9 expression may be

increased by some estrogenic compounds, and therefore represent a cardiovascular or other

hazard.

Currently, we are working towards understanding if this working hypothesis can be further

evaluated in humans, and explore the virtues of estrogen receptor specific modulators [11,34]. As

well, we are trying to better comprehend how HSP27 (or HSP25) alone or in combination with

its autoantibody, modulate LDLR and PCSK9 transcription. We do know that in the presence of

AAbs, HSP27 shows enhanced docking at the cell membrane, interacts with cell membrane

receptors and signaling pathways, and can undergo internalization [12]. As low HSP27 and AAb

levels are associated with atherosclerosis,[13] we want to develop immunotherapies that boost AAb

levels in order to potentiate the biological activity of existing (low) levels of HSP27, especially

in women post-MP.

Acknowledgments:

Author Contributions

N. Maarouf and E.R. O'Brien designed the studies. N. Maarouf performed all *in vivo*

experiments. N. Maarouf, Y.-X. Chen, C. Shi, J. Deng, C. Diao, M. Rosin, V. Shrivastava

performed or contributed to some of the *in vitro* and *ex vivo* research; N. Maarouf, C. Diao and J.

Liu performed or contributed to the design and analysis of the promoter studies; N. Maarouf, V.

Shrivastava, Z. Batulan and E.R. O'Brien analyzed all other data; N. Maarouf and E.R. O'Brien

wrote the manuscript.

Conflict of Interest

EOB and YXC are inventors on US patents 8343915B2 and 8343916B2; EOB, CS and YXC are

inventors on US patent application PCT/CA2016/051018; EOB is the inventor on US patent

application 63/031,6402020; all intellectual property pertains to HSP27 diagnostics /

therapeutics. EOB is the Scientific Co-Founder of Pemi31 Therapeutics Inc., a start-up company

that controls the aforementioned intellectual property. EOB, CS and YXC have equity interests

in Pemi31 Therapeutics Inc.

Figure Legends

Fig. 1. rHSP25 vaccination to prevent atherogenesis in ovariectomized *ApoE*[-/-] mice.

A) Schematic representation of rHSP25 (mouse ortholog of HSP27) and the biologically

4   inactive truncated C-terminal of HSP27 (rC1 control) proteins. The black box denotes the

5   alpha beta-crystallin domain that is important for protein oligomerization. The IXI box at

6   the C-terminus is a flexible domain involved in the formation of multiple inter-subunit

7   interactions.

8   B) Age matched, sexually mature female *ApoE*[-/-] mice were subjected to either OVX or

9   sham operation (no OVX). One week later, they were randomized to the following

treatments: rC1/OVX, rHSP25/OVX, E2/OVX, rHSP25+E2/OVX, rHSP25/sham and

rC1/sham. A HFD was initiated one week later and continued for 5 weeks until

euthanasia. There were 7-8 mice per group. See Supplemental Fig. 1 and 2 for additional

control mice data.

C) Anti-HSP25 IgG levels increased in all mice vaccinated with either rHSP25 or rC1. The

initiation of rHSP25 vaccination began at 7 weeks of age (one-week post-surgery) and

measurements were taken every week thereafter, rising to a plateau by 3-4 weeks after

the first vaccination and continuing at this level until euthanasia. Data are displayed as

a.u. for OD @ 450 nm.

D) Representative photographs of uteri and fallopian tubes from each treatment groups.

1   **Fig. 1.** (continued)

2   E) Combined uterine and fallopian tube weights at time of euthanasia. After OVX there was

3   marked atrophy of the uteri and fallopian tubes of mice in the rC1/OVX and

4   rHSP25/OVX groups compared to rC1/sham mice. There was no difference in weights

5   between the rC1/sham and the other groups (E2/OVX, [E2 + rHSP25/OVX],

6   rHSP25/sham), thereby suggesting that the E2 supplementation was in a physiological

7   hormonal range.

Fig. 2. Atherosclerosis and Hyperlipidemia after surgical MP

A) Photomicrographs of *en face* thoracic aortae stained with Oil Red O.

B) Quantitation of aortic atherosclerotic burden, as reflected by Oil Red O staining. Compared

to the rC1 sham group there was an increase in aortic lesion area with rC1/OVX. Separately,

rHSP25 and E2 reduced aortic atherosclerotic burden compared to rC1/OVX, but the largest

decrease was noted with combined [E2+RHSP25] therapy. The atherosclerotic burden was

lower in the rHSP25/sham *vs.* rC1/sham mice (p=0.04).

C) OVX resulted in an increase in total plasma cholesterol in the rC1 mice. Compared to

rC1/OVX, cholesterol levels were lower post-OVX with either rHSP25 or E2 therapies, and

lowest with combined [rHSP25+E2] treatment. Cholesterol levels in the rHSP25/sham *vs.*

rC1/sham mice (p<0.0001).

D) Cholesterol measured in FPLC plasma fractions from pooled blood samples from rC1/OVX,

rHSP25/OVX and E2/OVX group *ApoE$^{-/-}$* mice (n=8 per group). Lipid sub-fractions based on

size are indicated above the cholesterol graphs. Compared to the rC1/OVX mice, there are

reductions in the LDL subfraction in both rHSP25/OVX and E2/OVX groups.

Fig. 3 rHSP25 and E2 differentially regulate of LDLR

A) Compared to rC1/OVX mice, hepatic LDLR mRNA increased dramatically in the

rHSP25/OVX but not the E2/OVX mice. In the OVX mice, adding rHSP25 to E2 noticeably

increased hepatic LDLR mRNA expression compared to the E2/OVX group. There was a

marked increase in LDLR expression in the rHSP25/sham *vs.* the rC1/sham mice.

B) Hepatic LDLR protein expression in the OVX experiment involving *ApoE$^{-/-}$* mice, assessed

by Western blotting. Each LDLR protein band was normalized using β-actin as the loading

control.

C) LDLR protein expression was similar in the rC1/sham, rC1/OVX and E2/OVX mice, but

appreciably increased with rHSP25 vaccination or combined [rHSP25 + E2] treatment post-

OVX. LDLR protein expression was similarly increased in the rHSP25/sham *vs.* rC1/sham

mice.

Fig. 4 rHSP25 and E2 differentially regulate of PCSK9

A) Plasma PCSK9 protein levels increased with E2 treatment in the OVX mice, an effect that

was attenuated when co-treated with rHSP25. rHS25 vaccination (alone) had no appreciable

effects on plasma PCSK9 levels with or without OVX.

B) Hepatic PCSK9 mRNA expression dramatically increased with E2 treatment (alone), but did

not change when co-treated with rHSP25. Hepatic PCSK9 mRNA levels were unchanged in

rHSP25 vaccinated mice and were unaltered by OVX.

C) PCSK9 promoter activity in HepG2 cells was strongly upregulated by E2 in a dose dependent

manner, an effect that was partially attenuated with the pan-ER blocker, ICI 182,780.

D) Compared to control, rHSP27+PAb significantly reduced PCSK9 promoter activity, while E2

markedly upregulated PCSK9 promoter activity – an effect that was partially reduced by

adding treatment with rHSP27.

Fig. 5. Working Model for HSP27 IC Modulation of PCSK9 and LDL-R Pre- and Post-

Menopause

A) Pre-MP estrogen upregulate both PCSK9 and HSP27 expression. However, HSP27 combines

with its auto-antibody (AAb) to augment LDLR expression that then facilitates PCSK9

clearance. The HSP27 IC also has a direct effect on PCSK9 expression. The net effects are

reduced LDL-C and PCSK9 levels.

B) Post-MP estrogen levels plummet and there is no increase in HSP27 to modulate LDLR and

PCKS9, resulting in elevated LDL-C and PCSK9 levels.

1   **References**

1.   Miller H, Poon S, Hibbert B, Rayner K, Chen YX, O'Brien ER. Modulation of estrogen

signaling by the novel interaction of heat shock protein 27, a biomarker for

atherosclerosis, and estrogen receptor beta. *Arteriosclerosis Thrombosis and Vascular*

Biology. 2005;25:E10-E14

2.   Rayner K, Sun J, Chen YX, McNulty M, Simard T, Zhao X, Wells DJ, de Belleroche J,

O'Brien ER. Heat shock protein 27 protects against atherogenesis via an estrogen-

dependent mechanism: Role of selective estrogen receptor beta modulation.

Arteriosclerosis, thrombosis, and vascular biology. 2009;29:1751-1756

3.   Martin-Ventura JL, Duran MC, Blanco-Colio LM, Meilhac O, Leclercq A, Michel JB,

Jensen ON, Hernandez-Merida S, Tunon J, Vivanco F, Egido J. Identification by a

differential proteomic approach of heat shock protein 27 as a potential marker of

atherosclerosis. *Circulation*. 2004;110:2216-2219

4.   Park HK, Park EC, Bae SW, Park MY, Kim SW, Yoo HS, Tudev M, Ko YH, Choi YH,

Kim S, Kim DI, Kim YW, Lee BB, Yoon JB, Park JE. Expression of heat shock protein

27 in human atherosclerotic plaques and increased plasma level of heat shock protein 27

in patients with acute coronary syndrome. *Circulation*. 2006;114:886-893

5.   Lepedda AJ, Cigliano A, Cherchi GM, Spirito R, Maggioni M, Carta F, Turrini F,

Edelstein C, Scanu AM, Formato M. A proteomic approach to differentiate histologically

classified stable and unstable plaques from human carotid arteries. *Atherosclerosis*.

2009;203:112-118

6. Liang W, Ward LJ, Karlsson H, Ljunggren SA, Li W, Lindahl M, Yuan XM. Distinctive

proteomic profiles among different regions of human carotid plaques in men and women.

Sci Rep. 2016;6:26231

7. Rayner K, Chen YX, McNulty M, Simard T, Zhao X, Wells DJ, de Belleroche J, O'Brien

ER. Extracellular release of the atheroprotective heat shock protein 27 is mediated by

estrogen and competitively inhibits acldl binding to scavenger receptor-a. *Circulation*

research. 2008;103:133-141

8. Seibert TA, Hibbert B, Chen YX, Rayner K, Simard T, Hu T, Cuerrier CM, Zhao X, de

Belleroche J, Chow BJ, Hawken S, Wilson KR, O'Brien ER. Serum heat shock protein 27

levels represent a potential therapeutic target for atherosclerosis: Observations from a

human cohort and treatment of female mice. *J Am Coll Cardiol*. 2013;62:1446-1454

9. Raizman JE, Chen YX, Seibert T, Hibbert B, Cuerrier CM, Salari S, Zhao X, Hu T, Shi

C, Ma X, Simard T, Caravaggio J, Rayner K, Bowdish D, Moore K, O'Brien ER. Heat

shock protein-27 attenuates foam cell formation and atherogenesis by down-regulating

scavenger receptor-a expression via nf-kappab signaling. *Biochimica et biophysica acta*.

2013;1831:1721-1728

10. Cuerrier CM, Chen YX, Tremblay D, Rayner K, McNulty M, Zhao X, Kennedy CR, de

BJ, Pelling AE, O'Brien ER. Chronic over-expression of heat shock protein 27 attenuates

atherogenesis and enhances plaque remodeling: A combined histological and mechanical

assessment of aortic lesions. *PLoS ONE*. 2013;8:e55867

11.   Batulan Z, Maarouf N, Shrivastava V, O'Brien E. Prophylactic salpingo-oophorectomy &

2   surgical menopause for inherited risks of cancer: The need to identify biomarkers to

3   assess the theoretical risk of premature coronary artery disease. *Women's Midlife Health*.

4   2018;4:7

5   12.   Shi C, Deng J, Chiu MH, Chen Y-X, O'Brien ER. Heat shock protein 27 immune

6   complex altered signaling and transport (icast): Novel mechanisms of attenuating

7   inflammation. *bioRxiv*. 2020:2020.2005.2031.126581

8   13.   Chen Y-X, Shi C, Deng J, Diao C, Maarouf N, Rosin M, Shrivastava V, Hu A, Bharadwa

9   S, Adijiang A, Ulke-Lemee A, Gwilym BL, Hellmich A, Malozzi C, Batulan Z, Dean

JLE, Ramirez FD, Liu J, Gerthoffer WT, O'Brien ER. Heat shock protein 27 immune

complex upregulates ldlr expression thereby reducing plasma cholesterol and

atherogenesis. *bioRxiv*. 2020:2020.2005.2021.102350

14.   Chiu MH, Shi C, Rosin M, Batulan Z, O'Brien ER. Biophysical analyses and functional

implications of the interaction between heat shock protein 27 and antibodies to hsp27.

Biochimica et Biophysica Acta (BBA) - General Subjects. 2019

15.   Salari S, Seibert T, Chen YX, Hu T, Shi C, Zhao X, Cuerrier CM, Raizman JE, O'Brien

ER. Extracellular hsp27 acts as a signaling molecule to activate nf-kappab in

macrophages. *Cell Stress Chaperones*. 2013;18:53-63

16.   Pulakazhi Venu VK, Adijiang A, Seibert T, Chen YX, Shi C, Batulan Z, O'Brien ER.

Heat shock protein 27-derived atheroprotection involves reverse cholesterol transport that

is dependent on gm-csf to maintain abca1 and abcg1 expression in apoe(-/-) mice. *FASEB J*. 2017;31:2364-2379

17. Li H, Dong B, Park SW, Lee HS, Chen W, Liu J. Hepatocyte nuclear factor 1 alpha plays a critical role in pcsk9 gene transcription and regulation by the natural hypocholesterolemic compound berberine. *J Biol Chem*. 2009;284:28885-28895

18. Kovanen PT, Brown MS, Goldstein JL. Increased binding of low density lipoprotein to liver membranes from rats treated with 17 alpha-ethinyl estradiol. *J Biol Chem*. 1979;254:11367-11373

19. Ngo Sock ET, Chapados NA, Lavoie JM. Ldl receptor and pcsk9 transcripts are decreased in liver of ovariectomized rats: Effects of exercise training. *hmr*. 2014;46:3-3

20. Srivastava RAK, Baumann D, Schonfeld G. In vivo regulation of low☐density lipoprotein receptors by estrogen differs at the post☐transcriptional level in rat and mouse. *European Journal of Biochemistry*. 1993;216:527-538

21. van Vlijmen BJ, van 't Hof HB, Mol MJ, van der Boom H, van der Zee A, Frants RR, Hofker MH, Havekes LM. Modulation of very low density lipoprotein production and clearance contributes to age- and gender- dependent hyperlipoproteinemia in apolipoprotein e3-leiden transgenic mice. *J Clin Invest*. 1996;97:1184-1192

22. Shelton KA, Cline JM, Cann JA. 17-β estradiol reduces atherosclerosis without exacerbating lupus in ovariectomized systemic lupus erythematosus-susceptible ldlr–/– mice. *Atherosclerosis*. 2013;227:228-235

23.   Semenkovich CF, Ostlund RE, Jr. Estrogens induce low-density lipoprotein receptor

activity and decrease intracellular cholesterol in human hepatoma cell line hep g2.

Biochemistry. 1987;26:4987-4992

24.   Wade DP, Knight BL, Soutar AK. Hormonal regulation of low-density lipoprotein (ldl)

receptor activity in human hepatoma hep g2 cells. Insulin increases ldl receptor activity

and diminishes its suppression by exogenous ldl. *Eur J Biochem*. 1988;174:213-218

25.   Angelin B, Olivecrona H, Reihnér E, Rudling M, Ståhlberg D, Eriksson M, Ewerth S,

Henriksson P, Einarsson K. Hepatic cholesterol metabolism in estrogen-treated men.

Gastroenterology. 1992;103:1657-1663

26.   Zhang Y, Klein K, Sugathan A, Nassery N, Dombkowski A, Zanger UM, Waxman DJ.

Transcriptional profiling of human liver identifies sex-biased genes associated with

polygenic dyslipidemia and coronary artery disease. *PLOS ONE*. 2011;6:e23506

27.   Lakoski SG, Lagace TA, Cohen JC, Horton JD, Hobbs HH. Genetic and metabolic

determinants of plasma pcsk9 levels. *J Clin Endocrinol Metab*. 2009;94:2537-2543

28.   Peticca P, Raymond A, Gruslin A, Cousins M, Adetola E, Abujrad H, Mayne J, Ooi TC.

Human serum pcsk9 is elevated at parturition in comparison to nonpregnant subjects

while serum pcsk9 from umbilical cord blood is lower compared to maternal blood. *ISRN*

Endocrinol. 2013;2013:341632

29.   Abbassi-Ghanavati M, Greer LG, Cunningham FG. Pregnancy and laboratory studies: A

reference table for clinicians. *Obstetrics & Gynecology*. 2009;114:1326-1331

30.   Cui Q, Ju X, Yang T, Zhang M, Tang W, Chen Q, Hu Y, Haas JV, Troutt JS, Pickard RT,

Darling R, Konrad RJ, Zhou H, Cao G. Serum pcsk9 is associated with multiple

metabolic factors in a large han chinese population. *Atherosclerosis*. 2010;213:632-636

31.   Ruscica M, Ferri N, Fogacci F, Rosticci M, Botta M, Marchiano S, Magni P, D'Addato S,

Giovannini M, Borghi C, Cicero AFG, Brisighella Heart Study G. Circulating levels of

proprotein convertase subtilisin/kexin type 9 and arterial stiffness in a large population

sample: Data from the brisighella heart study. *Journal of the American Heart*

Association. 2017;6:e005764

32.   Jeenduang N. Circulating pcsk9 concentrations are increased in postmenopausal women

with the metabolic syndrome. *Clin Chim Acta*. 2019

33.   Shi C, Ulke-Lemee A, Deng J, Batulan Z, O'Brien ER. Characterization of heat shock

protein 27 in extracellular vesicles: A potential anti-inflammatory therapy. *FASEB J*.

2019;33:1617-1630

34.   Sun J, Ma X, Chen YX, Rayner K, Hibbert B, McNulty M, Dhaliwal B, Simard T,

Ramirez D, O'Brien E. Attenuation of atherogenesis via the anti-inflammatory effects of

the selective estrogen receptor beta modulator 8beta-ve2. *Journal of Cardiovascular*

Pharmacology. 2011;58:399-405

Fig. 1

A

Full length rodent (rHSP25)

Truncated C-terminus (rC1)

B

chow diet | 5 weeks on HFD + weekly rHSP25 vaccination (and blood sampling)

Surgery

E2 and Vaccination Initiation

Week 13 Euthanasia

AGE (weeks)

6 7 8 9 10 11 12 13

rC1/OVX   rHSP25/OVX   E2/OVX   E2 + rHSP25/OVX   rHSP25/Sham   rC1/Sham

Bilateral ovariectomy

Sham surgery

C

Vaccination #

1 2 3 4 5 6

Anti-HSP25 IgG (a.u.)

Age (weeks)

- rHSP25/Sham
- rC1/OVX
- rHSP25/OVX
- PBS/OVX
- PBS/Sham
- E2+rHSP25/OVX
- E2/OVX
- rC1/Sham

D

rC1/Sham rC1/OVX rHSP25/OVX E2/OVX E2+rHSP25/OVX rHSP25/Sham

E

Uterine + fallopian tube wts (mg)

p=NS

Atrophy: -4.5-fold

p=NS

rC1/Sham rC1/OVX rHSP25/OVX E2/OVX E2+rHSP25/OVX rHSP25/Sham

Fig. 2

A

B

C

D

Fig. 3

A

B

C

Fig. 4

Fig. 5

A

Pre-Menopause: HSP27 & Cholesterol Metabolism

⬆ Estrogen ➡ PCSK9 ⊖ (decreased PCSK9 transcription)

⬆ HSP27 --AAb-- AAb➡ ⬆ LDL-R

Increased PCSK9 clearance

Net Effect: "Reduced" LDL-C / PCSK9

B

Post-Menopause: HSP27 & Cholesterol Metabolism

⬇ Estrogen → PCSK9 "Normal" PCSK9 expression & clearance

✗

⬇ HSP27 --AAb ✗-- AAb→ LDL-R (basal)

⊖

Net Effect: "Elevated" LDL-C / PCSK9