Impact of trimethoprim on the river microbiome and antimicrobial resistance.

Delaney, J[1]., Raguideau, S[2]., Holden, J[1]., Zhang, L[3]., Tipper, H.J[4]., Hill, G.L[1]., Klümper, U[2]., Zhang, T[5].,

Cha, C.J.[3], Lee, K[3]., James, R[10]., Travis, E[1]., Bowes, M. J[4]., Hawkey, P.M[6]., Lindstrom H.S[7,8]., Tang, C[7,8].,

Gaze, W.H[3*]., Mead, A[9*]., Quince, C[10*]., Singer, A[4*]., Wellington, E.M.H[1*].

[1]School of Life Sciences, University of Warwick, Coventry, UK

[2]Warwick Medical School (WMS), University of Warwick, Coventry, UK.

[3]European Centre for Environment and Human Health, University of Exeter Medical School, ESI,

Penryn Campus, Cornwall, UK

[4]UK Centre for Ecology and Hydrology, Wallingford, UK.

[5]Department of Civil Engineering, The University of Hong Kong, Hong Kong, SAR, China.

[6]Institute of Microbiology and Infection, University of Birmingham, Edgbaston, UK.

[7]Department of Chemistry, Umeå University, 901 87 Umeå, Sweden.

[8]Department of Public Health and Clinical Medicine, Occupational and Environmental Medicine, Umeå

University, 901 87 Umeå, Sweden.

[9]Applied Statistics Group, Department of Computational and Systems Biology, Rothamsted

Research, Hertfordshire, UK.

[10]Earlham Institute, Norwich Research Park, Norfolk, UK.

<u>Introduction</u>

Antimicrobial resistance (AMR) is a global health issue, with forecasts predicting ten million deaths a

year by 2050 if resistance levels continue to rise (O'Neill and The Review on Antimicrobial Resistance,

2016). The struggle against AMR has conventionally been viewed as a medical problem (Bengtsson-

Palme et al, 2018). However, the environment is now being acknowledged as a substantial contributor

to the global spread of antibiotic resistance (Wellington et al, 2013 & Singer et al, 2016;2019). Rivers

and waterways are prominent examples of environmental settings that have been identified as a

major repository and route for dissemination of both low levels of antibiotics and antibiotic resistance

genes (ARGs) (Pärnänen et al, 2019; Amos et al, 2015).

A recent study quantified antibiotic concentrations in freshwater samples taken at a range of sites

located on the Thames river catchment; the highest levels of antibiotics (as well as many other drugs)

were recorded downstream of waste-water treatment plants (WWTPs) (Holden et al, 2020). While the

origin and fate of these antibiotics could be strongly linked to community and clinical use, the impact

that they have on the emergence of antimicrobial resistance in the environment remains unclear

(Aslam et al, 2018). It has been proposed that ARGs have emerged within environmental bacteria to

provide a protective effect against these polluting drugs (Märtinez et al, 2012). This is of concern to

human health, as the genes under selection may be clinically relevant and have the potential to be

transferred between environmental and clinical bacteria via horizontal gene transfer (HGT) (Baquero

et al, 2008). However, it is also widely recognised that the evolutionary arms race, or red queen

hypothesis: the concept that species need to continually adapt, evolve, and proliferate to survive plays

an important role in shaping the environmental resistome (Papot et al, 2017). The fact that ARGs have

continually emerged within the riverine environment to protect against antibiotic producing

organisms in the riverine environment supports this latter concept (Singer et al, 2016). As a result, the

primary drivers of AMR within complex riverine bacterial communities remain difficult to define.

We have previously shown that the prevalence of an integron borne ARG, *sul1,* correlates with the

concentration of several antibiotics across river sites on the Thames catchment (Holden et al, 2020).

sul1 provides resistance to sulfonamides, a bacteriostatic class of antibiotics that were originally

synthesised in the 1930s. Sulfonamides bind to dihydropteroate synthase (DHPS) in bacteria and

hinder dihydrofolic acid formation which is vital for vitamin B formation and bacterial growth (Sköld,

2000). One sulfonamide; sulfamethoxazole (SMX) can be prescribed with trimethoprim (TMP) as co-

trimoxazole to treat chronic bronchitis, shigellosis, and *Pneumocystis carinii* pneumonitis (Alsaad et

al, 2013). SMX and TMP act in a synergistic manner to target successive steps of the folate synthesis

pathway, allowing co-trimoxazole to be effective against most aerobic gram-negative and gram-

positive organisms (Macejko and Schaeffer, 2008) (Figure 1).

59

Figure 1- A) Chemical structure of SMX/TMP

B) SMX and TMP combined mechanism of action

Despite the synergistic effects of SMX and TMP, recent human antibiotic prescription surveillance data

suggests that TMP is more commonly prescribed as a single-dose antibiotic for clinical purposes

(Ashiru-Oredope et al, 2014). Furthermore, the recent veterinary medicines directorates report on

antibiotics showed that a combined volume of 66.2mg/kg of sulfonamides/TMP was used to treat

infections in pigs in 2015, which accounted for the highest proportion of all antibiotics prescribed for

pigs (24%) (Bos et al, 2019). Studies have identified several different ARGs relevant to TMP (*DfrA1,*

DfrE, DfrB2/3, DfrB1/6) (Šeputienė et al, 2010) and sulfonamides (*sul1, sul2, sul3, sul4*) (Razavi et al,

2017). Due to the prominence of sulfonamide resistance in the environment, the *sul1* gene has

become incorporated into the structure of many class 1 integrons (CL1s) (Na et al, 2014). Several

studies have noted the persistence of sulfonamide and TMP resistance genes in the same

environmental settings, often located on the same plasmids (Phuong Hoa et al, 2008; Kiiru et al, 2013;

Muziasari et al, 2014).

During our previous work, sites on the Thames catchment with elevated TMP levels were identified

(Holden et al, 2020). Despite this, there is still no evidence for a causative relationship between the

residue of antibiotics such as TMP and the prevalence of ARGs (Holden et al, 2020). It is possible that

the correspondence between ARGs and drugs downstream of WWTPs is purely correlative, as both

are being released from the treatment process at the same time. In fact, studies have suggested that

WWTPs themselves, as opposed to river sites are 'hotspots' of ARGs and therefore most of the

selection for resistance occurs before antibiotic residue enters the riverine environment (Lood et al,

2017; Pazda et al, 2019; Fouz et al, 2020).

The aim of the current study was to analyse a combination of large metagenomic, geospatial and

chemical datasets, to compare twenty sites across the Thames river catchment. Specifically, we aimed

to look at sites with the highest and lowest TMP levels and establish the role that factors such as

proximity to WWTPs, population equivalence (PE) and land coverage play in the accumulation of TMP

at river sites. A secondary aim was to investigate seasonal variation in the levels TMP and SMX at river

sites. Lastly, as TMP can be prescribed alongside SMX as co-trimoxazole, we aimed to establish if the

levels of TMP and SMX identified at these specific sites mirrored the prevalence of ARGs relevant to

TMP (*Dfr* genes) and SMX (*sul* genes).

Methods

A total of 69 sampling locations were chosen within the Thames catchment. However, metagenomic

data was only generated for twenty of these 69 sites, and these twenty sites were analysed during

this study. All locations provide a depth of understanding regarding long-term water quality

parameters. A subset of these locations also contributed to a previous effort on modelling AMR within

the Thames Catchment (Amos et al 2015). The remaining sampling locations were selected to

maximise the diversity of river environments, river inputs, and adjacent land uses. Specific sampling

locations within a river catchment were selected to capture upstream and downstream positions

from a known river discharge or confluence. All locations also had to consider site accessibility as a

factor. Key discharges and confluences which structured sampling location choice were wastewater

treatment plants, agricultural fields (e.g., animal grazing), urban runoff, canals, fish farms, and other

rivers (see Table 1 Figure 1).

In the specific case of WWTPs, small WWTP were plants serving a population less than 10,000 and

any plant serving over 10,000 was considered large. WWTP impact was determined based on size of

the nearest WWTP and distance from the plant. Impact was determined based on distance away

from the WWTP.

Table 1 Summary of the twenty sampling sites analysed in this study

Name of Site	Thames Tributary	Nearest upstream Thames WWTP and treatment type (Secondary (S) or Tertiary (T))	XY coordinates	Postcode
1. Coln1	River Coln	Withington (S)	X:410239 Y:207637	GL7 5NY
2. Coln2	River Coln	Withington (S)	X:411501 Y:206777	GL7 5NW

3. Coln3	River Coln	Bibury (S)	X:414943 Y:201324	GL7 4AF
4. Coln4	River Coln	Fairford (S)	X:416285 Y:200409	GL7 4LX
5. Cut1	River Cut	N/A	X:488447 Y:171940	BG42 6JQ
6. Cut2	River Cut	Ascot (S)	X:485223 Y:171833	RG42 5NR
7. Cut4	River Cut	Bracknell (T)	X:485544 Y:173282	RG42
8. Cut6	River Cut	Maidenhead (T)	X:490245 Y:179218	SL6 2AR
9. Leach1	River Leach	Northleach (S)	X:420266 Y:204232	GL7 3NS
10. Leach2	River Leach	Lechlade (S)	X:422552 Y:199090	GL7 3HA
11. Ray1	River Ray (Thames headwater)	N/A	X:410831 Y:187351	N/A
12. Ray3	River Ray	N/A	X:412867 Y:185375	SN5 7UJ
13. Ray4	River Ray	Swindon (T)	X:412403 Y:186741	SN2
14. Ray5	River Ray	N/A	X:412310 Y:187249	SN2
15. Ray6	River Ray	Swindon (T)	X:411910 Y:192617	SN6 6
16. RiverThames1	River Thames	Castle Eaton (S)	X:417451 Y:196098	SN6 7RX
17. RiverThames3	River Thames	Witney (T)	X:444319 Y:208614	OX29 4BT
18. Thame1	River Thame	Rowsham (S)	X:479984 Y:215448	HP19 9UA
19. Thame2	River Thame	Aylesbury (T)	X:477886 Y:213623	N/A
20. Thame7	River Thame	Little Milton (S)	X:460064 Y:197680	OX10 7AW

Sampling regime

Samples were collected on two- or three-day sampling campaigns in August 2015, early March 2016

and August 2016. Sampling was carried out using purpose-made expandable sampling poles to allow

access across the riverbed. The poles were fitted with a machined holder that secured a 50ml falcon

tube. Sediment samples were collected by extending the pole into the river and scraping the upper

sediment layer into the tube until it was largely full. Sediment tubes were stored on ice. Samples were

collected at each site in triplicate. Water samples were collected in 500 ml screw cap bottles and

stored on ice. Samples for nutrient analysis was prepared as previously described (Amos et al, 2015).

Sample Processing

DNA extraction was carried out using the FastDNA™ spin kit for soil following manufacturers protocol,

0.5 g of sediment (wet weight) was used. DNA was pooled after extraction in 75μl to give total DNA

in final volume of 150 μl.

Metagenome

PCR free shotgun libraries were prepared for each sample from 2.5 ug metagenomic DNAs by Exeter

Sequencing Service for the metagenomic sequencing using the TruSeq DNA library Prep Kit

(Illumina). The libraries were sequenced to approximately 7GBp using the HiSeq 2500 rapid run

mode (2X 250 bp paired end) by the Exeter Sequencing Service (http://sequencing.exeter.ac.uk/). A

repeat sequencing effort was done on six DNA samples to determine if the coverage was sufficient

and each library was sequenced to provide 35 GBp per sample and the total was assembled as

below.

Metagenome assembly, annotation and coverage

Metagenome assembly was performed using megahit v1.1.3 with option -k--min 29 (Dinghua et al,

2015). Three assembly schemes were implemented, assemblies per sample, co-assemblies of samples

from the same tributary and a unique co-assembly built from all samples. Comparison of the 3

schemes lead to the global co-assembly being chosen for the rest of the analysis. Contigs smaller than

500 nucleotides were discarded. Gene prediction was carried out using prodigal v2.6.3, with option -

p meta (Hyatt et al, 2015). A selection of 36 Single copy Core Genes (SCG) were annotated through

rpsblast v2.9.0 using pssm formatted COG database made available by the CDD (Tatusoy et al 2000;

Lu et al, 2020). Best hits were filtered so that hit span at least 50% of the query with an e-value smaller

than 1e-5. Additionally, ARG and *intI1* annotation was completed using diamond v0.9.9 (Buchfink et

al, 2015). Databases for both were sourced from (Lee et al, 2020), clustered respectively from of CARD

(Jia et al, 2017) and NCBI sequence search. Best hits were filtered using 80% query coverage, 80%

percentage identity and 20% subject coverage. Taxonomic annotation was carried out using diamond

blastp on a NR database reduced to prokaryotes (Pruitt et al, 2005).

Assembly was converted to contig graph using megahit toolkit, it was then translated to orf graph

using result from gene prediction in custom python script. Visualisation of annotations on orf graph is

done through bandage v0.8.1 (Wick et al, 2015).

Samples were mapped to contigs using bwa-mem v0.7.17-r1188 (Li, 2013). Contig and genes mean

coverage were computed through both samtools v1.5 (Li et al, 2009) and bedtools v2.28.0 (Quinlan

et al, 2013). Coverage of functional feature, such as SCG, AMR or *intI1* were obtained by summation

of coverage of genes sharing the same annotation. All coverages are normalised by sample, using

the median of the 36 SCG coverages.

Chemical Analysis

For each site 100 mL of each water sample was acidified to pH 3, internal standards were added and

the sample was extracted with a solid phase extraction (SPE) vacuum manifold system using Oasis HLB

(200mg) cartridges. Thereafter each sample was evaporated to <10 μL, reconstituted to a final volume

of 0.1 mL 5% acetonitrile with 0.1% formic acid and 10 μL of each water sample was analysed with

liquid chromatography-tandem mass spectrometry (LC-MS/MS). A detailed description of the two

analytical methods used to analyse most of the pharmaceuticals, and the validation of each method

is shown in (Holden et al, 2020). However, at the time of sample analysis, contamination in the LC-

MS/MS system interfered with the analysis of the six biocides included in the study. Instead, an

analytical method developed by (Östman et al, 2017) specifically for biocides, on the LC-MS/MS

system, was used to analyse the six biocides.

Geospatial analyses and meteorological data

All geospatial analyses were performed using ArcGIS v10.4 software (ESRI, Aylesbury, UK). To analyse

possible drivers of AMR for each sampling site, an upstream catchment of that location was delineated

using the hydrology tools to create a flow accumulation and flow direction layer per site, and

subsequently a watershed (upstream catchment), using a 50m grid interval hydrological digital terrain

model (the Centre for Ecology and Hydrology Integrated Hydrological Digital Terrain Model (IHDTM))

(Morris & Flavin, 1994). A Network Dataset was created from the Thames river data, using the

Ordnance Survey MasterMap Water Network 2016 (Ordnance Survey, 2016) and comprised all layers

needed for distance measurements, including WWTPs. Data on WWTPs was obtained from the

environment agency (Environment Agency, 2012; Environment Agency, 2013). To analyse upstream

land use, and more specifically to calculate the percentage land cover of each sample site watershed,

the Centre for Ecology and Hydrology Land Cover Map 2015 (Rowland et al, 2017) and Land Cover

Plus: Crops Map 2016 were used (NERC CEH, 2016).

Land-cover profiles were calculated for each sampling site based on the identified watershed,

comprising the percentage cover for 13 broad habitat classes (Arable horticulture, Acid Grassland,

Broadleaf woodland, Calcareous grassland, Coniferous woodland, Freshwater, Heather, Heather

grassland, Improved grassland, Inland rock, Neutral grassland, Suburban, Urban). The "Arable

horticulture" class was further sub-divided into 12 primary crop categories, and three further major

land-cover classes were also calculated by summing over broad habitats (All Grassland (sum of 5 broad

habitats), All Woodland (sum of 2 broad habitats), Urban-Suburban (sum of 2 broad habitats)).

For each WWTP in each watershed, data on the Population Equivalent (overall sewage load), resident

and non-resident populations and industry load (all in terms of the people producing the equivalent

sewage load assuming 200l of sewage flow containing 60g of BOD per person per day) were tabulated

together with the river distance (m) from the WWTP to the sampling site, the dry weather flow at the

WWTP (m^3 per day) and the general classification of sewage treatment plants (WWTP Type: SA -

Secondary Activated Sludge; SB - secondary biological filter; TA - Activated Sludge with tertiary

treatment; TB - Biological filter with tertiary treatment). Simple summaries of these data were

calculated for each watershed (sampling site), including total numbers of WWTPs and number for

each Type, mean, minimum and maximum distance, and mean and maximum PE. Based on the

modelling approach developed in (Amos et al, 2015), summaries of the potential aggregated impact

of upstream WWTPs were also calculated – these weighted the impact of the load (PE) from each

WWTP by the inverse of the distance raised to a power (0, 0.35 (close to the value estimated in the

model fitting in (Amos et al, 2015), 0.5 or 1), allowing for different decay functions (with a power of

zero indicating no impact of distance). Where (Amos et al, 2015) only considered WWTPs within a

10km radius of the sampling site, summaries of the numbers of WWTPs and the inverse distance-

weighted aggregated PE were calculated for WWTPs within 10km, 20km, 30km, 40km and 50km

upstream as well as for the whole upstream watershed. Data were also tabulated for each sampling

site (watershed) for the number and density of septic tanks, and for the number and mean upstream

distance for fish farms.

Statistical analysis and model development

Initial analysis of chemical data used a mixed-effects model with Tukey's multiple comparison tests

to assess differences in trimethoprim (TMP) and sulfamethoxazole (SMX) concentrations between

seasons (summer & winter) and between twenty sample sites (Coln1, Coln2, Coln3, Coln4, Leach1,

Leach2, Key1, Ray1, Ray6, Ray4, Ray3, Thames1, Thames3, Thame7, Thame2, Cut1, Cut2, Cut4, Cut6,

Thames6).

Analysis of geospatial data relating to the distance (metres) of the twenty sample sites downstream

from waste-water treatment plants (WWTPs) and the concentration of TMP was carried out using

Spearman's rank correlation coefficient due to the non-parametric nature of the data. Both variables

were log-transformed prior to analysis. Analysis of the difference in population equivalence (PE)

between sites with the highest and lowest TMP levels was carried out using an unpaired t-test with

Welch's correction.

Analysis of metagenomic data used multiple t-tests with Bonferri-Dunn correction to assess

differences in ARG (*sul1*, *sul2*, *dfrB1/6*, *dfrB2/3*, *dfrE*) and *intI1* prevalence between seasons (summer

and winter). Two-way ANOVA was also applied to assess differences in the prevalence of

aforementioned ARGs & *intI1* between seasons and sites with the four highest (Cut4, Ray4, Cut6,

Ray6) and four lowest (Coln1, Coln2, Coln3, Leach1) TMP concentrations.

All analysis was performed using GraphPad® Prism version 8 and R® software. For all tests, p values

of statistical significance are signified by * = $p<0.05$, ** = $p<0.005$, *** = $p<0.0005$ & **** $p<0.0001$.

Maps

Maps shown in Figure 4 were created using ArcGIS v10.4 software. Geographic coordinates of sites

were taken from (Ordnance Survey, 2016).

Results

Analysis of TMP and SMX levels

The levels of TMP and SMX identified in the water at the twenty different sites across Thames

catchment differed greatly. Furthermore, there was also a significant seasonal variation in the levels

of TMP and SMX identified at individual sites (Figure 2).

Figure 2- The concentrations of TMP and SMX at twenty sites across the Thames catchment in summer

(n=6) and winter (n=3) (SEM). The sites were ordered by geographical location across the catchment,

with Coln1 being furthest East and Thames6 being furthest West.

TMP concentrations were higher than SMX concentrations at all sites other than Thame7. Ten of the

twenty sites had significantly higher TMP in winter compared to summer (mixed effects model). Four

sites had levels of TMP exceeding 750ng/L (Ray4, Ray6, Cut4 and Cut6), with Ray4 providing a TMP

concentration of 3000ng/L. Aside from Cut4, the level of TMP was significantly higher in winter

compared to summer for these sites (mixed effects model).

For three sites (Ray4, Thame7 and Thame2), the concentration of SMX was higher in summer

compared to winter, but these differences were not significant (mixed effects model).

Figure 3- Relationship between distance of sample sites downstream from a WWTP and the

concentration of TMP across 20 sites on the Thames catchment.

A significant negative correlation was observed between the distance of sites downstream from the

nearest WWTP and the TMP concentration across 20 sites on the Thames catchment (Spearman r = -

0.456 & p<0.05). This relationship was further explored to determine where the highest and lowest

TMP levels were detected (Table 1).

Table 1- A sub-set of four sites that showed the highest and lowest TMP concentrations from the 20

sites. Of this sub-set, only one of the sites showed a recordable concentration of SMX (Ray4), so this

data was not shown in the table. TMP concentrations shown were averaged across summer/winter

seasons.

*Secondary: 'Treatment methods include biological filtration (including conventional filtration,

rotating biological contactors and root zone treatment (where used as a secondary treatment stage),

alternating double filtration and high rate filtration'.

(Sewage Treatment in the UK, 2020).

*Tertiary: 'Works with a secondary activated sludge process whose treatment methods also include

rapid-gravity sand filters, moving bed filters, pressure filters, nutrient control using physico-chemical

and biological methods, disinfection, hard COD and colour removal, where used as a tertiary

treatment stage' (Sewage Treatment in the UK, 2020).

River Site	Nearest upstream WWTP	Distance from WWTP (m)	WWTP Treatment Type*	Human population served	Trimethoprim concentration (ng/L)
Cut4	Bracknell	1361	Tertiary	90,000	1671
Cut6	Maidenhead	1540	Tertiary	87,400	934
Ray4	Swindon	1213	Tertiary	210,000	2037
Ray6	Swindon	10208	Tertiary	210,000	1384
Coln1	Withington	16677	Secondary	230	14
Coln2	Withington	18688	Secondary	230	4
Coln3	Bibury	9479	Secondary	600	5
Leach1	Northleach	20270	Secondary	1,670	4

Three of the four sites with the highest TMP concentrations were located within 2000 metres of

WWTPs (Table 1) as expected from the data in Figure 4. The four sites with the highest TMP

concentrations were located closest to tertiary WWTPs, whereas the all of the four sites with the

lowest TMP concentrations were located closest to secondary WWTPs. Two of the four sites with the

highest TMP concentrations (Ray4 and Ray6) were located closest to the same Swindon WWTP.

Analysis of land use and population equivalence (PE)

Land use and PE data were assembled for the four sites with the highest and lowest TMP

concentrations across the Thames catchment, as shown in Figures 4, 5 and 6.

Figure 4

A) Locations of the Rivers Cut, Ray, Coln and Leach on the Thames catchment.

B) Locations and land-use corresponding to the four sites with the lowest TMP concentrations.

C) Locations and land-use corresponding to the four sites with the highest TMP concentrations.

Examination of the land-use data in proximity to sites with highest and lowest TMP levels indicate

urbanization correlating with tertiary WWTPs and high TMP levels (Figure 4). The data also shows that

the proportion of arable and pasture-based land coverage was higher within the catchment of sites

furthest from WWTPs.

Figure 5- The average $\log_{10}$ population equivalence (PE) for four sites with the highest TMP concentrations (left) and four sites with the lowest TMP concentrations (right) (n=4) (SEM).

The average PE for the four sites with the highest TMP concentrations was significantly higher than the PE for the four sites with the lowest TMP concentrations (unpaired t-test).

Figure 6- Percentage of land coverage within the catchment of four sites with the highest TMP levels

(left) and the lowest TMP levels (right).

The proportion of urban/suburban land coverage was found to be higher in the catchment of the four

sites with the highest TMP concentrations compared to the four with the lowest TMP concentrations.

By contrast, the proportion of arable land coverage was higher across the catchment of sites with the

lowest TMP concentrations compared to sites with the highest TMP concentrations.

Prevalence of ARGs

Only two SMX ARGs and three TMP ARGs were found at detectable levels in the metagenomes derived

from river sediment (Figure 7) and of these *sul1* was detected at significantly higher levels in the

winter. These were compared with *intI1* (CL1 marker gene) prevalence (Figure 7).

Figure 7- Prevalence of ARGs relevant to TMP/SMX resistance across 20 sites on the Thames

catchment in summer (n=6) and winter (n=3) (SEM).

The *sul1* gene showed the highest overall prevalence compared to the four other ARGs relevant to

TMP and SMX and *intI1*. *sul1* was also the only gene to show significant variation between the summer

and winter seasons (multiple t-tests).

The two ARGs relevant to SMX and the three ARGs relevant to TMP were then analysed in comparison

with the levels of TMP antibiotic identified at the sites with the four highest and four lowest TMP

levels (same sub-set as Table 1). These were compared with *intI1* (CL1 marker gene) prevalence.

Figure 8- Prevalence of ARGs relevant to TMP/SMX and *intI1* at sites with the four highest and four

lowest TMP levels in summer (n=6) (above) compared to winter (n=3) (SEM).

The prevalence of *sul1* was higher at sites Ray4, Ray6 and Cut4 in winter compared to summer. At

these four sites, the prevalence of *sul1* was higher than any of the prevalence of any other genes in

both winter and summer. Coln2 and Coln3 showed significantly higher prevalence's of *dfrB1/6* in

summer compared to winter (two-way ANOVA). The prevalence of *sul2* at Coln2 was significantly

higher in summer compared to winter (two-way ANOVA).

Discussion

Over the last two decades, the occurrence of antibiotics in water bodies and the potential for the

development of antimicrobial resistance (AMR) as a result has become an issue of concern (Khan et

al, 2013). Particularly in urban areas, waste-water treatment plants (WWTPs) are among the main

recipients of antibiotics, some of which will persist after treatment and can be released into nearby

rivers (Rodriguez-Mozaz et al, 2020).

The levels of two clinically relevant antibiotics (trimethoprim (TMP) and sulfamethoxazole (SMX))

were analysed at river sites during this study. Results showed that nineteen of the twenty river sites

analysed showed higher levels of TMP compared to SMX. TMP and SMX can be co-prescribed as co-

trimoxazole, which might suggest that their levels identified in rivers might be similar. However,

clinical antibiotic surveillance data shows that TMP is more commonly prescribed as a single dose

antibiotic (Ashiru-Oredope et al, 2014). By contrast, veterinary surveillance data suggests that TMP is

co-prescribed with 'sulfonamides' in high levels in agricultural settings (Bos et al, 2019). As a result, it

is possible that TMP levels were higher than SMX levels at these sites because the two drugs are more

often being used for different purposes and therefore more likely to be found at different river sites.

Another possible reason why TMP was significantly higher than SMX at certain sites could be the

differing biodegradation profiles that the drugs display in the environment (Thiebault, 2020). One

study used batch reactors containing wastewater matrices to show that the biodegradation rate of

SMX was significantly higher than TMP over a 21-day period (Pérez et al, 2005). Most WWTPs are

designed to extract nutrients and easily degradable carbon compounds. However, it has been shown

that they have limited capacities to remove pharmaceuticals such as TMP, making them sources of

TMP in rivers and streams (Metcalfe et al, 2010; Sengupta et al, 2014). The presence of antibiotics in

the treatment process can inhibit the biological activity of activated sludge, meaning that the

antibiotics themselves are not degraded effectively (Al-Riyami et al, 2018). As a result, it is possible

that SMX is more readily degraded than TMP as part of the WWTP process and therefore was found

in higher concentrations at certain sites.

As a result of the incapacity of conventional treatment systems to remove emerging contaminants

such as antibiotics during treatment, studies have indicated that urban-associated river sites often

have the highest concentrations of antibiotics (Chen et al, 2018; Lundborg et al, 2017). This would

explain why TMP and SMX were both detected at river sites in this study, particularly as those sites

with the highest TMP levels all had a high level of urban/suburban land coverage associated with

them. The significant negative correlation between levels of TMP and the distance of a site from a

WWTP further suggests that the TMP being identified in rivers coming from WWTPs. Furthermore, in

areas of high human population, it has been shown that excretion by humans contributes to most of

the pharmaceutical waste that ends up in domestic sewers and is eventually transported to WWTPs

(Tehrani and Gilbrade, 2018). This would explain the fact that the PE was significantly higher at the

four sites with the highest TMP concentrations compared to those with the lowest TMP

concentrations.

During this study, four river sites (Cut4, Cut6, Ray4 and Ray6) were shown to have elevated levels of

TMP. Despite showing significantly higher levels of TMP compared to all other sites, they did not show

a significantly higher prevalence of the ARGs that are known to confer resistance to TMP (*Dfr* genes).

DfrB1/6 for example, was only detected at sites with the lowest TMP levels. This suggests that the

concentrations of TMP at river sites are not directly selecting for resistance, which contrasts with

previous reports which have shown that sub-Minimum inhibitory concentrations (MIC) of

antimicrobials can select for resistance in environmental settings (Ter Kuile et al, 2016; Winstrand-

Yuen et al, 2018; Opatowski et al, 2010). Interestingly, sites with the highest levels of the SMX

associated ARG *sul1* did not show high levels of SMX but did show high levels of TMP. One possible

explanation for why this happens is that TMP is indirectly selecting for *sul1* by selecting for the

presence of class 1 integrons (CL1s), as CL1s are known to carry the *sul1* gene. CL1s which have been

shown to confer resistance to several antibiotics, including SMX, TMP, amoxicillin (AMX), tetracycline

(TCN) and erythromycin (ERY) (Langata et al, 2019; Ndi and Barton, 2011; Callejas et al, 2019). Our

previous data showed no significant correlations between TCN and ERY concentrations with *sul1*

prevalence, but we do not have chemical data available relating to AMX (Holden et al, 2020). It is also

possible that concentrations of AMX are present at these river sites and they are also able to select

for CL1s and therefore *sul1*. This effect has been noted in a previous study (Ironmonger et al, 2018).

However, based solely on the data shown in this study, given that the prevalence of CL1s did not differ

significantly between river sites or seasons like *sul1* did, it seems unlikely that this is the case. Another

possible explanation is that both TMP and *sul1* are released to rivers following the waste-water

treatment process at the same time, and therefore were both detected by coincidence. This would

mean that the relationship between TMP and *sul1* noted in this paper and our previous study is purely

correlative (Holden et al, 2020).

Data from this study showed seasonal variation in the prevalence of individual ARGs and the

concentrations of TMP/SMX antibiotics at river sites. For example, the prevalence of *sul1* over the 20

sites was significantly higher in winter compared to summer. Also, the concentration of TMP was

significantly higher in winter compared to summer for ten of the twenty sites analysed. Increased

prescription/use of TMP in the population during the winter months could increase the concentrations

detected at river sites and potentially the prevalence of ARGs. A recent study indicated that

prescription rates for antibiotics (including TMP) by General Practitioners (GPs) in the West Midlands

UK increase consistently in winter compared to the three other seasons (Ironmonger et al, 2018). An

increase in antibiotic consumption can increase the antibiotic concentration found in sewage, which

will eventually be treated by WWTPs (Pärnänen et al, 2019). If antibiotics are not successfully removed

during sewage treatment processes, then the higher concentrations of TMP shown in winter

compared to summer could be due to increased antibiotic usage/consumption. However, with regards

the chemical analysis of TMP and SMX, it must be acknowledged that the freshwater samples used

were collected over two summer seasons (summer 2015 and summer 2016) and only one winter

season (winter 2016). Also, the concentrations of chemicals determined at each site are only reflective

of the concentration of that chemical at one particular time, with no corrections made for the flow

rate of the individual river site or data on sewage spills from WWTPs. As a result, the chemical data

shown is potentially not reflective of the whole seasons analysed and the differences between TMP

concentrations in summer and winter is potentially more of a temporal effect as opposed to a seasonal

one.

One site in this study; Coln2 showed a showed significantly increased prevalence of *sul2* in summer

despite having minimal concentrations of TMP and SMX, which contrasted with its low *sul1* prevalence.

Sul2 is a sulfonamide resistant dihydropteroate synthase of gram-negative bacteria, generally found on

small plasmids, whereas *sul1* is generally found linked to other resistance genes on CL1s (Antunes et al,

2005; Jia et al, 2017). According to the National Office of Animal Health (NOAH), there are two different

sulfonamide classes in addition to SMX that can be prescribed in combination with TMP to treat

infections in farm animals: sulfadiazine (SDZ) and sulfadoxine (SX) (NOAH, 2020). TMP-SDZ is a short-

acting bacteriostatic antibiotic that is typically applied to treat infections in horses (Ensink et al, 2003).

TMP-SX is a bacteriostatic antibiotic typically used to treat respiratory tract infections in cattle (Dunkley,

1994). Previous studies have noted increased abundance of the *sul2* in agricultural samples containing

SDZ (Jechalke et al, 2012; Kopmann et al, 2012). As a result, is possible that the increased prevalence of

sul2 shown at Coln2 in spite of the low SMX and TMP concentrations is driven by an agriculturally relevant

sulfonamide such as SDZ, which was not targeted by the chemical analysis of this study. This is further

emphasised by the fact that there was no relationship between *sul2* and *intI1* prevalence in this study,

as this shows that *sul2* genes are not emerging on CL1s alongside clinically relevant ARGs. However, it

must be acknowledged that the available UK antibiotic surveillance data for veterinary applications only

shows data for 'sulfonamides' as a general class, as opposed to distinguishing between individual drugs

such as SMX and SDZ (Bos et al, 2019). As a result, it is difficult to determine the actual levels of SDZ that

are being applied in agriculture. Also, despite the isolated high levels of *sul2* at Coln2, this study showed

that across all 20 sites on the Thames catchment, *sul1* was significantly more prevalent than *sul2.*

Conclusions

This study involved comparative analysis of large metagenomic, chemical and geospatial datasets to

investigate the impact of antimicrobials such as trimethoprim (TMP) on the profile of antimicrobial

resistance and the wider river microbiome. The following conclusions were drawn:

1) Nineteen of the twenty sites analysed showed higher levels of trimethoprim (TMP) than

sulfamethoxazole (SMX), except for Thame7.

2) There was a significant negative correlation between the distance of river sites from waste-water

treatment plants (WWTP) and the levels of TMP detected.

3) Of the twenty river sites analysed, four showed particularly high levels of TMP. For three of these

four sites, TMP levels were significantly higher in winter compared to summer.

4) *DfrE*, *DfrB1/6*, *DfrB2/3 and intI1* prevalence at the four sites with the highest TMP levels were not

significantly higher than those with the lowest TMP levels. *DfrE*, *DfrB1/6*, *DfrB2/3* was significantly

higher in summer compared to winter. *intI1* prevalence showed no significant seasonal variation.

5) The four sites with high TMP levels showed significantly higher PE (population equivalence) levels

than the four sites with the lowest TMP levels.

6) The land coverage of the four sites with the highest TMP levels were significantly more

urban/suburban than the four sites with the lowest TMP levels. By contrast, the land coverage of

the four sites with the lowest TMP levels was significantly more arable than those with the highest

TMP levels.

558
7) *sul1* prevalence at the four sites with the highest TMP levels were significantly higher than those

with the lowest TMP levels. *sul1* prevalence was also significantly higher in winter compared to

summer.

8) *sul2* prevalence at one of the sites with the lowest TMP levels (Coln2) was significantly higher than

those with the highest TMP levels. *sul2* prevalence at the Coln2 was also significantly higher in

summer compared to winter.

Future Work

Our previous work has shown evidence of a significant positive correlation between TMP levels and

the prevalence of *sul1* at sites across the Thames catchment (Holden et al, 2020). Data from this study

pinpointed the sites with the highest levels of TMP and the highest prevalence of *sul1.*

We have also shown that sites with the highest TMP concentrations and highest *sul1* prevalence are

generally located closer to WWTPs than those with low TMP concentrations. Furthermore, sites with

the highest TMP concentrations have a higher PE, are more urban/suburban and that their TMP

concentration/*sul1* prevalence is higher in winter compared to summer. Despite this, there is still no

previously explained mechanism as to why levels of TMP would select for *sul1* directly.

definitive evidence of a causative impact of TMP concentrations on the prevalence of ARGs such as

sul1. Future work will aim to look at whether or not the concentrations of TMP and SMX recorded in

this study can have a significant causative impact on the prevalence of relevant ARGs.

References

1.   Adelowo, O.O., Caucci, S., Banjo, O.A., Nnanna, O.C., Awotipe, E.O., Peters, F.B., Fagade,

O.E., and Berendonk, T.U. (2018) Extended Spectrum Beta-Lactamase (ESBL)-producing bacteria

isolated from hospital wastewaters, rivers and aquaculture sources in Nigeria. *Environ Sci Pollut Res*

Int **25**: 2744-2755.

2.   Alsaad, N., van Altena, R., Pranger, A.D., van Soolingen, D., de Lange, W.C., van der Werf,

T.S., Kosterink, J.G., and Alffenaar, J.W. (2013) Evaluation of co-trimoxazole in the treatment of

multidrug-resistant tuberculosis. *Eur Respir J* **42**: 504-512.

3.   Amos, G.C., Gozzard, E., Carter, C.E., Mead, A., Bowes, M.J., Hawkey, P.M., Zhang, L., Singer,

A.C., Gaze, W.H., and Wellington, E.M. (2015) Validated predictive modelling of the environmental

resistome. *ISME J* **9**: 1467-1476.

4.   Amos, G.C.A., Ploumakis, S., Zhang, L., Hawkey, P.M., Gaze, W.H., and Wellington, E.M.H.

(2018) The widespread dissemination of integrons throughout bacterial communities in a riverine

system. *ISME J* **12**: 681-691.

5.   Ashiru-Oredope, D., Hopkins, S., and Group, E.S.P.f.A.U.a.R.O. (2013) Antimicrobial

stewardship: English Surveillance Programme for Antimicrobial Utilization and Resistance (ESPAUR).

J Antimicrob Chemother **68**: 2421-2423.

6.   Aslam, B., Wang, W., Arshad, M.I., Khurshid, M., Muzammil, S., Rasool, M.H., Nisar, M.A.,

Alvi, R.F., Aslam, M.A., Qamar, M.U., Salamat, M.K.F., and Baloch, Z. (2018) Antibiotic resistance: a

rundown of a global crisis. *Infect Drug Resist* **11**: 1645-1658.

7.   Baharoglu, Z., and Mazel, D. (2014) SOS, the formidable strategy of bacteria against

aggressions. *FEMS Microbiol Rev* **38**: 1126-1145.

8.   Baquero, F., Martínez, J.L., and Cantón, R. (2008) Antibiotics and antibiotic resistance in

water environments. *Current Opinions in Biotechnology* **19**: 260-265

9.   Bengtsson-Palme, J., Kristiansson, E., and Larsson, D.G.J. (2018) Environmental factors

influencing the development and spread of antibiotic resistance. *FEMS Microbiol Rev* **42**.

10.   CEH(2016) https://www.ceh.ac.uk/sites/default/files/LCM2015_Dataset_Documentation.pdf

Accessed April 28, 2020.

620
621

11.   Chen, Y., Xi, X., Xu, J., Xie, R., and Jiang, J. (2018) Distribution patterns of antibiotic residues

in an urban river catchment. *Water and Environment Journal* **33**: 31-39.

12.   Creating a River Thames fit for our future: an updated strategic and economic case for the

Thames Tideway Tunnel Department for Environment, Food and Rural Affairs

(DEFRA)2015 https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attach

ment_data/file/471847/thames-tideway-tunnel-strategic-economic-case.pdf

13.   Dan A, Yang, Y., Dai, Y.N., Chen, C.X., Wang, S.Y., and Tao, R. (2013) Removal and factors

influencing removal of sulfonamides and trimethoprim from domestic sewage in constructed

wetlands. *Bioresour Technol* **146**: 363-370.

14.   Davies, J., and Davies, D. (2010) Origins and evolution of antibiotic resistance. *Microbiol Mol*

Biol Rev **74**: 417-433.

15.   Di Cesare, A., Eckert, E.M., Rogora, M., and Corno, G. (2017) Rainfall increases the

abundance of antibiotic resistance genes within a riverine microbial community. *Environ Pollut* **226**:

473-478.

16.   Fouz, N., Pangesti, K.N.A., Yasir, M., Al-Malki, A.L., Azhar, E.I., Hill-Cawthorne, G.A., and Abd

El Ghany, M. (2020) The Contribution of Wastewater to the Transmission of Antimicrobial

Resistance in the Environment: Implications of Mass Gathering Settings. *Trop Med Infect Dis* **5**.

17.   Frank, T., Gautier, V., Talarmin, A., Bercion, R., and Arlet, G. (2007) Characterization of

sulphonamide resistance genes and class 1 integron gene cassettes in Enterobacteriaceae, Central

African Republic (CAR). *J Antimicrob Chemother* **59**: 742-745.

18.   Hyatt, D., Chen, G.L., Locascio, P.F., Land, M.L., Larimer, F.W., and Hauser, L.J. (2010)

Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC*

Bioinformatics **11**: 119.

19.   Händel, N., Schuurmans, J.M., Brul, S., and ter Kuile, B.H. (2013) Compensation of the

metabolic costs of antibiotic resistance by physiological adaptation in Escherichia coli. *Antimicrob*

Agents Chemother **57**: 3752-3762.

20.   Händel, N., Schuurmans, J.M., Feng, Y., Brul, S., and ter Kuile, B.H. (2014) Interaction

between mutations and regulation of gene expression during development of de novo antibiotic

resistance. *Antimicrob Agents Chemother* **58**: 4371-4379.

21.   Holden, J., Raguideau, S., Zhang, L., Tipper, H.J, Delaney, J., Hill, G.L et al. (2020) How sewage

impacts antibiotic levels and bacterial resistance in the Thames catchment. *To be released*

22.   Ironmonger, D., Edeghere, O., Verlander, N.Q., Gossain, S., Hopkins, S., Hilton, B., and

Hawkey, P.M. (2018) Effect of general practice characteristics and antibiotic prescribing on

Escherichia coli antibiotic non-susceptibility in the West Midlands region of England: a 4 year

ecological study. *J Antimicrob Chemother* **73**: 787-794.

23.   Jia, B., Raphenya, A.R., Alcock, B., Waglechner, N., Guo, P., Tsang, K.K., Lago, B.A., Dave,

B.M., Pereira, S., Sharma, A.N., Doshi, S., Courtot, M., Lo, R., Williams, L.E., Frye, J.G., Elsayegh, T.,

Sardar, D., Westman, E.L., Pawlowski, A.C., Johnson, T.A., Brinkman, F.S., Wright, G.D., and

McArthur, A.G. (2017) CARD 2017: expansion and model-centric curation of the comprehensive

antibiotic resistance database. *Nucleic Acids Res* **45**: D566-D573.

24.   Kemnic T.R. Coleman M. (2019) Trimethoprim Sulfamethoxazole In: StatPearls [Internet].

Treasure Island (FL): *StatPearls Publishing*; https://www.ncbi.nlm.nih.gov/books/NBK513232/.

25.   Khan, G.A., Berglund, B., Khan K.M., Lindgren P.E & Fick, J. (2013) Occurrence and abundance

of antibiotics and resistance genes in rivers, canal and near drug formulation facilities--a study in

Pakistan. *PLoS One* **28**;8(6):e62712.

26.   Kiiru, J., Butaye, P., Goddeeris, B.M., and Kariuki, S. (2013) Analysis for prevalence and

physical linkages amongst integrons, ISEcp1, ISCR1, Tn21 and Tn7 encountered in Escherichia coli

strains from hospitalized and non-hospitalized patients in Kenya during a 19-year period (1992-

2011). *BMC Microbiol* **13**: 109.

27.   Lee, K., Kim, D.W., Lee, D.H., Kim, Y.S., Bu, J.H., Cha, J.H., Thawng, C.N., Hwang, E.M., Seong,

H.J., Sul, W.J., Wellington, E.M.H., Quince, C., and Cha, C.J. (2020) Mobile resistome of human gut

and pathogen drives anthropogenic bloom of antibiotic resistance. *Microbiome* **8**: 2.

28.   Li, B., and Webster, T.J. (2018) Bacteria antibiotic resistance: New challenges and

opportunities for implant-associated orthopedic infections. *J Orthop Res* **36**: 22-32.

29.   Li, D., Liu, C.M., Luo, R., Sadakane, K., and Lam, T.W. (2015) MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics **31**: 1674-1676.

30.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and Subgroup, G.P.D.P. (2009) The Sequence Alignment/Map format and SAMtools.

Bioinformatics **25**: 2078-2079.

31.   Long, H., Miller, S.F., Strauss, C., Zhao, C., Cheng, L., Ye, Z., Griffin, K., Te, R., Lee, H., Chen,

C.C., and Lynch, M. (2016) Antibiotic treatment enhances the genome-wide mutation rate of target

cells. *Proc Natl Acad Sci U S A* **113**: E2498-2505.

32.   Lood, R., Ertürk, G., and Mattiasson, B. (2017) Revisiting Antibiotic Resistance Spreading in

Wastewater Treatment Plants - Bacteriophages as a Much-Neglected Potential Transmission

Vehicle. *Front Microbiol* **8**: 2298.

33.   Lu, S., Wang, J., Chitsaz, F., Derbyshire, M.K., Geer, R.C., Gonzales, N.R., Gwadz, M., Hurwitz,

D.I., Marchler, G.H., Song, J.S., Thanki, N., Yamashita, R.A., Yang, M., Zhang, D., Zheng, C., Lanczycki,

C.J., and Marchler-Bauer, A. (2020) CDD/SPARCLE: the conserved domain database in 2020. *Nucleic*

Acids Res **48**: D265-D268.

34.   Lundborg, C.S., and Tamhankar, A.J. (2017) Antibiotic residues in the environment of South

East Asia. *BMJ* **358**: j2440.

35.   Martínez, J.L. (2012) Natural antibiotic resistance and contamination by antibiotic resistance

determinants: the two ages in the evolution of resistance to antimicrobials. *Front Microbiol* **3**: 1.

36.   Masters, P.A., O'Bryan, T.A., Zurlo, J., Miller, D.Q., and Joshi, N. (2003) Trimethoprim-

sulfamethoxazole revisited. *Arch Intern Med* **163**: 402-410.

37.   McCarthy, M., Spillane, S., Walsh, S., and Kendon, M. (2016) The meteorology of the

exceptional winter of 2015/2016 across the UK and Ireland. *Weather* **71**: 305-313.

38.   Metcalfe, C.D., Chu, S., Judt, C., Li, H., Oakes, K.D., Servos, M.R., and Andrews, D.M. (2010)

Antidepressants and their metabolites in municipal wastewater, and downstream exposure in an

urban watershed. *Environ Toxicol Chem* **29**: 79-89.

39.   Morris, D.G., Flavin, R.W., (1994) A digital terrain model for hydrology. Proc 4th International

Symposium on Spatial Data Handling. Sub-set of UK 50 m by 50 m hydrological digital terrain model

grids. Wallingford, NERC, Institute of Hydrology **1**: 250–262.

40.   Munro, K., Martins, C.P.B., Loewenthal, M., Comber, S., Cowan, D.A., Pereira, L., and Barron,

L.P. (2019) Evaluation of combined sewer overflow impacts on short-term pharmaceutical and illicit

drug occurrence in a heavily urbanised tidal river catchment (London, UK). *Sci Total Environ* **657**:

1099-1111.

41.   Muziasari, W.I., Managaki, S., Pärnänen, K., Karkman, A., Lyra, C., Tamminen, M., Suzuki, S.,

and Virta, M. (2014) Sulphonamide and trimethoprim resistance genes persist in sediments at Baltic

Sea aquaculture farms but are not detected in the surrounding environment. *PLoS One* **9**: e92702.

42.   Na, G., Zhang, W., Zhou, S., Gao, H., Lu, Z., Wu, X., Li, R., Qiu, L., Cai, Y., and Yao, Z. (2014)

Sulfonamide antibiotics in the Northern Yellow Sea are related to resistant bacteria: implications for

antibiotic resistance genes. *Mar Pollut Bull* **84**: 70-75.

43.   Nagulapally, S.R., Ahmad, A., Henry, A., Marchin, G.L., Zurek, L., and Bhandari, A. (2009)

Occurrence of ciprofloxacin-, trimethoprim-sulfamethoxazole-, and vancomycin-resistant bacteria in

a municipal wastewater treatment plant. *Water Environ Res* **81**: 82-90.

44.   O'Neill J., The Review on Antimicrobial Resistance (2016). *Tackling Drug-Resistant*

Infections Globally: Final Report and Recommendations. Available at: http://amr-

review.org/sites/default/files/160518_Final%20paper_with%20cover.pdf

45.   Opatowski, L., Mandel, J., Varon, E., Boëlle, P.Y., Temime, L., and Guillemot, D. (2010)

Antibiotic dose impact on resistance selection in the community: a mathematical model of beta-

lactams and Streptococcus pneumoniae dynamics. *Antimicrob Agents Chemother* **54**: 2330-2337.

46.   *Ordnancesurveycouk* (2016) https://www.ordnancesurvey.co.uk/documents/os-mastermap-

sites-layer-user-guide.pdf.

47.   Pallares-Vega, R., Blaak, H., van der Plaats, R., de Roda Husman, A.M., Hernandez Leal, L.,

van Loosdrecht, M.C.M., Weissbrodt, D.G., and Schmitt, H. (2019) Determinants of presence and

removal of antibiotic resistance genes during WWTP treatment: A cross-sectional study. *Water Res*

161: 319-328.

48.   Papot, C., Massol, F., Jollivet, D., and Tasiemski, A. (2017) Antagonistic evolution of an

antibiotic and its molecular chaperone: how to maintain a vital ectosymbiosis in a highly fluctuating

habitat. *Sci Rep* **7**: 1454.

49.   Pazda, M., Kumirska, J., Stepnowski, P., and Mulkiewicz, E. (2019) Antibiotic resistance genes

identified in wastewater treatment plant systems – A review. *Sci Total Environ* **697**: 134023.

50.   Phuong Hoa, P.T., Nonaka, L., Hung Viet, P., and Suzuki, S. (2008) Detection of the sul1, sul2,

and sul3 genes in sulfonamide-resistant bacteria from wastewater and shrimp ponds of north

Vietnam. *Sci Total Environ* **405**: 377-384.

51.   Pruden, A., Larsson, D.G., Amézquita, A., Collignon, P., Brandt, K.K., Graham, D.W.,

Lazorchak, J.M., Suzuki, S., Silley, P., Snape, J.R., Topp, E., Zhang, T., and Zhu, Y.G. (2013)

Management options for reducing the release of antibiotics and antibiotic resistance genes to the

environment. *Environ Health Perspect* **121**: 878-885.

52.   Pruitt, K.D., Tatusova, T., and Maglott, D.R. (2005) NCBI Reference Sequence (RefSeq): a

curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res*

33: D501-504.

53.   Pärnänen, K., Karkman, A., Hultman, J., Lyra, C., Bengtsson-Palme, J., Larsson, D.G.J.,

Rautava, S., Isolauri, E., Salminen, S., Kumar, H., Satokari, R., and Virta, M. (2018) Maternal gut and

breast milk microbiota affect infant gut antibiotic resistome and mobile genetic elements. *Nat*

Commun **9**: 3891.

54.   Pérez, S., Eichhorn, P., and Aga, D.S. (2005) Evaluating the biodegradability of

sulfamethazine, sulfamethoxazole, sulfathiazole, and trimethoprim at different stages of sewage

treatment. *Environ Toxicol Chem* **24**: 1361-1367.

55.   Quinlan, A.R., and Hall, I.M. (2010) BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**: 841-842.

56.   Razavi, M., Marathe, N.P., Gillings, M.R., Flach, C.F., Kristiansson, E., and Joakim Larsson,

D.G. (2017) Discovery of the fourth mobile sulfonamide resistance gene. *Microbiome* **5**: 160.

57.   Rowland, C.S., Morton, R.D., Carrasco, L., McShane, G., O'Neil, A.W., Wood, C.M. (2017).

Land Cover Map 2015 (25m raster, GB). NERC Environmental Information Data

Centre. https://doi.org/10.5285/bb15e200-9349-403c-bda9-b430093807c7.

58.   Sengupta, A., Lyons, J.M., Smith, D.J., Drewes, J.E., Snyder, S.A., Heil, A., and Maruya, K.A.

(2014) The occurrence and fate of chemicals of emerging concern in coastal urban rivers receiving

discharge of treated municipal wastewater effluent. *Environ Toxicol Chem* **33**: 350-358.

59.   Sewage Treatment in the UK

(2020) *Assetspublishingservicegovuk* https://assets.publishing.service.gov.uk/government/uploads/

system/uploads/attachment_data/file/69582/pb6655-uk-sewage-treatment-020424.pdf.

60.   Singer, A.C., Shaw, H., Rhodes, V., and Hart, A. (2016) Review of Antimicrobial Resistance in

the Environment and Its Relevance to Environmental Regulators. *Front Microbiol* **7**: 1728.

61.   Singer, A.C., Xu, Q., and Keller, V.D.J. (2019) Translating antibiotic prescribing into antibiotic

resistance in the environment: A hazard characterisation case study. *PLoS One* **14**: e0221568.

62.   Sköld, O. (2000) Sulfonamide resistance: mechanisms and trends. *Drug Resist Updat* **3**: 155-

160.

63.   SOAF (2018) *Waterorguk* http://www.water.org.uk/wp-

content/uploads/2018/12/SOAF.pdf. Accessed May 7, 2020.

64.   Stoodley, P., Wilson, S., Hall-Stoodley, L., Boyle, J.D., Lappin-Scott, H.M., and Costerton, J.W.

(2001) Growth and detachment of cell clusters from mature mixed-species biofilms. *Appl Environ*

Microbiol **67**: 5608-5613.

65.   Tehrani, A.H., and Gilbride, K.A. (2018) A closer look at the antibiotic-resistant bacterial

community found in urban wastewater treatment systems. *Microbiologyopen* **7**: e00589.

66.   Thiebault, T. (2020) Sulfamethoxazole/Trimethoprim ratio as a new marker in raw

wastewaters: A critical review. *Sci Total Environ* **715**: 136916.

67.   Thiebault, T., Fougère, L., Destandau, E., Réty, M., and Jacob, J. (2019) Impact of

meteorological and social events on human-excreted contaminant loads in raw wastewater: From

daily to weekly dynamics. *Chemosphere* **230**: 107-116.

68.   Tran, N.H., Hoang, L., Nghiem, L.D., Nguyen, N.M.H., Ngo, H.H., Guo, W., Trinh, Q.T., Mai,

N.H., Chen, H., Nguyen, D.D., Ta, T.T., and Gin, K.Y. (2019) Occurrence and risk assessment of

843 multiple classes of antibiotics in urban canals and lakes in Hanoi, Vietnam. *Sci Total Environ* **692**:

844 157-174.

846 69. Uyaguari-Díaz, M.I., Croxen, M.A., Luo, Z., Cronin, K.I., Chan, M., Baticados, W.N., Nesbitt,

847 M.J., Li, S., Miller, K.M., Dooley, D., Hsiao, W., Isaac-Renton, J.L., Tang, P., and Prystajecky, N. (2018)

848 Human Activity Determines the Presence of Integron-Associated and Antibiotic Resistance Genes in

849 Southwestern British Columbia. *Front Microbiol* **9**: 852.

851 70. Vilchèze, C., and Jacobs, W.R. (2012) The combination of sulfamethoxazole, trimethoprim,

852 and isoniazid or rifampin is bactericidal and prevents the emergence of drug resistance in

853 Mycobacterium tuberculosis. *Antimicrob Agents Chemother* **56**: 5142-5148.

855 71. Wellington, E.M., Boxall, A.B., Cross, P., Feil, E.J., Gaze, W.H., Hawkey, P.M., Johnson-

856 Rollings, A.S., Jones, D.L., Lee, N.M., Otten, W., Thomas, C.M., and Williams, A.P. (2013) The role of

857 the natural environment in the emergence of antibiotic resistance in gram-negative bacteria. *Lancet*

858 *Infect Dis* **13**: 155-165.

860 72. Wick, R.R., Schultz, M.B., Zobel, J., and Holt, K.E. (2015) Bandage: interactive visualization of

861 de novo genome assemblies. *Bioinformatics* **31**: 3350-3352.

863 73. Zhang, X., Li, Y., Liu, B., Wang, J., Feng, C., Gao, M., and Wang, L. (2014) Prevalence of

864 veterinary antibiotics and antibiotic-resistant Escherichia coli in the surface water of a livestock

865 production region in northern China. *PLoS One* **9**: e111026.

867   74.   Östman, M., Lindberg, R.H., Fick, J., Björn, E., and Tysklind, M. (2017) Screening of biocides,

868   metals and antibiotics in Swedish sewage sludge and wastewater. *Water Res* **115**: 318-328.

870   75.   Šeputienė, V., Povilonis, J., Ružauskas, M., Pavilonis, A., and Sužiedėlienė, E. (2010)

871   Prevalence of trimethoprim resistance genes in Escherichia coli isolates of human and animal origin

872   in Lithuania. *J Med Microbiol* **59**: 315-322.