Title: **Abiotic treatment to common bean plants results in an altered seed microbiome**

Authors: A. Fina Bintarti[1,2], Patrick J. Kearns[2]*, Abby Sulesky[2,3], and Ashley Shade[1,2,3]

1. Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing MI

48824

2. The Plant Resilience Institute, Michigan State University, East Lansing MI 48824

3. Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing MI

48824

current affiliation: Department of Natural and Applied Sciences, Bentley University, Waltham, MA
02452
11

Abstract

We performed a pilot study in a controlled growth chamber to investigate how the seed microbiome of

the common bean (*Phaseolus vulgaris L.* (var. Redhawk)) was altered under abiotic treatments relevant

for crop management with changing climate. Bean plants were subjected to one of three treatments:

66% water withholding to simulate mild drought, 50 % Hoagland nutrient solution to simulate

fertilization, or control with sufficient water and baseline nutrition. We performed 16S rRNA gene

amplicon sequencing and ITS amplicon sequencing of the endophytic metagenomic DNA to assess seed

bacterial/archaeal and fungal community structure, respectively. We found that variability in the seed

microbiome structure was high while alpha diversity was low, with tens of taxa present. Water

withholding and nutrient addition altered the seed microbiome structure for bacterial/archaeal

communities as compared to the untreated control, and each treatment resulted in a distinct

microbiome structure. There were no statistically supported differences in the fungal microbiome across

treatments. While we discuss several limitations of this study, the promising results suggest that further

and more comprehensive investigation is needed to better understand abiotic stress-induced changes in

the seed microbiome, the mechanisms that drive those changes, and the implications of seed

microbiome changes for the health and stress responses of the next plant generation.

Key words

plant microbiome, 16S rRNA gene, ITS, drought, fertilizer, legume, growth chamber, endophyte, vertical

transmission, abiotic stress

Introduction

The plant microbiome includes bacteria, archaea, fungi, and viruses that associate with the plant

and inhabit different plant compartments, including the rhizosphere, phyllosphere, and endosphere

(Compant et al., 2019). The plant microbiome plays important roles for plant fitness, including nutrient

acquisition (Herrera Paredes et al., 2018), secondary metabolite production (Gargallo-Garriga et al.,

2016), flowering time (Panke-Buisse et al., 2015), and resistance to abiotic (Lau & Lennon, 2012; Naylor

& Coleman-Derr, 2018) and biotic stresses (Mendes et al., 2011; Ritpitakphong et al., 2016). Plant

microbiota can interact with each other as well as with the host plant, and the plant is able to shape its

microbiome structure and composition, for example, by producing root exudates or allelochemicals

(Jones et al., 2019; Sasse et al., 2018). Because plants are also influenced by the environment, any

stresses that a plant experiences are likely to also affect the plant microbiome. These environmental

stresses include abiotic stressors, such as changes in water and nutrient availability (limitation or excess)

and exposure to high temperatures, and biotic stressors, such as pathogen infection and herbivory.

Regardless of the type of stressor, as the plant responds it may alter its microbiome, either as a direct or

indirect consequence of the stress. Thus, both the environmental conditions and the host plant can act

as important filters that contribute to the ultimate composition of the plant microbiome (Hacquard et

al., 2017; Vacher et al., 2016).

Managing or manipulating the plant microbiome is one promising strategy to support plant

tolerance to environmental stress. We are just beginning to understand how exactly the plant

microbiome structure is altered during particular stressors (e.g., drought (Xu et al., 2018), with a focus

on the root zone and rhizosphere microbiome. We conducted a controlled study in an environmental

growth chamber to characterize changes in the rhizosphere microbiome of the legume common bean

under two different stressors of water withholding and nutrient excess. The initial purpose of our pilot

study was to identify members of the root microbiome that were particularly resilient either to these

stressors. However, the plants set pods at the end of the experiment, and we realized the opportunity to

also assess the seed microbiome of the treated plants as compared to control plants. The purpose of

this brief report is to share the seed microbiome results from the pilot study, to discuss its limitations,

and to suggest immediate future directions based on the most promising results.

Methods

Plant growth conditions and harvest. Common bean seeds were surface-sterilized in a 10% bleach

solution and planted in 24 one-gallon pots filled with a mixture of agricultural topsoil, sphagnum peat

and sand that had been steam-sterilized at ~100°C. The plants were grown in controlled conditions

in a BioChambers FLEX™ LED growth chamber with a 16-hour day/8-hour night cycle at 29°C and 22°C,

respectively. The plants were divided into three groups: 8 control plants received ample water (300 mL

every other day), 8 plants were subjected to a mild "drought" during plant development, and received

66% less water (100 mL every other day) (water withholding), and 8 plants received half strength

Hoagland solution (300 mL every other day) provided by the growth chamber facility (nutrient addition).

The plants were grown until the R8 stage, when plants were fully developed, and the seed pods began

drying.

Harvesting was conducted by collecting the bean pods and plant biomass. Bean seeds were

removed from the pods and the remaining above ground biomass from each plant was placed in a

brown paper bag and dried at 70°C for one week. The root system was gently pulled from the pot,

cleaned of excess soil with deionized water and dried at 70°C for one week. Once dried, the above and

below ground dry weight was measured for each plant. The remaining soil was collected for soil

chemical analysis. One hundred grams of each soil sample was sent to the Michigan State University Soil

and Plant Nutrient Laboratory (SPNL) for soil chemical testing. Soil parameters including pH, lime index,

phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), nitrate (NO_3^-), ammonium (NH_4^+), and

organic matter (OM) were measured from all soil samples.

DNA extraction and amplicon sequencing. Twenty seeds from each plant were collected for DNA

extraction following the protocol of the previous study (Barret et al., 2015). Seeds were surfaced

sterilized in 10% bleach and placed in sterile 50 mL centrifuge tubes with 30 mL of sterile 1X PBS with

0.05% Tween 20 and shaken at 140 rpm at room temperature for 4 hours. After shaking, tubes were

centrifuged at 500 x g for 15 minutes and the supernatant and seeds were discarded. The pellet was

resuspended with 2 mL of sterile 1X PBS-Tween and transferred to a microcentrifuge tube and spun at

20,000 x g for 10 min. The supernatant was discarded, and the pellet was used for DNA extraction with

the PowerSoil®DNA Isolation Kit (MoBio Laboratories, Solana Beach, CA, United States) following

manufacturer's instructions. DNA extracted from seeds were quantified with Qubit™dsDNA BR Assay Kit

(ThermoFisher Scientific, Waltham, MA, United States) and verified with Polymerase Chain Reaction

(PCR) using 515f/806r universal primers (Caporaso et al., 2011) for amplification of the V4 region of the

16S rRNA gene for bacterial/archaeal community analysis. The 16S rRNA gene amplification was

conducted under the following conditions: 95°C for 5 min, followed by 30 cycles of 95°C (30 s), 54°C (45

s), and 72°C (90 s), with a final extension at 72°C (5 min). The amplification was performed in 12.5 µl

mixtures containing 6.25 µl GoTaq®Green Master Mix (Promega, Madison, WI, United States), 0.125 µl

of each primer (20 mM), 1 µl of DNA template (final concentration of 0.02 - 0.626 ng per µl), and 4.5 µl

nuclease free water. Seed DNA (concentration range of 5 - 20 ng per µl) was sequenced at the Research

Technology Support Facility (RTSF) Genomics Core, Michigan State sequencing facility using Illumina

MiSeq platform.

Fungal communities were assessed using PCR amplification of the Internal Transcribed Spacer 1

(ITS1) region with the ITS1f/ITS2 primer pair (Walters et al., 2015) with the addition of index adapters as

required by the RTSF Genomics Core, (https://rtsf.natsci.msu.edu/genomics/sample-

[requirements/illumina-sequencing-sample-requirements/](https://rtsf.natsci.msu.edu/genomics/sample-requirements/illumina-sequencing-sample-requirements/)). The PCR conditions of ITS gene amplification

were as follows: 95°C for 5 min, followed by 30 cycles of 95°C (30 s), 54°C (45 s), and 72°C (90 s), with a

final extension at 72°C (5 min). The amplification was performed in 50 µl mixtures containing 20 µl

GoTaq®Green Master Mix (Promega, Madison, WI, United States), 1 µl of each primer (20 mM), 4 µl of

DNA template (final concentration of 0.02 – 0.626 ng per µl), and 26 µl nuclease free water. The product

of the ITS gene amplification was cleaned and purified using the Wizard®SV Gel and PCR Clean-Up

System (Promega, Madison, WI, United States), following the manufacturer's protocol. Purified ITS gene

amplification products with the concentration range of 5 - 50 ng per µl were sequenced at the RTSF

Genomics Core using the Ilumina MiSeq platform. The 16S and ITS libraries were prepared using the

Illumina TruSeq®Nano DNA Library Prep Kit. Ilumina MiSeq was run using v2 Standard and paired end

reads sequencing format (2 x 250 bp).

Sequencing data analysis and OTU clustering. Bacterial/archaeal raw reads produced from Illumina

MiSeq were processed including merging the paired end reads, filtering the low-quality sequences,

dereplication to find unique sequence, singleton removal, denoising, and chimera checking using the

USEARCH pipeline (v.10.0.240) (Edgar & Flyvbjerg, 2015). Operational taxonomic unit (OTU) clustering

121 was conducted using an open reference strategy (Lee et al., 2017). First, closed reference OTU picking

122 was performed at 97 % identity by clustering quality filtered reads against the SILVA database (v.132)

123 (Quast et al., 2013) using USEARCH algorithm (-*usearch_global* command) (Edgar, 2010). Reads that

124 failed to match the SILVA reference were subsequently clustered *de novo* at 97% identity using UPARSE-

125 OTU algorithm (-*cluster_otus* command) (Edgar, 2013). Closed reference and *de novo* OTUs were

126 combined into a full set of representative sequences, and then all merged sequences were mapped back

127 to that set using the -*usearch_global* command.

128  The set of representative sequences were aligned on QIIME 1.9.1 (Caporaso et al., 2010b) using

129 PyNAST (Caporaso et al., 2010a) against the SILVA (v.132) reference database. The unaligned OTU

130 sequences were excluded from the OTU table and the representative sequences. Taxonomic assignment

131 was conducted on QIIME 1.9.1 using the SILVA (v.132) database and the UCLUST default classifier at a

132 minimum confidence of 0.9 (Edgar, 2010). Plant contaminants such as chloroplast and mitochondria;

133 and unassigned taxa and sequences were removed from the OTU table as well as the representative

134 sequences using *filter_taxa_from_otu_table.py* and *filter_fasta.py* command on QIIME. Rarefaction to

135 the lowest sequencing depth (Gihring et al., 2012; Weiss et al., 2017) (11,137 bacterial/archaeal reads)

136 was conducted on QIIME.

137  The processing of fungal ITS raw reads was also conducted using the USEARCH (v.10.0.240)

138 pipeline. Read processing included paired end read merging, primer removal using cutadapt (v.2.0)

139 (Martin, 2011), filtering the low-quality sequences, and dereplication to find unique sequence.

140 Operational taxonomic unit clustering was conducted using an open reference OTU picking strategy.

141 First, closed reference OTU picking was performed by clustering quality filtered reads against the UNITE

142 fungal ITS database (v.8.0) (Kõljalg et al., 2013) at 97% identity threshold using the USEARCH algorithm.

143 Reads that failed to match the reference were clustered *de novo* at 97% identity using the UPARSE-OTU

144 algorithm. Closed reference and *de novo* OTUs were combined into a full set of representative

sequences, and then all merged sequences were mapped back to that set using -*usearch_global*

command. Fungal taxonomic classification was performed in CONSTAX tool (Gdanetz et al., 2017) at a

minimum confidence of 0.8 using the UNITE reference database release 01-12-2017. Assigning

taxonomy in CONSTAX was conducted using three classifiers, including RDP Classifier (v.11.5) (Cole et al.,

2014; Wang et al., 2007), UTAX from USEARCH (v.8.1.1831) (Edgar, 2013), and SINTAX from USEARCH

(v.9.2) (Edgar, 2016). Any contaminants including mitochondria, chloroplast and other unwanted

lineages of eukaryotes were removed from the OTU table. Rarefaction was conducted to the lowest

number of sequences (21,329 fungal reads) on QIIME.

Microbial community analysis. Microbial community analyses were conducted in R (v.3.6.1) (R Core

Development Team). Microbial composition and relative abundance were analyzed using Phyloseq

package (v.1.28.0) on R (McMurdie & Holmes, 2013). Microbial richness (the number of taxa present)

was calculated on the rarefied OTU table using the vegan package (v.2.5-6) (Oksanen et al., 2019). The

normality and homoscedasticity of the data were tested using Saphiro-Wilk and Levene's test,

respectively. The one-way analysis of variance (one-way ANOVA) or non-parametric Kruskal-Wallis test

was then performed to analyze the data. Post hoc Dunn's test with false discovery rate (FDR) correction

using the Benjamini-Hochberg adjustment for multiple comparisons was performed to compare plant

biomass data among treatments.

Beta diversity was calculated on the rarefied OTU table using the vegan package using Jaccard

dissimilarity indices and visualized with a principal coordinate analysis (PCoA) plot. We used the Jaccard

index, which is based on presence-absence counts rather than relative abundance data, because we

reasoned that the seed microbiome members are unlikely to be actively growing inside the seed and

that any differences in relative abundances in the seed endophyte are unlikely attributable to

competitive growth outcomes *in situ*. Permutational multivariate analysis of variance (PERMANOVA)

using the function adonis (Oksanen et al., 2019) was performed to assess the effects of the treatments

to the microbial community structure. We performed multivariate analysis to check the homogeneity of

dispersion (variance) among groups using the function betadisper (Oksanen et al., 2019).

Data and code availability. The computational workflows for sequence processing and ecological

statistics are available on GitHub (https://github.com/ShadeLab/BioRxiv_Seed_Microbiome_2020). Raw

sequence data of bacteria/archaea and fungi have been deposited in the Sequence Read Archive (SRA)

NCBI database under Bioproject accession number PRJNA635871.

Results

There were overall differences in plant biomass among treatments (Kruskal-Wallis results Table

1). Specifically, plants receiving nutrient-addition were larger in shoot and root biomass than control or

mildly droughted plants (Fig. 1, Table 1). Nutrient-addition plants also had higher pod number and pod

mass compared to the water withholding and control plants (Fig. 1, Table 1). Rhizosphere soil from the

nutrient addition treatment had higher phosphorus and potassium content than the other two

treatments (Fig. 2, Table 2). Nutrient addition plants also had rhizosphere soils with higher nitrate

relative to control plants (Fig. 2, Table 2).

Analysis of bacterial/archaeal and fungal sequences from seed samples resulted in a total of 81

and 226 OTUs (97% sequence identity), respectively. Bacterial/archaeal communities in control, water

withholding, and nutrient addition seeds had different taxonomic compositions (Fig. 3A).

Bacterial/archaeal communities in the control seeds were almost exclusively dominated by the OTUs

within the genus *Bacillus,* with a mean relative genus-level abundance of more than 99%. Although the

bacterial/archaeal community in the water withholding and nutrient addition seeds were also

dominated by *Bacillus*, genus-level taxonomic diversity increased with the addition of other, non-

dominating lineages. Specifically, water with-holding and nutrient addition seed communities were also

composed of *Virgibacillus*, *Pseudomonas*, and several other bacterial/archaeal genera.

Similarly, different plant treatments had different seed fungal community compositions. Even

though *Aspergillus* dominated the fungal community in the control and treated seeds, in the treated

seeds there was a shift to include other fungal taxa, including some identified as *Penicillium* and

Wallemia (Fig. 3B). These observations indicate that the seed microbiome is altered when maternal

plants are exposed to abiotic stress or environmental alternation. Since the seed microbiomes contained

a relatively simple community of tens to dozens of taxa, even alterations in the composition or

abundances of few taxa may have consequence for microbiome assembly of the next plant generation.

Because we do not expect microbiome members to be actively doubling inside the seed, we

used a presence-absence assessment (Jaccard index) of beta diversity. There was a statistically

supported difference in bacterial/archaeal microbiome composition between treated seeds and control

seeds (Fig. 4A, permuted multivariate analysis of variance, PERMANOVA, F-stat = 4.73, R^2 = 0.31, P-val =

0.001). In contrast, there was no distinct clustering of fungal communities associated with different

treatments (Fig. 4B, PERMANOVA P-val > 0.05). These results indicate that abiotic treatments alter the

bacterial/archaeal but not fungal community composition in the common bean seed. Also, there were

differences variability in composition (multivariate dispersion) among treatments for bacterial/archaeal

communities (permutated analysis of multivariate dispersion, PERMDISP, F=7.553, P-val=0.0033), but

not for fungal composition (PERMDISP F=0.491, P-val=0.619). This provide additional evidence that

abiotic treatments can lead to increased variability in seed microbiome composition. Notably,

PERMANOVA was found to be largely unaffected by heterogeneity for balanced designs (Anderson &

Walsh, 2013).

Discussion

Seed microbiomes are an endpoint of microbiome assembly for the maternal plant's

reproductive compartment, but also the starting point of microbiome assembly for the new seedling

(Shade et al., 2017). Seed microbiomes are composed of early colonizers that are acquired from the

maternal plant either through the vascular system or floral stigma, and also of late colonizers that are

acquired via seed contact with its environment (Maude, 1996). Moreover, the seed microbiome also can

directly impact the seed and seedling in ways that are important for crop establishment, such as by

releasing the seed from dormancy and promoting seed germination and seedling emergence (Goggin et

al., 2015; Lamichhane et al., 2018). While the vast majority of plant-associated microorganisms are

acquired from the surrounding environment, (e.g., aerosols (Vacher et al., 2016) and soils (Edwards et

al., 2015)), vertical transmission of microbes via the seed has been reported for a variety of plant

species, as recently summarized (Shade et al., 2017) and reported (Bergna et al., 2018; Rodríguez et al.,

2020). For the majority of seed microbiome members, their identities, functions and persistence are

either not known or not well-understood. However, it is plausible that some of members of the seed

microbiome that are altered as a result of stress may have consequence for the health or resilience of

the next plant generation. Specifically, a depletion of beneficial members or enrichment of pathogens

could disadvantage the plant, while an enrichment of beneficial members could advantage the plant.

We highlight three important observations from this study. First, treatment of the parent plant

altered the seed microbiome structure and composition compared to control plants, especially for the

bacterial/archaeal community. Second, bacterial/archaeal community from the seeds of treated plants

had more variation as compared to the seeds from control plants. This suggests that abiotic stress

results in changes analogous to those observed during other types of microbiome "dysbiosis" (aka Anna

Karenina effects: higher variability across replicates, increased beta-dispersion, and higher contribution

of stochastic assembly processes (Zaneveld et al., 2017)). The expectation of high variability should be

taken into consideration for future studies, as sufficient replication will be needed to power statistical

tests. Third and finally, the fungal community was, on balance, stable relative to the bacterial/archaeal

community, suggesting that the persistence of fungal members is less sensitive to water-withholding

and nutrient addition.

We acknowledge that this is a pilot study and that these results are preliminary. We offer a

discussion of some of the major considerations and limitations in interpreting the results and for

planning future seed microbiome studies.

A first consideration is that there is an apparent maximum stress to plants that can be applied

when investigating its consequence for a seed microbiome. After stress exposure is released, plants

must be healthy enough to produce pods and seeds, and a balance must be achieved in which plants are

stressed but still able to become fully mature. This constraint in stress exposure will never

accommodate an experimental design of severe or prolonged stress. However, the investigation of a

mild or moderate stress is still valuable because it is pertinent to agriculture. There many situations in

which non-lethal stress occurs over part of a growing season, but then crops recover fully or partially to

produce some yield. Therefore, the result of mild or moderate stress for seed microbiomes has real-

world relevance.

Another consideration is the definition and directness of abiotic treatment, and whether an

abiotic treatment is expected to act on the plant, the microbiome, or both. In this pilot study, we applied

two different abiotic treatments: one that was expected to stress the plant directly (water-withholding

to simulate mild drought) and one that was expected to weaken a legume's relationship with its root-

associated microbiome and symbiotic nitrogen fixers as nitrogen fixers are down-regulated by nitrogen

application (Müller & Pereira, 1995; Wilker et al., 2019) (nutrient addition). Thus, the addition of

nutrients was a benefit to the plant, rather than a stress, as indicated by the increased root and shoot

masses. However, nutrient addition caused a clear shift in the seed microbiome, demonstrating the

potential of fertilizer use to have multi-generational for plant microbiome assembly. Therefore,

management practices that advantage the plant as far as yield and health in the short term could have

long-term consequences for plant-microbiome relationships.

A clear limitation of the study is the substrate used for plant growth, which, with the microbes in

and on the original seeds, serves as a starting source for the assembly of the new plant's microbiome

(Barret et al., 2015; Shade et al., 2017). For the pilot, used agricultural topsoil mixed with sphagnum

peat and sand and sterilized, and this mix is provided by the growth chamber facility. The exact origin

and physical/chemical characteristics of facility topsoil us unknown, and so it is unclear how

representative this soil may be of bean field soil. Additionally, the initial microbial community in the soil

was not analyzed before planting the common bean plants, so we cannot determine the origin of the

observed microbial consortia in the seeds and to what extent they overlap with the potting substrate.

Previous work suggests that soil type can have a large influence on the seed endophytic bacteria in rice

(Hardoim et al., 2012), and this is likely also true for other plant seeds. We observed a dominance of a

specific taxon from genus *Bacillus* in common bean seeds in all three treatments. While *Bacillus* taxa

were reported to be enriched in the green bean seedling (Barret et al., 2015), steam sterilization of the

growth chamber soil may have killed most of the indigenous microbial taxa that were not spore-formers

or otherwise resistant to heat (and, bacilli are known to be resistant to such treatments, as per

(Nicholson et al., 2000)). Therefore, the microbial composition that were observed in these bean seeds

may not be representative of or similar to those in bean seeds grown in the field, or even in different

substrates. We urge caution in generalizing from the compositional changes, but rather to focus on the

larger changes in beta-diversity and dispersion that were consistently observed across very different

abiotic treatments and may be more characteristic of seed microbiome responses. Future work should

focus efforts on using soil that is representative of the typical agricultural environment of common

bean, and the existing microbial community in the soil should be sequenced prior to planting for

comparison to the seed microbiome.

Another limitation of this study the absence of negative controls for DNA extraction controls.

Seed endophytes contain a very low total biomass of microbial cells. Here, we pooled twenty seeds to

use for one extraction to increase the microbial biomass yield for microbiome interrogation. However,

we did not perform a buffer-only control, which would allow for assessment of contaminants from the

DNA extraction process. While the surface-sterilization of the seeds prior to extraction and negative PCR

controls provide confidence that the starting material was not compromised and that we did not

unintentionally amplify contaminants from the PCR reagents, we cannot know if there were or

extraction kit or buffer contaminants that contributed to the observed seed microbiome composition.

A final minor limitation is in the choice of bacterial marker gene. We performed amplicon

sequencing of the 16S rRNA gene for the bacterial/archaeal community analysis in the seed. It is well

known that the variability of rRNA copy numbers among bacterial species can lead to an inflation of

species richness and obfuscate relative abundances of taxa (Větrovský & Baldrian, 2013). Moreover, the

most resolved taxonomic level achieved from 16S rRNA amplicon gene sequences often is genus, rather

than species or strain. Using a single-copy marker gene that has higher precision and sensitivity at the

species level, like the *gyr*B gene, may be a valuable alternative (Barret et al., 2015). The *gyr*B gene has

been successfully applied to other seed microbiome studies (Barret et al., 2015; Rezki et al., 2016; Rezki

et al., 2017). It may be valuable to consider use of both marker genes to the same seed microbiome

samples, so that taxonomic precision can be maximized and compared across seeds with *gyrB*, while

also maintaining an ability to source-track and compare composition across the many plant-microbiome

16S rRNA amplicon datasets that have been deposited publicly.

In summary, while this pilot study provides a key insight into the response of the seed

microbiome structure to abiotic treatment in the host plant, there is much more work to be done. Next

steps include exposing the plants to more severe drought and nutrient excess conditions, quantifying

the physiological status of plants to determine their experience of stress, using representative field soil

for plants and assessing the field soil microbiome to deduce seed taxon origins, using negative controls

for both DNA extractions to identify contaminants, and considering use of an alternative marker gene

for improved precision in microbial taxonomy and taxon abundances.

Despite noted considerations and limitations, we posit that this pilot study revealed an

important insight regarding how seed microbiomes may be altered after abiotic treatment of a plant.

Next, we need to understand the implications of this change for both the host plant and the microbial

community. An altered seed microbiome may have positive, negative, or entirely neutral outcomes for

the next plant generation. Additional work is needed to understand these outcomes over consecutive

plant generations to determine the effects on plant fitness and resilience. If positive or negative

outcomes are detected, this work opens a new direction of research that could spur exciting

applications in plant-microbiome management.

Acknowledgements

This work was supported by the Michigan State University Plant Resilience Institute and by the USDA

National Institute of Food and Agriculture award USDA 2019-67019-29305. AFB acknowledges support

from the Fulbright Foreign Student Program and Office for International Students and Scholars (OISS),

Michigan State University; and AS acknowledges support from the USDA National Institute of Food and

Agriculture and Michigan AgBioResearch (Hatch). We thank Matthieu Barret and the Shade Lab for

helpful discussions.

332
333

334 **Tables**

335 Table 1. Comparison of plant biomass among treatments. Bolded rows are significantly different as

336 determined by Kruskal-Wallis test and post-hoc Dunn's test with Benjamini-Hochberg false discovery

337 rate (FDR) correction

	Kruskal-Wallis test			Post-hoc Dunn's test with Benjamini-Hochberg FDR correction			
Parameter	Chi-squared	df	p-val	Comparison	Z	P. unadj	P. adj
Shoot mass	20.165	2	**0.00004**	Control – Nutrient addition	-2.29810	0.02156	**0.03233**
				Control – Water withholding	2.19203	0.02838	**0.02838**
				Nutrient addition – Water withholding	4.49013	0.00001	**0.00002**
Root mass	15.365	2	**0.00046**	Control – Nutrient addition	-3.42947	0.00060	**0.00181**
				Control – Water withholding	-0.07071	0.94363	0.94363
				Nutrient addition – Water withholding	3.35876	0.00078	**0.00117**
Pod mass	18.305	2	**0.00010**	Control – Nutrient addition	-2.08597	0.03698	**0.03698**
				Control –Water withholding	2.19203	0.02838	**0.04257**
				Nutrient addition – Water withholding	4.27800	0.00002	**0.00006**
Pod number	17.973	2	**0.00013**	Control – Nutrient addition	-2.53347	0.01129	**0.01694**
				Control – Water withholding	1.67708	0.09353	0.09353
				Nutrient addition – Water withholding	4.21055	0.00003	**0.00008**

340   Table 2. Comparison of rhizosphere soil chemistry of the three treatments using one-way ANOVA or

341   Kruskal-Wallis test. Bolded rows are significantly different as determined by Kruskal-Wallis test and post-

342   hoc Dunn's test with Benjamini-Hochberg false discovery rate (FDR) correction.

	Kruskal-Wallis test			Post-hoc Dunn's test with Benjamini-Hochberg FDR correction			
Parameter	Chi-squared	df	p-value	Comparison	Z	P. unadj	P. adj
pH	14.634	2	**0.00066**	Control – Nutrient addition	3.824263	0.00013	**0.00039**
				Control – Water withholding	1.83057	0.06716	0.06716
				Nutrient addition – Water withholding	-1.99369	0.04619	0.06928
P	15.613	2	**0.00041**	Control – Nutrient addition	-3.60033	0.00032	**0.00095**
				Control – Water withholding	-0.39018	0.69640	0.69640
				Nutrient addition – Water withholding	3.21014	0.00133	**0.00199**
K	19.172	2	**0.00007**	Control – Nutrient addition	-4.34149	0.00001	**0.00004**
				Control –Water withholding	-1.67822	0.09330	0.09330
				Nutrient addition – Water withholding	2.66327	0.00774	**0.01161**
NO_3^-	14.261	2	**0.0008**	Control – Nutrient addition	-3.67776	0.00024	**0.00071**
				Control –Water withholding	-2.58150	0.00984	**0.01476**
				Nutrient addition – Water withholding	1.09625	0.27297	0.27297
Mg	4.8514	2	0.08842				
OM	0.53985	2	0.7634				
	One-way ANOVA test						
Parameter	F-value	df	p-value				
Ca	2.035	2	0.156				
NH_4^+	0.343	2	0.713				

Figure legends

Fig 1. Plant aboveground (shoot) and belowground (root) biomass for control, water withholding, and nutrient addition treatments of common bean. Plant biomasses were calculated on eight plant replicates for each treatment. For each box plot, circles represent a single plant measurement within a treatment. The central horizontal lines represent the mean, the outer horizontal lines of the box represent the 25th and 75th percentiles. Boxes labelled with different letters were significantly different by a Kruskal-Wallis and post-hoc Dunn's test with a Benjamini-Hochberg false discovery rate correction.

Fig 2. Plant rhizosphere soil chemistry for control, water with-holding, and nutrient addition treatments of common bean. For each box plot, circles represent one rhizosphere measurement within a treatment. The central horizontal lines represent the mean of measurement, the outer horizontal lines of the box represent the 25th and 75th percentiles. Boxes labelled with different letters are identified as significantly different by a Kruskal-Wallis and post-hoc Dunn's test with a Benjamini-Hochberg false discovery rate correction.

Fig 3. Mean relative abundances of genera of bacterial/archaeal (A) and fungal (B) detected in the seed across control, water with-holding and nutrient addition treatments. Each bar represents the endophyte microbiome from DNA extracted from 20 seeds collected from one plant replicate within a treatment. Bacterial/archaeal and fungal genera with mean relative abundances of less than 1 and 10 %, respectively, were grouped into the 'Other' classification, which includes many lineages (not monophyletic).

Fig. 4 Principal coordinate analysis (PCoA) plot of the bacterial/archaeal (A) and fungal community (B) in

the common bean seed based on the Jaccard index. Symbol colors and shapes represent different

abiotic treatments.

References

Anderson, M. J., & Walsh, D. C. I. (2013). PERMANOVA, ANOSIM, and the Mantel test in the face of
heterogeneous dispersions: What null hypothesis are you testing? *Ecological Monographs,*
83(4), 557-574. doi:10.1890/12-2010.1
Barret, M., Briand, M., Bonneau, S., Préveaux, A., Valière, S., Bouchez, O., . . . Jacques, M.-A. (2015).
Emergence Shapes the Structure of the Seed Microbiota. *Applied and Environmental*
Microbiology, 81(4), 1257-1266. doi:10.1128/aem.03722-14
Bergna, A., Cernava, T., Rändler, M., Grosch, R., Zachow, C., & Berg, G. (2018). Tomato Seeds Preferably
Transmit Plant Beneficial Endophytes. *Phytobiomes Journal, 2*(4), 183-193.
doi:10.1094/pbiomes-06-18-0029-r
Caporaso, J. G., Bittinger, K., Bushman, F. D., DeSantis, T. Z., Andersen, G. L., & Knight, R. (2010a).
PyNAST: a flexible tool for aligning sequences to a template alignment. *Bioinformatics, 26*(2),
266-267. doi:10.1093/bioinformatics/btp636
Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., . . . Knight, R.
(2010b). QIIME allows analysis of high-throughput community sequencing data. *Nature*
Methods, 7(5), 335-336. doi:10.1038/nmeth.f.303
Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., . . . Knight,
R. (2011). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
Proceedings of the National Academy of Sciences, 108(Supplement 1), 4516-4522.
doi:10.1073/pnas.1000080107
Cole, J. R., Wang, Q., Fish, J. A., Chai, B., McGarrell, D. M., Sun, Y., . . . Tiedje, J. M. (2014). Ribosomal
Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res,*
42(Database issue), D633-642. doi:10.1093/nar/gkt1244
Compant, S., Samad, A., Faist, H., & Sessitsch, A. (2019). A review on the plant microbiome: ecology,
functions and emerging trends in microbial application. *Journal of advanced research, 19*, 29-97.
doi:https://doi.org/10.1016/j.jare.2019.03.004
Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics, 26*(19),
2460-2461. doi:10.1093/bioinformatics/btq461
Edgar, R. C. (2013). UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature*
Methods, 10(10), 996-998. doi:10.1038/nmeth.2604
Edgar, R. C. (2016). SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences.
bioRxiv, 074161. doi:10.1101/074161
Edgar, R. C., & Flyvbjerg, H. (2015). Error filtering, pair assembly and error correction for next-generation
sequencing reads. *Bioinformatics, 31*(21), 3476-3482. doi:10.1093/bioinformatics/btv401
Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N. K., Bhatnagar, S., . . . Sundaresan,
V. (2015). Structure, variation, and assembly of the root-associated microbiomes of rice.
Proceedings of the National Academy of Sciences, 112(8), E911-E920.
doi:10.1073/pnas.1414592112
Gargallo-Garriga, A., Sardans, J., Pérez-Trujillo, M., Guenther, A., Llusià, J., Rico, L., . . . Parella, T. (2016).
Shifts in plant foliar and floral metabolomes in response to the suppression of the associated
microbiota. *BMC plant biology, 16*(1), 78. doi:https://doi.org/10.1186/s12870-016-0767-7
Gdanetz, K., Benucci, G. M. N., Vande Pol, N., & Bonito, G. (2017). CONSTAX: a tool for improved
taxonomic resolution of environmental fungal ITS sequences. *BMC Bioinformatics, 18*(1), 538.
doi:10.1186/s12859-017-1952-x
Gihring, T. M., Green, S. J., & Schadt, C. W. (2012). Massively parallel rRNA gene sequencing exacerbates
the potential for biased community diversity comparisons due to variable library sizes.
Environmental Microbiology, 14(2), 285-290. doi:10.1111/j.1462-2920.2011.02550.x

Goggin, D. E., Emery, R. J. N., Kurepin, L. V., & Powles, S. B. (2015). A potential role for endogenous
microflora in dormancy release, cytokinin metabolism and the response to fluridone in Lolium
rigidum seeds. *Ann Bot, 115*(2), 293-301. doi:10.1093/aob/mcu231
Hacquard, S., Spaepen, S., Garrido-Oter, R., & Schulze-Lefert, P. (2017). Interplay Between Innate
Immunity and the Plant Microbiota. *Annu Rev Phytopathol, 55*, 565-589. doi:10.1146/annurev-
phyto-080516-035623
Hardoim, P. R., Hardoim, C. C. P., van Overbeek, L. S., & van Elsas, J. D. (2012). Dynamics of seed-borne
rice endophytes on early plant growth stages. *PloS one, 7*(2), e30438-e30438.
doi:10.1371/journal.pone.0030438
Herrera Paredes, S., Gao, T., Law, T. F., Finkel, O. M., Mucyn, T., Teixeira, P. J. P. L., . . . Castrillo, G.
(2018). Design of synthetic bacterial communities for predictable plant phenotypes. *PLOS
Biology, 16*(2), e2003962. doi:10.1371/journal.pbio.2003962
Jones, P., Garcia, B. J., Furches, A., Tuskan, G. A., & Jacobson, D. (2019). Plant Host-Associated
Mechanisms for Microbial Selection. *Frontiers in Plant Science, 10*(862).
doi:10.3389/fpls.2019.00862
Kõljalg, U., Nilsson, R. H., Abarenkov, K., Tedersoo, L., Taylor, A. F. S., Bahram, M., . . . Larsson, K.-H.
(2013). Towards a unified paradigm for sequence-based identification of fungi. *Molecular
Ecology, 22*(21), 5271-5277. doi:10.1111/mec.12481
Lamichhane, J. R., Debaeke, P., Steinberg, C., You, M. P., Barbetti, M. J., & Aubertot, J.-N. (2018). Abiotic
and biotic factors affecting crop seed germination and seedling emergence: a conceptual
framework. *Plant and Soil, 432*(1), 1-28. doi:10.1007/s11104-018-3780-9
Lau, J. A., & Lennon, J. T. (2012). Rapid responses of soil microorganisms improve plant fitness in novel
environments. *Proceedings of the National Academy of Sciences, 109*(35), 14058-14062.
doi:10.1073/pnas.1202319109
Lee, S.-H., Sorensen, J. W., Grady, K. L., Tobin, T. C., & Shade, A. (2017). Divergent extremes but
convergent recovery of bacterial and archaeal soil communities to an ongoing subterranean coal
mine fire. *The ISME Journal, 11*(6), 1447-1459. doi:10.1038/ismej.2017.1
Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *2011,
17*(1), 3. doi:10.14806/ej.17.1.200
Maude, R. (1996). *Seedborne diseases and their control: principles and practice*: CAB international.
McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. *PLOS ONE, 8*(4), e61217.
doi:10.1371/journal.pone.0061217
Mendes, R., Kruijt, M., de Bruijn, I., Dekkers, E., van der Voort, M., Schneider, J. H., . . . Raaijmakers, J. M.
(2011). Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science,
332*(6033), 1097-1100. doi:10.1126/science.1203980
Müller, S. H., & Pereira, P. A. A. (1995). Nitrogen fixation of common bean (Phaseolus vulgaris L.) as
affected by mineral nitrogen supply at different growth stages. *Plant and Soil, 177*(1), 55-61.
doi:10.1007/BF00010337
Naylor, D., & Coleman-Derr, D. (2018). Drought Stress and Root-Associated Bacterial Communities.
Frontiers in Plant Science, 8(2223). doi:10.3389/fpls.2017.02223
Nicholson, W. L., Munakata, N., Horneck, G., Melosh, H. J., & Setlow, P. (2000). Resistance of Bacillus
endospores to extreme terrestrial and extraterrestrial environments. *Microbiol Mol Biol Rev,
64*(3), 548-572. doi:10.1128/mmbr.64.3.548-572.2000
Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., . . . Wagner, H. (2019).
vegan: Community Ecology Package. R package version 2.5-6. Retrieved from https://CRAN.R-
project.org/package=vegan

Panke-Buisse, K., Poole, A. C., Goodrich, J. K., Ley, R. E., & Kao-Kniffin, J. (2015). Selection on soil
466   microbiomes reveals reproducible impacts on plant function. *The ISME Journal, 9*(4), 980-989.
467   doi:10.1038/ismej.2014.196

468   Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., . . . Glöckner, F. O. (2013). The SILVA
469   ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic*
470   *Acids Res, 41*(Database issue), D590-596. doi:10.1093/nar/gks1219

471   Rezki, S., Campion, C., Iacomi-Vasilescu, B., Preveaux, A., Toualbia, Y., Bonneau, S., . . . Barret, M. (2016).
472   Differences in stability of seed-associated microbial assemblages in response to invasion by
473   phytopathogenic microorganisms. *PeerJ, 4*, e1923-e1923. doi:10.7717/peerj.1923

474   Rezki, S., Campion, C., Simoneau, P., Jacques, M.-A., Shade, A., & Barret, M. (2017). Assembly of seed-
475   associated microbial communities within and across successive plant generations. *Plant and Soil,*
476   *422*, 1-13. doi:10.1007/s11104-017-3451-2

477   Ritpitakphong, U., Falquet, L., Vimoltust, A., Berger, A., Métraux, J.-P., & L'Haridon, F. (2016). The
478   microbiome of the leaf surface of Arabidopsis protects against a fungal pathogen. *New*
479   *Phytologist, 210*(3), 1033-1043. doi:10.1111/nph.13808

480   Rodríguez, C. E., Antonielli, L., Mitter, B., Trognitz, F., & Sessitsch, A. (2020). Heritability and Functional
481   Importance of the Setaria viridis Bacterial Seed Microbiome. *Phytobiomes Journal, 4*(1), 40-52.
482   doi:10.1094/pbiomes-04-19-0023-r

483   Sasse, J., Martinoia, E., & Northen, T. (2018). Feed Your Friends: Do Plant Exudates Shape the Root
484   Microbiome? *Trends Plant Sci, 23*(1), 25-41. doi:10.1016/j.tplants.2017.09.003

485   Shade, A., Jacques, M. A., & Barret, M. (2017). Ecological patterns of seed microbiome diversity,
486   transmission, and assembly. *Curr Opin Microbiol, 37*, 15-22. doi:10.1016/j.mib.2017.03.010

487   Vacher, C., Hampe, A., Porté, A. J., Sauer, U., Compant, S., & Morris, C. E. (2016). The Phyllosphere:
488   Microbial Jungle at the Plant–Climate Interface. *Annual Review of Ecology, Evolution, and*
489   *Systematics, 47*(1), 1-24. doi:10.1146/annurev-ecolsys-121415-032238

490   Větrovský, T., & Baldrian, P. (2013). The Variability of the 16S rRNA Gene in Bacterial Genomes and Its
491   Consequences for Bacterial Community Analyses. *PLOS ONE, 8*(2), e57923.
492   doi:10.1371/journal.pone.0057923

493   Walters, W., Hyde, E. R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A., . . . Knight, R. (2015).
494   Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker
495   Gene Primers for Microbial Community Surveys. *mSystems, 1*(1), e00009-00015.
496   doi:10.1128/mSystems.00009-15

497   Wang, Q., Garrity, G. M., Tiedje, J. M., & Cole, J. R. (2007). Naive Bayesian classifier for rapid assignment
498   of rRNA sequences into the new bacterial taxonomy. *Applied and environmental microbiology,*
499   *73*(16), 5261-5267. doi:10.1128/AEM.00062-07

500   Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., . . . Knight, R. (2017). Normalization
501   and microbial differential abundance strategies depend upon data characteristics. *Microbiome,*
502   *5*(1), 27. doi:10.1186/s40168-017-0237-y

503   Wilker, J., Navabi, A., Rajcan, I., Marsolais, F., Hill, B., Torkamaneh, D., & Pauls, K. P. (2019). Agronomic
504   Performance and Nitrogen Fixation of Heirloom and Conventional Dry Bean Varieties Under
505   Low-Nitrogen Field Conditions. *Frontiers in Plant Science, 10*(952). doi:10.3389/fpls.2019.00952

506   Xu, L., Naylor, D., Dong, Z., Simmons, T., Pierroz, G., Hixson, K. K., . . . Coleman-Derr, D. (2018). Drought
507   delays development of the sorghum root microbiome and enriches for monoderm bacteria. *Proc*
508   *Natl Acad Sci U S A, 115*(18), E4284-e4293. doi:10.1073/pnas.1717308115

509   Zaneveld, J. R., McMinds, R., & Vega Thurber, R. (2017). Stress and stability: applying the Anna Karenina
510   principle to animal microbiomes. *Nat Microbiol, 2*, 17121. doi:10.1038/nmicrobiol.2017.121

516   **Figure 2**

517
518

     Figure 3

Figure 4

A. Bacteria/archaea

B. Fungi

● Control   ▢ Water withholding   ▲ Nutrient addition