Title: **Abiotic treatment to common bean plants results in an altered endophytic seed microbiome**

Authors: A. Fina Bintarti[1,2], Patrick J. Kearns[2]*, Abby Sulesky-Grieb[2,3], and Ashley Shade[1,2,3]

1. Department of Plant, Soil and Microbial Sciences, Michigan State University, East Lansing MI

48824

2. The Plant Resilience Institute, Michigan State University, East Lansing MI 48824

3. Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing MI

48824

9
10 *current affiliation*: Department of Natural and Applied Sciences, Bentley University, Waltham, MA 02452
11

Abstract

There has been a growing interest in the seed microbiome due to its important role as an end

and starting point of plant microbiome assembly that can have consequences for plant health. However,

the effect of abiotic conditions on the seed microbial community remains unknown. We performed a

pilot study in a controlled growth chamber to investigate how the endophytic seed microbiome of the

common bean (*Phaseolus vulgaris L.* (var. Red Hawk)) was altered under abiotic treatments relevant for

crop management with changing climate. Bean plants were subjected to one of three treatments: 66%

water withholding to simulate mild drought, 50% Hoagland nutrient solution to simulate fertilization, or

control with sufficient water and baseline nutrition. We performed 16S rRNA gene amplicon sequencing

and ITS1 amplicon sequencing of the endophytic DNA to assess seed bacterial/archaeal and fungal

community structure, respectively. We found that variability in the seed microbiome structure was high

while alpha diversity was low, with tens of taxa present. Water withholding and nutrient addition

altered the seed microbiome structure for bacterial/archaeal communities as compared to the control,

and each treatment resulted in a distinct microbiome structure. There were no statistically supported

differences in the fungal microbiome across treatments. While we discuss several limitations of this

study, the promising results suggest that further investigation is needed to better understand abiotic or

stress-induced changes in the seed microbiome, the mechanisms that drive those changes, and their

implications for the health and stress responses of the next plant generation.

Importance

Seed microbiome members initiate the assembly of plant-associated microbial communities, but

the environmental drivers of endophytic seed microbiome composition are unclear. Here, we exposed

plants to short-term drought and fertilizer treatments during early vegetative growth and quantified the

microbiome composition of the seeds that were ultimately produced. We found that seeds produced by

plants stressed by water limitation or receiving nutrient addition had statistically different endophytic

bacterial/archaeal microbiome compositions from each other and from seeds produced by control

plants. This work suggests that the abiotic experience of a parental plant can influence the composition

of its seed microbiome, with unknown consequences for the next plant generation.

Key words

plant microbiome, 16S rRNA gene, ITS, drought, fertilizer, legume, growth chamber, endophyte, vertical

transmission, abiotic stress

Introduction

The plant microbiome includes bacteria, archaea, fungi, and viruses that associate with the plant

and inhabit different plant compartments, including the rhizosphere, phyllosphere, and endosphere (1).

The plant microbiome plays important roles for plant fitness, including nutrient acquisition (2),

secondary metabolite production (3), flowering time (4), and resistance to abiotic (5, 6) and biotic

stresses (7, 8). Plant microbiota can interact with each other as well as with the host plant, and the plant

is able to shape its microbiome structure and composition, for example, by producing root exudates or

allelochemicals (9, 10). Because of the close relationship between plants and their associated

microbiomes, it is likely that both or either could be affected by external stressors and, in turn, also

affect each other. These environmental stressors can include abiotic stress, such as changes in water

and nutrient availability (limitation or excess) or exposure to extreme temperatures, and biotic stress,

such as pathogen infection and herbivory. As plants respond to various stressors, their microbiome may

also be altered, either as a direct or indirect consequence of the stress. A recent study reported

increases in stress-related gene expression in the rhizosphere microbial community of plants treated

with high pH and high salinity wastewater, which suggested that the functional gene profile and

expression pattern of the plant microbiome under stressors can be used as an indicator tool to identify

stresses affecting host plants (11). Thus, both the environmental conditions and the host plant can act as

important filters that contribute to the ultimate composition of the plant microbiome (12, 13).

As a critical part of the plant microbiome, the seed microbiome can directly impact the seed and

seedling in ways that are important for crop establishment, such as by releasing the seed from

dormancy and promoting seed germination and seedling emergence (14, 15). However, there are

relatively fewer comprehensive studies of the seed microbiome as compared to studies of the

rhizosphere and phyllosphere microbiomes, which can acquire microbiota from the environment, (e.g.,

aerosols (12) and soils (16, 17)). Previously, investigation of seed microbiota employed culture-

dependent methods and focused on the transmission of plant pathogenic bacteria or fungi (18, 19).

More recently, seed microbiome studies have expanded and adopted cultivation-independent methods

(20-25). It is now recognized that the seed represents an endpoint of microbiome assembly for the

parental plant's reproductive compartment and the starting point of microbiome assembly for the new

seedling (26).

Vertical transmission of microbes via the seed has been reported for a variety of plant species,

as recently summarized (26) and reported (23, 27). Seeds acquire microbiota through different modes of

transmission, where early colonizers of the seed endophytic environment are acquired from the

parental plant either through the vascular system or floral stigma, while late colonizers are acquired on

external surfaces via seed contact with the environment (24, 28). A recent study of temporal dynamics

of the bean seed microbiome assembly reported that the vascular pathway is the dominant route for

seed microbiome transmission in common bean (24). Seed endophytes, acquired through the parental

plant vascular tissue, are of great interest because they are vertically transmitted to plant offspring, and

plants may preserve specific taxa through vertical transmission over generations. We hypothesized that

these preserved taxa may play vital roles in plant growth and tolerance to environmental stress. Thus, in

this study, we focused on seed endophytes and excluded seed epiphytes.

Managing or manipulating the plant microbiome is one promising strategy to support plant

tolerance to environmental stress. We are just beginning to understand how the plant microbiome

structure is altered during particular stresses (e.g., drought (29)), with a focus in the literature on the

root zone and rhizosphere microbiome. An important initial step of plant microbiome engineering to

enhance plant fitness and growth under environmental stresses is to understand the effect of these

stresses on the plant microbiome, and, next, to decipher underlying mechanisms involved in the

process. A previous study revealed that grass root microbiome diversity and structure was affected by

drought and there was enrichment for Actinobacteria (30). Drought resulted in reduced diversity of

sorghum root microbiome and increased abundance and activity of Gram-positive, and this shift was

correlated with altered plant metabolism and increased expression of bacterial ATP-binding cassette

transporter genes (29). Another recent study showed shifts in the wheat seed microbiome where

Actinobacteria were enriched and Gammaproteobacteria were depleted under drought conditions, and

these selected seed microbiome members demonstrated plant growth promoting ability on plants

undergoing drought (31). These studies that we highlight here are a few of many papers in this area of

the root and rhizosphere microbiome and its response to stress, but they do not investigate the seed

microbiome.

We conducted a controlled pilot study in an environmental growth chamber to determine

changes in the rhizosphere microbiome of the legume common bean under two different treatments of

water withholding and nutrient addition. The initial purpose of our pilot study was to identify members

of the root microbiome that were particularly resilient to either of these treatments. However, the

plants set pods at the end of the experiment, and we realized the opportunity to also assess the seed

microbiome of the treated plants as compared to control plants. The purpose of this brief report is to

share the surprising seed microbiome results from the pilot study, to discuss its limitations, and to

suggest immediate future directions based on the most promising results.

Methods

Plant growth conditions and harvest. Common bean seeds of the Red Hawk cultivar (32) were obtained

from a laboratory in the Michigan State University Plant Biology department. Seeds were surface-

sterilized in a 10% bleach solution followed by five rinses in sterile DI water. Three seeds per pot were

planted in 24 one-gallon pots filled with a steam-sterilized (~100°C) mixture of agricultural topsoil,

sphagnum peat and sand, and culled to one seedling per pot after the first unifoliate leaves had

emerged. The plants were grown in controlled conditions in a high-light BioChambers FLEX™ LED

growth chamber with a 16-hour day/8-hour night cycle at 26°C and 22°C, respectively. The plants were

divided into three groups: 8 control plants received ample water (300 mL every other day), 8 plants

were subjected to a mild "drought" during plant development, and received 66% less water (100 mL

every other day) (water withholding), and 8 plants received half strength Hoagland solution (300 mL

every other day) provided by the growth chamber facility (nutrient addition). Hoagland Solution details

can be found in Supplementary Table 1. The plants were grown for ~60 days until the R7 stage, when

plant pods were fully developed. All plants grew at relatively the same rate as expected for the cultivar,

so were harvested at the same time.

Harvesting was conducted by collecting the bean pods and plant biomass. Bean pods were

removed from the plants and the remaining above ground biomass from each plant was placed in a

paper bag and dried at 70°C for one week. The root system was gently pulled from the pot, cleaned of

excess soil with deionized water and dried at 70°C for one week. Once dried, the shoot and root dry

weight was measured for each plant. The remaining soil was collected for soil chemical analysis. One

hundred grams of each soil sample along with three replicates of pre-treatment bulk soil were sent to

the Michigan State University Soil and Plant Nutrient Laboratory (SPNL) for soil chemical testing. Soil

parameters including pH, lime index, phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg),

nitrate (NO_3^-), ammonium (NH_4^+), and organic matter (OM) were measured for all soil samples (testing

procedures available in Supplementary Methods, data available in Supplementary Table S2). Soil

chemistry differences among treatments were assessed using one-way analysis of variance (ANOVA).

The normality and homoscedasticity of the data were evaluated using Shapiro-Wilk and Levene's test,

respectively. We performed non-parametric Kruskal-Wallis test and post-hoc Dunn's test with

Benjamini-Hochberg correction for *p*-values if the ANOVA's assumptions were not met.

DNA extraction and amplicon sequencing. Twenty seeds from each plant were collected for DNA

extraction following the protocol of a previous study (20) with minor modifications to include surface

sterilization (33). Seeds were surfaced sterilized in 10% bleach, rinsed five times with sterile DI water,

and placed in sterile 50 mL centrifuge tubes with 30 mL of sterile 1X phosphate buffered saline (PBS)

with 0.05% Tween 20 and shaken at 140 rpm at room temperature for 4 hours. After shaking, tubes

were centrifuged at 500 x g for 15 minutes and the supernatant and seeds were discarded. The

remaining pellet was resuspended with 2 mL of sterile 1X PBS-Tween and transferred to a

microcentrifuge tube and spun at 20,000 x g for 10 min. The supernatant was discarded, and the pellet

was used for DNA extraction with the PowerSoil®DNA Isolation Kit (MoBio Laboratories, Solana Beach,

CA, United States) following manufacturer's instructions. DNA extracted from seed samples was

quantified with Qubit™dsDNA BR Assay Kit (ThermoFisher Scientific, Waltham, MA, United States) and

verified with Polymerase Chain Reaction (PCR), including a negative PCR control and blank extraction

reagents.

The bacterial/archaeal community PCR was conducted using the 515f (5'-

GTGCCAGCMGCCGCGGTAA-3') and 806r (5'-GGACTACHVGGGTWTCTAAT-3') primer pair (34) for

amplification of the V4 region of the 16S rRNA gene. The 16S rRNA gene amplification was conducted

under the following conditions: 94°C for 3 min, followed by 34 cycles of 94°C (45 s), 50°C (60 s), and 72°C

(90 s), with a final extension at 72°C (5 min). The amplification was performed in 25 µl mixtures

containing 12.5 µl GoTaq®Green Master Mix (Promega, Madison, WI, United States), 0.625 µl of each

primer (20mM), 1 µl of DNA template (final concentration of 0.02 - 0.626 ng per µl), and 4.5 µl nuclease

free water. Seed DNA (concentration range of 5 - 20 ng per µl) was sequenced at the Research

Technology Support Facility (RTSF) Genomics Core, Michigan State University sequencing facility using

the Illumina MiSeq platform.

Fungal communities were assessed using PCR amplification of the Internal Transcribed Spacer 1

(ITS1) region with the ITS1f (5⊠-CTTGGTCATTTAGAGGAAGTAA-3⊠) and ITS2 (5⊠-

GCTGCGTTCTTCATCGATGC-3⊠) primer pair (35) with the addition of index adapters as required by the

RTSF Genomics Core, (https://rtsf.natsci.msu.edu/genomics/sample-requirements/illumina-sequencing-

sample-requirements/). The PCR conditions of the ITS gene amplification conditions were as follows:

95°C for 5 min, followed by 30 cycles of 95°C (30 s), 54°C (45 s), and 72°C (90 s), with a final extension at

72°C (5 min). The amplification was performed in 50 µl mixtures containing 20 µl GoTaq®Green Master

Mix (Promega, Madison, WI, United States), 1 µl of each primer (20 mM), 4 µl of DNA template (final

concentration of 0.02 – 0.626 ng per µl), and 26 µl nuclease free water. The product of the ITS gene

amplification was cleaned and purified using the Wizard®SV Gel and PCR Clean-Up System (Promega,

Madison, WI, United States), following the manufacturer's protocol. Purified ITS gene amplification

products with the concentration range of 5 - 50 ng per µl were sequenced at the RTSF Genomics Core

using the Ilumina MiSeq platform. No amplification of the ITS gene was observed in one water

withholding and one nutrient addition seed sample, so only seven samples were sequenced for fungal

analysis in these treatment groups.

The 16S and ITS libraries were prepared by the sequencing facility using the Illumina

TruSeq®Nano DNA Library Prep Kit (Illumina, Inc., San Diego, CA, United States). Ilumina MiSeq was run

using a v2 Standard sequencing format with paired end reads (2 x 250 bp), and negative and mock

positive sequencing controls provided by the sequencing facility were included with each run.

Sequencing data analysis and OTU clustering. Bacterial/archaeal raw reads produced from Illumina

MiSeq were processed including merging the paired end reads, filtering the low-quality sequences,

dereplication to find unique sequence, singleton removal, denoising, and chimera checking using the

USEARCH pipeline (v.10.0.240) (36). Operational taxonomic unit (OTU) clustering was conducted using

an open reference strategy (37). First, closed reference OTU picking was performed at 97 % identity by

clustering quality filtered reads against the SILVA database (v.132) (38) using USEARCH algorithm (-

usearch_global command) (39). Reads that failed to match the SILVA reference were subsequently

clustered *de novo* at 97% identity using UPARSE-OTU algorithm (-*cluster_otus* command) (40). Closed

reference and *de novo* OTUs were combined into a full set of representative sequences, and then all

merged sequences were mapped back to that set using the -*usearch_global* command.

The set of representative sequences were aligned on QIIME 1.9.1 (41) using PyNAST (42) against

the SILVA (v.132) reference database. The unaligned OTU sequences were excluded from the OTU table

and the representative sequences. Taxonomic assignment was conducted on QIIME 1.9.1 using the

SILVA (v.132) database and the UCLUST default classifier at a minimum confidence of 0.9 (39). Plant

contaminants such as chloroplast and mitochondria; and unassigned taxa and sequences were removed

from the OTU table as well as the representative sequences using *filter_taxa_from_otu_table.py* and

filter_fasta.py command on QIIME. Rarefaction to the lowest sequencing depth (43, 44) (11,137

bacterial/archaeal reads) was conducted on QIIME.

The processing of fungal ITS raw reads was also conducted using the USEARCH (v.10.0.240)

pipeline. Read processing included paired end read merging, primer removal using cutadapt (v.2.0) (45),

filtering the low-quality sequences, and dereplication to find unique sequence. Operational taxonomic

unit clustering was conducted using an open reference OTU picking strategy. First, closed reference OTU

picking was performed by clustering quality filtered reads against the UNITE fungal ITS database (v.8.0)

(46) at 97% identity threshold using the USEARCH algorithm. Reads that failed to match the reference

were clustered *de novo* at 97% identity using the UPARSE-OTU algorithm. Closed reference and *de novo*

OTUs were combined into a full set of representative sequences, and then all merged sequences were

mapped back to that set using *-usearch_global* command. Fungal taxonomic classification was

performed in the CONSTAX tool (47) at a minimum confidence of 0.8 using the UNITE reference

database release 01-12-2017. Assigning taxonomy in CONSTAX was conducted using three classifiers,

including RDP Classifier (v.11.5) (48, 49), UTAX from USEARCH (v.8.1.1831) (40), and SINTAX from

USEARCH (v.9.2) (50). Any contaminants including mitochondria, chloroplast and other unwanted

lineages of eukaryotes were removed from the OTU table. Rarefaction was conducted to the lowest

number of sequences (21,329 fungal reads) on QIIME.

Microbial community analysis. Microbial community analyses were conducted in R (v.3.6.1) (R Core

Development Team). Microbial composition and relative abundance were analyzed using the Phyloseq

package (v.1.28.0) on R (51). Microbial richness (the number of taxa present) was calculated on the

rarefied OTU table using the vegan package (v.2.5-6) (52). The normality and homoscedasticity of the

data were tested using Saphiro-Wilk and Levene's test, respectively. The one-way ANOVA or non-

parametric Kruskal-Wallis test was then performed to analyze the data. Post hoc Dunn's test with false

discovery rate (FDR) correction using the Benjamini-Hochberg adjustment for multiple comparisons was

performed to compare plant biomass data among treatments.

Beta diversity was calculated on the rarefied OTU table with the vegan package using Jaccard

dissimilarity indices and visualized with a principal coordinate analysis (PCoA) plot. We used the Jaccard

index, which is based on presence-absence counts rather than relative abundance data, because we

reasoned that the seed microbiome members are unlikely to be actively growing inside the seed and

that any differences in relative abundances in the seed endophyte are unlikely attributable to

competitive growth outcomes *in situ*. Permutational multivariate analysis of variance (PERMANOVA)

using the function adonis (52) was performed to assess the effects of the treatments to the microbial

community structure. We performed multivariate analysis to check the homogeneity of dispersion

(variance) among groups using the function betadisper (52).

We analyzed shared microbial taxa between seeds from control and treated plants by

calculating the their occupancy (53). Microbial OTUs with occupancy value of 1 were those OTUs that

were detected in all samples from included treatments (control and nutrient addition, control and water

withholding).

Data and code availability. The computational workflows for sequence processing and ecological

statistics are available on GitHub (https://github.com/ShadeLab/BioRxiv_Seed_Microbiome_2020). Raw

sequence data of bacteria/archaea and fungi have been deposited in the Sequence Read Archive (SRA)

NCBI database under Bioproject accession number PRJNA635871.

Results

There were overall differences in plant biomass among treatments (Fig. 1, detailed Kruskal-

Wallis results can be found in Supplementary Table S3). Specifically, plants receiving nutrient addition

were larger in shoot and root biomass than control or mildly droughted plants (Fig. 1). Nutrient addition

plants also had higher pod number and pod mass compared to the water withholding and control plants

(Fig. 1). As expected, the addition of the Hoagland solution increased the nutrients available to the

plants in the nutrient addition treatment as compared to the control, as rhizosphere soil from the

nutrient addition treatment had higher phosphorus and potassium content than the other two

treatments,as well as higher nitrate content than the control treatment (Supplementary Figure S1,

Supplementary Table S4).

We removed less than 0.1 % of reads identified as plant and eukaryote contaminants from

bacterial/archaeal and fungal sequences, respectively. Analysis of contaminant-filtered

bacterial/archaeal and fungal sequences from seed samples resulted in a total of 81 and 226 OTUs (97%

sequence identity), respectively. Bacterial/archaeal communities in control, water withholding, and

nutrient addition seeds had different taxonomic compositions (Fig. 2A). Bacterial/archaeal communities

in the control seeds were almost exclusively dominated by the OTUs within the genus *Bacillus*, with a

mean relative genus-level abundance of more than 99%. Although the bacterial/archaeal community in

the water withholding and nutrient addition seeds were also dominated by *Bacillus*, genus-level

taxonomic diversity increased with the addition of other, non-dominating lineages. Specifically, seed

communities from water withholding and nutrient addition plants were also composed of *Virgibacillus*,

Pseudomonas, and several other bacterial/archaeal genera.

Similarly, different plant treatments resulted in different seed fungal community compositions.

Even though *Aspergillus* dominated the fungal community in the control and treated seeds, in the

treated seeds there was a shift to include other fungal taxa, including some identified as *Penicillium* and

Wallemia (Fig. 2B). These observations indicate that the seed microbiome is altered when parental

plants are exposed to abiotic stress or environmental alteration.

Analysis of overlapping taxa across treatments revealed that there were 4 and 3 bacterial taxa

shared between control and water withholding-treated seeds, and between control and nutrient

addition seeds, respectively (Supplementary Table S7). Bacterial taxa shared between samples belonged

exclusively to genus *Bacillus*. Fungal communities were dominated by genus *Aspergillus* that were

detected between control and water withholding-treated seeds, and *Penicillium* and *Aspergillus* that

were shared between control and nutrient addition seeds (Supplementary Table S8).

Because we do not expect microbiome members to be actively doubling inside the seed (and

therefore relativized abundances inside the seed to reflect fitness differences as an outcome of growth

therein, see (33) or a discussion of this), we used a presence-absence assessment (Jaccard index) of beta

diversity. There was a statistically supported differences in bacterial/archaeal microbiome composition

between treated seeds and control seeds (Fig. 3A, permutational multivariate analysis of variance,

PERMANOVA, F-stat = 4.73, R^2 = 0.31, P-val = 0.001). In contrast, there was no distinct clustering of

fungal communities associated with different treatments (Fig. 3B, PERMANOVA P-val > 0.05). These

results indicate that the abiotic treatments applied in this study altered the bacterial/archaeal, but not

fungal, community composition in the common bean seed. Also, there were differences in the

composition variability (multivariate dispersion) among treatments for bacterial/archaeal communities

(permutational analysis of multivariate dispersion, PERMDISP, F=7.553, P-val=0.003). Specifically, the

seeds from plants that experienced nutrient addition and water withholding had higher dispersion as

compared to seeds from control plants (TukeyHSD.betadisper, P-val=0.003 and 0.03, respectively).

Meanwhile, there were no differences in dispersion observed for fungal composition (PERMDISP,

F=0.491, P-val=0.628). This provides additional evidence that these abiotic treatments can lead to

increased variability in seed microbiome composition. Notably, PERMANOVA was found to be largely

unaffected by heterogeneity for balanced designs (54).

Discussion

The identities, functions and persistence of seed microbiome members are either not known or

not well-understood for many plant species of ecological or agricultural importance. This fundamental

information is the first step in understanding what constitutes a "typical" microbiome to initiate plant

microbiome assembly from seed. Changes to a typical seed microbiome because of parental plant

exposure to stress may have consequences for the health or resilience of the next plant generation. For

example, a depletion of beneficial members or enrichment of pathogens in the seed could disadvantage

the plant offspring, while an enrichment of beneficial members could provide a health advantage.

Therefore, observing a modification in the seed microbiome after parental exposure to stress has

potentially important implications and warrants report and continuation of research to understand any

consequences in functions for plant health.

We highlight three important observations from this study. First, treatment of the parent plant

altered the seed endophyte structure and composition compared to control plants, especially for the

bacterial/archaeal community. Although studies on the impact of drought on seed microbiota are still

new, especially as compared to studies on drought and the root-associated microbiome, our results are

consistent with a recent work which reported shifts in seed microbial communities of wheat under

drought conditions (31). Shifts in microbial communities have also been observed in various studies on

the root and rhizosphere of different plant species under drought stress (6, 55-57) with some reporting

that the effect of drought was stronger on the endophyte communities relative to rhizosphere

communities (30, 55, 57). This suggests an indirect effect of drought to endophyte communities through

physiological changes in drought-affected plants.

To our knowledge, this study is the first study assessing the impact of nutrient addition on seed

endophytes, as previous studies into the effect of fertilization have focused almost exclusively on the

rhizosphere and root-associated microbiome. We observed alterations in the seed endophyte

communities from nutrient addition parental plants. A previous study on the common bean rhizosphere

microbiome reported higher diversity in agricultural soil than native soil, suggesting that management

practices including fertilization may be a driver of the observed differences (58). Additionally, our recent

work on the biogeography of common bean rhizosphere microbiomes across different bean production

regions in the U.S revealed that fertilization differences between samples (including synthetic, organic

manure and no-input) had an explanatory value on the microbial community structure (59).Together,

these results suggest that any perturbations that affect the host plant may also affect its microbial

communities and these alterations may result in specific outcomes for plant stress tolerance and

resilience. Since the seed microbiomes contained a relatively simple community of tens to dozens of

taxa, even alterations in the composition or abundances of a few taxa may have consequences for

microbiome assembly of the next plant generation.

Our second key observation in this study is that bacterial/archaeal communities from the seeds

of treated plants had more variation as compared to the seeds from control plants. This suggests that

abiotic stress results in changes analogous to those observed during other types of microbiome

"dysbiosis" (aka Anna Karenina effects: higher variability across replicates, increased beta-dispersion,

and higher contribution of stochastic assembly processes (60)). As observed in dysbiosis of human or

animal microbiomes, the plant microbiome structure can be disrupted by disturbances and changing

environmental conditions that alter the composition and diversity of 'normal' microbiota (61, 62).

However, unlike dysbiosis in the human microbiome, any deviations from unaltered microbial

communities are not always associated with negative effects on plant health, and this alteration may be

necessary for maintaining plant health under stress conditions (63). Notably, there is no clear definition

of a "healthy" plant microbiome, though it has been suggested that high diversity and high evenness are

typical characteristics of healthy plant microbiomes and dysbiosis is correlated with reduced diversity

(64-66). However, we know that diversity metrics are aggregations of complex community data that are

intended to compare but ultimately directly cannot explain underlying biological mechanisms (67). We

speculate that the variations observed in the seed endophyte composition in our study, and the increase

in alpha and beta diversity of wheat seed microbiome of drought-exposed plants from another recent

study (31), are possibly due to selection of functionally beneficial microbial taxa. It has been

hypothesized that the selection and enrichment of particular taxa under drought stress is more likely

associated with functionality rather than taxonomy (56). There could be a microbial shift towards a state

associated with positive effects on plant health, as opposed dysbiosis (63)). However, the expectation of

high variability should be taken into consideration for future studies, as sufficient replication will be

needed to power statistical tests (e.g., (33)).

Third and finally, the fungal community was, on balance, stable relative to the bacterial/archaeal

community, suggesting that the persistence of fungal members is less sensitive to water withholding and

nutrient addition treatments. It has been reported that fungi can be very resistant to environmental

disturbances including drought due to several mechanisms including thick cell wall and osmolytes (68-

70). A previous study reported that drought had a more pronounced impact on bacteria than on fungal

networks in soil (71). Another study also reported that bacterial communities were significantly and

consistently altered not only in composition and diversity but also co-occurrence networks in forests

disturbed by clear-cutting or conversion to agricultural use relative to non-disturbed, meanwhile these

effects were marginal in fungal communities (72). All together, this suggests that fungal communities

are more stable over bacterial community under environmental changes.

We acknowledge that this is a pilot study and that these results are preliminary. We offer a

discussion of some of the major considerations and limitations in interpreting the results and for

planning future seed microbiome studies.

A first consideration is that there is an apparent maximum stress to plants that can be applied

when investigating its consequence for a seed microbiome. After stress exposure is released, plants

must be healthy enough to produce pods and seeds, and a balance must be achieved in which plants are

stressed but still able to fully mature. This constraint in stress exposure will never accommodate an

experimental design of severe or prolonged stress. However, the investigation of a mild or moderate

stress is still valuable because it is pertinent to agriculture. There are many situations in which non-

lethal stress occurs over part of a growing season, but then crops recover fully or partially to produce

some yield. Therefore, the result of mild or moderate stress for seed microbiomes has real-world

relevance.

Another consideration is the definition and directness of abiotic treatment, and whether an

abiotic treatment is expected to act on the plant, the microbiome, or both. In this pilot study, we applied

two different abiotic treatments: one that was expected to stress the plant directly (water withholding

to simulate mild drought) and one that was expected to weaken a legume's relationship with its root-

associated microbiome and symbiotic nitrogen fixers as nitrogen fixers are down-regulated by nitrogen

application (73, 74) (nutrient addition). Thus, the addition of nutrients was a benefit to the plant, rather

than a stress, as indicated by the increased root and shoot biomass. However, nutrient addition caused

a clear shift in the seed microbiome, demonstrating the potential of fertilizer use to have multi-

generational impacts on plant microbiome assembly. Therefore, management practices that advantage

the plant as far as yield and health in the short-term could have long-term consequences for plant-

microbiome relationships.

A clear limitation of the study is the substrate used for plant growth, which, with the microbes in

and on the original seeds, serves as a starting source for the assembly of the new plant's microbiome

(20, 26). For this pilot study, we used a sterilized mixture of agricultural topsoil, sphagnum peat and

sand provided by the growth chamber facility. The exact origin and physical/chemical characteristics of

the facility soil is unknown, and so it is unclear how representative this soil may be of agricultural field

soil. Additionally, the initial microbial community in the soil was not analyzed before planting the

common bean plants, so we cannot determine the origin of the observed microbial consortia in the

seeds and to what extent they overlap with the potting substrate. Previous work suggests that soil type

can have a large influence on the seed endophytic bacteria in rice (75), and this is likely also true for

other plant seeds. We observed a dominance of taxa from genus *Bacillus* in common bean seeds in all

three treatments, and *Bacillus* taxa have previously been reported to be enriched in the green bean

seedling (20). However, steam sterilization of the growth chamber soil may have killed many indigenous

microbial taxa in the soil that were not spore-formers or otherwise resistant to heat (and, bacilli are

known to be resistant to such treatments, as per (76)). Therefore, the microbial consortia available to

colonize the plant and seed microbiome from the sterilized soil was likely more limited as compared to

plants grown in the field or other substrates. We urge caution in generalizing from specific

compositional changes, but rather focus on the larger changes in beta-diversity and dispersion that were

consistently observed across very different abiotic treatments and may be more characteristic of seed

microbiome responses. Future work should focus efforts on using soil that is representative of the

typical agricultural environment of common bean, and the existing microbial community in the soil

should be sequenced prior to planting for comparison to the seed microbiome.

We included many standard controls in this study, including DNA extraction, PCR, and other

standard molecular biology controls. However, another limitation of this study the absence of negative

controls for DNA extraction that were followed the whole way through to sequencing. Seed endophytes

contain a very low total biomass of microbial cells. Here, we pooled twenty seeds to use for one

extraction to increase the microbial biomass yield for microbiome interrogation. We performed a buffer-

only control that was PCR-negative, but we did not save the material for sequencing, which would allow

for direct assessment of contaminants from the DNA extraction process. While the surface-sterilization

of the seeds prior to extraction and negative PCR controls provide confidence that the starting material

was not compromised and that we did not unintentionally amplify contaminants from the PCR reagents,

we cannot know if there were a few OTUs signatures from the extraction kit or buffer contaminants that

could have contributed to the observed seed microbiome composition (though we did not observe any

common contaminants reported in the literature). We now advocate for sequencing the DNA extraction

buffer control and using a package such as decontam (77) and microDecon (78) to ensure removal of

spurious contaminants, which are expected in low biomass samples (79).

A final minor limitation is in the choice of bacterial marker gene. We performed amplicon

sequencing of the 16S rRNA gene for the bacterial/archaeal community analysis in the seed. It is well

known that the variability of rRNA copy numbers among bacterial species can lead to an inflation of

species richness and obfuscate relative abundances of taxa (80). Moreover, the most resolved

taxonomic level achieved from 16S rRNA amplicon gene sequences often is genus, rather than species or

strain. Using a single-copy marker gene that has higher precision and sensitivity at the species level, like

the *gyr*B gene, may be a valuable alternative (20). The *gyr*B gene has been successfully applied to other

seed microbiome studies (20, 21, 81, 82). It may be valuable to consider use of both marker genes to the

same seed microbiome samples, so that taxonomic precision can be maximized and compared across

seeds with *gyrB*, while also maintaining an ability to source-track and compare composition across the

many plant-microbiome 16S rRNA amplicon datasets that have been deposited publicly.

In summary, while this pilot study provides a key insight into the response of the seed

microbiome structure to abiotic treatment in the host plant, there is much more work to be done. Next

steps include exposing the plants to more severe drought and nutrient excess conditions, quantifying

the physiological status of plants to determine their experience of stress, using representative field soil

for plant growth and assessing the field soil microbiome to deduce seed taxon origins, sequencing

negative controls from the DNA extractions to identify contaminants, and considering use of an

alternative marker gene for improved precision in microbial taxonomy and taxon abundances.

Despite noted considerations and limitations, we posit that this pilot study revealed an

important insight regarding how seed microbiomes may be altered after abiotic treatment of a plant.

Next, we need to understand the implications of this change for both the host plant and the microbial

community. An altered seed microbiome may have positive, negative, or entirely neutral outcomes for

the next plant generation. Additional work is needed to understand these outcomes over consecutive

plant generations to determine the effects on plant fitness and resilience. If positive or negative

outcomes are detected, this work opens a new direction of research that could spur exciting

applications in plant microbiome management.

Acknowledgements

This work was supported by the Michigan State University Plant Resilience Institute and by the USDA

National Institute of Food and Agriculture award USDA 2019-67019-29305. AFB acknowledges support

from the Fulbright Foreign Student Program and Office for International Students and Scholars (OISS),

Michigan State University; and AS acknowledges support from the USDA National Institute of Food and

Agriculture and Michigan AgBioResearch (Hatch). We thank Matthieu Barret and the Shade Lab for

helpful discussions.

454
455

Figure Legends

Fig 1. Plant aboveground (shoot) and belowground (root) biomass for control, water withholding, and nutrient addition treatments of common bean. Plant biomasses were calculated on eight plant replicates for each treatment. For each box plot, circles represent a single plant measurement within a treatment. The central horizontal lines represent the mean, the outer horizontal lines of the box represent the 25th and 75th percentiles. Boxes labelled with different letters were significantly different by a Kruskal-Wallis and post-hoc Dunn's test with a Benjamini-Hochberg false discovery rate correction (p-value significance ranges from <0.05 to <0.0001).

Fig 2. Mean relative abundances of genera of bacteria/archaea (A) and fungi (B) detected in the seed across control, water withholding and nutrient addition treatments. Each bar represents the endophytic microbiome identified in DNA extracted from 20 seeds collected from one plant replicate within a treatment. Bacterial/archaeal and fungal genera with mean relative abundances of less than 1 and 10 %, respectively, were grouped into the 'Other' classification, which includes many lineages (not monophyletic). Genera identified in the 'Other' classifications can be found in Supplementary Tables S5 and S6.

Fig. 3 Principal coordinate analysis (PCoA) plot of the bacterial/archaeal community (PERMANOVA, F-stat = 4.73, R^2 = 0.31, P-val = 0.001) (A) and fungal community (PERMANOVA P-val > 0.05) (B) in the common bean seed based on the Jaccard index. Symbol colors and shapes represent different abiotic treatments on the parent plant.

479   **References**

480   1.   Compant S, Samad A, Faist H, Sessitsch A. 2019. A review on the plant microbiome: ecology,
481   functions and emerging trends in microbial application. Journal of advanced research 19:29-97.
482   2.   Herrera Paredes S, Gao T, Law TF, Finkel OM, Mucyn T, Teixeira PJPL, Salas González I, Feltcher
483   ME, Powers MJ, Shank EA, Jones CD, Jojic V, Dangl JL, Castrillo G. 2018. Design of synthetic
484   bacterial communities for predictable plant phenotypes. PLOS Biology 16:e2003962.
485   3.   Gargallo-Garriga A, Sardans J, Pérez-Trujillo M, Guenther A, Llusià J, Rico L, Terradas J, Farré-
486   Armengol G, Filella I, Parella T. 2016. Shifts in plant foliar and floral metabolomes in response to
487   the suppression of the associated microbiota. BMC plant biology 16:78.
488   4.   Panke-Buisse K, Poole AC, Goodrich JK, Ley RE, Kao-Kniffin J. 2015. Selection on soil microbiomes
489   reveals reproducible impacts on plant function. The ISME Journal 9:980-989.
490   5.   Lau JA, Lennon JT. 2012. Rapid responses of soil microorganisms improve plant fitness in novel
491   environments. Proceedings of the National Academy of Sciences 109:14058-14062.
492   6.   Naylor D, Coleman-Derr D. 2018. Drought Stress and Root-Associated Bacterial Communities.
493   Frontiers in Plant Science 8.
494   7.   Mendes R, Kruijt M, de Bruijn I, Dekkers E, van der Voort M, Schneider JH, Piceno YM, DeSantis
495   TZ, Andersen GL, Bakker PA, Raaijmakers JM. 2011. Deciphering the rhizosphere microbiome for
496   disease-suppressive bacteria. Science 332:1097-100.
497   8.   Ritpitakphong U, Falquet L, Vimoltust A, Berger A, Métraux J-P, L'Haridon F. 2016. The
498   microbiome of the leaf surface of Arabidopsis protects against a fungal pathogen. New
499   Phytologist 210:1033-1043.
500   9.   Sasse J, Martinoia E, Northen T. 2018. Feed Your Friends: Do Plant Exudates Shape the Root
501   Microbiome? Trends Plant Sci 23:25-41.
502   10.   Jones P, Garcia BJ, Furches A, Tuskan GA, Jacobson D. 2019. Plant Host-Associated Mechanisms
503   for Microbial Selection. Frontiers in Plant Science 10.
504   11.   Zolti A, Green SJ, Sela N, Hadar Y, Minz D. 2020. The microbiome as a biosensor: functional
505   profiles elucidate hidden stress in hosts. Microbiome 8:71.
506   12.   Vacher C, Hampe A, Porté AJ, Sauer U, Compant S, Morris CE. 2016. The Phyllosphere: Microbial
507   Jungle at the Plant–Climate Interface. Annual Review of Ecology, Evolution, and Systematics
508   47:1-24.
509   13.   Hacquard S, Spaepen S, Garrido-Oter R, Schulze-Lefert P. 2017. Interplay Between Innate
510   Immunity and the Plant Microbiota. Annu Rev Phytopathol 55:565-589.
511   14.   Goggin DE, Emery RJN, Kurepin LV, Powles SB. 2015. A potential role for endogenous microflora
512   in dormancy release, cytokinin metabolism and the response to fluridone in Lolium rigidum
513   seeds. Annals of botany 115:293-301.
514   15.   Lamichhane JR, Debaeke P, Steinberg C, You MP, Barbetti MJ, Aubertot J-N. 2018. Abiotic and
515   biotic factors affecting crop seed germination and seedling emergence: a conceptual framework.
516   Plant and Soil 432:1-28.
517   16.   Truyens S, Weyens N, Cuypers A, Vangronsveld J. 2015. Bacterial seed endophytes: genera,
518   vertical transmission and interaction with plants. Environmental Microbiology Reports 7:40-50.
519   17.   Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA,
520   Sundaresan V. 2015. Structure, variation, and assembly of the root-associated microbiomes of
521   rice. Proceedings of the National Academy of Sciences 112:E911-E920.
522   18.   Nelson EB. 2018. The seed microbiome: Origins, interactions, and impacts. Plant and Soil 422:7-
523   34.
524   19.   Wassermann B, Cernava T, Müller H, Berg C, Berg G. 2019. Seeds of native alpine plants host
525   unique microbial communities embedded in cross-kingdom networks. Microbiome 7:108.

20. Barret M, Briand M, Bonneau S, Préveaux A, Valière S, Bouchez O, Hunault G, Simoneau P,
Jacques M-A. 2015. Emergence Shapes the Structure of the Seed Microbiota. Applied and
Environmental Microbiology 81:1257-1266.
21. Rezki S, Campion C, Simoneau P, Jacques M-A, Shade A, Barret M. 2017. Assembly of seed-
associated microbial communities within and across successive plant generations. Plant and Soil
422:1-13.
22. Torres-Cortés G, Bonneau S, Bouchez O, Genthon C, Briand M, Jacques M-A, Barret M. 2018.
Functional Microbial Features Driving Community Assembly During Seed Germination and
Emergence. Frontiers in Plant Science 9.
23. Rodríguez CE, Antonielli L, Mitter B, Trognitz F, Sessitsch A. 2020. Heritability and Functional
Importance of the Setaria viridis Bacterial Seed Microbiome. Phytobiomes Journal 4:40-52.
24. Chesneau G, Torres-Cortes G, Briand M, Darrasse A, Preveaux A, Marais C, Jacques M-A, Shade
A, Barret M. 2020. Temporal dynamics of bacterial communities during seed development and
maturation. FEMS Microbiology Ecology 96.
25. Bziuk N, Maccario L, Straube B, Wehner G, Sørensen SJ, Schikora A, Smalla K. 2021. The treasure
inside barley seeds: microbial diversity and plant beneficial bacteria. Environmental Microbiome
16:20.
26. Shade A, Jacques MA, Barret M. 2017. Ecological patterns of seed microbiome diversity,
transmission, and assembly. Curr Opin Microbiol 37:15-22.
27. Bergna A, Cernava T, Rändler M, Grosch R, Zachow C, Berg G. 2018. Tomato Seeds Preferably
Transmit Plant Beneficial Endophytes. Phytobiomes Journal 2:183-193.
28. Maude R. 1996. Seedborne diseases and their control: principles and practice. CAB international.
29. Xu L, Naylor D, Dong Z, Simmons T, Pierroz G, Hixson KK, Kim YM, Zink EM, Engbrecht KM, Wang
Y, Gao C, DeGraaf S, Madera MA, Sievert JA, Hollingsworth J, Birdseye D, Scheller HV, Hutmacher
R, Dahlberg J, Jansson C, Taylor JW, Lemaux PG, Coleman-Derr D. 2018. Drought delays
development of the sorghum root microbiome and enriches for monoderm bacteria. Proc Natl
Acad Sci U S A 115:E4284-e4293.
30. Naylor D, DeGraaf S, Purdom E, Coleman-Derr D. 2017. Drought and host selection influence
bacterial community dynamics in the grass root microbiome. The ISME Journal 11:2691-2704.
31. Hone H, Mann R, Yang G, Kaur J, Tannenbaum I, Li T, Spangenberg G, Sawbridge T. 2021.
Profiling, isolation and characterisation of beneficial microbes from the seed microbiomes of
drought tolerant wheat. Scientific Reports 11:11916.
32. Kelly JD, Hosfield GL, Varner GV, Uebersax MA, Long RA, Taylor J. 1998. Registration of 'Red
Hawk' Dark Red Kidney Bean. Crop Science 38:cropsci1998.0011183X003800010056x.
33. Bintarti AF, Sulesky-Grieb A, Stopnišek N, Shade A. 2021. Endophytic microbiome variation
among single plant seeds. Phytobiomes Journal 0:null.
34. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Lozupone CA, Turnbaugh PJ, Fierer N, Knight
R. 2011. Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
Proceedings of the National Academy of Sciences 108:4516-4522.
35. Smith DP, Peay KG. 2014. Sequence Depth, Not PCR Replication, Improves Ecological Inference
from Next Generation DNA Sequencing. PLOS ONE 9:e90234.
36. Edgar RC, Flyvbjerg H. 2015. Error filtering, pair assembly and error correction for next-
generation sequencing reads. Bioinformatics 31:3476-82.
37. Lee S-H, Sorensen JW, Grady KL, Tobin TC, Shade A. 2017. Divergent extremes but convergent
recovery of bacterial and archaeal soil communities to an ongoing subterranean coal mine fire.
The ISME Journal 11:1447-1459.

38. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The
SILVA ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res 41:D590-6.
39. Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. Bioinformatics
26:2460-1.
40. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nature
Methods 10:996-998.
41. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG,
Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA,
McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA,
Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010. QIIME allows analysis of high-throughput
community sequencing data. Nature Methods 7:335-336.
42. Caporaso JG, Bittinger K, Bushman FD, DeSantis TZ, Andersen GL, Knight R. 2010. PyNAST: a
flexible tool for aligning sequences to a template alignment. Bioinformatics 26:266-7.
43. Gihring TM, Green SJ, Schadt CW. 2012. Massively parallel rRNA gene sequencing exacerbates
the potential for biased community diversity comparisons due to variable library sizes.
Environmental Microbiology 14:285-290.
44. Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, Lozupone C, Zaneveld JR, Vázquez-
Baeza Y, Birmingham A, Hyde ER, Knight R. 2017. Normalization and microbial differential
abundance strategies depend upon data characteristics. Microbiome 5:27.
45. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 17:3.
46. Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AFS, Bahram M, Bates ST, Bruns TD,
Bengtsson-Palme J, Callaghan TM, Douglas B, Drenkhan T, Eberhardt U, Dueñas M, Grebenc T,
Griffith GW, Hartmann M, Kirk PM, Kohout P, Larsson E, Lindahl BD, Lücking R, Martín MP,
Matheny PB, Nguyen NH, Niskanen T, Oja J, Peay KG, Peintner U, Peterson M, Põldmaa K, Saag L,
Saar I, Schüßler A, Scott JA, Senés C, Smith ME, Suija A, Taylor DL, Telleria MT, Weiss M, Larsson
K-H. 2013. Towards a unified paradigm for sequence-based identification of fungi. Molecular
Ecology 22:5271-5277.
47. Gdanetz K, Benucci GMN, Vande Pol N, Bonito G. 2017. CONSTAX: a tool for improved
taxonomic resolution of environmental fungal ITS sequences. BMC Bioinformatics 18:538.
48. Wang Q, Garrity GM, Tiedje JM, Cole JR. 2007. Naive Bayesian classifier for rapid assignment of
rRNA sequences into the new bacterial taxonomy. Applied and environmental microbiology
73:5261-5267.
49. Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, Brown CT, Porras-Alfaro A, Kuske CR,
Tiedje JM. 2014. Ribosomal Database Project: data and tools for high throughput rRNA analysis.
Nucleic Acids Res 42:D633-42.
50. Edgar RC. 2016. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences.
bioRxiv doi:10.1101/074161:074161.
51. McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. PLOS ONE 8:e61217.
52. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB,
Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: Community Ecology
Package. R package version 2.5-6.
53. Shade A, Stopnisek N. 2019. Abundance-occupancy distributions to prioritize plant core
microbiome membership. Current Opinion in Microbiology 49:50-58.

54. Anderson MJ, Walsh DCI. 2013. PERMANOVA, ANOSIM, and the Mantel test in the face of
heterogeneous dispersions: What null hypothesis are you testing? Ecological Monographs
83:557-574.
55. Santos-Medellín C, Liechty Z, Edwards J, Nguyen B, Huang B, Weimer BC, Sundaresan V. 2021.
Prolonged drought imparts lasting compositional changes to the rice root microbiome. Nature
Plants 7:1065-1077.
56. Jochum MD, McWilliams KL, Pierson EA, Jo Y-K. 2019. Host-mediated microbiome engineering
(HMME) of drought tolerance in the wheat rhizosphere. PLOS ONE 14:e0225933.
57. Fitzpatrick CR, Copeland J, Wang PW, Guttman DS, Kotanen PM, Johnson MTJ. 2018. Assembly
and ecological function of the root microbiome across angiosperm plant species. Proceedings of
the National Academy of Sciences 115:E1157-E1165.
58. Pérez-Jaramillo JE, de Hollander M, Ramírez CA, Mendes R, Raaijmakers JM, Carrión VJ. 2019.
Deciphering rhizosphere microbiome assembly of wild and modern common bean (Phaseolus
vulgaris) in native and agricultural soils from Colombia. Microbiome 7:114.
59. Stopnisek N, Shade A. 2021. Persistent microbiome members in the common bean rhizosphere:
an integrated analysis of space, time, and plant genotype. The ISME Journal 15:2708-2722.
60. Zaneveld JR, McMinds R, Vega Thurber R. 2017. Stress and stability: applying the Anna Karenina
principle to animal microbiomes. Nat Microbiol 2:17121.
61. Liu H, Brettell LE, Singh B. 2020. Linking the Phyllosphere Microbiome to Plant Health. Trends in
Plant Science 25:841-844.
62. Berg G, Rybakova D, Fischer D, Cernava T, Vergès M-CC, Charles T, Chen X, Cocolin L, Eversole K,
Corral GH, Kazou M, Kinkel L, Lange L, Lima N, Loy A, Macklin JA, Maguin E, Mauchline T,
McClure R, Mitter B, Ryan M, Sarand I, Smidt H, Schelkle B, Roume H, Kiran GS, Selvin J, Souza
RSCd, van Overbeek L, Singh BK, Wagner M, Walsh A, Sessitsch A, Schloter M. 2020. Microbiome
definition re-visited: old concepts and new challenges. Microbiome 8:103.
63. Paasch BC, He SY. 2021. Toward understanding microbiota homeostasis in the plant kingdom.
PLOS Pathogens 17:e1009472.
64. Berg G, Kusstatscher P, Abdelfattah A, Cernava T, Smalla K. 2021. Microbiome Modulation-
Toward a Better Understanding of Plant Microbiome Response to Microbial Inoculants. Front
Microbiol 12:650610.
65. Berg G, Köberl M, Rybakova D, Müller H, Grosch R, Smalla K. 2017. Plant microbial diversity is
suggested as the key to future biocontrol and health trends. FEMS Microbiology Ecology 93.
66. Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. 2020. Plant–microbiome interactions: from
community assembly to plant health. Nature Reviews Microbiology 18:607-621.
67. Shade A. 2017. Diversity is the question, not the answer. The ISME journal 11:1-6.
68. Schimel J, Balser TC, Wallenstein M. 2007. Microbial stress-response physiology and its
implications for ecosystem function. Ecology 88:1386-94.
69. Treseder KK, Lennon JT. 2015. Fungal Traits That Drive Ecosystem Dynamics on Land.
Microbiology and Molecular Biology Reviews 79:243-262.
70. Treseder KK, Berlemont R, Allison SD, Martiny AC. 2018. Drought increases the frequencies of
fungal functional genes related to carbon and nitrogen acquisition. PloS one 13:e0206441-
e0206441.
71. de Vries FT, Griffiths RI, Bailey M, Craig H, Girlanda M, Gweon HS, Hallin S, Kaisermann A, Keith
AM, Kretzschmar M, Lemanceau P, Lumini E, Mason KE, Oliver A, Ostle N, Prosser JI, Thion C,
Thomson B, Bardgett RD. 2018. Soil bacterial networks are less stable under drought than fungal
networks. Nature Communications 9:3033.

72.   Osburn ED, McBride SG, Aylward FO, Badgley BD, Strahm BD, Knoepp JD, Barrett JE. 2019. Soil
665   Bacterial and Fungal Communities Exhibit Distinct Long-Term Responses to Disturbance in
666   Temperate Forests. Frontiers in Microbiology 10.
667   73.   Müller SH, Pereira PAA. 1995. Nitrogen fixation of common bean (Phaseolus vulgaris L.) as
668   affected by mineral nitrogen supply at different growth stages. Plant and Soil 177:55-61.
669   74.   Wilker J, Navabi A, Rajcan I, Marsolais F, Hill B, Torkamaneh D, Pauls KP. 2019. Agronomic
670   Performance and Nitrogen Fixation of Heirloom and Conventional Dry Bean Varieties Under
671   Low-Nitrogen Field Conditions. Frontiers in Plant Science 10.
672   75.   Hardoim PR, Hardoim CCP, van Overbeek LS, van Elsas JD. 2012. Dynamics of seed-borne rice
673   endophytes on early plant growth stages. PloS one 7:e30438-e30438.
674   76.   Nicholson WL, Munakata N, Horneck G, Melosh HJ, Setlow P. 2000. Resistance of Bacillus
675   endospores to extreme terrestrial and extraterrestrial environments. Microbiol Mol Biol Rev
676   64:548-72.
677   77.   Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. 2018. Simple statistical
678   identification and removal of contaminant sequences in marker-gene and metagenomics data.
679   Microbiome 6:226.
680   78.   McKnight DT, Huerlimann R, Bower DS, Schwarzkopf L, Alford RA, Zenger KR. 2019. microDecon:
681   A highly accurate read-subtraction tool for the post-sequencing removal of contamination in
682   metabarcoding studies. Environmental DNA 1:14-25.
683   79.   Eisenhofer R, Minich JJ, Marotz C, Cooper A, Knight R, Weyrich LS. 2019. Contamination in Low
684   Microbial Biomass Microbiome Studies: Issues and Recommendations. Trends Microbiol 27:105-
685   117.
686   80.   Větrovský T, Baldrian P. 2013. The Variability of the 16S rRNA Gene in Bacterial Genomes and Its
687   Consequences for Bacterial Community Analyses. PLOS ONE 8:e57923.
688   81.   Rezki S, Campion C, Iacomi-Vasilescu B, Preveaux A, Toualbia Y, Bonneau S, Briand M, Laurent E,
689   Hunault G, Simoneau P, Jacques M-A, Barret M. 2016. Differences in stability of seed-associated
690   microbial assemblages in response to invasion by phytopathogenic microorganisms. PeerJ
691   4:e1923-e1923.
692   82.   Rochefort A, Briand M, Marais C, Wagner M-H, Laperche A, Vallée P, Barret M, Sarniguet A.
693   2019. Influence of Environment and Host Plant Genotype on the Structure and Diversity of the
694   Brassica napus Seed Microbiota. Phytobiomes Journal 3:326-336.
695

Figure 1

Figure 2

Figure 3