AIM2 Inflammasome Mediates Hallmark Neuropathological Alterations and Cognitive Impairment in a Mouse Model of Vascular Dementia

Luting Poh[1], David Y. Fann[1,2,*], Peiyan Wong[3,4], Hong Meng Lim[1], Sok Lin Foo[1], Sung-Wook Kang[1], Vismitha Rajeev[1], Sharmelee Selvaraji[1], Rajagopal Iyer Vinaya[3], Nageiswari Parathy[3], Mohammad Badruzzaman Khan[5], Dong-Gyu Jo[6], Grant R. Drummond[7], Christopher G. Sobey[7], Mitchell K.P. Lai[3,8], Christopher Li-Hsian Chen[3,8], Lina H. K. Lim[1], Thiruma V. Arumugam[1,6,7,*]

[1]Department of Physiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[2]Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[3]Department of Pharmacology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore
[4]Neuroscience and Behavioural Disorders Programme, Duke-NUS Medical School, Singapore
[5]Department of Neurology, Medical College of Georgia, Augusta University, Augusta, GA, USA
[6]School of Pharmacy, Sungkyunkwan University, Suwon, Republic of Korea
[7]Department of Physiology, Anatomy and Microbiology, La Trobe University, Bundoora, VIC, Australia
[8]Memory, Aging and Cognition Centre, National University Health System, Singapore

*Correspondence: Thiruma V. Arumugam, Department of Physiology, Anatomy and Microbiology, La Trobe University, Bundoora, VIC, Australia. E-mail: g.arumugam@latrobe.edu.au; David Y. Fann, Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, Singapore. E-mail: bchfywd@nus.edu.sg

Abstract

Chronic cerebral hypoperfusion is associated with vascular dementia (VaD). Cerebral hypoperfusion may initiate complex molecular and cellular inflammatory pathways that contribute to long term cognitive impairment and memory loss. Here we used a bilateral common carotid artery stenosis (BCAS) mouse model of VaD to investigate its effect on the innate immune response – particularly the inflammasome signaling pathway. Comprehensive analyses revealed that chronic cerebral hypoperfusion induces a complex temporal expression and activation of inflammasome components and their products (IL-1β and IL-18) in different brain regions, and promotes activation of apoptotic and pyroptotic cell death pathways. Glial cell activation, white matter lesion formation and hippocampal neuronal loss also occurred in a spatiotemporal manner. Moreover, in AIM2 knockout mice we observed attenuated inflammasome-mediated production of proinflammatory cytokines, apoptosis and pyroptosis, as well as resistance to chronic microglial activation, myelin breakdown, hippocampal neuronal loss, and behavioural and cognitive deficits following BCAS. Hence, we have demonstrated that activation of the AIM2 inflammasome substantially contributes to the pathophysiology of chronic cerebral hypoperfusion-induced brain injury and may therefore represent a promising therapeutic target for attenuating cognitive impairment in VaD.

Introduction

Vascular dementia (VaD) is the second most common form of dementia that is associated with vascular diseases resulting in dysfunction and damage to the cerebral vasculature, which causes disruption of cerebral blood flow, brain injury, and ultimately cognitive impairment and memory loss[1]. Clinical findings in VaD include chronic cerebral hypoperfusion, white matter lesions (WMLs), strokes, hippocampal atrophy, decline in executive functioning and memory impairment[2]. Recent evidence indicates that chronic cerebral hypoperfusion is associated with systemic inflammation before WMLs and neurological symptoms develop in VaD[3-5]. Interleukin 1-beta (IL-1β), a potent cytokine that orchestrates inflammatory pathways, is likely to be a key contributor to VaD[5]. The primary molecular machinery for IL-1β production in response involves assembly and activation of the inflammasome.

Inflammasome is an intracellular multi-protein complex that initiates an innate immune response, and is involved in multiple acute and chronic neurological diseases such as ischemic stroke, Alzheimer's disease (AD), Parkinson's Disease (PD), and amyotrophic lateral sclerosis (ALS)[6-11]. Upon inflammasome assembly and activation, activity of cleaved caspase-1 is increased and downstream proinflammatory cytokines such as mature IL-1β and IL-18 are generated[6]. This leads to activation of glial cells, endothelial dysfunction, and oligodendrocyte injury and demyelination, eventually resulting in neurovascular unit disintegration, neuronal loss and brain circuit dysfunction. Cleaved caspase-1 may also promote cell death by activating apoptosis and pyroptosis pathways[12-13]. Inhibition of caspase-1 can prevent pyroptosis in experimental models of multiple sclerosis, reducing demyelination and neurodegeneration[14]. The inflammasome complex also contributes to cognitive decline in APP/PS1 mice[9,15]. Deletion of the AIM2 inflammasome in that model promoted spatial memory, synaptic plasticity and dendrite branching[16]. While evidence for direct involvement of the inflammasome complex in VaD is lacking, a cytokine profile of plasma from VaD patients found IL-1β to be the most abundant[17-20]. It was also reported that IL-1β impeded oligodendrocyte recruitment and inhibited white matter repair and functional recovery during the early stages of chronic cerebral hypoperfusion[5]. However, any potential involvement of inflammasome activation in chronic cerebral hypoperfusion-induced glial activation, white matter damage, neuronal cell death and cognitive dysfunction has not been studied.

Here, we have comprehensively assessed the activation of inflammasome signaling in the bilateral common carotid artery stenosis (BCAS) mouse model of VaD. We report an increased expression and activation of inflammasome complexes, production of mature proinflammatory cytokines, glial activation, WMLs, and neuronal cell death occurring in a time-dependent manner in the cerebral cortex, hippocampus and striatum following BCAS. Notably, these brain regions exhibited distinct profiles of inflammasome expression and signaling. We also demonstrate activation and a functional role of the AIM2 inflammasome in glial activation, WML formation and neuronal cell death. Furthermore, genetic deletion of AIM2 protects against BCAS-

107 induced inflammasome activation, proinflammatory cytokine production, WML and
108 neuronal loss, and improves cognitive outcome. Collectively, our findings reveal that
109 the AIM2 inflammasome may be a potential therapeutic target in VaD.
110
111

Materials and Methods

Experimental Animals and Genotyping

All *in vivo* experimental procedures were approved by the National University of Singapore, Singapore Animal Care and Use Committee and performed according to the guidelines set forth by the National Advisory Committee for Laboratory Animal Research (NACLAR), Singapore. Mice were housed in individual cages under standard laboratory conditions. All efforts were made to minimize suffering and numbers of animals used. All sections of the manuscript were performed and reported in accordance with ARRIVE (Animal Research: Reporting In Vivo Experiments) guidelines. Fourteen to sixteen weeks old wild type (WT) male C57BL/6 mice (weighing 24 to 30 grams) were obtained from In Vivos, Singapore. The Aim2 knockout (AIM2 KO) mice were generated by Ingenious Targeting Laboratory (Ronkonkoma, NY) on the C57BL/6 background via targeted replacement of the Aim2 coding region with a neomycin resistance gene through homologous recombination and obtained from our collaborator Professor Jenny P. Y. Ting[21]. Primer sequences used for PCR-based Aim2 deficient mouse genotyping are as follows: neomycin (KO) forward, GGAACTTCGCTAGACTAGTACGCGTG; neomycin (KO) reverse, CAACATTGTACAGATTGAGCAGG; AIM2 (WT) forward, GATGGAGAGTGAGTACCGGGAAATGCTGTT; and AIM2 (WT) reverse, TCTGCAAGTAGATTGGAGACAGACTCTGGTGA, resulting in a 450-bp band for the wild type and a 250-bp band for the targeted Aim2 knockout when PCR analysis is performed on the genomic DNA (Supplementary Fig. 1). All mice were given free access to food and water *ad libitum*. The experimental groups consist of 20-25 animals in each group.

Bilateral Common Carotid Artery Stenosis (BCAS) Mouse Model

Animals were anaesthetized with isoflurane and subjected to BCAS, and subsequent chronic hypoperfusion using microcoils specially designed for the mouse

(microcoil specifications: piano wire diameter 0.08mm, internal diameter 0.18mm,
coiling pitch 0.5mm, and total length 2.5mm; Sawane Spring Co Ltd, Japan) was
described elsewhere[22]. The left and right common carotid arteries (CCAs) were
exposed one-by-one, freed from their sheaths, and a microcoil was twined by rotating
it around each CCA. The site of surgery was closed, and the mice were observed
and taken care of post-surgery until conscious and recovered to freely access food
and water ad libitum. To study the disease progression, the experimental animals
were divided into eight time-point groups: Sham, 1, 3, 7, 15, 21, 30 and 60-days
BCAS. Sham animals were given a skin incision and their CCAs were exposed. All
animals were euthanized at their respective end-point after BCAS for subsequent
analysis. For the molecular work involving AIM2 KO and Wild Type control, animals
were divided into 3 experimental groups: Sham, 15 and 30-days BCAS. However,
only the Sham and 30-days timepoints were selected to challenge the cognitive
ability of both WT and AIM2 KO mice during behavioral analysis.

Measurements of Cerebral Blood Flow by Laser Speckle Contrast Imager

High-resolution Laser Speckle Contrast Imager (PSI system, Perimed Inc.) was
used to image cerebral blood perfusion and record cerebral blood flow (CBF) before
BCAS (baseline), immediately after BCAS surgery and finally at their respective end-
points of BCAS. As shown in Supplementary Fig. 2, regions of interest (ROIs)
between the bregma and lambda were selected for overall perfusion in the area of
two hemispheres. Body temperature was maintained at 37 ± 0.5 °C, and the skull
was shaved and exposed by a midline skin incision. The skull was cleaned gently
with sterile phosphate buffered saline (PBS) using a cotton applicator. Finally, the
image area was kept moist and a non-toxic silicon oil was applied on the skull, which
improved imaging. Perfusion images were acquired using the PSI system with a 70
mW built-in laser diode for illumination and a 1388 x 1038 pixels CCD camera
installed 10 cm above the skull (speed 19 Hz, and exposure time 6 mSec). Acquired
images were analyzed for changes in CBF (cerebral perfusion) using a dedicated
PIMSoft program (Perimed Inc.).

Sample Collection and Processing

Following each time points, mice were euthanized by administering a lethal dose
of inhaled carbon dioxide (CO_2) and the brains harvested. The different brain areas

(cerebral cortex, hippocampus and striatum) were immediately separated on ice and
were frozen in dry ice for tissue biochemical analysis (n = 7-8 in each experimental
group). A separate group of animals were sacrificed for histological analysis. Mice
were deeply anaesthetized, and perfusion fixation though the heart was performed
with 25 mL of chilled 1 x PBS (pH 7.4) then followed by 25 mL chilled
paraformaldehyde (4%). Once the perfusion was completed, the brains were
harvested and placed in vials containing 4% paraformaldehyde solution for
immersion-fixation overnight at 4°C (n = 5-7 in each experimental group).

Immunoblot Analysis

Immunoblot analysis was performed as described by us previously[23]. Briefly, brain
tissues were homogenized in lysis buffer and then combined with 2 x Laemelli buffer
(Bio-Rad Laboratories, Inc., Hercules, CA, USA). Protein samples were then
separated on 7.5 to 12.5% v/v sodium dodecyl sulfate (SDS) gels. The SDS-PAGE
gels were transferred onto nitrocellulose membranes to probe for proteins. Next, the
nitrocellulose membranes were incubated with the following primary antibodies:
NLRP1 (Santa Cruz, sc390133), NLRP3 (Adipogen, AG20B0014), NLRC4 (Milipore,
06-1125), AIM2 (Cell Signaling, #13095), IL-1β (Genetex, GTX74034), IL-18
(Biovision, 5180R-100), Total Caspase-1 and Cleaved Caspase-1 (p33 subunit)
(Adipogen, AG20B0042), Cleaved Caspase-1 (p20 subunit) (Cell Signaling, #67314),
Casapse-8 (Cell Signaling, #4927), Caspase-11 (Cell Signaling, #14340), Total
Caspase-3 (Cell Signaling, #9662), Cleaved Caspase-3 (Cell Signaling, #9664),
GSDMD (Cell Signaling, #93709), GSDMD-NT (Cell Signaling, #50928), GSDME
(Santa Cruz, sc393162), IBA-1 (Abcam, ab5076), GFAP (Cell Signaling, #12389) and
β-actin (Sigma-Aldrich, A5441) overnight at 4 °C with agitation. Following primary
antibody incubation, membranes were washed three times with 1 x TBST before
incubating with horseradish peroxidase (HRP)-conjugated secondary antibodies
(Goat Anti-Rabbit – Cell Signaling Technology, Danvers, MA, USA; Goat Anti-Mouse
– Sigma-Aldrich, St. Louis, MO, USA; Goat Anti-Rat – GE Healthcare Life Sciences,
Little Chalfont, UK) for 1 h at 24 °C with agitation. Following secondary antibody
incubation, membranes were washed three times with 1 x TBST, each time for 10
min. The substrate for HRP, enhanced chemiluminescence (ECL) (Bio-Rad
Laboratories, Inc., Hercules, CA, USA) was applied before the membranes were
imaged using ChemiDocXRS+imaging system (Bio-Rad Laboratories, Inc., Hercules,

CA, USA). Quantification of proteins was conducted using Image J software (Version
1.46; National Institute of Health, Bethesda, MD, USA), where protein densitometry
was expressed relative to the densitometry of the corresponding β-actin.
211

Luxol Fast Blue and Cresyl Violet Staining
Mouse brain tissues were fixed in 10% neutral buffered formalin and then
processed into paraffin wax blocks. Coronal sections (4 μm thick) were obtained via
microtome sectioning. Axonal fibre density of Luxol-fast-blue (LFB) staining was
performed to detect the severity of WM lesions. Briefly, de-waxed rehydrated
sections were immersed in the LFB solution (Abcam, UK) at 37°C overnight. Excess
staining was removed by 95% ethanol treatment followed by washing with deionized
water. Grey and white matter differentiation was initiated with the treatment of 0.05%
aqueous lithium carbonate (Abcam, UK) for 20 seconds, followed by 70% ethanol
until the nuclei are decolorized. Sections were immersed in Cresyl Violet solution
(Abcam, UK) for 5 min and washed in deionized water. They were dehydrated in a
ethanol gradient (70 – 100%), and finally cleared in xylene and mounted with a
mounting agent. The bright field images were taken under 4X and 60X magnification
using an Olympus upright Fluorescence Microscope BX53. The WM lesions were
evaluated in 5 brain regions: the optic tract, internal capsule, caudoputamen, corpus
callosum (Medial) and corpus callosum (Paramedian). The severity of the WM
lesions was graded as normal (grade 0), disarrangement of the nerve fibres (grade 1),
the formation of marked vacuoles (grade 2), and the disappearance of myelinated
fibres (grade 3). The severity of neuropathology was scored by three blinded
examiners and the number of neurons in CA1, CA2 and CA3 were counted as
previously described[24]. The general morphology and the neurodegeneration in the
hippocampus of BCAS animals were assessed by Cresyl violet staining.
234

Immunofluorescence Analysis
Paraffinized brain sections were cut from sham and BCAS animals and
immunostained with primary antibodies against Cleaved Caspase-1 (Affinity
Biosciences, AF4022), Cleaved Caspase-3 (Cell Signaling, #9661), GSDMD (Santa
Cruz, sc393581), MBP (Cell Signaling, #78896), MAP2 (Milipore, MAB3418), GFAP
(Cell Signaling, #3670), IBA-1 (Wako, 016-26721), OLIG2 (R&D System, AF2418)
and PECAM-1 (BD Pharmingen, 553370). Images were captured with an Olympus

FluoView FV1000 (Olympus, Japan) laser scanning confocal microscope using
20x/0.70 air objective and 100x/1.45 oil objective, with 488nm Argon ion and 543nm
HeNe laser as the excitation source. Single confocal images were converted to 512 x
512 pixel 12-bit TIFF images.
246

Behavioral Paradigm and Training of BCAS Animals

Visual Acuity Test: To assess visual detection, mice were placed on a platform
within a rotating drum covered with black and white vertical stripes at consistent
spatial frequency. The automated drum was then rotated in a clockwise direction for
30 sec, and subsequently an anti-clockwise direction for 30 sec, after a 15 sec inter-
trial interval. Both larger stripes (2°) and smaller stripes (1°) were tested for all
animals before conducting other behavioral tests.

Open Field Test: Locomotor activity was measured using an open field test. Each
subject was placed in the center of the open field apparatus (25×25×46 cm; Clever
Sys Inc.; VA, USA). Total distance traveled (cm) and average speed (mm/s) were
recorded and tracked via TopScan (Clever Sys Inc.; VA, USA). Data was collected
for 30 minutes.

Morris Water Maze: Morris water maze was performed 3 weeks after surgery (sham
or BCAS), as described previously[25]. Mice were trained 4 times a day at 10-min
intervals for 5 consecutive days in this fixed platform training. In each trial, mice were
given 60 secs to find the platform. Activity of the animal in the water maze was video-
tracked by EthoVision software (Noldus Information Technology, Leesburg, VA), and
the escape latency was recorded and analyzed. Long term spatial memory was
tested on the 6th day, whereby the platform was removed, and each animal was
given 60 secs in the water maze after being released from four different directions.
Time spent around the platform, number of visits to the targeted quadrant and time
spent in the targeted quadrant on the probe day were tracked and analyzed. The
probe day was designed to fall between 32 to 34 days of BCAS surgery.
270

Double Stranded DNA Measurement

Double stranded DNA from the plasma of BCAS animals were measured as
previously described using Quant-iT™ PicoGreen ® dsDNA Reagent and Kits. Briefly,
20μL of plasma sample collected was diluted with 1X TE solution. Subsequently,
equal volume of working solution of Quant-iT™ PicoGreen reagent was added. It was

then protected from light and incubated for 5 minutes at room temperature. Samples
were excited at 485nm and fluorescence intensity was measured at 520nm using a
Synergy HT multi-detection microplate reader (BioTek, Winnooski, VT).
279

Statistical Analysis
Experimental data were analysed by GraphPad Prism 5.02 software (GraphPad
Software, San Diego, CA, USA). All values are expressed as mean ± standard error
of the mean (S.E.M). One-way Analysis of Variance (ANOVA) was used, followed by
a Bonferroni post-hoc test to determine differences between groups. A P-value <0.05
was deemed to be statistically significant.
286
287

Results
Chronic cerebral hypoperfusion increases inflammasome signalling
We evaluated the protein expression of inflammasome receptors over 60 days of
BCAS in the cerebral cortex, hippocampus and striatum (Supplementary Figs. 3, 4 &
5). NLRP3, AIM2 and NLRC4 were all increased in the cerebral cortex, while only
AIM2 receptor expression was increased in the hippocampus compared to sham
controls (Supplementary Fig. 3a-d). NLRP1 expression was increased in the striatum
while other inflammasome receptors remained unchanged (Supplementary Figs. 4a
& b).
297

In the canonical inflammasome pathway, activation and homo-oligomerization of
inflammasome receptors initiates formation of the inflammasome complex,
recruitment and activation of adaptor (ASC) and effector (total caspase-1 and -8)
inflammasome components into biologically active cleaved caspases-1 and -8[6,26].
These effector proteins then catalyze the conversion of mature IL-1β and IL-18 from
their precursors. In the noncanonical inflammasome pathway, caspase-11 is
activated via auto-proteolysis upon dimerization to produce cleaved caspase-11.
Here, expression of total caspase-8 and precursor IL-1β were increased in the
cerebral cortex after BCAS, whereas a decrease in precursor IL-18 was evident, and
ASC and total caspases-1 and -11 were unchanged (Supplementary Fig. 3e-f). In the
hippocampus, expression of total caspases-1, -8 -11 and precursor IL-18 increased,

whereas ASC was decreased after BCAS (Supplementary Fig. 3g-h). ASC, total
caspase-8 and precursors for IL-1β and IL-18 were also increased in the striatum
following BCAS (Supplementary Fig. 5c & d).

To assess canonical inflammasome activation, expression of cleaved caspase-1
subunits (p33, transient active unit; p20, final by-product) and -8 were examined.
After BCAS, elevated expression of cleaved caspase-1 (p20) and -8 was detected in
the cerebral cortex (Fig. 1a & b) and also hippocampus (Fig. 1c & d). Mature IL-1β
and IL-18 – direct downstream markers of inflammasome activation – were also
higher in the cerebral cortex and hippocampus after BCAS versus sham controls
(Fig. 1a-d). Expression of non-canonical cleaved caspase-11 was also elevated in
the cerebral cortex and hippocampus after BCAS (Fig. 1a-d), whereas only cleaved
caspase-8, IL-1β and IL-18 were increased in the striatum (Supplementary Fig. 6).
Overall, the data reveal that inflammasome activation occurs in the brain in a spatial-
temporal manner as a result of chronic cerebral hypoperfusion.

Chronic cerebral hypoperfusion promotes apoptosis and pyroptosis
We next examined cell death mechanisms associated with inflammasome
activation. Cleaved caspases-1 and -8 can induce apoptosis and pyroptosis by
catalyzing the cleavage of total caspase-3 and full-length Gasdermin D (GSDMD-FL).
GSDMD-NT can oligomerize with other N-terminal units and cause pore formation in
the nuclear and plasma membranes, leading to leakage of cellular content and
proinflammatory cytokines[27-28]. GSDMD activation can also be facilitated by cleaved
caspase-11[29]. Thus, we assessed the expression of cleaved caspase-3
and GSDMD-NT, along with their precursor proteins after BCAS (Fig. 1e-
h; Supplementary Fig. 7). Our data show that in the cerebral cortex (Fig. 1e & f) and
hippocampus (Fig. 1g & h) the expression of cleaved caspase-3 and GSDMD-NT
were increased following BCAS compared to sham controls. Pyroptotic cell death
(i.e. GSDMD-NT expression) was more prominent in the cerebral cortex following
BCAS (Fig. 1e & f), whereas only apoptosis (i.e. increased cleaved caspase-3
expression) was detected in the striatum following BCAS (Supplementary Fig. 8).

We also assessed inflammasome-associated secondary necrosis, which can be
induced by cleaved caspase-3, as indicated by expression of N-Terminal Gasdermin
E (GSDME-NT). Similar to GSDMD-NT, inflammasome-generated GSDME-NT can
permeabilize nuclear and plasma membranes and mitochondria, linking
inflammasome activation to cell death[28]. We found GSDME-NT to be increased in the
cerebral cortex, but not in the hippocampus or striatum following BCAS
(Supplementary Fig. 7 & 8). Expression patterns of inflammasome-related proteins
and pathways following BCAS-induced chronic cerebral hypoperfusion are
summarized in Supplementary Fig. 9. Overall, the data indicate that BCAS induces
differential priming of inflammasome components, inflammasome activation, and
induction of apoptosis and pyroptosis pathways in the brain in a spatial-temporal
manner.
353
Chronic cerebral hypoperfusion induces glial cell activation, white matter
lesions and hippocampal neuronal death
We next analyzed for associations between inflammasome activity and hallmark
VaD pathologies involving glial cell activation, white matter integrity and neuronal
loss. Expression of ionized calcium binding adaptor molecule 1 (Iba1) and glial
fibrillary acidic protein (GFAP) indicated activation of microglia and astrocytes,
respectively. Myelin basic protein (MBP) and Luxol fast blue staining revealed myelin
integrity. Cresyl violet staining and microtubule-associated protein 2 (MAP2) enabled
assessment of neuronal loss in the hippocampus. Following BCAS, we found
increased Iba-1 in the cerebral cortex (Fig. 2a & b) and increased GFAP in the
hippocampus (Fig. 2c & d), and both markers were increased in the striatum
(Supplementary Fig. 10a & b). Staining for MBP and MAP2 was reduced in the
cerebral cortex (Fig. 2e & g), hippocampus (Fig. 2f & h) and striatum (Supplementary
Fig. 10c & d) following 30 days of BCAS.
368
White matter integrity of five brain regions was assessed after 7-60 days of
hypoperfusion, and found to be disrupted in the corpus callosum (Paramedian and
Medial), caudoputamen, internal capsule and optic tract (Fig. 5i & j). All five areas
exhibited time-dependent rarefaction of white matter (Fig. 5j) from 7 days, although
the white matter lesion (WML) index was not increased in the optic tract until 60 days
of BCAS (Fig. 5i & j). BCAS-induced neuronal loss was evident in hippocampal CA1,

CA2 and CA3 areas (Fig. 5k-n). Whereas hippocampal sections from sham controls

showed normal neuronal cell bodies with distinct nuclei, nucleoli, and densely packed

neurons in all three hippocampal areas, there were signs of widespread neuronal

loss evident after BCAS (Fig. 5k-n). Degeneration was more pronounced in the

hippocampal CA2 and CA3 areas with severe atrophy from 7 days (Fig. 5k-n).

AIM2 activation mediates apoptosis and pyroptosis during chronic cerebral

hypoperfusion

As AIM2 was upregulated in the cerebral cortex and hippocampus, we postulated

its involvement in BCAS-induced brain injury. Analysis of serum revealed an increase

in cell-free double stranded DNA (dsDNA) levels following BCAS, especially after 15

days (Fig. 3a). As dsDNA is the only known ligand that activates the AIM2 receptor[30-

31], the finding supports a potential involvement of the AIM2 inflammasome in BCAS-

induced brain injury.

To further examine the role of AIM2 inflammasome activation on injury following

chronic cerebral hypoperfusion, mice with AIM2 deficiency (AIM2 KO) were studied

(Supplementary Fig. 1). We first confirmed via laser speckle contrast imaging that

cerebral blood flow was equivalent under basal conditions and similarly reduced

during the surgery to perform BCAS in wild-type (WT) and AIM2 KO mice

(Supplementary Fig. 11). This was similar to previously published data from BCAS in

mice[32]. This BCAS-induced reduction in flow was sustained to a similar degree for at

least 15 days in both genotypes but was slightly higher in AIM2 KO than WT mice at

30 days (Supplementary Fig. 11a-c).

We found no difference in the basal expression levels of cleaved caspase-1, -8,

and -11, and both mature IL-1β and IL-18 (Fig. 3b-e) or in the levels of total caspases

-1, -8 and -11, or IL-1β and IL-18 precursors in the cerebral cortex and hippocampus

of WT and AIM2 KO mice following BCAS (Supplementary Fig. 12). However, there

was evidence of reduced inflammasome activation following BCAS in AIM2 KO mice

(Fig. 3b-e). Of the three active effector proteins, only canonical cleaved caspase-1

expression was reduced in AIM2 KO mice compared to WT control following BCAS

(Fig. 3b-e; Supplementary Fig. 13a & b). There was also reduced levels of mature IL-

1β and IL-18 in the cerebral cortex and hippocampus of AIM2 KO mice at 15 and 30 days following BCAS (Fig. 3b-e; Supplementary Fig. 13a & b). These data thus indicated less inflammasome activity occurs after BCAS in the absence of the AIM2 receptor, especially where mediated by canonical cleaved caspase-1. Furthermore, protein expression of both cleaved caspase-3 and GSDMD-NT was lower at 30 days after BCAS in the cerebral cortex (Fig. 3f & g) and hippocampus (Fig. 3h & i) of AIM2 KO mice.

Immunofluorescence studies indicated that cleaved caspase-1 (CC1) activity was lower in AIM2 KO mice in cortical (Fig. 3j; Supplementary Fig. 14a-e & 15a-e) and hippocampal (Fig. 3k; Supplementary Fig. 14f-j & 15f-j) neurons (MAP2 positive) and microglia (Iba-1 positive) compared to WT controls at 30 days following BCAS. Immunoreactivity against cleaved caspase-1 was similar in cortical and hippocampal oligodendrocytes (OLIG2 positive), astrocytes (GFAP positive) and endothelial cells (PECAM-1 positive) of WT and AIM2 KO mice. Cellular specificity of AIM2 inflammasome-mediated cell death following BCAS was assessed by immunoreactivity against cleaved caspase-3 (CC3) and GSDMD in the cerebral cortex (Fig. 3j; Supplementary Fig. 14a-e; Supplementary Fig. 16a-e; Supplementary Fig.17a-e) and hippocampus (Fig. 3k; Supplementary Fig. 14f-j; Supplementary Fig. 16f-j; Supplementary Fig.17f-j). The data indicate that pro-apoptotic cleaved caspase-3 immunoreactivity was lower in cortical (Fig. j; Supplementary Fig. 14a-e & 16a-e) and hippocampal (Fig. 3k; Supplementary Fig. 14f-j & 16f-j) neurons (MAP2 positive) and microglia (Iba-1 positive) of AIM2 KO versus WT mice. Similarly, immunoreactivity of pro-pyroptotic GSDMD was reduced in neurons and microglia in the cortex (Fig. 3j; Supplementary Fig. 14a-e & 17a-e) and hippocampus (Fig. 3k; Supplementary Fig. 14f-j & 17f-j) of AIM2 KO mice. These findings were consistent with more neurons and fewer microglial activation in AIM2 KO than WT mice following chronic cerebral hypoperfusion.

AIM2 KO mice are resistant to microglial activation, myelin breakdown, hippocampal neuronal loss and cognitive deficits following chronic cerebral hypoperfusion

There was a lower expression of Iba-1 in the cerebral cortex and hippocampus of AIM2 KO than WT mice after BCAS (Fig. 4a-d). These results suggest a role for

AIM2 in microglial activation during chronic cerebral hypoperfusion. We next
investigated if AIM2 KO mice are protected against BCAS-induced myelin injury.
Although sham-operated WT and AIM2 KO mice displayed no difference in
immunoreactivity for MBP, at 30 days after BCAS, there was weaker
immunoreactivity for MBP in the cerebral cortex of WT mice exhibited than AIM2 KO
mice (Fig. 4e). A similar difference in MBP immunoreactivity between WT and AIM2
KO was seen in CA2 and CA3 hippocampal regions (Fig. 4e). As myelin integrity was
superior in AIM2 KO mice following BCAS, we assessed whether these mice might
be resistant to WML. After BCAS, Luxol fast blue staining in WT mice indicated
myelin rarefaction and vacuole formation in several WM regions, but a smaller
changes were present in the corpus callosum (paramedian or medial),
caudoputamen or internal capsule regions of AIM2 KO mice (Fig. 5a-e). Neuronal
loss in hippocampal CA1, CA2 and CA3 regions was also less in AIM2 KO compared
to WT mice at 15 and 30 days of cerebral hypoperfusion (Fig. 5f-h). These data
confirm a deleterious role of the AIM2 inflammasome in mediating VaD pathologies
associated with chronic cerebral hypoperfusion.

Finally, we tested for a role of the AIM2 inflammasome in BCAS-induced cognitive
decline. It was previously shown that BCAS-induced chronic cerebral hypoperfusion
in mice resulted in cognitive impairment such as working memory dysfunction and
memory loss[32-33]. To examine the effects of AIM2 KO against BCAS-induced
cognitive deficit, we conducted the Morris Water Maze and two other control tests. To
ensure that the animals were able to detect the visual cues during the Morris Water
Maze, a visual acuity test was conducted. All animals showed a 100% passing rate,
and so proceeded to the subsequent tests. In the open field test conducted to
investigate explorative and locomotor ability, the average speed of AIM2 KO mice
was significantly higher than the WT BCAS controls (Fig. 5i). Several parameters
were assessed in the Morris Water Maze performance, including escape latency
during five training days. Generally, we observed a shorter time was taken to reach
the platform each day across all groups (Fig. 5j). Similar daily improvement was
exhibited in WT or AIM2 KO mice sham controls and AIM2 KO mice following BCAS,
whereas WT mice subjected to BCAS took approximately twice as long (Fig. 5j).
Following 5 days of training, spatial memory was assessed whereby WT mice
subjected to BCAS spent less time around the platform and fewer visits to the target

quadrant than WT controls (Fig. 5k). By contrast, AIM2 KO mice subjected to BCAS

spent a longer time around the platform and visited the target quadrant more often

than WT BCAS mice (Fig. 5k & l). Thus, the behavioral data indicates that BCAS-

induced cognitive impairments were reduced in mice deficient in AIM2, suggesting a

pivotal role for the AIM2 inflammasome in mediating VaD pathology associated with

chronic cerebral hypoperfusion.

Discussion

Our findings indicate that inflammasome receptors are expressed in the cortex and

hippocampus where the AIM2 inflammasome is activated in response to chronic

cerebral hypoperfusion, leading to cellular pathology and cognitive impairment in a

mouse model of vascular dementia. Genetic deficiency of AIM2 expression resulted

in less brain inflammation, white matter injury, neuronal loss and cognitive decline

during cerebral hypoperfusion. These data are the first to indicate that the AIM2

inflammasome promotes neuronal and white matter injury following cerebral

hypoperfusion, identifying AIM2 as a key contributor to sterile inflammatory

responses in vascular dementia.

Inflammasomes are intracellular multiprotein complexes composed of sensors for

various microbial components, viral RNA and damage- or danger-associated

molecular patterns (DAMPs) produced during cell injury[34-35]. These innate immune

complexes are categorized according to their structural characteristics as either

nucleotide-binding domain–like receptors (NLRs) or absent in melanoma 2 (AIM2)-

like receptors[35-36]. These receptors oligomerize in response to activation by external

stimuli and then recruit the adaptor protein, apoptosis-associated speck-like protein

containing a caspase recruitment domain (ASC)[37]. Ultimately, binding of caspase-1

results in its cleavage and activation, and the generation of mature pro-inflammatory

cytokines, IL-1β and IL-18, as well as pro-apoptotic cleaved caspase-3, and pro-

pyroptotic N-terminal GSDMD and GSDME[38].

We previously demonstrated that inflammasomes contribute to neuronal death

following ischemic stroke[7], in part via NF-κB and MAPK signaling[23]. Our present data

indicate that chronic cerebral hypoperfusion may also lead to expression and

activation of NLRP and AIM2 inflammasome receptors in multiple brain areas,

especially in the hippocampal region which is known to be particularly vulnerable to

injury after even mild ischemia[39]. Cerebral hypoperfusion in the BCAS model used

here is known to impact the hippocampus by mechanisms involving ionic imbalance,

excitotoxicity, mitochondrial dysfunction and oxidative stress[40-42].

It is important to elucidate the mechanism(s) by which the AIM2 inflammasome

mediates chronic hypoperfusion-induced brain injury. Double-stranded DNA (dsDNA)

is one candidate as a ligand to activate AIM2 in neurons and glia following chronic

hypoperfusion[30-31]. AIM2 is known to play a role in the host response to microbial

DNA[43], but several reports suggest that endogenous self dsDNA can also activate

the AIM2 inflammasome[30,44]. Here we found that the plasma concentration of dsDNA

progressively increased following BCAS-induced cerebral hypoperfusion, consistent

with the possibility that this ligand promoted AIM2-mediated amplification of brain

inflammation and injury. Thus it is plausible that AIM2 functions as a DAMP sensor to

promote sterile inflammation in vascular dementia.

We found that various white matter regions exhibited injury within 7 days of BCAS,

consistent with the known vulnerability of white matter to hypoperfusion. Reduced

blood flow to white matter can predict the clinical development of WML within 18

months[45]. We observed that AIM2 deficient animals subjected to BCAS exhibited

greater white matter integrity and lower WML severity scores than WT mice after

BCAS. Using magnetic resonance imaging, WML appeared as white matter

hyperintensities (WMH) in VaD patients arises due to demyelination and axonal loss

as well as microglial and endothelial activation[46-48]. Consistent with this, we found

that cell death, loss of myelin basic protein and microglia activation were present in

our BCAS mouse model and were reduced in mice lacking AIM2 inflammasomes.

CA1, CA2 and CA3 are the sub-regions of the dorsal hippocampus responsible for

memory encoding and retrieval[49-50]. We found that CA2 and CA3 neurons

experienced death during chronic hypoperfusion sooner than CA1 neurons. Reduced

levels of BCAS-induced neuronal loss occurred in hippocampal CA2 and CA3 areas

of AIM2-deficient animals compared to WT controls, suggesting that AIM2

inflammasome contributed to neuronal loss in the hippocampus during hypoperfusion.

The AIM2 receptor is widely expressed, with its highest levels occurring in

microglia under physiological conditions[51]. Furthermore, AIM2 is the most abundant

inflammasome receptor expressed in cortical and hippocampal neurons[52]. Our data

are consistent with a role of AIM2 in these brain resident cells and regions, but we

cannot exclude circulating leukocytes as another possible cell mediating the pro-

inflammatory actions of AIM2 in the brain in chronic hypoperfusion. Leukocytes

contribute to other forms of brain injury and neurodegeneration[53-55] their infiltration

from the circulation would be expected in association with reduced blood-brain

barrier integrity and blood vessel density known to occur in hypoperfusion-induced

VaD[56].

Previous studies have recently reported evidence that the AIM2 inflammasome

contributes to brain injury and chronic post-stroke cognitive impairment in mice[57], and

to reduced neuroplasticity and spatial memory in a mouse model of AD[16]. Similarly,

there is evidence supporting a role for the NLRP3 inflammasome in cognitive

impairment in models of dementia and diabetes[9,58]. Inflammation is known to

adversely impact cognitive function, and our data are consistent with inflammasomes

promoting an innate immune response to mediate such an effect during cerebral

hypoperfusion.

In summary, we have provided substantial evidence that activation of the AIM2

inflammasome plays a key role in promoting brain inflammation, white matter lesions,

neuronal cell death and cognitive impairment induced by chronic cerebral

hypoperfusion. These effects of AIM2 involve promotion of apoptosis and pyroptosis

of cortical and hippocampal neurons. The AIM2 inflammasome may therefore

represent a promising therapeutic target for attenuating cognitive impairment in VaD.

Acknowledgements

We thank Professor Jenny P. Ting (University of North Carolina, Chapel Hill, Chapel Hill, NC, USA) for providing the AIM2 deficient mice. We thank Professor David Hess (Augusta University) for allowing T.V.A to learn the BCAS model of VaD in his laboratory. Supplementary Figure 1 and Supplementary Figure 9a in this article was created using BioRender.

Funding

This work was supported by the National Medical Research Council Research Grants (NMRC-CBRG-0102/2016 and NMRC/OFIRG/0036/2017), Singapore.

Author Contributions

Study Conception & Design: T.V.A., L.P., and D.Y.F; Investigation-Performed Experiment or Data Collection: T.V.A., L.P., P.Y., H.M.L., S.L.F., S.W.K., V.R., S.S., D.Y.F., R.I.V., and N.P.; Data Analysis: L.P., P.Y., V.R., S.S., D.Y.F., and T.V.A.; Data Interpretation: L.P., T.V.A., and D.Y.F; Writing-Manuscript Preparation and intellectual input: T.V.A., L.P., D.Y.F., M.B.K., G.R.D., C.G.S., D.G.J., M.K.P.L., C.L.H.C., and L.H.K.L.; Supervision & Administration: T.V.A., D.Y.F., M.K.P.L., and C.L.H.C.

Conflicts Of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

FIGURE LEGENDS

Figure 1: Effect of chronic cerebral hypoperfusion on inflammasome activation and cell death in the cerebral cortex and hippocampus over time following BCAS. (a & b), representative immunoblots and quantification illustrating increased levels of activated inflammasome effector proteins such as cleaved caspase-1 (p20), -8, and -11, and maturation of downstream effector targets, IL-1β and IL-18, in the cerebral cortex. (c & d), representative immunoblots and quantification illustrating increased levels of activated inflammasome effector proteins such as cleaved caspase-1 (p20, p33), -8, and -11, and maturation of downstream effector targets, IL-1β and IL-18, in the hippocampus. (e & f), representative immunoblots and quantification illustrating increases in the expression of apoptotic marker cleaved caspase-3 and pyroptotic marker N-terminal GSDMD in the cerebral cortex. (g & h), representative immunoblots and quantification illustrating increases in the expression of apoptotic marker cleaved caspase-3 and pyroptotic marker N-terminal GSDMD in the hippocampus. β-actin was used as a loading control. Data are represented as mean ± S.E.M. n=6-7 mice in each experimental group. *P<0.05 compared with Sham; **P<0.01 compared with Sham; ***P<0.001 compared with Sham. Abbreviations: BCAS, bilateral common carotid artery stenosis; Cl, cleaved; FL, full length; NT, N-terminal; GSDMD, gasdermin D; GSDME, gasdermin E.

Figure 2: Effect of chronic cerebral hypoperfusion on the levels of glial activation, white matter integrity and hippocampal neuronal density in the cerebral cortex and hippocampus following BCAS. (a), representative immunoblots and quantification illustrating increased microglial activation due to increased levels of Iba-1 in the cerebral cortex over time following BCAS. β-actin was used as a loading control. (b), representative immunofluorescence analysis of Iba-1 and GFAP in the cerebral cortex provide supporting evidence of microglial activation following BCAS. (c), representative immunoblots and quantification illustrating increased astroglial activation due to increased levels of GFAP in the hippocampus over time following BCAS. (d), representative immunofluorescence analysis of Iba-1 and GFAP in the hippocampus provide supporting evidence of astroglial activation following BCAS. Magnification x 100. Scale bar, 20 μm. (e & f), representative immunofluorescence images illustrating a loss of myelin due to decreased levels of MBP immunoreactivity in the cerebral cortex and hippocampus following BCAS. (g & h), representative immunofluorescence images illustrating a loss of neurons due to decreased levels of MAP2 immunoreactivity in the cerebral cortex and hippocampus following BCAS. Magnification x 20. Scale bar, 120 μm. Data are represented as mean ± S.E.M. n=6-7 mice in each experimental group. β-actin was used as a loading control. *P<0.05 compared with Sham; **P<0.01 compared with Sham; ***P<0.001 compared with Sham. (i & j), representative Luxol fast blue stained images and quantification illustrating disruption of white matter integrity due to increased myelin rarefaction and white matter lesion formation in the corpus callosum (paramedian), corpus callosum (medial), caudoputamen, internal capsule and optic tract over time following BCAS. The severity of white matter disruption was graded accordingly: Grade 0 = no disruption; Grade 1 = disarrangement of nerve fibers; Grade 2 = formation of marked vacuoles; Grade 3 = disappearance of myelinated fibers. (k-n), representative crystal violet images and quantification illustrating loss of Nissl positively stained neurons in hippocampal CA1, CA2 and CA3 regions over time following BCAS. Data are represented as mean ± S.E.M. n=5-7 mice in each

experimental group. *P<0.05 compared with Sham; **P<0.01 compared with Sham;
***P<0.001 compared with Sham. Magnification x 60. Scale bar, 20 µm. Images
were taken under identical exposures and conditions. Abbreviations: BCAS, bilateral
common carotid artery stenosis; Iba-1, ionized calcium binding adaptor molecule-1;
GFAP, glial fibrillary acidic protein; MBP, myelin basic protein; MAP2, microtubule-
associated protein 2.

Figure 3: Effect of chronic cerebral hypoperfusion on inflammasome-mediated
programmed cell death in AIM2 KO mice following BCAS. (a), quantification
illustrating increased serum concentrations of the AIM2 receptor ligand, cell free
double stranded DNA (dsDNA), over time in wild-type mice following BCAS. Data are
represented as mean ± S.E.M. n=5-7 mice in each experimental group. *P<0.05
compared with Sham. (b-e), representative immunoblots and quantification
illustrating suppression of inflammasome activation due to decreased expression of
cleaved caspase-1 (p20) at Day 15 and 30, and maturation of IL-1β cytokine
production at Day 30 in the cerebral cortex (b & c) and hippocampus (d & e) of AIM2
KO mice compared to WT controls following BCAS. (f-i), representative immunoblots
and quantification illustrating attenuation of apoptotic and pyroptotic cell death due to
decreased expression of cleaved caspase-3 and GSDMD-NT, respectively, at Day
30 in the cerebral cortex and hippocampus of AIM2 KO mice compared to WT
controls following BCAS. β-actin was used as a loading control. Data are represented
as mean ± S.E.M. n=5-8 mice in each experimental group. *P<0.05 compared with
WT Sham; **P<0.01 compared with WT Sham; ***P<0.001 compared with WT Sham.
+P<0.05 compared with WT BCAS; ++P<0.01 compared with WT BCAS; +++P<0.001
compared with WT BCAS. (j & k), representative merged immunofluorescence
images of DAPI, cleaved caspase-1 p10 (CC1), cleaved caspase-3 (CC3) and
GSDMD co-localized within neurons (MAP2 positive), microglia (Iba-1 positive),
oligodendrocytes (OLIG2 positive) and endothelial cells (PECAM-1 positive) in the
cerebral cortex and hippocampus of WT controls following BCAS. No substantial co-
localization of cleaved caspase-1 p10, cleaved caspase-3 and GSDMD was
observed in astrocytes (GFAP positive) in the cerebral cortex and hippocampus of
WT controls following BCAS. (j 7 k), representative merged immunofluorescence
images illustrate a reduction in inflammasome activation, and apoptotic and
pyroptotic cell death due to decreased expression levels of cleaved caspase-1,
cleaved caspase-3 and GSDMD, respectively, in both neurons and microglia in the
cerebral cortex (j) and hippocampus (k) in AIM2 KO mice compared to WT controls
following BCAS. Zoom Magnification x 100. Scale bar, 20 µm. Abbreviations: BCAS,
bilateral common carotid artery stenosis; Cl, cleaved; FL, full length; NT, N-terminal;
WT, wild-type; KO, knock out; CC1, cleaved caspase-1; CC3, cleaved caspase-3;
GSDMD, gasdermin D; MAP2, microtubule-associated protein 2; Iba-1, ionized
calcium binding adaptor molecule-1; OLIG2, oligodendrocyte transcription factor 2;
GFAP, glial fibrillary acidic protein; PECAM-1, platelet endothelial cell adhesion
molecule-1.

Figure 4: Effect of chronic cerebral hypoperfusion on glial activation and
myelin expression in the cerebral cortex and hippocampus in AIM2 KO mice
following BCAS. (a-d), representative immunoblots and quantification illustrating
resistance to microglial activation due to decreased expression of Iba-1 in the
cerebral cortex and hippocampus in AIM2 KO mice compared to WT controls
following BCAS. No effect on astroglial activation was observed as the expression of

GFAP in the cerebral cortex and hippocampus remained unchanged in AIM2 KO mice compared to WT controls following BCAS. β-actin was used as a loading control. Data are represented as mean ± S.E.M. n=6-8 mice in each experimental group. *$P<0.05$ compared with WT Sham; **$P<0.01$ compared with WT Sham. +$P<0.05$ compared with WT BCAS; ++$P<0.01$ compared with WT BCAS; +++$P<0.001$ compared with WT BCAS. (e), representative immunofluorescence images illustrating resistance to myelin loss due to increased MBP immunoreactivity in the cerebral cortex, and hippocampal CA1, CA2 and CA3 regions in AIM2 KO mice compared to WT controls following BCAS. Magnification x 100. Scale bar, 20 µm. Abbreviations: BCAS, bilateral common carotid artery stenosis; KO, knock out; Iba-1, ionized calcium binding adaptor molecule-1; GFAP, glial fibrillary acidic protein.

Figure 5: Effect of chronic cerebral hypoperfusion on cortical white matter integrity, hippocampal neuronal density and explorative locomotive behavior and spatial memory in AIM2 KO mice following BCAS. (a-e), representative Luxol fast blue stained images and quantification illustrating preserved white matter integrity due to decreased myelin rarefaction and white matter lesion formation in the corpus callosum (paramedian), corpus callosum (medial), caudoputamen and internal capsule in AIM2 KO mice compared with WT controls following BCAS. No significant difference in white matter integrity was observed in the optic tract between AIM2 KO mice compared with WT controls following BCAS. The severity of white matter disruption was graded accordingly: Grade 0 = no disruption; Grade 1 = disarrangement of nerve fibres; Grade 2 = formation of marked vacuoles; Grade 3 = disappearance of myelinated fibres. (f-h), representative crystal violet images and quantification illustrating increased Nissl positively stained neurons in hippocampal CA1, CA2 and CA3 regions in AIM2 KO mice compared to WT controls following BCAS. Data are represented as mean ± S.E.M. n=5-7 mice in each experimental group. *$P<0.05$ compared with WT Sham; **$P<0.01$ compared with WT Sham; ***$P<0.001$ compared with WT Sham. +$P<0.05$ compared with WT BCAS; ++$P<0.01$ compared with WT BCAS; +++$P<0.001$ compared with WT BCAS. Magnification x 60. Scale bar, 20 µm. (i), quantification illustrating an open field test conducted on all mice to examine locomotor ability by measuring the average speed of movement (mm/s) of the animal in the open field. It was shown that WT BCAS mice had a higher average speed (mm/s) of movement when compared to WT Sham controls indicating no impairment of locomotor ability. (j-l), quantifications illustrating a Morris water maze test conducted on all mice to examine spatial learning and memory by measuring the escape latency (sec), time spent around the platform on probe day (sec) and number of visits in the target quadrant. (j), in general, a reduction in escape latency (sec) was observed in all mice across the training period. Specifically, WT Sham, AIM2 KO Sham and AIM2 KO BCAS groups displayed a trend with a shorter escape latency and steeper declining gradient, which reflected an improved learning ability when compared to the WT BCAS group. (k & l), in general, the AIM2 KO BCAS group displayed a trend with a longer time spent around the platform on probe day and a significantly higher number of visits in the target quadrant, which reflected retention of spatial memory when compared to the WT BCAS group. Data are represented as mean ± S.E.M. n=9-11 mice in each experimental group. *$P<0.05$ compared with WT Sham. #$P<0.05$ compared with WT BCAS. Abbreviations: BCAS, bilateral common carotid artery stenosis; WT, wild type; KO, knock out; WML, white matter lesion.

REFERENCES

1) Wolters FJ, Ikram MA. Epidemiology of Vascular Dementia. Arterioscler Thromb Vasc Biol 2019; **39:**1542-1549.
2) Van der Flier WM, Skoog I, Schneider JA, Pantoni L, Mok V, Chen CLH, Scheltens P. Vascular cognitive impairment. Nat Rev Dis Primers 2018; **4:** 18003.
3) Alber J, Alladi S, Bae HJ, Barton DA, Beckett LA, Bell JM, Berman SE, Biessels GJ, Black SE, Bos I, et al. White matter hyperintensities in vascular contributions to cognitive impairment and dementia (VCID): Knowledge gaps and opportunities. Alzheimers Dement (N Y) 2019; **5:** 107-117.
4) Schmitz M, Hermann P, Oikonomou P, Stoeck K, Ebert E, Poliakova T, Schmidt C, Llorens F, Zafar S, Zerr I. Cytokine profiles and the role of cellular prion protein in patients with vascular dementia and vascular encephalopathy. Neurobiol Aging 2015; **36:** 2597-2606.
5) Zhou Y, Zhang J, Wang L, Chen Y, Wan Y, He Y, Jiang L, Ma J, Liao R, Zhang X, et al. Interleukin-1β impedes oligodendrocyte progenitor cell recruitment and white matter repair following chronic cerebral hypoperfusion. Brain Behav Immun 2017; **60:** 93-105.
6) Fann DY, Lee SY, Manzanero S, Chunduri P, Sobey CG, Arumugam TV. Pathogenesis of acute stroke and the role of inflammasomes. Ageing Res Rev 2013; **12:** 941-966.
7) Fann DY, Lee SY, Manzanero S, Tang SC, Gelderblom M, Chunduri P, Bernreuther C, Glatzel M, Cheng YL, Thundyil J, et al. Intravenous immunoglobulin suppresses NLRP1 and NLRP3 inflammasome-mediated neuronal death in ischemic stroke. Cell Death Dis 2013; **4:** e790.
8) Gordon R, Albornoz EA, Christie DC, Langley MR, Kumar V, Mantovani S, Robertson AAB, Butler MS, Rowe DB, O'Neill LA, et al. Inflammasome inhibition prevents α-synuclein pathology and dopaminergic neurodegeneration in mice. Sci Transl Med 2018; **10:** 465.
9) Heneka MT, Kummer MP, Stutz A, Delekate A, Schwartz S, Vieira-Saecker A, Griep A, Axt D, Remus A, Tzeng TC, et al. NLRP3 is activated in Alzheimer's disease and contributes to pathology in APP/PS1 mice. Nature 2013; **493:** 674-678.
10) Johann S, Heitzer M, Kanagaratnam M, Goswami A, Rizo T, Weis J, Troost D, Beyer C. NLRP3 inflammasome is expressed by astrocytes in the SOD1 mouse model of ALS and in human sporadic ALS patients. Glia 2015; **63:** 2260-2273.
11) Poh L, Kang SW, Baik SH, Ng GYQ, She DT, Balaganapathy P, Dheen ST, Magnus T, Gelderblom M, Sobey CG et al. Evidence that NLRC4 inflammasome mediates apoptotic and pyroptotic microglial death following ischemic stroke. Brain Behav Immun 2019; **75:** 34-47.
12) Ball DP, Taabazuing CY, Griswold AR, Orth EL, Rao SD, Kotliar IB, Vostal LE, Johnson DC, Bachovchin DA. Caspase-1 interdomain linker cleavage is required for pyroptosis. Life Sci Alliance 2020; **3:** 3.
13) Yuan B, Zhou XM, You ZQ, Xu WD, Fan JM, Chen SJ, Han YL, Wu Q, Zhang X. Inhibition of AIM2 inflammasome activation alleviates GSDMD-induced pyroptosis in early brain injury after subarachnoid haemorrhage. Cell Death Dis 2020; **11:** 76.
14) McKenzie BA, Mamik MK, Saito LB, Boghozian R, Monaco MC, Major EO, Lu JQ, Branton WG, Power C. Caspase-1 inhibition prevents glial inflammasome

activation and pyroptosis in models of multiple sclerosis. Proc Natl Acad Sci USA
2018; **115:** E6065-E6074.

15) Dempsey C, Rubio Araiz A, Bryson KJ, Finucane O, Larkin C, Mills EL,
Robertson AAB, Cooper MA, O'Neill LAJ, Lynch MA. Inhibiting the NLRP3
inflammasome with MCC950 promotes non-phlogistic clearance of amyloid-β and
cognitive function in APP/PS1 mice. Brain Behav Immun 2017; **61:** 306-316.

16) Chen J, Shu S, Chen Y, Liu Z, Yu L, Yang L, Xu Y, Zhang M. AIM2 deletion
promotes neuroplasticity and spatial memory of mice. Brain Res Bull 2019; **152:**
85-94.

17) Paganelli R, Di Iorio A, Patricelli L, Ripani F, Sparvieri E, Faricelli R, Iarlori C,
Porreca E, Di Gioacchino M, Abate G. Proinflammatory cytokines in sera of
elderly patients with dementia: levels in vascular injury are higher than those of
mild-moderate Alzheimer's disease patients. Exp Gerontol 2002; **37:** 257-263.

18) Yasutake C, Kuroda K, Yanagawa T, Okamura T, Yoneda H. Serum BDNF, TNF-
alpha and IL-1beta levels in dementia patients: comparison between Alzheimer's
disease and vascular dementia. Eur Arch Psychiatry Clin Neurosci 2006; **256:**
402-406.

19) Zuliani G, Ranzini M, Guerra G, Rossi L, Munari MR, Zurlo A, Volpato S, Atti AR,
Blè A, Fellin R. Plasma cytokines profile in older subjects with late onset
Alzheimer's disease or vascular dementia. J Psychiatr Res 2007; **41:** 686-693.

20) Mulugeta E, Molina-Holgado F, Elliott MS, Hortobagyi T, Perry R, Kalaria RN,
Ballard CG, Francis PT. Inflammatory mediators in the frontal lobe of patients
with mixed and vascular dementia. Dement Geriatr Cogn Disord 2008; **25:** 278-
286.

21) Wilson JE, Petrucelli AS, Chen L, Koblansky AA, Truax AD, Oyama Y, Rogers
AB, Brickey WJ, Wang Y, Schneider M, et al. Inflammasome-independent role of
AIM2 in suppressing colon tumorigenesis via DNA-PK and Akt. Nat Med 2015;
21: 906-913.

22) Khan MB, Hafez S, Hoda MN, Baban B, Wagner J, Awad ME, Sangabathula H,
Haigh S, Elsalanty M, Waller JL, Hess DC. Chronic Remote Ischemic
Conditioning Is Cerebroprotective and Induces Vascular Remodeling in a VCID
Model. Transl Stroke Res 2018; **9:** 51-63.

23) Fann DY, Lim YA, Cheng YL, Lok KZ, Chunduri P, Baik SH, Drummond GR,
Dheen ST, Sobey CG, Jo DG, et al. Evidence that NF-κB and MAPK Signaling
Promotes NLRP Inflammasome Activation in Neurons Following Ischemic Stroke.
Mol Neurobiol 2018; **55:** 1082-1096.

24) Nishio K, Ihara M, Yamasaki N, Kalaria RN, Maki T, Fujita Y, Ito H, Oishi N,
Fukuyama H, Miyakawa T, Takahashi R, Tomimoto H. A Mouse Model
Characterizing Features of Vascular Dementia With Hippocampal Atrophy.
Stroke 2010; **41:** 1278–1284.

25) Madar R, Rotter A, Waldman Ben-Asher H, Mughal MR, Arumugam TV, Wood
WH 3rd, Becker KG, Mattson MP, Okun E. Postnatal TLR2 activation impairs
learning and memory in adulthood. Brain Behav Immun. 2015; **48:** 301-312.

26) Fritsch M, Günther SD, Schwarzer R, Albert MC, Schorn F, Werthenbach JP,
Schiffmann LM, Stair N, Stocks H, Seeger JM., et al, Caspase-8 is the molecular
switch for apoptosis, necroptosis and pyroptosis. Nature 2019; **575:** 683-687.

27) Liu X, Zhang Z, Ruan J, Pan Y, Magupalli VG, Wu H, Lieberman J.
Inflammasome-activated gasdermin D causes pyroptosis by forming membrane
pores. Nature 2016; **535:** 153-158.

28) Rogers C, Erkes DA, Nardone A, Aplin AE, Fernandes-Alnemri T, Alnemri ES.
Gasdermin pores permeabilize mitochondria to augment caspase-3 activation
during apoptosis and inflammasome activation. Nat Commun 2019; **10:** 1689.
29) Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, Cuellar T,
Haley B, Roose-Girma M, Phung QT. Caspase-11 cleaves gasdermin D for non-
canonical inflammasome signalling. Nature 2015; **526:** 666-671.
30) Hornung V, Ablasser A, Charrel-Dennis M, Bauernfeind F, Horvath G, Caffrey DR,
Latz E, Fitzgerald KA. AIM2 recognizes cytosolic dsDNA and forms a caspase-1-
activating inflammasome with ASC. Nature 2009; **458:** 514-518.
31) Xiao TS. The nucleic acid-sensing inflammasomes. Immunol Rev 2015; **265:**
103-111.
32) Shibata M, Ohtani R, Ihara M, Tomimoto H. White matter lesions and glial
activation in a novel mouse model of chronic cerebral hypoperfusion. Stroke
2004; **35:** 2598-2603.
33) Shibata M, Yamasaki N, Miyakawa T, Kalaria RN, Fujita Y, Ohtani R, Ihara M,
Takahashi R, Tomimoto H. Selective impairment of working memory in a mouse
model of chronic cerebral hypoperfusion. Stroke 2007; **38:** 2826-2832.
34) Feldman N, Rotter-Maskowitz A, Okun E. DAMPs as mediators of sterile
inflammation in aging-related pathologies. Ageing Res Rev 2015; **24:** 29-39.
35) Lamkanfi M, Dixit VM. Inflammasomes and their roles in health and disease.
Annu Rev Cell Dev Biol 2012; **28:** 137-161.
36) Schroder K, Tschopp J. The inflammasomes. Cell 2010; **140,** 821-832.
37) Lu A, Magupalli VG, Ruan J, Yin Q, Atianand MK, Vos MR, Schröder GF,
Fitzgerald KA, Wu H, Egelman EH. Unified polymerization mechanism for the
assembly of ASC-dependent inflammasomes. Cell 2014; **156:** 1193-1206.
38) Xue Y, Enosi Tuipulotu D, Tan WH, Kay C, Man SM. Emerging Activators and
Regulators of Inflammasomes and Pyroptosis. Trends Immunol 2019; **40:** 1035-
1052.
39) Shaw K, Bell L, Boyd K, Grijseels DM, Clarke D, Bonnar O, Crombag HS, Hall C
N. Hippocampus has lower oxygenation and weaker control of brain blood flow
than cortex, due to microvascular differences. BioRxiv, 2019; 835728.
40) Almeida A, Allen KL, Bates TE, Clark JB. Effect of Reperfusion Following
Cerebral Ischaemia on the Activity of the Mitochondrial Respiratory Chain in the
Gerbil Brain. Journal of Neurochemistry 2002; **65:** 1698-1703.
41) Belov Kirdajova D, Kriska J, Tureckova J, Anderova M. Ischemia-Triggered
Glutamate Excitotoxicity From the Perspective of Glial Cells. Frontiers in Cellular
Neuroscience, 2020; **14:** 51.
42) Heuser D, Guggenberger H. Ionic Changes in Brain Ischaemia and Alterations
Produced by Drugs. British Journal of Anaesthesia 1985; **57:** 23-33.
43) Briard B, Place DE, Kanneganti TD. DNA Sensing in the Innate Immune
Response. Physiology (Bethesda) 2020; **35:** 112-124.
44) Jakobs C, Perner S, Hornung V. AIM2 Drives Joint Inflammation in a Self-DNA
Triggered Model of Chronic Polyarthritis. PLoS One 2015; **10:** e0131702.
45) Bernbaum M, Menon BK, Fick G, Smith EE, Goyal M, Frayne R, Coutts SB.
Reduced blood flow in normal white matter predicts development of leukoaraiosis.
Journal of Cerebral Blood Flow & Metabolism 2015; **35:** 1610.
46) Fazekas F, Kleinert R, Offenbacher H, Schmidt R, Kleinert G, Payer F, Radner H,
Lechner H. Pathologic correlates of incidental MRI white matter signal
hyperintensities. Neurology 1993; **43,** 1683-1689.

47) Fernando MS, O'Brien JT, Perry RH, English P, Forster G, McMeekin W, Slade
JY, Golkhar A, Matthews FE, Barber R, Kalaria RN, Ince PG, Neuropathology
Group of MRC CFAS. Comparison of the pathology of cerebral white matter with
post-mortem magnetic resonance imaging (MRI) in the elderly brain.
Neuropathology and Applied Neurobiology 2004; **30:** 385-395.
48) Wardlaw JM, Valdés Hernández MC, Muñoz☐Maniega S. What are White Matter
Hyperintensities Made of? Journal of the American Heart Association 2015; **4:**
001140.
49) Mankin EA, Diehl GW, Sparks FT, Leutgeb S, Leutgeb JK. Hippocampal CA2
Activity Patterns Change over Time to a Larger Extent than between Spatial
Contexts. Neuron 2015; **85:** 190-201.
50) Mao H, Elkin BS, Genthikatti VV, Morrison B III, Yang KH. Why Is CA3 More
Vulnerable Than CA1 in Experimental Models of Controlled Cortical Impact-
Induced Brain Injury? Journal of Neurotrauma 2013; **30:** 1521.
51) Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, Phatnani HP,
Guarnieri P, Caneda C, Ruderisch N, et al. An RNA-Sequencing Transcriptome
and Splicing Database of Glia, Neurons, and Vascular Cells of the Cerebral
Cortex. The Journal of Neuroscience 2014; **34:** 11929.
52) Wu PJ, Liu HY, Huang TN, Hsueh YP. AIM 2 inflammasomes regulate neuronal
morphology and influence anxiety and memory in mice. Scientific Reports 2016;
6: 32405.
53) Brochard V, Combadière B, Prigent A, Laouar Y, Perrin A, Beray-Berthat V,
Bonduelle O, Alvarez-Fischer D, Callebert J, Launay JM, et al. Infiltration of
CD4+ lymphocytes into the brain contributes to neurodegeneration in a mouse
model of Parkinson disease. J Clin Invest 2009; **119:** 182-192.
54) Chu HX, Kim, HA, Lee S, Moore JP, Chan CT, Vinh A, Gelderblom M,
Arumugam TV, Broughton BR, Drummond GR, Sobey CG. Immune cell
infiltration in malignant middle cerebral artery infarction: comparison with
transient cerebral ischemia. J Cereb Blood Flow Metab 2014; **34:** 450-459.
55) Zhang SR, Piepke M, Chu HX, Broughton BR, Shim R, Wong CH, Lee S, Evans
MA, Vinh A, Sakkal S, et al. IL-33 modulates inflammatory brain injury but
exacerbates systemic immunosuppression following ischemic stroke. JCI Insight
2018; **3:** 18.
56) Roberts JM, Maniskas ME, Bix GJ. Bilateral carotid artery stenosis causes
unexpected early changes in brain extracellular matrix and blood-brain barrier
integrity in mice. PLoS One 2018; **13:** e0195765.
57) Kim H, Seo JS, Lee SY, Ha KT, Choi BT, Shin YI, Ju Yun Y, Shin HK. AIM2
inflammasome contributes to brain injury and chronic post-stroke cognitive
impairment in mice. Brain Behav Immun 2020; pii: S0889-1591(19)31515-6. doi:
10.1016/j.bbi.2020.03.011.
58) Ward R, Li W, Abdul Y, Jackson L, Dong G, Jamil S, Filosa J, Fagan SC, Ergul A.
NLRP3 inflammasome inhibition with MCC950 improves diabetes-mediated
cognitive impairment and vasoneuronal remodeling after ischemia. Pharmacol
Res 2019; **142:** 237-250.
952

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5