1 **AIM2 Inflammasome Mediates Hallmark Neuropathological Alterations**
2 **and Cognitive Impairment in a Mouse Model of Vascular Dementia**
3

4 Luting Poh[1,2], David Y. Fann[1,3,*], Peiyan Wong[2,4], Hong Meng Lim[1], Sok Lin Foo[1],
5 Sung-Wook Kang[1], Vismitha Rajeev[1,2], Sharmelee Selvaraji[1,2], Vinaya Rajagopal Iyer
6 [2], Nageiswari Parathy[2], Mohammad Badruzzaman Khan[5], David C. Hess[5], Dong-Gyu
7 Jo[6], Grant R. Drummond[7], Christopher G. Sobey[7], Mitchell K.P. Lai[2,8], Christopher Li-
8 Hsian Chen[2,8], Lina H. K. Lim[1], Thiruma V. Arumugam[1,6,7,*]
9
10
[1]Department of Physiology, Yong Loo Lin School of Medicine, National University of
Singapore, Singapore
[2]Department of Pharmacology, Yong Loo Lin School of Medicine, National University
of Singapore, Singapore
[3]Department of Biochemistry, Yong Loo Lin School of Medicine, National University
of Singapore, Singapore
[4]Neuroscience and Behavioural Disorders Programme, Duke-NUS Medical School,
Singapore
[5]Department of Neurology, Medical College of Georgia, Augusta University, Augusta,
GA, USA
[6]School of Pharmacy, Sungkyunkwan University, Suwon, Republic of Korea
[7]Department of Physiology, Anatomy and Microbiology, La Trobe University,
Bundoora, VIC, Australia
[8]Memory, Aging and Cognition Centre, National University Health System, Singapore
25
26
27 *Correspondence: Thiruma V. Arumugam, Department of Physiology, Anatomy and
28 Microbiology, La Trobe University, Bundoora, VIC, Australia. E-mail:
29 g.arumugam@latrobe.edu.au; David Y. Fann, Department of Biochemistry, Yong Loo
30 Lin School of Medicine, National University of Singapore, Singapore. E-mail:
31 bchfywd@nus.edu.sg

Abstract

Chronic cerebral hypoperfusion is associated with vascular dementia (VaD). Cerebral hypoperfusion may initiate complex molecular and cellular inflammatory pathways that contribute to long-term cognitive impairment and memory loss. Here we used a bilateral common carotid artery stenosis (BCAS) mouse model of VaD to investigate its effect on the innate immune response – particularly the inflammasome signaling pathway. Comprehensive analyses revealed that chronic cerebral hypoperfusion induces a complex temporal expression and activation of inflammasome components and their downstream products (IL-1β and IL-18) in different brain regions, and promotes activation of apoptotic and pyroptotic cell death pathways. Polarized glial cell activation, white matter lesion formation and hippocampal neuronal loss also occurred in a spatiotemporal manner. Moreover, in AIM2 knockout mice we observed attenuated inflammasome-mediated production of proinflammatory cytokines, apoptosis and pyroptosis, as well as resistance to chronic microglial activation, myelin breakdown, hippocampal neuronal loss, and behavioural and cognitive deficits following BCAS. Hence, we have demonstrated that activation of the AIM2 inflammasome substantially contributes to the pathophysiology of chronic cerebral hypoperfusion-induced brain injury and may therefore represent a promising therapeutic target for attenuating cognitive impairment in VaD.

Introduction

Vascular dementia (VaD) is the second most common form of dementia that is associated with vascular diseases resulting in dysfunction and damage to the cerebral vasculature, which causes disruption of cerebral blood flow, brain injury, and ultimately cognitive impairment and memory loss[1]. Clinical findings in VaD include chronic cerebral hypoperfusion, white matter lesions (WMLs), strokes, hippocampal atrophy, decline in executive functioning and memory impairment[2]. Recent evidence indicates that chronic cerebral hypoperfusion is associated with systemic inflammation before WMLs and neurological symptoms develop in VaD[3-5]. Interleukin 1-beta (IL-1β), a potent cytokine that orchestrates inflammatory pathways, is likely to be a key contributor to VaD[5]. The primary molecular machinery for IL-1β production in response involves assembly and activation of the inflammasome.

Inflammasome is an intracellular multi-protein complex that initiates an innate immune response, and is involved in multiple acute and chronic neurological diseases such as ischemic stroke, Alzheimer's disease (AD), Parkinson's Disease (PD), and amyotrophic lateral sclerosis (ALS)[6-11]. In the canonical inflammasome pathway, stimulation and homo-oligomerization of inflammasome receptors such as nucleotide-binding domain and leucine-rich repeat containing (NLR) family pyrin domain containing (NLRP) 1 (NLRP1), NLRP3, NLR family CARD domain containing 4 (NLRC4) and absent in melanoma 2 (AIM2) initiates formation of the inflammasome complex, recruitment and activation of adaptor (ASC) and effector (total caspase-1 and -8) inflammasome components into biologically active cleaved caspases-1 and -8[6,12]. In the noncanonical inflammasome pathway, caspase-11 is activated via auto-proteolysis upon dimerization to produce cleaved caspase-11. Upon inflammasome assembly and activation, activity of cleaved caspase-1 and -8 are increased and downstream proinflammatory cytokines such as mature IL-1β and IL-18 are generated[6]. This leads to activation of glial cells, endothelial dysfunction, and oligodendrocyte injury and demyelination, eventually resulting in neurovascular unit disintegration, neuronal loss and brain circuit dysfunction. Cleaved caspase-1 and -8 may also promote cell death by activating apoptosis and pyroptosis pathways[13-14]. Cleaved caspase-1 and -8 induces apoptosis and pyroptosis by catalyzing the cleavage of total caspase-3 and full-length Gasdermin D (GSDMD-FL) into cleaved caspase-3 and N-terminal Gasdermin D (GSDMD-NT), respectively. GSDMD-NT can oligomerize with other N-terminal units and cause pore formation in the nuclear and plasma membranes, leading to leakage of cellular content and proinflammatory cytokines[15,16]. GSDMD activation can also be facilitated directly by non-canonical cleaved caspase-11[17]. The inhibition of caspase-1 was demonstrated to prevent pyroptosis in experimental models of multiple sclerosis by reducing demyelination and neurodegeneration[18]. It was also established that the inflammasome complex also contributed to cognitive decline in a APP/PS1 mouse model of Alzheimer's disease[9,19]. Specifically, it was shown that deletion of the AIM2 inflammasome in that AD model promoted dendrite branching and synaptic plasticity, and improvement in spatial memory[20]. AIM2 is known to play a role in the host immune response to microbial DNA[21]; however, several reports have established that endogenous double-stranded DNA (dsDNA) can also activate AIM2 to form the AIM2

inflammasome[22,23]. Moreover, AIM2 also responds to dsDNA released from damaged
host cells and is a potential ligand that can activate AIM2 in neurons and glial cells
following chronic hypoperfusion[22, 24].

While evidence for direct involvement of the inflammasome complex in VaD is
lacking, a cytokine profile of plasma from VaD patients found IL-1β to be the most
abundant[25-28]. It was also reported that IL-1β impeded oligodendrocyte recruitment
and inhibited white matter repair and functional recovery during the early stages of
chronic cerebral hypoperfusion[5]. However, any potential involvement of
inflammasome activation in chronic cerebral hypoperfusion-induced glial activation,
white matter damage, neuronal cell death and cognitive dysfunction has not been
studied.

In the present study, we have comprehensively investigated the role of the
inflammasome signaling pathway in the bilateral common carotid artery stenosis
(BCAS) mouse model of VaD. The BCAS mouse model has been well established in
imitating a wide range of neurological clinical conditions of white matter rarefaction,
hippocampal atrophy and cognitive decline induced by chronic cerebral
hypoperfusion in humans[29,30]. Moreover, BCAS develops white matter lesions without
cerebral infarctions and optic nerve damage, while hippocampal neuronal death and
atrophy is induced making it the most appropriate animal model in studying VaD[31].
Using the BCAS mouse model, we report an increased expression and activation of
inflammasome complexes, production of mature proinflammatory cytokines,
polarized glial activation, WMLs, and neuronal cell death occurring in a time-
dependent manner in the cerebral cortex, hippocampus and striatum following BCAS.
Notably, these brain regions exhibited distinct profiles of inflammasome expression
and activation, and a specific prominent increase of AIM2 in the cerebral cortex and
hippocampus. Furthermore, we demonstrated that genetic deletion of AIM2 protects
against BCAS-induced inflammasome activation, proinflammatory cytokine
production, polarized glial activation, WML formation and neuronal cell death,
ultimately leading to an improvement in cognitive outcome. Overall, our findings
reveal that the AIM2 inflammasome may be a potential therapeutic target in VaD.

Materials and Methods

Experimental Animals and Genotyping

All *in vivo* experimental procedures were approved by the National University of Singapore, Singapore Animal Care and Use Committee and performed according to the guidelines set forth by the National Advisory Committee for Laboratory Animal Research (NACLAR), Singapore. Mice were housed in individual cages under standard laboratory conditions. All efforts were made to minimize suffering and numbers of animals used. All sections of the manuscript were performed and reported in accordance with ARRIVE (Animal Research: Reporting In Vivo Experiments) guidelines. Fourteen to sixteen weeks old wild type (WT) male C57BL/6 mice (weighing 24 to 30 grams) were obtained from In Vivos, Singapore. The AIM2 knockout (B6.129P2-Aim2<Gt(CSG445)Byg>/J; AIM2 KO) mice were generated by Ingenious Targeting Laboratory (Ronkonkoma, NY) on the C57BL/6 background mice via targeted replacement of the AIM2 coding region with a neomycin resistance gene through homologous recombination that was kindly obtained from our collaborator Professor Jenny P. Y. Ting[32]. Primer sequences used for PCR-based AIM2 deficient mouse genotyping are as follows: neomycin (KO) forward, GGAACTTCGCTAGACTAGTACGCGTG; neomycin (KO) reverse, CAACATTGTACAGATTGAGCAGG; AIM2 (WT) forward, GATGGAGAGTGAGTACCGGGAAATGCTGTT; and AIM2 (WT) reverse, TCTGCAAGTAGATTGGAGACAGACTCTGGTGA, resulting in a 450-bp band for the wild type and a 250-bp band for the targeted AIM2 knockout when PCR analysis was performed on the genomic DNA (Supplementary Fig. 1). All mice were given free access to food and water *ad libitum*. The experimental groups consisted of 20-25 animals in each group.

Bilateral Common Carotid Artery Stenosis (BCAS) Mouse Model

Animals were anaesthetized with isoflurane and subjected to BCAS, and subsequent chronic hypoperfusion using microcoils specifically designed for the mouse (microcoil specifications: piano wire diameter 0.08mm, internal diameter 0.18mm, coiling pitch 0.5mm, and total length 2.5mm; Sawane Spring Co Ltd, Japan) was described elsewhere[33]. The left and right common carotid arteries (CCAs) were exposed individually, freed from their sheaths, and a microcoil was twined by rotating

it around each CCA. The site of surgery was closed, and the mice were observed
and taken care of post-surgery until conscious and recovered to freely access food
and water ad libitum. To study the disease progression, the experimental animals
were divided into eight time-point groups: Sham, 1, 3, 7, 15, 21, 30 and 60-days
BCAS. Sham animals were given a skin incision and their CCAs were exposed. All
animals were euthanized at their respective end-point after BCAS for subsequent
analysis. For biochemical analysis involving Wild Type control and AIM2 KO mice,
the animals were divided into three experimental groups: Sham, 15 and 30-days
BCAS. However, only the Sham and 30-days timepoints were selected to challenge
the cognitive ability of both WT and AIM2 KO mice during behavioral analysis.
185

Measurements of Cerebral Blood Flow by Laser Speckle Contrast Imager
High-resolution Laser Speckle Contrast Imager (PSI system, Perimed Inc.) was
used to image cerebral blood perfusion and record cerebral blood flow (CBF) before
BCAS (baseline), immediately after BCAS surgery and finally at their respective end-
points of BCAS. As shown in Supplementary Fig. 2, regions of interest (ROIs)
between the bregma and lambda were selected for overall perfusion in the area of
two hemispheres. Body temperature was maintained at 37 ± 0.5 °C, and the skull
was shaved and exposed by a midline skin incision. The skull was cleaned gently
with sterile phosphate buffered saline (PBS) using a cotton applicator. Finally, the
image area was kept moist and a non-toxic silicon oil was applied on the skull, which
improved imaging. Perfusion images were acquired using the PSI system with a
70mW built-in laser diode for illumination and a 1388 x 1038 pixels CCD camera
installed 10cm above the skull (speed 19Hz, and exposure time 6mSec). Acquired
images were analyzed for changes in CBF (cerebral perfusion) using a dedicated
PIMSoft program (Perimed Inc.).
201

Sample Collection and Processing
Following each time points, mice were euthanized by administering a lethal dose
of inhaled carbon dioxide (CO_2) and the brains harvested. The different brain areas
(cerebral cortex, hippocampus and striatum) were immediately separated on ice and
were frozen in dry ice for tissue biochemical analysis (n = 7-8 in each experimental
group). A separate group of animals were sacrificed for histological analysis. Mice
were deeply anaesthetized, and perfusion fixation though the heart was performed

with 25mL of chilled 1xPBS (pH 7.4) then followed by 25mL chilled paraformaldehyde
(4%). Once the perfusion was completed, the brains were harvested and placed in
vials containing 4% paraformaldehyde solution for immersion-fixation overnight at
4°C (n = 5-7 in each experimental group).
213
Immunoblot Analysis
Immunoblot analysis was performed as described by us previously[34]. Briefly, brain
tissues were homogenized in lysis buffer and then combined with 2 x Laemelli buffer
(Bio-Rad Laboratories, Inc., Hercules, CA, USA). Protein samples were then
separated on 7.5 to 12.5% v/v sodium dodecyl sulfate (SDS) gels. The SDS-PAGE
gels were transferred onto nitrocellulose membranes to probe for proteins. Next, the
nitrocellulose membranes were incubated with the following primary antibodies:
NLRP1 (Santa Cruz, sc390133), NLRP3 (Adipogen, AG20B0014), NLRC4 (Millipore,
06-1125), AIM2 (Cell Signaling, #13095), IL-1β (Genetex, GTX74034), IL-18
(Biovision, 5180R-100), Total Caspase-1 and Cleaved Caspase-1 (p33 subunit)
(Adipogen, AG20B0042), Cleaved Caspase-1 (p20 subunit) (Cell Signaling, #67314),
Casapse-8 (Cell Signaling, #4927), Caspase-11 (Cell Signaling, #14340), Total
Caspase-3 (Cell Signaling, #9662), Cleaved Caspase-3 (Cell Signaling, #9664),
GSDMD (Cell Signaling, #93709), GSDMD-NT (Cell Signaling, #50928), GSDME
(Santa Cruz, sc393162), IBA-1 (Abcam, ab5076), GFAP (Cell Signaling, #12389),
and β-actin (Sigma-Aldrich, A5441) overnight at 4°C with agitation. Following primary
antibody incubation, membranes were washed three times with 1xTBST before
incubating with horseradish peroxidase (HRP)-conjugated secondary antibodies
(Goat Anti-Rabbit – Cell Signaling Technology, Danvers, MA, USA; Goat Anti-Mouse
– Sigma-Aldrich, St. Louis, MO, USA; Goat Anti-Rat – GE Healthcare Life Sciences,
Little Chalfont, UK) for 1hr at 24°C with agitation. Following secondary antibody
incubation, membranes were washed three times with 1xTBST, each time for 10 min.
The substrate for HRP, enhanced chemiluminescence (ECL) (Bio-Rad Laboratories,
Inc., Hercules, CA, USA) was applied before the membranes were imaged using
ChemiDocXRS+imaging system (Bio-Rad Laboratories, Inc., Hercules, CA, USA).
Quantification of proteins was conducted using Image J software (Version 1.46;
National Institute of Health, Bethesda, MD, USA), where protein densitometry was
expressed relative to the densitometry of the corresponding β-actin.

Luxol Fast Blue and Cresyl Violet Staining

Mouse brain tissues were fixed in 10% neutral buffered formalin and then processed into paraffin wax blocks. Coronal sections (4µm thick) were obtained via microtome sectioning. Axonal fibre density of Luxol-fast-blue (LFB) staining was performed to detect the severity of white matter (WM) lesions. Briefly, de-waxed rehydrated sections were immersed in the LFB solution (Abcam, UK) at 37°C overnight. Excess staining was removed by 95% ethanol treatment followed by washing with deionized water. Grey and white matter differentiation was initiated with the treatment of 0.05% aqueous lithium carbonate (Abcam, UK) for 20 seconds, followed by 70% ethanol until the nuclei are decolorized. Sections were immersed in Cresyl Violet solution (Abcam, UK) for 5 min and washed in deionized water. The sections were dehydrated in an ethanol gradient (70 – 100%), and finally cleared in xylene and mounted with a mounting agent. The bright field images were taken under 4X and 60X magnification using an Olympus upright Fluorescence Microscope BX53. The WM lesions were evaluated in five brain regions: the optic tract, internal capsule, caudoputamen, corpus callosum (Medial) and corpus callosum (Paramedian). The severity of the WM lesions was graded as normal (grade 0), disarrangement of the nerve fibres (grade 1), the formation of marked vacuoles (grade 2), and the disappearance of myelinated fibres (grade 3). The severity of neuropathology was scored by three blinded examiners and the number of neurons in CA1, CA2 and CA3 were counted as previously described[35]. The general morphology and the neurodegeneration in the hippocampus of BCAS animals were assessed by Cresyl violet staining.

Immunofluorescence Analysis

Paraffinized brain sections were cut from sham and BCAS animals and immunostained with primary antibodies against Cleaved Caspase-1 (Affinity Biosciences, AF4022), Cleaved Caspase-3 (Cell Signaling, #9661), GSDMD (Santa Cruz, sc393581), MBP (Cell Signaling, #78896), MAP2 (Millipore, MAB3418), GFAP (Cell Signaling, #3670; #12389), IBA-1 (Wako, 016-26721; Cell Signaling, #17198), CD86 (Santa Cruz, sc-28347), CD206 (eBioscience, 12-2069-42), C3 (Santa Cruz, sc-28294), S100S10 (Invitrogen, PA5-95505), OLIG2 (R&D System, AF2418) and PECAM-1 (BD Pharmingen, 553370). Images were captured with an Olympus FluoView FV1000 (Olympus, Japan) laser scanning confocal microscope using

20x/0.70 air objective and 100x/1.45 oil objective, with 488nm Argon ion and 543nm
HeNe laser as the excitation source. Single confocal images were converted to 512 x
512 pixel 12-bit TIFF images.

Behavioral Paradigm and Training of BCAS Animals

Visual Acuity Test: To assess visual detection, mice were placed on a platform
within a rotating drum covered with black and white vertical stripes at consistent
spatial frequency[36, 37]. The automated drum was then rotated in a clockwise direction
for 30sec, and subsequently an anti-clockwise direction for 30sec, after a 15sec inter-
trial interval. Both larger stripes (2°) and smaller stripes (1°) were tested for all
animals before conducting other behavioral tests.

Open Field Test: Locomotor activity was measured using an open field test[38].
Each subject was placed in the center of the open field apparatus (20×20×40cm;
Clever Sys Inc.; VA, USA). Total distance traveled (cm) and average speed (mm/s)
were recorded and tracked via TopScan (Clever Sys Inc.; VA, USA). Data was
collected for 35 minutes.

Morris Water Maze: Morris water maze was performed 3 weeks after surgery
(sham or BCAS), as described previously[39]. Mice were trained 4 times a day at 10-
min intervals for 5 consecutive days in this fixed platform training. In each trial, mice
were given 60secs to find the platform. Activity of the animal in the water maze was
video-tracked by EthoVision software (Noldus Information Technology, Leesburg,
VA), and the escape latency was recorded and analyzed. Long term spatial memory
was tested on the 6[th] day, whereby the platform was removed, and each animal was
given 60secs in the water maze after being released from four different directions.
Time spent around the platform, number of visits to the targeted quadrant and time
spent in the targeted quadrant on the probe day were tracked and analyzed. The
probe day was designed to fall between 32 to 34 days of BCAS surgery.

Double Stranded DNA Measurement

Double stranded DNA from the serum of BCAS animals were measured as
previously described using Quant-iT™ PicoGreen ® dsDNA Reagent and Kits. Briefly,
20µL of plasma sample collected was diluted with 1X TE solution. Subsequently,
equal volume of working solution of Quant-iT™ PicoGreen reagent was added. It was
then protected from light and incubated for 5 minutes at room temperature. Samples

were excited at 485nm and fluorescence intensity was measured at 520nm using a
Synergy HT multi-detection microplate reader (BioTek, Winnooski, VT).
312

Statistical Analysis
Experimental data were analyzed by GraphPad Prism 5.02 and 8.0 software
(GraphPad Software, San Diego, CA, USA). All values are expressed as mean ±
standard error of the mean (S.E.M). One-way Analysis of Variance (ANOVA) was
used, followed by a Bonferroni post-hoc test to determine differences between
groups. A P-value <0.05 was deemed to be statistically significant. For behavioral
data, non-parametric Kruskal-Wallis test was used to determine differences between
groups. A P-value <0.05 was deemed to be statistically significant.
321

Results
Chronic cerebral hypoperfusion increases inflammasome signalling
We first evaluated the protein expression of inflammasome receptors over 60 days
of BCAS in the cerebral cortex, hippocampus and striatum (Supplementary Figs. 3, 4
& 5). NLRP3, AIM2 and NLRC4 were all increased in the cerebral cortex, while only
AIM2 receptor expression was increased in the hippocampus compared to sham
controls (Supplementary Fig. 3a-d, 4). NLRP1 expression was increased in the
striatum while other inflammasome receptors remained unchanged (Supplementary
Figs. 5a & b). Expression of total caspase-8 and precursor IL-1β were increased in
the cerebral cortex after BCAS, whereas a decrease in precursor IL-18 was evident,
and ASC and total caspases-1 and -11 were unchanged (Supplementary Fig. 3e & f).
In the hippocampus, expression of total caspases-1, -8 -11 and precursor IL-18
increased, whereas ASC was decreased after BCAS (Supplementary Fig. 3g & h).
ASC, total caspase-8 and precursors for IL-1β and IL-18 were also increased in the
striatum following BCAS (Supplementary Fig. 5c & d).
338

To assess canonical inflammasome activation, expression of cleaved caspase-1
subunits (p33, transient active unit; p20, final by-product) and -8 were examined.
After BCAS, elevated expression of cleaved caspase-1 (p20) and -8 was detected in
the cerebral cortex (Fig. 1a & b) and also hippocampus (Fig. 1c & d). Mature IL-1β

and IL-18 – direct downstream markers of inflammasome activation – were also
higher in the cerebral cortex and hippocampus after BCAS versus sham controls
(Fig. 1a-d). Expression of non-canonical cleaved caspase-11 was also elevated in
the cerebral cortex and hippocampus after BCAS (Fig. 1a-d), whereas only cleaved
caspase-8, IL-1β and IL-18 were increased in the striatum (Supplementary Fig. 6).
Overall, the data reveal that inflammasome activation occurs in the brain in a spatial-
temporal manner as a result of chronic cerebral hypoperfusion.
350

Chronic cerebral hypoperfusion promotes apoptosis and pyroptosis
We next examined cell death mechanisms associated with inflammasome
activation. Thus, we assessed the expression of cleaved caspase-3 and GSDMD-NT,
along with their precursor proteins after BCAS (Fig. 1e-h; Supplementary Fig. 7 & 8).
Our data showed that in the cerebral cortex (Fig. 1e & f) and hippocampus (Fig. 1g &
h) the expression of cleaved caspase-3 and GSDMD-NT were increased following
BCAS compared to sham controls. Pyroptotic cell death (i.e. GSDMD-NT expression)
was more prominent in the cerebral cortex following BCAS (Fig. 1e & f), whereas
only apoptosis (i.e. increased cleaved caspase-3 expression) was detected in the
striatum following BCAS (Supplementary Fig. 8).
361

We also assessed inflammasome-associated secondary pyroptosis/necrosis,
which can be induced by cleaved caspase-3, as indicated by expression of N-
Terminal Gasdermin E (GSDME-NT). Similar to GSDMD-NT, inflammasome-
generated GSDME-NT can permeabilize nuclear and plasma membranes and
mitochondria, linking inflammasome activation to cell death[16]. We found GSDME-NT
to be increased in the cerebral cortex, but not in the hippocampus or striatum
following BCAS (Supplementary Fig. 7 & 8). Expression patterns of inflammasome-
related proteins and pathways following BCAS-induced chronic cerebral
hypoperfusion are summarized in Supplementary Fig. 9. Overall, the data indicate
that BCAS induces differential priming of inflammasome components, inflammasome
activation, and induction of apoptosis and pyroptosis pathways in the brain in a
spatial-temporal manner.
374
375

Chronic cerebral hypoperfusion induces glial cell activation, white matter lesions and hippocampal neuronal death

We next analyzed for associations between inflammasome activity and hallmark VaD pathologies involving glial cell activation, white matter integrity and neuronal loss. Expression of ionized calcium binding adaptor molecule 1 (Iba1) and glial fibrillary acidic protein (GFAP) indicated activation of microglia and astrocytes, respectively (Fig. 2a-d). Following BCAS, we found increased Iba-1 in the cerebral cortex (Fig. 2a & b) and increased GFAP in the hippocampus (Fig. 2c & d), and both markers increased in the striatum (Supplementary Fig. 10a & b). Subsequently, colocalization of both Iba1 and GFAP with respective polarization markers CD86, CD206, complement component 3 (C3) and S100A10 revealed activation of mainly M1 (CD86 positive) and M2 (CD206 positive) microglia upon chronic cerebral hypoperfusion in the cerebral cortex and hippocampus (Fig. 2e-h; Supplementary Fig. 11a-h) and striatum (Supplementary Fig. 10c-f). Myelin basic protein (MBP) and Luxol fast blue staining revealed myelin integrity, while cresyl violet and microtubule-associated protein 2 (MAP2) staining enabled assessment of neuronal loss in the hippocampus. Staining for MBP and MAP2 was reduced in the cerebral cortex (Fig. 2i & k), hippocampus (Fig. 2j & l) and striatum (Supplementary Fig. 10g & h) following 30 days of BCAS.

White matter integrity of five brain regions was assessed after 7-60 days of hypoperfusion, and found to be disrupted in the corpus callosum (Paramedian and Medial), caudoputamen, internal capsule and optic tract (Fig. 2m). All five areas exhibited time-dependent rarefaction of white matter (Fig. 2n) from 7 days, although the white matter lesion (WML) index was not increased in the optic tract until 60 days of BCAS (Fig. 2m & n). BCAS-induced neuronal loss was evident in hippocampal CA1, CA2 and CA3 areas (Fig. 2o-r). Whereas hippocampal sections from sham controls showed normal neuronal cell bodies with distinct nuclei, nucleoli, and densely packed neurons in all three hippocampal areas, there were signs of widespread neuronal loss evident after BCAS (Fig. 2o-r). Degeneration was more pronounced in the hippocampal CA2 and CA3 areas with severe atrophy from 7 days (Fig. 2o-r).

AIM2 activation mediates apoptosis and pyroptosis during chronic cerebral hypoperfusion

As AIM2 was upregulated in the cerebral cortex and hippocampus, we postulated its involvement in BCAS-induced brain injury. Analysis of serum revealed an increase in cell-free double stranded DNA (dsDNA) levels following BCAS, especially after 15 days (Fig. 3a). As dsDNA is the only known ligand that activates the AIM2 receptor[22, 24], the finding supports a potential involvement of the AIM2 inflammasome in BCAS-induced brain injury.

To further examine the role of AIM2 inflammasome activation on injury following chronic cerebral hypoperfusion, mice with AIM2 deficiency (AIM2 KO) were studied (Supplementary Fig. 1). We first confirmed via laser speckle contrast imaging that cerebral blood flow was equivalent under basal conditions and reduced following BCAS surgery in wild-type (WT) and AIM2 KO mice (Supplementary Fig. 12). This was similar to previously published data from BCAS in mice[29]. This BCAS-induced reduction in flow was sustained to a similar degree for at least 15 days in both genotypes but was slightly higher in AIM2 KO than WT mice at 30 days (Supplementary Fig. 12a-c).

We found no difference in the basal expression levels of cleaved caspase-1, -8, and -11, and both mature IL-1β and IL-18 (Fig. 3b-e) or in the levels of total caspases -1, -8 and -11, or IL-1β and IL-18 precursors in the cerebral cortex and hippocampus of WT and AIM2 KO mice (Supplementary Fig. 13a-d). However, there was evidence of reduced inflammasome activation following BCAS in AIM2 KO mice when compared to WT control (Fig. 3b-e). Of the three active effector proteins, only canonical cleaved caspase-1 expression was reduced in AIM2 KO mice compared to WT control following BCAS (Fig. 3b-e; Supplementary Fig. 14a & b). There was also reduced levels of mature IL-1β and IL-18 in the cerebral cortex and hippocampus of AIM2 KO mice at 15 and 30 days following BCAS (Fig. 3b-e; Supplementary Fig. 14a & b). These data thus indicated less inflammasome activity occurs after BCAS in the absence of the AIM2 receptor, especially mediated by canonical cleaved caspase-1. Furthermore, protein expression of both cleaved caspase-3 and GSDMD-NT was lower at 30 days after BCAS in the cerebral cortex (Fig. 3f & g) and hippocampus

(Fig. 3h & i) of AIM2 KO mice, indicating AIM2 inflammasome mediating apoptosis and pyroptosis under chronic cerebral hypoperfusion.

Immunofluorescence studies indicated that cleaved caspase-1 (CC1) activity was lower in AIM2 KO mice in cortical (Fig. 3j; Supplementary Fig. 15a-e & 16a-e) and hippocampal (Fig. 3k; Supplementary Fig. 15f-j & 16f-j) neurons (MAP2 positive) and microglia (Iba-1 positive) compared to WT controls at 30 days following BCAS. Immunoreactivity against cleaved caspase-1 was similar in cortical and hippocampal oligodendrocytes (OLIG2 positive), astrocytes (GFAP positive) and endothelial cells (PECAM-1 positive) of WT and AIM2 KO mice. Cellular specificity of AIM2 inflammasome-mediated cell death following BCAS was assessed by immunoreactivity against cleaved caspase-3 (CC3) and GSDMD in the cerebral cortex (Fig. 3j; Supplementary Fig. 15a-e; Supplementary Fig.17a-e & 18a-e) and hippocampus (Fig. 3k; Supplementary Fig. 15f-j; Supplementary Fig.17f-j & 18f-j). The data indicate that pro-apoptotic cleaved caspase-3 immunoreactivity was lower in cortical (Fig. 3j; Supplementary Fig. 15a-e & 17a-e) and hippocampal (Fig. 3k; Supplementary Fig. 15f-j & 17f-j) neurons (MAP2 positive) and microglia (Iba-1 positive) of AIM2 KO versus WT mice. Similarly, immunoreactivity of pro-pyroptotic GSDMD was reduced in neurons and microglia in the cortex (Fig. 3j; Supplementary Fig. 15a-e & 18a-e) and hippocampus (Fig. 3k; Supplementary Fig. 15f-j & 18f-j) of AIM2 KO mice. These findings were consistent with more neurons and fewer microglial activation in AIM2 KO than WT mice following chronic cerebral hypoperfusion.

AIM2 KO mice are resistant to microglial activation, myelin breakdown, hippocampal neuronal loss and cognitive deficits following chronic cerebral hypoperfusion

There was a lower expression of Iba-1 in the cerebral cortex and hippocampus of AIM2 KO than WT mice after BCAS (Fig. 4a-f), especially in CD86 positive inflammatory microglial cells (Fig. 4c and f and Supplementary Fig. 19a-h). These results suggest a role for AIM2 in microglial activation and polarization during chronic cerebral hypoperfusion. We next investigated if AIM2 KO mice are protected against BCAS-induced myelin injury. Although sham-operated WT and AIM2 KO mice

displayed no difference in protein expression and immunoreactivity of MBP, at 30

days after BCAS, there was lower expression and weaker immunoreactivity for MBP

in the cerebral cortex of WT mice than AIM2 KO mice (Fig. 4g-i). A similar difference

in MBP immunoreactivity between WT and AIM2 KO was seen in CA2 and CA3

hippocampal regions (Fig. 4j-l). As myelin integrity was healthier in AIM2 KO mice

following BCAS, we assessed whether these mice might be resistant to WML. After

BCAS, Luxol fast blue staining in WT mice indicated myelin rarefaction and vacuole

formation in several WM regions, but a smaller degree of damage was present in the

corpus callosum (paramedian or medial), caudoputamen or internal capsule regions

of AIM2 KO mice (Fig. 5a-e). Neuronal loss in hippocampal CA1, CA2 and CA3

regions was also less in AIM2 KO compared to WT mice at 15 and 30 days of

cerebral hypoperfusion (Fig. 5f-h). These data confirm a deleterious role of the AIM2

inflammasome in mediating VaD pathologies associated with chronic cerebral

hypoperfusion.

Finally, we tested for a role of the AIM2 inflammasome in BCAS-induced cognitive

decline. It was previously shown that BCAS-induced chronic cerebral hypoperfusion

in mice resulted in cognitive impairment such as working memory dysfunction and

memory loss[29,30]. To examine the effects of AIM2 KO against BCAS-induced

cognitive deficits, we conducted the Morris Water Maze and two other control tests

(Fig. 6a-h). To ensure that the animals were able to detect the visual cues during the

Morris Water Maze, a visual acuity test was conducted. All animals showed a 100%

passing rate, and so proceeded to the subsequent tests. In the open field test

conducted to investigate explorative and locomotor ability, WT BCAS mice did not

have a shorter distance travelled (Fig. 6a) or a lower average speed of movement

(Fig. 6b) when compared to WT Sham controls, indicating no impairment of

locomotor ability. Similarly, there was also no significant difference in the time spent

and the distance travelled in the outer and inner zone among the WT mice, showing

no BCAS-induced anxiety effects (Fig. 6c & d). However, it was observed that both

sham and BCAS AIM2 KO mice spent more time at the outer zone as compared to

the WT controls (Fig. 6c), indicating that AIM2 KO mice generally exhibit higher

levels of anxiety independent of BCAS. Several parameters were assessed in the

Morris Water Maze performance, including escape latency during five training days.

Generally, we observed a shorter time was taken to reach the platform each day

across all groups. Similar daily improvement was exhibited in WT or AIM2 KO mice
sham controls and AIM2 KO mice following BCAS, whereas WT mice subjected to
BCAS took approximately twice as long (Fig. 6e). Following five days of training,
spatial memory was assessed whereby WT mice subjected to BCAS spent less time
around the platform and significantly fewer visits to the target quadrant than WT
controls (Fig. 6f-h). By contrast, AIM2 KO mice subjected to BCAS spent a longer
time around the platform and visited the target quadrant more often than WT BCAS
mice (Fig. 6f-h). This suggest that AIM2 KO mice exhibit better memory retention as
compared to WT controls under similar BCAS conditions. Thus, the behavioral data
indicates that BCAS-induced cognitive impairments were reduced in mice deficient in
AIM2, suggesting a pivotal role for the AIM2 inflammasome in mediating VaD
pathology associated with chronic cerebral hypoperfusion.

Discussion

Our findings indicate that inflammasome receptors are expressed in the cortex and
hippocampus where the AIM2 inflammasome is activated in response to chronic
cerebral hypoperfusion, leading to cellular pathology and cognitive impairment in a
mouse model of vascular dementia. Genetic deficiency of AIM2 expression resulted
in less brain inflammation, white matter injury, neuronal loss and cognitive decline
during cerebral hypoperfusion. These data are the first to indicate that the AIM2
inflammasome promotes neuronal and white matter injury following cerebral
hypoperfusion, identifying AIM2 as a key contributor to sterile inflammatory
responses in vascular dementia.

Inflammasomes are intracellular multiprotein complexes composed of sensors for
various microbial components, viral RNA and damage- or danger-associated
molecular patterns (DAMPs) produced during cell injury[40,41]. These innate immune
complexes are categorized according to their structural characteristics as either
nucleotide-binding domain–like receptors (NLRs) or absent in melanoma 2 (AIM2)-
like receptors[41,42]. These receptors oligomerize in response to activation by external
stimuli and then recruit the adaptor protein, apoptosis-associated speck-like protein
containing a caspase recruitment domain (ASC)[43]. Ultimately, binding of caspase-1
results in its cleavage and activation, and the generation of mature pro-inflammatory

cytokines, IL-1β and IL-18, as well as pro-apoptotic cleaved caspase-3, and pro-
pyroptotic N-terminal GSDMD and GSDME[44]. While our study established the role of
AIM2 in chronic hypoperfusion-induced brain injury, it is still unknown whether other
inflammasomes might have a role in WML formation and neuronal cell death
following hypoperfusion. Although our data shows increased expression levels of
NLRP3 and NLRC4 in brain tissues following BCAS suggesting a potential
involvement of other inflammasome complexes, further studies are required to
establish the role of these inflammasomes in chronic hypoperfusion-induced brain
injury.

We previously demonstrated that inflammasomes contribute to neuronal cell death
following ischemic stroke[7], in part via NF-κB and MAPK signaling[34]. Our present data
indicate that chronic cerebral hypoperfusion may also lead to increased expression
and activation of NLRP and AIM2 inflammasome receptors in multiple brain areas,
especially in the hippocampal region, which is known to be particularly vulnerable to
injury following mild ischemia[45]. Cerebral hypoperfusion in the BCAS model used
here is known to impact the hippocampus by mechanisms involving ionic imbalance,
excitotoxicity, mitochondrial dysfunction and oxidative stress[46-48]. I'm not sure what
the purpose of this sentence is.

It is important to elucidate the mechanism(s) by which the AIM2 inflammasome
mediates chronic hypoperfusion-induced brain injury. Here we found that the plasma
concentration of dsDNA progressively increased following BCAS-induced cerebral
hypoperfusion, consistent with the possibility that this ligand promoted AIM2-
mediated amplification of brain inflammation and injury. Thus it is plausible that AIM2
functions as a DAMP sensor to promote sterile inflammation in vascular dementia.
We found that various white matter regions exhibited injury within 7 days of BCAS,
consistent with the known vulnerability of white matter to hypoperfusion. Reduced
blood flow to the white matter has been shown to predict the clinical development of
WMLs within 18 months[49]. We observed that AIM2 deficient animals subjected to
BCAS exhibited greater white matter integrity and lower WML severity scores than
WT mice after BCAS. Using magnetic resonance imaging, WMLs appear as white
matter hyperintensities (WMH) in VaD patients that arises due to demyelination and
axonal loss as well as microglial and endothelial activation[50-52]. In accordance with

previous studies, we found that neuronal cell death, loss of myelin basic protein and
microglial activation were present in our BCAS mouse model, which were reduced in
AIM2 deficient mice. Our data also showed that cerebral blood flow was significantly
improved in AIM2 KO mice compared to WT controls following 30 days of BCAS.
While we do not know the precise molecular and cellular mechanisms behind this
increase, it is possible that reduced inflammasome activation and brain injury may
protect the integrity of the neurovasculature to facilitate improved blood flow to the
brain.

The sub-regions of the dorsal hippocampus, CA1, CA2 and CA3, are responsible
for memory encoding and retrieval[53,54]. Our study showed that neurons in CA2 and
CA3 hippocampal regions experienced elevated levels of death sooner in
comparison to neurons in the CA1 hippocampal regions during chronic hypoperfusion.
Reduced levels of BCAS-induced neuronal loss occurred in hippocampal CA2 and
CA3 areas of AIM2-deficient animals when compared to WT controls, which is
consistent with reduced inflammasome-mediated cell death in the hippocampal
region of AIM2 KO mice, suggesting that the AIM2 inflammasome contributed to
neuronal loss in the hippocampus during cerebral hypoperfusion.

The AIM2 receptor is widely expressed in the brain, with its highest levels
occurring in microglia under physiological conditions[55]. Furthermore, specifically
within the cortex and hippocampus the AIM2 receptor is most abundantly expressed
in neurons.[56]. Our data are consistent with a role of AIM2 in these brain resident cells
and regions, but we cannot exclude circulating leukocytes as another possible cell
mediating the pro-inflammatory actions of AIM2 in the brain in chronic hypoperfusion.
Leukocytes contribute to other forms of brain injury and neurodegeneration[57-59] and
their infiltration from the circulation would be expected in association with reduced
blood-brain barrier integrity and blood vessel density known to occur in
hypoperfusion-induced VaD[60]. However, in this study we have not assessed the role
of leukocyte infiltration and related neutrophil extracellular traps (NETs), known as
NETosis, as a potential source of AIM2 ligand. It was previously established that
NETosis and its associated extracellular dsDNA is known to contribute to the
pathogenesis of number of inflammatory diseases[61]. In addition, recent evidences
suggest that neutrophils use an inflammasome- and GSDMD-dependent mechanism

to activate NETosis, suggesting that further studies are needed to understand the

role of NETosis in the pathology of VaD[62].

Previous studies have recently reported that the AIM2 inflammasome contributes

to brain injury and chronic post-stroke cognitive impairment in mice[63], and to reduced

neuroplasticity and spatial memory in a mouse model of AD[20]. Similarly, there is also

evidence supporting a role for the NLRP3 inflammasome in cognitive impairment in

models of dementia and diabetes[9,64]. Inflammation is known to adversely impact

cognitive function, and our data are consistent with inflammasomes promoting an

innate immune response to mediate such an effect during cerebral hypoperfusion.

Moreover, we observed in our data that AIM2 KO mice generally exhibited higher

levels of anxiety independent of BCAS. This was reported by a previous study that

established that AIM2 KO mice exhibited increased anxious behaviors and reduced

auditory fear memory at the physiological level[65].

In summary, we have provided substantial evidence that activation of the AIM2

inflammasome plays a key role in promoting brain inflammation, white matter lesions,

neuronal cell death and cognitive impairment induced by chronic cerebral

hypoperfusion. These effects of AIM2 involve promotion of apoptosis and pyroptosis

of cortical and hippocampal neurons. As inflammasome activation plays a major role

in a number of inflammatory diseases, there are substantial efforts to develop

inflammasome inhibitors. However, application of inflammasome inhibitors is limited

in clinical conditions as clinical trials have not yet been completed. While our study

establishes that the AIM2 inflammasome may therefore represent a promising

therapeutic target for attenuating cognitive impairment in VaD, further research is

warranted to develop specific AIM2 inhibitors to test their effect in experimental and

clinical studies.

Acknowledgements

We thank Professor Jenny P. Ting (University of North Carolina, Chapel Hill, Chapel Hill, NC, USA) for providing the AIM2 deficient mice. Supplementary Figure 1 and Supplementary Figure 9a in this article was created using BioRender.

Funding

This work was supported by the National Medical Research Council Research Grants (NMRC-CBRG-0102/2016 and NMRC/OFIRG/0036/2017), Singapore and the startup fund to TVA from La Trobe University, Melbourne, Australia.

Author Contributions

Study Conception & Design: T.V.A., L.P., and D.Y.F.; Experiment or Data Collection: T.V.A., L.P., P.W., H.M.L., S.L.F., S.W.K., V.R., S.S., D.Y.F., R.I.V., and N.P.; Data Analysis: L.P., P.W., V.R., S.S., D.Y.F., and T.V.A.; Data Interpretation: L.P., T.V.A., and D.Y.F; Writing-Manuscript Preparation and intellectual input: T.V.A., L.P., D.Y.F., M.B.K., G.R.D., C.G.S., D.C.H., D.G.J., M.K.P.L., C.L.H.C., and L.H.K.L.; Supervision & Administration: T.V.A., D.Y.F., M.K.P.L., and C.L.H.C.

Conflicts Of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

REFERENCES

679
680

1) Wolters FJ, Ikram MA. Epidemiology of Vascular Dementia. Arterioscler Thromb
Vasc Biol 2019; **39:**1542-1549.
2) Van der Flier WM, Skoog I, Schneider JA, Pantoni L, Mok V, Chen CLH, et al.
Vascular cognitive impairment. Nat Rev Dis Primers 2018; **4:** 18003.
3) Alber J, Alladi S, Bae HJ, Barton DA, Beckett LA, Bell JM, et al. White matter
hyperintensities in vascular contributions to cognitive impairment and dementia
(VCID): Knowledge gaps and opportunities. Alzheimers Dement (N Y) 2019; **5:**
107-117.
4) Schmitz M, Hermann P, Oikonomou P, Stoeck K, Ebert E, Poliakova T, et al.
Cytokine profiles and the role of cellular prion protein in patients with vascular
dementia and vascular encephalopathy. Neurobiol Aging 2015; **36:** 2597-2606.
5) Zhou Y, Zhang J, Wang L, Chen Y, Wan Y, He Y, et al. Interleukin-1β impedes
oligodendrocyte progenitor cell recruitment and white matter repair following
chronic cerebral hypoperfusion. Brain Behav Immun 2017; **60:** 93-105.
6) Fann DY, Lee SY, Manzanero S, Chunduri P, Sobey CG, Arumugam TV.
Pathogenesis of acute stroke and the role of inflammasomes. Ageing Res Rev
2013; **12:** 941-966.
7) Fann DY, Lee SY, Manzanero S, Tang SC, Gelderblom M, Chunduri P, et al.
Intravenous immunoglobulin suppresses NLRP1 and NLRP3 inflammasome-
mediated neuronal death in ischemic stroke. Cell Death Dis 2013; **4:** e790.
8) Gordon R, Albornoz EA, Christie DC, Langley MR, Kumar V, Mantovani S, et al.
Inflammasome inhibition prevents α-synuclein pathology and dopaminergic
neurodegeneration in mice. Sci Transl Med 2018; **10:** 465.
9) Heneka MT, Kummer MP, Stutz A, Delekate A, Schwartz S, Vieira-Saecker A, et
al. NLRP3 is activated in Alzheimer's disease and contributes to pathology in
APP/PS1 mice. Nature 2013; **493:** 674-678.
10) Johann S, Heitzer M, Kanagaratnam M, Goswami A, Rizo T, Weis J, et al.
NLRP3 inflammasome is expressed by astrocytes in the SOD1 mouse model of
ALS and in human sporadic ALS patients. Glia 2015; **63:** 2260-2273.
11) Poh L, Kang SW, Baik SH, Ng GYQ, She DT, Balaganapathy P, et al. Evidence
that NLRC4 inflammasome mediates apoptotic and pyroptotic microglial death
following ischemic stroke. Brain Behav Immun 2019; **75:** 34-47.
12) Fritsch M, Günther SD, Schwarzer R, Albert MC, Schorn F, Werthenbach JP, et
al. Caspase-8 is the molecular switch for apoptosis, necroptosis and pyroptosis.
Nature 2019; **575:** 683-687.
13) Ball DP, Taabazuing CY, Griswold AR, Orth EL, Rao SD, Kotliar IB, et al.
Caspase-1 interdomain linker cleavage is required for pyroptosis. Life Sci
Alliance 2020; **3:** 3.
14) Yuan B, Zhou XM, You ZQ, Xu WD, Fan JM, Chen SJ, et al. Inhibition of AIM2
inflammasome activation alleviates GSDMD-induced pyroptosis in early brain
injury after subarachnoid haemorrhage. Cell Death Dis 2020; **11:** 76.
15) Liu X, Zhang Z, Ruan J, Pan Y, Magupalli VG, Wu H, et al. Inflammasome-
activated gasdermin D causes pyroptosis by forming membrane pores. Nature
2016; **535:** 153-158.
16) Rogers C, Erkes DA, Nardone A, Aplin AE, Fernandes-Alnemri T, Alnemri ES.
Gasdermin pores permeabilize mitochondria to augment caspase-3 activation
during apoptosis and inflammasome activation. Nat Commun 2019; **10:** 1689.

17) Kayagaki N, Stowe IB, Lee BL, O'Rourke K, Anderson K, Warming S, et al.
Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling.
Nature 2015; **526:** 666-671.
18) McKenzie BA, Mamik MK, Saito LB, Boghozian R, Monaco MC, Major EO, et al.
Caspase-1 inhibition prevents glial inflammasome activation and pyroptosis in
models of multiple sclerosis. Proc Natl Acad Sci USA 2018; **115:** E6065-E6074.
19) Dempsey C, Rubio Araiz A, Bryson KJ, Finucane O, Larkin C, et al. Inhibiting the
NLRP3 inflammasome with MCC950 promotes non-phlogistic clearance of
amyloid-β and cognitive function in APP/PS1 mice. Brain Behav Immun 2017;
61: 306-316.
20) Chen J, Shu S, Chen Y, Liu Z, Yu L, Yang L, et al. AIM2 deletion promotes
neuroplasticity and spatial memory of mice. Brain Res Bull 2019; **152:** 85-94.
21) Briard B, Place DE, Kanneganti TD. DNA Sensing in the Innate Immune
Response. Physiology (Bethesda) 2020; **35:** 112-124.
22) Hornung V, Ablasser A, Charrel-Dennis M, Bauernfeind F, Horvath G, Caffrey DR,
et al. AIM2 recognizes cytosolic dsDNA and forms a caspase-1-activating
inflammasome with ASC. Nature 2009; **458:** 514-518.
23) Jakobs C, Perner S, Hornung V. AIM2 Drives Joint Inflammation in a Self-DNA
Triggered Model of Chronic Polyarthritis. PLoS One 2015; **10:** e0131702.
24) Xiao TS. The nucleic acid-sensing inflammasomes. Immunol Rev 2015; **265:**
103-111.
25) Paganelli R, Di Iorio A, Patricelli L, Ripani F, Sparvieri E, Faricelli R, et al.
Proinflammatory cytokines in sera of elderly patients with dementia: levels in
vascular injury are higher than those of mild-moderate Alzheimer's disease
patients. Exp Gerontol 2002; **37:** 257-263.
26) Yasutake C, Kuroda K, Yanagawa T, Okamura T, Yoneda H. Serum BDNF, TNF-
alpha and IL-1beta levels in dementia patients: comparison between Alzheimer's
disease and vascular dementia. Eur Arch Psychiatry Clin Neurosci 2006; **256:**
402-406.
27) Zuliani G, Ranzini M, Guerra G, Rossi L, Munari MR, Zurlo A, et al. Plasma
cytokines profile in older subjects with late onset Alzheimer's disease or vascular
dementia. J Psychiatr Res 2007; **41:** 686-693.
28) Mulugeta E, Molina-Holgado F, Elliott MS, Hortobagyi T, Perry R, Kalaria RN, et
al. Inflammatory mediators in the frontal lobe of patients with mixed and vascular
dementia. Dement Geriatr Cogn Disord 2008; **25:** 278-286.
29) Shibata M, Ohtani R, Ihara M, Tomimoto H. White matter lesions and glial
activation in a novel mouse model of chronic cerebral hypoperfusion. Stroke
2004; **35:** 2598-2603.
30) Shibata M, Yamasaki N, Miyakawa T, Kalaria RN, Fujita Y, Ohtani R, et al.
Selective impairment of working memory in a mouse model of chronic cerebral
hypoperfusion. Stroke 2007; **38:** 2826-2832.
31) Washida K, Hattori Y, Ihara M. Animal Models of Chronic Cerebral
Hypoperfusion: From Mouse to Primate. Int J Mol Sci. 2019; **20:** 6176.
32) Wilson JE, Petrucelli AS, Chen L, Koblansky AA, Truax AD, Oyama Y, et al.
Inflammasome-independent role of AIM2 in suppressing colon tumorigenesis via
DNA-PK and Akt. Nat Med 2015; **21:** 906-913.
33) Khan MB, Hafez S, Hoda MN, Baban B, Wagner J, Awad ME, et al. Chronic
Remote Ischemic Conditioning Is Cerebroprotective and Induces Vascular
Remodeling in a VCID Model. Transl Stroke Res 2018; **9:** 51-63.

34) Fann DY, Lim YA, Cheng YL, Lok KZ, Chunduri P, Baik SH, et al. Evidence that NF-κB and MAPK Signaling Promotes NLRP Inflammasome Activation in Neurons Following Ischemic Stroke. Mol Neurobiol 2018; **55**: 1082-1096.

35) Nishio K, Ihara M, Yamasaki N, Kalaria RN, Maki T, Fujita Y, et al. A Mouse Model Characterizing Features of Vascular Dementia With Hippocampal Atrophy. Stroke 2010; **41**: 1278–1284.

36) Thaung C, Arnold K, Jackson IJ, Coffey PJ. Presence of visual head tracking differentiates normal sighted from retinal degenerate mice. Neurosci Lett. 2002; **325:** 21-24.

37) Mitchiner JC, Pinto LH, Vanable JW Jr. Visually evoked eye movements in the mouse (Mus musculus). Vision Res. 1976; **16**: 1169-1171.

38) Wong P, Chang CC, Marx CE, Caron MG, Wetsel WC, Zhang X. Pregnenolone rescues schizophrenia-like behavior in dopamine transporter knockout mice. PLoS One. 2012; **7:** e51455.

39) Madar R, Rotter A, Waldman Ben-Asher H, Mughal MR, Arumugam TV, Wood WH 3rd, et al. Postnatal TLR2 activation impairs learning and memory in adulthood. Brain Behav Immun. 2015; **48:** 301-312.

40) Feldman N, Rotter-Maskowitz A, Okun E. DAMPs as mediators of sterile inflammation in aging-related pathologies. Ageing Res Rev 2015; **24:** 29-39.

41) Lamkanfi M, Dixit VM. Inflammasomes and their roles in health and disease. Annu Rev Cell Dev Biol 2012; **28:** 137-161.

42) Schroder K, Tschopp J. The inflammasomes. Cell 2010; **140,** 821-832.

43) Lu A, Magupalli VG, Ruan J, Yin Q, Atianand MK, Vos MR, et al. Unified polymerization mechanism for the assembly of ASC-dependent inflammasomes. Cell 2014; **156:** 1193-1206.

44) Xue Y, Enosi Tuipulotu D, Tan WH, Kay C, Man SM. Emerging Activators and Regulators of Inflammasomes and Pyroptosis. Trends Immunol 2019; **40:** 1035-1052.

45) Shaw K, Bell L, Boyd K, Grijseels DM, Clarke D, Bonnar O, et al. Hippocampus has lower oxygenation and weaker control of brain blood flow than cortex, due to microvascular differences. BioRxiv, 2019; 835728.

46) Almeida A, Allen KL, Bates TE, Clark JB. Effect of Reperfusion Following Cerebral Ischaemia on the Activity of the Mitochondrial Respiratory Chain in the Gerbil Brain. Journal of Neurochemistry 2002; **65:** 1698-1703.

47) Belov Kirdajova D, Kriska J, Tureckova J, Anderova M. Ischemia-Triggered Glutamate Excitotoxicity From the Perspective of Glial Cells. Frontiers in Cellular Neuroscience, 2020; **14:** 51.

48) Heuser D, Guggenberger H. Ionic Changes in Brain Ischaemia and Alterations Produced by Drugs. British Journal of Anaesthesia 1985; **57:** 23-33.

49) Bernbaum M, Menon BK, Fick G, Smith EE, Goyal M, Frayne R, et al. Reduced blood flow in normal white matter predicts development of leukoaraiosis. Journal of Cerebral Blood Flow & Metabolism 2015; **35:** 1610.

50) Fazekas F, Kleinert R, Offenbacher H, Schmidt R, Kleinert G, Payer F, et al. Pathologic correlates of incidental MRI white matter signal hyperintensities. Neurology 1993; **43,** 1683-1689.

51) Fernando MS, O'Brien JT, Perry RH, English P, Forster G, McMeekin W, et al. Comparison of the pathology of cerebral white matter with post-mortem magnetic resonance imaging (MRI) in the elderly brain. Neuropathology and Applied Neurobiology 2004; **30:** 385-395.

52) Wardlaw JM, Valdés Hernández MC, Muñoz☐Maniega S. What are White Matter
Hyperintensities Made of? Journal of the American Heart Association 2015; **4:**
001140.
53) Mankin EA, Diehl GW, Sparks FT, Leutgeb S, Leutgeb JK. Hippocampal CA2
Activity Patterns Change over Time to a Larger Extent than between Spatial
Contexts. Neuron 2015; **85:** 190-201.
54) Mao H, Elkin BS, Genthikatti VV, Morrison B III, Yang KH. Why Is CA3 More
Vulnerable Than CA1 in Experimental Models of Controlled Cortical Impact-
Induced Brain Injury? Journal of Neurotrauma 2013; **30:** 1521.
55) Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, et al. An
RNA-Sequencing Transcriptome and Splicing Database of Glia, Neurons, and
Vascular Cells of the Cerebral Cortex. The Journal of Neuroscience 2014; **34:**
11929.
56) Wu PJ, Liu HY, Huang TN, Hsueh YP. AIM 2 inflammasomes regulate neuronal
morphology and influence anxiety and memory in mice. Scientific Reports 2016;
6: 32405.
57) Brochard V, Combadière B, Prigent A, Laouar Y, Perrin A, Beray-Berthat V, et al.
Infiltration of CD4+ lymphocytes into the brain contributes to neurodegeneration
in a mouse model of Parkinson disease. J Clin Invest 2009; **119:** 182-192.
58) Chu HX, Kim HA, Lee S, Moore JP, Chan CT, Vinh A, et al. Immune cell
infiltration in malignant middle cerebral artery infarction: comparison with
transient cerebral ischemia. J Cereb Blood Flow Metab 2014; **34:** 450-459.
59) Zhang SR, Piepke M, Chu HX, Broughton BR, Shim R, Wong CH, et al. IL-33
modulates inflammatory brain injury but exacerbates systemic
immunosuppression following ischemic stroke. JCI Insight 2018; **3:** 18.
60) Roberts JM, Maniskas ME, Bix GJ. Bilateral carotid artery stenosis causes
unexpected early changes in brain extracellular matrix and blood-brain barrier
integrity in mice. PLoS One 2018; **13:** e0195765.
61) Liu J, Yang D, Wang X, Zhu Z, Wang T, Ma A, et al. Neutrophil extracellular traps
and dsDNA predict outcomes among patients with ST-elevation myocardial
infarction. Sci Rep. 2019; **9:** 11599.
62) Chen KW, Monteleone M, Boucher D, Sollberger G, Ramnath D, Condon ND, et
al. Noncanonical inflammasome signaling elicits gasdermin D-dependent
neutrophil extracellular traps. Sci Immunol. 2018; **3:** eaar6676.
63) Kim H, Seo JS, Lee SY, Ha KT, Choi BT, Shin YI, et al. AIM2 inflammasome
contributes to brain injury and chronic post-stroke cognitive impairment in mice.
Brain Behav Immun 2020; **87:** 765-776.
64) Ward R, Li W, Abdul Y, Jackson L, Dong G, Jamil S, et al. NLRP3 inflammasome
inhibition with MCC950 improves diabetes-mediated cognitive impairment and
vasoneuronal remodeling after ischemia. Pharmacol Res 2019; **142:** 237-250.
65) Wu PJ, Liu HY, Huang TN, Hsueh YP. AIM 2 inflammasomes regulate neuronal
morphology and influence anxiety and memory in mice. Sci Rep. 2016; **6:** 32405.

 FIGURE LEGENDS

Figure 1: Effect of chronic cerebral hypoperfusion on inflammasome activation

and cell death in the cerebral cortex and hippocampus over time following

BCAS. (a & b), representative immunoblots and quantification illustrating increased

levels of activated inflammasome effector proteins such as cleaved caspase-1 (p20),

-8, and -11, and maturation of downstream effector targets, IL-1β and IL-18, in the

cerebral cortex. (c & d), representative immunoblots and quantification illustrating

increased levels of activated inflammasome effector proteins such as cleaved

caspase-1 (p20, p33), -8, and -11, and maturation of downstream effector targets, IL-

1β and IL-18, in the hippocampus. (e & f), representative immunoblots and

quantification illustrating increases in the expression of apoptotic marker cleaved

caspase-3 and pyroptotic marker N-terminal GSDMD in the cerebral cortex. (g & h),

representative immunoblots and quantification illustrating increases in the expression

of apoptotic marker cleaved caspase-3 and pyroptotic marker N-terminal GSDMD in

the hippocampus. β-actin was used as a loading control. Data are represented as

mean ± S.E.M. n=6-7 mice in each experimental group. *P<0.05 compared with

Sham; **P<0.01 compared with Sham; ***P<0.001 compared with Sham.

Abbreviations: BCAS, bilateral common carotid artery stenosis; Cl, cleaved; FL, full

length; NT, N-terminal; GSDMD, gasdermin D; GSDME, gasdermin E.

Figure 2: Effect of chronic cerebral hypoperfusion on the levels of glial

activation, white matter integrity and hippocampal neuronal density in the

cerebral cortex and hippocampus following BCAS. (a), representative

immunoblots and quantification illustrating increased microglial activation due to

increased levels of Iba-1 in the cerebral cortex over time following BCAS. Data are

represented as mean ± S.E.M. n=6-7 mice in each experimental group. β-actin was

used as a loading control. *P<0.05 compared with Sham. (b), representative

immunofluorescence analysis of Iba-1 and GFAP in the cerebral cortex provide

supporting evidence of microglial activation following BCAS. Magnification x 100.

Scale bar, 20µm. Images were taken under identical exposures and conditions. (c),

representative immunoblots and quantification illustrating increased astroglia

activation due to increased levels of GFAP in the hippocampus over time following

BCAS. Data are represented as mean ± S.E.M. n=6-7 mice in each experimental

group. β-actin was used as a loading control. *P<0.05 compared with Sham;
**P<0.01 compared with Sham; ***P<0.001 compared with Sham. (d), representative
immunofluorescence analysis of Iba-1 and GFAP in the hippocampus provide
supporting evidence of astroglia activation following BCAS. Magnification x 100.
Scale bar, 20µm. Images were taken under identical exposures and conditions. (e &
f), representative merged immunofluorescence images of DAPI, CD86 and CD206
co-localized within microglia (Iba-1 positive) in the cerebral cortex and hippocampus
of WT controls following BCAS. Representative merged immunofluorescence images
illustrate the activation of CD86 (M1) and CD206 (M2) positive microglia in the
cerebral cortex (e) and hippocampus (f) following BCAS. (g & h), representative
merged immunofluorescence images of DAPI, Complement C3 (C3) and S100A10
within astrocytes (GFAP positive) in the cerebral cortex and hippocampus of WT
controls following BCAS. Other than colocalization of S100A10 (A2) within astrocytes
in cerebral cortex (g), no substantial co-localization of C3 (A1) was observed in the
cerebral cortex and hippocampus following BCAS. This illustrates activation of A2
astrocytes in the cerebral cortex (g) upon BCAS. Zoom Magnification x 100. Scale
bar, 20µm. (i & j), representative immunofluorescence images illustrating a loss of
myelin due to decreased levels of MBP immunoreactivity in the cerebral cortex and
hippocampus following BCAS. (k & l), representative immunofluorescence images
illustrating a loss of neurons due to decreased levels of MAP2 immunoreactivity in
the cerebral cortex and hippocampus following BCAS. Magnification x 20. Scale bar,
120µm. (m & n), representative Luxol fast blue stained images and quantification
illustrating disruption of white matter integrity due to increased myelin rarefaction and
white matter lesion formation in the corpus callosum (paramedian), corpus callosum
(medial), caudoputamen, internal capsule and optic tract over time following BCAS.
The severity of white matter disruption was graded accordingly: Grade 0 = no
disruption; Grade 1 = disarrangement of nerve fibers; Grade 2 = formation of marked
vacuoles; Grade 3 = disappearance of myelinated fibers. Images were taken under
identical exposures and conditions. (o-r), representative crystal violet images and
quantification illustrating loss of Nissl positively stained neurons in hippocampal CA1,
CA2 and CA3 regions over time following BCAS. Data are represented as mean ±
S.E.M. n=5-7 mice in each experimental group. *P<0.05 compared with Sham;
**P<0.01 compared with Sham; ***P<0.001 compared with Sham. Magnification x 60.
Scale bar, 20 µm. Images were taken under identical exposures and conditions.

Abbreviations: BCAS, bilateral common carotid artery stenosis; CD86, cluster of
differentiation 86; CD206, cluster of differentiation 206; C3, complement component
3; GFAP, glial fibrillary acidic protein; Iba-1, ionized calcium binding adaptor
molecule-1; MBP, myelin basic protein; MAP2, microtubule-associated protein 2;
S100A10, S100 calcium-binding protein A10.

Figure 3: Effect of chronic cerebral hypoperfusion on inflammasome-mediated
programmed cell death in AIM2 KO mice following BCAS. (a), quantification
illustrating increased serum concentrations of the AIM2 receptor ligand, cell free
double stranded DNA (dsDNA), over time in wild-type mice following BCAS. Data are
represented as mean ± S.E.M. n=5-7 mice in each experimental group. *P<0.05
compared with Sham. (b-e), representative immunoblots and quantification
illustrating suppression of inflammasome activation due to decreased expression of
cleaved caspase-1 (p20) at Day 15 and 30, and maturation of IL-1β cytokine
production at Day 30 in the cerebral cortex (b & c) and hippocampus (d & e) of AIM2
KO mice compared to WT controls following BCAS. (f-i), representative immunoblots
and quantification illustrating attenuation of apoptotic and pyroptotic cell death due to
decreased expression of cleaved caspase-3 and GSDMD-NT, respectively, at Day
30 in the cerebral cortex (f & g) and hippocampus (h & i) of AIM2 KO mice compared
to WT controls following BCAS. β-actin was used as a loading control. Data are
represented as mean ± S.E.M. n=5-8 mice in each experimental group. *P<0.05
compared with WT Sham; **P<0.01 compared with WT Sham; ***P<0.001 compared
with WT Sham. +P<0.05 compared with WT BCAS; ++P<0.01 compared with WT
BCAS; +++P<0.001 compared with WT BCAS. (j & k), representative merged
immunofluorescence images of DAPI, cleaved caspase-1 p10 (CC1), cleaved
caspase-3 (CC3) and GSDMD co-localized within neurons (MAP2 positive), microglia
(Iba-1 positive), oligodendrocytes (OLIG2 positive) and endothelial cells (PECAM-1
positive) in the cerebral cortex and hippocampus of WT controls following BCAS. No
substantial co-localization of cleaved caspase-1 p10, cleaved caspase-3 and
GSDMD was observed in astrocytes (GFAP positive) in the cerebral cortex and
hippocampus of WT controls following BCAS. (j & k), representative merged
immunofluorescence images illustrate a reduction in inflammasome activation, and
apoptotic and pyroptotic cell death due to decreased expression levels of cleaved
caspase-1, cleaved caspase-3 and GSDMD, respectively, in both neurons and

microglia in the cerebral cortex (j) and hippocampus (k) in AIM2 KO mice compared

to WT controls following BCAS. Zoom Magnification x 100. Scale bar, 20 µm.

Images were taken under identical exposures and conditions. Abbreviations: BCAS,

bilateral common carotid artery stenosis; Cl, cleaved; FL, full length; NT, N-terminal;

WT, wild-type; KO, knock out; CC1, cleaved caspase-1; CC3, cleaved caspase-3;

GSDMD, Gasdermin D; MAP2, microtubule-associated protein 2; Iba-1, ionized

calcium binding adaptor molecule-1; OLIG2, oligodendrocyte transcription factor 2;

GFAP, glial fibrillary acidic protein; PECAM-1, platelet endothelial cell adhesion

molecule-1.

Figure 4: Effect of chronic cerebral hypoperfusion on glial activation and

myelin expression in the cerebral cortex and hippocampus of AIM2 KO mice

following BCAS. (a & b, d & e), representative immunoblots and quantification

illustrating resistance to microglial activation due to decreased expression of Iba-1 in

the cerebral cortex and hippocampus of AIM2 KO mice compared to WT controls

following BCAS. No effect on astroglia activation was observed as the expression of

GFAP in the cerebral cortex and hippocampus remained unchanged in AIM2 KO

mice compared to WT controls following BCAS. β-actin was used as a loading control.

Data are represented as mean ± S.E.M. n=6-8 mice in each experimental group.

*$P<0.05$ compared with WT Sham; **$P<0.01$ compared with WT Sham. +$P<0.05$

compared with WT BCAS; ++$P<0.01$ compared with WT BCAS; +++$P<0.001$

compared with WT BCAS. (c & f), representative merged immunofluorescence

images of DAPI, CD86 and CD206 co-localized within microglia (Iba-1 positive) in the

cerebral cortex and hippocampus of WT controls following BCAS. Representative

merged immunofluorescence images illustrate a reduction in M1 (CD86 positive) and

M2 (CD206 positive) microglial activation due to decreased expression levels of

CD86 and CD206 respectively, in the cerebral cortex (c) and hippocampus (f) in

AIM2 KO mice compared to WT controls following BCAS. (c & f), representative

merged immunofluorescence images of DAPI, Complement C3 (C3) and S100A10

within astrocytes (GFAP positive) in the cerebral cortex and hippocampus of WT

controls following BCAS. Other than colocalization of S100A10 (A2) within astrocytes

in the cerebral cortex (c), no substantial co-localization of C3 (A1) was observed in

the cerebral cortex (c) and hippocampus (f) of following BCAS. Representative

merged immunofluorescence images illustrate a reduction in A2 (S100A10 positive)

astrocytes activation due to decreased expression levels of S100A10 in the cerebral
cortex (c) in AIM2 KO mice compared to WT controls following BCAS. (g & h, j & k),
representative immunoblots and quantification illustrating resistance to myelin loss
due to increased expression of MBP in the cerebral cortex (g) and hippocampus (j) in
AIM2 KO mice compared to WT controls following BCAS. β-actin was used as a
loading control. Data are represented as mean ± S.E.M. n=5-7 mice in each
experimental group. *P<0.05 compared with WT Sham; +P<0.05 compared with WT
BCAS; ++P<0.01 compared with WT BCAS (i & l), representative
immunofluorescence images illustrating resistance to myelin loss due to increased
MBP immunoreactivity in the cerebral cortex (i), and hippocampal CA1, CA2 and
CA3 regions (l) in AIM2 KO mice compared to WT controls following BCAS.
Magnification x 100. Scale bar, 20µm. Images were taken under identical exposures
and conditions. Abbreviations: BCAS, bilateral common carotid artery stenosis; CD86,
cluster of differentiation 86; CD206, cluster of differentiation 206; C3, complement
component 3; GFAP, glial fibrillary acidic protein; Iba-1, ionized calcium binding
adaptor molecule-1; KO, knock out; S100A10, S100 calcium-binding protein A10.

1028

Figure 5: Effect of chronic cerebral hypoperfusion on cortical white matter
integrity and hippocampal neuronal density in AIM2 KO mice following BCAS.
(a-e), representative Luxol fast blue stained images and quantification illustrating
preserved white matter integrity due to decreased myelin rarefaction and white
matter lesion formation in the corpus callosum (paramedian), corpus callosum
(medial), caudoputamen and internal capsule in AIM2 KO mice compared with WT
controls following BCAS. No significant difference in white matter integrity was
observed in the optic tract between AIM2 KO mice compared with WT controls
following 30 days of BCAS. The severity of white matter disruption was graded
accordingly: Grade 0 = no disruption; Grade 1 = disarrangement of nerve fibres;
Grade 2 = formation of marked vacuoles; Grade 3 = disappearance of myelinated
fibres. (f-h), representative crystal violet images and quantification illustrating
increased Nissl positively stained neurons in hippocampal CA1, CA2 and CA3
regions in AIM2 KO mice compared to WT controls following BCAS. Data are
represented as mean ± S.E.M. n=5-7 mice in each experimental group. *P<0.05
compared with WT Sham; **P<0.01 compared with WT Sham; ***P<0.001 compared
with WT Sham. +P<0.05 compared with WT BCAS; ++P<0.01 compared with WT

BCAS; [+++]P<0.001 compared with WT BCAS. Magnification x 60. Scale bar, 20 µm.

Images were taken under identical exposures and conditions. Abbreviations: BCAS,

bilateral common carotid artery stenosis; WT, wild type; KO, knock out; WML, white

matter lesion.

1050

1051

Figure 6: Effect of chronic cerebral hypoperfusion on explorative locomotive

behavior and spatial memory in AIM2 KO mice following BCAS. (a-d),

quantifications illustrating an open field test conducted on all mice to examine the

locomotor ability and level of anxiety by measuring the total distance travelled (mm)

and average speed of movement (mm/s); total time (sec) and distance travelled (mm)

in the outer and inner zone of the open field. It was shown that WT BCAS mice did

not have a shorter distance travelled (mm) (a) and a lower average speed (mm/s) of

movement (b) when compared to WT Sham controls, indicating no impairment of

locomotor ability. Similarly, there are also no significant difference in the time spent

and distance travelled in the outer and inner zone among WT mice, showing no

BCAS-induced anxiety effect (c & d). However, it was observed that in both sham

and BCAS AIM2 KO mice more time was spent at the outer zone when compared to

WT controls (c), indicating that AIM2 KO mice generally exhibit higher levels of

anxiety independent of BCAS. (e-h), quantifications illustrating a Morris water maze

test conducted on all mice to examine spatial learning and memory by measuring the

escape latency (sec), time spent around the platform on probe day (sec), number of

visits in the target quadrant and time spent in target quadrant (sec). (e), in general, a

reduction in escape latency (sec) was observed in all mice across the training period.

Specifically, WT Sham, AIM2 KO Sham and AIM2 KO BCAS groups displayed a

trend with a shorter escape latency and steeper declining gradient, which reflected

an improved learning ability when compared to the WT BCAS group. (f-h), in general,

the AIM2 KO BCAS group displayed a trend with a longer time spent around the

platform on probe day and a significantly higher number of visits and time spent in

the target quadrant, which reflected retention of spatial memory when compared to

the WT BCAS group. Data are represented as mean ± S.E.M. n=9-11 mice in each

experimental group. *P<0.05 compared with WT Sham. #P<0.05 compared with WT

BCAS; ##P<0.01 compared with WT BCAS; ###P<0.001 compared with WT BCAS.

Abbreviations: BCAS, bilateral common carotid artery stenosis; WT, wild type; KO,

knock out.

1081