Large scale metagenome assembly reveals novel animal-associated microbial genomes,
biosynthetic gene clusters, and other genetic diversity

Nicholas D. Youngblut*,[1], Jacobo de la Cuesta-Zuluaga[1], Georg H. Reischer[2,3], Silke

Dauser[1], Nathalie Schuster[2], Chris Walzer[4], Gabrielle Stalder[4], Andreas H.

Farnleitner[2,3,5] , Ruth E. Ley[1]

[1]Department of Microbiome Science, Max Planck Institute for Developmental Biology, Max Planck Ring 5,

72076 Tübingen, Germany

[2]TU Wien, Institute of Chemical, Environmental and Bioscience Engineering, Research Group for

Environmental Microbiology and Molecular Diagnostics 166/5/3, Gumpendorfer Straße 1a, A-1060

Vienna, Austria

[3]ICC Interuniversity Cooperation Centre Water & Health, 1160 Vienna, Austria

[4]Research Institute of Wildlife Ecology, University of Veterinary Medicine, Vienna, Austria.

[5]Research Division Water Quality and Health, Karl Landsteiner University for Health Sciences, 3500

Krems an der Donau, Austria

22 * Corresponding author: Nicholas Youngblut (nyoungblut@tuebingen.mpg.de)

23
24 **Running title:** Metagenome assembly reveals vertebrate microbiome diversity
25 **Key words:** animal microbiome, gut, metagenome assembly, novel diversity
26
27

Abstract

Large-scale metagenome assemblies of human microbiomes have produced a
vast catalogue of previously unseen microbial genomes; however, comparatively few
microbial genomes derive from other vertebrates. Here, we generated 5596
metagenome-assembled genomes from the gut metagenomes of 180 predominantly
wild animal species representing 5 classes, in addition to 14 existing animal gut
metagenome datasets. The MAGs comprised 1522 species-level genome bins (SGBs);
most of which were novel at the species, genus, or family levels, and the majority were
enriched in host versus environment metagenomes. Many traits distinguished SGBs
enriched in host or environmental biomes, including the number of antimicrobial
resistance genes. We identified 1986 diverse biosynthetic gene clusters; only 23
clustered with any MIBiG database references. Gene-based assembly revealed
tremendous gene diversity, much of it host- or environment-specific. Our MAG and gene
datasets greatly expand the microbial genome repertoire and provide a broad view of
microbial adaptations to the vertebrate gut.

Importance

Microbiome studies on a select few mammalian species (*e.g.,* humans, mice, and
cattle) have revealed a great deal of novel genomic diversity in the gut microbiome.
However, little is known of the microbial diversity in the gut of other vertebrates. We
studied the gut microbiome of a large set of mostly wild animal species consisting of
mammals, birds, reptiles, amphibians, and fish. Unfortunately, we found that existing
reference databases commonly used for metagenomic analyses failed to capture the
microbiome diversity among vertebrates. To increase database representation, we
applied advanced metagenome assembly methods to our animal gut data and to many
public gut metagenome datasets that had not been used to obtain microbial genomes.
Our resulting genome and gene cluster collections comprised a great deal of novel
taxonomic and genomic diversity, which we extensively characterized. Our findings
substantially expand what is known of microbial genomic diversity in the vertebrate gut.

Introduction

The vertebrate gut microbiome comprises a vast amount of genetic diversity, yet
even for the most well-studied species such as humans, the number of microbial
species lacking a reference genome was recently estimated to be 40-50%[1]. Uncovering
this "microbial dark matter" is essential to understanding the roles of individual
microbes, their intra- and inter-species diversity within and across host populations, and
how each microbe interacts with each other and the host to mediate host physiology in
a myriad number of ways[2]. On a more applied level, characterizing novel gut microbial
diversity aids in bioprospecting of novel bioactive natural products, catalytic and

carbohydrate-binding enzymes, probiotics, etc., along with aiding in the discovery and
tracking of novel pathogens and antimicrobial resistance (AMR)[3].
Recent advances in culturomic approaches have generated thousands of novel
microbial genomes[4–6], but the throughput is currently far outpaced by metagenome
assembly approaches[7]. However, such large-scale metagenome assembly-based
approaches have not been as extensively applied to most non-human vertebrates. The
low amount of metagenome reads classified in some recent studies of the rhinoceros,
chicken, cod, and cow gut/rumen microbiome suggests that databases lack much of the
genomic diversity in less-studied vertebrates[8–11]. Indeed, the limited number of studies
incorporating metagenome assembly hint at the extensive amounts of as-of-yet novel
microbial diversity across the >66,000 vertebrate species on our planet.
Here, we developed an extensive metagenome assembly pipeline and applied it
to a multi-species dataset of microbiome diversity across vertebrate species comprising
5 classes: Mammalia, Aves, Reptilia, Amphibia, and Actinopterygii, with >80% of
samples obtained from wild individuals[12] combined with data from 14 published animal
gut metagenomes. Moreover, we also applied a recently developed gene-based
metagenome assembly pipeline to the entire dataset in order to obtain gene-level
diversity for rarer taxa that would otherwise be missed by genome-based assembly[13,14].
Our assembly approaches yielded a great deal of novel genetic diversity, which we
found to be largely enriched in animals versus the environment, and to some degree,
enriched in particular animal clades.

Methods

Sample collection

Sample collection was as described in Youngblut and colleagues[12]. Table S1A shows
the dates, locations, and additional metadata of all samples collected. All fecal samples
were collected in sterile sampling vials, transported to a laboratory and frozen within 8
hours. DNA extraction was performed with the PowerSoil DNA Isolation Kit (MoBio
Laboratories, Carlsbad, USA).

"multi-species" vertebrate gut metagenomes

Metagenome libraries were prepared as described by Karasov and colleagues[15].
Briefly, 1 ng of input gDNA was used for Nextera Tn5 tagmentation. A BluePippin was
used to restrict fragment sizes to 400-700 bp. Barcoded samples were pooled and
sequenced on an Illumina HiSeq3000 with 2x150 paired-end sequencing. Read quality
control (QC) is described in the Supplemental Methods.
Post-QC reads were taxonomically profiled with Kraken2 and Bracken v.2.2[16]
against the Struo-generated GTDB-r89 Kraken2 and Bracken databases[17]. HUMAnN2

v.0.11.2[18] was used to profile genes and pathways against the Struo-generated
HUMAnN2 database created from GTDB-r89.

Publicly available animal gut metagenomes

Published animal gut metagenome reads were downloaded from the Sequence
Read Archive (SRA) between May and August of 2019. Table S1B lists all included
studies. We selected studies with Illumina paired-end metagenomes from gut contents
or feces. MGnify samples were downloaded from the SRA in Oct 2019 (Table S1C).
Read quality control is described in the Supplemental Methods.

Metagenome assembly of genomes pipeline

Assemblies were performed on a per-sample basis, with reads subsampled via
seqtk v.1.3 to ≤20 million read pairs. The details of the assembly pipeline are described
in the Supplemental Methods.
A multi-locus phylogeny of all SGB representatives was inferred with PhyloPhlAn
v.0.41[19]. Secondary metabolites were identified with AntiSMASH v.5.1.1[20] and
DeepBGC v.0.1.18[21] and then characterized with BiGSCAPE[22]. Abricate was used to
identify antimicrobial resistance genes. We used Krakenuniq v.0.5.8[23] for estimating
abundance of MAGs in metagenome samples. Details can be found in the
Supplemental Methods.

Metagenome assembly of genes pipeline

Assemblies performed on a per-sample basis, with reads subsampled via seqtk
v.1.3 to ≤20 million pairs. We used PLASS v.2.c7e35[14] and Linclust (mmseqs
v.10.6d92c)[13] to assemble and cluster contigs. A full description is in the Supplemental
Methods. DESeq2[24] was used to estimate enrichment of MAGs and gene clusters in
metagenomes from host and environment biomes.

Data availability

The raw sequence data are available from the European Nucleotide Archive
under the study accession number PRJEB38078. Fasta files for the 5596 non-
redundant MAGs, 1522 SGBs,and gene clusters (50, 90, and 100% sequence identity
clustering) can be found at
http://ftp.tue.mpg.de/ebio/projects/animal_gut_metagenome_assembly/, along with
genbank files for all BGCs. Code used for processing the data can be found at
https://github.com/leylabmpi/animal_gut_metagenome_assembly.

Results

Animal gut metagenomes from a highly diverse collection of animals

We generated animal gut metagenomes from a breadth of vertebrate diversity spanning five classes: Mammalia, Aves, Reptilia, Amphibia, and Actinopterygii (the "multi-species" dataset; Figure 1). In total, 289 samples passed our read quality control, with 3.4e6 ± 5e6 s.d. paired-end reads per sample, resulting in a mean estimated coverage of 0.54 ± 0.14 s.d. (Figure S1). 180 animal species were represented, with up to 6 individuals per species (mean of 1.6). Most individuals were wild (81%).

Our read-quality control pipeline included stringent filtering of host reads; some samples contained high amounts of reads mapping to vertebrate genomes (up to 74%; 6 ± 17% s.d.; Figure 1). Gut content samples contained a significantly higher amount of host reads (13.5 ± 21.6% s.d.) versus feces metagenomes (4.7 ± 12.7% s.d.; Wilcox, *P* < 1.8e-7; Table S1A). We mapped all remaining reads to a custom comprehensive Kraken2 database built from the GTDB (Release 89). Still, many samples had a low percentage of mapped reads (43 ± 22 s.d.; Figure 1), with 29% of the samples having <20% mapped reads.

Figure 1. A large percentage of unmapped reads, even when using multiple comprehensive metagenome profiling databases. The dated host species phylogeny was obtained from http://timetree.org, with branches colored by host class. From inner to outer, the data mapped onto the tree is host diet, host captive/wild status, and the mean number of metagenome reads mapped to various host-specific, non-microbial, and microbial databases. Note that captive/wild status sometimes differs among individuals of the same species. The databases are i) a representative of each publicly available genome from the host species ("vertebrata host genome"), ii) all entries in the NCBI nt database with taxonomy IDs matching host species ("vertebrata host nt"), iii) as the previous, but all vertebrata sequences included, iv) the Kraken2 "plant" database, v) the Kraken2 "fungi" database, vi) the Kraken2 "protozoa" database, vii) a custom bacteria and archaea database created from the Genome Taxonomy Database, Release 89 ("GTDB-r89"). Reads were mapped iteratively to each database in the order shown in the legend (top to bottom), with only unmapped reads included in the next iteration. "unclassified" reads did not map to any database, which were used along with reads mapping to GTDB-r89 for downstream analyses ("microbial + unclassified").

Discovery of novel diversity by large-scale metagenome assembly

Our comprehensive metagenome assembly pipeline generated 4374 non-redundant MAGs. After quality control and de-replication (see Methods), 296 MAGs remained, with a mean percent completeness and contamination of 84 ± 14 and 1.5 ± 1.2 s.d., respectively (Figure S2; Supplemental Results).

We expanded our MAG dataset by applying our assembly pipeline to 14
publically available animal gut metagenome datasets in which no MAGs have been
generated by *de novo* metagenome assembly (Table S1B). Our metagenome selection
included 554 samples from members of Mammalia (dogs, cats, woodrats, pigs, whales,
rhinoceroses, pangolins, and non-human primates), Aves (geese, kakapos, and
chickens), and Actinopterygii (cod). We applied our assembly pipeline to each individual
dataset and generated a total of 5301 non-redundant MAGs (Figure S3; Supplemental
Results). The substantially higher number of MAGs from these 14 datasets versus our
single multi-species dataset is likely due to the larger number of samples and the high
sequencing depth for many of those samples (*e.g.,* we used 2 billion paired-end reads
from the dog gut microbiome dataset[25]).
We combined all MAGs and de-replicated at 99.9 and 95% average nucleotide
identity (ANI) to produce 5596 non-redundant MAGs and 1522 species-level genome
bins (SGBs), respectively (Tables S2A & S2B). Of the 5596 MAGs, 2773 (50%) had a
completeness of ≥90%. Of the 1522 SGBs, 1184 (78%) lacked a ≥95% ANI match to
the GTDB-r89, 266 (17%) lacked a genus-level match, and 6 lacked a family-level
match (Figures 2 & S4). Mapping taxonomic novelty onto a multi-locus phylogeny of all
1522 SGBs revealed that novel taxa were rather dispersed across the phylogeny
(Figure 2).
We also assessed the novelty of our SGBs relative to UHGG, a comprehensive
human gut genome database, and found that only 31% of our SGBs had ≥95% ANI to
any of the 4644 UHGG representatives, and this overlap only increased to 34% at a
90% ANI cutoff.
Our SGB collection mostly consisted of MAGs assembled from a few species in
the multi-study dataset, suggesting that the SGBs may not be representative of taxa
found in other, more distantly related vertebrates. To assess the level of representation,
we determined the prevalence of all SGBs across all multi-species metagenomes
(Figure S5). The host species with the highest number of observed SGBs did tend to be
those comprising the multi-study dataset (*e.g.,* pigs and primates); however, SGBs were
frequently observed across the host phylogeny (41 ± 61 s.d. SGBs per host), indicating
that the SGB collection was generally representative of the vertebrate gut microbiome.
Integrating the 1522 SGBs into our custom GTDB Kraken2 database significantly
increased the percent reads mapped (paired t-test, $P < 0.005$; Figure S6). The percent
increase varied from <1 to 62.8% (mean of 5.3 ± 6.7 s.d.) among animal species but did
not appear biased to just pigs, dogs, or other vertebrate species in the multi-study
datasets that we incorporated (Figure S7), which corresponds with our analysis of SGB
prevalence across vertebrate hosts (Figure S5).
207

1) Phylum
- Bacteroidota
- Firmicutes
- Firmicutes_C
- Actinobacteriota
- Firmicutes_A
- Fusobacteriota
- Proteobacteria
- Campylobacterota
- Verrucomicrobiota_A
- Desulfobacterota_A
- Firmicutes_B
- Verrucomicrobiota
- Synergistota
- Cyanobacteria
- Elusimicrobiota
- Eremiobacterota
- Armatimonadota
- Halobacterota
- Patescibacteria
- Fibrobacterota
- Chloroflexota
- Thermoplasmatota
- Spirochaetota
- Euryarchaeota
- Deferribacterota
- Myxococcota
- Planctomycetota

3) GTDB novelty
- Species match
- No species match
- No genus match
- No family match

2a) Class: Firmicutes
- Bacilli
- Negativicutes
- Clostridia
- Peptococcia
- Dehalobacteriia

2b) Class: Actinobacteriota
- Actinobacteria
- Coriobacteriia

2c) Class: Proteobacteria
- Gammaproteobacteria
- Alphaproteobacteria

4) DESeq2 Adj P < 0.01
- Host
- Environment

5) DESeq2 Adj P < 0.01
- Mammals
- Other animals

Tree scale: 0.1

Figure 2. A phylogeny of all 1522 SGBs. From innermost to outermost, the data mapped onto the phylogeny is: GTDB phylum-level taxonomic classifications, class-level taxonomies for Actinobacteriota, class-level taxonomies for Firmicutes, class-level taxonomies for Proteobacteria, taxonomic novelty, significant enrichment in host gut or environmental metagenomes, and significant enrichment in Mammals or other animals in our multi-species gut metagenome dataset. The phylogeny was inferred from multiple conserved loci via PhyloPhlAn. Orange dots on the phylogeny denote bootstrap values in the range of 0.7 to 1. The phylogeny is rooted on the last common ancestor of Archaea and Bacteria.

Enrichment of SGBs among animal clades

While the MAGs generated here derive from animal gut metagenomes, many of these taxa might be transient in the host and actually more prevalent in the environment. We tested this by generating a "host-environment" metagenome dataset comprising 283 samples from 30 BioProjects (17 environmental and 13 host-associated; Figure 3A). We found 932 of the 1522 SGBs (61%) to be significantly enriched in the host metagenomes (DESeq2, *adj. P* < 0.01; Figure 3B). The host-enriched SGBs (host-SGBs) were taxonomically diverse, comprising 22 phyla. In contrast, only 15 SGBs (1%) were environment-enriched (env-SGBs), which all belonged to either Actinobacteriota or Proteobacteria (Figure 3B). The only SGBs that were not significantly enriched in either group belonged to Actinobacteriota or Proteobacteria, along with two SGBs from the

Firmicutes A phylum. Mapping these data onto the SGB phylogeny revealed
phylogenetic clustering of the environment-enriched SGBs (Figure 2).
We investigated the traits of the host- and environment-enriched SGBs and
found many predicted phenotypes to be more prevalent in one or the other group
(Figure 3C; Table S2C). A total of 67 traits were predicted based on genomic content of
certain pfam domains[26]. Almost all env-SGBs were aerobes (93%), which may aid in
transmission between the environment and host biomes. In contrast, 87% of host-SGBs
were anaerobes. Furthermore, all env-SGBs could generate catalase and were bile
susceptible, while both phenotypes were sparse in host-SGBs (Figure 3C).
Carbohydrate metabolism also differed, with most host-SGBs predicted to consume
various tri-, di-, and mono-saccharides. In contrast, env-SGBs were enriched in
phenotypes associated with motility, nitrogen metabolism, and breakdown of
heterogeneous substrates (*e.g.,* cellobiose metabolism).
We also compared SGB enrichment in mammals versus non-mammals in our
"multi-species" metagenome dataset and found 361 SGBs (24%) to be significantly
enriched in mammals, while 22 (1%) were enriched in non-mammals (DESeq2, *adj. P* <
0.01; Figure S2C. Interestingly, 100% of SGBs in the two archaeal phyla (Halobacteria
and Euryarchaeota) were enriched in mammals. Also of note, most of the
Verrucomicrobiota SGBs (87%) were enriched in mammals. The only 2 phyla with >10%
of SGBs enriched in non-mammals were Proteobacteria (29%) and Campylobacteria
(25%).
In contrast to our assessment of phenotypes distinct to host- or env-SGBs, we
did not observe such a distinction of phenotypes among SGBs enriched in Mammalia or
non-mammal gut metagenomes (Figure S8). Certain phenotypes such as anaerobic
growth and lactose consumption were more prevalent among mammal species, but they
were not found to be significantly enriched relative to the null model.
Little is known about the distribution of antimicrobial resistance genes in the gut
microbiomes of most vertebrate species[27]; therefore, we investigated the distribution of
AMR genes among MAGs enriched in the environment versus host biomes. We found a
mean of 35 ± 26 s.d. AMR markers per genome (Figure S9A). The high average was
largely driven by Proteobacteria and Campylobacter genomes, which had a mean of
387 and 161 AMR markers per genome, respectively. The 5 most abundant markers
were ruvB, galE, tupC, fabL (ygaA), and arsT (Figure S9A). The more abundant
markers predominantly belonged to Firmicutes A, while Proteobacteria comprised larger
fractions of the less abundant markers. Environment-enriched taxa contained
substantially more AMR genes than host-enriched taxa, and the same was true for non-
Mammalia versus Mammalia-enriched taxa (Figures S9B & S9C).
264

265
Figure 3. A) Summary of the number of samples per biome for our multi-environment metagenome
dataset selected from the MGnify database. B) Number of SGBs found to be significantly enriched in
relative abundances in host versus (positive log$_2$ fold change; "l2fc") environmental metagenomes
(negative l2fc). Values shown are the number of MAGs significantly enriched (blue) in either biome or not
found to be significant (red). C) Host- and environment-enriched SGBs have distinct traits. Phenotypes
predicted based on MAG gene content (via Traitar [26]) are summarized for the SGBs significantly enriched
in host or environmental metagenomes (DESeq2 *Adj. P* < 0.01) or neither biome ("Neither" in the x-axis
facet). Note the difference in x-axis scale. Asterisks denote phenotypes significantly more prevalent in
SGBs of the particular biome versus a null model of 1000 permutations in which biome labels were
shuffled among SGBs. See Table S3A for all DESeq2 results.

MAGs reveal novel secondary metabolite diversity

We identified 1986 biosynthetic gene clusters (BGCs) among all 1522 SGBs. A
total of 28 different products were predicted, with the most abundant being non-
ribosomal peptide synthetases (NPRS; *n* = 473), sactipeptides (*n* = 307), and
arylpolyenes (*n* = 291; Figure S10). BGCs were identified in 2 archaeal and 18 bacterial
phyla. MAGs in the Firmicutes A phylum contained the most BGCs (*n* = 764; 38%),
while Bacteroidota and Actinobacteriota phyla possessed 381 (19%) and 272 (14%),
respectively (Figure S10). Still, Actinobacteriota SGBs did possess the highest average
number of BGCs per genome (16.3), followed by Eremiobacterota (9), Proteobacteria
(7.7), and Halobacterota (5.1).
Clustering all 1986 BGCs by BiGSCAPE generated 1764 families and 1305
clans, with clans being a second, coarser level of clustering[22]. Only 8 clans (comprising
23 BGCs) included any MIBiG database reference, suggesting a high degree of novelty

(Figure S11). Mapping the BGCs on a genome phylogeny of all species containing ≥3
BGCs (233 SGBs) revealed that the number of BGCs per genome was somewhat
phylogenetically clustered: the five genomes with the most BGCs belonged either to the
Actinobacteria or Gammaproteobacteria (Figure 4). Notably, these clades contained a
high number of host-SGBs. Of these 233 SGBs, the majority were taxonomically novel,
with 62% lacking a species-level match to GTDB-r89, and 18% lacking a genus-level
match (Figure 4). To determine which of the BGCs are most prevalent across animal
hosts, we quantified the prevalence of each BGC family across our multi-species
metagenome dataset and mapped it to the genome phylogeny (Figures 4 & S12). Of the
1543 BGC families found in the 233 SGBs, 83 were present in ≥25% of the animal
metagenomes, with ribosomally synthesized and post-translationally modified peptides
(RIPPs) being by far the most prevalent (up to 98% prevalence of individual BGC
families) and also found in species from a number of phyla.

303
Figure 4. Phylogeny of all SGBs (*n* = 233) with ≥3 BGCs identified by AntiSMASH. From innermost to
outermost, the data mapped onto the phylogeny is: 1) GTDB phylum-level taxonomic classifications, 2)
taxonomic novelty, 3) significant enrichment in host or environmental metagenomes, 4) the prevalence of
BGC families across the multi-species metagenome dataset, and 5) the number of BGCs identified in the
MAG. Prevalence is the maximum of any BGC family for that BGC type, and only BGC families with a

prevalence of ≥25% are shown. The phylogeny is a pruned version of that shown in Figure 2. Orange
dots on the phylogeny denote bootstrap values in the range of 0.7 to 1.

Large-scale gene-based metagenome assembly reveals novel diversity

We applied gene-based assembly methods to our combined metagenome

dataset[14], which generated a total of 150,718,125 non-redundant coding sequences

(average length of 179 amino acids). Clustering at 90 and 50% sequence identity

resulted in 140,225,322 and 6,391,861 clusters, respectively. Only 16.9 and 11.3% of

each respective cluster set mapped to the UniRef50 database, indicating that most

coding sequences were novel. The clusters comprised 88 bacterial and 11 archaeal

phyla; 80 of which were represented by <100 clusters, and 60 lacking a cultured

representative. Proteobacteria (mostly Gammaproteobacteria), Firmicutes, and

Bacteroidetes made up 92.2% of all clusters (Figure 5A). The proportion of clusters

belonging to each COG functional category was largely the same for the more abundant

bacterial phyla (Figure 5B), while more variation was seen among Euryarchaeota

(Figure 5C). The dominant 7 phyla showed substantial variation in the number of

clusters associated with various KEGG pathway categories (Figure S14). For instance,

a high proportion of Fusobacteria and Tenericutes clusters were associated with the

"nucleotide metabolism", "replication and repair", and "translation" categories. A total of

87,573 clusters were annotated as CAZy families, with GT51, GH13, GH18, GT02, and

GT04 representing 48% of all CAZy-annotated clusters (Figure 5E). Of the 12 phyla with

the most CAZy family clusters, there were substantial differences in proportions of

clusters falling into each family (Figure 5F).

332
Figure 5. A summary of the 50% sequence identity clusters generated from the gene-based metagenome
assembly of the combined dataset. A) The total number of gene clusters per phylum. For clarity, only
phyla with ≥100 clusters are shown. Labels on each bar list the number of clusters (and percent of the
total). B) The number of bacterial gene clusters per phylum and COG category. The "P" facet label refers
to "poorly characterized". C) The number of archaeal gene clusters per class (all belonging to
Euryarchaeota) and COG category. D) The number of viral gene clusters per COG category. E) The
number clusters annotated as each CAZy family. For clarity, only phyla with ≥100 clusters are shown.
Labels next to each bar denote the number of clusters. F) The number of clusters per CAZy family,
broken down by phylum. CAZy families and phyla are ordered by most to least number of clusters. For
clarity, only CAZy families and phyla with ≥100 total clusters are shown.

Biome enrichment of gene clusters from specific phyla

We mapped reads from our host-environment metagenome dataset to each
cluster and used DESeq2 to identify those significantly enriched (*adj. P* < 1e-5) in each
biome. Most strikingly, the same functional groups were enriched in both biomes,
regardless of the grouping (*i.e.,* COG functional category, KEGG pathway, or CAZy
family); however, the gene clusters belonged to different microbial phyla (Figure 6;
Supplemental Results). For instance, nearly all COG categories for gene clusters
belonging to Proteobacteria were environment-enriched, while the same COG

categories for clusters belonging to Firmicutes and Bacteroidetes were host-enriched. In
contrast, functional groups of certain phyla were enriched in one biome, while different
groups were enriched in the other, indicating within-phylum differences in functional
content and habitat distributions. For instance, Fusobacteria KEGG pathways were
predominantly host-enriched, but protein export, bacteria secretion system, and
aminoacyl-tRNA biosynthesis were environment-enriched, indicating that these 3
pathways were more predominant in environment-enriched members of Fusobacteria
(Figure 6B). Overall, these results suggest that both biomes select for these same
microbial functions, but the microbes involved often differ at coarse taxonomic scales.
We also assessed gene cluster enrichment in Mammalia versus non-Mammalia
and found fewer significantly enriched features, which may be due to the smaller
metagenome sample size or less pronounced partitioning of functional groups among
biomes (Figure S15; Supplemental Results). Still, we again observed that both biomes
enriched for the same microbial functions, but these belonged to different coarse
taxonomic groups. To assess whether abundance estimations were substantially
erroneous due to mis-mapping of metagenome reads to the gene clusters, we reran the
analysis with stricter DIAMOND mapping parameters but observed similar findings,
even though 48% fewer gene clusters were detected in any metagenome (Figure S16).
369

370
Figure 6. Enrichment of gene clusters grouped by phylum and A) COG category B) KEGG pathway or C)
CAZy family. Only groupings significantly enriched in abundance (DESeq2, *adj. P* < 1e-5) in either biome
are shown. Only gene clusters observed in at least 25% of the metagenomes were included. For clarity,
only KEGG pathways enriched in >7 phyla are shown, and only CAZy families enriched in >1 phylum are
shown. Note that the axes are flipped in B) relative to A) and C). See Tables S5A, S5B, and S5C for all
DESeq2 results.

 Functional metagenome profiling benefits from our gene catalogue

Lastly, we integrated our gene catalogue into a custom HUMAnN2 database built
from the GTDB-r89 and found that this combined database substantially increased
mappability of reads from our multi-species metagenome dataset (Figure S17;
Supplemental Results).

Discussion

Our MAG and gene cluster datasets, derived from 289 newly generated
metagenomes from 180 vertebrate species, along with 544 metagenomes from 14
publicly available animal gut metagenome datasets, substantially helps to expand the
breadth of cross-species gut metagenome comparisons (Figures 1 & 5). While
metagenomics is rapidly expanding in popularity[7], most analyses of metagenomic data
suffer from a reliance on incomplete reference databases[17], which we show to be
acutely problematic for the gut microbiomes of most vertebrates in our dataset (Figure
1). Grossly incomplete surveys of microbial diversity can lead to incorrect findings on
community assembly in the vertebrate gut[28]. Although our dataset has only partially
revealed this unknown diversity, it does substantially improve reference database
coverage at both the genome and gene levels and also provides an estimate of the
incompleteness of existing reference databases.
A major contribution of this study is the extensive MAG collection that we
generated by assembling the metagenomes of our multi-species dataset together with
14 other animal gut metagenome datasets from understudied host species. This
collection includes 1184, 266, and 6 genomes from novel species, genera, and families,
respectively (Figures 2 & S4). Moreover, we found little overlap (31%) between our
MAG collection and the extensive human microbiome genome catalogue comprising the
UHGG, which underscores its taxonomic novelty. We also showed substantial SGB
prevalence across all 5 vertebrate taxonomic classes (Figure S5), indicating that our
MAG collection is representative microbes found across the vertebrate taxonomy. Our
MAG collection, once combined with the GTDB[29], improved our ability to classify reads
in our multi-species metagenome dataset (Figure S6), which is critical for accurately
assessing gut microbiome diversity across vertebrates. Although MAGs have been
criticized for their incompleteness and potentially high prevalence of misassemblies[30],
we note that i) the overall completeness of our MAGs was rather high (90% median
completeness), ii) complete genomes are not required for accurate taxonomic
profiling[31], iii) the prevalence of misassemblies among MAGs is likely quite low when
using state-of-the-art assembly and binning approaches[32]. Still, researchers who may
utilize this set of MAGs should use caution when analyzing individual SNPs, plasmids,
genomic islands, or other potentially missing or misassembled genomic features[33].
We investigated the distribution of our MAGs across environment and host
biomes to elucidate the diversity of host-microbe symbiosis in the vertebrate gut.

Microbe-host symbiosis spans the continuum from free-living microbes that can simply
survive passage through the host gut, to obligate symbioses[34]. Therefore, MAGs
enriched in the environment versus the host would indicate a weak association, while
the opposite enrichment would suggest a more obligate symbiosis. We provide
evidence of host specificity for the majority of SGBs, while a few Proteobacteria and
Actinobacteria SGBs were environment-enriched. When just considering host-
associated metagenomes, these env-SGBs were generally enriched in non-mammals
(Figures 2, 3, & S8). This is consistent with the hypothesis that mixed-mode
transmission, especially between environmental sources and hosts, is more
commonplace in non-mammalian gut microbiome community assembly versus in
mammals[35].
Our trait-based analysis of SGBs supports the notion that host-enriched taxa are
adapted for a symbiotic lifestyle, while environment-enriched taxa are adapted for a
free-living or facultative symbiosis lifestyle (Figure 3). For instance, anaerobes
comprised almost all host-enriched SGBs, while environment-enriched SGBs were
aerobes or facultative anaerobes and generally motile, which could be highly beneficial
for transmission between the environment and gut biomes. Indeed, a recent directed
evolution experiment showed that selecting for inter-host migration can generate
bacterial strains with increased motility[36], and a trait-based study of the human infant
gut microbiome showed that later stages of succession are dominated by taxa adapted
to the anoxic gut[37].
By assessing SGB enrichment in Mammalia versus non-Mammlia metagenomes,
we elucidated the specificity of host-microbe symbioses in the gut across large
evolutionary distances. More SGBs were enriched in mammals versus non-mammals
(Figures 2 & S8), as we observed in our previous 16S rRNA assessment of these
vertebrate clades[12]. Few traits differed among SGBs enriched in either biome (Figure
S8), which may indicate that the traits assessed are similarly required for adaptation to
each host clade, even at this coarse evolutionary scale.
Vertebrates both play a critical role in the spread of antimicrobial resistance and
also have been sources of novel antibiotics and other natural products[27,38]. We
investigated BGC and AMR diversity in our MAG collection and observed a high
diversity of BGC products, but very few of the BGCs clustered into families with
experimentally characterized BGCs from the MIBiG database (Figures S9 & S10). This
contrasts with findings that only ~10% of BGCs in the human microbiome are
uncharacterized[39], which is likely due to the limited study of natural products in the gut
microbiome of non-human vertebrates[40,41]. We found NRPS-producing BGCs to be
prevalent among the Firmicutes SGBs, which is similar to a recent assessment of 501
genomes from rumen isolates in which thousands of BGCs were identified[42]. Still, RiPPs
were most prevalent across all vertebrate clades, which expands upon observations of
high prevalence of this BGC class in the gut microbiome of humans[39] (Figures 2 & 4).

By combining our AMR marker screen with our SGB biome enrichment analysis,
we were able to characterize how AMR is associated with varying degrees of symbiosis
(Figure S9), which is important for understanding AMR reservoirs[27,43]. Our findings
indicate that the AMR reservoir may be greater for free-living and facultatively symbiotic
taxa relative to microbes with stronger host associations (Figure S9). Indeed, some of
the most abundant AMR markers were associated with metal resistance (*e.g.,* ruvB,
tupC, and arsT), which may reflect a lifestyle in which the microbe is exposed to
environmental sources of metals[44,45].
While MAGs provide a powerful means of investigating species and strain-level
diversity within the vertebrate gut microbiome, the approach is limited to only relatively
abundant taxa with enough coverage to reach adequate assembly contiguity[46]. Our
gene-based assembly approach allowed us to greatly expand the known gene
catalogue of the vertebrate gut microbiome beyond just the abundant taxa, with a total
of >150 million non-redundant coding sequences generated, comprising 88 bacterial
and 11 archaeal phyla (Figure 5). In comparison, recent large-scale metagenome
assemblies of the gut microbiome from chickens, pigs, rats, and dogs have generated
7.7, 9.04, 7.7, 5.1, and 1.25 million non-redundant coding sequences,
respectively[8,25,47,48]. It is also illustrative to consider that a recent large-scale
metagenome assembly of cattle rumen metagenomes generated 69,678 non-redundant
genes involved in carbohydrate metabolism[9], while our gene collection comprised
substantially more CAZy-annotated gene clusters (*n* = 87,573), even after collapsing at
50% sequence identity. The increased mappability that we achieved across all 5
vertebrate clades when incorporating our gene catalogue in our functional metagenome
profiling pipeline demonstrates how our gene collection will likely aid future vertebrate
gut metagenome studies (Figure S17).
Our assessment of gene cluster abundances in metagenomes from environment
and host-associated biomes illuminates how microbiome functioning and taxonomy is
distributed across the free-living to obligate symbiont spectrum. Most notably, nearly all
prominent functional groups were enriched in both the environment and host-associated
biomes, but the specific gene clusters belonged to different taxonomic groups in each
biome (Figure 6). For instance, almost all abundant CAZy families were enriched in both
the environment and host biomes, but the environment was dominated by
Proteobacteria, while Firmicutes, Bacteroidetes, and Actinobacteria gene clusters
comprised most host-enriched CAZy families. This suggests the same coarse-level
functional groups are present across the free-living to obligate microbe-vertebrate
symbiosis lifestyles, but coarse-level taxonomy strongly differs across this spectrum.
This pattern largely remained true when we compared enrichment between the
Mammalia and non-mammals, suggesting that taxonomic differences prevail over
functional differences in regards to host specificity, at least over broad-scale vertebrate
evolutionary distances. While comparing function to taxonomy is challenging due to

differing levels of resolution, we do not believe that our findings are simply due to using
functional groupings that are coarser than taxonomy, given that i) we assessed multiple
functional grouping (COG, KEGG, and CAZy), which all showed similar patterns, even
though they differ in functional resolution, and ii) we assessed taxonomy at the very
coarse phylum level but still found stark taxonomic differences across biomes.
In conclusion, our large-scale metagenome assembly of both MAGs and coding
sequences from a diverse collection of vertebrates substantially expands the known
taxonomic and functional diversity of the vertebrate gut microbiome. We have
demonstrated that both taxonomic and functional metagenome profiling of the
vertebrate gut is improved by our MAG and gene catalogues, which will aid future
investigations of the vertebrate gut microbiome. Moreover, our collection can help guide
natural product discovery and bioprospecting of novel carbohydrate-active enzymes,
along with modeling AMR transmission among reservoirs. By characterizing the
distribution of MAGs and microbial genes across environment and host biomes, we
gained insight into how taxonomy and function differ along the free-living to obligate
symbiosis lifestyle spectrum. We must note that our metagenome assembly dataset is
biased toward certain animal clades, which likely impacts these findings. As
metagenome assembly becomes more commonplace for studying the vertebrate gut
microbiome, bias toward certain vertebrates (*e.g.,* humans) will decrease, and thus
allow for a more comprehensive reassessment of our findings.

Acknowledgements

We thank Nadine Ziemert for helpful discussions in regards to bioinformatic
approaches for secondary metabolite detection and analysis. This work was supported
by the Department of Microbiome Science at the Max Planck Institute for
Developmental Biology. This study was supported by the Austrian Science Fund (FWF)
research projects P23900 granted to Andreas H. Farnleitner and P22032 granted to
Georg H. Reischer. Further support came from the Science Call 2015 "Resource und
Lebensgrundlage Wasser" Project SC15-016 funded by the Niederösterreichische
Forschungs- und Bildungsgesellschaft (NFB).
We would like to thank the following collaborators for their huge efforts in sample
and data collection: Mario Baldi, School of Veterinary Medicine, Universidad Nacional
de Costa Rica; Wolfgang Vogl and Frank Radon, Konrad Lorenz Institute of Ethology
and Biological Station Illmitz; Endre Sós and Viktor Molnár, Budapest Zoo; Ulrike
Streicher, Conservation and Wildlife Management Consultant, Vietnam; Katharina Mahr,
Konrad Lorenz Institute of Ethology, University of Veterinary Medicine Vienna and
Flinders University Adelaide, South Australia; Peggy Rismiller, Pelican Lagoon
Research Centre, Australia; Rob Deaville, Institute of Zoology, Zoological Society of
London; Alex Lécu, Muséum National d'Histoire Naturelle and Paris Zoo; Danny
Govender and Emily Lane, South African National Parks, Sanparks; Fritz Reimoser,

Research Institute of Wildlife Ecology, University of Veterinary Medicine Vienna; Anna
Kübber-Heiss and Team, Pathology, Research Institute of Wildlife Ecology, University of
Veterinary Medicine Vienna; Nikolaus Eisank, Nationalpark Hohe Tauern, Kärnten;
Attila Hettyey and Yoshan Moodley, Konrad Lorenz Institute of Ethology, University of
Veterinary Medicine Vienna; Mansour El-Matbouli and Oskar Schachner, Clinical Unit of
Fish Medicine, University of Veterinary Medicine; Barbara Richter, Institute of Pathology
and Forensic Veterinary Medicine, University of Veterinary Medicine Vienna; Hanna
Vielgrader and Zoovet Team, Schönbrunn Zoo; Reinhard Pichler, Herberstein Zoo. We
explicitly thank the Freek Venter of South African National Parks and the National
Zoological Gardens of South Africa for granting access to their Parks for sample
collection.

Author Contributions

547   G.H.R., R.E.L., and A.H.F. created the study concept. G.H.R., N.S., C.W., and
548 G.S. performed the sample collection and metadata compilation. G.H.R., N.S., and S.D.
549 performed the laboratory work. N.D.Y. and J.C. performed the data analysis. N.D.Y.,
550 J.C., and R.E.L. wrote the manuscript.

Competing Interest Statement

552 No conflicts of interest declared.

References

1.   Nayfach, S., Rodriguez-Mueller, B., Garud, N. & Pollard, K. S. An integrated
metagenomics pipeline for strain profiling reveals novel patterns of bacterial
transmission and biogeography. *Genome Res.* **26**, 1612–1625 (2016).
2.   Thomas, A. M. & Segata, N. Multiple levels of the unknown in microbiome
research. *BMC Biol.* **17**, 48 (2019).
3.   Wang, W.-L. *et al.* Application of metagenomics in the human gut microbiome.
World J. Gastroenterol. **21**, 803–814 (2015).
4.   Zou, Y. *et al.* 1,520 reference genomes from cultivated human gut bacteria enable
functional microbiome analyses. *Nat. Biotechnol.* **37**, 179–185 (2019).
5.   Forster, S. C. *et al.* A human gut bacterial genome and culture collection for
improved metagenomic analyses. *Nat. Biotechnol.* **37**, 186–192 (2019).
6.   Mukherjee, S. *et al.* 1,003 reference genomes of bacterial and archaeal isolates
expand coverage of the tree of life. *Nat. Biotechnol.* **35**, 676–683 (2017).
7.   Almeida, A. *et al.* A unified sequence catalogue of over 280,000 genomes obtained
from the human gut microbiome. *bioRxiv* 762682 (2019) doi:10.1101/762682.
8.   Huang, P. *et al.* The chicken gut metagenome and the modulatory effects of plant-
derived benzylisoquinoline alkaloids. *Microbiome* **6**, 211 (2018).
9.   Stewart, R. D. *et al.* Assembly of 913 microbial genomes from metagenomic
sequencing of the cow rumen. *Nat. Commun.* **9**, 870 (2018).

10. Riiser, E. S. *et al.* Switching on the light: using metagenomic shotgun sequencing
to characterize the intestinal microbiome of Atlantic cod. *Environ. Microbiol.* **21**,
2576–2594 (2019).
11. Gibson, K. M. *et al.* Gut microbiome differences between wild and captive black
rhinoceros - implications for rhino health. *Sci. Rep.* **9**, 7570 (2019).
12. Youngblut, N. D. *et al.* Host diet and evolutionary history explain different aspects
of gut microbiome diversity among vertebrate clades. *Nat. Commun.* **10**, 2200
(2019).
13. Steinegger, M. & Söding, J. Clustering huge protein sequence sets in linear time.
Nat. Commun. **9**, 2542 (2018).
14. Steinegger, M., Mirdita, M. & Söding, J. Protein-level assembly increases protein
sequence recovery from metagenomic samples manyfold. *Nat. Methods* **16**, 603–
606 (2019).
15. Karasov, T. L. *et al.* Arabidopsis thaliana and Pseudomonas Pathogens Exhibit
Stable Associations over Evolutionary Timescales. *Cell Host Microbe* **24**, 168–
179.e4 (2018).
16. Lu, J., Breitwieser, F. P., Thielen, P. & Salzberg, S. L. Bracken: estimating species
abundance in metagenomics data. *PeerJ Comput. Sci.* **3**, e104 (2017).
17. de la Cuesta-Zuluaga, J., Ley, R. E. & Youngblut, N. D. Struo: a pipeline for
building custom databases for common metagenome profilers. *Bioinformatics*
(2019) doi:10.1093/bioinformatics/btz899.
18. Franzosa, E. A. *et al.* Species-level functional profiling of metagenomes and
metatranscriptomes. *Nat. Methods* **15**, 962–968 (2018).
19. Segata, N., Börnigen, D., Morgan, X. C. & Huttenhower, C. PhyloPhlAn is a new
method for improved phylogenetic and taxonomic placement of microbes. *Nat.
Commun.* **4**, 2304 (2013).
20. Blin, K. *et al.* antiSMASH 5.0: updates to the secondary metabolite genome mining
pipeline. *Nucleic Acids Res.* **47**, W81–W87 (2019).
21. Hannigan, G. D. *et al.* A deep learning genome-mining strategy for biosynthetic
gene cluster prediction. *Nucleic Acids Res.* **47**, e110 (2019).
22. Navarro-Muñoz, J. C. *et al.* A computational framework to explore large-scale
biosynthetic diversity. *Nat. Chem. Biol.* **16**, 60–68 (2020).
23. Breitwieser, F. P., Baker, D. N. & Salzberg, S. L. KrakenUniq: confident and fast
metagenomics classification using unique k-mer counts. *Genome Biol.* **19**, 198
(2018).
24. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
25. Coelho, L. P. *et al.* Similarity of the dog and human gut microbiomes in gene
content and response to diet. *Microbiome* **6**, 72 (2018).
26. Weimann, A. *et al.* From Genomes to Phenotypes: Traitar, the Microbial Trait
Analyzer. *mSystems* **1**, (2016).
27. Allen, H. K. *et al.* Call of the wild: antibiotic resistance genes in natural
environments. *Nat. Rev. Microbiol.* **8**, 251–259 (2010).
28. Rodriguez-R, L. M., Gunturu, S., Tiedje, J. M., Cole, J. R. & Konstantinidis, K. T.
Nonpareil 3: Fast Estimation of Metagenomic Coverage and Sequence Diversity.
mSystems **3**, (2018).

29. Parks, D. H. *et al.* A standardized bacterial taxonomy based on genome phylogeny
substantially revises the tree of life. *Nat. Biotechnol.* **36**, 996–1004 (2018).
30. Chen, L.-X., Anantharaman, K., Shaiber, A., Eren, A. M. & Banfield, J. F. Accurate
and complete genomes from metagenomes. *Genome Res.* **30**, 315–333 (2020).
31. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database. *Bioinformatics* (2019)
doi:10.1093/bioinformatics/btz848.
32. Mineeva, O., Rojas-Carulla, M., Ley, R. E., Schölkopf, B. & Youngblut, N. D.
DeepMAsED: Evaluating the quality of metagenomic assemblies. *Bioinformatics*
(2020) doi:10.1093/bioinformatics/btaa124.
33. Maguire, F. *et al.* Metagenome-Assembled Genome Binning Methods with Short
Reads Disproportionately Fail for Plasmids and Genomic Islands.
2020.03.31.997171 (2020) doi:10.1101/2020.03.31.997171.
34. Sachs, J. L., Skophammer, R. G. & Regus, J. U. Evolutionary transitions in
bacterial symbiosis. *Proc. Natl. Acad. Sci. U. S. A.* **108 Suppl 2**, 10800–10807
(2011).
35. Ebert, D. The Epidemiology and Evolution of Symbionts with Mixed-Mode
Transmission. *Annu. Rev. Ecol. Evol. Syst.* **44**, 623–643 (2013).
36. Robinson, C. D. *et al.* Experimental bacterial adaptation to the zebrafish gut reveals
a primary role for immigration. *PLoS Biol.* **16**, e2006893 (2018).
37. Guittar, J., Shade, A. & Litchman, E. Trait-based community assembly and
succession of the infant gut microbiome. *Nat. Commun.* **10**, 512 (2019).
38. Adnani, N., Rajski, S. R. & Bugni, T. S. Symbiosis-inspired approaches to antibiotic
discovery. *Nat. Prod. Rep.* **34**, 784–814 (2017).
39. Donia, M. S. *et al.* A systematic analysis of biosynthetic gene clusters in the human
microbiome reveals a common family of antibiotics. *Cell* **158**, 1402–1414 (2014).
40. Donia, M. S. & Fischbach, M. A. Small molecules from the human microbiota.
Science **349**, 1254766 (2015).
41. Milshteyn, A., Colosimo, D. A. & Brady, S. F. Accessing Bioactive Natural Products
from the Human Microbiome. *Cell Host Microbe* **23**, 725–736 (2018).
42. Seshadri, R. *et al.* Cultivation and sequencing of rumen microbiome members from
the Hungate1000 Collection. *Nat. Biotechnol.* **36**, 359–367 (2018).
43. von Wintersdorff, C. J. H. *et al.* Dissemination of Antimicrobial Resistance in
Microbial Ecosystems through Horizontal Gene Transfer. *Front. Microbiol.* **7**, 173
(2016).
44. Pal, C., Bengtsson-Palme, J., Kristiansson, E. & Larsson, D. G. J. Co-occurrence of
resistance genes to antibiotics, biocides and metals reveals novel insights into their
co-selection potential. *BMC Genomics* **16**, 964 (2015).
45. Ben Fekih, I. *et al.* Distribution of Arsenic Resistance Genes in Prokaryotes. *Front.*
Microbiol. **9**, 2473 (2018).
46. Parks, D. H. *et al.* Recovery of nearly 8,000 metagenome-assembled genomes
substantially expands the tree of life. *Nat Microbiol* **2**, 1533–1542 (2017).
47. Xiao, L. *et al.* A reference gene catalogue of the pig gut microbiome. *Nat Microbiol*
1, 16161 (2016).
48. Pan, H. *et al.* A gene catalogue of the Sprague-Dawley rat gut metagenome.
Gigascience **7**, (2018).