Ancient viral genomes reveal introduction of HBV and B19V to Mexico during the transatlantic slave trade.

Axel A. Guzmán-Solís[1]*, Daniel Blanco-Melo[2]*‡, Viridiana Villa-Islas[1], Miriam J. Bravo-López[1], Marcela Sandoval-Velasco[3], Julie K. Wesp[4], Jorge A. Gómez-Valdés[5], María de la Luz Moreno-Cabrera[6], Alejandro Meraz-Moreno[6], Gabriela Solís-Pichardo[7], Peter Schaaf[8], Benjamin R. tenOever[2,9,10] & María C. Ávila-Arcos[1‡].

1.International Laboratory for Human Genome Research, Universidad Nacional Autónoma de México (México); 2.Department of Microbiology, Icahn School of Medicine at Mount Sinai (NY, USA); 3.Section for Evolutionary Genomics, The Globe Institute, Faculty of Health, University of Copenhagen (Denmark); 4.Department of Sociology and Anthropology, North Carolina State University (USA); 5.Escuela Nacional de Antropología e Historia (México); 6.Instituto Nacional de Antropología e Historia (México); 7.Laboratorio Universitario de Geoquímica Isotópica (LUGIS), Instituto de Geología, Universidad Nacional Autónoma de México (México); 8.LUGIS, Instituto de Geofísica, Universidad Nacional Autónoma de México (México); 9.Virus Engineering Center for Therapeutics and Research (VECToR), Icahn School of Medicine at Mount Sinai (NY, USA); 10.Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai (NY, USA). * These authors contributed equally to this work. ‡ Correspondence: daniel.blancomelo@mssm.edu (D.B-M.), mavila@liigh.unam.mx (M.C.A-A.).

ABSTRACT

After the European colonization of the Americas there was a dramatic population collapse of the Indigenous inhabitants caused in part by the introduction of new pathogens. Although there is much speculation on the etiology of the Colonial epidemics, direct evidence for the presence of specific viruses during the Colonial era is lacking. To uncover the diversity of viral pathogens during this period, we designed an enrichment assay targeting ancient DNA (aDNA) from viruses of clinical importance and applied it on DNA extracts from individuals found in a Colonial (16th c. – 18th c.) hospital and a Colonial chapel where records suggest victims of epidemics were buried during important outbreaks in Mexico City. This allowed us to reconstruct three ancient human parvovirus B19 genomes, and one ancient human hepatitis B virus genome from distinct individuals. The viral genomes are similar to African strains, consistent with the inferred morphological and genetic African ancestry of the hosts as well as with the isotopic analysis of the human remains, suggesting an origin on the African continent. This study provides direct molecular evidence of ancient viruses being transported to the Americas during the transatlantic slave trade and their subsequent introduction to New Spain. Altogether, our observations enrich the discussion about the etiology of infectious diseases during the Colonial period in Mexico.

INTRODUCTION

European colonization in the Americas resulted in a notable genetic exchange mainly between Native American populations, Europeans, and Africans[1–3]. Along

with human migrations, numerous new species were introduced to the Americas

including bacterial and viral pathogens, which played a major role in the dramatic

population collapse that afflicted the immunologically-naïve Indigenous

inhabitants[4,5]. Among these pathogens, viral diseases, such as smallpox, measles

and mumps have been proposed to be responsible for many of the devastating

epidemics during the Colonial period[4]. Remarkably, the pathogen(s) responsible

for the deadliest epidemics reported in New Spain (the Spanish viceroyalty that

corresponds to Mexico, Central America, and the current US southwest states)

remains unknown and is thought to have caused millions of deaths during the 16th

century[4]. Indigenous populations were drastically affected by these mysterious

epidemics, generically referred to as *Cocoliztli* ("pest" in Nahuatl)[6], followed by

Africans and to a lesser extent European people[4,6,7]. Symptoms of the 1576

Cocoliztli epidemic were described in autopsy reports of victims treated at the

"Hospital Real de San José de los Naturales" (HSJN)[6,8], the first hospital in Mexico

dedicated specifically to treat the Indigenous population[6,8] (Figure 1a-b). The

symptoms described included high fever, severe headache, neurological disorders,

internal and external bleeding, hepatitis and intense jaundice[4,6,7]. This has led

some scholars to postulate that the etiological agent of the *Cocoliztli* epidemic was

a hemorrhagic fever virus[4,9], although others have suggested that the symptoms

could be explained by bacterial infections[6,10].

The study of ancient viral genomes has revealed important insights into the

evolution of specific viral families[11–19], as well as their interaction with human

populations[20]. To explore the presence of viral pathogens in circulation during

epidemic periods in New Spain, we leveraged the vast historical and archeological

information available for the Colonial HSJN. These include the skeletal remains of

over 600 individuals recovered from mass burials associated with the hospital's

architectural remnants (Figure 1b). Many of these remains were retrieved from

burial contexts suggestive of an urgent and simultaneous disposal of the bodies, as

in the case of an epidemic[8,21]. Prior bioarcheological research has shown that the

remains of a number of individuals in the HSJN collection displayed dental

modifications and/or morphological indicators typical of African ancestry[21],

consistent with historical and archeological research that documents the presence

of a large number of both free and enslaved Africans and their descendants in

Colonial Mexico[3]. Indeed a recent paleogenomics study reported a Sub-Saharan

African origin of three individuals from this collection[22].

Here we describe the screening for viral pathogens that circulated in New Spain

during Colonial times, using ancient DNA (aDNA) techniques (Supplementary

Figure 1). For this work, we sampled skeletal human remains recovered from the

HSJN, and from a second site in Mexico City ("La Concepcion" chapel) belonging

to one of the first catholic conversion centers in New Spain[23], where records

suggest victims of epidemics were also buried (Figure 1a). We report the

reconstruction of ancient hepatitis B virus (HBV) and human parvovirus B19

(B19V) genomes recovered from these remains. Our findings provide a direct

molecular evidence of human viral pathogens of African origin being introduced to

New Spain during the transatlantic slave trade.

RESULTS

We sampled the skeletal remains from two archeological sites, a Colonial Hospital and a Colonial chapel in Mexico City (Figure 1a). For the HSJN (as a part of a broader ongoing study of African ancestry), 21 dental samples (premolar and molar teeth) were selected based on previous morphometric analyses and dental modifications that suggested an African ancestry[21,24–26]. The African presence in the Indigenous Hospital might reflect an urgent response to an epidemic outbreak, since hospitals treated patients regardless of the origin of the affected individuals during serious public health crises[21]. Dental samples of five additional individuals were selected (based on their conservation state) from "La Concepción" chapel (COY), which is located 10 km south of the HSJN in Coyoacán, a Pre-Hispanic Indigenous neighborhood that later became the first Spanish settlement in Mexico City after the fall of Tenochtitlan[23]. Following strict aDNA protocols, we processed these dental samples to isolate aDNA for next-generation sequencing (NGS) (Methods). Teeth roots (which are vascularized) can be assumed to be a good source of pathogen DNA[27], especially in the case of viruses that are widespread in the bloodstream during a systemic infection. Accordingly, a number of previous studies have successfully recovered ancient viral DNA from teeth roots[14–16,22].

Metagenomic analysis with MALT[10] on the NGS data using the Viral NCBI RefSeq database as a reference[28], revealed that sixteen samples contained traces of viral DNA, particularly of sequences related to *Hepadnaviridae*, *Herpesviridae*, *Parvoviridae* and *Poxviridae* (Figure 1c, Supplementary Figure 2, Methods). These viral hits revealed the potential to recover ancient viral genomes from these

samples. We selected thirteen samples for further screening based on the

availability of the NGS library and the quality of the detected viral hit (best hit when

using megaBLAST[29] to NCBI nr database) to a clinically important virus (HBV,

B19V, Papillomavirus, Smallpox). To isolate and enrich the viral DNA fraction in the

sequencing libraries, biotinylated single-stranded (ss) RNA probes designed to

capture sequences from diverse human viral pathogens were synthesized

(Supplementary Table 1). The selection of the viruses included in the capture

design considered the following criteria: 1) DNA viruses previously retrieved from

archeological human remains (i.e. Hepatitis B virus, Human Parvovirus B19,

Variola Virus), 2) representative viruses from families capable of integrating into

the human genome (i.e. *Herpesviridae*, *Papillomaviridae*, *Polyomaviridae*,

Circoviridae) or 3) RNA viruses with a DNA intermediate (i.e. *Retroviridae*).

Additionally, a virus-negative aDNA library, which showed no hits to any viral family

included in the capture assay (except for a frequent *Poxviridae*-like region

identified as an Alu repeat[30]), was captured and sequenced as a negative control

(HSJN177) to estimate the efficiency of our capture assay. Four post-capture

libraries had a ~100-300-fold increase of HBV-like hits or a ~50-100-fold increase

of B19V-like hits (Figure 1c, Supplementary Table 2) compared to their

corresponding pre-capture libraries (Methods). In contrast, the captured negative

control (HSJN177) presented a negligible enrichment of these viral hits (Figure 1c,

Supplementary Table 2).

We verified the authenticity of the viral sequences by querying the mapped reads

against the non-redundant (nr) NCBI database using megaBLAST[29]. We only

retained reads for which the top hit was to either B19V or HBV, respectively

(Supplementary Table 3). For each sample, the retained reads were mapped

simultaneously to different available genotype sequences for their respective virus,

and the genotype with the best sequence coverage was selected as the reference

for final mapping (Supplementary Methods). To confirm the ancient origin of these

viral reads, we evaluated the misincorporation damage patterns using the program

mapDamage 2.0[31], which revealed an accumulation of C to T mutations towards

their 5' terminal site with an almost symmetrical G to A pattern on the 3' end

(Figure 2a, Supplementary Figure 3), as expected for aDNA[32]. Three ancient B19V

genomes were reconstructed (Figure 2b, Supplementary Table 3) with sequence

coverages between 92.37% and 99.1%, and average depths of 2.98-15.36X along

their single stranded DNA (ssDNA) coding region, which excludes the double

stranded DNA (dsDNA) hairpin regions at each end of the genome[33]. These

inverse terminal repeats (ITRs) displayed considerably higher depth values

compared to the coding region (<218X) consistent with the better *post-mortem*

preservation of dsDNA compared to ssDNA[34] (Figure 2b). In addition, we

reconstructed one ancient HBV genome (Figure 2c, Supplementary Table 3) at

30.8X average depth and with a sequence coverage of 89.9%, including its ssDNA

region at a reduced depth (<10X). This genome presents a 6 nucleotide (nt)

insertion in the core gene, that is characteristic of the genotype A[35]. Further

phylogenetic analyses (Methods) revealed that the Colonial HBV genome clustered

with modern sequences corresponding to sub-genotype A4 (previously named

A6)[36] (Figure 3a, Supplementary Figure 4). The Genotype A (HBV/GtA) has a

broad diversity in Africa reflecting its long history in this continent[35,37], while the

sub-genotype A4 has been recovered uniquely from African individuals in

Belgium[38] and has never been found in the Americas. Regarding the three Colonial

B19V genomes from individuals HSJN240, COYC4 and HSJNC81, these were

phylogenetically closer to modern B19V sequences belonging to genotype 3

(Figure 3b, Supplementary Figure 5). This B19V genotype is divided into two sub-

genotypes: 3a that is mostly found in Africa, and 3b, which is proposed to have

spread outside Africa in the last decades[39]. The viral sequences from the

individuals HSJN240 and COYC4 are similar to sub-genotype 3b genomes

sampled from immigrants (Morocco, Egypt and Turkey) in Germany[40] (Figure 3b,

Supplementary Figure 5); while the sequence of the individual HSJNC81 is more

similar to a divergent sub-genotype 3a strain (Figure 3b, Supplementary Figure 5)

retrieved from a child with severe anemia from France[41].

We recalibrated the molecular clock of HBV (Supplementary Figure 4b) and B19V

(Supplementary Figure 5b) with a dated coalescent phylogeny analysis that

included our ancient Colonial viral genomes, as well as other published ancient

and modern HBV and B19V genomes from different genotypes in diverse

geographical regions. The median evolutionary rate for HBV was estimated to be

between 1.1×10^{-5} and 9.5×10^{-6} (mean of 7.1×10^{-6}) nucleotide substitutions per site

per year (s/s/y) with a divergence time from last common ancestor between 8.3

and 13.61 (mean of 10.68) thousand years ago (ka) under a relaxed molecular

clock model and a coalescent exponential population prior (Supplementary Figure

4b). For B19V, the estimated substitution rate is 1.3×10^{-5} and 9.9×10^{-6} (mean of

1.17×10^{-6}) s/s/y with a divergence time from last common ancestor between 9.45

and 11.83 (mean 10.6) ka under a strict molecular clock model and a coalescent

Bayesian skyline population prior (Supplementary Figure 5b). The recalibrated

estimations are similar to previous reports[15,16] that included only Eurasian viral

genomes considerably older (~0.2–24.0 ka) than our Colonial viral genomes.

The similarity of our ancient genomes to African strains is consistent with the

previously reported morphological indicators of African ancestry[21,24–26] of the

corresponding HSJN individuals, as well as to an ancient DNA analysis based on

three individuals from the HSJN collection[22]. We thus used the *de novo* generated

sequence data to determine the mitochondrial haplogroup of the hosts, as well as

their autosomal genetic ancestry using the 1000 Genomes Project[42] as a reference

panel (Figure 3c, Supplementary Table 4). The nuclear genetic ancestry analysis

showed that all three HSJN individuals, from which the reconstructed viral

genomes were isolated, fall within African genetic variation in a Principal

Component Analysis plot (Figure 3c), while their mitochondrial DNA belong to the L

haplogroup, which has high frequency in African populations (Supplementary Table

4). Additionally, we performed $^{87}Sr/^{86}Sr$ isotopic analysis on two of the HSJN

individuals using teeth enamel as well as phalange (HSJN240) or parietal bone

(HSJNC81) to provide insights on the places of birth (adult enamel) and where the

last years of life were spent (phalange/parietal). The $^{87}Sr/^{86}Sr$ ratios measured on

the enamel of the individual HSJNC81 (0.71098) and HSJN240 (0.71109) are

similar to average $^{87}Sr/^{86}Sr$ ratios found in soils and rocks from West Africa

(average of 0.71044, Supplementary Figure 6, Supplementary Tables 5 and 6). In

contrast, the ^{87}Sr/^{86}Sr ratios on the parietal and phalange bones from the HSJNC81

(0.70672) and HSJN240 (0.70755), show lower values similar to those observed in

the Trans Mexican Volcanic Belt where the Mexico City Valley is located (0.70420 -

0.70550, Supplementary Figure 6, Supplementary Tables 5 and 6). Moreover,

radiocarbon dating of HSJN240 (1442-1608 CE, years calibrated for 1σ) and

HSJN194 (1472-1625 CE, years calibrated for 1σ) (Supplementary Table 4,

Supplementary Figure 7) indicates that these individuals arrived during the first

decades of the Colonial period, when the number of enslaved individuals arriving

from Africa was particularly high[3]. Strikingly, Colonial individual COYC4, who was

also infected with an African B19V strain, clusters with present-day Latin-American

populations from the 1000 Genomes Project (Figure 3c), suggesting that following

introduction from Africa, the virus (B19V) spread and infected people of different

ancestries during Colonial times.

DISCUSSION

In this study we reconstructed one HBV and three B19V ancient genomes from

four different individuals using NGS metagenomics and in-solution targeted

enrichment methods (Figure 2b, c, Supplementary Figure 1). Several lines of

evidence support that these ancient viral genomes are authentic and not an

environmental contamination nor a capture artifact. First, our negative control did

not show a considerable enrichment for B19V or HBV hits (Figure 1c). For those

samples that showed an enrichment in viral sequences after capture, the reads

covered the reference genomes almost in their entirety and displayed deamination

patterns at the terminal ends of the reads as expected for aDNA (Figure 2a).

Moreover, it is important to notice that B19V and HBV are blood-borne human

pathogens that are not present in soil or the environment, and that DNA from these

viruses had never been extracted before in the aDNA facilities used in this study.

We also described an unusual coverage pattern on the B19V genome, where the

dsDNA hairpins at its terminal sites are highly covered reflecting a better stability of

these regions over time (Figure 2b). Similarly, the partially circular dsDNA genome

from HBV was poorly covered at the ssDNA region (Figure 2c), as in at least three

previous ancient HBV genomes[14] (Supplementary Discussion 1). The variable

coverage in both viruses argues against an integration event of these viruses, that

would result as an uniform dsDNA coverage; further analysis are needed to

elucidate if the aDNA retrieved in this and other studies is coming from circulating

virions or from cell-free DNA intermediates[43] produced after viral replication in bone

marrow or liver, for B19V and HBV, respectively[44,45].

The ancient B19V genomes were assigned to genotype 3. This genotype is

considered endemic and the most prevalent in West Africa (Ghana: 100%, 11/11;

Burkina Faso: 100%, 5/5)[39,46,47] and a potential African origin has been

suggested[46]. This genotype has been also sporadically found outside Africa[46,47] in

countries historically tied to this continent, like Brazil (50%, 6/12)[48,49], India (15.4%,

2/13)[50], France (11.4%, 9/79)[41,51], and USA (0.85%, 1/117)[47] as well as in

immigrants from Morocco, Egypt, and Turkey in Germany (6.7%, 4/59)[40]. Two

other genotypes, 1 and 2 exist for this virus. Genotype 1 is the most common and

is found worldwide, while the almost extinct genotype 2 is mainly found in elderly

people from Northern Europe[52]. Ancient genomes from genotypes 1 and 2 have

been recovered from Eurasian samples, including a genotype 2 B19V genome

from a 10th century Viking burial in Greenland[16]. $^{87}Sr/^{86}Sr$ isotopes on individuals

from the same burial revealed they were immigrants from Iceland[16], suggesting an

introduction of the genotype 2 to contiguous North America during Viking

explorations of Greenland. While serological evidence indicates that B19V

currently circulates in Mexico, only the presence of genotype 1 has been formally

described using molecular analyses[53]. Taken together, our results are consistent

with an introduction of the genotype 3 to New Spain as a consequence of the

transatlantic slave trade imposed by the European colonization. This hypothesis is

supported by the $^{87}Sr/^{86}Sr$ isotopic analysis, which indicates that the individuals

from the HSJN with B19V (HSJN240, HSJNC81) were born in West Africa and

spent their last years of life in New Spain (Supplementary Figure 6). Furthermore,

the radiocarbon ages of individuals HSJN240, HSJN194 (Supplementary Figure 7)

support this notion as they correspond to the Early Colonial period, during which

the number of enslaved Africans arriving was higher compared to later periods[3].

Remarkably, a similar B19V belonging to the genotype 3 was recovered from an

Indigenous individual (COYC4) in an independent archeological site 10 Km south

of the HSJN (Figure 1a), supporting that viral transmissions between African

individuals and people of different ancestries occurred during the Colonial period in

Mexico City.

The genotype A from HBV is highly diverse in Africa, reflecting its long evolutionary

history, originated somewhere between Africa, Middle East and Central Asia[37]. The

introduction of the genotype A from Africa to America has been proposed based on

phylogenetic analysis of modern strains for Brazil[37,49] and Mexico[54], and more

precisely to the sub-genotype A1 for sequences from Martinique[55], Venezuela[56],

Haiti[57] and Colombia[58]. Recently, a similar introduction pattern was proposed for

the quasi genotype A3 based on an ancient HBV genome recovered from African

individuals sampled in Mexico[22]. The Colonial ancient HBV genome reconstructed

in our work is assigned to the genotype A4 (Figure 3a, Supplementary Figure 4),

which has never been reported in the Americas. A similar introduction pattern has

been proposed for other human-infecting viruses such as smallpox[7,59], based on

historical records; or Yellow fever virus[60], HTLMV-1[61], Hepatitis C virus (genotype

2)[62] and human herpes simplex virus[63] based on phylogenetic analysis of modern

strains from Afro-descendant or admixed human populations.

Although we cannot assert where exactly the African-born individuals in this study

contracted B19V or HBV (Africa, America, or the Middle Passage) nor if the cause

of their deaths can be attributed to such infections, the identification of ancient

B19V and HBV in contexts associated with Colonial epidemics in Mexico City is still

relevant in light of their paleopathological marks and the clinical information

available for the closest sequences in the phylogenetic analyses. The

reconstructed ancient B19V genome from individual HSJNC81 is closest to the V9

strain, which was isolated from an individual with severe anemia[41] (AJ249437)

(Figure 3b, Supplementary Figure 5). In fact, individual HSJNC81 displayed cribra

orbitalia in the eye sockets and porotic hyperostosis on the cranial vault

(Supplementary Figure 8); these morphological changes to the skeleton are

typically associated with anemias of varying different causes[64]. It is acknowledged

that B19V can result in a severe or even fatal anemia due to the low level of

hemoglobin when present simultaneously with other blood disorders, as

thalassemia, sickle-cell anemia, malaria and iron deficiency[45,65]. Therefore, since

B19V infects precursors of the erythroid lineage[45], it is possible that the

morphological changes found in HSJNC81 might be the result of a severe anemia

caused or enhanced by a B19V infection (Supplementary Discussion 2). Moreover,

the identification of ancient B19V in a Colonial context is noteworthy considering

there are historical records that describe the treatment of an outbreak of measles

at the HSJN in 1531[21], that numerous cases of measles were reported during this

period[4,8,59], and that several recent reports reveal that measles-like cases were

actually attributable to B19V[66,67] (Supplementary Discussion 2). Nevertheless, this

hypothesis requires future and more comprehensive studies aimed to characterize

the presence of measles and rubella viruses from ancient remains. A task that

imposes difficult technical challenges given that RNA is known to degrade rapidly,

in fact the most ancient viral RNA genomes (1912 CE) have been recovered only

from formalin-fixed tissue[17,18]. Additionally, historical records of the autopsies of

victims of the 1576 *Cocoliztli* epidemic treated at the HSJN, describe the

observation of enlarged hard liver and jaundice[4,6,7,9,68], which could be explained by

severe liver damage or epidemic hepatitis[4,6]. This is noteworthy given both viruses

HBV and B19V proliferate in the liver, have been isolated from liver tissue and are

associated with hepatitis and jaundice[44,45]. However, it is important to acknowledge

that both viruses have also been previously identified in aDNA datasets not

necessarily associated with disease or epidemic contexts[11,12,14,15], thus establishing

a direct link would require additional samples and a more comprehensive pathogen

screening to rule out the involvement of other pathogens. Finally, although our data

is not a strong evidence that the reported manifestations of liver damage in

Cocolitzi autopsies were directly caused by HBV or B19V, the identification of

these viruses in likely victims of epidemic outbreaks in the Colonial period opens

up new opportunities for investigating the presence of these viruses in similar

contexts. This type of research is particularly relevant when considering previous

hypotheses favoring the synergistic action of different pathogens in these

devastating Colonial epidemics[7] (Supplementary Discussion 3).

It is important to emphasize that our findings should be interpreted with careful

consideration of the historical and social context of the transatlantic slave trade.

This cruel episode in history involved the forced transportation of millions of

individuals to the Americas (ca. 250,000 to New Spain[3]) under inhumane,

unsanitary and overcrowded conditions that, with no doubt, favored the spread of

infectious diseases[59]. Therefore, the introduction of these and other pathogens

from Africa to the Americas should be attributed to the brutal and harsh conditions

of the Middle Passage that enslaved Africans were subjected to by traders and

colonizers, and not to the African peoples themselves. Moreover, the adverse life

conditions for enslaved Africans and Native Americans, especially during the first

decades after colonization, surely favored the spread of diseases and emergence

of epidemics[59]. Integrative and multidisciplinary approaches are thus needed to

understand this phenomenon at its full spectrum.

In summary, our study provides direct aDNA evidence of HBV and B19V

introduced to the Americas from Africa during the transatlantic slave trade. The

isolation and characterization of these ancient HBV and B19V genomes represent

an important contribution to the recently reported ancient viral genome reported in

the Americas[22] (only one before the present study). Our results expand our

knowledge on the viral agents that were in circulation during Colonial epidemics

like *Cocoliztli*, some of which resulted in the catastrophic collapse of the

immunologically-naïve Indigenous population. Although we cannot assign a direct

causality link between HBV and B19V and *Cocoliztli*, our findings confirm that

these potentially harmful viruses were indeed circulating in individuals found in

archeological contexts associated with this epidemic outbreak. Further analyses

from different sites and samples will help understand the possible role of these and

other pathogens in Colonial epidemics, as well as the full spectrum of pathogens

that were introduced to the Americas during European colonization.

METHODS

Sample selection and DNA extraction

Dental samples (premolars and molars) were obtained from twenty-one individuals

from the skeletal collection of the HSJN, selected based on their African-related

skeletal indicators[21,24–26]. Five additional samples were taken from "La Concepción"

chapel, based on their conservation state. Permits to carry out this sampling and

aDNA analyses were obtained by the Archeology Council of the National Institute

of Anthropology and History (INAH).

DNA extraction and NGS library construction

Bone samples were transported to a dedicated ancient DNA clean-room laboratory

at the International Laboratory for Human Genome Research (LIIGH-UNAM,

Querétaro, Mexico), were DNA extraction and NGS-libraries construction was

performed under the guidelines on contamination control for aDNA studies[69].

Previously reported aDNA extraction protocols were used for the HSJN[70] and

COY[71] samples. Double-stranded DNA (dsDNA) indexed sequencing libraries were

constructed from the DNA extract, as previously reported[72]. In order to detect

contaminants in reagents or by human manipulation, extraction and library

constructions protocols included negative controls (NGS blanks) that were

analyzed in parallel with the same methodology. The resulting NGS dsDNA

indexed libraries were quantified with a Bioanalyzer 2100 (Agilent) and pooled into

equimolar concentrations.

NGS sequencing

Pooled libraries were paired-end sequenced on an Illumina NextSeq550 at the

"Laboratorio Nacional de Genómica para la Biodiversidad" (LANGEBIO, Irapuato,

Mexico), with a Mid-output 2x75 format. The reads obtained (R1 and R2) were

merged (>11bp overlap) and trimmed with AdapterRemoval 1.5.4[73]. Overlapping

reads (>30 bp in length) were kept and mapped to the human genome (hg19)

using BWA 0.7.13[74], mapped reads were used for further human analysis (genetic

ancestry, and mitochondrial haplogroup determination), whereas unmapped reads

were used for metagenomic analysis and viral genome reconstruction.

Metagenomic analyses

The NCBI Viral RefSeq database was downloaded on February 2018; this included

7530 viral genomes. MALT 0.4.0[10] software was used to taxonomically classify the

reads using the viral genomes database. The viral database was formatted

automatically with malt-build once, and not human (unmapped) reads were aligned

with malt-run (85 minimal percent identity). The produced RMA files with the viral

abundances were normalized (default parameters) and compared to all the

samples from the same archeological site with MEGAN 6.8.0[75].

Capture-enrichment Assay

Twenty-nine viruses were included in the in-solution enrichment design, the

complete list of NCBI IDs is provided in Supplementary methods 5 and

Supplementary Table 1. It contained viral genomes previously recovered from

archeological remains like B19V, B19V-V9, and HBV (consensus genomes),

selected VARV genes, as well as clinically important viral families that are able to

integrate into the human genome, have dsDNA genomes, or dsDNA intermediates.

The resulting design comprised 19,147 ssRNA 80 nt probes targeting with a 20 nt

interspaced distance the whole or partial informative regions of eight viral families

of clinical relevance (*Poxviridae, Hepadnaviridae, Parvoviridae, Herpesviridae,*

Retroviridae, Papillomaviridae, Polyomaviridae, Circoviridae). To avoid a

simultaneous false-positive DNA enrichment, low complexity regions and human-

like (hg38) sequences were removed (*in sillico*). The customized kit was produced

by Arbor Biosciences (Ann Arbor, MI, USA). Capture-enrichment was performed on

the indexed libraries based on the manufacturer's protocol (version 4) to pull-down

aDNA with minimal modifications. Libraries were amplified with 18-20 cycles

(Phusion U Hot Start DNA Polymerase by Thermo Fischer Scientific), purified with

SPRISelect Magnetic Beads (Beckman Coulter) and quantified with a Bioanalyzer

2100 (Agilent). Amplified libraries were then pooled in different concentrations and

deep sequenced yielding $>1 \times 10^6$ non-human reads (Supplementary Table 4) in

order to saturate the target viral genome. Reads generated from each enriched

library were analyzed exactly as the shotgun (not-enriched) libraries. Normalized

abundances between shotgun and captured libraries were compared in MEGAN

6.8.0[75] to evaluate the efficiency and specificity of the enrichment assay.

Viral datasets

The full list of accession numbers of the following datasets is given in

Supplementary Methods 8.

HBV-Dataset-1 (HBV/DS1): comprises 35 HBV genomes from A-J human

genotypes, 2 well-covered ancient HBV genomes (LT992443, LT992459) and a

wholly monkey genome.

HBV-Dataset-2 (HBV/DS2): comprises 110 genomes based on a previous

phylogenetic analysis[15], that included genomes from A-J genotypes as well as not-

human primates HBV genomes (gibbon, gorilla, and chimpanzee), 16 ancient

HBVgenomes[11,12,14,15] and one ancient HBV genome from this study (HSJN194).

B19V-Dataset-1 (B19V/DS1): comprises 13 B19V genomes from human

genotypes 1-3 as well as a bovine parvovirus.

B19V-Dataset-2 (B19V/DS2): comprises 112 genomes from 1 to 3 B19V

genotypes based on previous phylogenetic analysis[16], that included the 10 best-

covered ancient genomes from genotype 1 and 2[16] as well as 3 ancient B19V from

this study. Since many of the reported genomes are not complete, only the whole

coding region (CDS) was used.

Genome Reconstruction and authenticity

HBV: Non-human reads were simultaneously mapped to HBV/DS1 with BWA (aln

algorithm) with seedling disabled[76]. The reference sequence with the most hits was

used to map uniquely to this reference and generate a BAM alignment without

duplicates, from which damage patterns were determined and damaged sites

rescaled using mapDamage 2.0[31], the rescaled alignment was used to produce a

consensus genome. All the HBV mapped reads were analyzed through

megaBLAST[29] using the whole NCBI nr database, in order to verify they were

assigned uniquely to HBV (carried out with Krona 2.7[77]).

B19V: The reconstruction of the B19V ancient genome was done as previously

reported from archeological skeletal remains[16], but increasing the stringency of

some parameters as described next. Non-human reads were mapped against

B19V/DS1 with BWA (aln algorithm) with seedling disabled[76], if more than 50% of

the genome was covered, the sample was considered positive to B19V. Reads

from the B19V-positive libraries were aligned with blastn (-evalue 0.001) to

B19V/DS1 in order to recover all the parvovirus-like reads. To avoid local

alignments, only hits covering >85% of the read were kept and joined to the B19V

mapped reads (from BWA), duplicates were removed. The resulting reads were

analyzed with megaBLAST[29] using the whole NCBI nr database to verify the top hit

was to B19V (carried out with Krona 2.7[77]). This pipeline was applied for two

independent enrichments assays per sample and the filtered reads from the two

capture rounds were joined. The merged datasets per sample were mapped using

as a reference file the three known B19V genotypes with GeneiousPrime

2019.0.4[78] using median/fast sensibility and iterate up to 5 times. The genotype

with the longest covered sequence was selected as the reference for further

analysis. Deamination patterns for HBV and B19V were determined with

mapDamage 2.0[31] and damaged sites were rescaled in the same program to

produce a consensus whole genome using SAMtools 1.9[79].

Phylogenetic analyses

HBV/DS2 and B19V/DS2 were aligned independently in Aliview[80] (Muscle

algorithm), and evolutionary models were tested under an AICc and BIC in

jModelTest[81]. A neighbor joining tree with 1000 bootstraps was generated in

GeneiousPrime 2019.0.4[78] using a Hasegawa-Kishino-Yano (HKY) model for both

alignments. A maximum likelihood tree with 1000 bootstraps was produced in

RAxML 8.2.10[82] using as a model a Generalized Time-Reversible (GTR)+G and

GTR+G+I for B19V and HBV, respectively.

Since a temporal signal has been described for the coding region of B19V[16] and

the whole genome of HBV[14,15], a Bayesian tree was generated to estimate the

impact of the Colonial viral genomes on the divergence time from the most recent

common ancestor (MRCA). We used BEAST 1.8.4[83], a Bayesian skyline

population prior and a relaxed lognormal or strict molecular clock for B19V and a

Coalescent Exponential Population prior with relaxed lognormal molecular clock for

HBV as previously tested[15,16]. All parameters were mixed and converged into an

estimated sample size (ESS) >150 analyzed in Tracer 1.7[84]. The first 25% of trees

where discarded (burn in) and a Maximum Clade Credibility Tree was created with

TreeAnnotator[83]. The generated trees were visualized and edited in FigTree 1.4.3

with a midpoint root.

Human population genetic analyses

Human-mapped reads (BWA aln) obtained from the pre-capture sequence data of

viral-positive samples were used to infer the genetic ancestry of the hosts. A

Principal Components Analysis (PCA) was carried out using 16 populations from

the 1000 Genomes Project[42] reference panel including genotype data of 1,562,771

single nucleotide variants (SNVs) from 2,504 individuals (phase 3). Genomic

alignments of each ancient individual (HSJNC81, HSJN240, HSJN194 and

COYC4) were intersected with the positions of the SNVs present in the reference

panel genotype data. Pseudo haploid genotypes were called by randomly selecting

one allele at each intersected site and filtering by a base quality >30. Pseudo

haploid genotypes were also called for the complete reference panel. PCA was

performed on the merged ancient and modern dataset with smartpca (EIGENSOFT

package)[85,86] using the option *lsqproject* to project the ancient individuals into the

PC space defined by the modern individuals.

Mitochondrial haplogroup and sex determination

NGS reads were mapped to the human mitochondrial genome reference (rCRS) with BWA (aln algorithm, -l default), the alignment file was then used to generate a consensus mitochondrial genome with program Schmutzi[87] The assignment of the mitochondrial haplogroup was carried out with Haplogrep[88,89] using the consensus sequence as the input. Assignment of biological sex was inferred based on the fraction of reads mapped to the Y-chromosome (Ry) compared to those mapping to the Y and X-chromosome[90]. Ry<0.016 and Ry>0.075 were considered XX or XY genotype, respectively. The resulting sex was coherent with the inferred morphologically (Supplementary Methods).

ACKNOWLEDGMENTS

This work was funded by the Wellcome Trust Sanger grant number 208934/Z/17/Z, and by project IA201219 PAPIIT-DGAPA-UNAM. D.B-M is an Open Philanthropy Fellow of the Life Sciences Research Foundation (LSRF). We thank the INAH Archeology Council for the sample permissions for this study. We are grateful with Teodoro Hernández Treviño, Gerardo Arrieta García from the "Laboratorio Universitario de Geoquímica Isotópica" (LUGIS-UNAM) for their technical support in performing the ^{87}Sr/^{86}Sr analyses and to Luis Alberto Aguilar Bautista, Alejandro de León Cuevas, Carlos Sair Flores Bautista and Jair Garcia Sotelo from the "Laboratorio Nacional de Visualización Científica Avanzada" (LAVIS/UNAM) who stored our data and provided the computational resources to perform this study.

We thank Alejandra Castillo Carbajal and Carina Uribe Díaz for technical support

throughout the project.

DATA AVAILABILITY

Reconstructed genomes from this study are available in Genbank under accession

number MT108214, MT108215, MT108216, MT108217. Accession number of

sequences used in phylogenetic analysis are indicated in supplementary

information. NGS data is available upon reasonable request.

REFERENCES:

1.   Rotimi, C. N., Tekola-Ayele, F., Baker, J. L. & Shriner, D. The African Diaspora: History, Adaptation and Health. *Curr. Opin. Genet. Dev.* **41**, 77–84 (2016).

2.   Salas, A. *et al.* The African Diaspora: Mitochondrial DNA and the Atlantic Slave Trade. *Am. J. Hum. Genet.* **74**, 454–465 (2004).

3.   Aguirre-Beltrán, G. La presencia del negro en México. *Rev. CESLA* **7**, 351–367 (2005).

4.   Acuna-Soto, R., Stahle, D. W., Therrell, M. D., Griffin, R. D. & Cleaveland, M. K. When half of the population died: the epidemic of hemorrhagic fevers of 1576 in Mexico. *FEMS Microbiol. Lett.* **240**, 1–5 (2004).

5.   Lindo, J. *et al.* A time transect of exomes from a Native American population before and after European contact. *Nat. Commun.* **7**, 13175 (2016).

6.   Malvido, E. & Viesca, C. La epidemia de cocoliztli de 1576. in *Ensayos sobre la historia de las epidemias en México.* (eds. Florescano, E. & Malvido, E.) 27–32 (Instituto Mexicano del Seguro Social, 1982).

7.   Somolinos d'Árdois, G. Las epidemias en México durante el siglo XVI. in *Ensayos sobre la historia de las epidemias en México.* (eds. Florescano, E. & Malvido, E.) 138–143 (Instituto Mexicano del Seguro Social, 1982).

8.   Wesp, J. K. Caring for Bodies or Simply Saving Souls: The Emergence of Institutional Care in Spanish Colonial America. in *New Developments in the Bioarchaeology of Care: Further Case Studies and Expanded Theory* (eds. Tilley, L. & Schrenk, A. A.) 253–276 (Springer International Publishing, 2017).

9.   Marr, J. S. & Kiracofe, J. B. Was the huey cocoliztli a haemorrhagic fever? *Med. Hist.* **44**, 341–362 (2000).

10.   Vågene, Å. J. *et al.* Salmonella enterica genomes from victims of a major sixteenth-century epidemic in Mexico. *Nat. Ecol. Evol.* **2**, 520 (2018).

11.   Kahila Bar-Gal, G. *et al.* Tracing hepatitis B virus to the 16th century in a Korean mummy. *Hepatol. Baltim. Md* **56**, 1671–1680 (2012).

12.   Patterson Ross, Z. *et al.* The paradox of HBV evolution as revealed from a

16th century mummy. *PLoS Pathog.* **14**, e1006750 (2018).

13. Duggan, A. T. *et al.* 17th Century Variola Virus Reveals the Recent History of Smallpox. *Curr. Biol. CB* **26**, 3407–3412 (2016).

14. Krause-Kyora, B. *et al.* Neolithic and medieval virus genomes reveal complex evolution of hepatitis B. *eLife* **7**, (2018).

15. Mühlemann, B. *et al.* Ancient hepatitis B viruses from the Bronze Age to the Medieval period. *Nature* **557**, 418–423 (2018).

16. Mühlemann, B. *et al.* Ancient human parvovirus B19 in Eurasia reveals its long-term association with humans. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 7557–7562 (2018).

17. Xiao, Y.-L. *et al.* High-throughput RNA sequencing of a formalin-fixed, paraffin-embedded autopsy lung tissue sample from the 1918 influenza pandemic. *J. Pathol.* **229**, 535–545 (2013).

18. Düx, A. *et al. The history of measles: from a 1912 genome to an antique origin.* http://biorxiv.org/lookup/doi/10.1101/2019.12.29.889667 (2019) doi:10.1101/2019.12.29.889667.

19. Pajer, P. *et al.* Characterization of Two Historic Smallpox Specimens from a Czech Museum. *Viruses* **9**, (2017).

20. Spyrou, M. A., Bos, K. I., Herbig, A. & Krause, J. Ancient pathogen genomics as an emerging tool for infectious disease research. *Nat. Rev. Genet.* **20**, 323–340 (2019).

21. Meza, A. Presencia africana en el cementerio del Hospital Real de San Jóse de los Naturales. *Arqueol. Mex.* **119**, 40–44 (2013).

22. Barquera, R. *et al.* Origin and Health Status of First-Generation Africans from Early Colonial Mexico. *Curr. Biol.* S0960982220304826 (2020) doi:10.1016/j.cub.2020.04.002.

23. Moreno-Cabrera, M. de la L., Meraz-Moreno, A. & Cervantes-Rosado, J. Relleno aligerado con vasijas cerámicas en el templo de la Inmaculada Concepción, en Coyoacán. in *Boletín de monumentos históricos 35* vol. 35 121–134 (Instituto Nacional de Antropología e Historia, 2015).

24. Ruíz-Albarrán, P. *Estudio de variabilidad biológica en la colección*

esquelética Hospital Real de San José de los Naturales. Un acercamiento a tráves

de la técnica de morfometría geométrica. (Escuela Nacional de Antropología e

Historia., 2012).

25.   Hernández-Lopez, P. E. & Negrete, S. *¿Realmente Eran Indios? Afinidad*

biológica entre las personas atendidas en el Hospital Real San Jose de los

Naturales, siglos XVI - XVIII. (Escuela Nacional de Antropología e Historia., 2012).

26.   Karam-Tapia, C. E. *Estimación del Mestizaje Mediante la Morfología Dental*

en la Ciudad de México (Siglo XVI al XIX). (Escuela Nacional de Antropología e

Historia., 2012).

27.   Key, F. M., Posth, C., Krause, J., Herbig, A. & Bos, K. I. Mining

Metagenomic Data Sets for Ancient DNA: Recommended Protocols for

Authentication. *Trends Genet. TIG* **33**, 508–520 (2017).

28.   Pruitt, K. D., Tatusova, T. & Maglott, D. R. NCBI reference sequences

(RefSeq): a curated non-redundant sequence database of genomes, transcripts

and proteins. *Nucleic Acids Res.* **35**, D61-65 (2007).

29.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local

alignment search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

30.   Tithi, S. S., Aylward, F. O., Jensen, R. V. & Zhang, L. FastViromeExplorer: a

pipeline for virus and phage identification and abundance profiling in

metagenomics data. *PeerJ* **6**, e4227 (2018).

31.   Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F. & Orlando, L.

mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage

parameters. *Bioinforma. Oxf. Engl.* **29**, 1682–1684 (2013).

32.   Briggs, A. W. *et al.* Patterns of damage in genomic DNA sequences from a

Neandertal. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 14616–14621 (2007).

33.   Luo, Y. & Qiu, J. Human parvovirus B19: a mechanistic overview of infection

and DNA replication. *Future Virol.* **10**, 155–167 (2015).

34.   Lindahl, T. Instability and decay of the primary structure of DNA. *Nature*

362, 709–715 (1993).

35.   Kramvis, A. Genotypes and Genetic Variability of Hepatitis B Virus.

Intervirology **57**, 141–150 (2014).

36.   Pourkarim, M. R., Amini-Bavil-Olyaee, S., Kurbanov, F., Van Ranst, M. &
Tacke, F. Molecular identification of hepatitis B virus genotypes/subgenotypes:
revised classification hurdles and updated resolutions. *World J. Gastroenterol.* **20**,
7152–7168 (2014).

37.   Kostaki, E.-G. *et al.* Unravelling the history of hepatitis B virus genotypes A
and D infection using a full-genome phylogenetic and phylogeographic approach.
eLife **7**, (2018).

38.   Pourkarim, M. R., Lemey, P., Amini-Bavil-Olyaee, S., Maes, P. & Van Ranst,
M. Novel hepatitis B virus subgenotype A6 in African-Belgian patients. *J. Clin.*
Virol. Off. Publ. Pan Am. Soc. Clin. Virol. **47**, 93–96 (2010).

39.   Hübschen, J. M. *et al.* Phylogenetic analysis of human parvovirus b19
sequences from eleven different countries confirms the predominance of genotype
1 and suggests the spread of genotype 3b. *J. Clin. Microbiol.* **47**, 3735–3738
(2009).

40.   Schneider, B. *et al.* Simultaneous persistence of multiple genome variants of
human parvovirus B19. *J. Gen. Virol.* **89**, 164–176 (2008).

41.   Nguyen, Q. T. *et al.* Novel Human Erythrovirus Associated with Transient
Aplastic Anemia. *J. Clin. Microbiol.* **37**, 2483–2487 (1999).

42.   1000 Genomes Project Consortium *et al.* A global reference for human
genetic variation. *Nature* **526**, 68–74 (2015).

43.   Cheng, A. P. *et al.* A cell-free DNA metagenomic sequencing assay that
integrates the host injury response to infection. *Proc. Natl. Acad. Sci.* **116**, 18738–
18744 (2019).

44.   Yuen, M.-F. *et al.* Hepatitis B virus infection. *Nat. Rev. Dis. Primer* **4**, 18035
(2018).

45.   Broliden, K., Tolfvenstam, T. & Norbeck, O. Clinical aspects of parvovirus
B19 infection. *J. Intern. Med.* **260**, 285–304 (2006).

46.   Candotti, D., Etiz, N., Parsyan, A. & Allain, J.-P. Identification and
Characterization of Persistent Human Erythrovirus Infection in Blood Donor
Samples. *J. Virol.* **78**, 12169–12178 (2004).

47.   Rinckel, L. A., Buno, B. R., Gierman, T. M. & Lee, D. C. Discovery and

analysis of a novel parvovirus B19 Genotype 3 isolate in the United States.

Transfusion (Paris) **49**, 1488–1492 (2009).

48.   Sanabani, S., Neto, W. K., Pereira, J. & Sabino, E. C. Sequence Variability

of Human Erythroviruses Present in Bone Marrow of Brazilian Patients with

Various Parvovirus B19-Related Hematological Symptoms. *J. Clin. Microbiol.* **44**,

604–606 (2006).

49.   Freitas, R. B. *et al.* Molecular characterization of human erythrovirus B19

strains obtained from patients with several clinical presentations in the Amazon

region of Brazil. *J. Clin. Virol. Off. Publ. Pan Am. Soc. Clin. Virol.* **43**, 60–65 (2008).

50.   Jain, P. *et al.* Prevalence and genotypic characterization of human

parvovirus B19 in children with hemato-oncological disorders in North India. *J.*

Med. Virol. **87**, 303–309 (2015).

51.   Servant, A. *et al.* Genetic Diversity within Human Erythroviruses:

Identification of Three Genotypes. *J. Virol.* **76**, 9124–9134 (2002).

52.   Pyöriä, L. *et al.* Extinct type of human parvovirus B19 persists in tonsillar B

cells. *Nat. Commun.* **8**, 14930 (2017).

53.   Valencia Pacheco, G. *et al.* Serological and molecular analysis of parvovirus

B19 infection in Mayan women with systemic lupus erythematosus in Mexico.

Colomb. Médica **v48**, 105–112 (2017).

54.   Roman, S. *et al.* Occult hepatitis B in the genotype H-infected Nahuas and

Huichol native Mexican population. *J. Med. Virol.* **82**, 1527–1536 (2010).

55.   Brichler, S. *et al.* African, Amerindian and European hepatitis B virus strains

circulate on the Caribbean Island of Martinique. *J. Gen. Virol.* **94**, 2318–2329

(2013).

56.   Quintero, A. *et al.* Molecular epidemiology of hepatitis B virus in Afro-

Venezuelan populations. *Arch. Virol.* **147**, 1829–1836 (2002).

57.   Andernach, I. E., Nolte, C., Pape, J. W. & Muller, C. P. Slave trade and

hepatitis B virus genotypes and subgenotypes in Haiti and Africa. *Emerg. Infect.*

Dis. **15**, 1222–1228 (2009).

58.   Alvarado-Mora, M. V. *et al.* Phylogenetic analysis of complete genome

sequences of hepatitis B virus from an Afro-Colombian community: presence of

HBV F3/A1 recombinant strain. *Virol. J.* **9**, 244 (2012).

59.   Mandujano-Sánchez, A., Solache, L. C. & Mandujano, M. A. Historia de las Epidemias en el México Antiguo: Algunos Aspectos Biológicos y Sociales. in *Ensayos sobre la historia de las epidemias en México.* (eds. Florescano, E. & Malvido, E.) 9–21 (Instituto Mexicano del Seguro Social, 1982).

60.   Bryant, J. E., Holmes, E. C. & Barrett, A. D. T. Out of Africa: A Molecular Perspective on the Introduction of Yellow Fever Virus into the Americas. *PLoS Pathog.* **3**, (2007).

61.   Gadelha, S. R. *et al.* The origin of HTLV-1 in the South Bahia by phylogenetic, mitochondrial DNA and β-globin analysis. *Retrovirology* **11**, P49 (2014).

62.   Markov, P. V. *et al.* Phylogeography and molecular epidemiology of hepatitis C virus genotype 2 in Africa. *J. Gen. Virol.* **90**, 2086–2096 (2009).

63.   Forni, D. *et al.* Recent out-of-Africa migration of human herpes simplex viruses. *Mol. Biol. Evol.* msaa001 (2020) doi:10.1093/molbev/msaa001.

64.   Angel, J. L. Porotic hyperostosis, anemias, malarias, and marshes in the prehistoric Eastern Mediterranean. *Science* **153**, 760–763 (1966).

65.   Heegaard, E. D. & Brown, K. E. Human Parvovirus B19. *Clin. Microbiol. Rev.* **15**, 485–505 (2002).

66.   Rezaei, F. *et al.* Prevalence and genotypic characterization of Human Parvovirus B19 in children with measles- and rubella-like illness in Iran. *J. Med. Virol.* **88**, 947–953 (2016).

67.   De Los Ángeles Ribas, M. *et al.* Identification of human parvovirus B19 among measles and rubella suspected patients from Cuba. *J. Med. Virol.* **91**, 1351–1354 (2019).

68.   Acuna-Soto, R., Stahle, D. W., Cleaveland, M. K. & Therrell, M. D. Megadrought and Megadeath in 16th Century Mexico. *Emerg. Infect. Dis.* **8**, 360–362 (2002).

69.   Warinner, C. *et al.* A Robust Framework for Microbial Archaeology. *Annu. Rev. Genomics Hum. Genet.* **18**, 321–356 (2017).

70.   Dabney, J. *et al.* Complete mitochondrial genome sequence of a Middle

Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 15758–15763 (2013).

71. Rohland, N. & Hofreiter, M. Ancient DNA extraction from bones and teeth. *Nat. Protoc.* **2**, 1756–1762 (2007).

72. Meyer, M. & Kircher, M. Illumina Sequencing Library Preparation for Highly Multiplexed Target Capture and Sequencing. *Cold Spring Harb. Protoc.* **2010**, 1–10 (2010).

73. Schubert, M., Lindgreen, S. & Orlando, L. AdapterRemoval v2: rapid adapter trimming, identification, and read merging. *BMC Res. Notes* **9**, (2016).

74. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

75. Huson, D. H. *et al.* MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale Microbiome Sequencing Data. *PLoS Comput. Biol.* **12**, e1004957 (2016).

76. Schubert, M. *et al.* Improving ancient DNA read mapping against modern reference genomes. *BMC Genomics* **13**, 178 (2012).

77. Ondov, B. D., Bergman, N. H. & Phillippy, A. M. Interactive metagenomic visualization in a Web browser. *BMC Bioinformatics* **12**, 385 (2011).

78. Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649 (2012).

79. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

80. Larsson, A. AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**, 3276–3278 (2014).

81. Darriba, D., Taboada, G. L., Doallo, R. & Posada, D. jModelTest 2: more models, new heuristics and high-performance computing. *Nat. Methods* **9**, 772 (2012).

82. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

83. Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian

Phylogenetics with BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* **29**, 1969–1973 (2012).

84.   Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. Posterior Summarization in Bayesian Phylogenetics Using Tracer 1.7. *Syst. Biol.* **67**, 901–904 (2018).

85.   Price, A. L. *et al.* Principal components analysis corrects for stratification in genome-wide association studies. *Nat. Genet.* **38**, 904–909 (2006).

86.   Patterson, N., Price, A. L. & Reich, D. Population Structure and Eigenanalysis. *PLoS Genet.* **2**, e190 (2006).

87.   Renaud, G., Slon, V., Duggan, A. T. & Kelso, J. Schmutzi: estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol.* **16**, 224 (2015).

88.   Kloss-Brandstätter, A. *et al.* HaploGrep: a fast and reliable algorithm for automatic classification of mitochondrial DNA haplogroups. *Hum. Mutat.* **32**, 25–32 (2011).

89.   Weissensteiner, H. *et al.* HaploGrep 2: mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Res.* **44**, W58–W63 (2016).

90.   Skoglund, P., Storå, J., Götherström, A. & Jakobsson, M. Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* **40**, 4477–4482 (2013).

Figure 1. Metagenomic analysis of Colonial individuals reveal HBV-like and B19V-like hits. a, Location of the archeological sites used in this study, HSJN (19.431704, -99.141740) is shown as a triangle and COY (19.347079, -99.159017) as a circle, lines in pink map show current division of Mexico City. **b**, Several individuals discovered in massive burials archaeologically associated with the HSJN and colonial epidemics (photo courtesy of "Secretaria de Cultura INAH, SINAFO, Fototeca DSA"). **c**, Metagenomic analysis performed with MALT 0.4.0 based on the Viral NCBI RefSeq. Viral abundancies were compared and normalized automatically in MEGAN between shotgun (samplename) and capture (samplename-c) NGS data. Only targeted viral families are shown for HBV or B19V positive samples (all samples analysis shown in Supplementary Figure 3), a capture negative control (HSJN177) is shown as well that presented one B19V-like hit and three HBV-like hits after capture.

Figure 2. Ancient B19V and HBV ancient genomes. a, Superimposed damage patterns of ancient HBV (HSJN194) and B19V (HSJNC81, HSJN240, COYC4), X axis shows the position (nt) on the 5' (left) and 3' (right) end of the read, Y axis shows the damage frequency (raw damage patterns are shown on Supplementary Figure 4). **b**, B19V ssDNA linear genome, X axis shows position (nt) based on the reference genome (AB550331), and Y axis shows depth (as number of reads), GC content is shown as a percentage of each 100 bp windows, CDS coverage and average depth are shown under each individual ID. Schematic of the B19V genome is shown at the bottom. Highly covered regions correspond to dsDNA ITRs shown as a red hairpin. **c**, HBV circular genome, outer numbers show position (nt) based on reference genome (GQ331046), outer bars show genes with names, blue bars represent coverage and gray bars shows GC content each 10 bp windows. Coverage and average depth are shown in the center. Low covered region between S and X overlaps with ssDNA region.

Figure 3. Viral Colonial genomes and their human hosts cluster with modern African genetic diversity. Maximum likelihood tree performed on RAxML 8.2.10 (1000 bootstraps) with a midpoint root based on the HBV whole genome (**a**) and B19V CDS (**b**), previously reported ancient genomes are colored, Colonial samples are shown in red. Sequences are named as follows: genotype_ID_sampling.year_country.of.origin_area.of.origin_host. HBV genotype nomenclature is based on letters, while for B19V is on numbers. **c**, PCA showing genetic affinities of ancient human hosts compared to 1000 Genomes Project reference panel, "x" show individuals from the reference panel while other shapes show human hosts from which ancient HBV (HSJN194) and B19V (HSJNC81, HSJN240, COYC4) were recovered.