1

A molecular phylogeny of forktail damselflies (genus *Ischnura*) reveals a dynamic macroevolutionary history of female colour polymorphisms

Rachel Blow[1], Beatriz Willink[2], and Erik I. Svensson[3*]

[1] Department of Zoology, University of Cambridge, Cambridge, UK

[2] School of Biology, University of Costa Rica, San José, Costa Rica 11501-2060

[3] Evolutionary Ecology Unit, Department of Biology, Lund University, SE-223 62 Lund, Sweden

*Corresponding author email: erik.svensson@biol.lu.se

Keywords: ancestral state reconstruction, geographic range, morphic speciation, sexual conflict, StarBEAST2, trans-species polymorphism.

Abstract

Colour polymorphisms are popular study systems among biologists interested in evolutionary dynamics, genomics, sexual selection and sexual conflict. In many damselflies, such as in the globally distributed genus *Ischnura* (forktails), female colour polymorphisms occur in some species. Female-polymorphic species contain two or three female morphs, one of which is male-coloured (androchrome or male mimic) and co-exists with sexually dimorphic (heterochrome) females. These female colour polymorphisms are considered to be maintained by frequency-dependent sexual conflict, but their macroevolutionary histories are unknown, due to the lack of a robust molecular phylogeny. Here, we present the first time-calibrated phylogeny of *Ischnura*, using a multispecies coalescent approach (StarBEAST2), incorporating both molecular data and fossil information of 41 extant species (55% of the genus). We estimate the age of *Ischnura* to be between 13.8 and 23.4 millions of years, i.e. Miocene. We infer the ancestral state of this genus as female monomorphism with heterochrome females, with multiple gains and losses of polymorphisms, evidence of trans-species polymorphisms and a significant positive relationship between polymorphism incidence and current geographic range size. Our study provides a robust phylogenetic framework for future research on the dynamic macroevolutionary history of this clade with its extraordinary diversity of sex-limited female polymorphisms.

Introduction

The evolutionary processes shaping within- and between-population genetic diversity are classical research themes in population genetics and evolutionary biology that go back to the early days of the Modern Synthesis (Crow and Kimura, 1970; Dobzhansky, 1970; Fisher, 1930; Haldane, 1949, 1932; Hartl and Clark, 1997; Provine, 1986; Wright, 1932, 1931). Before the development of molecular tools to quantify genetic variation in natural populations, ecological geneticists and evolutionary biologists often studied discrete and heritable phenotypic polymorphisms governed by one or a few loci (Dobzhansky, 1970; Ford, 1975; Provine, 1986; Schemske and Bierzychudek, 2001; Svensson, 2017; Turelli et al., 2001). Since these early days, genetic colour polymorphisms became popular study systems to address questions about the maintenance of genetic and phenotypic variation (Gordon et al., 2015; Llaurens et al., 2017; Mitchell-Olds et al., 2007; Svensson, 2017; Svensson et al., 2009) and the molecular basis behind suites of correlated traits (Andrade et al., 2019; Gangoso et al., 2011; McKinnon and Pierotti, 2010; Sinervo and Svensson, 2002; Willink et al., 2020; Woronik et al., 2019). Colour polymorphisms are widespread among many animal taxa and their genetic basis is often due to multiple segregating alleles at a single locus, with different degrees of dominance in relation to each other (Svensson et al., 2009; Wellenreuther et al., 2014). Such colour polymorphisms provide excellent opportunities to link the microevolutionary processes that maintain genetic variation within populations (Le Rouzic et al., 2015; Svensson, 2017; Takahashi et al., 2014) to macroevolutionary dynamics of discrete and heritable traits (Jamie and Meier, 2020; Willink et al., 2019).

In many colour polymorphic animals, different male morphs are favoured under different ecological and social conditions and the male morphs are often maintained by balancing natural

or sexual selection, such as negative frequency-dependent selection or overdominance (Blows

et al., 2003; Formica and Tuttle, 2009; Hughes et al., 2013; Lank et al., 1995; Sinervo and

Lively, 1996; Tuttle et al., 2016; Wellenreuther et al., 2014). In contrast to many previous

studies on male colour polymorphisms, there are still relatively few studies on female colour

polymorphisms and other heritable female polymorphisms (Iversen et al., 2019; Karlsson et al.,

2013; Svensson et al., 2009). Whereas male colour morphs are often associated with different

reproductive strategies and sexual competition over access to females, female colour

polymorphisms have been linked to avoidance of male mating harassment and as female

defenses in sexual conflict (Lee et al., 2019; Neff and Svensson, 2013; Willink et al., 2019).

For instance, in the insect order Odonata (dragonflies and damselflies), female colour

polymorphisms occur in several genera and many species (Fincke et al., 2005). Female colour

polymorphisms are particularly common in the globally distributed genus *Ischnura* (Gering,

2017; Gosden and Svensson, 2009; Sirot and Brockmann, 2001; Takahashi and Watanabe,

2010) (Fig. 1) within the superfamily Coenagrionoidea, which includes pond damselflies and

their relatives (Dijkstra et al., 2014).

Within the damselfly genus *Ischnura,* there are both female-monomorphic and female-

polymorphic species (Fincke et al., 2005; Sanchez-Guillen et al., 2011; Willink et al., 2019).

Of the approximately 75 species in the genus *Ischnura*, c.a. 40% are female-monomorphic,

having only one distinct female phenotype, whilst another 40% have two, and some species

even have three female morphs (Willink *et al.* 2019). The female-morph status of the remaining

20% of species is currently unknown (Willink *et al.* 2019). In female-polymorphic species of

Ischnura, females are either phenotypically similar in colouration and sometimes also body

shape to males (referred to here as androchrome females, or male mimics) (Gosden and

Svensson, 2009), or they are phenotypically different from males (referred to here as

heterochrome females). These colour polymorphisms are limited in their expression to females,

and males are monomorphic (Svensson et al., 2020, 2009a). Such female colour polymorphisms

are particularly suited for phylogenetic comparative studies of the macroevolution of discrete

and heritable morphs.

The classical explanation for the evolutionary origin and adaptive function of female colour

polymorphisms in *Ischnura,* and other damselflies is the "male-mimicry hypothesis"

(Robertson 1985; Cordero 1989). Male-mimics are thought to have evolved as a response to

selection from sexual conflict over mating, which favoured visual features that made such male-

like females less recognizable as potential mates to males (Cordero et al., 1998; Gosden and

Svensson, 2009; Willink et al., 2019; Xu et al., 2014). The mimicry hypothesis predicts that the

common ancestor of polymorphic damselflies was a monomorphic species with heterochrome

females, which was invaded after a novel male mimicking female phenotype emerged that

subsequently increased in frequency due to a frequency-dependent fitness advantage of male

mimicry. An alternative, though not mutually exclusive hypothesis for the evolution and

maintenance of such female polymorphisms are flexible search images in males (Fincke, 2004;

Miller and Fincke, 1999). Under this hypothesis, males are expected to switch mating

preferences between morphs, depending on their frequency, resulting in increased male

preferences for currently common morphs (Van Gossum et al., 2001). The most common

female morph is then expected to be disproportionally harassed, resulting in negative

frequency-dependent selection (NFDS) and the stable maintenance of the polymorphism across

many generations (Gosden and Svensson, 2009; Iserbyt et al., 2013, 2011; Le Rouzic et al.,

2015; Svensson et al., 2005; Takahashi et al., 2014, 2010). Unlike a pure version of the male-

mimicry hypothesis, female colour polymorphisms in *Ischnura* could under this latter scenario

have evolved from an ancestral state of either heterochrome or androchrome females and with

female monomorphism. We re-emphasize that these two hypotheses are not mutually exclusive,

and both can operate simultaneously, with a crucial role for NFDS maintaining female

polymorphisms under both scenarios. Regardless of mechanism, NFDS is therefore thought to

be the main force maintaining such female polymorphisms (Le Rouzic et al., 2015; Svensson

et al., 2005; Takahashi et al., 2010).

Female-monomorphic species of *Ischnura* have received less attention compared to female-

polymorphic taxa, and consequently we know very little about why there is such a striking

variation in the presence or absence of female-limited polymorphisms in different subclades

within this genus. One explanation could be that female-monomorphic lineages reflect an

ancestral state with low intensity of sexual conflict. If so, female polymorphisms may have

subsequently evolved in lineages experiencing novel selective regimes of more intense sexual

conflict. This could occur, for instance, due to a rapid increase in population density or

transition to a new mating system. Alternatively, female-monomorphic lineages could reflect

transitions from an ancestral state of female-limited polymorphism, via founder effects or

genetic drift, or through the selective fixation of one of the alternative previously sympatric

morphs (Corl et al., 2010; West-Eberhard, 1986). Such speciation by morph loss has been

termed "morphic speciation" (Corl et al., 2010; West-Eberhard, 1986). The morphic speciation

hypothesis explicitly predicts that monomorphic taxa should primarily be located at the terminal

tips on a phylogeny, with polymorphism being the ancestral state (Corl et al., 2010). Regardless

of the generality and validity of the morphic speciation hypothesis, our knowledge about the

relationship between polymorphism andspeciation is limited and there are only a few

phylogenetic comparative studies in this area (Hugall and Stuart-Fox, 2012; McLean and

Stuart-Fox, 2014; Schwander and Leimar, 2011). Recently, it has also been emphasized that

some colour polymorphisms can be maintained across speciation events resulting in

transspecies polymorphisms (Jamie and Meier, 2020).

There have been a few previous comparative studies in the genus *Ischnura*, aiming to

understand the ecological correlates of colour polymorphisms and their evolutionary origin.

These previous studies suggested that female-limited colour polymorphisms are more common

in species with female polyandry, than in species with (putative) female monandry, consistent

with polymorphism being promoted by sexual conflict (Mattern and Van Gossum, 2008;

Robinson and Allgeyer, 1996). However, these previous comparative studies suffer from

several limitations, as they did not take into account the shared evolutionary history of related

species (Robinson and Allgeyer, 1996) or they used highly sensitive methods to infer ancestral

character states such as maximum parsimony (Mattern and Van Gossum, 2008). The use of

maximum parsimony is particularly problematic as it provides poor estimations of tree

topologies (Wright and Hillis, 2014) and unreliable inferences of ancestral character states,

especially for rapidly evolving characters like behavioural traits (Cunningham et al., 1998;

Schluter et al., 1997). For these reasons, inferences of ancestral character states should

preferably come from Bayesian or Maximum Likelihood (ML) approaches (Cunningham et al.,

1998; Schluter et al., 1997; Wright and Hillis, 2014). A previous study claimed, based on the

results from from maximum parsimony, that female colour polymorphism was the ancestral

state in *Ischnura* (Mattern and Van Gossum, 2008). Such an inferred polymorphic ancestral

state in *Ischnura* is, however, unlikely to be robust given the problems with maximum

parsimony, and this hypothesis would therefore need to be re-evaluated, using updated

phylogenetic information and more reliable modern comparative methods.

Here, we present novel phylogenetic and comparative data and analyses from the genus

Ischnura aimed to shed light on the origin of female colour polymorphisms, and we also

estimated the age and studied the evolutionary history of this genus. We use a multispecies

coalescent model to simultaneously estimate a posterior distribution of species tree topologies

and divergence times for 55% of species in the genus *Ischnura*. We then investigate the

macroevolutionary history of female-limited polymorphisms by reconstructing the ancestral

state of this trait. We proceed by examining whether the presence of female-limited colour

polymorphisms are associated with large geographic range sizes, a factor that is likely to reflect

population size and the potential for morphic speciation by neutral processes. A positive

association between the presence of female colour polymorphisms and geographic range would

be expected in both of the following scenarios: 1) if morphs are lost by neutral or selective

processes after a reduction of population (and range) size or 2) if the effects of female colour

polymorphisms on population fitness allow female-polymorphic species to sustain larger

populations over a greater geographic range.

Our study contributes to our understanding the macroevolutionary history of this genus, the

origin of female colour polymorphism and the demographic factors promoting the maintenance

of these polymorphisms. In presenting these results, we hope to stimulate future empirical

investigations on these and other damselfly clades, and we discuss some promising hypotheses

for follow-up studies. We note that many damselfly groups, although popular in much

ecological and evolutionary research (Cordoba-Aguilar, 2008) are still largely unknown with

respect to phylogenetic relationships and evolutionary history, although some recent progress

has been made, particularly within the family Coenagrionidae (Beatty et al., 2017; Callahan and

McPeek, 2016; Dijkstra et al., 2014; Swaegers et al., 2014; Torres-Cambas et al., 2019; Vega-

Sánchez et al., 2019).

1. Materials and Methods

2.1 Data collection

We have gathered DNA-sequence data, phenotypic data and geographic range size data for 41 species of *Ischnura*. Two of the taxa in this dataset are nominally in the genus *Pacificagrion*, however recent molecular data shows they are nested within the *Ischnura* clade (Willink et al. 2019). Likewise, two other taxa, including a yet undescribed species, were classified in the genus *Amorphostigma* (Marinov et al., 2015). However, Dijkstra et al. (2014) subsumed *Amorphostigma* into *Ischnura*, and recent molecular data support this change (Willink et al. 2019). Here, we use the generic names *Ischnura* and *Pacificagrion*, according to the "World Odonata List" (https://www.pugetsound.edu/academics/academic-resources/slater-museum/biodiversity-resources/dragonflies/world-odonata-list2/), which maintains a taxonomic classification of Odonata with currently valid names.

We obtained sequences for five molecular markers including nuclear (H3: 28.86 kb, PMRT: 51.68 kb, 28S: 67.91 kb) and mitochondrial regions (COI: 57.72 kb, 16S: 47.64 kb) from 147 specimens, of which 64 were sequenced for this study (Table S1). Genomic DNA was extracted from 1-2 legs using a Qiagen DNeasy tissue kit and following the manufacturer's instructions, except that we increased the amount of proteinase K and incubation time for museum samples collected before 2005, in order to obtain a higher DNA yield. Polymerase chain reaction (PCR) primers and amplification programs used for the molecular markers were the same as in Willink et al. (2019). Amplification products were purified using the Affymetrix ExoSAP-IT reagent (Cat. No. 78200) and sequenced on an ABI 3100 capillary sequencer at Lund University. These novel sequence data were complemented with already-published sequences (Table S1),

downloaded   from   the   National   Centre   for   Biotechnology   Information

(https://www.ncbi.nlm.nih.gov/genbank/). There were no indels in the sequences of two

molecular markers (COI and H3), which were aligned using MUSCLE (Edgar, 2004). To

reduce gap overmatching in ribosomal sequences (16S and 28S) and intronic regions (present

in PMRT), we used the phylogenetically informed algorithm PRANK (Löytynoja and

Goldman, 2008, 2005) to align these data.

Data on female colour morph status (i.e. female-monomorphic, or female-polymorphic) has

been recently compiled for *Ischnura* damselflies, based on published literature, field

observations, examination of museum specimens and online resources (Willink et al. 2019).

We complemented these data with information on whether female-polymorphic species had

two or three morphs (i.e. female-dimorphic or female-trimorphic) (Table S2). Here, species

with a single female morph were classified as female-monomorphic with androchrome females,

if they had only one female morph with colouration and patterning similar to that of conspecific

males, or as female-monomorphic with heterochrome females, if they had only one female

morph with colouration and patterning differing from the conspecific males. If a species

contained two distinct female morphs (always one androchrome and one heterochrome), we

classified it as female-dimorphic. Finally, species that had three morphs (which were always

one androchrome and two heterochrome female morphs), were classified as female-trimorphic.

We complemented these colour-state data with coarse-grained estimates of geographic range

size (Table S2). Because accurate geographic distribution data is lacking for many species, we

estimated geographic range using occurrence data from administrative regions, mainly

countries, and where available, states or provinces (Table S2). Presence data was translated to

geographic range size (km^2) by adding the areas of all the countries or other administrative

regions where a species is known to occur (Table S2), as has been done previously in

comparative studies where geographic range is taken as a proxy for population size (Ross et al.,

2013). The areas of administrative regions were extracted from the GADM database of global

administrative areas v. 3.6 (Global Administrative Areas, 2018).

1.2 *Time-calibrated species tree*

We analysed sequence data under a multispecies coalescent model using StarBEAST2 (Ogilvie

et al., 2017). As the model assumes free recombination amongst unlinked loci, we linked the

tree models between mitochondrial sequences. We placed a birth-death prior, conditional on

the root, on the species tree to allow for extinction, set the gene ploidy to 0.5 for mitochondrial

sequences and 2.0 for nuclear sequences and allowed population-size to be analytically

integrated to avoid over-parameterization. We used a relaxed uncorrelated log normal (UCLN)

clock per locus to allow for variation in evolutionary rates between lineages. Substitution

models for each locus were inferred during the analysis using the package "bModelTest"

(Bouckaert and Drummond, 2017) to allow for site model uncertainty. We placed a lognormal

prior with a mean of 0.0115 and standard deviation of 1.0 on each relaxed mitochondrial clock

and a broader lognormal prior with a mean of 0.0115 and standard deviation of 2.0 on each

relaxed nuclear clock to centre clock rates on the suggested insect mitochondrial divergence

rate of 2.3% per million years (Brower, 1994) whilst allowing for clock rate uncertainty,

particularly in the nuclear loci.

Data for *Ischnura* fossils were downloaded from the Paleobiology database (www.pbdb.org)

on 14[th] May 2018. There were three fossils classified as either in or closely related to the genus

Ischnura. Two of these fossils were relatively young (7.2 - 5.3 mya), each one consisting of a

single wing with missing fragments (Gentilini and Bagli, 2004). The third fossil was preserved

in Dominican amber (20.44 – 13.82 mya) and was identified and described as *Ischnura velteni*

based on a combination of wing venation and morphological traits in other body parts (Bechly,

2000). The combination of these characters suggests that *I. velteni* belongs to the extant genus

Ischnura (Bechly, 2000), and we therefore considered it the most appropriate crown fossil with

which to calibrate the root age of the *Ischnura* tree. We placed an exponentially distributed

prior on the root age of the species tree with an offset of 13.8 to ensure the most recent common

ancestor (MRCA) of *Ischnura* could not have appeared after the minimum age of this known

member of the genus. The mean of the exponential was drawn from a gamma hyperprior with

a shape parameter $\alpha = 5.0$ and an inverse scale parameter $\beta = 2.0$. This hyperprior specification

centred the root age prior distribution around 10 mya prior to the offset, allowing a moderately

high level of uncertainty in the age difference between the fossil and the MRCA of *Ischnura.*

We ran two separate Markov Chain Monte Carlo (MCMC) algorithms for 200 million iterations

with a sampling rate of 20,000. We confirmed that independent chains were stationary and had

converged for these and all subsequent analyses in R v.3.6.1 (R Core Team 2019) using the

"coda" package (Plummer et al., 2006). We combined posterior trees using LogCombiner

(available as part of BEAST v2.4.7)(Bouckaeret al., 2014), simultaneously discarding 20% of

the posterior as burnin, and summarized them into a maximum clade credibility (MCC) tree

using TreeAnnotator (available as part of BEAST v2.4.7). The MCC tree and all further

phylogenetic plots were annotated in R v.3.6.1 (R Core Team 2019) using the packages "ape"

(Paradis and Schliep, 2019), "ggtree" (Yu et al., 2017) and "ggplot2" (Wickham, 2016).

1.3 *Reconstruction of ancestral character states of female colour polymorphism*

We used the Multistate-method in BayesTraits version 3.0 (Pagel et al., 2004) to infer female-

colour character states at ancestral nodes, given this new *Ischnura* phylogeny and the female-

colour character states of extant species. To account for phylogenetic uncertainty, we ran the

model using a sample of 1000 trees from the combined species-tree posterior of the two

independent runs. Additionally, we ran the model using reversible jump MCMC to account for

uncertainty in the number of model parameters. In this model, the number of different transition

rate parameters and their values are sampled in proportion to their posterior probability.

Consequently, in each posterior sample a number of transition rates are set to zero, and the non-

zero rates belong to one or more categories. We placed an exponential prior on all rate

parameters, with the mean seeded from a uniform hyperprior bounded between 1 and 100. We

ran two separate chains for 10 million iterations, sampling every 1000 with 20% posterior

burnin. Trees were scaled to an average branch length of 0.01 to prevent various rates from

becoming too small. Although our main focus is on ancestral state reconstructions, we present

the posterior distributions of transition rate parameters in the Supporting Material (Fig. S1).

1.4 *Linking female colour polymorphism to geographic range size*

We investigated if the occurrence of female colour polymorphisms could be influenced by

demographic changes by using geographic range size of extant species as a proxy for population

size. Geographic range size is also a biologically interesting variable in the light of recent work

showing that female colour polymorphisms can increase population fitness (Takahashi et al.,

2014). e used a Bayesian phylogenetic mixed-effect model (BPMM)(Hadfield and Nakagawa,

2010) implemented in the package MCMCglmm v. 2.29 (Hadfield, 2010) to test for such

phylogenetic correlation. Female-morph status was treated as a binary response variable, with

species being classified as either female-monomorphic or female-polymorphic. Since the

occurrences of female-trimorphic species and female-monomorphic species with androchrome

females were low, we classified these species-states as being female-polymorphic and female-

monomorphic, respectively. We thus investigated if the probability of a species being female-

polymorphic were significantly and positively related to geographic range size (a continuous

predictor variable).

We accounted for phylogenetic uncertainty by using a random sample of 2100 species-trees

generated in StarBEAST2. To estimate the phylogenetic random effect, each tree was sampled

for 10000 iterations, of which only the last iteration was saved to the posterior, and the first 100

were also discarded as burnin (O'Connor et al., 2018). We used a Knonecker prior (mu=0, V=

σ2units + π2/3) for the fixed effects as it is relatively flat when a logit link function is used

(Hadfield, 2015). We used a parameter-expanded χ^2 distribution (V=1 nu=1000, alpha.mu=0,

alpha.V = 1) for the prior on the phylogenetic variance components (Villemereuil et al., 2013).

The residual variance was fixed to 1, as it cannot be identified for binary response models. As

a statistical test of whether the probability of female colour polymorphism increases with range

size, we report the fraction of the posterior distribution in which increasing range size does not

result in an increasing probability of female colour polymorphism. The results of this analysis

were visualized by plotting the expected probability of female colour polymorphism against

estimated geographic range size.

2. Results

3.1 *Divergence time estimation*

Our time-calibrated phylogeny suggests that the age of the MRCA of the genus *Ischnura* lies

in a 95% highest posterior density (HPD) interval between 13.8 and 23.4 mya, i.e. during the

early Miocene (Fig. 2). A comparison of the posterior root age in our model to a model ran

without molecular data suggests the sequence data were sufficiently informative to narrow this

posterior distribution (Fig. 3), although age estimates had quite broad error margins (Fig. 2).

Nonetheless, divergence dating revealed several interesting features, such as the very recent

divergence of two of the most intensely studied polymorphic species in Europe (*I. elegans* and

I. graellsii) (Fig. 2). Our phylogenetic analysis also showed that the species that feature in most

evolutionary and ecological studies (*I. elegans*, *I. graellsii*, *I. ramburii* and *I. senegalensis*) are

part of awell supported clade (Clade II excluding *I. capreolus* and *I. barberi;* Fig. 2), with a

common ancestor less than 10 mya (Fig. 2). We found that all species from Australia and the

South Pacific islands, except *I. heterosticta,* form a single clade (Clade I; Fig. 2). Similarly, we

recovered a strongly supported Palearctic clade (Clade III; Fig. 2) and strongly supported clades

distributed primarily in the Neotropical (*I. prognata* to *I. cruzi*) and Nearctic (*I. verticalis* to *I.*

kellicotti) regions (Clade IV; Fig. 2).

2.2 Ancestral state reconstruction of female colour polymorphism

Out of the 41 species included in this study, 13 were female-monomorphic with heterochrome

females (**category 1**), 22 were female-dimorphic (**category 2**), three were female-trimorphic

(**category 3**) and finally three were female-monomorphic with androchrome females (**category**

4) (Fig. 4). The MRCA of *Ischnura* was reconstructed as being female-monomorphic with

heterochrome females, i.e. the ancestor was sexually dimorphic, with an mean posterior

probability (PP) of 81% (Fig. 4). Clade I (Fig. 2), which includes most taxa from the Pacific

islands was also inferred as having a heterochrome female ancestor (PP = 81%; Fig. 4). All

female-trimorphic species shared a female-trimorphic common ancestor (PP = 100%; Fig.

4).These taxa are included in Clade II, for which the most probable ancestral state if female

dimorphism (PP = 73%; Fig. 4).

Two evolutionary transitions occurred with high posterior probability during the diversification

of *Ischnura*, according to the RJ MCMC analysis. The transition rate from female colour

dimorphism (**category 2**) to female colour monomorphism with androchrome females

(**category 4**) (i.e. the evolutionary loss of the heterochrome female morph) was above zero in

99.7% of the posterior (Fig. S1). Similarly, the transition from female colour monomorphism

with heterochrome females (**category 1**) to female colour dimorphism (**category 2**) (i.e. the

evolutionary gain of an androchrome female morph) occurred with non-zero rate in 94% of the

posterior (Fig. S1). In contrast, the least probable transition, from female colour monomorphism

with heterochrome females (**category 1**) to female colour trimorphism (**category 3**) occurred

in only 7% of the posterior (Fig. S1). The extant data on female-colour states and the posterior

distribution of *Ischnura* time trees are therefore consistent with multiple evolutionary histories

and different combinations of other character-state transitions being present or absent.

3.3 *Geographic range size and female-colour polymorphism*

Current geographic range sizes varied from 4.3 x 10^7 Km^2 in the largely subtropical species *I.*

senegalensis, which is distributed from sub-Saharan Africa to Japan, to 222 Km^2 to species

endemic to American Samoa (Table S2). We found that all island-endemic species are female-

monomorphic, except for *I. foylei*, which is female-dimorphic and only known from the Jambi

province in Sumatra, Indonesia (Table S2). In contrast, the three species distributed across the

longitudinal range of the Old World, *I. senegalensis*, *I. pumilio* and *I. elegans,* are either female-

dimorphic or -trimorphic. The probability of a species being female-polymorphic increased

significantly with its geographic range size (PMCMC = 0.003; Fig. 5). The probability of

female colour polymorphism increased by ~ 5% (posterior mean = 0.051, 95% HPD interval =

0.008 – 0.990) with an increase of 1 x10^6 Km^2 in the geographic range size.

4. Discussion

Here, we have presented the first molecular time-calibrated phylogeny for the damselfly genus *Ischnura*, an insect group that has been subject to much interest in evolutionary and ecological research, particularly sexual selection, sexual conflict and female colour polymorphism (Fig. 1). The general branching patterns in this new phylogeny are largely and qualitatively in agreement with previous molecular phylogeniescovering a subset of the taxa included here (Chippindale et al., 1999; Willink, 2018; Willink et al., 2019). Both dragonflies (suborder Anisoptera) and damselflies (suborder Zygoptera) have long been popular study organisms in ecology, evolution and conservation biology (Cordoba-Aguilar, 2008). However, time-calibrated molecular phylogenies that could facilitate phylogenetic comparative studies in these two insect groups are still relatively few, although recently the situation has been improving (Callahan and McPeek, 2016; Damm et al., 2010; Letsch et al., 2016; Waller and Svensson, 2017).

Using a UCLN molecular clock and fossil calibration, we inferred the MRCA of *Ischnura* as living between 13.8 and 23.4 mya, that is, the early Miocene. This is earlier than the MRCA of the closely related damselfly genus *Enallagma*, which has an estimated age of about 10 my (mid- to late Miocene), and where many speciation events seem to have been driven by glacial cycles during the Pleistocene (Callahan and McPeek, 2016). An early Miocene origin of *Ischnura* is also slightly earlier than the inferred origin of the closely related Palearctic damselfly genus *Coenagrion,* during the later part of Miocene 15 mya (Swaegers et al., 2014) and the two genera *Melanosobasis* and *Nesobasis* in Fiji originating during late-middle Miocence (8-12 mya)(Beatty et al., 2017). We emphasize that our age estimate for the MRCA of *Ischnura* rests on the assumption that *I. velteni* is a crown fossil (Bechly 2000). Future studies

based on alternative dating strategies will be important to validate this inferred early Miocene

origin of *Ischnura*.

According to our phylogeny and ancestral state reconstruction, all of the well-studied species

in *Ischnura*, *I. senegalensis*, *I. graellsii, I. elegans* and *I. ramburii* (Gering, 2017; Robertson,

1985; Sirot et al., 2003; L.K. Sirot and Brockmann, 2001) are part of a single clade, likely

descended from a female-dimorphic ancestor (Fig. 4), that lived less than 10 mya (Figs. 2,4).

While the genus *Ischnura* has increasingly attracted attention in ecological an

microevolutionary studies, there is much untapped potential in understanding its

macroevolutionary history and repeated transitions between female colour states. Our ancestral

state reconstruction suggests that the ancestor of the genus *Ischnura* was female-monomorphic

with heterochrome females, although we caution that the support for this inference was 81%

(Fig. 4). Such a sexually dimorphic ancestor would imply multiple evolutionary transitions to

obtain female-dimorphism, through the evolutionary emergence of a male-mimic. This

evolutionary transition occurred with non-zero rate in 94% of the posterior distribution of

transition rate estimates in our RJ MCMC analysis (Fig. S1). This suggests that female colour

polymorphism, with heterochrome and androchrome females coexisting within species, has

evolved convergently multiple times in this genus, presumably as an evolutionary response to

increased intensity of sexual conflict. Specifically, the emergence of a novel male-mimic might

have have been possible if it gained an initial fitness benefit due to advantages in intersexual

mimicry through reduced male mating harassment (Robertson, 1985; Svensson et al., 2009;

Willink et al., 2019). Such male-mimics might have subsequently increased to high frequencies

in several populations and species, and later became maintained as stable polymorphisms by

frequency- and/or density dependent sexual selection caused by male mating harassment

(Gosden and Svensson, 2009; Le Rouzic et al., 2015; Svensson et al., 2005; Takahashi et al.,

2014, 2010).

We further note that the female-trimorphic clade, containing the well-studied European species

I. elegans and *I. graellsii,* most likely evolved from a female-dimorphic ancestor (Fig. 4). The

evolutionary emergence of a third female morph in an already female-polymorphic system

might indicate that sexual conflict through male mating harassment was particularly intense in

this clade, which would fit with available empirical evidence from previous field and

experimental studies (Cordero, 1992; Cordero et al., 1998; Gosden and Svensson, 2009;

Takahashi et al., 2014; Willink et al., 2019). In support of this, *I. graellsii* and *I. elegans* have

some of the longest copulation durations of all investigated species so far and mate for several

hours (Robinson and Allgeyer, 1996). The long copulation duration in these two female-

trimorphic species contrasts with the much shorter copulations and lower mating rates of some

of the North American monomorphic female species, some of which have also been suggested

to be monandrous (Robinson and Allgeyer, 1996). Thus, high mating rates, high degree of

density-dependent male mating harassment (Gosden and Svensson, 2009) and long copulation

duration (Robinson and Allgeyer, 1996) in the female-trimorphic clade containing *I. elegans*

and *I. graellsii* might all have contributed to the evolution and maintenance of a new morph in

these species. Models of sexual conflict suggest that an evolutionary outcome is the emergence

of multiple genetic clusters of both males and females (Gavrilets and Waxman, 2002; Hayashi

et al., 2007). There is now empirical evidence for female phenotypic polymorphisms that might

have evolved as a response to sexual conflict in a small but growing number of taxa (Iversen

et al., 2019; Karlsson et al., 2013; Lee et al., 2019; Moon and Kamath, 2019; Reinhardt et al.,

2007; Svensson et al., 2005).

The results of the present study run counter to the conclusions in a previous comparative study,

where the authors used maximum parsimony, and claimed that the ancestral state in *Ischnura*

was female polymorphism, rather than female monomorphism (Mattern and Van Gossum,

2008). However, this previous parsimony-based study did not take into account phylogenetic

uncertainty in the inference of ancestral character states and did not include some of tropical

taxa, the majority of which are female-monomorphic in our dataset (Table S2). Applying

statistical models of character evolution that take in to account phylogenetic uncertainty and

aim for a complete or at least unbiased taxonomic sampling, as we have done here, are important

steps to produce robust ancestral state estimations that appropriately represent uncertainty

(Cunningham et al., 1998; Heath et al., 2008; Losos, 2011; Pagel et al., 2004; Revell, 2013).

This is particularly important when inferring the ancestral states of rapidly evolving characters,

such as behavioural traits and the female colour polymorphisms studied here, which show a

dynamic macroevolutionary history with several gains and losses across the phylogeny (Fig. 4).

Polymorphisms have been suggested to promote speciation if different morphs diverge

phenotypically in sympatry, followed by selective or random loss in allopatry, resulting in the

fixation of alternative morphs in different populations (Jamie and Meier, 2020; West-Eberhard,

1986). This process – termed "morphic speciation" – predicts that monomorphic taxa are

commonly found at the terminal tips on a phylogeny (Corl et al., 2010; West-Eberhard, 1986).

There is some evidence for such a process among populations of colour polymorphic side-

blotched lizards (*Uta stansburiana*)(Corl et al., 2010). Our results do not lend support for

widespread morphic speciation, but are consistent with losses of heterochrome females in

female-dimorphic taxa, as this evolutionary transition occurred in 99.7% of the posterior in the

RJ MCMC analysis. The most likely candidate for such potential morphic speciation is the

common ancestor between *I. verticalis* and *I. posita*. These taxa are part of a North American

clade that diversified rapidly during the Pleistocene (*I. damula* to *I. cervula;* Fig. 2). Such rapid

speciation and diversification could have resulted from repeated glacial cycles, as in the closely

related genus *Enallagma* that contains many sympatric young species in North America

(Callahan and McPeek, 2016; McPeek and Gavrilets, 2006). Such glacial cycles and repeated

Pleistocene bottlenecks could have led to morph fixation of a male-like female androchrome

morph in the ancestor of *I. posita* during a period of allopatry (Fig. 4). *I. posita* is currently a

widely distributed species, ranging from Nova Scotia in Canada to Central America (Paulson,

2012; Table S2). If morph loss in the ancestor of *I. posita* resulted from genetic drift during a

period of allopatry and population contraction, it must have then been followed by rapid and

recent range expansion. While this hypothesis remains to be investigated for *I. posita,* closely

related and young sympatric taxa in *Enallagma* are thought to be the result of rapid speciation

and range expansion after glacial retreat in the Quaternary (McPeek and Gavrilets, 2006;

Turgeon et al., 2005)*.*

Another potential outcome of morphic speciation leading to fixation of male-like androchrome

females is *I. kellicotti*, although it is less clear whether the lineage leading to *I. kellicotti* had a

female-polymorphic ancestor (Fig. 4). Nevertheless, the potential for morphic speciation in *I.*

kellicotti is interesting because females display an interesting pattern of ontogenetic colour

changes. Sexually immature females start with a bright orange colouration that turns blue and

male-like in mature females, and develops increasing pruinescence with age thereafter

(Paulson, 2009, 2012). If used as a signal of reproductive availability (cf. Willink et al. 2019),

the evolution of distinct developmental colour phases may have reduced male mating

harassment and facilitated morphic speciation. We also note that in several relatively distantly

related species of *Ischnura*, immature females express orange colouration prior to reproductive

maturity (e. g. in *I. ramburii, I. pumilio* and *I. senegalensis*; Fig. 1E-G) which could be such a

signal of reproductive unsuitability. Alternatively, an evolutionary transition in habitat and

mating system could have led to morph loss and morphic speciation in the ancestor of *I.*

kellicotti. Individuals of *I. kelicotti* ("Lilypad forktail") are typically found sitting on lilypad

leaves in ponds (Paulson, 2012, 2009), consistent with a switch from non-territorial male-male

scrambling competition over females with high mating harassment, to a new mating system of

partial male territoriality with a lower degree of mating harassment. Such relaxed pressure from

sexual conflict may have lead to the loss of female polymorphism. We propose that a

combination of "phylogenetic natural history" (Uyeda et al., 2018) and more detailed

knowledge about the molecular and genetic basis of female colour development (M. Takahashi

et al., 2018; Willink et al., 2020) could shed light on this issue and increase our general

understanding of other macroevolutionary events in the genus *Ischnura*.

The presence of female colour polymorphism was significantly and positively correlated with

geographic range size (Fig. 5), which could come about in at least three different, but non-

mutually exclusive, ways. First, species with large ranges are also likely to have large effective

population sizes and this will decrease the effects of genetic drift and morph fixation. It is

interesting to note here that several of the female-monomorphic taxa are endemic to small

islands in the Pacific Ocean (Table S2). Second, and certainly not mutually exclusive, species

with larger ranges and associated larger population sizes, might experience stronger sexual

conflict due to higher population densities, consistent with more intense male mating

harassment under such conditions (Gosden and Svensson, 2009). Finally, female-polymorphic

species might also be able exploit a broader range of niches and thereby also achieve higher

population mean fitness compared to female-monomorphic species, with positive effects on

both population size and geographic range size (Forsman, 2016; Svensson, 2017; Takahashi et

al., 2014; Y. Takahashi et al., 2018; Takahashi and Noriyuki, 2019; West-Eberhard, 1986).

Although the data in this study does not allow us to firmly distinguish between these non-mutually exclusive explanations for the positive relationship between the incidence of colour polymorphism and geographic range size (Fig. 5), these different scenarios clearly deserved attention in future studies. A positive relationship between geographic range size and presence of colour polymorphism in *Ischnura* and other related damselflies was reported in a recent study, although without considering phylogenetic relatedness among species, as no molecular phylogeny was available at that time for *Ischnura* (Takahashi and Noriyuki, 2019).

Interestingly, the least supported nodes of our MCC tree are in Clade II and the North American clade within Clade IV (*I. kellicotti* to *I. verticalis*; Fig. 2), which are both dominated by female-polymorphic taxa (Fig. 4). These subsets of taxa were not particularly underrepresented in our dataset (Table S1). There are at least two potential biological explanations for this pattern. First, interspecific reproductive interference may be boosted by sexual conflict over mating rates (McPeek and Gavrilets, 2006; Svensson et al., 2009) and might therefore be more common between female-polymorphic species. This could result in more frequent hybridization and introgression in polymorphic clades (Okude et al., 2020; Sanchez-Guillen et al., 2005; Sánchez-Guillén et al., 2011). Second, faster speciation when population sizes are large can result in deep coalescence among lineages and thus increase gene tree discordance (Maddison, 1997).If female-polymorphic species have larger population sizes as our results suggest (Fig. 5), and colour-polymorphisms accelerate speciation (Hugall and Stuart-Fox, 2012), then the speciation dynamics of female-polymorphic clades may render them particularly challenging for phylogenetic inference. These alternative evolutionary scenarios are clearly worthy of further research.

Finally, in one, and potentially two, of the female-polymorphic clades, the female

polymorphism seem to have arisen early and was subsequently maintained across multiple

speciation events (Fig. 4). Therefore, the *Ischnura* female colour polymorphisms is an example

of trans-species polymorphisms (Jamie and Meier, 2020), when multiple allelic variants

become established in a population prior to speciation and are subsequently passed on to

descendant taxa across multiple speciation events (Klein, 1987). Other examples of trans-

species polymorphisms are rare, the most well-known being immune system polymorphisms

and blood group variation in great apes (Azevedo et al., 2015) and self-incompatibility allelic

systems in plants (Goldberg et al., 2010; Richman et al., 1996; Richman and Kohn, 2000).

However, such trans-species polymorphisms may be more common than the limited current

evidence suggests, particularly in colour polymorphic taxa (Corl et al., 2010; Jamie and Meier,

2020). In the damselfly genus *Ischnura*, some female-polymorphic species have higher

population-level fitness when female morph frequencies are even, compared to when they are

more biased (Takahashi *et al.* 2014) and in this study we also showed that polymorphic species

have larger range sizes than monomorphic species (Fig. 5). Although these female colour

polymorphisms are maintained by negative frequency-dependent selection at the individual

level (Le Rouzic et al., 2015; Svensson et al., 2005; Takahashi et al., 2010) the higher-level

population effects mentioned above might also indicate an additional potential role for species-

level selection and biased extinction of female-monomorphic taxa. Such higher level-effects

would be similar to how self-incompatability alleles are maintained in plant mating systems by

species selection (Goldberg et al., 2010).

Conclusions

580   The results in this study suggest that the damselfly genus *Ischnura* originated in the early

581   Miocene, with ancestral females being monomorphic and markedly distinct from males in their

colour pattern (heterochrome). The female colour polymorphisms in this genus arose on several occasions through repeated origins of a male-mimic in different species. Our ancestral state reconstruction also suggests that female colour polymorphisms have survived through multiple speciation events in at least two clades, contrary to the expectation of speciation by selective or random morph loss in allopatric populations (morphic speciation). However, we find at least one potential loss of the female colour polymorphism, and specifically of heterochrome females, that could have resulted in morphic speciation. The multiple transitions in female-morph states across the phylogeny imply that the selective pressures from sexual conflict differ betweentaxa, and that the strength of frequency-dependent conflict might be related to local population sizes and densities, with consequences for geographic range size and potentially also for extinction risk. This study provides a foundation for future micro- and macroevolutionary studies on how selective and neutral processes have shaped the evolution of discrete female-morphs in *Ischnura* damselflies, by providing a phylogeny for future comparative studies in this genus.

Acknowledgements

We are grateful to Debora Goedert and Masahito Tsuboi for constructive criticisms of an early draft of this manuscript. We wish to thank Tammy Ho, Chiara De Pasqual and Eva Friman for assistance in obtaining molecular sequence data. We also wish to thank all the following odonatologists who kindly provided samples for the phylogenetic analyses and shared their observations on colour variation in *Ischnura* species: Sylvain Charlat, Peter and Elaine Cowan, Nathalia von Ellenrieder, Sonia Ferreira, Kawsar Khan, Oleg Kosterin, Viktor Nilsson-Örtman, Ryo Futahashi, Roselyn and David Sparrow, Oliver Flint, curator of Odonata in the National Entomological Collection at the Smithsonian Institute, and Bill Mauffray, curator of Odonata at the Florida State Collection of Arthropods. Specimens were collected with permits from the

Guyana Environmental Protection Agency (Reference No. 010715 BR001 and 012615 SP:

002), from Secretaría de Ambiente y Desarrollo Sustentable, Argentina (Ref No. 11084/16) and

from the Cameroon Ministry of Forestry and Wildlife (Ref. No. 0000034). We acknowledge

the logistic support from the Iwokrama International Centre for Rain Forest Conservation and

Development, Guyana, the Karanambu Trust, Guyana, and the National Park Administration,

Argentina. Funding for this study was provided by research grants from The Swedish Research

Council (VR: grant no. 2016-03356), Carl Tryggers Foundation (CTS), Gyllenstiernska

Krapperupstiftelsen (grant no. KR2018-0038), Lunds Djurskyddsfond, The Royal

Physiographic Society in Lund, Stiftelsen Olle Enqvist Byggmästare and Stina Werners

Foundation to E.I.S. and from The Royal Physiographic Society of Lund, the Jörgen Lindströms

Fund and "Lunds Djurskyddsfond" to B.W.

Data Availability

Novel sequence data will be made available on the National Centre for Biotechnology

Information (https://www.ncbi.nlm.nih.gov/genbank/). All raw data, command files and

scripts are available at GitHub

(https://rachelblow.github.io/IschnuraPolymorphismEvolutionaryHistory/) and Dryad Digital

Repository (https://datadryad.org/stash) including convergence and stationarity checks of

MCMC analyses.

Competing Interests

628   The authors declare that they have no competing interests.

References

Andrade, P., Pinho, C., Lanuza, G.P. i de, Afonso, S., Brejcha, J., Rubin, C.-J., Wallerman, O., Pereira, P., Sabatino, S.J., Bellati, A., Pellitteri-Rosa, D., Bosakova, Z., Bunikis, I., Carretero, M.A., Feiner, N., Marsik, P., Paupério, F., Salvi, D., Soler, L., While, G.M., Uller, T., Font, E., Andersson, L., Carneiro, M., 2019. Regulatory changes in pterin and carotenoid genes underlie balanced color polymorphisms in the wall lizard. Proc. Natl. Acad. Sci., USA 116, 5633–5642. https://doi.org/10.1073/pnas.1820320116

Azevedo, L., Serrano, C., Amorim, A., Cooper, D.N., 2015. Trans-species polymorphism in humans and the great apes is generally maintained by balancing selection that modulates the host immune response. Hum. Genomics 9, 21. https://doi.org/10.1186/s40246-015-0043-1

Beatty, C.D., Herrera, M.S., Skevington, J.H., Rashed, A., Gossum, H.V., Kelso, S., Sherratt, T.N., 2017. Biogeography and systematics of endemic island damselflies: The Nesobasis and Melanesobasis (Odonata: Zygoptera) of Fiji. Ecol. Evol. 7, 7117–7129. https://doi.org/10.1002/ece3.3175

Bechly, G., 2000. A new fossil damselfly species (Insecta: Odonata: Zygoptera: Coenagrionidae: Ischnurinae) from Dominican Amber. Stuttgarter Beitr. Naturk. Ser. B. 299, 1–9.

Blows, M.W., Brooks, R., Kraft, P.G., 2003. Exploring complex fitness surfaces: multiple ornamentation and polymorphism in male guppies. Evolution 57, 1622–1630.

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M.A., Rambaut, A., Drummond, A.J., 2014. BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. PLOS Comput. Biol. 10, e1003537. https://doi.org/10.1371/journal.pcbi.1003537

Bouckaert, R.R., Drummond, A.J., 2017. bModelTest: Bayesian phylogenetic site model

averaging and model comparison. BMC Evol. Biol. 17, 42.

https://doi.org/10.1186/s12862-017-0890-6

Brower, A.V., 1994. Rapid morphological radiation and convergence among races of the

butterfly Heliconius erato inferred from patterns of mitochondrial DNA evolution. Proc.

Natl. Acad. Sci., USA 91, 6491–6495.

Callahan, M.S., McPeek, M.A., 2016. Multi-locus phylogeny and divergence time estimates of

Enallagma damselflies (Odonata: Coenagrionidae). Mol. Phylogenet. Evol. 94, 182–

195. https://doi.org/10.1016/j.ympev.2015.08.013

Chippindale, P.T., Dave, V.K., Whitmore, D.H., Robinson, J.V., 1999. Phylogenetic

relationships of North American damselflies of the genus Ischnura (Odonata:

Zygoptera: Coenagrionidae) based on sequences of three mitochondrial genes. Mol.

Phylogenet. Evol. 11, 110–121.

Cordero, A., 1992. Density-dependent mating success and colour polymorphism in females of

the damselfly Ischnura graellsii (Odonata: Coenagrionidae). J. Anim. Ecol. 61, 769–

780.

Cordero, A., Carbone, S.S.., Utzeri, C., 1998. Mating opportunities and mating costs are

reduced in androchrome female damselflies, Ischnura elegans. Anim. Behav. 55, 185–

187.

Cordoba-Aguilar, A., 2008. Dragonflies and damselflies: model organisms for ecological and

evolutionary research. Oxford University Press, Oxford.

Corl, A., Davis, A.R., Kuchta, S.R., Sinervo, B., 2010. Selective loss of polymorphic mating

types is associated with rapid phenotypic evolution during morphic speciation. Proc.

Natl. Acad. Sci., USA 107, 4254–4259. https://doi.org/10.1073/pnas.0909480107

Crow, J.F., Kimura, M., 1970. An Introduction to population genetic theory. Harper & Row.,

New York.

Cunningham, C.W., Omland, K.E., Oakley, T.H., 1998. Reconstructing ancestral character

states: a critical reappraisal. Trends Ecol. Evol. 13, 361–366.

Damm, S., Dijkstra, K.D., Hadrys, H., 2010. Red drifters and dark residents: the phylogeny and

ecology of a Plio-Pleistocene dragonfly radiation reflects Africa's changing

environment (Odonata, Libellulidae, Trithemis). Mol. Phylogenet. Evol. 54, 870–882.

https://doi.org/10.1016/j.ympev.2009.12.006

Dijkstra, K.-D.B., Kalkman, V.J., Dow, R.A., Stokvis, F.R., Van Tol, J., 2014. Redefining the

damselfly families: a comprehensive molecular phylogeny of Zygoptera (Odonata).

Syst. Entomol. 39, 68–96. https://doi.org/10.1111/syen.12035

Dobzhansky, T., 1970. Genetics of the evolutionary process. Columbia University Press, New

york.

Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

Fincke, O.M., 2004. Polymorphic signals of harassed female odonates and the males that learn

them support a novel frequency-dependent model. Anim. Behav. 67, 833–845.

Fincke, O.M., Jödicke, R., Paulson, D., Schultz, D.T., 2005. The evolution and frequency of

female color morphs in Holarctic Odonata: why are male-like morphs typically the

minority? Int. J. Odonatol. 8, 183–212.

Fisher, R.A., 1930. The Genetical Theory of Natural Selection. Clarendon Press, Oxford.

Ford, E.B., 1975. Ecological genetics. Chapman and Hall, London.

Formica, V.A., Tuttle, E.M., 2009. Examining the social landscapes of alternative

reproductive strategies. J. Evol. Biol. 22, 2395–2408. https://doi.org/10.1111/j.1420-

9101.2009.01855.x

Forsman, A., 2016. Is colour polymorphism advantageous to populations and species? Mol.

Ecol. https://doi.org/10.1111/mec.13629

Gangoso, L., Grande, J.M., Ducrest, A.-L., Figuerola, J., Bortolotti, G.R., Andrés, J.A., Roulin,

A., 2011. MC1R-dependent, melanin-based colour polymorphism is associated with

cell-mediated response in the Eleonora's falcon. J. Evol. Biol. 24, 2055–2063.

https://doi.org/10.1111/j.1420-9101.2011.02336.x

Gavrilets, S., Waxman, D., 2002. Sympatric speciation by sexual conflict. Proc. Natl. Acad.

Sci., USA 99, 10533–10538.

Gentilini, G., Bagli, L., 2004. Fossil Zygoptera and Anisoptera from the Upper Miocene of

Monte Castellaro (Pesaroo Marches, Central Italy). Quad. Studi Nat. Romagna 19, 17–

44.

Gering, E.J., 2017. Male-mimicking females increase male-male interactions, and decrease

male survival and condition in a female-polymorphic damselfly. Evolution 71, 1390–

1396. https://doi.org/10.1111/evo.13221

Global Administrative Areas, 2018. GADM database of Global Administrative Areas, version

3.6. [online] URL: www.gadm.org.

Goldberg, E.E., Kohn, J.R., Lande, R., Robertson, K.A., Smith, S.A., Igić, B., 2010. Species

Selection Maintains Self-Incompatibility. Science 330, 493–495.

https://doi.org/10.1126/science.1194513

Gordon, S.P., Kokko, H., Rojas, B., Nokelainen, O., Mappes, J., 2015. Colour polymorphism

torn apart by opposing positive frequency-dependent selection, yet maintained in space.

J. Anim. Ecol. 84, 1555–1564. https://doi.org/10.1111/1365-2656.12416

Gosden, T.P., Svensson, E.I., 2009. Density-dependent male mating harassment, female

resistance, and male mimicry. Am. Nat. 173, 709–721. https://doi.org/10.1086/598491

Hadfield, J.D., 2015. Increasing the efficiency of MCMC for hierarchical phylogenetic models

of categorical traits using reduced mixed models. Methods Ecol. Evol. 6, 706–714.

https://doi.org/10.1111/2041-210X.12354

Hadfield, J.D., 2010. MCMC Methods for Multi-Response Generalized Linear Mixed

Models: The MCMCglmm R Package. J. Stat. Softw. 33, 1–22.

https://doi.org/10.18637/jss.v033.i02

Hadfield, J.D., Nakagawa, S., 2010. General quantitative genetic methods for comparative

biology: phylogenies, taxonomies and multi-trait models for continuous and

categorical characters. J. Evol. Biol. 23, 494–508. https://doi.org/10.1111/j.1420-

9101.2009.01915.x

Haldane, J.B.S., 1949. Disease and evolution. Ric. Sci. Suppl. A 19, 68–76.

Haldane, J.B.S., 1932. The causes of evolution. Longmans, Green & Co., London.

Hartl, D.L., Clark, A.G., 1997. Principles of population genetics. Sinauer Associates, Inc.,

Sunderland, MA.

Hayashi, T.I., Vose, M., Gavrilets, S., 2007. Genetic differentiation by sexual conflict.

Evolution 61, 516–529.

Heath, T.A., Hedtke, S.M., Hillis*, D.M., 2008. Taxon sampling and the accuracy of

phylogenetic analyses. J. Syst. Evol. 46, 239–257.

https://doi.org/10.3724/SP.J.1002.2008.08016

Hugall, A.F., Stuart-Fox, D., 2012. Accelerated speciation in colour-polymorphic birds. Nature

485, 631–634. https://doi.org/10.1038/nature11050

Hughes, K.A., Houde, A.E., Price, A.C., Rodd, F.H., 2013. Mating advantage for rare males in

wild guppy populations. Nature 503, 108–110. https://doi.org/10.1038/nature12717

Iserbyt, A., Bots, J., Van Dongen, S., Ting, J.J., Van Gossum, H., Sherratt, T.N., 2011.

Frequency-dependent variation in mimetic fidelity in an intraspecific mimicry system.

Proc. R. Soc. B-Biol. Sci. 278, 3116–3122. https://doi.org/10.1098/rspb.2011.0126

Iserbyt, A., Bots, J., Van, G.H., Sherratt, T.N., 2013. Negative frequency-dependent selection

or alternative reproductive tactics: maintenance of female polymorphism in natural

populations. BMC Evol. Biol. 13, 139. https://doi.org/10.1186/1471-2148-13-139

Iversen, L.L., Svensson, E.I., Christensen, S.T., Bergsten, J., Sand-Jensen, K., 2019. Sexual

conflict and intrasexual polymorphism promote assortative mating and halt population

differentiation. Proc. R. Soc. B-Biol. Sci. 286, 20190251.

https://doi.org/10.1098/rspb.2019.0251

Jamie, G.A., Meier, J.I., 2020. The Persistence of Polymorphisms across Species Radiations.

Trends Ecol. Evol. https://doi.org/10.1016/j.tree.2020.04.007

Karlsson, G.K., Kovalev, A., Svensson, E.I., Gorb, S.N., 2013. Male clasping ability, female

polymorphism and sexual conflict: fine-scale elytral morphology as a sexually

antagonistic adaptation in female diving beetles. J. R. Soc. Interface 10, 20130409.

https://doi.org/10.1098/rsif.2013.0409

Klein, J., 1987. Origin of major histocompatibility complex polymorphism: The trans-species

hypothesis.   Hum.   Immunol.   19,   155–162.   https://doi.org/10.1016/0198-

8859(87)90066-8

Lank, D.B., Smith, C.M., Hanotte, O., Burke, T., Cooke, F., 1995. Genetic Polymorphism for

Alternative Mating-Behavior in Lekking Male Ruff Philomachus-Pugnax. Nature 378,

59–62.

Le Rouzic, A., Hansen, T.F., Gosden, T.P., Svensson, E.I., 2015. Evolutionary time-series

analysis reveals the signature of frequency-dependent selection on a female mating

polymorphism. Am. Nat. 185E, E182–E196.

Lee, J.-W., Kim, H.-N., Yoo, S., Yoo, J.-C., 2019. Common cuckoo females may escape male

sexual   harassment   by   color   polymorphism.   Sci.   Rep.   9,   1–9.

https://doi.org/10.1038/s41598-019-44024-6

Letsch, H., Gottsberger, B., Ware, J.L., 2016. Not going with the flow: a comprehensive time-

calibrated phylogeny of dragonflies (Anisoptera: Odonata: Insecta) provides evidence

for the role of lentic habitats on diversification. Mol. Ecol. 25, 1340–1353.

https://doi.org/10.1111/mec.13562

Llaurens, V., Whibley, A., Joron, M., 2017. Genetic architecture and balancing selection: the

life   and   death   of   differentiated   variants.   Mol.   Ecol.   26,   2430–2448.

https://doi.org/10.1111/mec.14051

Losos, J.B., 2011. Seeing the Forest for the Trees: The Limitations of Phylogenies in

Comparative Biology. Am. Nat. 177, 709–727. https://doi.org/10.1086/660020

Löytynoja, A., Goldman, N., 2008. Phylogeny-Aware Gap Placement Prevents Errors in

Sequence   Alignment   and   Evolutionary   Analysis.   Science   320,   1632–1635.

https://doi.org/10.1126/science.1158395

Löytynoja, A., Goldman, N., 2005. An algorithm for progressive multiple alignment of

sequences   with   insertions.   Proc.   Natl.   Acad.   Sci.,   USA.   102,   10557–10562.

https://doi.org/10.1073/pnas.0409137102

Maddison, W.P., 1997. Gene Trees in Species Trees. Syst. Biol. 46, 523–536.

https://doi.org/10.1093/sysbio/46.3.523

Marinov, M., Schmaedick, M., Polhemus, D., Stirne, R.L., 2015. Faunistic and taxonomic

investigations on the Odonata fauna of the Samoan archipelago with particular focus on

taxonomic ambiguities in the "Ischnurine complex. J. Int. Dragonfly Fund 91, 1–56.

Mattern, M., Van Gossum, H., 2008. A phylogenetic perspective on absence and presence of a

sex-limited polymorphism. Anim. Biol. 58, 257–273.

https://doi.org/10.1163/157075608X328099

McKinnon, J.S., Pierotti, M.E., 2010. Colour polymorphism and correlated characters: genetic

mechanisms and evolution. Mol. Ecol. 19, 5101–5125. https://doi.org/10.1111/j.1365-

294X.2010.04846.x

McLean, C.A., Stuart-Fox, D., 2014. Geographic variation in animal colour polymorphisms

and its role in speciation. Biol. Rev. 89, 860–873. https://doi.org/10.1111/brv.12083

McPeek, M.A., Gavrilets, S., 2006. The evolution of female mating preferences: Differentiation

from species with promiscuous males can promote speciation. Evolution 60, 1967–

1980.

Miller, M.N., Fincke, O.M., 1999. Cues for mate recognition and the effect of prior experience

on mate recognition in Enallagma damselflies. J. Insect Behav. 12, 801–814.

Mitchell-Olds, T., Willis, J.H., Goldstein, D.B., 2007. Which evolutionary processes influence

natural genetic variation for phenotypic traits? Nat. Rev. Genet. 8, 845–856.

https://doi.org/10.1038/nrg2207

Moon, R.M., Kamath, A., 2019. Re-examining escape behaviour and habitat use as correlates

of dorsal pattern variation in female brown anole lizards, Anolis sagrei (Squamata:

Dactyloidae). Biol. J. Linn. Soc. 126, 783–795.

https://doi.org/10.1093/biolinnean/blz006

Neff, B.D., Svensson, E.I., 2013. Polyandry and alternative mating tactics. Philos. Trans. R.

Soc. Lond. Biol. Sci. 368, 20120045. https://doi.org/10.1098/rstb.2012.0045

O'Connor, E.A., Cornwallis, C.K., Hasselquist, D., Nilsson, J.-Å., Westerdahl, H., 2018. The

evolution of immunity in relation to colonization and migration. Nat. Ecol. Evol. 2, 841–

849. https://doi.org/10.1038/s41559-018-0509-3

Ogilvie, H.A., Bouckaert, R.R., Drummond, A.J., 2017. StarBEAST2 Brings Faster Species

825   Tree Inference and Accurate Estimates of Substitution Rates. Mol. Biol. Evol. 34, 2101–

826   2114. https://doi.org/10.1093/molbev/msx126

827   Okude, G., Fukatsu, T., Futahashi, R., 2020. Interspecific crossing between blue-tailed

828   damselflies Ischnura elegans and I. senegalensis in the laboratory. Entomol. Sci. 23,

829   165–172. https://doi.org/10.1111/ens.12408

830   Pagel, M., Meade, A., Barker, D., 2004. Bayesian Estimation of Ancestral Character States on

831   Phylogenies. Syst. Biol. 53, 673–684. https://doi.org/10.1080/10635150490522232

832   Paradis, E., Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics and

833   evolutionary analyses in R. Bioinformatics 35, 526–528.

834   https://doi.org/10.1093/bioinformatics/bty633

835   Paulson, D., 2012. Dragonflies and Damselflies of the East. Princeton University Press, NJ.

836   Paulson, D., 2009. Dragonflies and Damselflies of the West, Princeton Field Guides. Princeton

837   University Press, Princeton, N.J.

838   Plummer, M., Best, N., Cowles, K., Vines, K., 2006. CODA: convergence diagnosis and output

839   analysis for MCMC. R News 6, 7–11.

840   Provine, W.B., 1986. Sewall Wright and evolutionary biology. University of Chicago Press,

841   Chicago.

842   Reinhardt, K., Harney, E., Naylor, R., Gorb, S., Siva-Jothy, M.T., 2007. Female-Limited

843   Polymorphism in the Copulatory Organ of a Traumatically Inseminating Insect. The

844   Am. Nat. 170, 931–935. https://doi.org/10.1086/522844

845   Revell, L.J., 2013. Two new graphical methods for mapping trait evolution on phylogenies.

846   Methods Ecol. Evol. 4, 754–759. https://doi.org/10.1111/2041-210X.12066

847   Richman, A.D., Kohn, J.R., 2000. Self-incompatability alleles from Physalis : Implications for

848   historical inference from balanced genetic polymorphisms. Proc. Natl. Acad. Sci., USA

849   96, 168–172.

850   Richman, A.D., Uyenoyama, M.C., Kohn, J.R., 1996. Allelic diversity and gene genealogy at

851   the self-incompatability locus in solanaceae. Science 273, 1212–1216.

852   Robertson, H.M., 1985. Female dimorphism and mating behaviour in a damselfly, Ischnura

853   ramburi: females mimicking males. Anim. Behav. 33, 805–809.

854   Robinson, J.V., Allgeyer, R., 1996. Covariation in life-history traits, demographics and

855   behaviour in ischnuran damselflies: The evolution of monandry. Biol. J. Linn. Soc. 58,

856   85–98.

857   Ross, L., Hardy, N.B., Okusu, A., Normark, B.B., 2013. Large Population Size Predicts the

858   Distribution of Asexuality in Scale Insects. Evolution 67, 196–206.

859   https://doi.org/10.1111/j.1558-5646.2012.01784.x

860   Sanchez-Guillen, R.A., Hansson, B., Wellenreuther, M., Svensson, E.I., Cordero-Rivera, A.,

861   2011. The influence of stochastic and selective forces in the population divergence of

862   female colour polymorphism in damselflies of the genus Ischnura. Heredity (Edinb.)

863   107, 513–522. https://doi.org/10.1038/hdy.2011.36

864   Sanchez-Guillen, R.A., Van Gossum, H., Rivera, A.C., 2005. Hybridization and the inheritance

865   of female colour polymorphism in two ischnurid damselflies (Odonata :

866   Coenagrionidae). Biol. J. Linn.Soc. 85, 471–481.

867   Sánchez-Guillén, R.A., Wellenreuther, M., Cordero-Rivera, A., Hansson, B., 2011.

868   Introgression and rapid species turnover in sympatric damselflies. BMC Evol. Biol. 11,

869   210. https://doi.org/10.1186/1471-2148-11-210

870   Schemske, D.W., Bierzychudek, P., 2001. Perspective: Evolution of flower color in the desert

871   annual Linanthus parryae: Wright revisited. Evolution 55, 1269–1282.

Schluter, D., Price, T., Mooers, A.O., Ludwig, D., 1997. Likelihood of ancestor states in
adaptive radiation. Evolution 51, 1699–1711.

Schwander, T., Leimar, O., 2011. Genes as leaders and followers in evolution. Trends Ecol.
Evol. 26, 143–151. https://doi.org/10.1016/j.tree.2010.12.010

Sinervo, B., Lively, C.M., 1996. The rock-paper-scissors game and the evolution of alternative
male strategies. Nature 380, 240–243.

Sinervo, B., Svensson, E., 2002. Correlational selection and the evolution of genomic
architecture. Heredity 16, 948–955.

Sirot, Laura K., Brockmann, H.J., 2001. Costs of sexual interactions to females in rambur's
forktail damselfly, Ischnura ramburi (Zygoptera: Coenagrionidae). Anim Behav. 61,
415–424. https://doi.org/10.1006/anbe.2000.1605

Sirot, L.K., Brockmann, H.J., 2001. Costs of sexual interaction to females in Rambur's forktail
damselfly, Ischnura ramburi (Zygoptera: Odonata). Anim. Behav. 61, 415–424.

Sirot, L.K., Brockmann, H.J., Marinis, C., Muschett, G., 2003. Maintenance of a female-limited
polymorphism in Ischnura ramburi (Zygoptera : Coenagrionidae). Anim. Behav. 66,
763–775.

Svensson, E.I., 2017. Back to basics: using colour polymorphisms to study evolutionary
processes. Mol. Ecol. 26, 2204–2211. https://doi.org/10.1111/mec.14025

Svensson, E.I., Abbott, J., Hardling, R., 2005. Female polymorphism, frequency dependence,
and rapid evolutionary dynamics in natural populations. Am. Nat. 165, 567–576.

Svensson, E.I., Abbott, J.K., Gosden, T.P., Coreau, A., 2009. Female polymorphisms, sexual
conflict and limits to speciation processes in animals. Evol. Ecol. 23, 93–108.

Svensson, E.I., Willink, B., Duryea, M.C., Lancaster, L.T., 2020. Temperature drives pre-
reproductive selection and shapes the biogeography of a female polymorphism. Ecol.
Lett. 23, 149–159. https://doi.org/10.1111/ele.13417

Swaegers, J., Janssens, S.B., Ferreira, S., Watts, P.C., Mergeay, J., McPeek, M.A., Stoks, R.,

2014. Ecological and evolutionary drivers of range size in Coenagrion damselflies. J.

Evol. Biol. 27, 2386–2395. https://doi.org/10.1111/jeb.12481

Takahashi, M., Takahashi, Y., Kawata, M., 2018. Candidate genes associated with color morphs

of female-limited polymorphisms of the damselfly Ischnura senegalensis. Heredity 122,

81–92. https://doi.org/10.1038/s41437-018-0076-z

Takahashi, Y., Kagawa, K., Svensson, E.I., Kawata, M., 2014. Evolution of increased

phenotypic diversity enhances population performance by reducing sexual harassment

in damselflies. Nat. Commun. 5, 4468. https://doi.org/10.1038/ncomms5468

Takahashi, Y., Noriyuki, S., 2019. Colour polymorphism influences species' range and

extinction risk. Biol. Lett. 15, 20190228. https://doi.org/10.1098/rsbl.2019.0228

Takahashi, Y., Tanaka, R., Yamamoto, D., Noriyuki, S., Kawata, M., 2018. Balanced genetic

diversity improves population fitness. Proc. R. Soc. B-Biol. Sci. 285, 20172045.

https://doi.org/10.1098/rspb.2017.2045

Takahashi, Y., Watanabe, M., 2010. Female reproductive success is affected by selective male

harassment in the damselfly Ischnura senegalensis. Anim Behav. 79, 211–216.

https://doi.org/10.1016/j.anbehav.2009.10.032

Takahashi, Y., Yoshimura, J., Morita, S., Watanabe, M., 2010. Negative frequency-dependent

selection in female color polymorphism of a damselfly. Evolution 64, 3620–3628.

https://doi.org/10.1111/j.1558-5646.2010.01083.x

Torres-Cambas, Y., Ferreira, S., Cordero-Rivera, A., Lorenzo-Carballa, M.O., 2019.

Mechanisms of allopatric speciation in an Antillean damselfly genus (Odonata,

Zygoptera): Vicariance or long-distance dispersal? Mol. Phylogenet. Evol. 137, 14–21.

https://doi.org/10.1016/j.ympev.2019.04.018

Turelli, M., Schemske, D.W., Bierzychudek, P., 2001. Stable two-allele polymorphisms
maintained by fluctuating fitnesses and seed banks: Protecting the blues in Linanthus
parryae. Evolution 55, 1283–1298.

Turgeon, J., Stoks, R., Thum, R.A., Brown, J.M., McPeek, M.A., 2005. Simultaneous
Quaternary radiations of three damselfly clades across the Holarctic. Am. Nat. 165, E78-
107. https://doi.org/10.1086/428682

Tuttle, E.M., Bergland, A.O., Korody, M.L., Brewer, M.S., Newhouse, D.J., Minx, P., Stager,
M., Betuel, A., Cheivron, Z.A., Warren, W.C., Gonser, R.A., Balakrishnan, C.N., 2016.
Divergence and functional degradation of a sex chromosome-like supergene. Curr. Biol.
26, 344–350.

Van Gossum, H., Stoks, R., De Bruyn, L., 2001. Reversible frequency-dependent switches in
male mate choice. Proc. R.Soc. Lond., B. 268, 83–85.

Vega-Sánchez, Y.M., Mendoza-Cuenca, L.F., González-Rodríguez, A., 2019. Complex
evolutionary history of the American Rubyspot damselfly, Hetaerina americana
(Odonata): Evidence of cryptic speciation. Mol. Phylogenet. Evol. 139, 106536.
https://doi.org/10.1016/j.ympev.2019.106536

Villemereuil, P. de, Gimenez, O., Doligez, B., 2013. Comparing parent–offspring regression
with frequentist and Bayesian animal models to estimate heritability in wild
populations: a simulation study for Gaussian and binary traits. Methods Ecol. Evol. 4,
260–275. https://doi.org/10.1111/2041-210X.12011

Waller, J.T., Svensson, E.I., 2017. Body size evolution in an old insect order: No evidence for
Cope's Rule in spite of fitness benefits of large size. Evolution 71, 2178–2193.
https://doi.org/10.1111/evo.13302

Wellenreuther, M., Svensson, E.I., Hansson, B., 2014. Sexual selection and genetic colour

polymorphisms in animals. Mol. Ecol. 23, 5398–5414.

https://doi.org/10.1111/mec.12935

West-Eberhard, M.J., 1986. Alternative adaptations, speciation, and phylogeny (A Review).

Proc. Natl. Acad. Sci, USA 83, 1388–1392. https://doi.org/10.1073/pnas.83.5.1388

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis, 2nd ed, Use R! Springer

International Publishing. https://doi.org/10.1007/978-3-319-24277-4

Willink, B., 2018. The descent of damselflies and variation in relation to sex, PhD-thesis. Lund

University (Sweden).

Willink, B., Duryea, M.C., Svensson, E.I., 2019. Macroevolutionary Origin and Adaptive

Function of a Polymorphic Female Signal Involved in Sexual Conflict. Am. Nat. 194,

707–724. https://doi.org/10.1086/705294

Willink, B., Duryea, M.C., Wheat, C., Svensson, E.I., 2020. Changes in gene expression during

female reproductive development in a color polymorphic insect. Evolution

https://doi.org/10.1111/evo.13979

Woronik, A., Tunström, K., Perry, M.W., Neethiraj, R., Stefanescu, C., Celorio-Mancera, M.

de la P., Brattström, O., Hill, J., Lehmann, P., Käkelä, R., Wheat, C.W., 2019. A

transposable element insertion is associated with an alternative life history strategy. Nat

Commun 10, 1–11. https://doi.org/10.1038/s41467-019-13596-2

Wright, A.M., Hillis, D.M., 2014. Bayesian Analysis Using a Simple Likelihood Model

Outperforms Parsimony for Estimation of Phylogeny from Discrete Morphological

Data. PLOS ONE 9, e109210. https://doi.org/10.1371/journal.pone.0109210

Wright, S., 1932. The roles of mutation, inbreeding, crossbreeding and selection in evolution.

Proc. Sixth Internat. Congr. Gen. 1, 356–366.

Wright, S., 1931. Evolution in Mendelian Populations. Genetics 16, 97–159.

Xu, M., Cerreta, A.L., Schultz, T.D., Fincke, O.M., 2014. Selective use of multiple cues by

males reflects a decision rule for sex discrimination in a sexually mimetic damselfly.

Anim. Behav. 92, 9–18. https://doi.org/10.1016/j.anbehav.2014.03.016

Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.-Y., 2017. ggtree: an r package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. Methods Ecol. Evol- 8, 28–36. https://doi.org/10.1111/2041-

210X.12628

Figure Legends

Figure 1. Some representative species of the damselfly genus *Ischnura*, from both the Old and the New World, exemplifying the extensive phenotypic and colour variation within and between species in this genus. All species in this panel are included in the molecular phylogeny we present in this paper. **A.** An immature female *Ischnura capreolus*, revealing the yellow colour females of this species exhibit prior to sexual maturation (Brazil). **B.** A sexually mature female of *I. graellsii* of the infuscans-morph (Portugal). **C.** Sexually mature male *I. graellsii* (Portugal). **D.** Sexually mature male *I. ramburii* (Texas, USA). **E-G:** Females in several both closely and distantly related species show an orange colour phase in females prior to sexual maturation. **E.** Immature female *I. ramburii* (Texas, USA). **F.** Immature female *I. pumilio* (Sweden). **G.** Immature female *I. senegalensis* (Japan). **H.** Mature male *I. senegalensis* (Japan). **I-K:** In addition to ontogenetic colour variation in females, some species of *Ischnura* also show heritable female colour polymorphism at the adult stage. Here are the three different female colour morphs in the trimorphic species *I. elegan*s (all photos from Sweden). **I.** Male *I. elegans* mating with an androchrome (A-female; male mimic) female morph. **J.** Male *I. elegans* mating with the *infuscans-obsoleta* (O-female) morph. **K.** Male *I. elegans* mating with the *infuscans* (I-female) morph. All photographs taken by E.I. Svensson.

Figure 2. Time calibrated *Ischnura* phylogeny of 41 species of damselflies from the genus *Ischnura* as estimated in StarBEAST2. Time axis represents millions of years from the present and grey bars at nodes indicate 95% HPD intervals for age estimates. Values at nodes indicate posterior probability of internal nodes.

1015 **Figure 3.** Posterior divergence time estimation at the root node sampling from the prior only

1016 (gold) and sampling with data (grey). Age estimation followed an exponentially distributed

1017 prior with an offset of 13.8 my to prevent improbable age estimation (see Methods).

1018

1019 **Figure 4.** Ancestral state reconstruction across the *Ischnura* phylogeny. Extant morph states

1020 are represented by circles at tips and reconstructed ancestral states are represented by piecharts

1021 at nodes (female-heterochromatic only, green; female-andromorphic only, blue; female-

1022 dimorphic, red; female-trimorphic, pink). Piecharts show the mean posterior probability for

1023 each morph state and are plotted on the MCC tree.

1024

Figure 5. Probability of species having female polymorphism in relation to the species

geographic range size, estimated in millions of squared kilometres. The left panel shows the

MMC tree for the genus *Ischnura*. Bars on the centre panel are proportional to the geographic

range of each species, except for species with very small ranges ($<10^5$ Km^2) which are all

depicted by a similarly thin line. Light grey bars represent female-monomorphic (M) species

and dark grey bars correspond to female-polymorphic (P) species. On the right panel, each

circle represents an extant species. The black line shows the prediction from a Bayesian

phylogenetic mixed-effect model and the grey area shows the 95% credible interval of the

prediction. Species with larger geographic ranges are more likely to have polymorphic females

(see Results).

Female Morph Status

- Female−monomorphic with androchrome females
- Female−dimorphic
- Female−monomorphic with heterochrome females
- Female−trimorphic

Ischnura verticalis
Ischnura posita
Ischnura perparva
Ischnura cervula
Ischnura demorsa
Ischnura damula
Ischnura erratica
Ischnura gemina
Ischnura denticollis
Ischnura kellicotti
Ischnura prognata
Ischnura hastata
Ischnura ultima
Ischnura cruzi
Ischnura ezoin
Ischnura asiatica
Ischnura pumilio
Ischnura intermedia
Ischnura senegalensis
Ischnura foylei
Ischnura heterosticta
Ischnura ramburii
Ischnura nursei
Ischnura evansi
Ischnura graellsii
Ischnura elegans
Ischnura saharensis
Ischnura fountaineae
Ischnura fluviatilis
Ischnura rufostigma
Ischnura barberi
Ischnura capreolus
Ischnura taitensis
Ischnura cardinalis
Ischnura pamelae
Ischnura aurora
Ischnura pruinescens
Pacificagrion sp.
Pacificagrion lachrymosa
Ischnura sp.
Ischnura armstrongi