Cell adhesion molecule IGPR-1 activates AMPK connecting cell adhesion to energy sensing and autophagy

Razie Amraie, Tooba Alwani, Rachel Xi-Yeen Ho, Zahra Aryan, Shawn Wang, Nader Rahimi[1]

Department of Pathology, School of Medicine, Boston University Medical Campus, Boston, MA 02118, USA.

Running title: IGPR-1 is activated by and regulates autophagy

[1]**Corresponding Author**:
Nader Rahimi
Departments of Pathology
Boston University Medical Campus
670 Albany St., Room 510, Boston, MA 02118
Tel: 617-638-5011
Fax: 617-414-7914
E-mail: nrahimi@bu.edu.

ACKNOWLEDGMENTS AND FUNDING:
Funding: This work was supported in part through grants from the NIH/NCI (R21CA191970, R21CA193958 and CTSI grant 1UL1TR001430 to N.R.).

CONFLICTS OF INTEREST: The authors declare no conflicts of interest

ABSTRACT:

Cellular stress caused by loss of extracellular matrix (ECM)-cell interaction involving integrins induces autophagy and autophagy is considered to protect cells from anoikis (i.e., cell death due to loss of cell adhesion). Immunoglobulin (Ig) and proline-rich receptor-1 (IGPR-1) is a newly identified cell adhesion molecule that mediates cell-cell adhesion and supports cell survival in response to anoikis. However, whether IGPR-1 is activated in response to and mediates autophagy is not known. In this study, we demonstrate that IGPR-1 is activated by autophagy inducing stimuli, such as nutrient deprivation, rapamycin and lipopolysaccharide (LPS). We have identified IκB kinaseβ (IKKβ) as a key serine/threonine kinase activated by autophagy stimuli and mediates phosphorylation of IGPR-1 at Ser220. Activation of IGPR-1, in turn, stimulates phosphorylation of AMP-activated protein kinase (AMPK), which leads to phosphorylation of key pro-autophagy proteins, ULK1 and Beclin-1, increased LC3-II levels and accumulation of LC3 punctum. This study demonstrates that IGPR-1 is activated by and regulates autophagy, connecting cell adhesion to autophagy, a finding that has important significance for autophagy-driven pathologies such cardiovascular diseases and cancer.

Introduction:

Autophagy (also called macroautophagy), the lysosomal degradation of cytoplasmic organelles or cytosolic components, is an evolutionarily conserved cytoprotective mechanism that is induced in response to cellular stress, such as nutrient withdrawal, loss of cell adhesion, and flow shear stress, or by therapeutic genotoxic agents and others[1, 2]. Cell-extracellular matrix (ECM) adhesion through engagement of cell adhesion molecules (CAMs) (*e.g.*, integrins) is thought to regulate autophagy[3, 4] and more importantly loss of cell adhesion to ECM is linked to induction of autophagy[4, 5].

Upon autophagy, unc-51-like kinase 1 (ULK1 also known as ATG1) associates with autophagy-related protein 13 (ATG13), and focal adhesion kinase family interacting protein of 200 kD (FIP200) to form the ULK1 complex. ULK1 interaction with ATG13 and FIP200 is critical for ULK1 kinase activity and stability[6]. The ULK1 complex translocates to autophagy initiation sites and recruits the class III phosphatidylinositol 3-kinase, vacuolar protein sorting 34 (VPS34) complex consisting of BECLIN-1 (the mammalian orthologue of the yeast autophagy protein Apg6/Vps30[7] and multiple other ATGs leading to the phagophore formation[8]. The serine/threonine protein kinase mTOR (mechanistic target of rapamycin) complex 1 (mTORC1) is a key regulator of autophagy in response to nutrient availability. In the presence of amino acids, mTORC1 is activated and suppresses autophagy through phosphorylation of ULK1 and ATG13. However, upon nutrient deprivation, mTORC1 activity is inhibited leading to the activation of ULK1 that induces the autophagy program[9, 10]. Suppression of mTORC1 activity by AMP-activated protein kinase (AMPK) is central to regulation of autophagy. AMPK inactivates mTORC1 through phosphorylation of RAPTOR, a key protein present within the mTORC1 and more importantly directly phosphorylates ULK1 at multiple serine residues and activates it[11, 12].

Commonly known autophagy inducing conditions or agents such as nutrient withdrawal, immunosuppressant rapamycin/Sirolimus, and lipopolysaccharide (LPS) all activate several key kinases,

including the IκB kinase (IKK) complex[13]. IKK complex is composed of at least three proteins, including two catalytic subunits (IKKα and IKKβ) and the scaffold protein NF-κB essential modulator (NEMO; also called IKKγ)[14]. In addition to its pivotal role in mediating phosphorylation of IκB[14], IKKβ also regulates autophagy in IκB-independent manner[15, 16] by mechanisms that are not fully understood.

Immunoglobulin (Ig) and proline-rich receptor-1 (IGPR-1) was recently identified as a novel cell adhesion molecule expressed in various human cell types including, endothelial and epithelial cells and mediates cell-cell adhesion[17]. IGPR-1 regulates angiogenesis, endothelial barrier function[17, 18], decreases sensitivity of tumor cells to genotoxic agent, doxorubicin, and supports tumor cell survival in response to anoikis[19]. IGPR-1 is localized to adherens junctions and, is activated through trans-homophilic dimerization[18]. Additionally, IGPR-1 responds to various cellular stresses, as its phosphorylation (*i.e.*, Ser220) is significantly increased by flow shear stress[20] and doxorubicin[19, 20]. Curiously, both shear stress[21] and doxorubicin[22, 23] are well-known potent inducer of autophagy, raising a possibility for the involvement of IGPR-1 in autophagy. In this study, we demonstrate that upon induction of autophagy, IGPR-1 is phosphorylated at Ser220 via a mechanism that involves activation of IKKβ. IKKβ-dependent phosphorylation of IGPR-1 stimulates phosphorylation of AMPK leading to activation of BECN1 and ULK1 connecting cell adhesion and energy sensing to autophagy.

Results:

IKKβ is activated by autophagy and phosphorylates IGPR-1

Homophilic transdimerization induces phosphorylation of IGPR-1 at Ser220[24]. Moreover, genotoxic agent, doxorubicin[19] and flow shear stress[20] also simulate phosphorylation of IGPR-1 at Ser220. As both shear stress and genotoxic agents are well-known for their role in induction of autophagy, we asked whether IGPR-1 is activated in response to autophagy. We used human embryonic kidney epithelial-293

(HEK-293) cells ectopically expressing IGPR-1 as a model system to study the role of IGPR-1 in autophagy, as these cells do not express IGPR-1 at the detectable level[25]. To this end, HEK-293 cells expressing IGPR-1 were either kept in 10% FBS or serum-starved for 15 or 30 minutes. Cells were lysed and whole cell lysates was subjected to Western blot analysis and immunoblotted for phospho-Ser220 and total IGPR-1. Phosphorylation of IGPR-1 at Ser220 was significantly increased by brief serum-starvation of HEK-293 cells (**Figure 1A**). In an additional strategy we determined the effect of rapamycin and LPS in phosphorylation of IGPR-1 at Ser220, as both LPS and rapamycin are known to induce autophagy[13, 26, 27]. We found that treatment of IGPR-1/HEK-293 cells with rapamycin (**Figure 1B**) and LPS (**Figure 1C**) stimulated phosphorylation of IGPR-1 at Ser220.

Activation of serine/threonine kinases represents a salient mechanistic feature of autophagy. Particularly, activation of IKKβ by nutrient deprivation[28], LPS[29, 30], and rapamycin[31] is known to play a central role in autophagy. Therefore, we asked whether activation of IKKβ by serum-starvation, LPS or rapamycin can mediate phosphorylation of IGPR-1 at Ser220. To this end, we over-expressed wild type IKKβ or kinase inactive IKKβ (IKKβ-A44) in IGPR-1/HEK-293 cells. After 48hours transfection, cells were either kept in 10%FBS or serum starved for 15, 30 or 60 minutes. Cells were lysed and whole cell lysates were immunoblotted for phospho-Ser220, total IGPR-1 and phospho-IKKβ. Expression of wild type IKKβ in IGPR-1/HEK-293 cells resulted in a robust phosphorylation of IGPR-1 in the presence of 10%FBS and this was further increased in response to serum starvation (**Figure 2A**). In contrast, over-expression of kinase inactive IKKβ-A44 in IGPR-1/HEK-293 cells markedly reduced phosphorylation of IGPR-1 at Ser220 (**Figure 2A**), indicating that IKKβ kinase activity is required for the serum-starvation dependent phosphorylation of IGPR-1 at Ser220. Similarly, LPS-induced phosphorylation of Ser220 on IGPR-1 was inhibited by a selective IKKβ inhibitor, IKK inhibitor III/BMS-345541[32] (**Figure 2B**). Furthermore, LPS stimulated phosphorylation of IKKβ and IKK inhibitor blocked its phosphorylation (**Figure 2B**). Conversely over-expression of a constitutively active IKKβ (IKKβ-S177E/S181E) in IGPR-1/HEK293 cells

augmented LPS-induced phosphorylation of Ser220 (**Figure 2C**), indicating that IKKβ activity also is required for LPS-induced phosphorylation of IGPR-1 at Ser220.

IGPR-1 is a substrate for IKKβ, and is phosphorylated by IKKβ *in vitro* and *in vivo*

Ser220 and the surrounding amino acids in IGPR-1 are strongly conserved both in human and non-human primates (**Figure 3A**), suggesting an evolutionary conserved mechanism for phosphorylation of Ser220. IKKβ phosphorylates peptides with aromatic residues at the −2 position, hydrophobic residues at the +1 position, and acidic residues at the +3 position[33], suggesting that IKKβ is a likely candidate kinase involved in the phosphorylation of IGPR-1 at Ser220 (**Figure 3B**). Therefore, we asked whether IKKβ can phosphorylate IGPR-1 at Ser220 independent of the autophagy inducing factors like serum-starvation or LPS and rapamycin. We over-expressed wild-type IKKβ or kinase inactive IKKβ-A44 in IGPR-1/HEK-293 and 48 hours after transfection, cells were lysed and phosphorylation of IGPR-1 was determined by Western blot analysis. Over-expression of wild type IKKβ increased phosphorylation of IGPR-1, whereas kinase inactive IKKβ-A44 inhibited phosphorylation of IGPR-1 at Ser220 (**Figure 3C**). Similarly, IKKβ inhibitor inhibited both phosphorylation of IGPR-1 and IKKβ (**Figure 3D**).

In an additional approach, we knocked down IKKβ via CRISPR-Cas9 system and examined the effect of loss of IKKβ in IGPR-1 phosphorylation. To this end, cells were either treated with a control vehicle or AMPK activator, Oligomycin and cells were lysed and phosphorylation of IGPR-1 at Ser220 was determined. Stimulation of IGPR-1/Ctr.sgRNA/HEK-293 cells with oligomycin stimulated AMPK activation and phosphorylation of IGPR-1 at Ser220 (**Figure 3E**). However, in IGPR-1/IKKβ.sgRNA/HEK-293 cells phosphorylation of Ser220 was not detected and treatment with oligomycin also did not stimulate phosphorylation of IGPR-1 at Ser220 (**Figure 3E**). Expression of IKKβ.sgRNA knocked down IKKβ (**Figure 3E**).

We next asked whether IKKβ can directly phosphorylates IGPR-1 at Ser220. To examine the direct involvement of IKKβ in catalyzing the phosphorylation of IGPR-1, we carried out an *in vitro* kinase assay using a purified recombinant GST-IGPR-1 protein encompassing the cytoplasmic domain of IGPR-1 and demonstrated that IKKβ phosphorylates IGPR-1 at Ser220 (**Figure 3F**). Taken together, our data identifies IGPR-1 as a novel substrate of IKKβ.

IGPR-1 activates AMPK and stimulates phosphorylation of Beclin-1 and ULK1

We sought to determine whether IGPR-1 could activate AMPK, which is considered a master regulator of autophagy[34]. HEK-293 cells expressing an empty vector (EV), IGPR-1 or A220-IGPR-1 were either maintained in 10%FBS or serum starved for 30 minutes or 12 hours. The Phospho-Thr172-AMPK immunoblot of whole cell lysates showed that expression of IGPR-1 in HEK-293 cells bypassed the requirement for serum starvation-dependent activation of AMPK as AMPK was strongly phosphorylated at Thr172 in IGPR-1/HEK-293 cells in 10%FBS compared to EV/HEK-293 cells (**Figure 4A**). Phosphorylation of AMPK was further augmented in IGPR-1/HEK-23 cells, particularly at 30min compared to control EV/HEK-293 cells (**Figure 4A**). Interestingly, A220-IGPR-1/HEK-293 cells showed also an increase in phosphorylation of AMPK, notwithstanding significantly less than the wild type IGPR-1, but more than the control EV/HEK-293 cells (**Figure 4A**). Phosphorylation of AMPKα at Thr172 in the activation loop is required for AMPK activation[35, 36], indicating that expression of IGPR-1 in HEK-293 cells activated AMPK. We next asked whether IGPR-1 induced AMPK activation also stimulates phosphorylation of Beclin-1 (BECN-1) and ULK1, which are the best-known substrates of AMPK and play central roles in autophagy[37]. Similar to AMPK activation, phosphorylation of BECN at Ser93 was significantly increased in IGPR-1/HEK-293 cells compared to EV/HEK-293 or A220-IGPR-1 cells (**Figure 4A**), indicating that IGPR-1 induced activation of AMPK in HEK-293 cells also induced phosphorylation of BECN1 (**Figure 4A**) as previously reported[38]. Interestingly, although phosphorylation of ULK1 at Ser555

was increased in IGPR-1/HEK-293 cells compared to control EV/HEK-293 cells, ULK1 phosphorylation, however, remained relatively high in A220-IGPR-1 cells (**Figure 4A**). The data indicates that phosphorylation of Ser220 on IGPR-1 plays an important role in the phosphorylation of AMPK and BECN-1, however its phosphorylation appears particularly not necessary for phosphorylation of ULK-1.

IGPR-1 induces autophagy in HEK-293 cells

We next examined the role of IGPR-1 in the autophagosome formation by measuring expression of LC3-phosphatidylethanolamine conjugate (LC3-II) and p62 endogenously expressed in HEK-293 cells, which are required for autophagosome development during autophagy[39], by Western blot analysis. HEK-293 cells expressing EV or IGPR-1 were either kept at 10%FBS or serum-starved for overnight. Cells were lysed and expression of LC3 and p62 levels was determined by immunoblotting with LC3 and p62 antibodies. While expression of LC3II was increased in response to serum-starvation in EV/HEK-293 cells, however expression of LC3II was significantly higher both in 10%FBS and serum-starved conditions in IGPR-1/HEK-293 cells (**Figure 5A**), indicating that IGPR-1 through increased in expression of LC3II regulates autophagosome formation. Additionally, as expected p62 level was markedly decreased in response to serum-starvation (**Figure 5A**).   To further elucidate the role of IGPR-1 in induction of autophagy, we established autophagic flux reporter cell lines by creating GFP-LC3 (microtubule-associated protein 1 light chain 3β)-RFP/HEK-293 and IGPR-1/GFP-LC3-RFP/HEK-293 cell lines via a retroviral expression system as previously reported[40]. During autophagy, GFP-LC3-RFP labeled autophagosomes fuse with lysosomes. While the GFP signals are quenched due to the acidic environment (GFP is acid-sensitive) in the autolysosomes, the RFP signals remain stable as RFP is acid-stable and hence increased in number of RFP-LC3 (red only) puncta is considered a reflection of autophagic flux[41]. In the presence of 10%FBS, only a few RFP-LC3 positive puncta were observed in GFP-LC3-RFP/HEK-293 cells (**Figure 5B**). However, we observed a significant baseline RFP-LC3 positive puncta, but not GFP-LC3 positive puncta, in IGPR-1/GFP-LC3-RFP/HEK-293 cells (**Figure 5C**). Moreover, when

IGPR-1/GFP-LC3-RFP/HEK-293 cells were induced to undergo autophagy by serum starvation, they displayed a substantial increase in RFP-LC3 positive puncta (**Figure 5C**). Taken together the data demonstrate that IGPR-1 is activated by and regulates autophagy program.

Discussion:

Activation of IKKβ plays an essential role in autophagy as the loss of function of IKKβ in mice and cell culture both blocked autophagy[42]. Likewise, IKKβ null cells are deficient in their ability to undergo autophagy in response to cellular starvation[16], which further underscores the critical role of IKKβ in autophagy. Activation of IKKβ is thought to regulate autophagy by mechanisms that involve expression of pro-autophagic genes via NF-κB independent pathway[43, 44] and phosphorylation of the p85 subunit of PI3K, which leads to inhibition of mTOR[45]. In this study we provide a mechanistic link between activation of IKKβ and phosphorylation of IGPR-1 at Ser220. IKKβ is activated by autophagy inducting factors and conditions such as LPS, rapamycin and serum-starvation and phosphorylates IGPR-1 at Ser220 both in vivo and in vitro. Mechanistically, IKKβ mediated phosphorylation of IGPR-1 at Ser220 leads to activation of AMPK, which plays a central role in autophagy.

AMPK is believed to exert its effect in autophagy by multiple mechanisms including, inactivating mTORC1 through phosphorylation of RAPTOR, a key protein present within the mTORC1, phosphorylating ULK1 at multiple serine residues including Ser555 that leads to its activation[11, 12] and more importantly phosphorylating BECN1 at Ser93 and Ser96[38]. IGPR-1 mediated activation of AMPK in HEK-293 cells increased phosphorylation of ULK1 at Ser555 and BECN1 at Ser93. Activation of AMPK and phosphorylation of BECN1 requires phosphorylation of IGPR-1 at Ser220, however ULK1 phosphorylation by IGPR-1 appears to be less dependent on the phosphorylation of Ser220. Activation of ULK1 and phosphorylation of BECN1 both play central roles in autophagy. ULK1 phosphorylation enables ULK1 to form a complex with ATG13 and focal adhesion kinase family interacting protein of 200 kd (FIP200) that leads to its translocation to autophagy initiation sites and subsequent recruitment of

the class III phosphatidylinositol 3-kinase, vacuolar protein sorting 34 (VPS34) complex consisting of BECN1 and multiple other autophagy related proteins leading to the phagophore formation[7]. Phosphorylation of BECN1 plays a key role in the initial steps in the assembly of autophagosomes from pre-autophagic structures is the recruitment and activation of VPS34 complex [37].

IGPR-1 is newly identified cell adhesion molecule that mediates cell-cell adhesion and its activation regulates cell morphology and actin stress fiber alignment[25, 46]. The finding that IGPR-1 is activated by and regulates autophagy by stimulating activation of AMPK not only suggest a significant role for IGPR-1 in autophagy, but also links cell-cell adhesion to energy sensing and autophagy. We have previously demonstrated that IGPR-1 can support tumor cell survival under anoikis (*i.e*., in the absence cell attachment to extracellular matrix)[19]. Intriguingly, loss of cell-matrix interaction is linked to induction of autophagy[47], suggesting that anoikis-induced autophagy could be regulated by IGPR-1. Additionally, cellular stress induced by flow shear stress[21, 48] and exposure of cells to chemotherapeutic agent, doxorubicin[22, 23] are linked to induction of autophagy, the conditions where IGPR-1 also is activated[19, 49]. Moreover, autophagy is associated with therapeutic resistance, including exposure to chemotherapeutic agents (e.g., cisplatin, doxorubicin, temozolomide and etoposide), metabolic stresses, and small molecule inhibitors, suggesting a pro-tumor function for autophagy[50, 51]. Curiously, IGPR-1 is strongly phosphorylated by doxorubicin and regulates sensitivity of tumor cells to doxorubicin[19], indicating that IGPR-1 through induction of autophagy program could contribute to development of resistance in cancer cells. Taken together, the emerging evidence suggest a significant role for IGPR-1 in autophagy and autophagy-associated diseases such as cancer and cardiovascular diseases. We propose IGPR-1 as a pro-autophagy cell adhesion molecule that upon activation stimulates AMPK activation, leading to phosphorylation of BECN1 and ULK1, key proteins involved in autophagy (**Figure 6**), linking cell adhesion to autophagy, a finding that has important significance for autophagy-driven pathologies such cardiovascular diseases and cancer.

Materials and Methods:

Antibodies, Plasmids, sgRNAs and chemicals: Anti–IGPR-1 and anti-phospho-Ser220 antibodies are homemade rabbit polyclonal antibodies previously described [25, 46]. Phospho-AMPK (T172), total AMPK, phospho-ULK1 (S555), total ULK1, phosph-Beclin-1 (S93), total Beclin-1, phospho-IKK (Ser176/180), total IKKβ, and LC3A/B, GAPDH antibodies all were purchased from Cell Signaling Technologies (Danvers, MA). The following plasmids were all purchased from Addgene (Watertown, MA); pcDNA3.FLAG-ULK1 (Cat#27636), pMRX-IP-GFP-LC3-RFP (Cat#84573), pcDNA-IKKβ-FLAG (Cat#23298), pcDNA-IKKβ-A44 (Cat#23299), and constitutive active IKKβ (S177E S181E, Cat#11105). IGPR-1 constructs including wild-type IGPR-1 and Ser220 mutant, A220-IGPR-1 constructs were cloned into retroviral vector, pQCXIP with C-terminal Myc tag as previously described[19, 25, 49]. Retroviruses were produced in 293-GPG cells as described[52]. IKK inhibitor III and rapamycin were purchased from Calbiochem and LPS was purchased from Sigma. Oligomycin was purchased from Cell Signaling Technologies (Danvers, MA). A set of three Human IKKβ sgRNAs (cat# GSGH-11938-16EG3551) were purchased from Dharmacon (Chicago, IL).

Cell culture assays: HEK-293 cells expressing empty vector (EV), IGPR-1 or A220-IGPR-1 were maintained in DMEM supplemented with 10% fetal bovine serum and penicillin/streptomycin. To measure phosphorylation of IGPR-1 in response to serum-starvation, cells were plated in 60cm plates with 10%FBS DMEM for overnight with approximate 80-90% confluency. Cells were washed twice with PBS and cells were starved for 15 and 30 minutes or as described in the figure legends. Cells were lysed and whole cell lysates were mixed with sample buffer (5X) and boiled for 5minutes. Whole cell lysates were subjected to Western blot analysis and immunoblotted with antibody of the interest as described in the figure legends. In some experiments, cells were treated with a specific chemical inhibitor or transfected with a particular construct as indicated in the figure legends.

Recombinant GST fusion protein production: The generation of GST-fusion cytoplasmic domain of IGPR-1 cloned into pGEX-2T vector was previously described[46]. The purified GST fusion IGPR-1 protein was subsequently used to measure the ability of IKKβ to phosphorylate IGPR-1 in an in vitro kinase.

In vitro kinase assay: To detect phosphorylation of IGPR-1 at Ser220, the purified recombinant GST-IGPR-1 encompassing the cytoplasmic domain of IGPR-1 was mixed with wild type or kinase inactive IKKβ expressed in HEK-293 cells in 1× kinase buffer plus 0.2mM ATP, and incubated at 30°C for 15 min. The samples were mixed in 2X sample buffer and after boiling at 95°C for 5min were resolved on 12% SDS–PAGE followed by Western blot analysis using anti-phospho-Ser220 antibody.

Western blotting analysis: The cells were prepared as described in the figures legends, lysed, and whole cell lysates were subjected to Western blot analysis. Normalized whole cell lysates were subjected to Western blotting analysis using IGPR-1 antibody, phospho-Ser220 antibodies or with appropriate antibody as indicated in the figure legends. Proteins were visualized using streptavidin–horseradish peroxidase–conjugated secondary antibody via chemiluminescence system. For each blot, films were exposed multiple times and films that showed within the linear range detection of protein bands were selected, scanned and subsequently used for quantification. Blots from at least three independent experiments were used for quantification purposes and representative data are shown. Image J software, an open source image processing program, used to quantify blots.

Immunofluorescence microscopy: Cells expressing IGPR-1 or other constructs were seeded (1.5×10^6 cells) onto coverslips and grown overnight in 60-mm plates to 90–100% confluence. The coverslips were mounted in Vectashield mounting medium with DAPI onto glass microscope slides. The slides were examined using a fluorescence microscope.

Statistical analyses: Experimental data were subjected to Student t-test or One-way analysis of variance analysis where appropriate with representative of at least three independent experiments. $p < 0.05$ was

considered significant or indicated with two stars (**) if a p-value was less than 0.01 and with three stars

(***) If a p-value was less than 0.001.

References:

1.   Kroemer G, Marino G, Levine B. Autophagy and the integrated stress response. Molecular cell 2010; 40:280-93.
2.   Vlahakis A, Debnath J. The Interconnections between Autophagy and Integrin-Mediated Cell Adhesion. Journal of molecular biology 2017; 429:515-30.
3.   Neill T, Schaefer L, Iozzo RV. Instructive roles of extracellular matrix on autophagy. The American journal of pathology 2014; 184:2146-53.
4.   Kenific CM, Wittmann T, Debnath J. Autophagy in adhesion and migration. Journal of cell science 2016; 129:3685-93.
5.   Fung C, Lock R, Gao S, Salas E, Debnath J. Induction of autophagy during extracellular matrix detachment promotes cell survival. Molecular biology of the cell 2008; 19:797-806.
6.   Hurley JH, Young LN. Mechanisms of Autophagy Initiation. Annu Rev Biochem 2017; 86:225-44.
7.   Kametaka S, Okano T, Ohsumi M, Ohsumi Y. Apg14p and Apg6/Vps30p form a protein complex essential for autophagy in the yeast, Saccharomyces cerevisiae. J Biol Chem 1998; 273:22284-91.
8.   Huang WP, Klionsky DJ. Autophagy in yeast: a review of the molecular machinery. Cell Struct Funct 2002; 27:409-20.
9.   Jung CH, Jun CB, Ro SH, Kim YM, Otto NM, Cao J, et al. ULK-Atg13-FIP200 complexes mediate mTOR signaling to the autophagy machinery. Mol Biol Cell 2009; 20:1992-2003.
10.   Hosokawa N, Hara T, Kaizuka T, Kishi C, Takamura A, Miura Y, et al. Nutrient-dependent mTORC1 association with the ULK1-Atg13-FIP200 complex required for autophagy. Mol Biol Cell 2009; 20:1981-91.
11.   Kim J, Kundu M, Viollet B, Guan KL. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. Nat Cell Biol 2011; 13:132-41.
12.   Egan DF, Shackelford DB, Mihaylova MM, Gelino S, Kohnz RA, Mair W, et al. Phosphorylation of ULK1 (hATG1) by AMP-activated protein kinase connects energy sensing to mitophagy. Science 2011; 331:456-61.
13.   Kroemer G, Marino G, Levine B. Autophagy and the integrated stress response. Mol Cell 2010; 40:280-93.
14.   Perkins ND. Integrating cell-signalling pathways with NF-kappaB and IKK function. Nature reviews Molecular cell biology 2007; 8:49-62.
15.   Criollo A, Senovilla L, Authier H, Maiuri MC, Morselli E, Vitale I, et al. The IKK complex contributes to the induction of autophagy. The EMBO journal 2010; 29:619-31.
16.   Comb WC, Cogswell P, Sitcheran R, Baldwin AS. IKK-dependent, NF-kappaB-independent control of autophagic gene expression. Oncogene 2011; 30:1727-32.
17.   Rahimi N, Rezazadeh K, Mahoney JE, Hartsough E, Meyer RD. Identification of IGPR-1 as a novel adhesion molecule involved in angiogenesis. Molecular biology of the cell 2012; 23:1646-56.
18.   Wang YHW, Meyer RD, Bondzie PA, Jiang Y, Rahimi I, Rezazadeh K, et al. IGPR-1 Is Required for Endothelial Cell-Cell Adhesion and Barrier Function. Journal of molecular biology 2016; 428:5019-33.
19.   Woolf N, Pearson BE, Bondzie PA, Meyer RD, Lavaei M, Belkina AC, et al. Targeting tumor multicellular aggregation through IGPR-1 inhibits colon cancer growth and improves chemotherapy. Oncogenesis 2017; 6:e378.

20.   Ho RX, Tahboub R, Amraei R, Meyer RD, Varongchayakul N, Grinstaff MW, et al. The cell adhesion molecule IGPR-1 is activated by, and regulates responses of endothelial cells to shear stress. J Biol Chem 2019.

21.   Liu J, Bi X, Chen T, Zhang Q, Wang SX, Chiu JJ, et al. Shear stress regulates endothelial cell autophagy via redox regulation and Sirt1 expression. Cell death & disease 2015; 6:e1827.

22.   Chen H, Zhao C, He R, Zhou M, Liu Y, Guo X, et al. Danthron suppresses autophagy and sensitizes pancreatic cancer cells to doxorubicin. Toxicol In Vitro 2019; 54:345-53.

23.   Sui X, Chen R, Wang Z, Huang Z, Kong N, Zhang M, et al. Autophagy and chemotherapy resistance: a promising therapeutic target for cancer treatment. Cell Death Dis 2013; 4:e838.

24.   Wang YH, Meyer RD, Bondzie PA, Jiang Y, Rahimi I, Rezazadeh K, et al. IGPR-1 Is Required for Endothelial Cell-Cell Adhesion and Barrier Function. J Mol Biol 2016; 428:5019-33.

25.   Wang YHW, Meyer RD, Bondzie PA, Jiang Y, Rahimi I, Rezazadeh K, et al. IGPR-1 Is Required for Endothelial Cell-Cell Adhesion and Barrier Function. J Mol Biol 2016; 428:5019-33.

26.   Xu Y, Jagannath C, Liu X-D, Sharafkhaneh A, Kolodziejska KE, Eissa NT. Toll-like receptor 4 is a sensor for autophagy associated with innate immunity. Immunity 2007; 27:135-44.

27.   Dunlop EA, Tee AR. mTOR and autophagy: a dynamic relationship governed by nutrients and energy. Seminars in cell & developmental biology 2014; 36:121-9.

28.   Comb WC, Hutti JE, Cogswell P, Cantley LC, Baldwin AS. p85alpha SH2 domain phosphorylation by IKK promotes feedback inhibition of PI3K and Akt in response to cellular starvation. Molecular cell 2012; 45:719-30.

29.   Yang F, Tang E, Guan K, Wang CY. IKK beta plays an essential role in the phosphorylation of RelA/p65 on serine 536 induced by lipopolysaccharide. J Immunol 2003; 170:5630-5.

30.   Dauphinee SM, Karsan A. Lipopolysaccharide signaling in endothelial cells. Lab Invest 2006; 86:9-22.

31.   Dan HC, Cooper MJ, Cogswell PC, Duncan JA, Ting JPY, Baldwin AS. Akt-dependent regulation of NF-{kappa}B is controlled by mTOR and Raptor in association with IKK. Genes & development 2008; 22:1490-500.

32.   Burke JR, Pattoli MA, Gregor KR, Brassil PJ, MacMaster JF, McIntyre KW, et al. BMS-345541 is a highly selective inhibitor of I kappa B kinase that binds at an allosteric site of the enzyme and blocks NF-kappa B-dependent transcription in mice. The Journal of biological chemistry 2003; 278:1450-6.

33.   Hutti JE, Turk BE, Asara JM, Ma A, Cantley LC, Abbott DW. IkappaB kinase beta phosphorylates the K63 deubiquitinase A20 to cause feedback inhibition of the NF-kappaB pathway. Mol Cell Biol 2007; 27:7451-61.

34.   Kim J, Kundu M, Viollet B, Guan K-L. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. Nature cell biology 2011; 13:132-41.

35.   Lizcano JM, Goransson O, Toth R, Deak M, Morrice NA, Boudeau J, et al. LKB1 is a master kinase that activates 13 kinases of the AMPK subfamily, including MARK/PAR-1. EMBO J 2004; 23:833-43.

36.   Hawley SA, Davison M, Woods A, Davies SP, Beri RK, Carling D, et al. Characterization of the AMP-activated protein kinase kinase from rat liver and identification of threonine 172 as the major site at which it phosphorylates AMP-activated protein kinase. J Biol Chem 1996; 271:27879-87.

37.   Menon MB, Dhamija S. Beclin 1 Phosphorylation - at the Center of Autophagy Regulation. Front Cell Dev Biol 2018; 6:137.

38.   Kim J, Kim YC, Fang C, Russell RC, Kim JH, Fan W, et al. Differential regulation of distinct Vps34 complexes by AMPK in nutrient stress and autophagy. Cell 2013; 152:290-303.

39.   Schaaf MB, Keulers TG, Vooijs MA, Rouschop KM. LC3/GABARAP family proteins: autophagy-(un)related functions. FASEB J 2016; 30:3961-78.

40.   Kaizuka T, Morishita H, Hama Y, Tsukamoto S, Matsui T, Toyota Y, et al. An Autophagic Flux Probe that Releases an Internal Control. Mol Cell 2016; 64:835-49.

41.   Ni H-M, Bockus A, Wozniak AL, Jones K, Weinman S, Yin X-M, et al. Dissecting the dynamic turnover of GFP-LC3 in the autolysosome. Autophagy 2011; 7:188-204.

42.   Criollo A, Senovilla L, Authier H, Maiuri MC, Morselli E, Vitale I, et al. The IKK complex contributes to the induction of autophagy. EMBO J 2010; 29:619-31.

43.   Baldwin AS. Regulation of cell death and autophagy by IKK and NF-kappaB: critical mechanisms in immune function and cancer. Immunological reviews 2012; 246:327-45.

44.   Criollo A, Senovilla L, Authier H, Maiuri MC, Morselli E, Vitale I, et al. IKK connects autophagy to major stress pathways. Autophagy 2010; 6:189-91.

45.   Comb WC, Hutti JE, Cogswell P, Cantley LC, Baldwin AS. p85alpha SH2 domain phosphorylation by IKK promotes feedback inhibition of PI3K and Akt in response to cellular starvation. Mol Cell 2012; 45:719-30.

46.   Rahimi N, Rezazadeh K, Mahoney JE, Hartsough E, Meyer RD. Identification of IGPR-1 as a novel adhesion molecule involved in angiogenesis. Mol Biol Cell 2012; 23:1646-56.

47.   Fung C, Lock R, Gao S, Salas E, Debnath J. Induction of autophagy during extracellular matrix detachment promotes cell survival. Mol Biol Cell 2008; 19:797-806.

48.   Guo F, Li X, Peng J, Tang Y, Yang Q, Liu L, et al. Autophagy regulates vascular endothelial cell eNOS and ET-1 expression induced by laminar shear stress in an ex vivo perfused system. Ann Biomed Eng 2014; 42:1978-88.

49.   Ho RX, Tahboub R, Amraei R, Meyer RD, Varongchayakul N, Grinstaff M, et al. The cell adhesion molecule IGPR-1 is activated by and regulates responses of endothelial cells to shear stress. J Biol Chem 2019; 294:13671-80.

50.   Amaravadi RK, Thompson CB. The roles of therapy-induced autophagy and necrosis in cancer treatment. Clin Cancer Res 2007; 13:7271-9.

51.   Degtyarev M, De Maziere A, Orr C, Lin J, Lee BB, Tien JY, et al. Akt inhibition promotes autophagy and sensitizes PTEN-null tumors to lysosomotropic agents. J Cell Biol 2008; 183:101-16.

52.   Rahimi N, Dayanir V, Lashkari K. Receptor chimeras indicate that the vascular endothelial growth factor receptor-1 (VEGFR-1) modulates mitogenic activity of VEGFR-2 in endothelial cells. J Biol Chem 2000; 275:16986-92.

Figure legends:

Figure 1. IGPR-1 is activated by autophagy stimuli LPS, serum starvation and rapamycin. (**A**) IGPR-1/HEK-293 cells were treated with different concentrations of LPS for one hour as indicated. Whole cell lysates was prepared and subjected to Western blot analysis and blotted with anti-phosphoserine 220

IGPR-1 antibody (pSer220), anti-IGPR-1 antibody and anti-GAPDH antibody for protein loading control. Graph is a representative of three independent experiments. (**B**) IGPR-1/HEK-293 cells were seeded for 48hr starved for 15 and 30 minutes by replacing the 10% DMEM media with serum-free DMEM media or cells left in 10% FBS DMEM media as a control (0). Cells were lysed and whole cell lysates blotted with the same antibodies as panel A. Graph is a representative of three independent experiments. (**C**) IGPR-1/HEK-293 cells were treated with vehicle (0) or with rapamycin (10nM) for one hour. Whole cell lysates prepared and subjected to Western blot analysis using the same antibodies as panel A. Graph is a representative of three independent experiments.

Figure 2: **IKKβ mediates LPS- and serum starvation-dependent activation of IGPR-1**. (**A**) IGPR-1/HEK-293 cells were treated with LPS (5µg/ml) alone or co-treated with LPS and IKKβ inhibitor, IKK inhibitor III/BMS-345541 (100ηM) for 60min. Cells were lysed and whole cell lysates was subjected to Western blot analysis and probed for pSer220, total IGPR-1, phospho-IKKβ, total IKKβ and GAPDH for protein loading control. Graph is a representative of three independent experiments. (**B**) IGPR-1/HEK-293 cells were transfected with an empty vector (-) or constitutively active IKKβ (IKKβ-S177E/S181E). After 48 hours cells were either left untreated (-) or treated with LPS (5µg/ml) for 60min (+). Whole cell lysates was subjected to Western blot analysis and probed for pSer220, total IGPR-1, total IKKβ and GAPDH for protein loading control. Graph is a representative of three independent experiments. (**C**) IGPR-1/HEK293 cells transfected with an empty vector, wild-type IKKβ (Wt.IKKβ) or with kinase-dead IKKβ (IKKβ-A44). After 48 hours transfection, cells were either left in 10% FBS DMEM or serum starved for 15 or 30 minutes. Cells were lysed and whole cell lysates immunoblotted for pSer220, total IGPR-1, phospho-IKKβ, total IKKβ or GAPDH for protein loading control. Graph is a representative of three independent experiments.

Figure 3: IGPR-1 is a substrate of and phosphorylated by IKKβ. (**A**) Ser220 and the surrounding amino acids on IGPR-1 in human and non-human primates is conserved. (**B**) IKKβ phosphorylation motif and IGPR-1 Ser220 phosphorylation site. (**C**) IGPR-1/HEK-293 cells were transfected with wild type IKKβ (Wt. IKKβ) or kinase inactive IKKβ (IKKβ-A44). After 48 hours cells were lysed and immunoblotted for pSer220, total IGPR-1, phospho-IKKβ, total IKKβ and GAPDH. Graph is a representative of three independent experiments. (**D**) IGPR-1/HEK-293 cells were treated with IKK inhibitor III/BMS-345541 (100ηM) for 60min. Cells were lysed and immunoblotted for pSer220, total IGPR-1, phospho-IKKβ, and total IKKβ. (**E**) Ctr.sgRNA/IGPR-1/HEK-293 cells and IKKβ-sgRNA/IGPR-1/HEK-293 cells were stimulated with IKKβ activator, Oligomycin (0.5μM), cells were lysed and subjected to Western blot analysis and immunoblotted for the indicated proteins. (**F**) In vitro kinase assay was performed by incubation of purified GST-cytoplasmic domain of IGPR-1 and purified wild type IKKβ or kinase inactive IKKβ-A44 in the presence of ATP. In vitro kinase reaction was stopped after 30min by heating the samples at 90°C for 5 min. Samples were subjected to Western blotting analysis and phosphorylation of Ser220 on GST-IGPR-1 was detected by immunoblotting with pSer220 antibody.

Figure 4: IGPR-1 stimulates phosphorylation of AMPK and mediates phosphorylation of ULK1. (**A**) HEK-293 cells expressing empty vector (EV), IGPR-1 or A220-IGPR-1 were either kept in 10%FBS DMEM or starved for 30 min or 12hours. Cells were lysed and immunoblotted for pT172-AMPK, pSer555-ULK1, pSer15-Beclin1, pSer220-IGPR-1, total IGPR-1 and GAPDH. (**B**) HEK-293 cells expressing empty vector (EV), IGPR-1 were either kept in 10%FBS or serum-starved for 12hours. Cells were lysed and immunoblotted for LC3 and p26, IGPR-1 and GAPDH. Graph is a presentative of three experiments.

Figure 5. IGPR-1 mediates serum starvation-induced autophagy: HEK-293 cells expressing GFP-LC3-RFP/HEK-293 alone or co-expressing GFP-LC3-RFP and IGPR-1 were either kept in 10%FBS DMEM (**A**) or in serum-free DMEM for 6hours (**B**). Cells were fixed and stained with DAPI (nucleus) and viewed under

a fluorescence microscope and pictures were taken. Image magnification (40X). (**C**) EV/HEK-293 and IGPR-1/HEK-293 cells were kept in 10% FBS or serum-starved for overnight, cells were lysed and whole cell lysates were blotted for LC3, p62, total IGPR-1, phospho-Ser220. Graph is a presentative of three independent experiments.

Figure 6. Proposed model of IGPR-1 in autophagy. Upon activation by autophagy stimuli, IGPR-1 is phosphorylated at Ser220 and acts as a pro-autophagy signaling receptor. Upon activation, it stimulates AMPK activation, leading to phosphorylation of BECN1 and ULK1, key proteins involved in autophagy. At the plasma membrane, IGPR-1 is present as a dimer and undergoes homophilic transdimerization in a cell density dependent manner[25] (not shown).

Amraie et al., Figure 1

Amraie et al., Figure 2

A

Amraie et al., Figure 6