1   When to be a male? Role of resource-limitation and pollinators in determining gender in an

2   andromonoecious spiderwort.

6   Asawari Albal [1,3,4], Azad G[1], Saket Shrotri[1], Vinita Gowda[1,2]

[1] Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences, Indian

Institute of Science Education and Research Bhopal, Bhopal, 462066, India

[2] Corresponding author: Vinita Gowda, gowdav@iiserb.ac.in

Present address:

[3] Department of Biology, University of Toronto Mississauga, Mississauga, ON L5L 1C6, Canada

[4] Ecology and Evolutionary Biology, University of Toronto, Toronto, ON M5S 2Z9, Canada

Abstract

The evolution and maintenance of sexual systems in plants is often driven by resource allocation and pollinator preferences. Gender or sex in plants, therefore, represents a physiological, functional, and behavioral response of a plant to its environment. Annual, entomophilous plants have three major constraints towards optimal reproduction: 1) nutrient resources obtained from the environment, 2) nutrient resources allocated towards reproduction, i.e., fruits vs. flowers, and 3) pollinator visitations. We know very little about the role of nutrient resources and pollinators in driving gender expression in flowers in a natural environment.

Andromonoecy is a sexual system where the plant bears both staminate and hermaphrodite flowers on the same inflorescence and it occurs in ~2% of flowering plants. Although it has been shown that andromonoecy has evolved multiple times within the angiosperms, our understanding of the underlying mechanisms of what determines gender expression in nature is limited. The optimal resource allocation hypothesis, one of the most widely accepted hypotheses which describes evolution of andromonoecy, suggests that under nutrient constraints, plants will produce more male flowers since they require lesser energy investment than the more expensive hermaphrodite flowers. We test this hypothesis in an andromonoecious plant *Murdannia simplex* (Commelinaceae) by measuring gender distributions in a large natural population and contrasting the gender distributions between two different resource conditions (stream population vs. plateau population). We also carried out choice experiments to quantify if pollinators showed any preference towards a gender.

We found that in *M. simplex*, production of hermaphrodite flowers is resource-dependent and under resource constraints not only fewer numbers of flowers were produced but also there were a higher number of male flowers. We failed to observe pollinator preference towards either

gender but *Amegilla spp. (Zonamegilla)* and *Apis cerana* showed frequency-dependent visitation

which suggests that selection on inflorescence size may be an indirect form of selection pressure

exerted by pollinators resulting in gender-bias in the plants. Thus, we conclude that

environmentally driven resource constraints play a bigger role in driving floral gender expression

in *Murdannia* over direct pollinator-driven constraints.

Keywords

Andromonoecy, Commelinaceae, floral gender, male, hermaphrodite, *Murdannia simplex*,

resource limitation, pollinator preference, Western Ghats

Introduction

Andromonoecy refers to the sexual system where both staminate (male) flowers and perfect

(hermaphrodite) flowers are present on the same plant (Bertin 1982a, Miller and Diggle 2003,

Vallejo-Marín and Rausher 2007a, 2007b). In andromonoecious plants, the hermaphrodite flower

fulfills the female function in the absence of a true 'female' flower, and it has been suggested

that andromonoecy probably evolved from hermaphroditism by the loss of female reproductive

structure (Lloyd 1980, Bertin 1982a). One of the most widely accepted hypotheses on the

evolution of andromonoecy is the optimal resource allocation hypothesis which suggests that

under resource limitation, male flowers will be produced instead of hermaphrodite flowers. This

hypothesis rests upon the premise that male flowers are energetically cheaper to produce because

reproductive allocation towards them ends with pollen production (Anderson and Symon 1989,

Narbona et al. 2002, Verdú et al. 2007), while females are expensive because fruit development

and seed germination are energy-intensive physiological processes (Bertin 1982a, Kaul et al.

2002, Obeso 2002, Verdú et al. 2007).

Examples of reallocation of resources to the cheaper sex (i.e. male bias) have been shown

in several plant species where staminate flowers are produced when nutrient resources become

scarce due to its utilization by the costlier gynoecia towards fruit production (Bertin 1982a,

1982b, May and Eugene Spears Jr 1988, Miller and Diggle 2003, Venkatesan 2017). A well-

studied species is *Solanum hirtum* (Solanaceae) in which it has been shown that as a result of

successful pollination, as the ovary develops, resources are relocated within an inflorescence to

the male flowers, thus influencing gender expression and distribution of gender in successive

flowers (Diggle 1993, Miller and Diggle 2003). Another example comes from *Raphanus*

raphanistrum, where Stanton et al. (1987) showed that in plants which were heavily pollinated,

the number of ovules per flower decreased and in contrast, unpollinated plants did not show any

significant decrease in number of ovules. This suggests that plants can modify their reproductive

output to adjust for the resource limitation faced by them.

In plants, resource limitation or nutrient limitation that can affect floral gender expression

can be identified as: a) environmental nutrient limitation (Charnov and Bull 1977, Primack and

Lloyd 1980a, Stephenson 1981, Bertin 1982a, Solomon 1985, Diggle 1993), and/or b) within-

plant nutrient limitation (Stephenson 1981, May and Eugene Spears Jr 1988, Diggle 1993, Miller

and Diggle 2003, Ortiz et al. 2003, Vallejo-Marín and Rausher 2007a). In *Aesculus californica*,

A. pavia, *Leptospermum scoparium* and *Passiflora incarnata*, phenotypic gender expression

(hermaphroditic inflorescences or hermaphroditic flowers) has been shown to vary as a response

to environmental conditions or status of the nutrient resources, especially during fruit

development when resources are limited (Benseler 1975, Primack and Lloyd 1980b, May and

Eugene Spears Jr 1988). Further, it has been shown that the amount of resources that a plant

acquires in the proximal/distal ends of the inflorescence can also vary. The proximal/ basal end

of the inflorescence has more resources than its distal end, which results in more hermaphrodite

flowers to be present at basal ranks of an inflorescence, while male flowers are relegated to the

distal end (Diggle 1994, 1997, Lewis and Gibbs 1999, Ashman and Hitchens 2000, Miller and

Diggle 2003, Kaul and Koul 2008).

Bateman's principle (Wilson et al. 1994) asserts that- "Fitness gain through male function

is limited primarily by mating opportunity, while fitness gain through female function is limited

primarily by resource availability for offspring production". To explain the role of male flowers

in andromonoecious plants, two not-mutually exclusive hypotheses have been proposed (Zhang

and Tan 2009): i) Pollen donor hypothesis which approaches the problem from the perspective of

male reproductive success. It predicts that in a resource-limited condition the wastage of nutrient

resources by investment in the female reproductive structure is reduced by the production of only

male flowers. ii) Pollinator attractor hypothesis which is based on female reproductive success. It

predicts that male flowers increase the floral display and attract more pollinators at a lower cost

thereby increasing the stigmatic load of the hermaphrodite flowers, which increases female

reproductive success at lower resource investment. The sex allocation theory predicts that by

attracting pollinators which promote outcrossing, floral attractiveness (e.g., large petals, nectar

availability) gains more fitness through male function than female function (Elle and Meagher

2000). This is because male success is correlated with mating opportunities, while one or a few

visits by pollinators can adequately fertilize all ovules. Thus, it is not expected that female fitness

will be strongly affected by floral attractiveness.

In entomophilous plants, pollinator dependence is often driven by floral characteristics

and for successful pollination the gender of the flowers and associated floral characters may be

very important (Armbruster 2001, Fenster et al. 2004). In many hermaphroditic plant systems, it

has been shown that several characteristics of an inflorescence such as number of flowers, size of

flowers, and reward of flowers play a critical role in attracting pollinators to the plant (Ortiz et al.

2003, Christopher et al. 2019). While the role of pollinators is very well known in

hermaphroditic systems, the perception of gender, pollinator preference, and reproductive output

in andromonoecious systems are not well studied (see Diggle 1993, Ashman et al. 2000, Ashman

and Morgan 2004).

Previous studies in sexually dimorphic plants have shown that pollinators typically show

a preference for the functionally male flowers over the conspecific functional female flowers

owing to different floral traits like petal length (Ashman et al. 2000), levels of pollen production

(Bierzychudek 1987, Eckhart 1991) and/or floral scents (Ashman et al. 2005), where these

preferences are learned through foraging experience (Cresswell and Galen 1991). Zhang and Tan

(Zhang and Tan 2009) were the first to examine the function of male flowers in the

andromonoecious *Capparis spinosa* where they showed that the pollinators did not discriminate

between the floral gender morphs, even when they were morphologically distinct. However,

similar studies investigating gender-specific pollinator preferences and more specifically gender-

specific pollinator preferences in sexually dimorphic plants that have near-identical

morphological features are not known.

In this study, we explore the effect of nutrient resources as well as pollinator preference

in driving gender expression in a wild population of the andromonoecious *Murdannia simplex*

(Commelinaceae, Spiderworts). The pantropical plant family Commelinaceae (Spiderworts) is

known to have several genera that display andromonoecy. Despite the dominance of

andromonoecy in this family, to date, most pollination studies within this family have focused

only on identifying pollinators (Kaul et al. 2002, Williams and Walker 2003, Oziegbe et al. 2013,

Sigrist and Sazima 2015), and only one study has explored the role of floral guides in pollinator

attraction (Ushimaru et al. 2007). Based on predictions from the optimal resource allocation

hypothesis and pollinator attractor hypothesis we address the following questions:

A) Does the proportion of staminate and perfect flowers in a population represent resource

limitation?

B) Do pollinators show preference to any one of the genders?

C) Do staminate flowers improve floral display to maintain visitation rate and function as a

cheaper source of outcrossing pollen?

Materials and methods

Murdannia simplex (Vahl) Brenan (Commelinaceae) is a mass-flowering, sub-erect, annual,

andromonoecious herb which is widely distributed in moist deciduous forests and grasslands of

the Western Ghats of India (Nandikar and Gurav 2015; Fig. 1a). The plant is seasonal, 40-65 cm

tall, and the shoot emerges once monsoonal rains begin. Flowering lasts from September to

November (Nandikar and Gurav 2015) and flowers are borne on a cymose inflorescence. The

inflorescence bears 5-20 purple-colored, three-petaled male and hermaphrodite flowers

(andromonoecy), and each flower measures ~15-25 mm in diameter. The hermaphrodite flowers

are characterized by the presence of a lateral pistil, with two upper fertile stamens, curved

downwards and a third lower sterile stamen. Filaments have long purple, bearded hair. The

flower also bears three trilobed staminodes with glabrous filaments. The male flowers are

characterized only by the lack of a pistil (Fig. 1b - f). The flowers are defined by a short floral

anthesis time, from 12:30 hrs. to 16:00 hrs., with the principal pollinator reward being pollen

grains only (Faden 2000, Nandikar and Gurav 2015).

Study site

Western Ghats is one of the four biodiversity 'hotspots' in India. The northern part of Western

Ghats is famous for high elevation laterite plateaus and slopes, one of them being Kaas plateau in

Maharashtra (17° 43′ 12″ N, 73° 49′ 22″ E), which is located at an altitude of ~1225 m above sea

level (Fig. 1a). Kaas plateau has an area of ~10 km^2 and is situated northwest, about 25 km away

from the city of Satara in the western state of Maharashtra. It was declared as a UNESCO World

Heritage site in 2012 due to its highly endemic floral diversity, which is most abundant in the

months of August to November.

Capped with red lateritic crusts, the upper Kaas Plateau provides arid habitats except

during the monsoon season (Fig. 1a). It receives over 2500 mm of rain every year, mainly during

the monsoon months (June to September), and the daily mean temperatures are over 22°C. Floral

diversity on the Kaas plateau is a model representation of flora that is associated with seasonal

monsoonal rains in the Western Ghats, as well as the flora of lateritic plateaus of the Western

Ghats. All field experiments were carried out across three consecutive years - 2017, 2018 and

2019, between 15th September and 5th November. The first two years focused on the study of

plant-pollinator interactions and the third year involved the study of the effect of nutrient

resources on the gender of the flower, and also monitored possible yearly fluctuations in gender

expression.

Gender distribution in the population

To determine the extent of sexual dimorphism between genders, flower sizes were compared.

Pairs of male and hermaphrodite flowers were collected from 15 random inflorescences and then

arranged and photographed on a laminated graph sheet. From these images, petal length and

width (widest diameter) were measured manually (Appendix S1: Fig. S1).

In order to identify if gender distribution within a plant and within an inflorescence was

naturally dominant for male or hermaphrodite flowers, a multi-year (2018 and 2019) census was

conducted within the study site. All census were carried out at an interval of seven days, across 4

weeks (spanning over one and a half months) covering the peak flowering season of *M. simplex*.

Gender census were conducted between 12:30 hrs. to 16:00 hrs. (anthesis time of *M. simplex*

flowers), on marked inflorescences (N = 30). Since a single individual can bear 2-5

inflorescences, a pooled census was taken from all the inflorescences of a plant as our initial

observations found no difference among the inflorescences within a plant. To identify if there is

a positional-gender bias i.e. from basal to the apical position within an inflorescence (see

Stephenson 1981, Diggle 1997, Cuevas and Polito 2004) the positions of each flower within an

inflorescence were recorded along with its gender details, over the census duration mentioned

above. The positions within an inflorescence were divided into three categories- apical, middle,

and basal as shown in Fig. 1c.

To test the effect of nutrient resources on gender expression, a census of floral gender

was conducted between two populations that differed in the availability of water and other

resources (presence of a stream and a deep soil substrate), which we predicted to be primary

limiting resources on an otherwise homogeneous plateau. The first population is henceforth

referred to as the plateau population - 'Plateau_pop', while the second population is referred to

as the stream population - 'Stream_pop'. Plateau_pop was selected as the resource-poor

population whereas Stream_pop was selected as the resource-rich population. The two locations

were separated by ~576 meters and other than the presence of a stream and a deep soil substrate,

can be viewed as a single population. The census was carried out in October 2019 on randomly

selected individuals within these two populations (N = 100 per location).

Pollination biology

Pollen is the only pollinator reward present in the genus *Murdannia* (Faden 2000, Nandikar and

Gurav 2015). To identify any quantifiable difference in the pollen or paternal contribution

between the male and hermaphroditic flowers, the amount of pollen per anther in virgin flowers

of *M. simplex* was quantified using a hemocytometer.

Pollinator visitations on *M. simplex* in the wild were quantified by observing pollinators

in 2 ft. x 2 ft. observation arenas (N = 49) in multiple intervals of 15 minutes between 12:30 hrs.

to 16:00 hrs. (anthesis time of *M. simplex*). Each arena consisted of 4-10 inflorescences having

15-30 flowers per area. Visitation by a pollinator was recorded when a pollinator was observed

to land on the flower and actively collect pollen. The following three features were recorded and

quantified: total number of flowers present, type of pollinator, and number of pollinator visits in

a 15-minute interval. Pollinator visitation rate was then calculated as the total number of visits

per flower per hour. At the end of the flowering season, fruitset was quantified for *M. simplex* by

measuring the number of fruits in randomly selected inflorescences from the study population in

2017 (N = 26) and 2019 (N = 40).

Gender-specific pollinator preference experiment

Manipulated choice experiments were designed and conducted in the wild to test a) presence of

visitation bias by pollinators towards male flowers (M) and/or hermaphrodite flowers (H), and b)

presence of floral-density dependent bias by pollinators. It was hypothesized that if the pollinator

has an inherent gender preference, irrespective of the gender density in a population, pollinators

will show high visitation to the same gender. Floral distributions within an inflorescence were

defined to have binary outcomes i.e., male abundant or hermaphrodite abundant, or equally male

and hermaphroditic. The experimental setup consisted of two pots that were kept one foot apart

with a total of 5-15 flowers in each pot and pollinator visits were quantified by eye between

12:30 hrs. to 16:00 hrs. Three gender-density treatments: H < M (T1), H = M (T2) and H > M

(T3) were tested. The density of flowers of the two genders in each experimental set up was

manually controlled and each treatment was repeated for a total of eight trials and a total

observation time of ~21.5 hours (Appendix S1: Fig. S2). Unmanipulated, wild set up

observations served as the control. The mean visitation rate per flower to male flowers and

hermaphrodite flowers were calculated for each pollinator type, by dividing the total number of

visits by a specific pollinator with the total number of flowers present in the experimental set up.

Pollinator visitation rate was calculated as the total number of visits per flower per hour.

239   The overall visitation observed in all three different density treatments in the

manipulative choice experiment were compared to test the effect of change in density on the

visitation pattern of pollinators in this andromonoecious species. The visitation rates in the

control treatments (natural visitations) were calculated and compared with the visitation rates

obtained in our choice experiment (H < M, H = M and H > M). The mean visitation rates across

all three treatments were also compared with the total number of flowers present in the trial to

identify any effect of floral display size.

Statistical analysis

All analyses were performed in R (R Core Team 2019) and raw data was tested for normality

before proceeding with statistical analysis using Shapiro-Wilk's method. When Gaussian type

distribution could not be assumed, non-parametric Kruskal-Wallis rank-sum tests were run using

the dplyr package (Wickham et al. 2020). The floral diameters and pollen densities per flower

between male and hermaphrodite flowers were compared using unpaired two-sample t-test. The

variation in proportion of male and hermaphrodite flowers was quantified using Kruskal-Wallis

test, and post hoc Dunn's test (Dinno 2017) was used to identify the significant differences in

proportion of genders between the four weeks. Simple linear regression was used to check the

effect of mean number of male flowers, hermaphrodite flowers and total flowers on the fruitset

respectively, and the effect of total floral display size on the mean visitation rate across the three

treatments in the choice experiments. The alpha was set at 0.5 (P<0.025) for all statistical

analysis. All graphical plots were produced using the ggplot2 package in R (Wickham 2016).

Results

Gender distribution in the population

Due to the small size and fragile nature of the flower, floral size was measured as the widths of

the flowers. The floral morphometry between the size of male and hermaphrodite flowers

showed no significant difference (t = 0.11072, df = 28, *p*-value = 0.9126; Appendix S1: Fig. S1).

Throughout the flowering season, male flowers were present in higher proportions than the

hermaphrodite flowers in all the study plots by at least 70-80% (Kruskal-Wallis chi-squared =

38.707, df = 7, *p*-value = 2.223^{e-06}; Fig. 2a; Appendix S1: Table S1). A slight decrease in the

proportion of male flowers was observed only in week 4 (Mean±SE: Week 1 = 1.33±0.31, Week

2 = 1.6±0.29, Week 3 = 1.73±0.35, Week 4 = 0.87±0.22; Fig. 2a; Appendix S1: Table S1). In

contrast, the number of hermaphrodite flowers remained similar across all four weeks of

observation (Mean±SE: Week 1 = 0.60±0.21 , Week 2 = 0.33±0.12 , Week 3 = 0.50±0.27 , Week

4 = 0.40±0.22 ; Fig. 2a; Appendix S1: Table S1) and only plants that were closer to the water

source i.e. stream population, had a significantly higher number of hermaphrodite flowers

(Mean±SE: Plateau_pop H = 0.43±0.07 vs. Stream_pop H = 1.75±0.14; *p*-values = 0.0000; Fig.

3; Appendix S1: Table S2). Overall, the mean number of total flowers remained consistent for

the first two weeks (Mean±SE: Week 1 = 1.93±0.39, Week 2 = 1.93±0.37 flowers; Fig. 2a inset),

increased in week 3 (2.23±0.41 flowers) and then reduced in week 4 (1.26±0.28 flowers;

Kruskal-Wallis chi-squared = 3.2847, df = 3, *p*-value = 0.3498; Appendix S1: Table S1).

Within an inflorescence, male and hermaphrodite flowers were present in all the three

positions i.e. apical, middle and basal positions (Fig. 1c), and the mean number of male flowers

varied across all these three positions during our four-week observation period. The number of

male flowers were relatively consistent in the basal position throughout the plants' flowering

season, than in the middle position or apical position (Fig. 2 b - e; Appendix S1: Table S3, S4).

We documented the highest number of male flowers in the middle position of the inflorescence,

whereas the number of hermaphrodite flowers was consistent at all positions over the flowering

season (Fig. 2 b - e; Appendix S1: Table S3, S4).

The total number of flowers was significantly higher in the stream population than in the

plateau population (Mean±SE: Plateau_pop = 2.78±0.12 vs. Stream_pop = 3.32±0.17; t = -

2.5646, df = 176.65, *p*-value = 0.0116; Fig. 3 inset) and the stream population also showed

similar proportions of male and hermaphrodite flowers (Mean±SE: Stream_pop H = 1.75±0.14

vs. Stream_pop M = 1.57±0.14; *p*-value = 0.1614; Fig. 3). Proportion of male flowers was

significantly greater in the plateau population over the stream population (Mean±SE:

Plateau_pop M = 2.35±0.12 vs. Stream_pop M = 1.57±0.14; *p*-value = 0.0000; Fig. 3; Appendix

S1: Table S2).

The mean fruit set per plant showed no effect with the total number of male,

hermaphrodite, or total number of flowers present per plant, respectively in the plateau_pop

(hermaphrodite: r^2 = -0.0457, df = 18, *p*-value = 0.6852; Appendix S1: Fig. S3a; male: r^2 = -

0.04999, df = 18, *p*-value = 0.761; Appendix S1: Fig. S3b; total flowers: r^2 = -0.04254, df = 18,

p-value = 0.6412; Appendix S1: Fig. S3c), and in the stream_pop (hermaphrodite: r^2 = -

0.0.05543, df = 18, p-value = 0.9637; Appendix S1: Fig. S3d; male: r^2 = 0.1125 , df = 18, p-

value = 0.08134; Appendix S1: Fig. S3e; total flowers: r^2 = 0.04828, df = 18, p-value = 0.1781;

Appendix S1: Fig. S3f), respectively.

Pollination biology

Flowers of both the genders opened between 12:30 hrs. to 16:00 hrs., and the mean pollen count

for both the genders ranged from 5000-10000 pollen grains per anther and no significant

difference in the pollen quantity between the two genders was observed (t = -0.67913, df = 60,

p-value = 0.4997; Azad 2018, Albal 2019, Nawge 2019). The major pollinators to *M. simplex*, in

the order of visitation rates from highest to lowest were *Apis dorsata* (Mean number of visits per

flower per hour±SE = 1.01±0.18), *Apis cerana* (0.86±0.10)*, Apis florea* (0.76±0.22), and

Amegilla spp. (Zonamegilla, 0.3±0.09) respectively. *Amegilla spp.* showed significantly lower

visitation than all the other pollinators (Kruskal-Wallis chi-squared = 6.5702, df = 3, p-value =

0.09; p-values: *Amegilla spp.* vs *A. cerana* = 0.0074, *Amegilla spp.* vs *A. dorsata* = 0.0058,

Amegilla spp. vs *A. florea* = 0.0469), whereas the visitation rates among *A. cerana, A. florea* and

A. dorsata were not significantly different (p-values: *A. cerana* vs *A. dorsata* = 0.3822, *A. cerana*

vs *A. florea* = 0.4278, *A. dorsata* vs *A. florea* = 0.3748; Azad 2018). Other pollinators observed

were *Paragus spp.* (Hoverfly), *Lasioglossum spp.*, *Pseudapis spp.* and two more species that we

could not identify, henceforth referred to as UnID sp. 3 and UnID sp. 4 (Appendix S1: Fig. S4).

Mean pollinator visitation rates were observed to be high at the beginning of the anthesis (1.74-

1.94 flowers from 13:00 hrs. to 15:00 hrs.) and it decreased with time (0.94-1.32 flowers from

15:00 hrs. till anthesis end; Azad 2018).

Gender specific pollinator preference experiment

In all the experimental setups in the manipulated choice experiment, pollinator visitations were

highest to the most abundant gender (Kruskal-Wallis chi-squared = 65.3103, df = 7, p-value =

1.303^{e-11}; Fig. 4; Appendix S1: Table S5). Pollinator visitation rates were significantly different

between hermaphrodite flowers in the unmanipulated, wild set up (control) and hermaphrodite

flowers in all the experimental setups (Fig. 4; Appendix S1: Table S5). However, the visitation

rates to the male flowers in the control were not different from those to male flowers in all the

experimental setup (Fig. 4; Appendix S1: Table S5). Display size (total number of flowers per

trial) had no effect on the visitation rates in any of the manipulated treatments (r^2 = -0.03765, df

= 21, p-value = 0.6309; Appendix S1: Fig. S5).

Quantification of pollinator-wise visitation rates in the manipulated experiments show

that visitations by *Apis cerana, Amegilla spp.* and UnID sp. 4 were higher to male flowers in

treatment 1 where male flowers were more abundant than hermaphrodite flowers (Mean

visitation rate per flower per hr±SE:- H < M: *A. cerana* = 3.29±1.11, *Amegilla spp.* = 1.81±0.91,

UnID sp. 4 = 2.49±1.89 flowers; Kruskal-Wallis chi-squared = 13.8517, df = 9, p-value =

0.1277; Fig. 5a; Appendix S1: Table S6). In treatment 3, *Apis cerana*, *Amegilla spp.*, and UnID

sp. 4 showed high visitation to hermaphrodite flowers (H > M: *A. cerana* = 2.76±0.74, *Amegilla*

spp. = 0.89±0.61, UnID sp. 4 = 0.66±0.56 flowers; Kruskal-Wallis chi-squared = 9.2848, df = 5,

p-value = 0.09823; Fig. 5c; Appendix S1: Table S6). We observed visitation by hoverflies only

in treatment 2 (H = M: Visitation rate per hermaphrodite flower = 0.56, visitation rate per male

flower = 0.75; Kruskal-Wallis chi-squared = 15.605, df = 11, p-value = 0.1565; Fig. 5b;

Appendix S1: Table S6).

Discussion

Gender distribution in the population

The optimal allocation hypothesis proposes that between a costly and a cheaper gender, in a

resource-limited environment, the gender which requires the least resources will be preferred.

From a plant's perspective, the male flowers are considered a cheaper gender to produce because

there is no investment towards fruit development, unlike the hermaphrodite flowers which

produces both viable pollen and ovules (Bertin 1982a, Anderson and Symon 1989, Kaul et al.

2002, Narbona et al. 2002, Obeso 2002, Verdú et al. 2007, Zhang and Tan 2009). Here, we show

that the *M. simplex* population on Kaas plateau has a higher density of male flowers and a slight

reduction in the number of male flowers and an increase in hermaphrodite flowers were observed

only among the individuals present along the stream. We consider the stream population of *M.*

simplex to have better resources available to them than the plateau population because of

presence of water, possible higher nutrient availability due to dissolved/associated nutrients in

the water, and deeper soil substrate for the plants (Thorpe et al. 2018). This suggests that

although the general tendency of *M. simplex* is to produce more male flowers over

hermaphrodites, water and water-dissolved minerals may be major limiting resources and in the

absence of resource constraints the plants can produce not only more flowers but also higher

number of hermaphrodite flowers.

Gender shift with time

Resources available and utilized by plants often fluctuate with time and age of the plant

(Stephenson 1982, Solomon 1985). We observed an initial increase in the number of flowers

followed by a decreasing trend in the total number of flowers being produced per plant

throughout its flowering period (Fig. 2a inset). Within this trend, male flowers were always

produced in higher proportions than the hermaphrodite flowers and interestingly, in the end of

the flowering season (i.e. fourth week) we observed a reduction in the number of male flowers

instead of the number of hermaphrodite flowers (Fig. 2a). Thus, although resource constraints do

affect the production of total flowers as observed in the stream vs. plateau population, resource

limitation due to the flowering stage of the plant only affected the number of male flowers in *M.*

simplex. It has been shown that under declining resource levels plants may opt to increase their

fitness by producing the relatively inexpensive male flowers over the expensive female flowers

(May and Eugene Spears Jr 1988). Further, it has been reported in *C. benghalensis* that a plant

will opt to drop a flower rather than dropping a fruit, under resource limiting conditions, because

the nutrient resources that would get committed to fruits that is aborted would be saved halfway

(Kaul and Koul 2008). We propose that a similar strategy may be operational in *M. simplex*

wherein under availability of resources *M. simplex* may be producing excess of male flowers

which are the first ones to be sacrificed when resource limitation starts to be an important factor

as the phenology of the plant progresses (Fig. 2a). The lack of change in the number of

hermaphrodite flowers in different stages of plant's flowering phenology also suggests that the

hermaphrodite flowers may already be produced at an optimal amount, resulting in the

maintenance of their number despite declining resources at the later stages (Fig. 2a).

Gender distribution of flowers on an inflorescence has been shown to be dependent on

resource allocation (Diggle 1997, Miller and Diggle 2003), and it has been observed that the

differential positioning of the genders within an inflorescence can affect outcrossing rates (Orth

and Waddington 1997, Lin and Forrest 2019). Studies in *Solanum* (Diggle 1994, 1997, Miller

and Diggle 2003), *Caesalpinia* (Lewis and Gibbs 1999), *F. virginiana* (Ashman and Hitchens

2000) and in other Commelinaceae genera (Kaul et al. 2002, Kaul and Koul 2008) have shown

that higher accessibility to nutrient resources at the base of an inflorescence (than the distal end)

can result in the presence of higher resource-demanding hermaphrodite flowers in the basal

position while males are relegated to the middle and apical positions of an inflorescence. We

found a similar pattern in *M. simplex,* where a greater number of male flowers were produced at

the middle and the apical positions on an inflorescence. However, unlike other plants where

hermaphrodites exclusively occupy the basal positions in an inflorescence, in *M. simplex*

hermaphrodites were also present in the apical position. In fact, the distribution of hermaphrodite

flowers was not significantly different across all the three positions, over the course of the plant's

flowering season. This implies that any manipulation in display size is carried out by increasing

or decreasing the male flowers thus regulating the floral display at a much lesser cost (as

proposed in *Solanum* by Anderson and Symon 1989).

Pollination biology

Pollinators have been shown to display higher preference to larger inflorescences, ultimately

resulting in higher reproductive success (Stephenson 1981, Harder and Barrett 1995, Ohashi and

Yahara 2001). Hence the disproportionate gender distribution within the inflorescence of an

andromonoecious species may be a result of indirect pollinator selection of inflorescence size:

that is larger inflorescences are better at attracting pollinators but under nutrient resource

limitation, this can be achieved only by producing more male flowers, which are cheaper to

produce (Sandring and Ågren 2009). Thus our results support the pollinator attractor hypothesis

as proposed by Zhang and Tan (2009) and sex allocation theory as proposed by Elle and

Meagher (2000) where they predicted that male flowers can increase the floral display and their

attractiveness to pollinators at a lower cost through male function than female function and

thereby increase the stigmatic load of the hermaphrodite flowers, which potentially increases

female reproductive success at lower resource investment.

The dominant pollinator assemblage visiting *M. simplex* consisted mainly of different

species of bees from the two genera common in the Indian tropics - *Apis* and *Amegilla*. All

pollinators that visit *M. simplex* are generalists and are known to be common pollinators in

grassland habitats (Corlett 2011, Dhargalkar 2019, Nawge 2019). There are at least 68 plants co-

flowering with *M. simplex* on the Kaas plateau (Dhargalkar 2019) and parallel studies on the

pollination of plants in the Kaas community showed that both *Apis cerana* and *Apis florea* switch

to *M. simplex* from *M. lanuginosa* (anthesis time 10.30 hrs.-13.00 hrs.) and various *Impatiens*

species as soon as the flowers of *M. simplex* open i.e. at 12:30 hrs. up to 16:00 hrs. (Albal 2019).

This suggests that although the pollinators are generalists, these pollinators may be displaying

temporal specialization towards *M. simplex*. Future studies will focus on temporal specialization

of the bee communities on Kaas plateau, which will provide insight on pollinator preference

towards *M. simplex* in a multi-species environment.

Gender-specific pollinator preference

Several studies have shown that pollinators avoid functionally female flowers because of

decreased pollen or absence of pollen (Bierzychudek 1987, Eckhart 1991), floral size (Ashman et

al. 2000), floral scents (Ashman et al. 2005) and pollen-pistil interference (Solomon 1985, Elle

and Meagher 2000). However, in *M. simplex* both male and hermaphrodite flowers showed

comparable pollen production. This has been shown in other andromonoecious systems as well

such as in *Solanum carolinense* (Solomon 1986), *Olea europaea* (Cuevas and Polito 2004) and

Dichorisandra spp. (Sigrist and Sazima 2015). As previously discussed, we observed that

nutrient resources affected the number of male flowers and not the number of hermaphrodite

flowers, and the rate of fruit sets were observed to be similar between the plateau and the stream

populations (Mean±SE: Plateau_pop = 3.22±0.23 vs. Stream_pop = 3.55±0.21). This suggests

that reproductive success is limited by the availability of hermaphrodite flowers and not

pollinator visitations. The lack of preference for either of the gender by the pollinators on Kaas

plateau is further supported by our observations from the manipulative experiment where all

pollinators except *A. cerana* displayed lack of preference towards either of the floral genders.

Further, this also fails to support the hypothesis that male-dominant inflorescences of *M. simplex*

may be under pollinator-mediated selection due to direct pollinator preference for any gender

(see Solomon 1985, Eckhart 1991, Elle and Meagher 2000). However, since pollinator visitations

were observed to be density dependent (i.e. floral density, compare treatment 3 with treatment 1,

Fig. 5c) it is possible that pollinator-mediated selection is acting on male flowers through

selection of larger inflorescence sizes.

Our understanding of pollinator services by bees primarily come from *Bombus* species

(Chittka et al. 1997) and *Apis mellifera* (Martin 2004) and studies on other bees from India are

very rare (see Somanathan and Borges 2001). At least five species of bees visited *M. simplex* and

in the choice experiment, we observed inter-species behavioral differences among bees both in

their floral preferences and visitation patterns, respectively. Almost all bees displayed higher

visitation to any gender which was present in greater density, suggesting that pollinator

visitations are predominantly gender-nonspecific and density-dependent. A slight preference for

gender was only observed in *A. cerana* and *Amegilla* in treatment 2 (H = M) where male flowers

were preferred over hermaphrodite flowers (Fig. 5b). Bees are known for both their short-term

and long-term memory retention and they can show learning through foraging experience

(Cresswell and Galen 1991, Schiestl and Johnson 2013) and we propose that the higher visitation

to male flowers in treatment 2 (Fig. 5b) may be a result of short-term memories and habituation

in bees since in natural conditions bees may encounter more male flowers than hermaphrodites.

We conclude that in *M. simplex* environmentally driven resource constraints play a bigger

role than pollinator preference towards a specific gender in determining gender expression.

However, we cannot dismiss an indirect role of pollinator-driven constraints which may occur

through selection on the display size as proposed in the pollinator attractor hypothesis (Zhang

and Tan 2009). In a resource-limited environment such as the one observed at Kaas plateau, *M.*

simplex can thus increase its display size by producing the less expensive male flowers over

hermaphrodite flowers.

Energy constraints will have a direct cost on floral production if genders require

differential resource allocation. As a result, it can be expected that the gender of the flowers will

be selected accordingly. Using the *M. simplex* system, we show that gender expression is directly

influenced by the availability of nutrient resources in the plants' environment and indirectly

affected by the pollinators' preference for larger display sizes of the inflorescence. Thus, a

combination of both environmental factors as well as ecological factors may be responsible for

the evolution and maintenance of andromonoecy in mass-flowering plants such as *M. simplex* in

the tropics.

Acknowledgements

The authors thank the Ministry of Human Resource Development, India (MHRD) and the

National Geographic Society for research funds to VG, IISER Bhopal for the infrastructure, DST

for INSPIRE fellowship to AA, AG, and SS. We thank the PCCF of Maharashtra Forest

Department, DFO of Satara Division (Forest Department) and the Joint Forest Management

Committee (JFMC), Kaas, Satara for permitting our fieldwork. We also thank Dr. Natapot Warrit

(Chulalongkorn University, Thailand) and Dr. Ximo Mengual (Research Museum Alexander

Koenig, Germany) for their help with taxonomic identification of the pollinators. Finally, we

thank all fellow TrEE lab members for their advice, discussion, and support.

Author contributions

VG conceived the idea, AA, AG, and VG designed the experiments. AA and AG conducted the

fieldwork with help from SS. AA performed the statistical analyses with inputs from SS and VG.

AA and VG wrote the manuscript and AG and SS helped with edits. All authors read and

approved the final draft of the manuscript.

Literature cited

Albal, A. 2019. When does a flower become male? Exploring the role of ecological factors and

energy resources in defining gender expression in sympatric plants. Master's thesis, IISER

Bhopal, India R00529:14030.

Anderson, G. J., and D. E. Symon. 1989. Functional dioecy and andromonoecy in *Solanum*.

Evolution 43:204.

Armbruster, W. S. 2001. Evolution of floral form: Electrostatic forces, pollination, and adaptive

compromise. New Phytologist 152:181–186.

Ashman, T. L., M. Bradburn, D. H. Cole, B. H. Blaney, and R. A. Raguso. 2005. The scent of a

male: The role of floral volatiles in pollination of a gender dimorphic plant. Ecology

86:2099–2105.

Ashman, T. L., and M. S. Hitchens. 2000. Dissecting the causes of variation in intra-

inflorescence allocation in a sexually polymorphic species, *Fragaria virginiana* (Rosaceae).

American Journal of Botany 87:197–204.

Ashman, T. L., and M. T. Morgan. 2004. Explaining phenotypic selection on plant attractive

characters: Male function, gender balance or ecological context? Proceedings of the Royal

Society B: Biological Sciences 271:553–559.

Ashman, T. L., J. Swetz, and S. Shivitz. 2000. Understanding the basis of pollinator selectivity in

sexually dimorphic *Fragaria virginiana*. Oikos 90:347–356.

Azad, G. 2018. Resource allocation and pollinator preference: role of male flowers in the

andromonoecious herb *Murdannia simplex* (Commelinaceae). Master's thesis, IISER

Bhopal, India R00335:13035.

Benseler, R. W. 1975. Floral biology of California buckeye. Madroño 23:41–53.

Bertin, R. I. 1982a. The evolution and maintenance of andromonoecy. Evolutionary Theory

6:25–32.

Bertin, R. I. 1982b. The ecology of sex expression in red buckeye. Ecology 63:445–456.

Bierzychudek, P. 1987. Pollinators increase the cost of sex by avoiding female flowers. Ecology

68:444–447.

Charnov, E., and J. Bull. 1977. When is sex environmentally determined? Nature 266:828–832.

Chittka, L., A. Gumbert, and J. Kunze. 1997. Foraging dynamics of bumble bees: Correlates of

movements within and between plant species. Behavioral Ecology 8:239–249.

Christopher, D. A., R. J. Mitchell, D. W. Trapnell, P. A. Smallwood, W. R. Semski, and J. D.

Karron. 2019. Hermaphroditism promotes mate diversity in flowering plants. American

Journal of Botany 106:1131–1136.

Corlett, R. T. 2011. Honeybees in natural ecosystems. In: Hepburn, H. and Radloff, S.E. (Eds)

Honeybees of Asia. Springer- Verlag Berlin Heidelberg Pages 215-225

Cresswell, J. E., and C. Galen. 1991. Frequency-dependent selection and adaptive surfaces for

floral character combinations: The pollination of *Polemonium viscosum*. The American

Naturalist 138:1342–1353.

Cuevas, J., and V. S. Polito. 2004. The role of staminate flowers in the breeding system of *Olea*

europaea (Oleaceae): An andromonoecious, wind-pollinated taxon. Annals of Botany

93:547–553.

Dhargalkar, R. K. 2019. A study of the temporal patterns in flowering phenology and plant-

pollinator network of a grassland community in the northern Western Ghats, Maharashtra.

Master's thesis, IISER Bhopal, India R00534:14054.

Diggle, P. K. 1993. Developmental plasticity, genetic variation, and the evolution of

andromonoecy in *Solanum hirtum* (Solanaceae). American Journal of Botany 80:967–973.

Diggle, P. K. 1994. The expression of andromonoecy in *Solanum hirtum* (Solanaceae):

Phenotypic plasticity and ontogenetic contingency. American Journal of Botany 81:1354–

1365.

Diggle, P. K. 1997. Ontogenetic contingency and floral morphology: The effects of architecture

and resource limitation. International Journal of Plant Sciences 158:99–107.

Dinno, A. 2017. dunn.test: Dunn's test of Multiple comparisons using rank sums.

Eckhart, V. M. 1991. The effects of floral display on pollinator visitation vary among

populations of *Phacelia linearis* (Hydrophyllaceae). Evolutionary Ecology 5:370–384.

Elle, E., and T. R. Meagher. 2000. Sex allocation and reproductive success in the

andromonoecious perennial *Solanum carolinense* (Solanaceae). II. Paternity and functional

gender. American Naturalist 156:622–636.

Faden, R. B. 2000. Floral biology of Commelinaceae. Pages 309–317 Monocots: Systematics

and Evolution.

Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004.

Pollination Syndromes and Floral Specialization. Annual Review of Ecology, Evolution,

and Systematics 35:375–403.

Harder, L. D., and S. C. H. Barrett. 1995. Mating costs of large floral displays in hermaphrodite

plants. Nature 373:512–515.

Kaul, V., and A. K. Koul. 2008. Floral phenology in relation to pollination and reproductive

output in *Commelina caroliniana* (Commelinaceae). Australian Journal of Botany 56:59–

66.

Kaul, V., N. Sharma, and A. K. Koul. 2002. Reproductive effort and sex allocation strategy in

Commelina benghalensis L., a common monsoon weed. Botanical Journal of the Linnean

Society 140:403–413.

Lewis, G., and P. Gibbs. 1999. Reproductive biology of *Caesalpinia calycina* and *C. pluviosa*

(Leguminosae) of the Caatinga of north-western Brazil. Plant Systematics and Evolution

217:43–53.

Lin, S., and J. R. K. Forrest. 2019. The function of floral orientation in bluebells: interactions

with pollinators and rain in two species of *Mertensia* (Boraginaceae). Journal of Plant

Ecology 12:113–123.

Lloyd, D. G. 1980. Sexual strategies in Plants: I. A hypothesis of serial adjustment of maternal

investment during one reproductive session. New Phytologist 86:69–79.

Martin, N. H. 2004. Flower size preferences of the honeybee (*Apis mellifera*) foraging on

Mimulus guttatus (Scrophulariaceae). Evolutionary Ecology Research 6:777–782.

May, P. G., and E. Eugene Spears Jr. 1988. Andromonoecy and variation in phenotypic gender

of *Passiflora incarnata* (Passifloraceae). American Journal of Botany 75:1830–1841.

Miller, J. S., and P. K. Diggle. 2003. Diversification of andromonoecy in *Solanum* section

Lasiocarpa (Solanaceae): The roles of phenotypic plasticity and architecture. American

Journal of Botany 90:707–715.

Nandikar, M. D., and R. V. Gurav. 2015. Revision of the genus *Murdannia* (Commelinaceae) in

India. Phytodiversity 2:56–112.

Narbona, E., P. L. Ortiz, and M. Arista. 2002. Functional Andromonoecy in *Euphorbia*

(Euphorbiaceae). Annals of Botany 89:571–577.

Nawge, V. V. 2019. Understanding pollination networks on Kaas plateau. Master's thesis, IISER

Bhopal, India R00543:14097.

Obeso, J. R. 2002. The costs of reproduction in plants. New Phytologist 155:321–348.

Ohashi, K., and T. Yahara. 2001. Behavioural responses of pollinators to variation in floral

display size and their influences on the evolution of floral traits. In: Chittka, L., Thomson,

J.D. Cognitive Ecology of Pollination- Animal behaviour and floral evolution. Cambridge

University Press Pages 274-296.

Orth, A. I., and K. D. Waddington. 1997. The movement patterns of carpenter bees *Xylocopa*

micans and bumblebees *Bombus pennsylvanicus* on *Pontederia cordata* inflorescences.

Journal of Insect Behavior 10:79-86.

Ortiz, P. L., M. Arista, P. E. Oliveira, and S. Talavera. 2003. Pattern of flower and fruit

production in *Stryphnodendron adstringens*, an andromonoecious legume tree of Central

Brazil. Plant Biology 5:592–599.

Oziegbe, M., I. O. Akinlua, and A. A. Olalekan. 2013. Comparative pollination role of stamens

and breeding system in three species of *Commelina* (Commelinaceae) in Ile-Ife, Nigeria.

Acta Botanica Brasilica 27:543–550.

Primack, R. B., and D. G. Lloyd. 1980a. Andromonoecy in the New Zealand montane shrub

Manuka, *Leptospermum scoparium* (Myrtaceae). American Journal of Botany 67:361–368.

Primack, R. B., and D. G. Lloyd. 1980b. Sexual strategies in plants IV. The distributions of

gender in two monomorphic shrub populations. New Zealand Journal of Botany 18:109–

114.

R Team, C. 2019. R: A language and environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria. https://www.r-project.org/

Sandring, S., and J. Ågren. 2009. Pollinator-mediated selection on floral display and flowering

time in the perennial herb *Arabidopsis lyrata*. Evolution 63:1292–1300.

Schiestl, F. P., and S. D. Johnson. 2013. Pollinator-mediated evolution of floral signals. Trends

in Ecology and Evolution 28:307–315.

Sigrist, M. R., and M. Sazima. 2015. Phenology, reproductive biology and diversity of buzzing

bees of sympatric *Dichorisandra* species (Commelinaceae): breeding system and

performance of pollinators. Plant Systematics and Evolution 301:1005–1015.

Solomon, B. P. 1985. Environmentally influenced changes in sex expression in an

andromonoecious plant. Ecology 66:1321–1332.

Solomon, B. P. 1986. Sexual allocation and andromonoecy: Resource investment in male and

hermaphrodite flowers of *Solanum carolinense* (Solanaceae). American Journal of Botany

73:1215–1221.

Somanathan, H., and R. M. Borges. 2001. Nocturnal pollination by the carpenter bee *Xylocopa*

tenuiscapa (Apidae) and the effect of floral display on fruit set of *Heterophragma*

quadriloculare (Bignoniaceae) in India. Biotropica 33:78–89.

Stanton, M. L., J. K. Bereczky, and H. D. Hasbrouck. 1987. Pollination thoroughness and

maternal yield regulation in wild radish, *Raphanus raphanistrum* (Brassicaceae). Oecologia

74:68–76.

Stephenson, A. G. 1981. Flower and fruit abortion: proximate causes and ultimate functions.

Annual Review of Ecology and Systematics 12:253–279.

Stephenson, A. G. 1982. When does outcrossing occur in a mass-flowering plant? Evolution

36:762–767.

Thorpe, C. J., T. R. Lewis, S. Kulkarni, A. Watve, N. Gaitonde, D. Pryce, L. Davies, D. T.

Bilton, and M. E. Knight. 2018. Micro-habitat distribution drives patch quality for sub-

tropical rocky plateau amphibians in the northern Western Ghats, India. PLoS ONE 13:1–

20.

Ushimaru, A., T. Watanabe, and K. Nakata. 2007. Colored floral organs influence pollinator

behavior and pollen transfer in *Commelina communis* (Commelinaceae). American Journal

of Botany 94:249–258.

Vallejo-Marín, M., and M. D. Rausher. 2007a. The role of male flowers in andromonoecious

species: Energetic costs and siring success in *Solanum carolinense* L. Evolution 61:404–

412.

Vallejo-Marín, M., and M. D. Rausher. 2007b. Selection through female fitness helps to explain

the maintenance of male flowers. The American Naturalist 169:563.

Venkatesan, A. 2017. Why be a male? Resource allocation and floral display in a wild

population of *Solanum surattense*. Master's thesis, IISER Bhopal, India R00064:10015.

Verdú, M., K. Spanos, I. Čaňová, B. Slobodník, and L. Paule. 2007. Similar gender dimorphism

in the costs of reproduction across the geographic range of *Fraxinus ornus*. Annals of

Botany 99:183–191.

Wickham, H. 2016. ggplot2: Elegant graphics for Data Analysis. Springer-Verlag NY.

https://ggplot2.tidyverse.org

Wickham, H., François, R., Henry, L., and Müller, K. (2020). dplyr: A grammar of data

manipulation. R package version 0.8.4. https://cran.r-project.org/package=dplyr

Williams, G., and K. Walker. 2003. Pollination of the wet forest herb *Pollia crispata*

(Commelinaceae). Cunninghamia 8(1):141–146.

Wilson, P., J. D. Thomson, M. L. Stanton, and L. P. Rigney. 1994. Beyond floral Batemania:

Gender biases in slection for pollination success. The American Naturalist 143:283-296.

Zhang, T., and D. Y. Tan. 2009. An examination of the function of male flowers in an

andromonoecious shrub *Capparis spinosa*. Journal of Integrative Plant Biology 51:316–

324.

Figure legends

Fig. 1 Habitat and floral features of *Murdannia simplex* (a) habitat on Kaas Plateau, Maharashtra

during flowering season (Inset: Location in India), (b) *M. simplex* habit, (c) inflorescence image

showing positions identified from base to the apex, (d) *M. simplex* flower front view, (e) *M.*

simplex male flower, (f) *M. simplex* hermaphrodite flower. Photo courtesy: (a), (c), (d) - Asawari

Albal, (b), (e), (f) - Saket Shrotri.

Fig. 2 Variation in gender in *M. simplex* (a) by week from first flowering date (Kruskal-Wallis

chi-squared = 38.7067, df = 7, *p*-value = 0.00; P<0.025) (Inset: Week-wise proportion of total

number of flowers; Kruskal-Wallis chi-squared = 3.2847, df = 3, *p*-value = 0.3498, N = 30;

P<0.025) (See Appendix S1: Table S1) and (b) - (e) within-inflorescence from Week 1 - Week 4

respectively (Kruskal-Wallis chi-square test; df = 5, P<0.025). For (a) columns to be compared

within the week and within gender (Mean±SE). For (b) columns to be compared within the

gender and within positions, across all 4 weeks (Mean±SE). Significant *p*-values depicted by

different alphabets (Appendix S1: Table S3; P<0.025).

Fig. 3 Natural gender distribution observed in *M. simplex* in 2019 between two populations

(Mean±SE) (Kruskal-Wallis chi-squared = 123.53, df = 3, *p*-value <$2.2^{e\text{-}16}$, N = 100 per

population; Appendix S1: Table S2; P<0.025) (Inset: Total number of flowers across the two

populations; t = -2.5646, df = 176.65, *p*-value = 0.0116). Male flowers are represented by open

squares, hermaphrodite flowers represented by filled squares and the gray squares represent total

(male + hermaphrodite) flowers. Columns to be compared within population and within genders

across the two populations only.

Fig. 4 Mean pollinator visitation rate per flower per hour (Mean±SE) observed in the

manipulative experiment where three treatments with varying numbers of hermaphrodite and

male flowers were used to test pollinator preference towards a gender; (Kruskal-Wallis chi-

squared = 63.4196, df = 7, *p*-value = 3.122^{e-11}, N = 8 trials for each treatment; P<0.025). Male

flowers are represented by open squares and hermaphrodite flowers are represented by the filled

squares. The first two columns represent the natural visitation rate observed per flower per hour

for both male and hermaphrodite, respectively. Different alphabets depict significantly different

values (Appendix S1: Table S5; P<0.025).

Fig. 5 Pollinator-wise mean visitation rate per flower per hour in the manipulative experiment on

M. simplex (Mean±SE) (a) Treatment 1- H < M (Kruskal-Wallis chi-squared = 13.8517, df = 9,

p-value = 0.1277; P<0.025; Appendix S1: Table S6), (b) Treatment 2- H = M (Kruskal-Wallis

chi-squared = 15.605, df = 11, *p*-value = 0.1565; P<0.025; Appendix S1: Table S6),

(c) Treatment 3- H > M (Kruskal-Wallis chi-squared = 9.2848, df = 5, *p*-value = 0.09823;

P<0.025; Appendix S1: Table S6). Male flowers are represented by open squares and

hermaphrodite flowers are represented by filled squares. Different alphabets depict significantly

different values (Appendix S1: Table S6; P<0.025).

Figures

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5