1   When to be a male? Role of resource-limitation and pollinators in determining gender in an

2   andromonoecious spiderwort.

6   Asawari Albal [1,3,4], Azad G[1], Saket Shrotri[1], Vinita Gowda[1,2]

[1] Tropical Ecology and Evolution (TrEE) Lab, Department of Biological Sciences, Indian

Institute of Science Education and Research Bhopal, Bhopal, 462066, India

[2] Corresponding author: Vinita Gowda, gowdav@iiserb.ac.in

Present address:

[3] Department of Biology, University of Toronto Mississauga, Mississauga, ON L5L 1C6,

Canada

[4] Ecology and Evolutionary Biology, University of Toronto, Toronto, ON M5S 2Z9, Canada

Abstract

The evolution and maintenance of sexual systems in plants is often driven by resource

allocation and pollinator preferences. Gender or sex in plants, therefore, represents a

physiological, functional, and behavioral response of a plant to its environment. Annual,

entomophilous plants have three major constraints towards optimal reproduction: 1) nutrient

resources obtained from the environment, 2) nutrient resources allocated towards

reproduction, i.e., fruits vs. flowers, and 3) pollinator visitations. We know very little about

the role of nutrient resources and pollinators in driving gender expression in flowers in a

natural environment.

Andromonoecy is a sexual system where the plant bears both staminate and

hermaphrodite flowers on the same inflorescence and it occurs in ~2% of flowering plants.

Although it has been shown that andromonoecy has evolved multiple times within the

angiosperms, our understanding of the underlying mechanisms of what determines gender

expression in nature is limited. The optimal resource allocation hypothesis, one of the most

widely accepted hypotheses which describes evolution of andromonoecy, suggests that under

nutrient constraints, plants will produce more male flowers since they require lesser energy

investment than the more expensive hermaphrodite flowers. We test this hypothesis in an

andromonoecious plant *Murdannia simplex* (Commelinaceae) by measuring gender

distributions in a large natural population and contrasting the gender distributions between

two different resource conditions (stream population vs. plateau population). We also carried

out choice experiments to quantify if pollinators showed any preference towards a gender.

We found that in *M. simplex*, production of hermaphrodite flowers is resource-

dependent and under resource constraints not only fewer numbers of flowers were produced

but also there were a higher number of male flowers. We failed to observe pollinator

preference towards either gender but *Amegilla spp. (Zonamegilla)* and *Apis cerana* showed

frequency-dependent visitation which suggests that selection on inflorescence size may be an

indirect form of selection pressure exerted by pollinators resulting in gender-bias in the

plants. Thus, we conclude that environmentally driven resource constraints play a bigger role

in driving floral gender expression in *Murdannia* over direct pollinator-driven constraints.

Keywords

Andromonoecy, Commelinaceae, floral gender, male, hermaphrodite, *Murdannia simplex*,

resource limitation, pollinator preference, Western Ghats

Introduction

Andromonoecy refers to the sexual system where both staminate (male) flowers and perfect

(hermaphrodite) flowers are present on the same plant (Yampolsky and Yampolsky 1922,

Bertin 1982a, Miller and Diggle 2003, Vallejo-Marín and Rausher 2007a, 2007b). In

andromonoecious plants, the hermaphrodite flower fulfills the female function in the absence

of a true 'female' flower, and it has been suggested that andromonoecy probably evolved

from hermaphroditism by the loss of female reproductive structure (Lloyd 1980, Bertin

1982a). One of the most widely accepted hypotheses on the evolution of andromonoecy is the

optimal resource allocation hypothesis which suggests that under resource limitation, male

flowers will be produced instead of hermaphrodite flowers. This hypothesis rests upon the

premise that male flowers are energetically cheaper to produce because reproductive

allocation towards them ends with pollen production (Anderson and Symon 1989, Narbona et

al. 2002, Verdú et al. 2007), while females are expensive because fruit development and seed

germination are energy-intensive physiological processes (Bertin 1982a, Kaul et al. 2002,

Obeso 2002, Verdú et al. 2007).

Examples of reallocation of resources to the cheaper sex (i.e. male bias) have been

shown in several plant species where staminate flowers are produced when nutrient resources

become scarce due to its utilization by the costlier gynoecia towards fruit production (Bertin

1982a, 1982b, May and Eugene Spears Jr 1988, Miller and Diggle 2003, Venkatesan 2017).

A well-studied species is *Solanum hirtum* (Solanaceae) in which it has been shown that as a

result of successful pollination, as the ovary develops, resources are relocated within an

inflorescence to the male flowers, thus influencing gender expression and distribution of

gender in successive flowers (Diggle 1993, Miller and Diggle 2003). Another example comes

from *Raphanus raphanistrum*, where Stanton et al. (1987) showed that in plants which were

heavily pollinated, the number of ovules per flower decreased and in contrast, unpollinated

plants did not show any significant decrease in number of ovules. This suggests that plants

can modify their reproductive output to adjust for the resource limitation faced by them.

In plants, resource limitation or nutrient limitation that can affect floral gender

expression can be identified as: a) environmental nutrient limitation (Charnov and Bull 1977,

Primack and Lloyd 1980a, Stephenson 1981, Bertin 1982a, Solomon 1985, Diggle 1993),

and/or b) within-plant nutrient limitation (Stephenson 1981, May and Eugene Spears Jr 1988,

Diggle 1993, 1995, Miller and Diggle 2003, Ortiz et al. 2003, Vallejo-Marín and Rausher

2007a). In *Aesculus californica*, *A. pavia*, *Leptospermum scoparium* and *Passiflora*

incarnata, phenotypic gender expression (hermaphroditic inflorescences or hermaphroditic

flowers) has been shown to vary as a response to environmental conditions or status of the

nutrient resources, especially during fruit development when resources are limited (Benseler

1975, Primack and Lloyd 1980b, May and Eugene Spears Jr 1988). Further, it has been

shown that the amount of resources that a plant acquires in the proximal and distal ends of the

inflorescence can also vary. The proximal/ basal end of the inflorescence has more resources

than its distal end, which results in more hermaphrodite flowers to be present at basal ranks

of an inflorescence, while male flowers are relegated to the distal end (Faden 1991, Diggle

1994, 1995, 1997, Lewis and Gibbs 1999, Ashman and Hitchens 2000, Miller and Diggle

2003, Kaul and Koul 2008).

Bateman's principle (Bateman 1948, Wilson et al. 1994) asserts that- "Fitness gain

through male function is limited primarily by mating opportunity, while fitness gain through

female function is limited primarily by resource availability for offspring production". To

explain the role of male flowers in andromonoecious plants, two, not mutually exclusive

hypotheses have been proposed (Zhang and Tan 2009): i) Pollen donor hypothesis

approaches the problem from the perspective of male reproductive success. It predicts that the

wastage of nutrient resources by investment in the female reproductive structure is reduced

by the production of only male flowers, in a resource-limited condition. ii) Pollinator attractor

hypothesis is based on female reproductive success. It predicts that male flowers increase the

floral display and attract more pollinators at a lower cost and thereby increase the stigmatic

load of the hermaphrodite flowers, which potentially increases female reproductive success at

lower resource investment. The sex allocation theory predicts that by attracting pollinators

which promote outcrossing, floral attractiveness (e.g., large petals, nectar availability) gains

more fitness through the male function than female function (Elle and Meagher 2000).

Although male success should be correlated with mating opportunities, one or a few visits by

pollinators are known to be adequate to fertilize all ovules. Therefore, it is not expected that

female fitness would be strongly affected by floral attractiveness. Thus, in sexually

reproducing plants, investment in the production of flowers is very important since this can

directly translate into reproductive success, and hence reproductive fitness. In entomophilous

plants, pollinator dependence is often driven by floral characteristics, and for successful

pollination, the gender of the flowers and associated floral characters may be very important

(Armbruster 2001, Fenster et al. 2004). In many hermaphroditic plant systems, it has been

shown that several characteristics of an inflorescence such as number of flowers, size of

flowers, and reward of flowers play a critical role in attracting pollinators to the plant (Ortiz

et al. 2003, Christopher et al. 2019). While the role of pollinators is very well known in

hermaphroditic systems, the perception of gender, pollinator preference, and reproductive

output in andromonoecious systems are not well studied (see Diggle 1993, Ashman et al.

2000, Ashman and Morgan 2004).

Previous studies in sexually dimorphic plants have shown that pollinators typically

show a preference for the functionally male flowers over the conspecific functional female

flowers owing to different floral traits like petal length (Ashman et al. 2000), levels of pollen

production (Bierzychudek 1987, Eckhart 1991) and/or floral scents (Ashman et al. 2005),

where these preferences are learned through foraging experience (Cresswell and Galen 1991).

Zhang and Tan (Zhang and Tan 2009) were the first to examine the function of male flowers

in the andromonoecious *Capparis spinosa* where they showed that the pollinators did not

discriminate between the floral gender morphs, even when they were morphologically

distinct. However, similar studies investigating gender-specific pollinator preferences and

more specifically gender-specific pollinator preferences in sexually dimorphic plants that

have near-identical morphological features are not known.

In this study, we explore the effect of nutrient resources as well as pollinator

preference in driving gender expression in a wild population of the andromonoecious

Murdannia simplex (Commelinaceae, Spiderworts). The pantropical plant family

Commelinaceae (Spiderworts) is known to have several genera that display andromonoecy.

Despite the dominance of andromonoecy in this family, to date, most pollination studies

within this family have focused only on identifying pollinators (Kaul et al. 2002, Williams

and Walker 2003, Oziegbe et al. 2013, Sigrist and Sazima 2015), and only one study has

explored the role of floral guides in pollinator attraction (Ushimaru et al. 2007). Based on

predictions from the optimal resource allocation hypothesis and pollinator attractor

hypothesis we address the following questions:

A) Does the proportion of staminate and perfect flowers in a population represent resource

limitation?

B) Do pollinators show preference to any one of the genders?

C) Do staminate flowers improve floral display to maintain visitation rate and function as a

cheaper source of outcrossing pollen?

Materials and methods

Murdannia simplex (Vahl) Brenan (Commelinaceae) is a mass-flowering, sub-erect, annual,

andromonoecious herb which is widely distributed in moist deciduous forests and grasslands

of the Western Ghats of India (Nandikar and Gurav 2015; Fig. 1a). The plant is seasonal, and

the shoot emerges once the monsoonal rains begin, and the flowering lasts from September to

November (Nandikar and Gurav 2015). The plant is 40-65 cm tall and bears a cymose

inflorescence which is a panicle. The inflorescence bears 5-20 purple-colored, three-petaled

male and hermaphrodite flowers (andromonoecy), and each flower measures ~15-25 mm in

diameter. The hermaphrodite flowers are characterized by the presence of a lateral pistil, with

two upper fertile stamens, curved downwards and a third lower sterile stamen. Filaments

have long purple, bearded hair. The flower also bears three trilobed sterile staminodes with

glabrous filaments. The male flowers are characterized only by the lack of a pistil (Fig. 1b -

f). The flowers are defined by a short floral anthesis time, from 12:30 hrs. to 16:00 hrs., with

the principal pollinator reward being pollen grains only (Faden 2000, Nandikar and Gurav

2015).

Study site

Western Ghats is one of the four biodiversity 'hotspots' in India. The northern part of

Western Ghats is famous for high elevation laterite plateaus and slopes, one of them being

Kaas plateau in Maharashtra (17° 43′ 12″ N, 73° 49′ 22″ E), which is located at an altitude of

~1225 m above sea level (Fig. 1a). Kaas plateau has an area of ~10 km^2 and is situated

northwest, about 25 km away from the city of Satara in the western state of Maharashtra. It

was declared as a UNESCO World Heritage site in 2012 due to its highly endemic floral

diversity, which is most abundant in the months of August to November.

178   Capped with red lateritic crusts, the upper Kaas Plateau provides arid habitats except

during the monsoon season (Fig. 1a). It receives over 2500 mm of rain every year, mainly

during the monsoon months (June to September), and the daily mean temperatures are over

22°C (Bhattarai et al. 2012). Floral diversity on the Kaas plateau is a model representation of

flora that is associated with seasonal monsoonal rains in the Western Ghats, as well as the

flora of lateritic plateaus of the Western Ghats. All field experiments were carried out across

three consecutive years - 2017, 2018 and 2019, between 15th September and 5th November.

The first two years focused on the study of plant-pollinator interactions and the third year

involved the study of the effect of nutrient resources on the gender of the flower, and also

monitored possible yearly fluctuations in gender expression.

Gender distribution in the population

To determine the extent of sexual dimorphism in morphological characters of *M. simplex*,

flower sizes were compared. Pairs of male and hermaphrodite flowers were collected from 15

random inflorescences and then arranged and photographed on a laminated graph sheet. From

these images, petal length and width (widest diameter) were measured manually (Appendix

S1: Fig. S1).

194   In order to identify if the natural gender distribution within a plant and within an

inflorescence were male dominant or hermaphrodite dominant, a multi-year census was

conducted within the Kaas plateau. All census were carried out in the years 2018 and 2019 at

an interval of seven days, across 4 weeks (spanning over one and a half months) covering the

peak flowering season of *M. simplex*. Gender census were conducted between 12:30 hrs. to

16:00 hrs. (anthesis time of *M. simplex* flowers), on marked inflorescences (N = 30). Since a

single individual can bear 2-5 inflorescences, a pooled census was taken from all the

inflorescences of a plant as our initial observations found no difference among the

inflorescences within a plant. To identify if there is a positional-gender bias i.e. from basal to

the apical position within an inflorescence (see Stephenson 1981, Diggle 1995, 1997, Cuevas

and Polito 2004) the positions of each flower within an inflorescence were recorded along

with its gender details, over the census duration mentioned above. The positions within an

inflorescence were divided into three categories- apical, middle, and basal as shown in Fig.

1c.

To test the effect of nutrient resources on gender expression, a floral gender census

was conducted among two populations that differed in the availability of water and other

resources (presence or absence of a stream and a deep soil substrate), which we observed to

be primary limiting resources on an otherwise homogeneous plateau. The first population is

henceforth referred to as the plateau population - 'Plateau_pop', while the second population

is referred to as the stream population - 'Stream_pop'. Plateau_pop was selected as the

resource-poor population whereas Stream_pop was selected as the resource-rich population.

The two locations were separated by ~576 meters and other than the presence of a stream and

a deep soil substrate, can be viewed as a single population. The census was carried out in

October 2019 on randomly selected individuals within these two populations (N = 100 per

location).

Pollination biology

Pollen is the only pollinator reward present in the genus *Murdannia* (Faden 2000, Nandikar

and Gurav 2015). To identify any quantifiable difference in the pollen or paternal

contribution between the male and hermaphroditic flowers, the amount of pollen per anther in

virgin flowers of *M. simplex* was quantified using a hemocytometer (Bechar et al. 1997).

Pollinator visitations on *M. simplex* in the wild were quantified by observing

pollinators in 2 ft. x 2 ft. observation arenas (N = 49) in multiple intervals of 15 minutes

between 12:30 hrs. to 16:00 hrs. (anthesis time of *M. simplex*). Each arena consisted of 4-10

inflorescences having 15-30 flowers per area. Visitation by a pollinator was recorded when a

pollinator was observed to land on the flower and actively collect pollen. The following three

features were recorded and quantified: total number of flowers present, type of pollinator, and

number of pollinator visits in a 15-minute interval. Pollinator visitation rate was then

calculated as the total number of visits per flower per hour. At the end of the flowering

season, fruitset was quantified for *M. simplex* by measuring the number of fruits in randomly

selected inflorescences from the study population in 2017 (N = 26) and 2019 (N = 40).

Gender-specific pollinator preference experiment

Manipulated choice experiments were designed and conducted in the wild in an open

manipulated setup to test: a) presence of visitation bias by pollinators towards male flowers

(M) and/or hermaphrodite flowers (H), and b) presence of floral-density dependent bias by

pollinators. It was hypothesized that if the pollinator has an inherent gender preference,

irrespective of the gender density in a population, pollinators will show high visitation to the

same gender. Floral distributions within an inflorescence were defined to have binary

outcomes i.e., male abundant or hermaphrodite abundant, or equally male and

hermaphroditic. The experimental setup consisted of two pots that were kept one foot apart

with a total of 5-15 flowers in each pot and pollinator visits were quantified by eye between

12:30 hrs. to 16:00 hrs. Three gender-density treatments: H < M (T1), H = M (T2) and H > M

(T3) were tested. The density of flowers of the two genders in each experimental set up was

manually controlled and each treatment was repeated for a total of eight trials and a total

observation time of ~21.5 hours (Appendix S1: Fig. S2). Unmanipulated, wild set up

observations served as the control. The mean visitation rate per flower to male flowers and

hermaphrodite flowers were calculated for each pollinator type, by dividing the total number

of visits by a specific pollinator with the total number of flowers present in the experimental

set up. Pollinator visitation rate was calculated as the total number of visits per flower per

hour.

The overall visitation observed in all three different density treatments in the

manipulative choice experiment were compared to test the effect of change in density on the

visitation pattern of pollinators in this andromonoecious species. The visitation rates in the

control treatments (natural visitations) were calculated and compared with the visitation rates

obtained in our choice experiment (H < M, H = M and H > M). The mean visitation rates

across all three treatments were also compared with the total number of flowers present in the

trial to identify any effect of floral display size.

Statistical analysis

All analyses were performed in R (R Core Team 2019) and raw data was tested for normality

before proceeding with statistical analysis using Shapiro-Wilk's method. When Gaussian

type distribution could not be assumed, non-parametric Kruskal-Wallis rank-sum tests were

run using the dplyr package (from tidyverse package; Wikcham et al. 2019). The floral

diameters and pollen densities per flower between male and hermaphrodite flowers were

compared using unpaired two-sample t-test. The variation in proportion of male and

hermaphrodite flowers was quantified using Kruskal-Wallis test, and post hoc Dunn's test

(Dinno 2017) was used to identify the significant differences in proportion of genders

between the four weeks. Simple linear regression was used to check the effect of mean

number of male flower, hermaphrodite flowers and total flowers on the fruitset respectively,

and the effect of total floral display size on the mean visitation rate across the three

treatments in the choice experiments. The alpha was set at 0.5 (P<0.025) for all statistical

analysis. All graphical plots were produced using the ggplot2 package in R (from tidyverse

package; Wikcham et al. 2019).

Results

Gender distribution in the population

Due to the small size and fragile nature of the flower, floral size was measured as the widths

of the flowers. The floral morphometry between the size of male and hermaphrodite flowers

showed no significant difference (t = 0.11072, df = 28, *p*-value = 0.9126; Appendix S1: Fig.

S1). Throughout the flowering season, male flowers were present in higher proportions than

the hermaphrodite flowers in all the study plots by at least 70-80% (Kruskal-Wallis chi-

squared = 38.707, df = 7, *p*-value = 2.223^{e-06}; Fig. 2a; Appendix S1: Table S1). A slight

decrease in the proportion of male flowers was observed only in week 4 (Mean±SE: Week 1

= 1.33±0.31, Week 2 = 1.6±0.29, Week 3 = 1.73±0.35, Week 4 = 0.87±0.22; Fig. 2a;

Appendix S1: Table S1). In contrast, the number of hermaphrodite flowers remained similar

across all four weeks of observation (Mean±SE: Week 1 = 0.60±0.21 , Week 2 = 0.33±0.12 ,

Week 3 = 0.50±0.27 , Week 4 = 0.40±0.22 ; Fig. 2a; Appendix S1: Table S1) and only plants

that were closer to the water source i.e. stream population, had a significantly higher number

of hermaphrodite flowers (Mean±SE: Plateau_pop H = 0.43±0.07 vs. Stream_pop H =

1.75±0.14; *p*-values = 0.0000; Fig. 3; Appendix S1: Table S2). Overall, the mean number of

total flowers remained consistent for the first two weeks (Mean±SE: Week 1 = 1.93±0.39,

Week 2 = 1.93±0.37 flowers; Fig. 2a inset), increased in week 3 (2.23±0.41 flowers) and then

reduced in week 4 (1.26±0.28 flowers; Kruskal-Wallis chi-squared = 3.2847, df = 3, *p*-value

= 0.3498; Appendix S1: Table S1).

Within an inflorescence, male and hermaphrodite flowers were present in all the three

positions i.e. apical, middle and basal positions (Fig. 1c), and the mean number of male

flowers varied across all these three positions during our four-week observation period. The

number of male flowers were relatively consistent in the basal position throughout the plants'

flowering season, than in the middle position or apical position (Fig. 2 b - e; Appendix S1:

Table S3, S4). We documented the highest number of male flowers in the middle position of

the inflorescence, whereas the number of hermaphrodite flowers was consistent at all

positions over the flowering season (Fig. 2 b - e; Appendix S1: Table S3, S4).

The total number of flowers was significantly higher in the stream population than in

the plateau population (Mean±SE: Plateau_pop = 2.78±0.12 vs. Stream_pop = 3.32±0.17; t =

-2.5646, df = 176.65, p-value = 0.0116; Fig. 3 inset) and the stream population also showed

similar proportions of male and hermaphrodite flowers (Mean±SE: Stream_pop H =

1.75±0.14 vs. Stream_pop M = 1.57±0.14; p-value = 0.1614; Fig. 3). Proportion of male

flowers was significantly greater in the plateau population over the stream population

(Mean±SE: Plateau_pop M = 2.35±0.12 vs. Stream_pop M = 1.57±0.14; p-value = 0.0000;

Fig. 3; Appendix S1: Table S2).

The mean fruit set per plant showed no effect with the total number of male,

hermaphrodite, or total number of flowers present per plant, respectively in the plateau_pop

(hermaphrodite: r^2 = -0.0457, df = 18, p-value = 0.6852; Fig. 4a; male: r^2 = -0.04999, df =

18, p-value = 0.761; Fig. 4b; total flowers: r^2 = -0.04254, df = 18, p-value = 0.6412; Fig. 4c),

and in the stream_pop (hermaphrodite: r^2 = -0.0.05543, df = 18, p-value = 0.9637; Fig. 4d;

male: r^2 = 0.1125 , df = 18, p-value = 0.08134; Fig. 4e; total flowers: r^2 = 0.04828, df = 18,

p-value = 0.1781; Fig. 4f), respectively.

Pollination biology

Flowers of both the genders opened between 12:30 hrs. to 16:00 hrs., and the mean pollen

count for both the genders ranged from 5000-10000 pollen grains per anther and no

significant difference in the pollen quantity between the two genders was observed (t = -

0.67913, df = 60, p-value = 0.4997; Azad 2018, Albal 2019, Nawge 2019). The major

pollinators to *M. simplex*, in the order of visitation rates from highest to lowest were *Apis*

dorsata (Mean number of visits per flower per hour±SE = 1.01±0.18), *Apis cerana*

(0.86±0.10)*, Apis florea* (0.76±0.22), and *Amegilla spp. (Zonamegilla*, 0.3±0.09)

respectively. *Amegilla spp.* showed significantly lower visitation than all the other pollinators

(Kruskal-Wallis chi-squared = 6.5702, df = 3, *p*-value = 0.09; *p*-values: *Amegilla spp.* vs *A.*

cerana = 0.0074, *Amegilla spp.* vs *A. dorsata* = 0.0058, *Amegilla spp.* vs *A. florea* = 0.0469),

whereas the visitation rates among *A. cerana, A. florea* and *A. dorsata* were not significantly

different (*p*-values: *A. cerana* vs *A. dorsata* = 0.3822, *A. cerana* vs *A. florea* = 0.4278, *A.*

dorsata vs *A. florea* = 0.3748; Azad 2018). Other pollinators observed were *Paragus spp.*

(Hoverfly), *Lasioglossum spp.*, *Pseudapis spp.* and two more species that we could not

identify, henceforth referred to as UnID sp. 3 and UnID sp. 4 (Fig. 5). Mean pollinator

visitation rates were observed to be high at the beginning of the anthesis (1.74-1.94 flowers

from 13:00 hrs. to 15:00 hrs.) and it decreased with time (0.94-1.32 flowers from 15:00 hrs.

till anthesis end; Azad 2018).

Gender specific pollinator preference experiment

In all the experimental setups in the manipulated choice experiment, pollinator visitations

were highest to the most abundant gender (Kruskal-Wallis chi-squared = 65.3103, df = 7, *p*-

value = 1.303^{e-11}; Fig. 6; Appendix S1: Table S5). Pollinator visitation rates were

significantly different between hermaphrodite flowers in the unmanipulated, wild set up

(control) and hermaphrodite flowers in all the experimental setups (Fig. 6; Appendix S1:

Table S5). However, the visitation rates to the male flowers in the control were not different

from those to male flowers in all the experimental setup (Fig. 6; Appendix S1: Table S5).

Display size (total number of flowers per trial) had no effect on the visitation rates in any of

the manipulated treatments (r^2 = -0.03765, df = 21, *p*-value = 0.6309; Appendix S1: Fig. S3).

Quantification of pollinator-wise visitation rates in the manipulated experiments show

that visitations by *Apis cerana, Amegilla spp.* and UnID sp. 4 were higher to male flowers in

treatment 1 where male flowers were more abundant than hermaphrodite flowers (Mean

visitation rate per flower per hr±SE:- H < M: *A. cerana* = 3.29±1.11, *Amegilla spp.* =

1.81±0.91, UnID sp. 4 = 2.49±1.89 flowers; Kruskal-Wallis chi-squared = 13.8517, df = 9, *p*-

value = 0.1277; Fig. 7a; Appendix S1: Table S6). In treatment 3, *Apis cerana*, *Amegilla spp.*,

and UnID sp. 4 showed high visitation to hermaphrodite flowers (H > M: *A. cerana* =

2.76±0.74, *Amegilla spp.* = 0.89±0.61, UnID sp. 4 = 0.66±0.56 flowers; Kruskal-Wallis chi-

squared = 9.2848, df = 5, *p*-value = 0.09823; Fig. 7c; Appendix S1: Table S6). We observed

visitation by hoverflies only in treatment 2 (H = M: Visitation rate per hermaphrodite flower

= 0.56, visitation rate per male flower = 0.75; Kruskal-Wallis chi-squared = 15.605, df = 11,

p-value = 0.1565; Fig. 7b; Appendix S1: Table S6).

Discussion

Gender distribution in the population

The optimal allocation hypothesis proposes that between a costly and a cheaper gender, in a

resource-limited environment, the gender which requires the least resources will be preferred.

From a plant's perspective, the male flowers are considered a cheaper gender to produce

because there is no investment towards fruit development, unlike the hermaphrodite flowers

which produces both viable pollen and ovules (Bertin 1982a, Anderson and Symon 1989,

Kaul et al. 2002, Narbona et al. 2002, Obeso 2002, Verdú et al. 2007, Zhang and Tan 2009).

Here, we show that the *M. simplex* population on Kaas plateau has a higher density of male

flowers and the only individuals that showed a slight reduction in the number of male flowers

and an increase in hermaphrodite flowers were the ones present along the stream. We

consider the stream population of *M. simplex* to have better resources available to them than

the plateau population because of the presence of water (a limited resource otherwise on the

plateau) and possible higher nutrient availability due to dissolved/associated nutrients in the

water and deeper soil substrate available for the plants (Watve 2013, Ramey and Richardson

2017, Thorpe et al. 2018). This suggests that although the general tendency of *M. simplex* is

to produce more male flowers over hermaphrodites, water and water-dissolved minerals may

be major limiting resources and in the absence of resource constraints the plants can produce

not only more flowers but also higher number of hermaphrodite flowers.

Gender shift with time

Resources available and utilized by plants often fluctuate with time and age of the plant

(Stephenson 1982, Solomon 1985). We observed an initial increase in the number of flowers

followed by a decreasing trend in the total number of flowers being produced per plant

throughout its flowering period (Fig. 2a inset). Within this trend, male flowers were always

produced in higher proportions than the hermaphrodite flowers and interestingly, in the end

of the flowering season (i.e. fourth week) we observed a reduction in the number of male

flowers instead of the number of hermaphrodite flowers (Fig. 2a). Thus, although resource

constraints do affect the production of total flowers as observed in the stream vs. plateau

population, resource limitation due to the flowering stage of the plant only affected the

number of male flowers in *M. simplex*. It has been shown that under declining resource levels

plants may opt to increase their fitness by producing the relatively inexpensive male flowers

over the expensive female flowers (May and Eugene Spears Jr 1988). Further, it has been

reported in *C. benghalensis* that a plant will opt to drop a flower rather than dropping a fruit,

under resource limiting conditions, because the nutrient resources that would get committed

to fruits that is aborted would be saved halfway (Kaul and Koul 2008). We propose that a

similar strategy may be operational in *M. simplex* wherein under availability of resources *M.*

simplex may be producing excess of male flowers which are the first ones to be sacrificed

when resource limitation starts to be an important factor as the phenology of the plant

progresses (Fig. 2a). The lack of change in the number of hermaphrodite flowers in different

stages of plant's flowering phenology also suggests that the hermaphrodite flowers may

already be produced at an optimal amount, resulting in the maintenance of their number

despite declining resources at the later stages (Fig. 2a).

Gender distribution of flowers on an inflorescence has been shown to be dependent on

resource allocation (Diggle 1995, 1997, Miller and Diggle 2003), and it has been observed

that the differential positioning of the genders within an inflorescence can affect outcrossing

rates (Orth and Waddington 1997, Lin and Forrest 2019). Studies in *Solanum* (Diggle 1994,

1995, 1997, Miller and Diggle 2003), *Caesalpinia* (Lewis and Gibbs 1999), *F. virginiana*

(Ashman and Hitchens 2000) and in other Commelinaceae genera (Faden 1991, Kaul et al.

2002, Kaul and Koul 2008) have shown that higher accessibility to nutrient resources at the

base of an inflorescence (than the distal end) can result in the presence of higher resource-

demanding hermaphrodite flowers in the basal position while males are relegated to the

middle and apical positions of an inflorescence. We found a similar pattern in *M. simplex,*

where a greater number of male flowers were produced at the middle and the apical positions

on an inflorescence. However, unlike other plants where hermaphrodites exclusively occupy

the basal positions in an inflorescence, in *M. simplex* hermaphrodites were also present in the

apical position. In fact, the distribution of hermaphrodite flowers was not significantly

different across all the three positions, over the course of the plant's flowering season. This

implies that any manipulation in display size is carried out by increasing or decreasing the

male flowers thus regulating the floral display at a much lesser cost (as proposed in *Solanum*

by Anderson and Symon 1989).

Pollination biology

For inflorescence size to affect reproductive success, the number of flowers visited per

inflorescence by pollinators must increase with display size because pollinators have been

shown to display higher preference to larger inflorescences (Stephenson 1981, Harder and

Barrett 1995, Ohashi and Yahara 2001). Hence the disproportionate gender distribution

within the inflorescence of an andromonoecious species may be a result of indirect pollinator

selection through inflorescence size: that is larger inflorescences are better at attracting

pollinators but under nutrient resource limitation, this can be achieved only by producing

more male flowers, which are cheaper to produce (Sandring and Ågren 2009). Thus our

results support the pollinator attractor hypothesis as proposed by Zhang and Tan (2009) and

sex allocation theory as proposed by Elle and Meagher (2000) where they predicted that male

flowers can increase the floral display and their attractiveness to pollinators at a lower cost

through male function than female function and thereby increase the stigmatic load of the

hermaphrodite flowers, which potentially increases female reproductive success at lower

resource investment.

The dominant pollinator assemblage of *M. simplex* consisted mainly of different

species of bees from the two genera of bees common in the Indian tropics - *Apis* and

Amegilla. All pollinators that visit *M. simplex* are generalists and are known to be common

pollinators in grassland habitats (Corlett 2011, Dhargalkar 2019, Nawge 2019). There are at

least 68 plants co-flowering with *M. simplex* on the Kaas plateau (Dhargalkar 2019) and

parallel studies on the pollination of plants in the Kaas community showed that both *Apis*

cerana and *Apis florea* switch to *M. simplex* from *M. lanuginosa* (anthesis time 10.30 hrs.-

13.00 hrs.) and various *Impatiens* species as soon as the flowers of *M. simplex* open i.e. at

12:30 hrs. up to 16:00 hrs. (Albal 2019). This suggests that although the pollinators are

generalists, these pollinators may be displaying temporal specialization towards *M. simplex*.

Future studies will focus on temporal specialization of the bee communities on Kaas plateau,

which will provide insight on pollinator preference towards *M. simplex* in a multi-species

environment.

Gender-specific pollinator preference

Several studies have shown that pollinators avoid functionally female flowers because of

decreased pollen or absence of pollen (Bierzychudek 1987, Eckhart 1991), floral size

(Ashman et al. 2000), floral scents (Ashman et al. 2005) and pollen-pistil interference

(Solomon 1985, Elle and Meagher 2000). However, in *M. simplex* both male and

hermaphrodite flowers showed comparable total pollen production. This has been shown in

other andromonoecious systems as well such as in *Solanum carolinense* (Solomon 1986),

Olea europaea (Cuevas and Polito 2004) and *Dichorisandra spp.* (Sigrist and Sazima 2015),

where the male flowers play a crucial role in the reproductive success of the plant by

increasing the total number of pollen grains available per plant. As previously discussed, we

observed that nutrient resources affected the number of male flowers and not the number of

hermaphrodite flowers, and the rate of fruit sets were observed to be similar between the

plateau and the stream populations (Mean±SE: Plateau_pop = 3.22±0.23 vs. Stream_pop =

3.55±0.21). This suggests that reproductive success is limited by the availability of

hermaphrodite flowers and not pollinator visitations. The lack of preference for either of the

gender by the pollinators on Kaas plateau is further supported by our observations from the

manipulative experiment where all pollinators except *A. cerana* displayed lack of preference

towards either of the floral genders. Further, this also fails to support the hypothesis that

male-dominant inflorescences of *M. simplex* may be under pollinator-mediated selection due

to direct pollinator preference for any gender (see Solomon 1985, Eckhart 1991, Elle and

Meagher 2000). However, since pollinator visitations were observed to be density dependent

(i.e. floral density, compare treatment 3 with treatment 1, Fig. 7c) it is possible that

pollinator-mediated selection is acting on male flowers through selection of larger

inflorescence sizes.

Our understanding of pollinator services by bees primarily come from *Bombus*

species (Chittka et al. 1997, 1999) and *Apis mellifera* (Martin 2004, Howard et al. 2019) and

studies on other bees from India are very rare (see Somanathan and Borges 2001). At least

five species of bees visited *M. simplex* and in the choice experiment, we observed inter-

species behavioral differences among bees both in their floral preferences and visitation

patterns, respectively. Almost all bees displayed higher visitation to any gender which was

present in greater density, suggesting that pollinator visitations are predominantly gender-

nonspecific and density-dependent. A slight preference for gender was only observed in *A.*

cerana and *Amegilla* in treatment 2 (H = M) where male flowers were preferred over

hermaphrodite flowers (Fig. 7b). Bees are known for both their short-term and long-term

memory retention and they can show learning through foraging experience (Cresswell and

Galen 1991, Schiestl and Johnson 2013) and we propose that the higher visitation to male

flowers in treatment 2 (Fig. 7b) may be a result of short-term memories and habituation in

bees since in natural conditions bees may encounter more male flowers than hermaphrodites.

We conclude that in *M. simplex* environmentally driven resource constraints play a

bigger role than pollinator preference towards a specific gender in determining gender

expression. However, we cannot dismiss an indirect role of pollinator-driven constraints

which may occur through selection on the display size as proposed in the pollinator attractor

hypothesis (Zhang and Tan 2009). In a resource-limited environment such as the one

observed at Kaas plateau, *M. simplex* can thus increase its display size by producing the less

expensive male flowers over hermaphrodite flowers.

Energy constraints will have a direct cost on floral production if genders require

differential resource allocation. As a result, it can be expected that the gender of the flowers

will be selected accordingly. Using the *M. simplex* system, we show that the gender

expression is directly influenced by the availability of nutrient resources in the plants'

environment and indirectly affected by the pollinators' preference for larger display sizes of

the inflorescence. Thus, a combination of both environmental factors as well as ecological

factors may be responsible for the evolution and maintenance of andromonoecy in mass-

flowering plants such as *M. simplex* in the tropics.

Acknowledgements

The authors thank the Ministry of Human Resource Development, India (MHRD) and the

National Geographic Society for research funds to VG, IISER Bhopal for the infrastructure,

DST for INSPIRE fellowship to AA, AG, and SS. We also thank the PCCF of Maharashtra

Forest Department, DFO of Satara Division (Forest Department) and the Joint Forest

Management Committee (JFMC), Kaas, Satara for permitting our fieldwork. We also thank

Dr. Natapot Warrit (Chulalongkorn University, Thailand) and Dr. Ximo Mengual (Research

Museum Alexander Koenig, Germany) for their help with taxonomic identification of the

pollinators. Finally, we thank all fellow TrEE lab members for their advice, discussion, and

support.

Author contributions

VG conceived the idea, AA, AG, and VG designed the experiments. AA and AG conducted

the fieldwork with help from SS. AA performed the statistical analyses with inputs from SS

and VG. AA and VG wrote the manuscript and AG and SS helped with edits. All authors

read and approved the final draft of the manuscript.

Literature cited

Albal, A. 2019. When does a flower become male? Exploring role of ecological factors and

energy resources in defining gender expression in sympatric plants. Master's thesis,

IISER Bhopal, India R00529:14030.

Anderson, G. J., and D. E. Symon. 1989. Functional dioecy and andromonoecy in *Solanum*.

Evolution 43:204.

Armbruster, W. S. 2001. Evolution of floral form: Electrostatic forces , pollination, and

adaptive compromise. New Phytologist 152:181–186.

Ashman, T. L., M. Bradburn, D. H. Cole, B. H. Blaney, and R. A. Raguso. 2005. The scent of

a male: The role of floral volatiles in pollination of a gender dimorphic plant. Ecology

86:2099–2105.

Ashman, T. L., and M. S. Hitchens. 2000. Dissecting the causes of variation in intra-

inflorescence allocation in a sexually polymorphic species, *Fragaria virginiana*

(Rosaceae). American Journal of Botany 87:197–204.

Ashman, T. L., and M. T. Morgan. 2004. Explaining phenotypic selection on plant attractive

characters: Male function, gender balance or ecological context? Proceedings of the

Royal Society B: Biological Sciences 271:553–559.

Ashman, T. L., J. Swetz, and S. Shivitz. 2000. Understanding the basis of pollinator

selectivity in sexually dimorphic *Fragaria virginiana*. Oikos 90:347–356.

Azad, G. 2018. Resource allocation and pollinator preference : role of male flowers in the

andromonoecious herb *Murdannia simplex* (Commelinaceae). Master's thesis, IISER

Bhopal, India R00335:13035.

Bateman, A. J. 1948. Intra-sexual selection in *Drosophila*. Heredity 2:349–368.

Bechar, A., S. Gan-Mor, Y. Vaknin, I. Shmulevich, B. Ronen, and D. Eisikowitch. 1997. An

image-analysis technique for accurate counting of pollen on stigmas. New Phytologist

137:639–643.

Benseler, R. W. 1975. Floral biology of California buckeye. Madroño 23:41–53.

Bertin, R. I. 1982a. The evolution and maintenance of andromonoecy. Evolutionary Theory

6:25–32.

Bertin, R. I. 1982b. The ecology of sex expression in red buckeye. Ecology 63:445–456.

Bhattarai, U., P. Tetali, and S. Kelso. 2012. Contributions of vulnerable hydrogeomorphic

habitats to endemic plant diversity on the Kas Plateau, Western Ghats. SpringerPlus 1:1-

9.

Bierzychudek, P. 1987. Pollinators increase the cost of sex by avoiding female flowers.

Ecology 68:444–447.

Charnov, E., and J. Bull. 1977. When is sex environmentally determined? Nature 266:828–

832.

Chittka, L., A. Gumbert, and J. Kunze. 1997. Foraging dynamics of bumble bees: Correlates

of movements within and between plant species. Behavioral Ecology 8:239–249.

Chittka, L., J. D. Thomson, and N. M. Waser. 1999. Flower constancy, insect psychology,

and plant evolution. Naturwissenschaften 86:361–377.

Christopher, D. A., R. J. Mitchell, D. W. Trapnell, P. A. Smallwood, W. R. Semski, and J. D.

Karron. 2019. Hermaphroditism promotes mate diversity in flowering plants. American

Journal of Botany 106:1131–1136.

Corlett, R. T. 2011. Honeybees in natural ecosystems. In: Hepburn, H. and Radloff, S.E.

(Eds) Honeybees of Asia. Springer- Verlag Berlin Heidelberg Pages 215-225

Cresswell, J. E., and C. Galen. 1991. Frequency-dependent selection and adaptive surfaces

for floral character combinations: The pollination of *Polemonium viscosum*. The

American Naturalist 138:1342–1353.

Cuevas, J., and V. S. Polito. 2004. The role of staminate flowers in the breeding system of

Olea europaea (Oleaceae): An andromonoecious, wind-pollinated taxon. Annals of

Botany 93:547–553.

Dhargalkar, R. K. 2019. A study of the temporal patterns in flowering phenology and plant-

pollinator network of a grassland community in the northern Western Ghats ,

Maharashtra. Master's thesis, IISER Bhopal, India R00534:14054.

Diggle, P. K. 1993. Developmental plasticity , genetic variation , and the evolution of

andromonoecy in *Solanum hirtum* (Solanaceae). American Journal of Botany 80:967–

973.

Diggle, P. K. 1994. The expression of andromonoecy in *Solanum hirtum* (Solanaceae):

Phenotypic plasticity and ontogenetic contingency. American Journal of Botany

81:1354–1365.

Diggle, P. K. 1995. Architectural effects and the interpretation of patterns of fruit and seed

development. Annual Review of Ecology and Systematics 26:531–552.

Diggle, P. K. 1997. Ontogenetic contingency and floral morphology : The effects of

architecture and resource limitation. International Journal of Plant Sciences 158:99–107.

Dinno, A. 2017. dunn.test: Dunn's test of Multiple comparisons using rank sums.

Eckhart, V. M. 1991. The effects of floral display on pollinator visitation vary among

populations of *Phacelia linearis* (Hydrophyllaceae). Evolutionary Ecology 5:370–384.

Elle, E., and T. R. Meagher. 2000. Sex allocation and reproductive success in the

andromonoecious perennial *Solanum carolinense* (Solanaceae). II. Paternity and

functional gender. American Naturalist 156:622–636.

Faden, R. B. 1991. The morphology and taxonomy of *Aneilema* R. Brown (Commelinaceae).

Smithsonian contributions to Botany 76.

Faden, R. B. 2000. Floral biology of Commelinaceae. Pages 309–317 Monocots: Systematics

and Evolution.

Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004.

Pollination Syndromes and Floral Specialization. Annual Review of Ecology, Evolution,

and Systematics 35:375–403.

Harder, L. D., and S. C. H. Barrett. 1995. Mating costs of large floral displays in

hermaphrodite plants. Nature 373:512–515.

Howard, S. R., M. Shrestha, J. Schramme, J. E. Garcia, A. Avarguès-Weber, A. D. Greentree,

and A. G. Dyer. 2019. Honeybees prefer novel insect-pollinated flower shapes over bird-

pollinated flower shapes. Current Zoology 65:457–465.

Kaul, V., and A. K. Koul. 2008. Floral phenology in relation to pollination and reproductive

output in *Commelina caroliniana* (Commelinaceae). Australian Journal of Botany

56:59–66.

Kaul, V., N. Sharma, and A. K. Koul. 2002. Reproductive effort and sex allocation strategy in

Commelina benghalensis L., a common monsoon weed. Botanical Journal of the

Linnean Society 140:403–413.

Lewis, G., and P. Gibbs. 1999. Reproductive biology of *Caesalpinia calycina* and *C .

pluviosa* (Leguminosae) of the caatinga of north-western Brazil. Plant Systematics and

Evolution 217:43–53.

Lin, S., and J. R. K. Forrest. 2019. The function of floral orientation in bluebells : interactions

with pollinators and rain in two species of *Mertensia* (Boraginaceae). Journal of Plant

Ecology 12:113–123.

Lloyd, D. G. 1980. Sexual strategies in Plants: I. An hypothesis of serial adjustment of

maternal investment during one reproductive session. New Phytologist 86:69–79.

Martin, N. H. 2004. Flower size preferences of the honeybee (*Apis mellifera*) foraging on

Mimulus guttatus (Scrophulariaceae). Evolutionary Ecology Research 6:777–782.

May, P. G., and E. Eugene Spears Jr. 1988. Andromonoecy and variation in phenotypic

gender of *Passiflora incarnata* (Passifloraceae). American Journal of Botany 75:1830–

1841.

Miller, J. S., and P. K. Diggle. 2003. Diversification of andromonoecy in *Solanum* section

Lasiocarpa (Solanaceae): The roles of phenotypic plasticity and architecture. American

Journal of Botany 90:707–715.

Nandikar, M. D., and R. V. Gurav. 2015. Revision of the genus *Murdannia* (Commelinaceae)

in India. Phytodiversity 2:56–112.

Narbona, E., P. L. Ortiz, and M. Arista. 2002. Functional Andromonoecy in *Euphorbia*

(Euphorbiaceae). Annals of Botany 89:571–577.

Nawge, V. V. 2019. Understanding pollination networks on Kaas plateau. Master's thesis,

IISER Bhopal, India R00543:14097.

Obeso, J. R. 2002. The costs of reproduction in plants. New Phytologist 155:321–348.

Ohashi, K., and T. Yahara. 2001. Behavioural responses of pollinators to variation in floral

display size and their influences on the evolution of floral traits. In: Chittka, L.,

Thomson, J.D. Cognitive Ecology of Pollination- Animal behaviour and floral evolution.

Cambridge University Press Pages 274-296.

Orth, A. I., and K. D. Waddington. 1997. The movement patterns of carpenter bees *Xylocopa*

micans and bumblebees *Bombus pennsylvanicus* on *Pontederia cordata* inflorescences.

Journal of Insect Behavior 10:79-86.

Ortiz, P. L., M. Arista, P. E. Oliveira, and S. Talavera. 2003. Pattern of flower and fruit

production in *Stryphnodendron adstringens*, an andromonoecious legume tree of Central

Brazil. Plant Biology 5:592–599.

Oziegbe, M., I. O. Akinlua, and A. A. Olalekan. 2013. Comparative pollination role of

stamens and breeding system in three species of *Commelina* (Commelinaceae) in Ile-Ife,

Nigeria. Acta Botanica Brasilica 27:543–550.

Primack, R. B., and D. G. Lloyd. 1980a. Andromonoecy in the New Zealand montane shrub

Manuka, *Leptospermum scoparium* (Myrtaceae). American Journal of Botany 67:361–

368.

Primack, R. B., and D. G. Lloyd. 1980b. Sexual strategies in plants IV. The distributions of

gender in two monomorphic shrub populations. New Zealand Journal of Botany 18:109–

114.

R Team, C. 2019. R: A language and environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.r-project.org/

Ramey, T. L., and J. S. Richardson. 2017. Terrestrial Invertebrates in the Riparian Zone:

Mechanisms Underlying Their Unique Diversity. BioScience 67:808–819.

Sandring, S., and J. Ågren. 2009. Pollinator-mediated selection on floral display and

flowering time in the perennial herb *Arabidopsis lyrata*. Evolution 63:1292–1300.

Schiestl, F. P., and S. D. Johnson. 2013. Pollinator-mediated evolution of floral signals.

Trends in Ecology and Evolution 28:307–315.

Sigrist, M. R., and M. Sazima. 2015. Phenology, reproductive biology and diversity of

buzzing bees of sympatric *Dichorisandra* species (Commelinaceae): breeding system

and performance of pollinators. Plant Systematics and Evolution 301:1005–1015.

Solomon, B. P. 1985. Environmentally influenced changes in sex expression in an

andromonoecious plant. Ecology 66:1321–1332.

Solomon, B. P. 1986. Sexual allocation and andromonoecy: Resource investment in male and

hermaphrodite flowers of *Solanum carolinense* (Solanaceae). American Journal of

Botany 73:1215–1221.

Somanathan, H., and R. M. Borges. 2001. Nocturnal pollination by the carpenter bee

Xylocopa tenuiscapa (Apidae) and the effect of floral display on fruit set of

Heterophragma quadriloculare (Bignoniaceae) in India. Biotropica 33:78–89.

Stanton, M. L., J. K. Bereczky, and H. D. Hasbrouck. 1987. Pollination thoroughness and

maternal yield regulation in wild radish, *Raphanus raphanistrum* (Brassicaceae).

Oecologia 74:68–76.

Stephenson, A. G. 1981. Flower and fruit abortion: proximate causes and ultimate functions.

Annual Review of Ecology and Systematics 12:253–279.

Stephenson, A. G. 1982. When does outcrossing occur in a mass-flowering plant? Evolution

36:762–767.

Thorpe, C. J., T. R. Lewis, S. Kulkarni, A. Watve, N. Gaitonde, D. Pryce, L. Davies, D. T.

Bilton, and M. E. Knight. 2018. Micro-habitat distribution drives patch quality for sub-

tropical rocky plateau amphibians in the northern Western Ghats, India. PLoS ONE

13:1–20.

Ushimaru, A., T. Watanabe, and K. Nakata. 2007. Colored floral organs influence pollinator

behavior and pollen transfer in *Commelina communis* (Commelinaceae). American

Journal of Botany 94:249–258.

Vallejo-Marín, M., and M. D. Rausher. 2007a. The role of male flowers in andromonoecious

species: Energetic costs and siring success in *Solanum carolinense* L. Evolution 61:404–

412.

Vallejo-Marín, M., and M. D. Rausher. 2007b. Selection through female fitness helps to

explain the maintenance of male flowers. The American Naturalist 169:563.

Venkatesan, A. 2017. Why be a male ? Resource allocation and floral display in a wild

population of Solanum surattense. Master's thesis, IISER Bhopal, India R00064:10015.

Verdú, M., K. Spanos, I. Čaňová, B. Slobodník, and L. Paule. 2007. Similar gender

dimorphism in the costs of reproduction across the geographic range of *Fraxinus ornus*.

Annals of Botany 99:183–191.

Watve, A. 2013. Status review of Rocky plateaus in the northern Western Ghats and Konkan

region of Maharashtra, India with recommendations for conservation and management.

Journal of Threatened Taxa 5:3935–3962.

Wikcham, H., M. Averick, J. Bryan, W. Chang, L. D. McGowan, R. Francois, G. Grolemund,

A. Hayes, L. Henry, J. Hester, M. Kuhn, T. L. Pedersen, E. Miller, S. M. Bache, K.

Muller, J. Ooms, D. Robinson, D. P. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C.

Wilke, K. Woo, and H. Yutani. 2019. tidyverse: Welcome to tidyverse.

Williams, G., and K. Walker. 2003. Pollination of the wet forest herb *Pollia crispata*

(Commelinaceae). Cunninghamia 8(1):141–146.

Wilson, P., J. D. Thomson, M. L. Stanton, and L. P. Rigney. 1994. Beyond floral Batemania:

Gender biases in slection for pollination success. The American Naturalist 143:283-296.

Yampolsky, E., and H. Yampolsky. 1922. Distribution of sex forms in the phanerogamic

flora. Bibliographia genetica 3:1–62.

Zhang, T., and D. Y. Tan. 2009. An examination of the function of male flowers in an

andromonoecious shrub *Capparis spinosa*. Journal of Integrative Plant Biology 51:316–

324.

Figure legends

Fig. 1 Habitat and floral features of *Murdannia simplex* (a) habitat on Kaas Plateau, Maharashtra during flowering season (Inset: Location in India), (b) *M. simplex* habit, (c) inflorescence image showing positions identified from base to the apex, (d) *M. simplex* flower front view, (e) *M. simplex* male flower, (f) *M. simplex* hermaphrodite flower. Photo courtesy: (a), (c), (d) - Asawari Albal, (b), (e), (f) - Saket Shrotri.

Fig. 2 Variation in gender in *M. simplex* (a) by week from first flowering date (Kruskal-Wallis chi-squared = 38.7067, df = 7, *p*-value = 0.00; P<0.025) (Inset: Week-wise proportion of total number of flowers; Kruskal-Wallis chi-squared = 3.2847, df = 3, *p*-value = 0.3498, N = 30; P<0.025) (See Appendix S1: Table S1) and (b) - (e) within-inflorescence from Week 1 - Week 4 respectively (Kruskal-Wallis chi-square test; df = 5, P<0.025). For (a) columns to be compared within the week and within gender (Mean±SE). For (b) columns to be compared within the gender and within positions, across all 4 weeks (Mean±SE). Significant *p*-values depicted by different alphabets (Appendix S1: Table S3; P<0.025).

Fig. 3 Natural gender distribution observed in *M. simplex* in 2019 between two populations (Mean±SE) (Kruskal-Wallis chi-squared = 123.53, df = 3, *p*-value <2.2^{e-16}, N = 100 per population; Appendix S1: Table S2; P<0.025) (Inset: Total number of flowers across the two populations; t = -2.5646, df = 176.65, *p*-value = 0.0116). Male flowers are represented by open squares, hermaphrodite flowers represented by filled squares and the gray squares represent total (male + hermaphrodite) flowers. Columns to be compared within population and within genders across the two populations only.

Fig. 4 For plateau_pop in 2019, effect on mean fruitset by (a) mean number of male flowers (r^2 = -0.04999, df = 18, *p*-value = 0.761, N = 20), (b) mean number of hermaphrodite flowers (r^2 = -0.0457, df = 18, *p*-value = 0.6852, N = 20) and (c) mean number of flowers per plant (r^2 = -0.04254, df = 18, *p*-value = 0.6412, N = 20). For stream_pop in 2019, effect on mean fruitset by (f) mean number of male flowers (r^2 = 0.1125, df = 18, *p*-value = 0.08134, N = 20), (e) mean number of hermaphrodite flowers (r^2 = -0.05543, df = 18, *p*-value = 0.9637, N = 20) and (f) mean number of flowers per plant (r^2 = 0.04828, df = 18, *p*-value = 0.1781, N = 20).

Fig. 5 Pollinators of *M. simplex* from Kaas plateau: (a) and (b) *Amegilla spp. (Zonamegilla)*; (c) *Apis florea*; (d) *Apis cerana*; (e) *Apis dorsata*; (f) *Paragus sp.* (Hoverfly); (g) *Lasioglossum sp.*; (h) and (i) *Pseudapis sp.* Photo courtesy- (a) - (d), (f), (h), (i) : Asawari Albal and (e), (g): Saket Shrotri.

Fig. 6 Mean pollinator visitation rate per flower per hour (Mean±SE) observed in the manipulative experiment where three treatments with varying numbers of hermaphrodite and male flowers were used to test pollinator preference towards a gender; (Kruskal-Wallis chi-squared = 63.4196, df = 7, *p*-value = 3.122^{e-11}, N = 8 trials for each treatment; P<0.025). Male flowers are represented by open squares and hermaphrodite flowers are represented by the filled squares. The first two columns represent the natural visitation rate observed per flower per hour for both male and hermaphrodite, respectively. Different alphabets depict significantly different values (Appendix S1: Table S5; P<0.025).

Fig. 7 Pollinator-wise mean visitation rate per flower per hour in the manipulative experiment on *M. simplex* (Mean±SE) (a) Treatment 1- H < M (Kruskal-Wallis chi-squared = 13.8517, df = 9, *p*-value = 0.1277; P<0.025; Appendix S1: Table S6), (b) Treatment 2- H = M (Kruskal-Wallis chi-squared = 15.605, df = 11, *p*-value = 0.1565; P<0.025; Appendix S1: Table S6),

(c) Treatment 3- H > M (Kruskal-Wallis chi-squared = 9.2848, df = 5, p-value = 0.09823; P<0.025; Appendix S1: Table S6). Male flowers are represented by open squares and hermaphrodite flowers are represented by filled squares. Different alphabets depict significantly different values (Appendix S1: Table S6; P<0.025).

Figures

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7