1

Phylogenetic comparative analyses of the determinants of anti-predator distraction behavior in shorebirds

Miki Saijo*, Nobuyuki Kutsukake

The Graduate University for Advanced Studies, SOKENDAI, Hayama, Miura, Kanagawa 240-0193, Japan

*corresponding author

email: saijo.0830@gmail.com

Acknowledgments

We thank the members of our laboratory for useful discussions and comments on the research plan and analyses. We thank Yamashina Institute for Ornithology for permission to do literature survey.

Abstract

Predation risk exerts a strong selective pressure on anti-predator behavior, resulting in

behaviors to achieve defense of offspring and the individual. In shorebirds, some

species perform distraction behavior that is attracting the attention of a predator. This

behavior evolved, and were lost multiple times, independently and the behavioral

repertoire varies among species. Although defense of offspring is critical for parents, the

determinants of inter-specific variation in the distraction behavior remain unstudied. We

surveyed the literature and conducted phylogenetic comparative analyses (n = 169

species) to test predictions regarding nest site, body mass, and coloniality. We found

that small species were more likely to perform distraction behavior than large species.

Solitary species were more likely to perform distraction behavior than colonial nesting

species. Previous studies suggested that colonial nesting and large species commonly

perform aggressive anti-predator behavior, implying that distraction behavior is an

alternative anti-predator strategy to aggressive ones.

Keywords: offspring defense, predation, phylogeny, alternative strategy, ground-nesting

birds, body mass

Introduction

Predation risk exerts a strong selection pressure on animal behaviors, producing

anti-predator behavior among others (Lack 1954; Caro 2005). Offspring predation is the

most important reason for nest failure in birds (Clark & Wilson, 1981). Therefore,

parents have evolved various anti-predator behaviors to counter offspring predation,

thus increasing their fitness (Royle et al. 2012).

Shorebirds (Charadriiformes) are an ideal taxon for assessing the determinants

of the anti-predator behavioral repertoire. Most shorebird species nest on open ground,

exposing the chicks and eggs to strong predation pressure (Gochfeld, 1984; Kubelka et

al., 2018). Shorebirds engage in two broad types of anti-predator behavior: attack and

distraction (Gochfeld, 1984). Attack behavior includes any type of attack on predators,

such as mobbing and scolding. Distraction behavior aims to attract the attention of a

predator via injury-feigning, false breeding, and "rodent run" behaviors, for example

(Caro, 2005; Gochfeld, 1984). When a predator was deceived by the display and

chasing a parental bird, the bird moves away from their offspring (Ristau, 1991). As a

result, predators lost sight of the nest or chick. When the predator was lured and far

away from the nest by the display, the parental bird ends its display and runs away

(Gómez-Serrano, 2018). This behavior seems to have evolved multiple times

independently under similar ecological and social conditions (Humphreys & Ruxton,

2020). However, only a few studies have examined the determinants of inter-species

differences in the anti-predator behavioral repertoire (Larsen et al. 1996; da Cunha et al.

2017). Larsen et al. (1996) tested factors of aggressive nest protection behavior in

waders. This paper revealed that body mass and number of parents present on the nest

territory explain the variation of aggressive anti-predator behavior. But they did not use

phylogenetic comparative analysis. da Cunha et al. (2017) showed that small species

were more likely to perform aggressive mobbing than large species. They included anti-

predator behavior during both the breeding and non-breeding seasons.

In contrast to those studies analyzing aggressive anti-predator behavior, the

determinants of distraction behavior remained unstudied. Armstrong (1954) listed six

factors that could explain the performance of distraction behavior, which could be

summarized that distraction display is likely to evolve in species that may be preyed

upon by terrestrial predators during the daytime (Humphreys & Ruxton, 2020).

However, a formal test of those ideas using a modern phylogenetic comparative analysis

has not been done. We assessed three possible determinants of distraction behavior:

nesting site, body mass, and coloniality. First, we predicted that species that nest in trees

and on cliffs are less likely to perform anti-predator behavior, because their nests are

less likely to be predated due to the difficulty of access compared with ground-nesting

species (Coulson & Thomas, 1985). Differences in other traits according to nesting site

were suggested in a detailed comparative study of two closely related sympatric species

(Cullen, 1957). Cliff-nesting kittiwakes have weak anti-predator traits, such as rare

production of alarm calls, unconcealed body color of chicks, and low response to an

attack by a predator. By contrast, the parents of the ground-nesting black-headed gull

(*Larus ridibundus*) attack predators, and their chicks escaped when attacked by a

predator. We predicted that this behavioral difference between the two species is

commonly seen in other taxa. Second, we predicted that body size would affect

distraction behavior. Because of the physical advantages of large species, it has been

supported such species are more likely to perform attack behavior than smaller species

(Larsen et al. 1996), whereas smaller species are more likely to perform distraction

behavior. Third, we predicted that solitary nesting species are more likely to perform

distraction behavior because they cannot attack predators effectively compared with

colonial species. Previous studies showed that colonial species are likely to perform

attack behavior because they can attack a predator effectively as a group (Hoogland &

Sherman, 1976; Robinson, 1985) and because they can minimize the per capita risk of

predation via the dilution effect (Hamilton, 1971; Hogan, Hildenbrandt, Scott-Samuel,

Cuthill, & Hemelrijk, 2017).

Methods

Data collection

We searched the literature for studies on anti-predator behavior during the breeding

season using Google Scholar, with the key words of 'shorebirds', 'plover', 'sandpiper',

'wader', 'gull', 'anti-predator', 'behavior', and 'nesting', and the scientific or species

names of Charadriiformes. We also added the key words 'distraction', 'injury-feigning',

'false-brooding', and 'rodent run' for distraction behaviors (Gochfeld, 1984). We

classified a given species in terms of performance of distraction behavior (0 = does not

perform; 1 = performs) if the paper described those behaviors as anti-predator behaviors

performed during the breeding season. We obtained data for 169 species from 87 papers

(Appendix). We also added data for three species (Kentish plover, *Charadrius*

alexandrinus, 6 nests; little ringed plover, *C. dubius*, 9 nests; and little tern, *Sternula*

albifrons, up to 892 nests) based on direct behavioral observations undertaken at the

Morigasaki Water Recycling Center in Japan (35°57.1'S 139°75.3'W) between May and

July 2017. In order to control the sampling bias by the number of papers and

observations, we used the number of papers retrieved by Google Scalar as an

explanatory variable. The searched word was "anti-predator", "behavior", and scientific

name.

Data on nesting sites and colonies were obtained from del Hoyo et al. (1998).

Data on body mass were obtained from Dunning (2007). We calculated the body mass

of sexual dimorphic species by sex, and used the body mass data of individuals that

incubate eggs (del Hoyo et al. 1998; Paton et al. 1994; Székely and Reynolds 1997;

Székely et al. 2000). In cases where both sexes defend eggs, we used the average of the

male and female weights. We also used the average of the male and female weights if

there were no data on the sexes that defend eggs,. We obtained 100 phylogenetic trees

of the study species from http://birdtree.org/ (Jetz, Thomas, Joy, Hartmann, & Mooers,

2012).

Comparative analysis

We performed all analysis in R software (ver. 3.5.3; R Development Core Team 2019).

We estimated D, a phylogenetic signal for discrete traits (Fritz & Purvis, 2010), for the

presence of distraction behaviors using the "caper" package (Orme, 2013). We used 100

phylogenetic trees and estimated 100 D values for each. For all trees, we found that the

phylogenetic signals of this behavior differed significantly from random ($p < 0.001$).

We used phylogenetic generalized linear mixed models with the MCMCglmm

(Hadfield, 2010) and mulTree packages (Guillerme & Healy, 2014). We included the

100 phylogenetic trees as a random effect to control for the effect of phylogeny. Markov

chain Monte Carlo (MCMC) simulations were run for 240,000 iterations with a 40,000

iteration burn-in period. Prior was uniform default distribution. The posterior

distribution was estimated based on samples drawn after every 100 iterations.

We included three species-specific explanatory variables in the model: body

mass (log-transformed; in grams), coloniality (colonial *vs*. solitary; categorical variable

with two levels), nesting site (in trees or on cliffs *vs*. ground; categorical variable with

two levels) and sampling bias (number of publications). We ran models in which

whether a given species performed distraction was set as an independent variable.

Results

Distraction behavior were noted in 55.62% (94/169) of the species, respectively, and

evolved, and were lost multiple times, independently (Fig. 1).

Species with a smaller body mass were more likely to perform distraction

behavior than larger species (Table 2, Fig. 2b). In addition, solitary-nesting species

commonly performed distraction behavior (78/111 species), while species that colonial-

nesting did not so (18/58 species; Table 2) Although distraction behavior was observed

only in one species nesting on cliffs or in trees (1/17 species), the nesting site was not

statistically associated with the occurrence of distraction behavior (93/152 ground-

nesting species).

Discussion

Our analyses suggested that distraction behavior evolved, and were lost, multiple times

independently, although the phylogenetic signal was significant. This suggests that

species-specific patterns of anti-predator behavior are evolutionarily flexible.

Contrary to our prediction, our analysis did not show a significant effect of nest

site on distraction behavior. We do not know why we failed to detect an effect of nest

site, but it might be that an evolutionary transition of nest site (*i.e*., ground-nesting *vs*.

others) was not associated with an evolutionary change in distraction; in other words,

distraction may have evolved within a clade in which all species nest on the ground.

As predicted, body mass explained the inter-specific variation in distraction

behavior. First, We found that small species performed distraction behavior more

frequently than large species. Distraction behavior may have evolved as a substitute for

attack behavior. In line with this idea, it is believed that aggressive nest defense carries

a higher risk of predation for parents than distractive nest defense (D. H. Brunton, 1986;

Gochfeld, 1984; Gómez-Serrano & López-López, 2017; Humphreys & Ruxton, 2020;

Sordahl, 1990b).

As predicted, we found that coloniality explained the inter-specific variation in

distraction behavior, with colonial species are rare to perform distraction behavior.

Previously, Larsen et al. (1996) analyzed factors affecting the performance of

aggressive nest defense across species, and found a relationship between aggressive

defense and coloniality. Colonial species can attack a predator effectively as a group

(Hoogland & Sherman, 1976; Robinson, 1985). By contrast, we found that distraction

behavior was common in solitary-nesting species. Distraction behavior is as an

alternative strategy because solitary species cannot do effective attacks.

In summary, this is the first study to analyze inter-specific variation in the

repertoire of anti-predator behavior during the breeding season using phylogenetic

comparative analyses. Although we succeeded in showing some determinants of anti-

predator behavior, our analyses had several limitations. First, it is true that birds change

anti-predator behavior type according to the type of predator, nesting season, nesting

stage, parental sex, and condition of the parents (Andersson, Wiklund, & Rundgren,

1980; D. Brunton, 1990; Burger et al., 1989; Byrkjedal, 1987, 1989; Caro, 2005;

Courchamp, Clutton-Brock, & Grenfell, 1999; Ghalambor & Martin, 2001; Gochfeld,

1984; Humphreys & Ruxton, 2020; Redondo, 1989; Sordahl, 1990a; Vincze et al.,

2017). We could not control for these factors, as we focused on species-specific

characteristics of the anti-predator behavior. Second, predation pressure on shorebirds

varies with latitude. It is known that species nesting in high latitude area suffer lower

predation risk than low latitude one (Kubelka et al., 2018; McKinnon et al., 2010).

Therefore, the latitude of the nesting site may affect anti-predator behavior. This study,

however, did not examined the effect of latitude because of multi-collinearity to body

weight. That is, it is known that species breeding in higher latitudes have larger bodies

(Olson et al., 2009). Third, our database could have underestimated the occurrence of

distraction behavior, as we treated papers that did not describe anti-predator behavior as

showing an absence of anti-predator behavior. The accumulation of detailed behavioral

data for individual species, and comparative analyses based on an updated database, will

be necessary to elucidate the details of interspecific variation in anti-predator behavior

in shorebirds. Similar studies should also examine the determinants of anti-predator

behavior in taxa in which multiple types of anti-predator behavior have evolved.

Compliance with ethical standards

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethical approval

Ethical approval is not applicable to the literature survey. Behavioral observation at the

Morigasaki Water Recycling Center was conducted with the permission from the

Ministry of the Environment of Japan and the Tokyo Metropolitan Government Bureau

of Environment. This study was conducted under Japanese laws and the guidelines of

Japanese Ethological Society.

References

Andersson, M., Wiklund, C. G., & Rundgren, H. (1980). Parental defence of offspring:

A model and an example. *Animal Behaviour*, *28*(2), 536–542.

https://doi.org/10.1016/S0003-3472(80)80062-5

Brunton, D. (1990). The effects of nesting stage, sex, and type of predator on parental

defense by killdeer (Charadrius vociferous): testing models of avian parental

defense. *Behavioral Ecology and Sociobiology*, *26*(3), 181–190.

https://doi.org/10.1007/BF00172085

Brunton, D. H. (1986). Fatal antipredator behavior of a Killdeer. *Wilson Bulletin*, *98*(98),

605–607.

Burger, J., Saliva, J. E., Gochfeld, M., Gochfeld, D., Gochfeld, D., & Morales, H.

(1989). Antipredator Behaviour In Nesting Zenaida Doves (Zenaida Aurita):

Parental Investment Or Offspring Vulnerability. *Behaviour*, *111*(1–4), 129–143.

https://doi.org/10.1163/156853989X00628

Byrkjedal, I. (1987). Antipredator Behavior and Breeding Success in Greater Golden-

Plover and Eurasian Dotterel. *The Condor*, *89*(1), 40.

https://doi.org/10.2307/1368758

Byrkjedal, I. (1989). Nest defense behavior of lesser golden-plovers. *Wilson Bulletin*,

101(4), 579–590. https://doi.org/10.2307/4162787

Caro, T. (2005). *Antipredator defenses in birds and mammals*.

Clark, A. B., & Wilson, D. S. (1981). Avian Breeding Adaptations: Hatching

Asynchrony, Brood Reduction, and Nest Failure. *The Quarterly Review of Biology*,

56(3), 253–277. https://doi.org/10.1086/412316

Coulson, J. C., & Thomas, C. S. (1985). Changes in the biology of the kittiwake Rissa

tridactyla: a 31-year study of a breeding colony. *Journal of Animal Ecology*, *54*(1),

9–26. https://doi.org/10.2307/4617

Courchamp, F., Clutton-Brock, T., & Grenfell, B. (1999, October 1). Inverse density

dependence and the Allee effect. *Trends in Ecology and Evolution*. Elsevier

Current Trends. https://doi.org/10.1016/S0169-5347(99)01683-3

Cullen, E. (1957). Adaptations in the Kittiwake to cliff-nesting. *Ibis*, *99*(2), 275–302.

https://doi.org/10.1111/j.1474-919X.1957.tb01950.x

da Cunha, F. C. R., Fontenelle, J. C. R., & Griesser, M. (2017). Predation risk drives the

expression of mobbing across bird species. *Behavioral Ecology*, *28*(6), 1517–1523.

https://doi.org/10.1093/beheco/arx111

del Hoyo, J., Elliott, A., & Sargatal, J. (1998). *Handbook of the birds of the world, vol.*

3. Hoatzin to Auks. Biological Conservation (Vol. 86).

https://doi.org/10.1016/S0006-3207(98)00049-4

Dunning, J. B. (2007). *CRC handbook of avian body masses, second edition. CRC*

Handbook of Avian Body Masses, Second Edition. CRC Press.

https://doi.org/10.1201/9781420064452

Fritz, S. A., & Purvis, A. (2010). Selectivity in mammalian extinction risk and threat

types: A new measure of phylogenetic signal strength in binary traits.

Conservation Biology, *24*(4), 1042–1051. https://doi.org/10.1111/j.1523-

1739.2010.01455.x

Ghalambor, C. K., & Martin, T. E. (2001). Fecundity-survival trade-offs and parental

risk-taking in birds. *Science*, *292*(5516), 494–497.

https://doi.org/10.1126/science.1059379

Gochfeld, M. (1984). Antipredator behavior: Aggressive and distraction displays of

shorebirds. In *Shorebirds: Breeding Behaviour and populations. Behavior of*

Marine Animals. Vol. 5 (pp. 289–377). Boston, MA: Springer US.

https://doi.org/10.1007/978-1-4684-4691-3_8

Gómez-Serrano, M. Á. (2018). Broken Wing Display. *Encyclopedia of Animal*

Cognition and Behavior, 1–3. https://doi.org/10.1007/978-3-319-47829-6_2007-2

Gómez-Serrano, M. Á., & López-López, P. (2017). Deceiving predators: linking

distraction behavior with nest survival in a ground-nesting bird. *Behavioral*

Ecology, *28*(1), 260–269. https://doi.org/10.1093/beheco/arw157

Guillerme, T., & Healy, K. (2014). mulTree: a package for running MCMCglmm

analysis on multiple trees. *ZENODO. 10.5281/Zenodo.12902.*

https://doi.org/10.5281/zenodo.12902

Hadfield, J. (2010). MCMC methods for Multi–response Generalised Linear Mixed

Models: The MCMCglmm R Package. *Journal of Statistical Software*, (33), 1–22.

https://doi.org/10.1002/ana.22635

Hamilton, W. D. (1971). Geometry for the selfish herd. *Journal of Theoretical Biology*,

31(2), 295–311. https://doi.org/10.1016/0022-5193(71)90189-5

Hogan, B. G., Hildenbrandt, H., Scott-Samuel, N. E., Cuthill, I. C., & Hemelrijk, C. K.

(2017). The confusion effect when attacking simulated three-dimensional starling

flocks. *Royal Society Open Science*, *4*(1), 160564.

https://doi.org/10.1098/rsos.160564

Hoogland, J. L., & Sherman, P. W. (1976). Advantages and Disadvantages of Bank

Swallow (Riparia riparia) Coloniality. *Ecological Monographs*, *46*(1), 33–58.

https://doi.org/10.2307/1942393

Humphreys, R. K., & Ruxton, G. D. (2020). Avian distraction displays: a review. *Ibis*.

Blackwell Publishing Ltd. https://doi.org/10.1111/ibi.12814

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global

diversity of birds in space and time. *Nature*, *491*(7424), 444–448.

https://doi.org/10.1038/nature11631

Kubelka, V., Šálek, M., Tomkovich, P., Végvári, Z., Freckleton, R. P., & Székely, T.

(2018). Global pattern of nest predation is disrupted by climate change in

shorebirds. *Science (New York, N.Y.)*, *362*(6415), 680–683.

https://doi.org/10.1126/science.aat8695

Lack, D. (1954). *The natural regulation of animal numbers*. *Journal of Human*

Evolution (Vol. 1). https://doi.org/https://doi.org/10.1163/156853989X00646

Larsen, T., Sordahl, T. A., & Byrkjedal, I. (1996). Factors related to aggressive nest

protection behavior: A comparative study of Holarctic waders. *Biological Journal*

of the Linnean Society, *58*(4), 409–439. https://doi.org/10.1006/bijl.1996.0045

McKinnon, L., Smith, P. A., Nol, E., Martin, J. L., Doyle, F. I., Abraham, K. F., … Bêty,

J. (2010). Lower predation risk for migratory birds at high latitudes. *Science*,

327(5963), 326–327. https://doi.org/10.1126/science.1183010

Olson, V. A., Davies, R. G., Orme, C. D. L., Thomas, G. H., Meiri, S., Blackburn, T. M.,

… Bennett, P. M. (2009). Global biogeography and ecology of body size in birds.

Ecology Letters, *12*(3), 249–259. https://doi.org/10.1111/j.1461-

0248.2009.01281.x

Orme, D. (2013). The caper package: comparative analysis of phylogenetics and

evolution in R. http://cran.r-project.org/web/packages/caper/vignettes/caper.pdf. *R*

Package Version, *5*(2), 1–36.

Paton, P. W. C., Messina, F. J., & Griffin, C. R. (1994). A Phylogenetic Approach to

Reversed Size Dimorphism in Diurnal Raptors. *Oikos*, *71*(3), 492–498.

https://doi.org/10.2307/3545837

R Development Core Team. (2019). R: A Language and Environment for Statistical

Computing. *R Foundation for Statistical Computing*, *1*(2.11.1), 409.

https://doi.org/10.1007/978-3-540-74686-7

Redondo, T. (1989). Avian Nest Defence: Theoreticals Models and Evidence.

Behaviour, *111*(1–4), 161–195. https://doi.org/10.1163/156853989X00646

Ristau, C. A. (1991). Aspects of the Cognitive Ethology of an Injury-Feigning Bird, The

Piping Plover. In *Cognitive ethology: The minds of other animals: Essays in honor*

of Donald R Griffin Comparative cognition and neuroscience (pp. 111–146).

Psychology Press. https://doi.org/10.4324/9780203761700-11

Robinson, S. K. (1985). Coloniality in the yellow-rumped cacique as a defence against

nest predators. *Auk*, *102*(3), 506–519. Retrieved from

http://www.jstor.org/stable/4086646

Royle, N. J., Smiseth, P. T., & Kölliker, M. (2012). *The evolution of parental care*.

Oxford University Press.

https://doi.org/10.1093/acprof:oso/9780199692576.003.0001

Sordahl, T. A. (1990a). Sexual Differences in Antipredator Behavior of Breeding

American Avocets and Black-Necked Stilts. *The Condor*, *92*(2), 530.

https://doi.org/10.2307/1368255

Sordahl, T. A. (1990b). The Risks of Avian Mobbing and Distraction Behavior: An

Anecdotal Review. *The Wilson Bulletin*, *102*(2), 349–352.

https://doi.org/10.2307/4162880

Szekely, T., & Reynolds, J. D. (1997). The evolution of parental care in shorebirds: life

histories, ecology, and sexual selection. *Behavioral Ecology*, *8*(2), 126–134.

Székely, T., Reynolds, J. D., & Figuerola, J. (2000). Sexual size dimorphism in

shorebirds, gulls, and alcids: The influence of sexual and natural selection.

Evolution, *54*(4), 1404–1413. https://doi.org/10.1111/j.0014-3820.2000.tb00572.x

Vincze, O., Kosztolányi, A., Barta, Z., Küpper, C., Alrashidi, M., Amat, J. A., …

Székely, T. (2017). Parental cooperation in a changing climate: fluctuating

environments predict shifts in care division. *Global Ecology and Biogeography*,

26(3), 347–358. https://doi.org/10.1111/geb.12540

Table 1. Phylogenetically controlled MCMC generalized linear mixed models

examining the effect of body mass (log-transformed; in grams), nesting site (0 = trees or

cliffs, 1 = ground) and coloniality (0 = solitary, 1 = colonial) of distraction behavior.

Means and 95% credible intervals (CIs; in parentheses) of the posterior distribution are

shown. Results in bold are statistically significant.

	Posterior mean Z	95% CI
Intercept	1.158	0.512 to 1.805
Body mass	-0.120	-0.217 to -0.021
Nesting site	0.192	-0.051 to 0.438
Coloniality	-0.218	-0.412 to -0.024
Sampling bias	0.000	-0.001 to 0.001
Random effect: Phylogeny	0.133	0.032 to 0.299

Figure legends

Fig. 1

Presence of distraction behavior layered on a phylogenetic tree. Solid, performs; white,

does not perform.

356
357

363   Fig. 2

364   Effect of body mass (log-transformed; in grams) on distraction behavior. The number of

365   species is indicated by the thickness of the violin plot.

367