TBPL2/TFIIA complex establishes the maternal transcriptome by an oocyte-specific promoter usage

Changwei Yu[1,2,3,4]**, Nevena Cvetesic**[5]**, Vincent Hisler**[1,2,3,4]**, Kapil Gupta**[6]**, Tao Ye**[1,2,3,4]**,**

Emese Gazdag[1,2,3,4]**, Luc Negroni**[1,2,3,4]**, Petra Hajkova**[5]**, Imre Berger**[6]**, Boris Lenhard**[5]**,**

Ferenc Müller[7]**, Stéphane D. Vincent**[1,2,3,4,8,*]** and László Tora**[1,2,3,4,8,*]

[1]Institut de Génétique et de Biologie Moléculaire et Cellulaire, 67404 Illkirch, France;

[2]Centre National de la Recherche Scientifique (CNRS), UMR7104, 67404 Illkirch, France;

[3]Institut National de la Santé et de la Recherche Médicale (INSERM), U1258, 67404 Illkirch, France;

[4]Université de Strasbourg, 67404 Illkirch, France;

[5]Imperial College London, South Kensington Campus, London SW7 2AZ, UK;

[6]Max Planck Bristol Centre for Minimal Biology, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK.

[7]Institute of Cancer and Genomic Sciences, College of Medical and Dental Sciences, University of Birmingham, UK;

[8] These authors contributed equally to this work

[*]Co-corresponding authors: S.D.V. vincent@igbmc.fr and L.T. laszlo@igbmc.fr

Abstract

During oocyte growth, transcription is required to create RNA and protein reserves to achieve maternal competence. During this period, the general transcription factor TATA binding protein (TBP) is replaced by its paralogue, TBPL2 (TBP2 or TRF3), which is essential for RNA polymerase II transcription. We show that in oocytes TBPL2 does not assemble into a canonical TFIID complex. Our transcript analyses demonstrate that TBPL2 mediates transcription of oocyte-expressed genes, including mRNA survey genes, as well as specific endogenous retroviral elements. Transcription start site (TSS) mapping indicates that TBPL2 has a strong preference for TATA-like motif in core promoters driving sharp TSS selection, in contrast with canonical TBP/TFIID-driven TATA-less promoters that have broader TSS architecture. Thus, we show a role for the TBPL2/TFIIA complex in the establishment of the oocyte transcriptome by using a specific TSS recognition code.

    Introduction

Regulation of transcription initiation by RNA Polymerase II (Pol II) is central to all

developmental processes. Pol II transcription requires the stepwise assembly of multi-protein

complexes called general transcription factors (GTFs) and Pol II[1]. The evolutionary conserved

TFIID complex plays a major role in transcription initiation as it is the first GTF to initiate the

assembly of the pre-initiation complex (PIC) by recognizing the core promoter[2]. TFIID is a

large multi-protein complex composed of the TATA box-binding protein (TBP) and 13 TBP-

associated factors (TAFs) in metazoa[3]. The model suggesting that transcription is always

regulated by the same transcription complexes has been challenged in metazoans by the

discovery of cell-type specific complexes containing specialized GTF-, TBP- or TAF-

paralogs[4]. Two TBP paralogues have been described in vertebrates: TBPL1 (TBP like factor;

TLF, also known as TRF2) has been identified in all metazoan species[5-10], while TBPL2 (also

known as TRF3 or TBP2) has only been described in vertebrates[11,12]. Remarkably, while

Tbpl1 and *Tbpl2* mutants display embryonic phenotypes in non-mammalian species[7-10,12,13],

Tbpl1 and *Tbpl2* loss of function in mouse results in male and female sterility, respectively[14-

16], suggesting that in mammals, these two TBP-like proteins are involved in cell specific

transcription. While TBPL2 shares high degree of identity (92%) within the conserved saddle-

shaped C-terminal DNA binding core domain of TBP[17], the C-terminus of TBPL1 is more

distant with only 42% identity[12]. A consequence is that TBPL2, but not TBPL1, is able to

bind canonical TATA-box sequences *in vitro*[5,12,18]. The N-terminal domains of the three

vertebrate TBP related factors do not show any conservation. All three vertebrate TBP related

factors can interact with the GTFs TFIIA and TFIIB, and can mediate Pol II transcription

initiation *in vitro*[12,13,18-20]. However, how alternative initiation complexes form, how they

regulate cell type-specific transcription and how they recognize promoter sequences remain

unknown.

Mapping of transcription start sites (TSSs), at single-nucleotide by Cap Analysis of

Gene Expression (CAGE) revealed two main modes for transcription start site (TSS) usage[21].

Transcription initiation within a narrow region, called "sharp" (or focused) TSS-type, is

common in highly active, tissue-specific gene promoters containing TATA boxes, while

transcription initiation with multiple initiation positions within an about 100 bp region, called

"broad" TSS promoter architecture[21], are more characteristic to ubiquitously expressed and

developmental regulated genes (reviewed in [22]). During zebrafish maternal to zygotic

transition it was described that two TSS-defining grammars coexist, in core promoters of

constitutively expressed genes to enable their expression in the two regulatory

environments[23]. Maternally active promoters in zebrafish tend to be sharp, with TATA-like,

AT-rich (W-box) upstream elements guiding TSS selection, while embryonically active broad

promoter architectures of the same genes appear to be regulated by nucleosome positioning

ques. Although a number of germ cell specific as well as somatic transcriptional regulators

have been well characterized during folliculogenesis (reviewed in [24]), the exact actors and

mechanisms required for setting up the oocyte specific transcriptome have not yet been

identified in vertebrates.

Female germ cells develop during oogenesis leading to the formation of a highly

differentiated and specialised cell, the oocyte. In females, oocytes enter meiosis during

embryonic life. Quiescent primordial follicles composed of meiotically arrested oocytes at the

late diplotene stage and surrounded by granulosa cells, are formed perinatally in mice

(reviewed in [24]). Shortly after birth, some primordial follicles enter folliculogenesis and

undertake a growth phase during which a specific oocyte-specific transcriptome is set up,

oocytes increase their size until the pre-antral follicular stage[25]. A remarkable feature of

oocyte is the very high expression of retrotransposons driven by Pol II transcription. These

elements are interspersed repetitive elements that can be mobile in the genome. One of the 3

major classes of retrotransposons in mammals is the long terminal repeat (LTR)

retrotransposons derived from retroviruses, also known as endogenous retroviruses (ERVs)

that is subdivided in 3 sub-classes: ERV1, ERVK and endogenous retrovirus like ERVL-

MaLR (mammalian apparent LTR retrotransposons) (reviewed in [26]). Transcription of mobile

elements in specific cell types depends on the presence of a competent promoter recognition

transcription machinery and/or the epigenetic status of the loci where these elements have

been incorporated. Remarkably, MaLRs encode no known proteins, but MaLR-dependent

transcription is key in initiating synchronous developmentally regulated transcription to

reprogram the oocyte genome during growth[27].

Remarkably, during oocyte growth TBP is absent and replaced by TBPL2[28]. Indeed,

TBP is only expressed up to the primordial follicular oocytes and becomes undetectable at all

subsequent stages during oocyte growth. In contrast, TBPL2 is highly expressed in the

growing oocytes, suggesting that TBPL2 is replacing TBP for its transcription initiating

functions during folliculogenesis[28]. In agreement with its oocyte-specific expression, a crucial

role of TBPL2 for oogenesis was demonstrated in *Tbpl2*[-/-] females, which show sterility due

to defect in secondary follicle production[16,29]. In the absence of TBPL2, immunofluorescent

staining experiments showed that elongating Pol II and histone H3K4me3 methylation signals

were abolished between the primary and secondary follicle stage oocytes suggesting that Pol

II transcription was impaired[16]. Initially TBPL2/TRF3 was suggested to be expressed during

muscle differentiation[30], but this observation was later invalidated[16,29]. Altogether, the

available data suggested that TBPL2 is playing a specialized role during mouse oocyte

development. However, how does TBPL2 regulate oocyte-specific transcription and what is

the composition of associated transcription machinery, remained unknown.

Here we demonstrate that in oocytes TBPL2 does not assemble into a canonical TFIID

complex, while it stably associates with TFIIA. The observation that the oocyte specific

deletion of *Taf7*, a TFIID specific TAF, does not influence oocyte growth and maturation,

corroborates the lack of TFIID in growing oocytes. Our transcriptomics analyses in wild type

and *Tbpl2*$^{-/-}$ oocytes show that TBPL2 mediates transcription of oocyte-expressed genes,

including mRNA destabilisation factor genes, as well as MaLR ERVs. Our transcription start

site (TSS) mapping from wild-type and *Tbpl2*$^{-/-}$ growing oocytes demonstrates that TBPL2

has a strong preference for TATA-like motif in gene core promoters driving specific sharp

TSS selection. This is in marked contrast with TBP/TFIID-driven TATA-less gene promoters

in preceding stages that have broad TSS architecture. Our results show a role for the TBPL2-

TFIIA transcription machinery in a major transition of the oocyte transcriptome mirroring the

maternal to zygotic transition that occurs after fertilization, completing a full germ line cycle.

Results

Oocyte-specific TBPL2/TFIIA complex distinct from TFIID

To characterize TBPL2-containing transcription complexes we prepared whole cell

extracts (WCE) from growing 14 days post-natal (P14) mouse ovaries and analysed TBPL2-

associated proteins by anti-mTBPL2 immunoprecipitation (IP) coupled to label free mass

spectrometry (Fig. 1a, Supplementary Fig. 1a, b). To determine the stoichiometry of the

composition of the immunoprecipitated complexes, normalized spectral abundance factor

(NSAF) values were calculated[31]. In the anti-TBPL2 IPs we identified TFIIA-αβ and TFIIA-γ

subunits as unique GTF subunits associated with TBPL2 (Fig. 1a, Supplementary Data 1). As

ovaries contain many other non-oocyte cell types which express TBP, in parallel from the

same extracts we carried out an anti-TBP IP. The mass spectrometry of the anti-TBP IP

indicated that TBP assembles into the canonical TFIID complex in non-oocyte cells (Fig. 1b,

Supplementary Data 2). As growing oocytes represent only a tiny minority of ovary cells, we

further tested the TBPL2-TFIIA interaction by a triple IP strategy (Fig. 1c): first, we depleted

TAF7-containing TFIID complexes with an anti-TAF7 IP, second, the remaining TFIID and

SAGA complexes, which contain also shared TAFs[32], were depleted with an anti-TAF10 IP

using the anti-TAF7 IP flow-through as an input, third we performed an anti-TBPL2 IP on the

anti-TAF7/anti-TAF10 flow-through fraction (Fig. 1d-f, Supplementary Data 3). The analysis

of this third consecutive IP further demonstrated that TBPL2 forms a unique complex with

TFIIA-αβ, and TFIIFA-γ, but without any TFIID subunits.

To further analyse the requirement of TFIID during oocyte growth, we carried out a

conditional depletion of TFIID-specific *Taf7* gene during oocyte growth using the *Zp3-Cre*

transgenic line[33] (Supplementary Fig. 1c-g). Remarkably, TAF7 is only detected in the

cytoplasm of growing oocyte (Supplementary Fig. 1c). The oocyte-specific deletion of *Taf7*

did not affect the presence of secondary and antral follicles and the numbers of collected

mature oocytes after superovulation (Fig. 1g, h, Supplementary Fig. 1f). The lack of

phenotype is not due to an inefficient deletion of *Taf7*, as TAF7 immunolocalization is

impaired (Supplementary Fig. 1d, e), and as oocyte-specific *Taf7* mutant females are severely

hypofertile (Supplementary Fig. 1g). The observations that TBP is not expressed in growing

oocytes, and that the oocyte specific deletion of *Taf7* abolishes the cytoplasmic localization of

TAF7, but does not influence oocyte growth, show that canonical TFIID does not assemble in

the nuclei of growing oocytes. Thus, our results together demonstrate that during oocyte

growth a stable TBPL2-TFIIA complex forms, and may function differently from

TBP/TFIID.

In order to further characterize the composition of the TBPL2-TFIIA complex, we

took advantage of NIH3T3 cells artificially overexpressing TBPL2 (NIH3T3-II10 cells[28]). In

this context where TBP and TAFs are present, TFIID is efficiently pulled down by an anti-

TBP IP, but no interaction with TFIIA could be detected (Fig. 2a). Interestingly, the anti-

TBPL2 IP showed that the artificially expressed TBPL2 can incorporate in TFIID-like

complexes as TAFs were co-IP-ed (Fig. 2a), however with much lower stoichiometry (NSAF

values) than that of TBP (Fig. 2b). In contrast, strong interaction with TFIIA-$\alpha\beta$ and TFIIFA-

γ were detected, suggesting that the TBPL2-TFIIA complex can be formed in the NIH3T3-

II10 cells and that TBPL2, to the contrary to TBP has the intrinsic ability to interact with

TFIIA. Remarkably, in spite of the high similarity between the core domains of TBP and

TBPL2, no interaction with SL1 (TAF1A-D) and TFIIIB (BRF1) complexes that are involved

in Pol I and Pol III transcription, respectively[34]) could be detected in the anti-TBPL2 IP, to the

contrary to the anti-TBP IP in the NIH3T3-II10   (Fig. 2a, b) as well as in the ovary

(Supplementary Data 1, 2), suggesting that TBPL2 is not involved in Pol I and Pol III

transcription initiation in the growing oocytes.

To analyze whether TBPL2 associates with TFIID TAFs and TFIIA in the same

complex, we performed a gel filtration analysis of NIH3T3-II10 WCE. The profile indicated

that most of the TBPL2 and TFIIA could be found in the same fractions (22-26) eluting

around 150-200 kDa, while TBPL2 protein was below the detection threshold of the western

blot assay in the TAF6-containing fractions 9-15 (Fig. 2c). To verify that TBPL2 and TFIIA

are part of a same complex in fractions 22-26, we IP-ed TBPL2 from these pooled fractions

and subjected them to mass spectrometric analysis. Our data confirmed that in these fractions

eluting around 170 kDa, TBPL2 and TFIIA form a stable complex that does not contain any

TAFs (Fig. 2d, supplementary Data 4). Thus, all these experiments together demonstrate that

TBPL2/TFIIA form a stable complex in oocytes, where TBP is not expressed and

TBPL2/TFIIA is the only promoter recognizing transcription complex that could direct Pol II

transcription initiation (see the summary of all these IPs in Fig. 2e).

TBPL2-dependent oocyte transcriptome

To characterize the growing oocyte-specific transcriptome and its dependence on

TBPL2, we have performed a transcriptomic analysis of wild-type (WT) and *Tbpl2*[-/-] oocytes

isolated from primary (P7) and secondary follicles and (P14) (Fig. 3, 4, Supplementary Fig. 2,

Supplementary Data 5). We observed down-regulation of a high number of oocyte-specific

genes, out of which *Bmp15* and *Gdf9* served as internal controls[35,36], as they were already

described to be regulated by TBPL2[16] (Fig. 3a, b and Supplementary Fig. 2a). Principal

component analysis showed that the four distinct RNA samples clustered in individual groups

and that the main explanation for the variance is the genotype, and then the stage

(Supplementary Fig. 2b). Comparison of the RNA level fold changes between mutant and WT

oocytes showed that in *Tbpl2*[-/-], there is a massive down-regulation of the most highly

expressed transcripts, both at P7 and P14 (Supplementary Fig. 2c). The Pearson correlation

between the P7 and P14 fold change data sets for transcripts expressed above 100 normalized

reads was close to 0.8 (Supplementary Fig. 2c), indicating that *Tbpl2* loss of function

similarly altered RNA levels at P7 and P14 stages. We therefore focused on the P14 stage for

the rest of the study.

In WT P14 oocytes transcripts corresponding to 10791 genes were detected.

Importantly, many of these detected transcripts have been transcribed at earlier stages and are

stored in growing oocytes[37]. As there is no Pol II transcription in *Tbpl2*[-/-] growing oocytes[16],

RNAs detected in the *Tbpl2*[-/-] mutant oocytes represent mRNAs transcribed by a TBP/TFIID-

dependent mechanism and deposited into the growing oocytes independently of TBPL2

activity at earlier stages, i.e. at primordial follicular stage, where TBP is still expressed. The

proportion of genes (1396) upregulated following *Tbpl2* deletion (Fig. 3c) can be explained

by two mutually not exclusive ways: i) the consequence of the normalization to the library

size resulting in a slight over-estimation of up-regulated transcripts, and ii) under-estimation

of down-regulated transcripts and/or by transcript buffering mechanisms due to mRNA

stabilization[38]. Validation of the up-regulation of some candidate transcripts levels

(Supplementary Fig. 2d, e) strongly supports the latter hypothesis (but see also the next

paragraph).

Nevertheless, we detected 1802 significantly downregulated transcripts in the *Tbpl2*[-/-]

oocytes (Fig. 3c). Key genes known to be expressed during oocyte growth, such as *Bmp15,*

Eloc, Fgf8, Gdf9 and *Zar1*[35,36,39], were confirmed by RT-qPCR to be down-regulated

(Supplementary Fig. 2f, g). These results suggest that TBPL2 has an important role in gene

expression in the growing oocytes. Gene Ontology (GO) analyses of biological process of the

identified down regulated categories of genes (Supplementary Data 6) indicated that many

genes, involved in meiosis II and distinct cell cycle processes, were significantly down-

regulated (Supplementary Fig. 2h). The most enriched molecular function GO category was

"poly(A)-specific ribonuclease activity" containing many genes coding for factors or subunits

of complexes contributing to deadenylation/decapping/decay activity in eukaryotes (Fig. 3d)

(i.e. CCR4-NOT, PAN2/PAN3[40]; DCP1A/DCP2[41], or BTG4[39]). In good agreement with the

transcriptome analyses, transcripts coding for these "poly(A)-specific ribonuclease activity"

factors were significantly down regulated in *Tbpl2*[-/-] mutant P14 oocytes when tested by RT-

qPCR (Fig. 3e, Supplementary Fig. 2i). Thus, in P14 oocytes TBPL2 is regulating the

transcription of many genes coding for factors, which are in turn crucial in regulating the

stability and translation of the mRNA stock deposited during early oogenesis, as well as

transcription of meiosis II- and cell cycle-related genes to prepare the growing oocytes for the

upcoming meiotic cell division.

A remarkable feature of oocyte is the very high expression of retrotransposons driven

by Pol II transcription (see Introduction). As expected, in WT P7 and P14 oocytes the

expression of ERVs was found to be the most abundant[27,42] (Supplementary Fig. 3a-c).

Importantly, the transcription of the vast majority of MaLR elements was the most affected in

Tbpl2[-/-] mutant oocytes at P7 and P14 (Fig. 4). Among them, three highly expressed members,

MT-int, *MTA_Mm*, and *MTA_Mm-int*, were dramatically down-regulated in P7 and P14

Tbpl2[-/-] mutant oocytes (Supplementary Fig. 3d, e). As in P14 oocytes TBPL2 depletion is

reducing transcription more than 4-fold from MaLR ERVs, which often serve as promoters

for neighbouring genes[27,42], TBPL2 could seriously deregulate oocyte-specific transcription

and consequent genome activation.

Therefore, this is the first demonstration that TBPL2 is orchestrating the *de novo*

restructuration of the maternal transcriptome and that TBPL2 is crucial for indirectly silencing

the translation of the earlier deposited TBP-dependent transcripts.

TBPL2-driven promoters contain TATA box and are sharp

The promoter usage changes during zebrafish maternal to zygotic transition revealed

different rules of transcriptional initiation in oocyte and in embryo, driven by independent and

often overlapping sets of promoter "codes" [23]. Importantly, this switch has not yet been

demonstrated in mammals and the role of TBPL2 in this switch during oogenesis remained to

be investigated. To this end, we mapped the TSS usage by carrying out super-low input

carrier-CAGE (SLIC-CAGE) [43] from WT and *Tbpl2*[-/-] P14 oocytes. To characterize only the

TBPL2-driven promoters, we removed the CAGE tags present in the *Tbpl2*[-/-] dataset from the

WT P14 dataset, to eliminate transcripts that have been deposited at earlier stages (hereafter

called "TBPL2-dependent"). Conversely, the *Tbpl2*[-/-] dataset corresponds to the TBP/TFIID-

dependent, or TBPL2-independent TSSs (hereafter called "TBPL2-independent").

Next, we analysed the genome-wide enrichment of T- and/or A-rich (WW) dinucleotide

motifs within the -250/+250 region centred on the dominant TSSs of the TBPL2-specific-only

and TBPL2-independent-only oocyte TSS clusters (Fig. 5a, b). TBPL2-dependent TSS

clusters are strongly enriched in a well-defined WW motif around their -30 bp region (Fig. 5a,

red arrowhead) [44]. In contrast, only about 1/3[rd] of the TBPL2-independent TSS clusters

contained WW-enriched motifs at a similar position (Fig. 5b, red arrowhead), as would be

expected from promoters that lack maternal promoter code determinants[23,44]. As canonical

TATA boxes are often associated with tissue-specific gene promoters, we investigated

whether the above observed WW motif densities correspond to TATA boxes using the TBP

position weight matrix (PWM) from the JASPAR database as a reference. To this end the

presence of TATA boxes was analysed in the TSS clusters of the two data sets and revealed

that TBPL2-dependent TSS clusters were enriched in high quality TATA boxes, including a

clear increase in the proportion of canonical TATA boxes, when compared to TBPL2-

independent TSS clusters (Fig. 5c). Genome browser view snapshots indicate that TSS

clusters in P14 WT oocytes tend to be sharp and are associated with TATA-like motifs

(Supplementary Fig. 4a, b). Analysis of the global distribution of the number of TSSs and of

the width of the TSS clusters in the above defined two categories confirmed that TBPL2-

dependent TSS are sharper compared to the TBPL2-independent TSS clusters (Supplementary

Fig. 4c, d).

In order to test whether TBPL2 controls transcription initiation from maternal promoter

code determinants, we grouped the expression profiles corresponding to each consensus TSS

clusters, to characterise promoter activity profiles among datasets by performing self-

organizing maps (SOM) [45] (Supplementary Fig. 4e). We then focussed on the two most

distinct SOM groups: the down-regulated promoters (blue group, containing 9442 consensus

TSS clusters) (Fig. 5d) and the up-regulated promoters (red group, with 6900 consensus TSS

clusters) (Fig. 5e). Motif analyses of these two categories of promoters in their -35/+5 regions

relative to the different dominant TSSs indicated that only the core promoters associated to

TBPL2-dependent dominant TSSs belonging to the down-regulated gene promoters contain a

well-defined 7 bp long TATA box-like motif (W-box) in their -31 to -24 regions (Fig. 5f, g,

Supplementary Fig. 4f-i). Importantly, W-box associated TSSs architecture usage distribution

for these TBPL2-dependent dominant TSSs was sharp (Supplementary Fig. 4j, l), as expected

for motif-dependent transcriptional initiation[23,44]. In contrast, TBPL2-independent TSSs

belonging to the up-regulated promoters exert a much broader TSS pattern (Supplementary

Fig. 4k, m). Interestingly, GO analyses of the genes associated with the down-regulated

promoters revealed a strong association with deadenylation/decapping/decay activity

(Supplementary Fig. 4n-p, Supplementary Data 7), further confirming our initial RNA-seq

analysis observations (Fig. 3).

Importantly, TSS architecture analyses of the TBPL2-dependent MaLR ERV TSSs

indicated that the majority of MaLR core promoters contain high quality TATA box motif

(median of the TATA box PWM match is 85%, Fig. 5h-j). These observations together

demonstrate that the TBPL2/TFIIA complex drives transcription initiation primarily from

core promoters that contain a TATA box-like motif in their core promoter and directs sharp

transcription initiation from the corresponding promoter regions to overhaul the growing

oocyte transcriptome.

In addition, we observed that TSS usage can shift within the promoter of individual

genes depending on the genetic background (Supplementary Fig. 4b). To get more insights in

these promoter architecture differences, we identified genome-wide 6429 shifting promoters

by comparing either TBPL2-dependent to TBPL2-independent TSS data. These results are

consistent with TSS shifts between somatic and maternal promoter codes occurring either in

5', or 3' directions (Fig. 6a, Supplementary Fig. 4q) [44]. WW motif analysis indicated that on

each shifting promoter, TBPL2-dependent dominant TSSs are associated with WW motifs,

while TBPL2-independent dominant TSSs are not (Fig. 6b). In addition, the TATA box PWM

match analyses indicated that these WW motifs are enriched in TATA box like elements

compared to the corresponding TBPL2-independent shifting TSSs (Fig. 6c). Thus, our

experiments provide a direct demonstration that TBP/TFIID and TBPL2/TFIIA machineries

recognize two distinct sequences co-existing in promoters of the same genes with TBPL2

directing a stronger WW/TATA box-dependent sharp TSS selection in them.

Discussion

In this study, we showed that a unique basal transcription machinery composed of TBPL2 associated with TFIIA is controlling transcription initiation during oocyte growth, orchestrating a transcriptome change prior to fertilization using an oocyte-specific TTS usage.

TBPL2 expression in mouse is limited to the oocytes and in its absence, oocytes fail to grow and *Tbpl2*[-/-] mouse females are sterile[16,28]. In a mirroring situation, TBPL1 (TRF2) expression is enriched during spermatogenesis and male germ cells lacking TBPL1 are blocked between the transition from late round spermatids to early elongating spermatids[14,15]. An interesting parallel between TBPL2 and TBPL1 is that both TBP-type factors form endogenous stable complexes with TFIIA. The beginning of TBPL2 accumulation in the oocyte nuclei, or TRF2 accumulation in male germ cell nuclei coincides with the phase of meiosis I[15,28,46]. It is thus conceivable that TBPL2-TFIIA in oocytes, or TRF2-TFIIA during spermatogenesis are involved in the control of gene expression in a meiotic context to set up the corresponding transcriptome. Interestingly, both transcription complexes seem to function in a compacted chromatin environment in which TBP/TFIID probably cannot. However, while TBPL2 and TBP show contrasting expression patterns in the oocytes[28], TBPL1 and TBP are co-expressed in spermatids[46,47] and it has been suggested that TBPL1 is a testis-specific subunit of TFIIA that is recruited to PIC containing TFIID and might not primarily act independently of TFIID/TBP to control gene expression in round spermatids[48]. While TBPL1 forms a complex also with the TFIIA-αβ paralogue, ALF, in testis[48-50], TBPL2 does not stably associate with ALF, in spite of the fact that it is expressed in oocytes[50].

TBP-like factors are bipartite proteins with variable N-terminal domains and a relatively well conserved shared C-terminal domains (core domains) forming a saddle-like structure with a concave surface that is known to bind to DNA[17]. Interestingly, TBPL1 has a very short N-terminal domain[5,18] suggesting that it lost some abilities to interact with partners.

Our data suggest that despite their very high similarity (92% identity between the core

domains of TBP and TBPL2 (reviewed in [51]), TBP and TBPL2 display different properties as

they seem to recognize different DNA sequences to regulate gene promoters with different

promoter architectures. Our IP-MS analyses from ovary WCE indicate that contrary to TBP,

TBPL2 does not interact with TAFs in growing oocytes. Our analyses in the NIH3T3-II10

cells that overexpresses TBPL2 showed that TBPL2 can interact with TAFs in this artificial

situation, albeit with less affinity compared to TFIIA, or TBP-TAFs interactions. Our

transcriptomic data indicates that all *Taf* mRNAs, except *Taf7l* are detected in growing

oocytes (Supplementary Data 5). However, whether they are also expressed in oocytes at the

protein level is not yet known, except for TAF4B that has been detected in female neonate

oocytes[52]. Nevertheless, our data suggest that TAF7 is expressed, but localized to the

cytoplasm. It could be conceivable that, similarly to *Tbp* mRNA that is transcribed, but not

translated in oocytes[53], *Taf* mRNA translations (other than *Taf7*) are also inhibited and as a

result the canonical TFIID, or its building blocks, cannot be assembled, and as a result the

canonical TFIID is not present in the nuclei of growing oocytes. Another reason why TBPL2

does not interact with TAFs, or ALF, but rather interacts with TFIIA could be its N-terminal

domain that is very different from TBP (only 23% identity[51]).

TBPL2 proteins from different vertebrates show a high degree of similarity in their C-

terminal core domains amongst themselves; but display very little conservation in their N-

terminal domains[12]. It is interesting to note that TBPL2 deficiency leads to an embryonic

phenotypes in Xenopus[13] and zebrafish[12], because, contrary to the mouse, TBPL2 is still

present in the embryo after fertilization and thus may act in parallel with TBP in the

transcription of specific embryonic genes[10,54]. The molecular mechanism by which TBPL2

controls the transcription of these specific sets of genes in frogs and in fish has not been

studied. In contrary, TBPL2 in mammals are only expressed in growing oocytes and the only

phenotype that can be observed in mammals is female sterility[16,29].

LTR retrotransposons, also known as ERVs, constitute ~10% of the mouse genome

(reviewed in [55]). While their expression is generally suppressed by DNA methylation and/or

repressive histone modifications, a subset of ERV subfamilies retain transcriptional activity in

specific cell types [56]. ERVs are especially active in germ cells and early embryos (reviewed in

[26]). Indeed, many genome-wide transcripts are initiated in LTRs, such as for example of

MaLRs in mouse oocytes, which constitute ~5% of the genome[57]. Members of the MT

subfamily of MaLRs are particularly active in oocytes and hundreds of MT LTRs have been

co-opted as oocyte-specific gene promoters[27,58]. As LTR-initiated transcription units shape

also the oocyte methylome, it will be important to analyse also how TBPL2 influences DNA

methylation in oocytes.

Oocytes display remarkable post-transcriptional regulatory mechanisms that control

mRNA stability and translation. During oogenesis, the oocyte genome is transcriptionally

active, and the newly synthesized maternal mRNAs are either translated or stored in a

dormant form (reviewed in [37]). The newly synthesized transcripts receive a long poly(A) tail

and subsequently undergo poly(A) shortening in the oocyte cytoplasm, preventing translation.

Until resumption of meiosis, mRNAs with a short poly(A) tail are stored in the cytoplasm in a

dormant form (for a review, see [59]). Thus, poly(A) tail deadenylation, amongst other

activities, coordinates post-transcriptional regulation of the oocyte mRNA pool. Interestingly,

TBPL2 is regulating the activity of several deadenylation/decapping/decay complexes and in

the absence of TBPL2, we observed apparent stabilization of a significant number of

transcripts, suggesting that in wild type oocytes TBPL2 is indirectly inhibiting the translation

of mRNAs, and/or inducing the degradation of the mRNAs, previously deposited by

TFIID/TBP in the primordial follicular oocytes (Fig. 7). To put in place the growing oocyte

specific maternal transcriptome TBPL2 is controlling the production of new mRNAs using a

maternal specific TSS grammar, as most of these transcripts will remain in the oocyte after

transcriptional quiescence. Remarkably, as TBPL2 does not interact with Pol I and Pol III

transcription machineries in the growing oocytes, this strongly suggest that rRNA and tRNA

are deposited very early during oogenesis in amounts sufficient for the initiation of

development.

Therefore, it seems that TBPL2 contributes to establish a novel TBPL2-dependent

growing oocyte transcriptome and consequent proteome required for further development and

oocyte competence for fertilization (Fig. 7). The indirect regulation of previously deposited

mRNAs by a global transcription regulator resembles the well characterised maternal to

zygotic transition (MZT), during which a clearance of inherited transcriptome is mediated by

de novo gene products generated by newly activated transcription machinery (reviewed in [59]).

At hundreds of gene promoters, two distinct TSS-defining "grammars" co-exist in close

proximity genome-wide and are differentially utilised in either TBPL2/TFIIA in

primary/secondary follicular oocytes, or by TBP/TFIID in primordial follicular oocytes or in

the fertilized embryo. This again shows a striking parallel to MZT[23], where multiple layers of

information are embedded in the same promoter sequence, each representing a different type

of regulatory grammar interpreted by dedicated transcription machinery depending on the

cellular environment.

Methods

Cell lines and cell culture

The NIH3T3-II10 line overexpressing TBPL2 and the control NIH3T3-K2 have already been

described[28] and were maintained in high glucose DMEM supplemented with 10% of new-

born calf serum at 37°C in 5% CO_2.

Whole cell extracts

NIH3T3-II10 and NIH3T3-K2 cells cultured in 15 cm dish were washed twice with 1x PBS,

subsequently harvested by scrapping on ice. Harvested cells were centrifuged 1000 g at 4°C

for 5 min and then resuspended in 1 packed cell volume of whole cell extraction buffer (20

mM Tris HCl pH7.5, 2 mM DTT, 20% Glycerol, 400 mM KCl, 1x Protease inhibitor cocktail

(PIC, Roche)). Cell lysates were frozen in liquid nitrogen and thawed on ice for 3 times,

followed by centrifugation at 20817 g, at 4°C for 15 min. The supernatant was collected and

protein concentration was measured by Bradford protein assay (Bio-Rad). The cell extracts

were used directly for immunoprecipitation and western blot, or stored at −80°C.

Ovaries collected from post-natal day 14 (P14) CD1 and C57BL/6N female mice were

homogenized in whole cell extraction buffer [20 mM Tris HCl pH7.5, 2 mM DTT, 20%

Glycerol, 400 mM KCl, 5x PIC (Roche)]. Cell lysates were frozen in liquid nitrogen and

thawed on ice for 3 times, followed by centrifugation at 20817 g, at 4°C for 15 min. The

supernatant extracts were used directly for immunoprecipitation.

Antibodies and antibody purification

The antibodies are listed in Supplementary Table 1. The IGBMC antibody facility raised the

anti-TBPL2 polyclonal 3024 serum against the CPDEHGSELNLNSNSSPDPQ peptide

(amino acids 111-129) coupled to ovalbumin and injected into one 2 months-old female New-

Zeland rabbit. The resulting serum was affinity purified by using the Sulfolink Coupling Gel

(Pierce) following the manufacturer's recommendations.

Immunoprecipitation

Ovary extract were incubated with anti-GST (10 µg per IP), anti-TBP (10 µg per IP), anti-

TBPL2 (3024, 12 µg (36 µg for gel filtration) per IP), anti-TAF7 (10 µg per IP) or anti-

TAF10 (10 µg per IP) coated Dynabeads (Invitrogen) at 4°C overnight. After incubation,

beads were washed 3 x 5min at 4°C with 500 mM KCl buffer [25 mM Tris-HCl (pH 7.9), 5

mM MgCl2, 10% glycerol, 0.1% NP40, 2 mM DTT, 500 mM KCl and 1x PIC (Roche)], then

washed 3 x 5min at 4°C with 100 mM KCl buffer (25 mM Tris-HCl pH 7.9, 5 mM MgCl2,

10% glycerol, 0.1% NP40, 2 mM DTT, 100 mM KCl and 1x). Immunoprecipitated proteins

were eluted with 0.1 M glycine pH 2.8 and neutralized with 1.5 M Tris-HCl pH 8.8.

Immunoprecipitation performed from whole cell extracts of NIH3T3-II10 and NIH3T3-K2

cells were following the same procedures with protein G Sepharose beads (GE Healthcare):

18 µg rabbit anti-TBPL2 (3024) and 15 µg anti-TBP per IP.

Western blot analyses

Protein samples (15-25 µg of cell extracts or 15 µL of IP elution) were mixed with $1/4^{th}$

volume of loading buffer (100 mM Tris-HCl pH 6.8, 30% glycerol, 4% SDS, 0.2%

bromophenol blue and freshly added 100 mM DTT) and boiled for 10 min. Samples were

then resolved on a 10 % SDS-PAGE and transferred to nitrocellulose membrane (Protran,

Amersham). Membranes were blocked in 3% non-fat milk in 1x PBS at room temperature

(RT) for 30 min, and subsequently incubated with the primary antibody overnight at 4°C

(dilution 1/1000). Membranes were washed three times (10 min each) with 1x PBS - 0.05%

Tween20. Membranes were then incubated with HRP-coupled goat anti Mouse Ig (Jackson

ImmunoResearch, #115-036-071, dilution 1/10000) or HRP-coupled goat anti-Rabbit Ig

(Jackson ImmunoResearch, #115-035-144, dilution 1/10000) for 1 hour at RT, followed by

ECL detection (Thermo Fisher). The signal was acquired with the ChemiDoc imaging system

(Bio-Rad).

Mass spectrometry analyzes and NSAF calculations

Samples were TCA precipitated, reduced, alkylated and digested with LysC and Trypsin at

37°C overnight. After C18 desalting, samples were analyzed using an Ultimate 3000 nano-

RSLC (Thermo Scientific, San Jose, California) coupled in line with a linear trap Quadrupole

(LTQ)-Orbitrap ELITE mass spectrometer via a nano-electrospray ionization source (Thermo

Scientific). Peptide mixtures were loaded on a C18 Acclaim PepMap100 trap column (75 μm

inner diameter × 2 cm, 3 μm, 100 Å; Thermo Fisher Scientific) for 3.5 min at 5 μL/min with

2% acetonitrile (ACN), 0.1% formic acid in H_2O and then separated on a C18 Accucore nano-

column (75 μm inner diameter × 50 cm, 2.6 μm, 150 Å; Thermo Fisher Scientific) with a 240

minutes linear gradient from 5% to 50% buffer B (A: 0.1% FA in H_2O / B: 80% ACN, 0.08%

FA in H_2O) followed with 10 min at 99% B. The total duration was set to 280 minutes at a

flow rate of 200 nL/min.

Proteins were identified by database searching using SequestHT with Proteome Discoverer

1.4 software (Thermo Fisher Scientific) a combined Mus musculus database generated using

Uniprot   [https://www.uniprot.org/uniprot/?query=proteome:UP000000589&sort=score]

(Swissprot, release 2015_11, 16730 entries) where 5 interesting proteins sequences (TrEMBL

entries: TAF4, ATXN7L2, TADA2B, BTAF1 and SUPT3) were added. Precursor and

fragment mass tolerances were set at 7 ppm and 0.5 Da respectively, and up to 2 missed

cleavages   were   allowed.   Oxidation   (M)   was   set   as   variable   modification,   and

Carbamidomethylation© as fixed modification. Peptides were filtered with a false discovery

rate (FDR) at 5 %, rank 1 and proteins were identified with 1 unique peptide. Normalized

spectral abundance factor (NSAF)[31] were calculated using custom R scripts (R software

version 3.5.3). Only proteins detected in at least 2 out of 3 of the technical or biological

replicates were considered for further analyses.

Gel filtration

A Superose 6 (10/300) column was equilibrated with buffer consisting of 25mM Tris HCl pH

7.9, 5 mM MgCl2, 150 mM KCl, 5% Glycerol, 1 mM DTT and 1x PIC (Roche). Five

hundred μL of whole cell extracts containing ∼5 mg of protein were injected in an ÄKTA

avant chromatography system (Cytiva) and run at 0.4 mL/min. Protein detection was

performed by absorbance at 280 nm and 260 nm. Five hundred μL fractions were collected

and analysed by Western blot and IP-MS.

Animal experimentation

Animal experimentations were carried out according to animal welfare regulations and

guidelines of the French Ministry of Agriculture and procedures were approved by the French

Ministry for Higher Education and Research ethical committee C2EA-17 (project

n°2018031209153651). The Tg(*Zp3-Cre*), *Taf7*flox and *Tbpl2*- mouse lines have already been

described[16,33,60].

Histology analyses of ovaries

Ovaries were collected from 6 weeks-old Tg(*Zp3-Cre*/+);*Taf7*flox/+ and Tg(*Zp3-

Cre*/+);*Taf7*flox/Δ oocyte specific mutant females, fixed in 4% paraformaldehyde (Electron

Microscopy Sciences) over-night at 4°C, washed 3 times in PBS at room temperature and

embedded in paraffin. Five μm-thick sections were stained with haematoxylin and eosin and

images were acquired using a slide scanner Nanozoomer 2.0HT (Hamamatsu Photonics).

Immunolocalization of TAF7 in the oocytes

Ovaries were dissected in PBS, fixed overnight in 4%PFA/PBS at 4°C, rinsed three times in

PBS, equilibrated in 30% sucrose/PBS and embedded in Cryomatrix (Thermo Scientific) in

liquid nitrogen vapor. Fifteen μm-thick sections were obtained on a Leica cryostat and stored

at -80°C. Sections were rehydrated in TBS (50 mM Tris, 150 mM NaCl, pH7.5), and

permeabilized with 0.5% Triton-X-100 (Sigma) and rinsed twice again in TBS before

blocking in 3% BSA, 1% Goat serum, 0.1% Tween-20 (Sigma). Immunolabeling was then

performed using M.O.M® Immunodetection Kit, Basic (Vector Laboratories, BMK-2202).

Purified anti-TAF7 rabbit polyclonal antibody (dilution 1/300) was revealed using an Alexa

Fluor 488 goat anti-rabbit IgG (Invitrogen #A-11108, dilution 1/1000). Sections were

counterstained with DAPI (4′,6-diamidino-2-phenylindole dihydrochloride; Molecular

Probes). Pictures were taken using a TCS SP5 Inverted confocal (Leica) with a 40x Plan APO

objective (CX PL APO 40x/1.25-0.75 OIL CS) and analyzed using Fiji 2.0.

Supervovulation

Five units of pregnant mare serum (PMS) was injected intraperitoneally in 4-week-old female

mice between 2-4 pm. After 44-46 hours, GV oocytes were collected from the ovaries by

puncturing with needles.

Oocytes collection

After dissection, ovaries are freed from adhering tissues in 1x PBS. Series of 6 ovaries were

digested in 500 μL of 2 mg/mL Collagenase (SIGMA), 0.025% Trypsin (SIGMA) and 0.5

mg/mL type IV-S hyaluronidase (SIGMA), on a ThermoMixer (Eppendorf) with gentle

agitation for 20 minutes. The digestion was then stopped by the addition of 1 mL of 37°C pre-

warmed αMEM - 5% FBS. The oocytes were then size-selected under a binocular.

RNA preparation

Pool of 100-200 oocytes collected were washed through several M2 drops, and total RNA

was isolated using NucleoSpin RNAXS kit (Macherey-Nagel) according to the user manual.

RNA quality and quantity were evaluated using a Bioanalyzer. Between 5-10 ng of RNA was

obtained from each pool of oocytes.

RNA-seq analyses

PolyA+ RNA seq libraries were prepared using the SMART-Seq v4 UltraLow Input RNA kit

(Clonetch) followed by the Nextera XT DNA library Prep kit (Illumina) according to the

manufacturer recommendations from 3 biological replicates for each condition (P7 wild-type

(WT), P7 *Tbpl2*[-/-] mutant, P14 WT and P14 *Tbpl2*[-/-] mutant oocytes) and sequenced 50 pb

single end using an Illumina HiSeq 4000 (GenomEast platform, IGBMC).

Reads were preprocessed in order to remove adapter, polyA and low-quality sequences (Phred

quality score below 20). After this preprocessing, reads shorter than 40 bases were discarded

for further analysis. These preprocessing steps were performed using cutadapt version 1.10[61].

Reads were mapped to spike sequences using bowtie version 2.2.8[62], and reads mapping to

spike sequences were removed for further analysis. Reads were then mapped onto the mm10

assembly of *Mus musculus* genome using STAR version 2.7.0f[63]. Gene expression

quantification was performed from uniquely aligned reads using htseq-count version 0.9.1[64],

with annotations from Ensembl version 96 and "union" mode. Read counts were normalized

across samples with the median-of-ratios method, to make these counts comparable between

samples and differential gene analysis were performed using the DESeq2 version 1.22.2[65]. All

the figures were generated using R software version 3.5.3.

RT-qPCR

Complementary DNA was prepared using random hexamer oligonucleotides and SuperScript

IV Reverse Transcriptase (Invitrogen) and amplified using LightCycler® 480 SYBR Green I

Master (Roche) on a LightCycler® 480 II (Roche). Primers used for qPCR analysis are listed

in Supplementary Table 2.

Repeat element analyses

Data were processed as already described[66] using Bowtie1 (version 1.2.2) [67] instead of Maq.

The repeatMasker annotation was used to identified the different types of repeat elements

(Smit, AFA, Hubley, R & Green, P. RepeatMasker Open-4.0. 2013-2015

http://www.repeatmasker.org). Differential expression analyses were performed using

DESeq2 (version 1.22.2) [65]. All the figures were generated using R custom scripts (version

3.5.3).

SLIC-CAGE analyses

Twenty-eight and 13 ng of total RNA isolated from P14 oocytes (biological replicate 1 and

replicate 2, approximately 500-1000 oocytes pooled for each replicate) and 15 ng of total

RNA isolated from P14 *Tbpl2*[-/-] mutant oocytes (approximately 550 pooled oocytes) were

used for SLIC-CAGE TSS mapping[43]. Briefly, 5 µg of the carrier RNA mix were added to

each sample prior to reverse transcription, followed by the cap-trapping steps designed to

isolate capped RNA polymerase II transcripts. The carrier was degraded from the final library

prior to sequencing using homing endonucleases. The target library derived from the oocyte

RNA polymerase II transcripts was PCR-amplified (15 cycles for P14 WT, 16 cycles for P14

Tbpl2[-/-] mutant) and purified using AMPure beads (Beckman Coulter) to remove short PCR

artifacts (< 200bp, size selection using 0.8 x AMPure beads to sample ratio). The libraries

were sequenced using HiSeq2500 Illumina platform in single-end, 50 bp mode (Genomics

Facility, MRC, LMS).

Sequenced SLIC-CAGE reads were mapped to the reference *M. musculus* genome (mm10

assembly) using the Bowtie2[62] with parameters that allow zero mismatches per seed sequence

(22 nucleotides). Uniquely mapped reads were kept for downstream analyses using CAGEr

Bioconductor package (version 1.20.0) [68] and custom R/Bioconductor scripts. Bam files were

imported into R using the CAGEr package, where the mismatching additional G, if added

through the template-free activity of the reverse transcriptase, was removed. Same samples

sequenced on different lanes and biological replicates were merged prior to final analyses.

Promoter analyses

In order to consider only the CAGE TSS dependent only on TBPL2, we removed all the P14

WT CAGE tags at position where CAGE tags were also present in the P14 *Tbpl2*[-/-] mutant

CAGE tags dataset: for the rest of the analysis, this dataset was called "TBPL2-dependent"

and we compared it to the P14 *Tbpl2*[-/-] mutant CAGE data (hereafter called "TBPL2-

independent"). Briefly, a CAGE set object was created from the TBPL2-dependent and

TBPL2-independent CTSS files using CAGEr Bioconductor package (version 1.20.0) [68], data

were normalized using normalizeTagCount (fitInRange = c(5,1000), alpha = 1.53, T = 1e6)

and the powerLaw option. Cluster of CTSS were collected using clusterCTSS (threshold = 1,

thresholdIsTpm = TRUE, nrPassThreshold = 1, method = "distclu", maxDist = 20,

removeSingletons = TRUE, keepSingletonsAbove = 5). Width of the TSS regions was

calculated using cumulativeCTSSdistribution and quantilePositions (clusters = "tagClusters",

qLow = 0.1, qUp = 0.9): interquantile width corresponds to the 10^{th}-90^{th} percentile of the total

tag cluster signal. In order to compare the different samples, consensus promoters were

computed using aggregateTagCluster (tpmThreshold = 3, qLow = 0.1, qUp = 0.9, maxDist =

100). Self-organizing map (SOM) expression profiling was performed using

getExpressionProfiles using a tpmThrshold of 3, the method "som", xDim = 3 and yDim = 2.

Shifting TSS were obtained after calculation of the cumulative distribution along the

consensus clusters using cumulativeCTSSdistribution and calculation of the shift score using

scoreShift with the Kolomogorov-Smirnov test. Shifting promoters were extracted using

getShiftingPromoters (tpmThreshold = 3, scoreThreshold = -Inf, fdrThreshold = 0.01).

TSSs corresponding to the MaLR-ERVS were identified after annotation using HOMER

(version 4.10) [69].

Sequences analyses were performed using Bioconductor R seqPattern (version 1.14) and R

custom scripts. WW dinucleotides enrichment was computed using plotPatternDensityMap on

-250/+250 regions centered on the dominant TSSs. TATA box position weight matrix (PWM)

matches analyses was performed using the MotifScanScores function applied on the -35/-20

sequences centered on the dominant TSSs, using the TBP PWM provided in the SeqPattern

package (derived from the JASPAR data base). Distribution of the best match for each

sequence was then plotted. Sequence Logo were created using Bioconductor R package

SeqLogo.

Data availability

The datasets generated during the current study are available in different repositories: proteomic data; ProteomeXchange PRIDE database (PXD0316347 [http://www.ebi.ac.uk/pride/archive/projects/PXD016347]), RNA-seq data; Gene Expression Omnibus database (GSE140090 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc= GSE140090]") and SLIC-CAGE data; ArrayExpress (E-MTAB-8866 [https://www.ebi.ac.uk/arrayexpress/experiments/E-MTAB-8866/]).

Code availability

RNA-seq data was analysed using Bioconductor package DESeq2, SLIC-CAGE data was analysed using Bioconductor package CAGEr. All custom code is available upon request.

References

1. Roeder, R. G. 50+ years of eukaryotic transcription: an expanding universe of factors
and mechanisms. *Nat Struct Mol Biol* **26,** 783–791 (2019).
2. Haberle, V. & Stark, A. Eukaryotic core promoters and the functional basis of
transcription initiation. *Nat Rev Mol Cell Biol* **19,** 621–637 (2018).
3. Tora, L. A unified nomenclature for TATA box binding protein (TBP)-associated
factors (TAFs) involved in RNA polymerase II transcription. *Genes Dev* **16,** 673–675
(2002).
4. Levine, M., Cattoglio, C. & Tjian, R. Looping back to leap forward: transcription
enters a new era. *Cell* **157,** 13–25 (2014).
5. Ohbayashi, T., Makino, Y. & Tamura, T. A. Identification of a mouse TBP-like protein
(TLP) distantly related to the drosophila TBP-related factor. *Nucleic Acids Res* **27,**
750–755 (1999).
6. Teichmann, M. *et al.* Human TATA-binding protein-related factor-2 (hTRF2) stably
associates with hTFIIA in HeLa cells. *Proceedings of the National Academy of*
Sciences **96,** 13720–13725 (1999).
7. Dantonel, J. C., Quintin, S., Lakatos, L., Labouesse, M. & Tora, L. TBP-like factor is
required for embryonic RNA polymerase II transcription in C. elegans. *Mol Cell* **6,**
715–722 (2000).
8. Kaltenbach, L., Horner, M. A., Rothman, J. H. & Mango, S. E. The TBP-like factor
CeTLF is required to activate RNA polymerase II transcription during C. elegans
embryogenesis. *Mol Cell* **6,** 705–713 (2000).
9. Veenstra, G. J. C. Distinct Roles for TBP and TBP-Like Factor in Early Embryonic
Gene Transcription in Xenopus. *Science* **290,** 2312–2315 (2000).
10. Müller, F., Lakatos, L., Dantonel, J., Strähle, U. & Tora, L. TBP is not universally
required for zygotic RNA polymerase II transcription in zebrafish. *Curr Biol* **11,** 282–
287 (2001).
11. Persengiev, S. P. *et al.* TRF3, a TATA-box-binding protein-related factor, is
vertebrate-specific and widely expressed. *Proceedings of the National Academy of*
Sciences **100,** 14887–14891 (2003).
12. Bártfai, R. *et al.* TBP2, a vertebrate-specific member of the TBP family, is required in
embryonic development of zebrafish. *Curr Biol* **14,** 593–598 (2004).
13. Jallow, Z., Jacobi, U. G., Weeks, D. L., Dawid, I. B. & Veenstra, G. J. C. Specialized
and redundant roles of TBP and a vertebrate-specific TBP paralog in embryonic gene
regulation in Xenopus. *Proceedings of the National Academy of Sciences* **101,** 13525–
13530 (2004).
14. Zhang, D., Penttila, T. L., Morris, P. L., Teichmann, M. & Roeder, R. G.
Spermiogenesis deficiency in mice lacking the Trf2 gene. *Science* **292,** 1153–1155
(2001).
15. Martianov, I. *et al.* Late arrest of spermiogenesis and germ cell apoptosis in mice
lacking the TBP-like TLF/TRF2 gene. *Mol Cell* **7,** 509–515 (2001).
16. Gazdag, E. *et al.* TBP2 is essential for germ cell development by regulating
transcription and chromatin condensation in the oocyte. *Genes Dev* **23,** 2210–2223
(2009).
17. Hernandez, N. TBP, a universal eukaryotic transcription factor? *Genes Dev* **7,** 1291–
1308 (1993).
18. Rabenstein, M. D., Zhou, S., Lis, J. T. & Tjian, R. TATA box-binding protein (TBP)-
related factor 2 (TRF2), a third member of the TBP family. *Proceedings of the*
National Academy of Sciences **96,** 4791–4796 (1999).

19. Hansen, S. K., Takada, S., Jacobson, R. H., Lis, J. T. & Tjian, R. Transcription
properties of a cell type-specific TATA-binding protein, TRF. *Cell* **91,** 71–83 (1997).
20. Ohbayashi, T. *et al.* Vertebrate TBP-like protein (TLP/TRF2/TLF) stimulates TATA-
less terminal deoxynucleotidyl transferase promoters in a transient reporter assay, and
TFIIA-binding capacity of TLP is required for this function. *Nucleic Acids Res* **31,**
2127–2133 (2003).
21. Carninci, P. *et al.* Genome-wide analysis of mammalian promoter architecture and
evolution. *Nat Genet* **38,** 626–635 (2006).
22. Lenhard, B., Sandelin, A. & Carninci, P. Metazoan promoters: emerging characteristics
and insights into transcriptional regulation. *Nat Rev Genet* **13,** 233–245 (2012).
23. Haberle, V. *et al.* Two independent transcription initiation codes overlap on vertebrate
core promoters. *Nature* **507,** 381–385 (2014).
24. Jagarlamudi, K. & Rajkovic, A. Oogenesis: Transcriptional regulators and mouse
models. *Mol Cell Endocrinol* **356,** 31–39 (2012).
25. Pedersen, T. & Peters, H. Proposal for a classification of oocytes and follicles in the
mouse ovary. *J. Reprod. Fertil.* **17,** 555–557 (1968).
26. Thompson, P. J., Macfarlan, T. S. & Lorincz, M. C. Long Terminal Repeats: From
Parasitic Elements to Building Blocks of the Transcriptional Regulatory Repertoire.
Mol Cell **62,** 766–776 (2016).
27. Peaston, A. E. *et al.* Retrotransposons regulate host genes in mouse oocytes and
preimplantation embryos. *Dev Cell* **7,** 597–606 (2004).
28. Gazdag, E., Rajkovic, A., Torres-Padilla, M.-E. & Tora, L. Analysis of TATA-binding
protein 2 (TBP2) and TBP expression suggests different roles for the two proteins in
regulation of gene expression during oogenesis and early mouse development.
Reproduction **134,** 51–62 (2007).
29. Malecova, B. *et al.* TBP/TFIID-dependent activation of MyoD target genes in skeletal
muscle cells. *Elife* **5,** 23 (2016).
30. Deato, M. D. E. *et al.* MyoD targets TAF3/TRF3 to activate myogenin transcription.
Mol Cell **32,** 96–105 (2008).
31. Zybailov, B. *et al.* Statistical analysis of membrane proteome expression changes in
Saccharomyces cerevisiae. *J. Proteome Res.* **5,** 2339–2347 (2006).
32. Helmlinger, D. & Tora, L. Sharing the SAGA. *Trends Biochem. Sci.* **42,** 850–861
(2017).
33. Lewandoski, M., Wassarman, K. M. & Martin, G. R. Zp3–cre, a transgenic mouse line
for the activation or inactivation of loxP-flanked target genes specifically in the female
germ line. *Current Biology* **7,** 148–151 (1997).
34. Vannini, A. & Cramer, P. Conservation between the RNA polymerase I, II, and III
transcription initiation machineries. *Mol Cell* **45,** 439–446 (2012).
35. Dong, J. *et al.* Growth differentiation factor-9 is required during early ovarian
folliculogenesis. *Nature* **383,** 531–535 (1996).
36. Galloway, S. M. *et al.* Mutations in an oocyte-derived growth factor gene (BMP15)
cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nat Genet*
25, 279–283 (2000).
37. Bettegowda, A. Mechanisms of maternal mRNA regulation: implications for
mammalian early embryonic development. *Frontiers in Bioscience* **12,** 3713 (2007).
38. Timmers, H. T. M. & Tora, L. Transcript Buffering: A Balancing Act between mRNA
Synthesis and mRNA Degradation. *Mol Cell* **72,** 10–17 (2018).
39. Yu, C. *et al.* BTG4 is a meiotic cell cycle-coupled maternal-zygotic-transition licensing
factor in oocytes. *Nat Struct Mol Biol* **23,** 387–394 (2016).

40.   Wolf, J. & Passmore, L. A. mRNA deadenylation by Pan2-Pan3. *Biochem. Soc. Trans.*
746   **42,** 184–187 (2014).
747   41.   Ma, J., Flemr, M., Strnad, H., Svoboda, P. & Schultz, R. M. Maternally recruited
748   DCP1A and DCP2 contribute to messenger RNA degradation during oocyte maturation
749   and genome activation in mouse. *Biol Reprod* **88,** 11 (2013).
750   42.   Veselovska, L. *et al.* Deep sequencing and de novo assembly of the mouse oocyte
751   transcriptome define the contribution of transcription to the DNA methylation
752   landscape. *Genome Biol.* **16,** 209 (2015).
753   43.   Cvetesic, N. *et al.* SLIC-CAGE: high-resolution transcription start site mapping using
754   nanogram-levels of total RNA. *Genome Res.* **28,** 1943–1956 (2018).
755   44.   Cvetesic, N. *et al.* Global regulatory transitions at core promoters demarcate the
756   mammalian germline cycle. *bioRxiv* **11,** 2020.10.30.361865 (2020).
757   45.   Törönen, P., Kolehmainen, M., Wong, G. & Castrén, E. Analysis of gene expression
758   data using self-organizing maps. *FEBS Letters* **451,** 142–146 (1999).
759   46.   Zhang, D., Penttila, T. L., Morris, P. L. & Roeder, R. G. Cell- and stage-specific high-
760   level expression of TBP-related factor 2 (TRF2) during mouse spermatogenesis. *Mech*
761   *Dev* **106,** 203–205 (2001).
762   47.   Martianov, I. *et al.* Distinct functions of TBP and TLF/TRF2 during spermatogenesis:
763   requirement of TLF for heterochromatic chromocenter formation in haploid round
764   spermatids. *Development* **129,** 945–955 (2002).
765   48.   Martianov, I., Velt, A., Davidson, G., Choukrallah, M.-A. & Davidson, I. TRF2 is
766   recruited to the pre-initiation complex as a testis-specific subunit of TFIIA/ALF to
767   promote haploid cell gene expression. *Sci Rep* **6,** 32069 (2016).
768   49.   Upadhyaya, A. B., Lee, S. H. & DeJong, J. Identification of a general transcription
769   factor TFIIAalpha/beta homolog selectively expressed in testis. *J Biol Chem* **274,**
770   18040–18048 (1999).
771   50.   Xiao, L., Kim, M. & DeJong, J. Developmental and cell type-specific regulation of
772   core promoter transcription factors in germ cells of frogs and mice. *Gene Expression*
773   *Patterns* **6,** 409–419 (2006).
774   51.   Müller, F., Zaucker, A. & Tora, L. Developmental regulation of transcription initiation:
775   more than just changing the actors. *Curr Opin Genet Dev* **20,** 533–540 (2010).
776   52.   Falender, A. E., Shimada, M., Lo, Y. K. & Richards, J. S. TAF4b, a TBP associated
777   factor, is required for oocyte development and function. *Developmental Biology* **288,**
778   405–419 (2005).
779   53.   Akhtar, W. & Veenstra, G. J. C. TBP2 is a substitute for TBP in Xenopus oocyte
780   transcription. *BMC Biol.* **7,** 45 (2009).
781   54.   Jacobi, U. G. *et al.* TBP paralogs accommodate metazoan- and vertebrate-specific
782   developmental gene regulation. *EMBO J* **26,** 3900–3909 (2007).
783   55.   Crichton, J. H., Dunican, D. S., Maclennan, M., Meehan, R. R. & Adams, I. R.
784   Defending the genome from the enemy within: mechanisms of retrotransposon
785   suppression in the mouse germline. *Cell Mol Life Sci* **71,** 1581–1605 (2014).
786   56.   Faulkner, G. J. *et al.* The regulated retrotransposon transcriptome of mammalian cells.
787   *Nat Genet* **41,** 563–571 (2009).
788   57.   McCarthy, E. M. & McDonald, J. F. Long terminal repeat retrotransposons of Mus
789   musculus. *Genome Biol.* **5,** R14 (2004).
790   58.   Franke, V. *et al.* Long terminal repeats power evolution of genes and gene expression
791   programs in mammalian oocytes and zygotes. *Genome Res.* **27,** 1384–1394 (2017).
792   59.   Vastenhouw, N. L., Cao, W. X. & Lipshitz, H. D. The maternal-to-zygotic transition
793   revisited. *Development* **146,** dev161471 (2019).

60. Gegonne, A. *et al.* The general transcription factor TAF7 is essential for embryonic
development but not essential for the survival or differentiation of mature T cells. *Mol*
Cell Biol **32,** 1984–1997 (2012).

61. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17,** 10–12 (2011).

62. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat.*
Methods **9,** 357–359 (2012).

63. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29,** 15–21
(2012).

64. Anders, S., Pyl, P. T. & Huber, W. HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31,** 166–169 (2015).

65. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15,** 31 (2014).

66. Fadloun, A. *et al.* Chromatin signatures and retrotransposon profiling in mouse
embryos reveal regulation of LINE-1 by RNA. *Nat Struct Mol Biol* **20,** 332–338
(2013).

67. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.* **10,** R25
(2009).

68. Haberle, V., Forrest, A. R. R., Hayashizaki, Y., Carninci, P. & Lenhard, B. CAGEr:
precise TSS data retrieval and high-resolution promoterome mining for integrative
analyses. *Nucleic Acids Res* **43,** e51–e51 (2015).

69. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38,**
576–589 (2010).

Acknowledgements

We thank D. Singer and A. Gegonne for the gift of the *Taf7*flox mouse line and H. Stunnenberg for TFIIA antibodies. We would also like to thank D. Devys for critically reading the manuscript, all members of the Tora lab for thoughtful discussions and suggestions throughout the course of the work. We are grateful to I. Kukhtevich, M. Borsos, M.E. Torres Padilla, T. Gupta, L. Casini, G. Barzaghi and A. Krebs for help in preliminary experiments. We thank C. Hérouard and M. Jung from the GenomEAST platform for library preparation and preliminary analyses, P. Eberling for peptide synthesis, F. Ruffenach for proteomic analyses, G. Duval for polyclonal antibody generation, the histology platform, the IGBMC cell culture facility and S. Falcone, M. Poirot and F. Memedov of the IGBMC animal facility for animal care taking. This work was supported by funds from CNRS, INSERM, and Strasbourg University. This study was also supported by the European Research Council (ERC) Advanced grant (ERC-2013-340551, Birtoaction) and grant ANR-10-LABX-0030-INRT (to LT) and a French State fund managed by the Agence Nationale de la Recherche under the frame program Investissements d'Avenir ANR-10-IDEX-0002-02. IB and FM acknowledge support by Wellcome Trust Senior Investigator awards (106115/Z/14/Z and 106955/Z/15/Z, respectively).

Authors' contribution

CY, SDV and LT designed the study; SDV and LT supervised the project; CY performed all molecular lab and mouse experiments, VH performed immunolocalizations and helped for the histological analysis of ovaries, EG generated the anti-TBPL2 polyclonal antibodies, LN carried out the proteomic analyses, KG and IB carried out preliminary analyses, PH organized the SLIC-CAGE, NC carried out SLIC-CAGE analyses, NC and BL analysed the SLIC-CAGE data, and SDV analysed the proteomic, RNA-seq and SLIC-CAGE data. FM oriented the promoter analyses.

CY, FM, SDV and LT wrote the manuscript with contributions to manuscript text and figure

legends from all authors. All authors gave final approval for publication.

Competing interests

The authors declare that they have no competing interests.

Additional Information

Supplementary information is available for this paper.

Correspondence and request for materials should be addressed to S. D. Vincent

(vincent@igbmc.fr) or L. Tora (laszlo@igbmc.fr).

 Figure legends

 Fig. 1 TBPL2 does not assemble in a TFIID-like complex in growing oocytes. **a** Anti-TBPL2

 immunoprecipitation followed by mass spectrometry (IP-MS) analysis from three biological

 replicates of mouse ovarian whole cell extracts (WCE). The color code for the different

 proteins or complexes is indicated on the right. NSAF; normalized spectral abundance factor.

 b Anti-TBP IP-MS from ovarian WCE (three technical triplicates). The color code is the same

 as in (**a**). **c-f** Sequential IP-MS experiment from ovarian WCE (three technical triplicates).

 Strategy of the sequential immunoprecipitation (**c**), anti-TAF7 IP-MS (**d**), followed by an

 anti-TAF10 IP-MS (**e**) and then an anti-TBPL2 IP-MS (**f**). The color code is the same as in

 (**a**). **g, h** Representative views of haematoxylin and eosin stained ovaries section from control

 (**e**) and oocyte-specific *Taf7* mutant (Tg(*Zp3-Cre*/+);*Taf7*$^{flox/\Delta}$, **f**) ovaries (analyses of 3

 sections from 3 biological replicates). The presence of antral follicles is indicated by an

 asterisk. Scale bars: 500 µm. Grey dots indicate replicates and error bars, +/- standard error of

 the mean in (**a**, **b**, **d-f**).

 Fig. 2 TBPL2 assembles into a TBPL2/TFIIA and TFIID-like complexes in NIH3T3 cells

 over-expressing TBPL2. **a** Anti-TBPL2 immunoprecipitation followed by mass spectrometry

 (IP-MS) analysis (three technical replicates) of NIH3T3 over-expressing TBPL2 (NIH3T3-

 II10) whole cell extracts (WCE). The color code for the different proteins or complexes is

 indicated on the right. **b** Anti-TBP IP-MS analysis (three technical replicates) of NIH3T3-II10

 WCE. Color legend for the different proteins is the same as in (**a**). **c** Western blot of a

 Superose 6 gel filtration analysis of NIH3T3-II10 WCE probed with anti-TAF6 (top panel),

 anti-TBPL2 and anti-TFIIA-α (middle panels), and anti-TBP (bottom panel) antibodies.

 Fraction numbers are shown above each lane and the elution of known molecular mass

markers is indicated above the panels. The pooled fractions used for mass spectrometry

analysis are indicated in red. **d** Anti-TBPL2 IP-MS analysis (three technical replicates) of the

gel filtration fraction indicated in (**c**). The color code for the different proteins or complexes is

indicated on the right. NSAF; normalized spectral abundance factor. **e** Schematic

representation of the fundamental differences existing between TBPL2- and TBP-containing

complexes in growing oocytes and NIH3T3-II10 cells. Grey dots indicate replicates and error

bars, +/- standard error of the mean in (**a**, **b**, **d**).

Fig. 3 Expression of genes related to the mRNA deadenylation/decapping/decay pathways in

growing *Tbpl2*$^{-/-}$ mutant oocytes. **a, b** Normalized Integrative Genomic Viewer (IGV)

snapshots of *Bmp15* (**a**) and *Gdf9* (**b**) loci. Exons and introns are indicated. **c** Expression

comparison between wild-type (WT) and *Tbpl2*$^{-/-}$ mutant post-natal day 14 (P14) oocytes

(biological triplicates). Expression was normalized to the median size of the transcripts in kb.

Grey dots correspond to non-significant genes and genes with high Cook's distance, light-

blue dots to significant genes for an adjusted p value ≤ 0.05 and dark-blue dots to significant

genes for an adjusted p value ≤ 0.05 and an absolute log2 fold change > 1, after two-sided

Wald test and Benjamini-Hochberg correction for multiple comparison. The number of up- or

down-regulated genes is indicated on the graph. **d** Down-regulated genes GO category

analyses for the molecular functions (MF). The top ten most enriched significant GO

categories for a FDR ≤ 0.05 are represented. **e** Heatmap of selected genes involved in mRNA

decay, decapping or deadenylation pathways. Expression levels in fold-change (compared to

the mean of WT) of three biological replicates of P14 WT and P14 *Tbpl2*$^{-/-}$ mutant oocytes are

indicated. The fold change color legend is indicated at the bottom. The first column on the left

corresponds to the log10 of expression, scale on the left.

Fig. 4 Expression of the mammalian apparent LTR retrotransposons (MaLR) endogenous

retroviral elements (ERV) are down regulated in growing *Tbpl2*$^{-/-}$ mutant oocytes. **a-h**

Differential expression between wild type and *Tbpl2*$^{-/-}$ mutant post-natal day (P) 7 (**a-d**) and

P14 (**e-h**) oocytes of the different transposon classes; RNA transposon classes [LINEs (**a**, **e**),

SINEs (**b**, **f**) and ERVs (**c**, **g**)] and DNA transposons (DNAs (**d**, **h**)). The ERV sub-class III

mammalian apparent LTR retrotransposon (MaLR) family is the most severely affected in

Tbpl2$^{-/-}$ mutant oocytes at P7 and P14. Dot size and color are explained on the right side,

adjusted *p* value after two-sided Wald test and Benjamini-Hochberg correction for multiple

comparison.

Fig. 5 Core promoter regions of TBPL2 specific transcription units in post-natal day 14

oocytes are enriched in TATA-like elements and are sharp. **a, b** Genome-wide A/T-rich

dinucleotide (WW) motif analyses of -250/+250 sequences centred on the dominant

transcription start sites (TSS, position +1, dashed red line) of TBPL2-dependent (**a**, n =

32448) and TBPL2-indendent (**b**, n = 23886) TSS clusters. Sequences have been ordered by

increasing size of the interquantile width of each. The red arrowheads indicate the WW

enrichment at position -30 in the TBPL2-dependent TSS clusters (**a**) and the equivalent

position in the TBPL2-independent TSS clusters (**b**). The number of TSS clusters is indicated

in brackets. **c** Distribution of the best TATA box position weight matrix (PWM) matches

within a -35 to -20 region upstream of the dominant TSSs (+1) of TBPL2-dependent (blue)

compared to the TBPL2-independent (orange) TSS clusters. The dashed lines indicate the

median of the TATA box PWM matches for the TBPL2-dependent (blue) and the TBPL2-

independent (orange) TSS clusters (*p* value after a two-tailed Wilcoxon rank-sum test). **d, e**

Two selected self-organizing map (SOM) groups of the consensus TSS clusters: the down-

regulated promoters (blue, **d**) and the up-regulated promoters (red, **e**) groups. **f, g** Sequence

logos of the -35/+5 sequence of the TBPL2-dependent dominant TSSs from the down-

regulated promoters (**f**) and of the TBPL2-independent dominant TSSs from the up-regulated

promoters (**g**). **h** Distribution of the best TATA box PWM matches within a -35 to -20 region

upstream of the TBPL2-dependent (blue) and TBPL2-independent (orange) mammalian

apparent LTR retrotransposons (MaLR) endogenous retroviral elements (ERV) dominant

TSS. The dashed lines indicate the median of the TATA box PWM matches for the TBPL2-

dependent (blue) and the TBPL2-independent (orange) TSS clusters (*p* value after a two-

tailed Wilcoxon rank-sum test). **i, j** Sequence logo of the -35/+5 sequence of the MaLR ERVs

TBPL2-dependent (**i**) and TBPL2-independent (**j**) dominant TSSs.

Fig. 6 Shifting promoters. **a, b** Analysis of the TBPL2-dependent versus TBPL2-independent

shifting promoters within a -250/+250 region centred on the position of the TBPL2-

independent transcription start sites (TSS) clusters (position +1 in (**a, b**) and red dashed line

in (**b**)). Position of the TBPL2-dependent (blue) and of the TBPL2-independent (orange)

dominant TSSs for each shifting promoter sequence (**a**) and WW dinucleotide enrichment

heat map (**l**) from the same set of sequences ordered as in (**b**). **c** Distribution of the best

TATA box PWM matches within a -35 to -20 region upstream of the TBPL2-dependent

(blue) and TBPL2-independent (orange) dominant TSS of the shifting promoters. The dashed

lines indicate the median of the TATA box PWM matches for the TBPL2-dependent (blue)

and the TBPL2-independent (orange) shifting TSS clusters (*p* value after a two-tailed

Wilcoxon rank-sum test).

Fig. 7 Transcriptome overhaul controlled by TBPL2/TFIIA during oocyte growth. At the

beginning of oocyte growth, the transcriptome in primordial and early primary follicles (blue

cell) depends on TFIID/TBP (blue complex) transcription from broad promoters (blue line).

As TBP protein disappears, Pol II transcription initiation is mediated (red line) only by the

oocyte-specific TFIIA/TBPL2 complex (red complex) from sharp promoters. At the growing

oocyte stage (red cell) TFIIA/TBPL2 complex is responsible for initiating high levels of Pol II

transcription, such as mammalian apparent LTR retrotransposons (MaLR) endogenous

retroviral elements (ERV) expression and also expression of genes of which the gene products

survey/limit the stability and/or translatability of transcripts previously deposited by

TFIID/TBP-dependent Pol II transcription.

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7