EEG in Motion: Using an Oddball Task to Explore Motor Interference in Active

Skateboarding

Daniel Robles, Jonathan W. P. Kuziek, Nicole A. Wlasitz, Nathan T. Bartlett, Pete L. Hurd, Kyle

E. Mathewson

Department of Psychology
Faculty of Science
University of Alberta

Corresponding Author:

Kyle E. Mathewson, Department of Psychology
(780) 492-2662
P-217 Biological Sciences Building, University of Alberta
Edmonton, Alberta
Canada T6G 2R3
kmathews@ualberta.ca

Number of pages: 33. Total number of figures: 7. Total number of tables: 3

Total number of words in manuscript: 9550. Total number of words in abstract: 157

Keywords: Mobile EEG, ERP, outdoor cognition, portable research

Abstract

Recent advancements in portable computer devices have opened new avenues in the study of human cognition outside research laboratories. This flexibility in methodology has led to the publication of several Electroencephalography (EEG) studies recording brain responses in real world scenarios such as cycling and walking outside. In the present study, we tested the classic auditory oddball task while participants moved around a running track using an electric skateboard. This novel approach allows for the study of attention in motion while virtually removing body movement. Using the skateboard auditory oddball paradigm, we found reliable and expected standard-target differences in the P3 and MMN/N2b event-related potentials (ERPs). We also recorded baseline EEG activity and found that, compared to this baseline, alpha power is attenuated in frontal and parietal regions during skateboarding. We also tested for the effect of stance preference to understand how changes in task difficulty modulate the P3 and alpha magnitude but found no differences in either.

Introduction

For decades, the study of cognitive electrophysiology using Electroencephalography (EEG) has taken place inside highly controlled research facilities as EEG signals are easily contaminated by a myriad of environmental factors (Luck, 2014). EEG research has informed our understanding of human attention, yet, this knowledge generally comes from paradigms that isolate participants in faraday cages to avoid electromagnetic fields and other sources of noise that can compromise data quality (Puce & Hämäläinen, 2017). Over recent years, developments in portable computer devices have allowed such studies to move outside the lab and into the real world, resulting in a growth of mobile EEG studies within ecologically rich environments (Askamp & van Putten, 2014; Kontson et al., 2015; Cruz-Garza et al., 2017; Kuziek et al., 2018).

Mobile EEG research allows us to further understand attention and brain function by exploring classic paradigms in real world scenarios. For example, an oddball task in which participants respond to rare stimuli and ignore frequent distractors has classically been used to study attentional allocation. The ERP (event-related potential) known as the P3, a positive voltage increase following infrequent stimuli (Squires et al., 1975), has been widely reported in EEG studies for decades since its amplitude is inversely related to task difficulty. This robust ERP is ideal to explore brain responses in more complex real-world environments due to its replicability in the oddball task (Zamrini et al., 1991). Furthermore, the oddball paradigm allows for a dual-task approach where a primary task is presented concurrently with the oddball stimuli. These paradigms assume that increases in primary task difficulty compete with cognitive resources simultaneously devoted to the oddball task, leading to measurable reductions in P3 amplitude (Kok, 2001). There is a long history of using the P3 as a measure of resource allocation in realistic tasks such as studies of pilots (Sirevaag et al., 1993), as well as video game

training studies showing that increases in task resources from training are related to P3 amplitude increases to gaming events (Maclin et al., 2011, Mathewson et al., 2012).

The past decade has seen an increased interest in Mobile brain/body imaging (MoBI) studies linking brain activity to naturalistic behaviors in 3D spaces (Makeig et al., 2009; Gramann et al., 2014; Jungnickel et al., 2016). This integrative approach collects EEG, motor behavior, and environmental events to study action and participant/environment interactions (Ojeda, et al., 2014). MoBI studies involving naturalistic behaviors such as walking and running generally use high density recording while applying independent component analysis (ICA) to systematically separate the sources of movement artifacts from the EEG signal (Gwin et al., 2010). Previous studies from our laboratory have successfully recorded ERP components during active physical performance and in real-world scenarios. For example, using an auditory oddball paradigm, Scanlon et al., (2017) reliably recorded the P3 and MMN/N2b components, as well as a frontal alpha peak, during stationary cycling. Differences in ERP components in indoor vs outdoor tasks have also been shown in these mobile paradigms: In a follow-up study where participants completed the oddball task while cycling outdoors, Scanlon et al., (2019) showed that the P3 and MMN/N2b are present during outside cycling. Additionally, it was found that compared to indoor cycling, there is a reduction in the oddball P3, P2, and alpha peak when cycling outside. This P3 reduction is likely attributed to an increase in cognitive processing due to being outdoors, where participants are continuously processing complex, external auditory and visual stimuli that compete with resources devoted to the oddball task while focusing on bicycle riding and direction. Additionally, relative to indoors, cycling outside was found to maximize the N1 amplitude for standard and target tones. Such an increase in N1 amplitude could reflect an increase in auditory filtering required to complete the oddball task in the noisier outdoor

environment. These cycling studies suggest that certain ERP components show a different topography outdoors, likely due to the influence of external stimuli bombarding the brain while being outside. Other studies comparing indoor vs outdoor cycling have also found a reduction in the P3 while participants cycle outside vs indoors (Zink et al., 2016). Results like these highlight the importance of outdoor paradigms for several reasons. First, replicating classic findings under less controlled scenarios (e.g., a busy road) is important for the field of cognitive electrophysiology since, ultimately, we want to predict and understand brain functioning in everyday tasks outside laboratories. Second, it allows for exploratory and novel paradigms to be developed for research questions about complex naturalistic behaviors where we can learn more about the outdoor brain. Third, mobile EEG is a promising tool that can offer affordable and flexible medical diagnosis (Krigolston et al., 2017), and can offer flexibility in measuring cognitive performance (e.g., fatigue/cognitive load) in real life tasks (Darari et al., 2017). Mobile EEG paradigms have also been employed to investigate the relationship between cognitive load and motor task demands in attentional paradigms involving physical action. For example, Liebherr et al. (2018) demonstrated that increases in both motor task difficulty and cognitive difficulty lead to reductions in P3 amplitude in a cognitive task. The increase in motor difficulty was manipulated by having participants stand on one leg while completing the task. In accordance with these results, Reiser et al. (2019) employed an oddball paradigm outdoors to measure the influence of movement complexity in P3 amplitude. They showed that movement complexity was associated with increases in cognitive load followed by a reduction in P3 amplitude to the target tones. Likewise, Ladouce et al. (2019) used an oddball task to show that, relative to standing, walking is associated with a reduction in P3 amplitude to targets. This P3

reduction in walking was linked to an increase in processing demands related to increased visual stimulation.

In addition to ERPs, oscillatory rhythms have also been used to estimate resource allocation and cognitive engagement in experimental tasks. For example, power in the alpha range (8-12 Hz) has been measured in studies of active visuospatial biasing and suppression (Kelly et al., 2009), selective inhibition, and other anticipatory mechanisms (Foxe et al., 1998; Rihs et al., 2007). Previous studies involving motor behaviors, such as walking and cycling, have found differences in alpha power based on attentional demands in these movement tasks. For example, Storzer et al., (2016) found that walking is associated with a stronger alpha decrease than stationary cycling due an increase in sensory processing and motor planning in the walking condition. Wagner et al., (2014) have found decreases in alpha power in walking when motor planning and intention is required. Cycling was associated with decreases in alpha power when participants completed an oddball task riding a bicycle relative to a sitting position indoors (Scanlon et al., 2019). This decrease in power was attributed to increased stimulus processing from doing the task in an outdoor environment.

In the current experiment we used an active task of electric-skateboarding (operating a self-propelling skateboard with a wireless Bluetooth handheld controller) on an indoor 200-meter track as a primary task while participants simultaneously completed an auditory oddball task. We adopted a skateboard paradigm because it allows participants to be in active interaction with the environment while greatly minimizing body movements that lower signal-to-noise ratio in EEG recordings (Oliveira et al., 2016). We employed the portable EEG methodology previously used in our laboratory in cycling paradigms (e.g., Scanlon et al., 2019) where participants could freely move while wearing the EEG system in a backpack (Figure 1). From the results of these cycling

studies, we were motivated to devise this basic skateboard task to test for the allocation of

cognitive resources while participants interact with their environment and minimize body

movements. This paradigm allows for flexible navigation, which can be beneficial for future

studies interested in brain functioning outdoors. Using skateboarding as the primary task, our

focus was to use an auditory oddball paradigm to test whether we could record a reliable P3 and

MMN effect between standard and target tones. Furthermore, we wanted to investigate whether

an increase in primary task difficulty (an increase in skateboarding difficulty) would produce

measurable changes in P3 amplitude. E-skateboarding difficulty can be increased by

manipulating stance preference: since skateboarding requires individuals to move laterally, it is

common for learners to develop a stance preference while learning the task. Preference is

reflected in which foot goes forward on the board or which shoulder they look over while riding.

Right handed skateboarders generally ride with their left leg forward, however, some right

handed skaters generally prefer to ride with their right foot forward. The former riding style is

generally called "regular" whereas the latter is referred to as "goofy".   Due to the stability and

safety of the e-skateboard, participants can be asked to switch their stance preference in order to

increase riding difficulty without turning the task into a falling hazard. In this context, riding in

the non-preferred stance implies an increase in primary task difficulty, which has been

previously associated with a decrease in P3 amplitude in the secondary oddball task (Kramer &

Strayer, 1988; Kida et al., 2012). We tested the robustness of the oddball P3 effect while

participants rode a skateboard on a running track in four different conditions (preferred stance

clockwise direction, preferred stance counterclockwise direction, non-preferred stance clockwise

direction, and non-preferred stance counterclockwise direction). For the ERP analysis, since dual

task processes modulate P3 amplitude, we predicted a decrease in P3 amplitude in the non-preferred stance due to increases in riding difficulty while completing the task.

Using the skateboard paradigm, we tested whether an increase in task difficulty (by having participants e-skateboard in their non-preferred stance) would be reflected in alpha power decreases during completion of the oddball task in the non-preferred condition. We also recorded participant's resting state EEG to test whether we could measure a typical alpha peak (an increase of spectra power in the 8-12 range) right by the running track prior to completing the skateboard task. Decreases in resting state alpha peak are associated with increased cognitive arousal (Barry, Clarke & Johnstone, 2011). Since the experiment was taking place in a busy environment, we were motivated to assess resting alpha levels prior to the start of the task. Additionally, we computed the averaged alpha power between the left and right electrodes to conduct an exploratory analysis on alpha lateralization during skateboarding. Spatial attention paradigms (Sauseng et al., 2005; Ikkai et al., 2016; Kelly et al., 2006) demonstrated that alpha power increases over areas to be ignored. Such a shift in alpha power, with enhanced power in the hemisphere contralateral to the distractors, has been identified as a mechanism of sensory inhibition (Mathewson et al., 2011). In the current paradigm, participants ride an e-skateboard around a running track clockwise and counterclockwise. This allows us to spatially separate the source of incoming distractors based on the direction participants ride. In the clockwise condition (regardless of riding preference) the largest source of distractors (joggers, walkers, and track users) come from the right side of space while in the counterclockwise condition the source of distraction is always located on the left visual field of the participant. This is because participants were set to only ride in the outermost track during the task. We conducted exploratory analyses to test whether we could get an increase of alpha power in the parieto-occipital region

contralateral to the source of distractors in the clockwise and counterclockwise riding directions. Using this indoor track and mobile EEG paradigm, we expect to reliably measure ERPs using the oddball task while exploring the influence of riding difficulty and distractors in ERP and oscillatory rhythms.

Methods

Participants

A total of 29 individuals from the university community completed the study (mean age = 20.96, age range = 18-27, goofy-footed proportion: 41.37%, female proportion = 37.93%). Participants received either an honorarium of $20 or credits towards Research Participation in an undergraduate psychology class. All participants had normal or corrected-to-normal vision and did not have neurological antecedents. Each participant was comfortable with basic skateboarding in both preferred and non-preferred stances. This study was approved by the Internal Research Ethics Board at the University of Alberta (Pro00050069) and participants gave informed consent before completing the study. This study conforms with the World Medical Association Declaration of Helsinki.

Materials and Procedure

The experiment was completed in the Universiade Pavilion (aka Butterdome) on a 200-meter indoor track at the University of Alberta. Participants first completed two, three-minute, resting-state tasks (beginning with either eyes closed or eyes open) to measure a baseline level of resting spectral activity while they were seated in the bleachers of the track. Participants were instructed to breathe calmly and either keep their eyes closed, or to fixate on a nearby object. Following both baseline tasks, participants were given a basic tutorial on using the Boosted V3 Stealth Electric Board (Palo Alto, California, United States; https://boostedboards.com/ca/)

before starting the main skateboard experiment. The Boosted skateboard is a two-motor electric skateboard controlled by a joystick held in the right hand, with throttle and clutch controls on the thumb and trigger finger respectively. The joystick was held in the right hand while a response button was held in the left hand. Participants began the task by standing on the skateboard and pressing the response button, starting a 10-second countdown. Participants were expected to be in motion around the Universiade Pavilion track when the countdown finished. To ensure clean data, participants were instructed to remain as still as possible, blink only when necessary, and keep their gaze directed forward. A sound cue was used to instruct the participants when the task was completed. Participants rode the skateboard using a limited acceleration mode with a maximum speed of 17 kph. Participants were trained to ride at a speed in which they felt most comfortable around the track for safety purposes.

Based on the participant's preferred skateboarding stance and the direction the participant travelled around the Pavilion track, four conditions were generated: 'preferred clockwise', 'preferred counterclockwise', 'non-preferred clockwise', and 'non-preferred counterclockwise'. Condition order was randomly determined and counter-balanced. In each condition, participants completed an auditory oddball task where two tones were consistently presented through headphones (either a 1000 Hz or 1500 Hz tone). Participants were instructed to press the response button each time the 1500 Hz tone was played (target tone) and to withhold a response following the 1000 Hz tone (standard tone). A delay, randomly selected from a distribution between 1000 and 1500 ms, followed each tone. Response times were collected during this delay period. In all, 20% of trials contained a target tone and 80% contained a standard tone, with 250 trials in each condition. Each condition was approximately six minutes long. Between

conditions, the participant returned to the starting position, the computers were reset, and the

participant was given a small break.

The tones and response button task were programmed and administered using a

Raspberry Pi 2 model B computer (Raspberry Pi Foundation, Cambridge, UK) running version 9

of the Raspbian Stretch operating system and version 2.7.13 of Python. The Raspberry Pi 2 was

powered via a micro-USB cable connected to a Microsoft Surface Pro 3 laptop. Audio output

was via Sony earbuds connected to the 3.5mm audio connector on the Raspberry Pi 2.

Coincident in time with tone onset, 8-bit TTL pulses were sent to the EEG amplifier via a cable

connected to the GPIO pins of the Raspberry Pi 2. These trigger markers were used to mark the

EEG data for ERP averaging. EEG data was collected from participants using active, wet, low-

impedance electrodes (actiCAP active electrodes kept below 5 kΩ). Inter-electrode impedances

were measured at the start of the experiment. The following electrode locations were used,

arranged in the typical 10-20 electrode positions (F3, F4, T7, T8, C3, C4, P7, P8, P3, P4, O1, O2,

Fz, Cz, Pz). A ground electrode was used, positioned at AFz. Ag/AgCl disk electrodes with

SuperVisc electrolyte gel and mild abrasion with a blunted syringe tip were used to lower

impedances. EEG data was recorded online, referenced to an electrode attached to the left

mastoid. Offline, the data were re-referenced to the arithmetically derived average of the left and

right mastoid electrodes.

EEG data was recorded with a Brain Products V-Amp 16-channel amplifier (Brain

Products GmbH), connected to the same Microsoft Surface Pro 3 laptop powering the Raspberry

Pi, running BrainVision Recorder software (Brain Products GmbH, Gilching, Germany). In

addition to the 15 EEG sensors, two reference electrodes, and the ground electrode, vertical and

horizontal bipolar EOG was recorded from passive Ag/AgCl easycap disks using Bip2Aux

adapters connected to the auxiliary ports of the amplifier. EOG electrodes were affixed vertically above and below the left eye and affixed horizontally 1 cm lateral from the outer canthus of each eye. The participant's skin was cleaned using Nuprep (an exfoliating cleansing gel) (Weaver & Co, Aurora, Colorado USA) before the placement of the electrodes, electrolyte gel was used to lower the impedance of these electrodes to under 5 kΩ in the same manner as previously mentioned. Data were digitized at 1000 Hz with a resolution of 24 bits and hardware filtered online between 0.1 Hz and 30 Hz, with a time constant of 1.5155 s and a notch filter at 60 Hz. All aforementioned equipment was held within a 2-pocket backpack worn by the participant, as shown in Figure 1.

EEG Analysis

Analyses were computed in MATLAB R2018a (Mathworks, Natick, Massachusetts, USA) using EEGLAB (Delorme & Makeig, 2004) and custom scripts (https://github.com/kylemath/MathewsonMatlabTools). Statistical analyses were computed on JASP (JASP Team, Amsterdam, Netherlands). The EEG markers were used to construct 1200 ms epochs (200 ms pre-stimulus baseline + 1000 ms post-stimulus) time locked to the onset of standard and target tones, with the average voltage in the first 200 ms baseline period subtracted from the data for each electrode and trial. To remove artifacts due to amplifier blocking and other non-physiological factors, any trials with a voltage difference from baseline larger than +/- 1000 µV on any channel (including eyes) were removed from further analysis. At this time, a regression-based eye-movement correction procedure was used to estimate and remove the artifact-based variance in the EEG due to blinks as well as horizontal and vertical eye movements (Gratton et al., 1983). After identifying blinks with a template-based approach, this technique computes propagation factors as regression coefficients predicting the vertical and

horizontal eye channel data from the signals at each electrode. The eye channel data is then

subtracted from each channel, weighted by these propagation factors, removing any variance in

the EEG predicted by eye movements. Artifact rejection was again performed, using a voltage

threshold of +/- 500µV. These artifact rejection thresholds were chosen to be relatively lenient,

similar to other mobile EEG studies we have done (Scanlon et al., Scanlon et al., 2017, Scanlon

et al., 2019, Scanlon et al., 2020), in order to quantify how much noise (electrical, muscle, or

otherwise) was present in the ERP data and to ensure an adequate number of trials were available

for analysis. Baseline correction was performed again following the second artifact rejection.

Table 1 shows the mean trial count for target and standard tones used for each condition after

artifact rejection. The rejected number of trials for targets and standards respectively was similar

for all conditions($F(3,112) = 0.84$, $p = 0.48$).

ERP analysis

Appropriate time window cut-offs for the MMN/N2-B and P3 waveforms were

determined by creating grand-average ERP waveforms, averaged across all participants and

conditions, to create a single ERP waveform for electrodes Fz and Pz (Figure 2). These grand-

average waveforms were used to avoid biasing the selected time windows towards any one

condition. The negative deflection between 195-295 ms for electrode Fz was used for the

MMN/N2-B time window while the large positive deflection between 375-525 ms for electrode

Pz was used for the P3 time window. Trial-averaged ERPs were derived in each of the four

conditions ('preferred clockwise', 'preferred counterclockwise', 'non-preferred clockwise', and

'non-preferred counterclockwise') and waveforms were compared. A 2x2x2 ANOVA was run to

compare tone type (standard and target), stance preference (preferred and nonpreferred), and

clock orientation (clockwise and counterclockwise) during the MMN/N2-B and P3 time

windows, at electrodes Fz and Pz respectively. Two-way repeated measures ANOVAs were also

conducted on the difference waveforms for the MMN/N2b and P3 time windows at electrode Fz

and Pz respectively. Stance preference and clock orientation were the factors used for these

analyses. These waveforms were derived by subtracting the standard ERP waveforms from the

target ERP waveforms for both the preferred/non-preferred, and clockwise-counterclockwise

conditions, with α set to 0.05 for all analyses.

Spectral Analysis

Average frequency spectra across the baseline and oddball epochs was also computed

using the wavelet routine from the Better Oscillation Method (BOSC; Whitten et al., 2011) with

a 6-cycle wavelet transform across a frequency range of 0.1 Hz to 30 Hz, increasing in 0.5 Hz

steps. We chose this wavelet cycle parameter because we were interested in the overall sustained

alpha power instead of the differences in power between target and standard tones. Spectral data

was further analyzed using the "Fitting Oscillations & One Over F" (FOOOF) algorithm (Haller

et al., 2018) to represent the data as two distinct components; the aperiodic background $1/f$ and

periodic oscillations which may contain greater spectral power than the aperiodic component.

This analysis was performed using version 0.1.1 of the FOOOF Matlab wrapper with the

following settings: peak width limits = [0.5,12]; max number of peaks = Inf; minimum peak

amplitude = 0.0; peak threshold = 2.0; background mode = fixed; verbose = true; frequency

range = 0.5-30 Hz. The background $1/f$ spectra was then subtracted from the periodic component

to better compare changes in spectral power between 0.5 Hz and 30 Hz across our conditions. All

spectral analyses were done using this calculated FOOOF spectral data (Figure 6 C-D).

For the baseline task, spectra were computed for each participant by first dividing the

data into 3000 ms segments. Spectra were calculated for each chunk, which were then averaged

across the 3000 ms duration for each chunk. Then each averaged-chunk was combined together within each participant, and finally averaged across participants to generate grand-averaged spectra for both eyes open and eyes closed conditions. For the oddball task, we generated 3000 ms epochs around the onset of each standard trial (1000 ms pre and 2000 ms post standard onset) for each participant. Spectra were calculated for each epoch, then averaged across time and number of standard trials to generate spectra for each participant. These spectra were then averaged across participants to create grand-average spectra for each condition in the oddball task. Spectra were calculated using electrodes Fz and Pz. Resting state EEG was not recorded from two participants and therefore their data was excluded from this main spectral analysis. Furthermore, in order to explore the role of left-right visual field distractors in hemispheric alpha, power was computed by combining the following parieto-occipital electrodes: left hemisphere (P7, P3, O1), right hemisphere (P8, P4, O2).

Results

Behavioral Results

Figure 3A shows the mean accuracy in response to standard tones across all four task conditions and by global stance preference (collapsed over direction of travel). Results from a two-way repeated measures ANOVA on behavioral accuracy show no significant main effect for either preference, ($F(1, 28) = 0.74$, $p = 0.40$, $\eta^2_p = 0.03$), or clock orientation (F(1, 28) = 0.85, p = 0.36, $\eta^2_p = 0.03$) and no significant interaction ($F(1, 28) = 0.01$, $p = 0.91$, $\eta^2_p = 4.61e-4$). Figure 3B shows mean reaction times in milliseconds for all four task conditions and by global stance preference. A two-way repeated measures ANOVA on reaction times show no significant main effect for either preference, ($F(1, 28) = 2.56e-4$, $p = .99$, $\eta^2_p = 9.15e-6$), or clock orientation

($F(1, 28) = .03$, $p = 0.87$, $\eta^2_p = 9.29e-4$), and no significant interaction ($F(1, 28) = 0.27$, $p = 0.61$,

$\eta^2_p = 0.01$).

ERP Morphology and Topography

Figure 4A shows the grand average ERPs for target and standard tones at electrode Fz for

all study conditions. Target tones are depicted in colored lines while standard tones are depicted

in black lines. Standard errors at each time point are depicted by the shaded regions. Compared

to standard tones, there is an increase in negative voltage for the MMN/N2 time window between

195 and 295 ms following target tones. Figure 4C shows the grand average ERP at electrode Pz

for all conditions, showing a clear positive increase in voltage in the P3 window (350 and 550

ms) following target tone onset. Additionally, topographies were computed for the MMN/N2b

and P3 time windows across conditions to show the overall scalp activation distribution for both

time windows. Based on the time window chosen for these topographies, one can observe a more

anterior distribution for the MMN/N2b time window and a more posterior distribution for the P3

time window. We conducted a 2x2x2 repeated measures ANOVA in order to understand the

relationship between target/standard tones in all four conditions and whether the MMN/N2-B

and P3 are influenced by riding condition. For the MMN/N2b time window at electrode Fz, there

was a significant main effect for tone type ($F(1, 28) = 87.72$, $p < .001$, $\eta^2_p = 0.76$). There was no

significant effect of preference type ($F(1, 28) = 0.63$, $p = 0.43$, $\eta^2_p = 0.02$) or orientation ($F(1,$

$28) = 2.78$, $p = 0.11$, $\eta^2_p = 0.09$), and there were no significant interactions: tone type*preference

($F(1, 28) = 2.04$, $p = 0.17$, $\eta^2_p = 0.07$), tone type*orientation ($F(1, 28) = 0.92$, $p = 0.35$, $\eta^2_p =$

0.03), preference*orientation ($F(1, 28) = 1.87$, $p = 0.18$, $\eta^2_p = 0.06$), tone

type*preference*orientation ($F(1, 28) = 0.01$, $p = 0.91$, $\eta^2_p = 4.77e-4$). Table 2 shows the

significant pairwise comparisons where it can be seen that the target-standard difference at the

MMN/N2b window was reliable in all four conditions. All p-values for the pairwise comparisons were adjusted for multiple comparisons using Bonferroni correction.

For the P3 window at electrode Pz, there was a significant main effect for tone type ($F(1, 28) = 50.86$, $p < .001$, $\eta^2_p = 0.65$), no significant effect of preference type ($F(1, 28) = 0.08$, $p = 0.78$, $\eta^2_p = 3\text{e-}3$) or orientation ($F(1, 28) = 0.11$, $p = 0.74$, $\eta^2_p = 4\text{e-}3$), and there were no significant interactions: tone type*preference ($F(1, 28) = 0.41$, $p = 0.53$, $\eta^2_p = 0.01$), tone type*orientation ($F(1, 28) = 1.42$, $p = 0.24$, $\eta^2_p = 0.05$), preference*orientation ($F(1, 28) = 0.17$, $p = 0.68$, $\eta^2_p = 6\text{e-}3$), tone type*preference*orientation ($F(1, 28) = 0.83$, $p = 0.37$, $\eta^2_p = 0.03$). Table 2 shows the pairwise comparisons at electrode Pz where, across all conditions, the oddball P3 difference between targets and standards was highly reliable across all combinations of orientation and preference.

We also analyzed the difference waveforms of our ERPs across the preference and orientation conditions. These waveforms were calculated by subtracting standard trial ERPs from target trial ERPs, allowing us to better understand the differences in evoked activity between our two tone types. Figure 5A shows the difference-wave ERP for the preferred and non-preferred stance in the MMN/N2b time window. Figure 5B shows the difference-wave ERP for the clockwise and counterclockwise conditions in the MMN/N2b time window. A 2x2 repeated measures ANOVA was calculated for electrodes Fz and Pz and their respective time windows, as described previously. Our results at electrode Fz show no significant effect of preference ($F(1, 28) = 2.04$, $p = 0.17$, $\eta^2_p = 0.07$) or orientation ($F(1, 28) = 0.92$, $p = 0.35$, $\eta^2_p = 0.03$), and no significant interaction ($F(1, 28) = 0.01$, $p = 0.91$, $\eta^2_p = 4.78\text{e-}4$).

Figure 5C shows the difference-wave ERP by preference conditions at electrode Pz for the P3 time window. Figure 5D shows the difference-wave ERP for the clockwise and

counterclockwise conditions in the P3 time window. Similar results also occur for electrode Pz, with no significant effect of preference ($F(1, 28) = 0.41$, p = 0.53, $\eta^2_p = 0.02$), or orientation ($F(1, 28) = 1.42$, p = 0.24, $\eta^2_p = 0.05$), and no significant interaction ($F(1, 28) = 0.83$, p = 0.37, $\eta^2_p = 0.03$).

Spectral Results

Figure 6A shows the power spectra plots for all four conditions for frequency range 1-30 Hz at electrode Fz. Resting state baseline spectra for eyes open-close conditions is also depicted. These resting-state power spectra show a predominant peak within the alpha frequency range (8-12 Hz) which is larger during the eyes closed condition. Power topographies for the preferred and non-preferred clockwise conditions (upper region) and preferred and non-preferred counterclockwise conditions (lower region) were generated for the alpha frequency range. Topographies show higher power distribution across central parietal-occipital regions particularly in the non-preferred conditions. Figure 6B shows the power spectra for all conditions and baseline at electrode Pz. Figures 6C-D depict the Fooof - background EEG spectra used for statistical analysis. This spectra plot shows a clear peak in the alpha frequency in the resting state condition that is more predominant in the eyes closed condition. These spectra plots also reveal an increase in the Beta activity (13-30 Hz) in all the riding conditions relative to the eyes open spectra at electrodes Fz and Pz likely due to muscle activity. Relative to the resting state, there is a general decrease of alpha power in the riding conditions. A one-way repeated measures ANOVA was conducted in order to assess the differences in alpha peak for all conditions at electrodes Fz and Pz. A Greenhouse-Geisser correction was applied to the analysis ($\varepsilon_{Fz} = 0.38$, $\varepsilon_{Pz} = 0.44$). Significant effects of condition type were observed for electrode Fz ($F(1.91, 49.66) = 84.84$, p < 0.001, $\eta^2_p = 0.76$) and electrode Pz ($F(2.12, 57.50) = 58.01$, p < 0.001, $\eta^2_p = 0.69$).

Table 3 shows the significant pairwise comparisons. All p-values were adjusted for multiple comparisons using Bonferroni correction. This analysis shows that both resting state conditions (eyes open and eyes closed) are significantly different from all skateboard conditions at electrode Pz and marginally significant at electrode Fz. There are no further significant differences between conditions. We conducted a two-way repeated measures ANOVA using preference and clock orientation as factors at electrode Fz and Pz. We did not find a main effect for either riding preference ($F(1, 26) = 0.38$, $p = 0.57$, $\eta^2_p = 0.01$) or riding orientation ($F(1, 26) = 3.46$, $p = 0.07$, $\eta^2_p = 0.009$). No significant main effects were found at electrode Pz for preference ($F(1, 26) = 1.65$, $p = 0.20$, $\eta^2_p = 0.060$) or riding orientation ($F(1, 26) = 0.2$, $p = 0.603$, $\eta^2_p = 0.01$).

Left-Right Hemispheric Alpha by Preference and Riding Orientation

We conducted an additional exploratory analysis to measure whether peripheral distractors influence alpha lateralization (due to the role of alpha synchronization in suppressing irrelevant areas of the visual field), we compared the left minus right hemisphere difference in alpha band power based on riding orientation (clockwise vs counterclockwise).

Figure 7 shows the left and right parieto-occipital alpha power by riding direction in the clockwise (left side panel) and counterclockwise (right panel) conditions. We conducted a t-test of the power difference between left-right hemispheres in the clockwise and counterclockwise conditions. Such t-test revealed no differences in hemispheric alpha power between conditions ($t(28) = 0.12$, $p = .90$ $d = 0.02$).

Discussion

The present study employed a mobile EEG paradigm to successfully record ERP components and oscillatory activity from participants moving on an e-skateboard. Participants responded to an auditory oddball task while riding around a complex real world scenario in their

preferred and nonpreferred stances. To our knowledge, this is the first portable EEG experiment using active skateboarding. We chose to use a skateboarding paradigm for several reasons: We wanted to deploy the mobile EEG methodology previously developed in our laboratory (Kuziek et al, 2017; Scanlon et al, 2017) to a novel scenario using a new task, as it is important to demonstrate that this mobile EEG methodology can help researchers design and test attentional paradigms away from the laboratory. Our study further demonstrates the robustness of the oddball-P3 effect under a novel task, providing further evidence that classic attentional paradigms can be replicated under less controlled environments.

Additionally, the implementation of the e-skateboards (and e-scooters) offers a simple solution to the problem of motion artifacts during EEG data collection while allowing participants to freely move in space. Considerable efforts have been made to ensure that the contribution of motion artifacts in the EEG data is managed effectively, such as the integration of portable EEG in experiments involving movement and action. For example, the MoBI approach (Makeig et al., 2009) simultaneously records EEG, muscle movements and environmental events in order to understand cognition using naturalistic behaviors. Other studies involving data collection during movement (Gramman et al., 2010; Gwin et al,.2010) have shown that independent component analysis (ICAs) can be a practical solution to accounting for movement artifacts during complex behaviors. Scanlon et al., (2017) compared the amount of single trial EEG noise between stationary pedaling and sitting and found that while cycling adds more noise to the data, an adequate amount of trials can yield reliable ERP recordings. Taken together, these mobile EEG studies involving complex behavior and outdoor recordings demonstrate that it is possible to obtain valid recordings under less controlled settings. The e-skateboard mobile paradigm presented in the current report offers great flexibility for studies looking to explore

voluntary navigation in complex spaces, particularly when body movements need to be controlled.

Behavioral Results

In the current study, participants responded to target tones by pressing a button with their left hand. We calculated response accuracy and reaction time based on the four experimental conditions (preferred-clockwise, preferred-counterclockwise, non-preferred-clockwise, and non-preferred-counterclockwise) and by the grand averaged preferred and non-preferred conditions. We found no significant differences in accuracy or reaction time across conditions, even though accuracy was marginally higher in the preferred stance condition. Additionally, there was larger standard deviation in the non-preferred condition for both accuracy and reaction time, suggesting that participants performed more uniformly in the preferred stance. These results contrast previous findings, such as Reiser et al. (2019), where increases in motor task difficulty in an oddball paradigm were associated with poorer task performance. It is possible that behavioral effects could not be achieved by our oddball task due to a ceiling effect since mean accuracy was quite high in all conditions.

P3 & MMN/N2b Results

We hypothesized that the target-related P3 would be larger in the preferred stance condition due to differences in riding difficulty. Previous research shows that an increase in task difficulty (i.e., riding in one's least comfortable stance) should be reflected in a reduction of P3 amplitude (Kok, 2001; De Sanctic et al., 2014). While we were able to measure significant, and expected, differences between ERPs following standard and target tones (we were able to detect a P3 response following rare stimuli as shown in Figure 2B and Table 2) we found no significant P3 differences between our conditions. Previous mobile EEG studies in naturalistic

environments have shown a P3 amplitude decrease during naturalistic behaviors such as walking

(Ladouce et al., 2019) and cycling (Zink et al., 2016; Scanlon et at., 2019; Scanlon et al., 2020).

We expected that the increase in difficulty in the non-preferred riding condition was going to be

marked by a reduction in P3 amplitude. However, we failed to find differences in our current

paradigm. Similar results were reported by Gramman et al. (2010), where they show no

differences in P3 amplitude based on increases in motor demands in a visual oddball task.

Considering that no behavioral differences in accuracy and reaction time were found by

increasing riding difficulty, it is possible that the change in stance preference did not interfere

with the cognitive resources employed by the oddball task. Future paradigms should consider

increasing the oddball task difficulty (e.g., manipulating tone frequency), or the non-preferred

task difficulty (e.g. wobbly skateboards).

We also found a MMN/N2b difference between target-standard tones in all our

experimental conditions. These results replicate previous EEG findings from other cycling

studies in our laboratory (Scanlon et al., 2017; Scanlon et al., 2019). This ERP component is

elicited by sudden changes or deviations following the repeated presentation of a stimulus, such

as the change from target to standard tones (Patel & Azzam, 2005). While the standard-target

amplitude difference is present in all our conditions, we did not find differences in this

component based on riding preference. The lack of differences in MMN/N2b amplitude is likely

due to the environmental noise being the same in all task conditions for participants, reflected in

similar ERP amplitudes across conditions.

Spectral Results

We recorded each participant's eyes open/close baseline spectra prior to completing the

oddball task. We found an expected and significant increase in power within the alpha range (8-

12 Hz) at electrodes Fz and Pz for the eyes-closed condition compared to eyes-open, as shown in

Table 3. Additionally, we found a complete attenuation of alpha power during the skateboard

task relative to both baseline conditions at electrode Pz. At electrode Fz, this general reduction of

alpha power between resting state and riding was found to be overall marginally significant at

electrode Fz (table 3). These results support the findings of previous studies showing a decrease

in alpha during complex behaviors such as walking (Beurskens et al., 2016; Storzer et al., 2016)

and bicycle riding (Zink et al., 2016; Scanlon et al., 2020). However, it is important to note that

this reduction in alpha power should not be taken as evidence that the experimental manipulation

leads to quantifiable reductions in alpha since participants did not complete the oddball task

during the resting conditions. Instead, the overall reduction observed in the present study is the

result of the overall engagement with the task in a busy outdoor environment. One of the goals of

the present study was to assess whether increases in riding difficulty would be reflected in

reduction in the alpha band, based on our understanding that alpha power is modulated by

increases in task demand (Foxe, et al., 2011). However, there were no significant differences in

alpha power based on riding preference. Since we found an overall decrease in power during the

task relative to their resting state power, it is possible that the desynchronization in alpha power

we found is likely due to an overload of incoming stimuli from the task and environment

(Klimesch et al., 2011). We found an unexpected increase of power in the beta band (13-30 Hz)

in the skateboarding conditions relative to the resting spectra (Figure 6). We attribute this

increase of power to increased muscle movements. Similar results were reported by Scanlon et

al., (2019), showing an overall increase in beta power during outdoor cycling. This beta increase

during the riding conditions contradicts previous findings linking beta desynchronization to

active movements (Jain et al., 2013).

An exploratory goal in the present study was to measure the potential influence of peripheral distractors in alpha lateralization suppression. Generally speaking, participants consistently experienced a higher number of distractors in the inner tracks of the running pavilion due to joggers and track users present while participants completed the experimental task. Based on previous findings showing shifts in hemispheric alpha based on the location of spatial distractors (Sauseng et al., 2005; Kelly et al, 2006; Ikkai et al., 2016), we were motivated to compare parieto-occipital alpha power across hemispheres to measure whether we could get an effect in overall lateralization without systematically manipulating distractor location. We found no significant differences in mean alpha power in parieto-occipital regions. This is likely due to the overall decrease in alpha power also observed at Fz and Pz in the main spectral analysis. We did not find any reliable effects in terms of alpha lateralization, but more suitable cueing paradigms that fully isolate distractor location could help in answering whether alpha lateralization to visual distractors during active motion occurs.

While this research was able to successfully elicit the expected effects related to the ERP and spectral differences for the oddball and baseline tasks respectively, there are still several limitations of this study that must be further discussed. One such limitation is that we failed to quantify the participants' perceived riding difficulty between the preferred and non-preferred conditions. This could have informed us about the degree of riding interference experienced by participants in the non-preferred condition and the relationship between subjective riding difficulty and overall ERP amplitude. It could be possible that the expected increase in difficulty between the preferred and nonpreferred conditions is being nullified by the electric skateboard in a way that participants can easily adapt to riding in their non-preferred stance since the board is self-propelling and very stable. We also did not measure torso muscle activity and acceleration

on the body. Skateboarding has little overt movement but relies on balance using the torso and proximal muscles of the legs and arms. Further research could correlate changes in these muscle groups during different riding conditions with brain activity and sensory input in order to objectively quantify muscle involvement and motor task interference. Additionally, while this skateboard paradigm offers the opportunity to greatly minimize covert movements (in comparison to walking or bicycle riding paradigms), it does not eliminate muscle movements methodically. Another limitation in the present study is the lack of an experimental condition where participants complete the oddball task while not moving. Our initial motivation to conduct this study was to test whether a change in riding difficulty leads to measurable changes in the ERP and oscillatory domains. While the inclusion of a non riding oddball condition would have improved the present results, we believe that the preferred vs nonpreferred comparison is suitable for the present research question.

The auditory oddball task used did not result in significant behavioral, ERP, or spectral alpha power differences between conditions, contrary to our expectations. Given that the field of mobile EEG is developing, obtaining behavioral effects in line with ERP results is an important step in understanding brain processes in real-life scenarios. This limitation may be attributable to our sample size, but we don't believe this is the case. Given that we are able to measure the expected oddball and spectral alpha differences, with large effect sizes, we believe the lack of difference between our skateboarding conditions is due to other limitations. The auditory oddball task we used may be too easy, with any behavioral or electrophysiological changes hitting a ceiling regardless of condition. A possible solution would be to utilise a more difficult auditory oddball task where the stimuli are more similar or introduce distractor stimuli throughout the

task. Given that participants are on a constantly moving skateboard, with a constantly changing visual environment, a visual oddball task may be more appropriate and attentionally demanding.

Conclusion

In the current study we developed a skateboard EEG paradigm and replicated the classic oddball P3 effect while participants freely skated through a busy running track. While we found consistent target-standard tone differences in the P3 and MMN/N2b in frontal and parietal regions, we did not observe a reduction in P3 amplitude following increases in riding difficulty. We also present evidence that the classic peak in resting state alpha completely diminishes while skateboarding, and this overall desynchronization is the result of an increased load of incoming stimuli from the environment and task. The results in the current study support the notion that EEG paradigms are suitable for the study of human cognition under high ecological validity.

Acknowledgements

This research was supported by start-up funds from the faculty of science and an NSERC Discovery grant (#RES0024267) awarded to Kyle Elliot Mathewson. The authors thank Pat Makhacheva for assistance in data collection, and Sarah Sheldon for analysis feedback.

Conflicts of Interest

No conflicts of interests have been declared by author(s).

Author contribution

Robles: design, data acquisition, analysis/interpretation, drafting

Kuziek: design, data acquisition, analysis/interpretation, drafting

Wlasitz: design, data acquisition, drafting

Bartlett: study conception, design, data acquisition

Hurd: study conception, design, critical revisions

Mathewson: design, analysis/interpretation, drafting, critical revisions

Data Accessibility Statement

This study's supporting data and analysis material is open to access using the following link:

https://github.com/APPLabUofA/Skateboard

Abbreviations

ERP: Event-Related Potential, EEG: Electroencephalography

References

Askamp, J., & van Putten, M. J. A. M. (2014) Mobile EEG in epilepsy. *Int. J. Psychophysiol.*, **91**, 30–35.

Barry, R. J., Clarke, A. R., & Johnstone, S. J. (2011) Caffeine and opening the eyes have additive effects on resting arousal measures. *Clin Neurophysiol*, **122**, 2010-2015.

Beurskens, R., Steinberg, F., Antonieeqicz, F., Wolff, W., & Granacher, U. (2016) Neural Correlates of Dual-Task Walking: Effects of Cognitive Versus Motor Interference in Young Adults. *Neural Plast.*, **2016**.

Caplan, J. B., Bottomley, M., Kang, P., & Dixon, R. A. (2015) Distinguishing rhythmic from non-rhythmic brain activity during rest in healthy neurocognitive aging. *NeuroImage*, **112**, 341–352.

Cruz-Garza, J. G., Brantley, J. A., Nakagome, S., Kontson, K., Megjhani, M., Robleto, D., & Contreras-Vidal, J. L. (2017) Deployment of mobile EEG technology in an art museum setting: Evaluation of signal quality and usability. *Front. Hum. Neurosci.*, **11**.

Dasari, D., Shou, G., & Ding, L. (2017). ICA-derived EEG correlates to mental fatigue, effort, and workload in a realistically simulated air traffic control task. *Front. Hum. Neurosci.*, **11**., 297.

D'Atri, A., De Simoni, E., Gorgoni, M., Ferrara, M., Ferlazzo, F., Rossini, P. M., & De Gennaro, L. (2015) Frequency-dependent effects of oscillatory-tDCS on EEG oscillations: A study with better oscillation detection method (BOSC). *Arch. Ital. Biol.*, **153**, 2–3.

De Sanctis, P., Butler, J. S., Malcolm, B. R., & Foxe, J. J. (2014) Recalibration of inhibitory control systems during walking-related dual-task interference: A Mobile Brain-Body Imaging (MOBI) Study. *NeuroImage*, **94**, 55–64.

Delorme, A., & Makeig, S. (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J. Neurosci. Methods*, **134**, 9–21.

Foxe, J. J., & Snyder, A. C. (2011) The role of alpha-band brain oscillations as a sensory suppression mechanism during selective attention. *Front. Psychol.*, **2**.

Gratton, G., Coles, M. G. ., & Donchin, E. (1983) A new method for off-line removal of ocular artifact. *Electroencephalogr. Clin. Neurophysiol.*, **55**, 468–484.

Gramann, K., Gwin, J. T., Bigdely-Shamlo, N., Ferris, D. P., & Makeig, S. (2010) Visual evoked responses during standing and walking. *Front. Hum. Neurosci.*, **4**, 202.

Gramann, K., Ferris, D. P., Gwin, J., & Makeig, S. (2014) Imaging natural cognition in action. *Int. J. Psychophysiol.*, **9**, 22-29.

Gwin, J. T., Gramann, K., Makeig, S., & Ferris, D. P. (2010) Removal of movement artifact from high-density EEG recorded during walking and running. *J. Neurophysiol.*, **103**, 3526–3534.

Haller, M., Donoghue, T., Peterson, E., Varma, P., Sebastian, P., Gao, R., Noto, T., Knight, R. T., Shestyuk, A., & Voytek, B. (2018) Parameterizing neural power spectra. *BioRxiv*, 299859.

Hughes, A. M., Whitten, T. A., Caplan, J. B., & Dickson, C. T. (2012) BOSC: A better oscillation detection method, extracts both sustained and transient rhythms from rat hippocampal recordings. *Hippocampus*, **22**, 1417–1428.

Ikkai, A., Dandekar, S., & Curtis, C. E. (2016) Lateralization in Alpha-Band Oscillations Predicts the Locus and Spatial Distribution of Attention. *PLOS ONE*, **11**, e0154796.

Jungnickel, E., & Gramann, K. (2016) Mobile brain/body imaging (MoBI) of physical interaction with dynamically moving objects. *Front. Hum. Neurosci.*, **10**, 306.

Kelly, S. P., Gomez-Ramirez, M., & Foxe, J. J. (2009) The strength of anticipatory spatial biasing predicts target discrimination at attended locations: A high-density EEG study. *Eur. J. Neurosci.*, **30**, 2224–2234.

Jain, S., Gourab, K., Schindler-Ivens, S., & Schmit, B. D. (2013). EEG during pedaling: evidence for cortical control of locomotor tasks. *Clin. Neurophysio*, **124**, 379-390.

Kelly, S. P., Lalor, E. C., Reilly, R. B., & Foxe, J. J. (2006) Increases in alpha oscillatory power reflect an active retinotopic mechanism for distracter suppression during sustained visuospatial attention. *J. Neurophysiol.*, **95**, 3844–3851.

Klimesch, W., Doppelmayr, M., Russegger, H., Pachinger, T., & Schwaiger, J. (1998) Induced alpha band power changes in the human EEG and attention. *Neurosci. Lett.*, **244**, 73–76.

Kok, A. (2001) On the utility of P3 amplitude as a measure of processing capacity. *Psychophysiology*, **38**, 557–577.

Kontson, K. L., Megjhani, M., Brantley, J. A., Cruz-Garza, J. G., Nakagome, S., Robleto, D., White, M., Civillico, E., & Contreras-Vidal, J. L. (2015) Your brain on art: Emergent cortical dynamics during aesthetic experiences. *Front. Hum. Neurosci.*, **9**.

Krigolson, O. E., Williams, C. C., Norton, A., Hassall, C. D., & Colino, F. L. (2017). Choosing MUSE: Validation of a low-cost, portable EEG system for ERP research. *Front. Neurosci.*, **11**, 109.

Kuziek, J. W. P., Redman, E. X., Splinter, G. D., & Mathewson, K. E. (2018) Increasing the mobility of EEG data collection using a Latte Panda computer. *J. Neurosci. Methods*, **308**, 34–47.

Ladouce, S., Donaldson, D. I., Dudchenko, P. A., & Ietswaart, M. (2019) Mobile EEG identifies the re-allocation of attention during real-world activity. *Sci. Rep.*, **9**, 1-10

Liebherr, M., Weiland-Breckle, H., Grewe, T., & Schumacher, P. B. (2018). Cognitive performance under motor demands–On the influence of task difficulty and postural control. *Brain Res J.*, **1684**, 1-8.

Makeig, S., Gramann, K., Jung, T. P., Sejnowski, T. J., & Poizner, H. (2009) Linking brain, mind and behavior. *Int. J. Psychophysiol.*, **73**, 95–100.

Packheiser, J., Schmitz, J., Pan, Y., El Basbasse, Y., Friedrich, P., Güntürkün, O., & Ocklenburg, S. (2020) Using Mobile EEG to Investigate Alpha and Beta Asymmetries During Hand and Foot Use. *Front. Neurosci.*, **14**.

Patel, S. H., & Azzam, P. N. (2005) Characterization of N200 and P300: Selected studies of the Event-Related Potential. *Int. J. Med. Sci.,* **2**.

Picton, T. W. (1992) The P300 wave of the human event-related potential. *J. Clin. Neurophysiol.*, **9**, 456–479.

Puce, A., & Hämäläinen, M. S. (2017) A review of issues related to data acquisition and analysis in EEG/MEG studies. *Brain Sci.,* **7.**

Reiser, J. E., Wascher, E., & Arnau, S. (2019) Recording mobile EEG in an outdoor environment reveals cognitive-motor interference dependent on movement complexity. *Sci. Rep.*, **9**, 1-14.

Sauseng, P., Klimesch, W., Stadler, W., Schabus, M., Doppelmayr, M., Hanslmayr, S., Gruber, W. R., & Birbaumer, N. (2005) A shift of visual spatial attention is selectively associated with human EEG alpha activity. *Eur. J. Neurosci.*, **22**, 2917–2926.

Scanlon, J. E. M., Redman, E. X., Kuziek, J. W. P., & Mathewson, K. E. (2020) A ride in the park: Cycling in different outdoor environments modulates the auditory evoked potentials. *Int. J. Psychophysiol.*, **151**, 59–69.

Scanlon, J. E. M., Sieben, A. J., Holyk, K. R., & Mathewson, K. E. (2017) Your brain on bikes: P3, MMN/N2b, and baseline noise while pedaling a stationary bike. *Psychophysiology*, **54**, 927–937.

Scanlon, J. E. M., Townsend, K. A., Cormier, D. L., Kuziek, J. W. P., & Mathewson, K. E. (2019) Taking off the training wheels: Measuring auditory P3 during outdoor cycling using an active wet EEG system. *Brain Res.*, **1716**, 50–61.

Sirevaag, E. J., Kramer, A. F., Wickens, C. D., Reisweber, M., Strayer, D. L., & Grenell, J. F. (1993) Assessment of pilot performance and mental workload in rotary wing aircraft. *Ergonomics*, **36**, 1121–1140.

Squires, N. K., Squires, K. C., & Hillyard, S. A. (1975) Two varieties of long-latency positive waves evoked by unpredictable auditory stimuli in man. *Electroencephalogr. Clin. Neurophysiol.*, **38**, 387–401.

Storzer, L., Butz, M., Hirschmann, J., Abbasi, O., Gratkowski, M., Saupe, D., Schnitzler, A., & Dalal, S. S. (2016) Bicycling and walking are associated with different cortical oscillatory dynamics. *Front. Hum. Neurosci.*, **10**.

Wagner, J., Solis-Escalante, T., Scherer, R., Neuper, C., & Müller-Putz, G. (2014) It's how you get there: walking down a virtual alley activates premotor and parietal areas. *Front. Hum. Neurosci.*, **8**, 93.

Whitten, T. A., Hughes, A. M., Dickson, C. T., & Caplan, J. B. (2011) A better oscillation detection method robustly extracts EEG rhythms across brain state changes: The human alpha rhythm as a test case. *NeuroImage*, **54**, 860–874.

Zamrini, E. Y., Meador, K. J., Thompson, W. O., & Lee, G. P. (1991) Reproducibility of P3. *Int. J. Neurosci.*, **61**, 113-120.

Zink, R., Hunyadi, B., Van Huffel, S., & De Vos, M. (2016) Mobile EEG on the Bike: Disentangling Attentional and Physical Contributions to Auditory Attention Tasks. *J. Neural Eng.*, **13**.

Tables

Table 1: Trial count information for targets and standards in each condition.

Condition	Stimulus Type	Mean Trial Count	Standard Deviation	Trial Count Range (Minimum:Maximum)
Preferred Clockwise	Targets	46.90	4.87	29:50
	Standards	185.83	21.61	104:200
Preferred CounterClockwise	Targets	47.79	5.06	26:50
	Standards	191.52	17.35	115:200
Non-preferred Clockwise	Targets	47.90	4.47	30:50
	Standards	190.34	17.59	131:200
Non-preferred CounterClockwise	Targets	47.76	4.56	29:50
	Standards	189.79	20.75	92:200

Table 2: Pairwise results comparing target and standard ERP waveforms in each condition.

Comparison	Electrode/ Time Window	Mean Voltages (µV) (Targets:Standards)	$t(28)$	p	Confidence Interval	Effect Size (Cohen's d)
Preferred Clockwise	Fz/MMN-N2b	-3.23:-0.58	-5.43	<0.001	-4.23:-1.09	-1.23
	Pz/P3	4.71:-0.47	5.98	<0.001	2.36:8.01	1.32
Preferred CounterClockwise	Fz/MMN-N2b	-2.86:-0.72	-4.37	<0.001	-3.71:-0.57	-0.90
	Pz/P3	4.35:-0.54	5.64	<0.001	2.07:7.71	0.94
Non-preferred Clockwise	Fz/MMN-N2b	-3.96:-0.75	-6.56	<0.001	-4.78:-1.64	-0.97
	Pz/P3	4.59:-0.78	6.20	<0.001	2.55:8.20	1.18
Non-preferred CounterClockwise	Fz/MMN-N2b	-3.10:-0.30	-5.71	<0.001	-4.37:-1.23	-1.10
	Pz/P3	3.95:-0.05	4.61	<0.001	1.17:6.82	0.82

Table 3: Pairwise results comparing A) Baseline Eyes Closed and B) Baseline Eyes Open spectra to each of the other five conditions.

A

Comparison	Electrode	Mean Spectra Difference (μV^2)	$t(26)$	p	Confidence Interval	Effect Size (Cohen's d)
Baseline	Fz	-0.51	-13.46	<0.001	-0.62:-0.39	2.59
Eyes Open	Pz	-0.50	-9.79	<0.001	-0.22:-0.32	1.88
Preferred	Fz	0.62	16.39	<0.001	0.50:0.73	3.15
Clockwise	Pz	0.67	13.20	<0.001	0.52:0.83	2.54
Preferred	Fz	0.61	16.39	<0.001	0.50:0.73	3.13
CounterClockwise	Pz	0.68	13.23	<0.001	0.52:0.83	2.54
Non-preferred	Fz	0.61	16.26	<0.001	0.50:0.73	3.12
Clockwise	Pz	0.70	13.72	<0.001	0.55:0.85	2.64
Non-preferred	Fz	0.61	16.16	<0.001	0.50:0.72	3.11
CounterClockwise	Pz	0.69	13.52	<0.001	0.54:0.84	2.60

B

Comparison	Electrode	Mean Spectra Difference (μV^2)	$t(26)$	p	Confidence Interval	Effect Size (Cohen's d)
Baseline	Fz	-0.51	-13.46	<0.001	-0.62:-0.39	-2.59
Eyes Closed	Pz	-0.50	-9.79	<0.001	-0.65:-0.35	-1.88
Preferred	Fz	0.11	2.93	0.060	-0.002:0.22	0.56
Clockwise	Pz	0.17	3.41	0.013	0.02:0.32	0.65
Preferred	Fz	0.10	2.82	0.083	-0.006:0.22	0.54
CounterClockwise	Pz	0.17	3.90	0.012	0.02:0.33	0.66
Non-preferred	Fz	0.10	2.80	0.088	-0.007:0.22	0.53
Clockwise	Pz	0.20	3.93	0.002	0.04:0.35	0.75
Non-preferred	Fz	0.10	2.70	0.11	-0.011:0.21	0.52
CounterClockwise	Pz	0.19	3.72	0.004	0.03:0.34	0.71

Figure Captions

Figure 1. Setup task and Location. A: Mobile EEG setup. B: Task diagram. C: Experiment location

Figure 2. Grand-averaged ERP across conditions used to select time windows for analyses.

Figure 3. Behavioral Task Results. A: Mean target accuracy. B: Median target reaction time. C: Target accuracy by grand preference. Target reaction time by grand preference

Figure 4: A: ERPs at electrode Fz. Conditions (left to right): Preferred clockwise, preferred counterclockwise, non-preferred clockwise, non-preferred counterclockwise. Target tones: colored lines. Standard tones: black lines. Shaded regions represent standard errors. Topographies for MMN/N2 window. B: ERPs at Electrode Pz. and topographies for P3 time windows.

Figure 5. Difference-wave ERPs. A: Difference-wave ERP by grand preference for the MMN/N2b time window at electrode Fz. B: Difference-wave ERP by grand clock orientation for the MMN/N2b time window at electrode Fz. C: Difference-wave ERP by grand preference for the P3 time window at electrode Pz. D: Difference-wave ERP by grand clock orientation for the P3 time window at electrode Pz.

Figure 6. A: Power spectra at electrode Fz for all conditions plus resting state baseline (eyes opened/closed). Shaded regions represent standard errors for all conditions. Power topographies for alpha band range (8-12Hz) for preferred clockwise and counterclockwise conditions (upper region) and non-preferred clockwise and counterclockwise conditions (lower region). B: Power spectra at electrode Pz for all conditions. C: Fooof - background data used for statistical analysis at electrode Fz. D: Fooof - background data use for statistical analysis at electrode Pz.

Figure 7. Power spectra for left and right parieto-occipital hemispheres for clockwise direction (left) and counterclockwise direction (right).

1

A

B

Pref-CW, Pref-CCW, NonPref-CW,
NonPref-CCW
(Randomized order)

| 250 trials | 250 trials | 250 trials | 250 trials |

1-1.5
sec

1000 Hz (standard; 4:1 ratio)   2000 Hz (target; 1:4 ratio)

C

2

3

Difference-Wave - Preference

Difference-Wave - Clock Orientation

Clockwise left-right difference

Counterclockwise left-right difference