A novel assay for tracking carboxylesterase gene amplifications conferring organophosphate resistance in the mosquito *Aedes aegypti*: from experimental evolution to field application

Running head: genomic amplification associated with insecticide resistance

Julien Cattel[1][§][*] | Chloé Haberkorn[1] | Fréderic Laporte[1] | Thierry Gaude[1] | Tristan Cumer[1] | Julien Renaud[1] | Ian W. Sutherland[2] | Jeffrey C. Hertz[3] | Jean-Marc Bonneville[1] | Victor Arnaud[1] | Camille Noûs[4] | Bénédicte Fustec[5,6] | Sébastien Boyer[7] | Sébastien Marcombe[8] | Jean-Philippe David[1]

[1]Laboratoire d'Ecologie Alpine (LECA), UMR 5553 CNRS – Université Grenoble-Alpes, 2233 rue de la piscine, Grenoble, France.
[2]United States Navy Entomology. Center of Excellence, NAS Jacksonville, Florida, United States of America.
[3]US Naval Medical Research Unit No. 2, Singapore.
[4]Laboratoire Cogitamus, 1 ¾ rue Descartes, 75005 Paris, France.
[5]Department of Microbiology, Khon Kaen University, Khon Kaen, Thailand.
[6]Institut de Recherche pour le Développement, UMR IRD 224-CNRS 5290-Université Montpellier. 911, avenue Agropolis, BP 64501, 34394 Montpellier Cedex 5, FRANCE.
[7]Medical and Veterinary Entomology, Institut Pasteur du Cambodge, P.O Box. 983 Phnom Penh, Cambodia.
[8]Medical Entomology and Vector-Borne Disease Laboratory, Institut Pasteur du Laos, Vientiane, Laos.
[§]Current address: Symbiosis Technologies for Insect Control (SymbioTIC). Plateforme de Recherche Cyroi, 2 rue Maxime Rivière, 97490 Ste Clotilde, France.

*Correspondence:
Julien Cattel,
Symbiosis Technologies for Insect Control (SymbioTIC)
Plateforme de recherche Cyroi, 2 rue Maxime Rivière, 97490 Ste Clotilde, France
E-mail: juliencattel@gmail.com

Email of all authors:
JC: juliencattel@gmail.com
CH: chloehbk@gmail.com
FL: frederic.laporte@univ-grenoble-alpes.fr
TG: thierry.gaude@univ-grenoble-alpes.fr
TC: t.cumer.sci@gmail.com
JR: julien.renaud@univ-grenoble-alpes.fr
IWS: ian.w.sutherland.mil@mail.mil
JCH: jeffrey.c.hertz.mil@mail.mil
JMB: jean-marc.bonneville@univ-grenoble-alpes.fr
VA: victor.arnaud19@gmail.com
CN: camille.nous@cogitamus.fr
BF: b.fustec@gmail.com
SB: sboyer@pasteur-kh.org
SM: s.marcombe@pasteur.la
JPD: jean-philippe.david@univ-grenoble-alpes.fr

Article type: Original Research article

Abstract

By altering gene expression and offering a hold for selection by creating paralogs, genomic duplications represent a key component of short-term adaptive processes. In insects, the use of insecticides can select gene duplications causing an increased expression of detoxification enzymes, supporting their usefulness for monitoring the dynamics of resistance alleles in the field. In this context, the present study aimed at characterising a genomic amplification event associated with resistance to organophosphate insecticides in the mosquito *Aedes aegypti* and developing a molecular assay to monitor the associated resistance alleles in the field. An experimental evolution experiment using a composite population from Laos confirmed the association between the over-transcription of multiple contiguous carboxylesterase genes on chromosome 2 and resistance to multiple organophosphate insecticides. Combining whole genome sequencing and qPCR on specific genes confirmed the presence of a large gene amplification at this locus with the co-existence of multiple structural duplication haplotypes. Field data confirmed their circulation in South-East Asia and revealed high copy number variations among and within populations suggesting a trade-off between this resistance mechanism and associated fitness costs. A dual-colour multiplex TaqMan assay allowing the rapid detection and copy number quantification of this duplication event in *Ae. aegypti* was developed and validated on field populations. The routine use of this novel assay shall improve resistance tracking in this major arbovirus vector.

Key words: genomic amplification, insecticide resistance, mosquito, *Aedes aegypti*, carboxylesterase, diagnostic assay

1 | INTRODUCTION

The duplication of genomic regions overlapping genes can affect fitness by creating paralogs that can diverge to generate novel functions or by altering gene expression levels (Iskow, Gokcumen, & Lee, 2012). Often called genomic amplification when multiple copies are present, such mechanism has been shown to be a major driver of short-term adaptation (Kondrashov, 2012). Indeed, duplication events are frequent in natural populations (Campbell & Eichler, 2013) and can be subjected to positive selection (Chatonnet, Lenfant, Marchot, & Selkirk, 2017; Cooper, Nickerson, & Eichler, 2007; Zhang, 2003).

In insects, genomic amplifications have been shown to play a key role in the evolution of insecticide resistance by two distinct mechanisms. First, the duplication of genes encoding insecticide targets can allow resistant individuals to reduce the fitness costs associated with target-site mutations by carrying both the susceptible and the resistant alleles (Assogba et al., 2015; Labbé et al., 2007). Second, genomic amplifications affecting genes encoding detoxification enzymes leading to their over-expression can confer the insect a higher capacity to degrade or sequester insecticides (Bass & Field, 2011). This later mechanism has been reported in all three mosquitoes genera of high medical importance, *Aedes*, *Anopheles* and *Culex*, affecting most detoxification enzymes including cytochrome P450 monooxygenases (*P450*s) and carboxylesterases (*CCE*s) (Cattel et al., 2019; Lucas et al., 2019; Weetman, Djogbenou, & Lucas, 2018).

Genomic amplifications affecting *CCE*s have even been described as the most common route of enzyme over-production in mosquitoes (Bass & Field, 2011). A classic example comes from the house mosquito *Culex pipiens* in which amplified carboxylesterases genes (in conjunction with the *ace-1* G119S target-site mutation) have spread across the globe, providing high resistance to organophosphate insecticides (Raymond, Berticat, Weill, Pasteur, & Chevillon, 2001). In *Aedes* mosquitoes, the low chance of occurrence of the G119S *ace-1*

mutation because of strong genetic constraints (Weill et al., 2004) suggest that *CCE*

amplifications play a central role in organophosphate resistance and are thus of high interest for

resistance monitoring.

In the tiger mosquito *Aedes albopictus*, the over-expression of two *CCE* genes (*CCEae3A* and

CCEae6A) through gene amplification was associated with resistance to the organophosphate

insecticide temephos (Grigoraki et al., 2017). In the yellow fever mosquito *Aedes aegypti*,

AAEL023844 (formerly *CCEAE3A*) and other *CCE* genes belonging to the same genomic

cluster were also found overexpressed through gene amplification in temephos-resistant

populations (Faucon et al., 2015, 2017; Poupardin, Srisukontarat, Yunta, & Ranson, 2014).

Further functional studies confirmed that *CCEAE3A* is able to sequester and metabolize the

active form of temephos in both *Ae. aegypti* and *Ae. albopictus* (Grigoraki et al., 2016).

Although the genomic structure and polymorphism of this *CCE* amplification was studied in

Ae. albopictus (Grigoraki et al., 2017) such work has not been conducted in *Ae. aegypti*. In

addition, the role of this *CCE* amplification in resistance to other insecticides remains unclear.

Finally, no high-throughput assay has yet been developed to track this resistance mechanism in

natural populations although such tool would significantly ease resistance monitoring and

management.

In this context, we combined an experimental evolution experiment with RNA-seq and

whole genome sequencing data to confirm the association between this genomic amplification,

the overexpression of *CCE* genes and resistance to organophosphate insecticides in *Ae. aegypti*.

We also showed that this *CCE* amplification confers resistance to multiple organophosphates

insecticides. Comparing gene Copy Number Variations (CNV) between the different genes of

this genomic cluster suggested the presence of at least two distinct haplotypes occurring in

South-East Asia (SEA), both associated with resistance. Investigating their spatial dynamics in

natural populations confirmed their co-occurrence in SEA with a high copy number

polymorphism within and among populations. Based on these results, we developed a novel

high-throughput multiplex TaqMan assay allowing the quantitative detection of this *CCE*

amplification in hundreds of individual mosquitoes within a few hours. By reducing the human

power and infrastructure needs associated with bioassays, this molecular assay will improve the

tracking and management of organophosphate resistance in natural mosquito populations.

2 | MATERIAL AND METHODS

2.1 | Field sampling and mosquito lines

Ae. *aegypti* larvae and pupae were collected in households and temples of eleven villages

belonging to five provinces of Laos in 2014 (Table S1). Previous work confirmed the

circulation of organophosphate and pyrethroid resistance alleles in these populations together

with the presence of duplication affecting AAEL023844 (formerly *CCEAE3A*) (Marcombe et

al., 2019). These populations were maintained under controlled conditions (27 ± 2 ℃ and $80 \pm$

10% relative humidity) at the Institut Pasteur of Laos for 5 generations and used for

experimental evolution (see below).

A second round of mosquito collection was conducted in 2017 for studying the spatial dynamics

of *CCE* genomic amplifications in SEA. Adult females were sampled in 14 different

populations in Laos, Thailand and Cambodia (see details in Table S1) and stored individually

at -20°C in silica gel until molecular analyses.

2.2 | Experimental evolution

A Laos composite population was created by pooling 50 virgin males and 50 virgin females of

each population in a single cage (Table S1). This population was then maintained for 2

generations without insecticide selection to allow genetic mixing before initiating insecticide

selection. The Laos composite population was then split in 2 lines (N > 1000 in each line): one line being maintained without insecticide selection (NS line) while the second line (Mala line) was artificially selected with malathion at the adult stage for 4 consecutive generations (from G1 to G5). For this, batches of thirty-three-days old non-blood fed adult mosquitoes (~1000 individuals of mixed sex) were exposed at each generation to filters papers impregnated with malathion using WHO test tubes. A constant dose of 5% malathion coupled with an exposure time of 10 min (leading to ~90% mortality at G1) were used through the whole selection process. Surviving females were collected 48h after insecticide exposure, blood fed on mice and allowed to lay eggs to generate the next generation.

Three days-old non-blood fed adult females (not exposed to insecticide) were sampled after four generations and used for bioassays and molecular work. Mosquitoes were identified as follows. G1: initial composite population, G5-NS: line maintained without insecticide pressure for four generations, G5-Mala: line maintained under malathion selection for four generations. Sampled mosquitoes were stored at -20°C until molecular analyses.

2.3 | Bioassays

Bioassays were used to monitor the dynamics of malathion resistance during the selection process. Four replicates of 20 calibrated three days-old non-blood fed females not previously exposed to insecticide were sampled at each generation and exposed to the insecticide as described above using the same dose and exposure time as for artificial selection. Mortality was recorded 48h after exposure.

Cross resistance to other insecticide was investigated in G5 individuals (G5-Mala and G5-NS) not previously exposed to insecticide. Calibrated individuals were exposed to three distinct insecticides: the organophosphates fenitrothion and temephos, and the pyrethroid deltamethrin. For the adulticides fenitrothion and deltamethrin, bioassays were performed on eight replicates

of fifteen three days old non-blood-fed females with the following doses and exposure times:

fenithrotion 1% for 30 min, deltamethrin 0.05% for 20 min. Mortality rates were recorded 48h

after exposure. For the larvicide temephos, bioassays were performed on eight replicates of

twenty calibrated third instar larvae exposed to 0.08 mg/μL temephos for 24h in 200 ml tap

water and mortality was recorded at the end of exposure.

2.4 | RNA sequencing

RNA-sequencing was performed to compare gene expression levels between the G6-NS, G6-

Mala lines and the fully susceptible reference line Bora-Bora. For each line, four RNA-seq

libraries were prepared from distinct batches of individuals not exposed to insecticide. For each

library, total RNA was extracted from 25 calibrated 3-days old non-blood fed females using

Trizol® (Thermo Fisher Scientific) following manufacturer's instructions. RNA samples were

then treated with the RNase-free DNase set (Qiagen) to remove gDNA contaminants and QC

checked using Qubit (Thermo Fisher Scientific) and bioanalyzer (Agilent). RNA-seq libraries

were prepared from 500 ng total RNA using the NEBNext® Ultra^{TM} II directional RNA library

Prep Kit for Illumina (New England Biolabs) following manufacturer's instructions. Briefly,

mRNAs were captured using oligodT magnetic beads and fragmented before being reverse

transcribed using random primers. Double stranded cDNAs were synthesized end-repaired and

adaptors were incorporated at both ends. Libraries were then amplified by PCR for 10 cycles

and purified before QC check using Qubit fluorimeter and Bioanalyzer. Libraries were then

sequenced in multiplex as single 75 bp reads using a NextSeq500 sequencer (Illumina).

After unplexing and removing adaptors, sequenced reads from each library were loaded into

Strand NGS (Strand Life Science) and mapped to the latest *Ae. aegypti* genome assembly (Aaeg

L5) using the following parameters: min identity = 90%, max gaps = 5%, min aligned read

length = 25, ignore reads with >5 matches, 3'end read trimming if quality <20, Kmer size = 11,

mismatch penalty = 4, gap opening penalty = 6, gap extension penalty = 1. Mapped reads were

then filtered based on their quality and alignment score as follows: mean read quality > 25, max

N allowed per read = 5, mapping quality ≥120, no multiple match allowed, read length ≥ 35.

Quantification of transcription levels was performed on the 14626 protein-coding genes using

the DESeq method with 1000 iterations (Anders & Huber, 2010). Only the 11825 genes

showing a normalized expression level ≥ 0.5 (~0.05 RPKM) in all replicates for all lines were

retained for further analysis. Differential gene transcription levels between each line across all

replicates were then computed using a one-way ANOVA followed by a Tukey post-hoc test

and P values were corrected using the Benjamini and Hockberg multiple testing correction

(Benjamini & Hochberg, 1995). Genes showing a fold change (FC) ≥ 3 (in any direction) and

a corrected P value ≤ 0.001 in the G6-Mala line versus both susceptible lines (G6-NS and Bora-

Bora) were considered as differentially transcribed in association with insecticide resistance.

2.5 | Whole genome sequencing

The occurrence of a genomic amplification affecting the *CCE* cluster on chromosome 2 at ~174

Mb was investigated by sequencing the whole genome of an organophosphate-resistant

population from Thailand (Nakh population as described in Faucon et al., 2015, 2017) as

compared to the fully susceptible line (Bora-Bora). For each population, genomic DNA was

extracted from 100 adult females as described in Faucon et al., 2015 and whole genome

sequencing was performed from 200 ng gDNA. Sequencing libraries were prepared according

to the TruSeq DNA Nano Reference guide for Illumina Paired-end Indexed sequencing (version

oct 2017) with a mean insert size of 550 bp. Sequencing was performed on a NextSeq 550

(Illumina) as 150 bp paired-reads. After unplexing and adaptor removal, reads were mapped to

the latest *Ae. aegypti* genome assembly (Aaeg L5) using BWA-MEM with default parameters

(version 0.7.12). Sequenced reads were then sorted using samtools sort (version 1.2), annotated

using Picard FixMateInformation (version 1.137) and PCR duplicates were identified using

Picard MarkDuplicates (version 1.137). Coverage profiles between the resistant and the

susceptible lines were then compared using non-duplicated reads with a mapping quality score

above 60 were considered.

2.6 | Quantification of Copy Number Variations

Among the six genes located within the genomic amplification detected on chromosome 2,

three genes clearly annotated as *CCE* and distributed throughout the cluster were studied:

AAEL019678, AAEL023844 (formerly *CCEAE3A*) and AAEL005113 (*CCEAE1A*). For each

gene, specific primer pairs were designed using NCBI primer Blast (Table S2). In order to

quantify CNV in natural populations and in individuals, genomic DNA was extracted either

from seven pools of five adult females (mean CNV comparison between lines) or from single

adult females (estimation of duplication prevalence) using the cetyltrimethylammonium

bromure (CTAB) method (Collins et al., 1987) and diluted to 0.5 ng/µL prior amplification.

Pooled samples were amplified in duplicates while individual mosquito samples were amplified

only once. Quantitative PCR reactions consisted of 3 µL gDNA template, 3,6 µL nuclease free

water, 0,45 µL of each primer (10µM), and 7,5 µL of iQ SYBR Green Supermix (Bio-Rad).

PCR amplification were performed on a CFX qPCR system (Bio-Rad) with cycles as follows:

95°C 3 min followed by 40 cycles of 95°C 15 secs and 30 secs for hybridization. A dilution

scale made from a pool of all gDNA samples was used for assessing PCR efficiency. Data were

analyzed using the $\Delta\Delta$Ct method (Pfaffl, 2001) taking into account PCR efficiency. Two control

genes (the *P450* AAEL007808 and the chloride channel protein AAEL005950) shown to be

present as single copies in multiple *Ae. aegypti* strains and populations (Faucon et al., 2015)

were used for normalization. For each gene, CNV were expressed as mean relative gDNA

quantity as compared to the fully susceptible line Bora-Bora. For assessing genomic

amplification frequencies, all individuals showing a CNV ≥ 2.5-fold as compared to the Bora-

Bora line were considered positive.

Individual CNV levels obtained for the *CCE* gene AAEL023844 by qPCR were cross-validated

by digital droplet PCR (ddPCR). Briefly, each sample was partitioned into ~20,000 nanoliter-

sized droplets using the QX 200 droplet generator (Bio-Rad) by mixing synthetic oil with 20

µL PCR mix containing 2X ddPCR Evagreen supermix (Bio-Rad), 0.9 mM of each primer and

5 µL of template gDNA at 0.5 ng/µL. Emulsified reaction mixtures were then amplified with a

C1000 thermal cycler (Bio-Rad) for 40 cycles. After amplification, the number of positive and

negative droplets were quantified for each sample using the QX 200 droplet reader (Bio-Rad)

and the positive/negative ratio was used to estimate the initial DNA assuming a Poisson

distribution. A similar procedure was applied to the control gene AAEL007808 present as a

single copy. After normalizing for initial gDNA quantity, CNV were expressed as relative

gDNA quantity as compared to the fully susceptible line Bora-Bora.

2.7 | CNV quantification using TaqMan multiplex assay

A TaqMan multiplex assay allowing the concomitant quantification of the *CCE* gene

AAEL023844 (present in both duplication haplotypes) and the control gene AAEL007808 from

single mosquitoes within the same qPCR reaction was developed. For each gene, primers and

probes were designed using Primer3web v 4.1.0 (Rozen & Skaletsky, 2000) with the AaegL5

assembly as reference genome for assessing specificity. For each gene, exonic regions were

targeted in order to limit amplification variations associated with natural polymorphism (Table

S2). The assay was then tested on all individuals detected as positive by qPCR representing 27

individuals belonging to seven populations from three countries. Each reaction mixture

contained 12.5 µL of qPCR probe Master Mix (Bio-Rad), 2.25 µL of each primer (10µM),

0.625 µL of each probe (10 µM), 1.25 µL of nuclease free water and 1 µL of template DNA

(0.5 ng/µL). PCR amplifications were performed on a CFX qPCR system (Bio-Rad) with cycles

set as follows: 95°C for 10 min followed by 40 cycles of 95°C for 10 secs and 60°C for 45 secs

followed by FAM and HEX levels reading (see Supplementary File 1 for a user guideline on

this TaqMan assay).

2.8 | Statistical analysis

All statistical analyses were performed with R v3.6.2 (R Core Team, 2013), using the package

lme4 for all mixed models (Bates et al., 2015). Mortality data were statistically compared across

conditions by using a Generalized Linear Model (GLM) with mixed effects (binomial family)

in which the replicates were included as a random factor. For comparing mean CNV obtained

from pools of mosquitoes, normalized gDNA levels obtained for each gene were Log_2

transformed and compared across conditions using a GLM with mixed-effects in which the

replicates were included as a random factor. For comparing CNV obtained from individual

mosquitoes, normalized gDNA levels were Log_2 transformed and compared between genes

using a one-way ANOVA. A Pearson's product moment correlation coefficient test was used

to compare normalized gDNA quantities obtained from qPCR and ddPCR.

3 | RESULTS

3.1 | Dynamics of organophosphate resistance during experimental selection

Maintaining the Laos composite population under selection with malathion resulted in the rapid

rise of resistance (Figure 1).

FIGURE 1. Dynamics of malathion resistance during the selection process. Black: Laos composite population selected with malathion; Grey: Laos composite population maintained without selection. Stars indicate a significant mortality difference as compared to the initial population (N=4, GLM mixed effects binomial family, *p<0.05).

Mortality to malathion dropped gradually from 90.3% in G1 to 52.2 % after four generations of selection (GLMER test: z=2.058, P<0.05 for G5-Mala vs G1). Conversely, mortality increased to 99.1% after four generations in without selection.

Bioassays performed with different insecticides revealed that selection with malathion for four generations also select resistance to other organophosphate insecticides (Figure 2).

FIGURE 2. Cross resistance of the Malathion-resistant line to other insecticides. G5-NS: Composite population maintained without selection. G5-Mala: composite population selected with malathion. For each insecticide, stars indicate a significant mortality difference between G5-NS and G5-Mala individuals (N=8, GLM mixed effects binomial family, *p<0,05, ***p<0,001, NS: non-significant).

As compared to the G5-NS line, the G5-Mala line showed a significant increased resistance to

the organophosphates fenitrothion at the adult stage (GLMER: z=-3.455, P<0.005) and

temephos at the larval stage (GLMER: z=-2.194, P<0.05). Conversely, no significant difference

was observed between G5-NS and G5-Mala individuals for susceptibility to the pyrethroid

deltamethrin at the adult stage suggesting that malathion selection did not select for pyrethroid

resistance alleles.

3.3 | Genes associated with malathion resistance

RNA-seq analysis identified 84 and 124 genes over- and under-transcribed, respectively, in G6-

Mala adult females (adjusted P value ≤ 0.001 and FC ≥ 3-fold) as compared to the susceptible

lines G6-NS and Bora-Bora (Table S3). Among them, 24 genes encoded proteins potentially

associated with known resistance mechanisms including cuticle alteration (14 genes) and

detoxification (10 genes). Only seven candidate genes were over-transcribed in the G5-Mala

line, all being associated with detoxification (Figure 3A). This included a microsomal

glutathione S-transferase on chromosome 1 (AAEL006818, 13-fold *versus* G6-NS), the

cytochrome P450 *CYP6N17* on chromosome 2 (AAEL010158, 8-fold *versus* G6-NS) and five

contiguous *CCE* genes at ~174 Mb on chromosome 2 (AAEL015304, AAEL019679,

AAEL019678, AAEL005123 and AAEL023844 (formerly *CCEAE3A*, up to 10-fold *versus* G6-

NS). A closer look at this genomic region revealed the presence of an additional *CCE* gene on

the 5' side of the cluster (AAEL005113 *CCEAE1A*) which was not significantly over-

transcribed in the resistant line. Although the *GST* AAEL006818 and the P450 *CYP6N17* were

significantly over-transcribed in the G6-Mala line, these two genes were also found over-

transcribed in two other Laos lines selected with unrelated insecticides (data not shown),

invalidating their specific association with malathion resistance.

FIGURE 3. Genomic amplification associated with carboxylesterase overexpression. A: Detoxification genes over-transcribed in the malathion-resistant line. Transcription levels were quantified by RNA-seq using 4 biological replicates per line. Detoxification genes showing a 3-fold over-transcription and a corrected P value <0.001 in the Malathion-resistant line G6-Mala versus both the unselected line NS (grey) and the fully susceptible line Bora-Bora (black) are shown. The genomic location of each gene is indicated. B: Comparison of read coverage profiles at the carboxylesterase locus between a Thai resistant line (grey) and the fully susceptible line Bora-Bora (black). Coverage profiles were obtained from whole genome DNA-seq performed on pools of 100 individuals. The genomic location of carboxylesterase genes is indicated according to Aaeg L5.1 annotation. Genes found overexpressed by RNA-seq are shown in orange. Genes targeted by qPCR are shown in bold. Repeated elements associated with coverage gaps are indicated as grey boxes.

Cross-comparing RNA-seq data with read coverage profiles obtained from whole genome

sequencing of a Thai resistant population known to over-express these CEE genes (Nakh

population, Faucon et al., 2015, 2017) revealed a ~30-fold increased coverage in this genomic

region as compared to the susceptible line (Figure 3B). In addition, this region revealed several low-coverage sections associated with the presence of repeated elements (mostly due to unresolved read assembly). The pattern of genomic amplification observed in this Thai resistant population, matches the expression pattern observed in the Laos resistant line with the same five CCE genes, but not *CCEAE1A*, found amplified.

The occurrence of this genomic amplification in Laos was then confirmed by qPCR (Figure 4A). Indeed, a slight elevation of gene copy number was observed for three *CCE* genes belonging to this genomic cluster before selection (G1) with important variations suggesting a high inter-individual heterogeneity in the initial line. Gene copy number were even lower after four generations without selection (G5-NS) with less variations observed. Conversely, four generations of selection with malathion lead to a strong increase in gene copy number for the *CCE* genes AAEL019678 and AAE023844 in G5-Mala individuals (up to 14-fold). A lower increase (4-fold) associated with a higher variance was observed for the gene *CCEAE1A*, suggesting that not all G5-Mala individuals carry multiple copies of this gene.

Quantification of *CCE* genes copy number in individual mosquitoes by qPCR confirmed the presence of two distinct structural duplication haplotypes in Laos with haplotype A including the three *CCE* genes and haplotype B not including *CCEAE1A* (Figure 4B). While the prevalence of these two *CCE* haplotypes was low in the initial composite population (7%), their cumulated frequency reached 84% in G5-Mala individuals with haplotype B being more frequent (67%) than haplotype A (33%).

FIGURE 4. Duplication haplotypes at the carboxylesterase locus. A: CNV of three selected CCE genes located in different positions of the carboxylesterase locus. For each gene, the position of the qPCR amplification product is indicated (dashed lines). Mean CNV were estimated by qPCR on pools of mosquitoes and are expressed as gDNA quantity relative to the fully susceptible line Bora-Bora (horizontal grey line). G1: unselected line at generation 1, G5-NS: unselected line at generation 5, G5-Mala: malathion-resistant line after 4 generations of selection. Distinct letters indicate significant mean CNV variations between populations (GLM mixed effects, N=7, $p \leq 0.05$). The two structural duplication haplotypes deduced from CNV data are represented. B: Frequencies of each haplotype in the different lines. Haplotypes frequencies were deduced from individual CNV data obtained by qPCR from 28 mosquitoes per line for the three genes AAEL019678, AAEL023844 (formerly *CCEAE3A*) and AAEL005113 (*CCEAE1A*).

3.4 | Prevalence and copy number polymorphism in SEA

The spatial dynamics of this genomic amplification event was investigated in field populations from Laos, Cambodia and Thailand. A total of 302 mosquitoes belonging to 14 field populations were genotyped for the presence of *CCE* amplifications using qPCR. Seven populations distributed across the three countries were found positive with at least one individual carrying an amplification haplotype A or B (Figure 5). The prevalence of *CCE* amplifications was low in most studied populations, high in two populations from Cambodia (26% and 29%) and very high in a Thai population resistant to organophosphates (79 %). Although both haplotypes were detected through the study area, all positive individuals from populations showing a high prevalence only carried haplotype B.

FIGURE 5. Prevalence of the carboxylesterase gene amplification in South-East Asia. For each population, the frequency of each haplotype is shown. The number of individuals genotyped is shown above each pie chart. Haplotypes frequencies were deduced from CNV data obtained by qPCR on individual mosquitoes for the three genes AAEL019678, AAEL023844 and AAEL005113 (*CCEAE1A*). Only individuals showing a CNV ≥ 2.5-fold (as compared to the susceptible line Bora-Bora) for any gene were considered positive. The description of the studied populations is presented in Table S1.

Cross comparing CNV data obtained for the *CCE* gene AAEL023844 between standard qPCR

and digital droplet PCR (ddPCR) indicated a good correlation between the two techniques

($r=0.85$, $P<0.001$, Figure S1) suggesting that despite the technical variations inherent to qPCR

on single mosquitoes this approach provides a relatively good estimation of gene copy numbers.

Comparing the number of copies of each *CCE* gene between all positive individuals revealed

an important copy number polymorphism in the SEA with estimated copy numbers ranging

from 3 to ~80 copies (Figure 6). No significant differences were observed between field

populations and the different lines (G1, G5-NS, G5-Mala), suggesting that insecticide selection

rather select for positive individuals than for individuals carrying a higher number of copies.

Overall, the mean copy number observed for the *CCE* gene AAEL023844 (present in both

haplotypes) was significantly higher than for the two other *CCE* genes AAEL019678 and

CCEAE1A ($P<0.001$ and $P<0.01$ respectively) possibly reflecting additional structural

haplotypes affecting this gene.

FIGURE 6. Genes copy number variations in experimental lines and field populations. For each gene, CNV were estimated by qPCR on individual mosquitoes and are expressed as gDNA quantity relative to the fully susceptible line Bora-Bora. Only positive individuals showing a CNV ≥ 2.5-fold for any gene are shown. Names of laboratory lines and field populations are indicated.

3.5 | A novel TaqMan assay to track *CCE* gene amplification in *Ae. aegypti*

A multiplex TaqMan qPCR assay allowing the concomitant amplification of the *CCE* gene AAEL023844 (included in both haplotypes) and a control gene within a single reaction was developed and tested against 27 positive individuals and 7 negative individuals belonging to all populations from which *CCE* amplification were detected (Figure 7). This assay showed a good specificity and sensitivity for detecting *CCE* amplifications in *Ae. aegypti*. All samples identified as positive by qPCR (i.e. showing a CNV higher than 2.5-fold) were also identified as positive using the TaqMan assay and no false negative was observed. A good amplification specificity was observed for both the *CCE* gene AAEL023844 and the control gene. A similar PCR efficiency of ~ 95% was observed for both the *CCE* gene AAEL023844 and the control gene leading to a C_q of ~30 cycles for both genes in absence of duplication with 0.5 ng/µL template gDNA (see Supplementary File 1 for a user guideline on this TaqMan assay).

FIGURE 7. Overview of the TaqMan multiplex assay. A: Amplification profiles obtained for a positive individual (red) and a negative individual (blue). Solid line: amplification profile of the target gene AAEL023844 (formerly *CCEAE3A*, FAM probe), Dashed line: amplification profile of the control gene (AAEL007808, HEX probe). B: Comparison of CNV obtained with standard qPCR assay (SybrGreen, dual reactions) and TaqMan assay (FAM/Hex probes, single multiplex reaction). For both methods, CNV were estimated using the ΔΔCt method and are expressed as normalized gDNA quantity relative to the fully susceptible line Bora-Bora. Blue: negative individuals, Red: positive individuals. Each dot type stands for a different population.

Comparing gene copy number estimated from standard qPCR and TaqMan assays revealed a

good correlation between the two techniques (r=0.84, P<0.001) (Figure 7B) although CNV

levels obtained with the TaqMan assay were lower as compared to those obtained with qPCR

and dd qPCR using primers targeting a different fragment.

4 | DISCUSSION

Chemical insecticides remain a key component of integrated strategies aiming at preventing the

transmission of mosquito-borne diseases worldwide but the selection and spread of resistance

threaten their efficacy (Moyes et al., 2017). Controlling resistance by alternating selection

pressures is theoretically possible but requires an efficient monitoring of the dynamics of

resistance alleles in the field (Dusfour et al., 2019). Resistance to organophosphate insecticides

is common in the mosquito *Ae. aegypti* and particularly frequent in SEA following their massive

use for decades (Boyer et al., 2018; Marcombe et al., 2019; Pethuan et al., 2007; Ranson,

Burhani, Lumjuan, & Black IV, 2008; Vontas et al., 2012). Although several *CCE* genes are

known to be involved, the genomic changes underlying resistance are not fully understood. As

a result, no rapid diagnostic assay is available to monitor the frequency of resistance alleles in

the field. By combining experimental evolution and multiple sequencing approaches, we

confirmed the key role of *CCE* amplification in resistance to organophosphate insecticides in

Ae. aegypti and further characterize the associated genomic event. The prevalence of resistant

alleles was then investigated in SEA and a TaqMan multiplex qPCR assay allowing their rapid

detection in single mosquito specimens was developed.

4.1 | *CCE* amplifications play a key role in organophosphate resistance in *Ae. aegypti*

Our experimental selection approach confirmed the presence of organophosphate resistance

alleles in *Ae. aegypti* populations in Laos and their rapid selection with malathion. These results

are consistent with the continuous use of organophosphates for vector control for 30 years in

Laos and the detection of resistance throughout the country (Marcombe et al., 2019). Bioassays

with other insecticides revealed that resistance alleles selected by malathion also confer cross-

resistance to other organophosphates at both larval and adult stage but not to the pyrethroid

deltamethrin suggesting a resistance spectrum restricted to the organophosphate family. This

confirms previous findings suggesting that the over-production of non-specific

carboxylesterases is a common adaptive response to organophosphates in mosquitoes (Cuany

et al., 1993; Hemingway, Hawkes, McCarroll, & Ranson, 2004; Naqqash, Gökçe, Bakhsh, &

Salim, 2016). However, higher resistance levels were observed for malathion and fenitrothion

at the adult stage as compared to temephos at the larval stage possibly due to the differential

expression of these *CCE* genes across life stages (Harker et al., 2013) or a lower specificity

toward temephos.

RNA-seq analysis identified seven detoxification genes over-transcribed in association with

malathion resistance including five consecutive *CCE* genes on chromosome 2, one microsomal

GST on chromosome 1, and one P450 (*CYP6N17*) on chromosome 3. The over-transcription of

this *P450* and this *GST* in two other sister lines selected with insecticides from different families

and not displaying malathion resistance does not support their key role in resistance (data not

shown). Conversely, the over-transcription of *CCE* genes was expected as *CCE*s have often

been associated with organophosphate resistance in mosquitoes. In *Cx pipiens* their

overproduction in response to organophosphate selection is well documented with distinct loci

having spread worldwide (Raymond, Chevillon, Guillemaud, Lenormand, & Pasteur, 1998). In

this species, high resistance levels were associated to the co-occurrence of carboxylesterases

over-production through genomic amplification and the presence of the *ace-1* G119S target-

site mutation affecting the acetylcholinesterase (Raymond et al., 2001). In *Aedes* mosquitoes,

the G119S *ace1* mutation is submitted to a strong genetic constraint (Weill et al., 2004) and has

thus not been reported, suggesting the central role of carboxylesterases over-production in

resistance.

Whole genome sequencing and quantification of gene copy number confirmed the role of

genomic amplifications in the over-production of carboxylesterases associated with

organophosphate resistance in *Ae. aegypti*, as previously suggested (Faucon et al., 2015, 2017;

Poupardin et al., 2014). In addition, our data revealed the co-existence of at least two distinct

structural duplication haplotypes, one including the three *CCE* genes AAEL019678,

AAEL023844 (formerly *CCEAE3A*) and AAEL005113 (haplotype A) and the other one not

including AAEL00113 (haplotype B), the *CCE* gene located at 5' side the *CCE* cluster.

Structural polymorphism of genomic duplications in clustered detoxification genes has been

recently reported in *Anopheles gambiae,* with twelve different alleles identified in a cluster of

*P450*s and eleven in a cluster of *GST*s (Lucas et al., 2019). In the tiger mosquito *Ae. albopictus,*

a structural polymorphism affecting a similar *CCE* cluster duplication was identified with at

least two distinct haplotypes: one including two *CCE* genes and the second one with only the

gene AALF007796, the best orthologue of *Ae. aegypti* AAEL023844 (Grigoraki et al., 2017).

These striking similarities between the two sister species *Ae. albopictus* and *Ae. aegypti*, likely

resulting from a convergent adaptation, further supports the key role of *CCE* amplifications in

the adaptation of *Aedes* mosquitoes to organophosphate insecticides.

The genetic mechanism underlying the amplification of these orthologous *loci* has not been

characterized yet. Previous studies suggested the existence of "hot spots" of recombination

favoring structural polymorphisms (Bass & Field, 2011). In insects the presence of transposable

elements is also known to favor duplication events associated with their rapid adaptation to

insecticides (Bass & Field, 2011; Grigoraki et al., 2017; Schmidt et al., 2010). Our genomic

data confirm the presence of multiple repeated elements in the vicinity of this duplication

though further genomic analyses are required to decipher their relative involvement in this genomic event.

4.2 | Evolutionary dynamics of *CCE* amplifications

The screening of this *CCE* gene amplification by qPCR on field-collected mosquitoes confirmed its occurrence in *Ae. aegypti* populations from SEA. Its prevalence in natural populations was globally low except in Cambodia and in one Thai population from which high organophosphate resistance was previously described (Paeporn et al., 2013; Pethuan et al., 2007; Poupardin et al., 2014; Saelim et al., 2005). Although our sampling campaign was restricted to a few populations in Thailand, Laos and Cambodia, the frequent elevated esterase activities detected in association with temephos resistance in SEA suggests that this *CCE* amplification is widely spread in the region (Paeporn et al., 2013; Pethuan et al., 2007). Previous studies also support the occurrence of this *CCE* amplification in the Caribbean region with high expression levels detected for AAEL023844 from multiple islands and the presence of gene amplification validated in Guadeloupe and Saint-Martin (Goindin et al., 2017; Marcombe et al., 2012, 2009). Although this needs to be confirmed, the frequent association between elevated esterase activities and organophosphate resistance in South-America (i.e. French Guiana, Brazil, Colombia and Costa-Rica) (Bisset et al., 2013; Gambarra et al., 2013; Melo-Santos et al., 2010; Paiva et al., 2016) and New Caledonia (Dusfour et al., 2015), suggests that this *CCE* amplification is distributed worldwide.

Despite the low frequency of *CCE* amplifications in most field populations, our experimental evolution approach demonstrated that the frequency of these resistance alleles increases rapidly in populations submitted to insecticide selection pressure while a susceptible allele is favoured in absence of selection. These findings support the highly beneficial effect of these *CCE* amplifications in the presence of insecticides but also their association with significant fitness

costs in the absence of selective pressure. Fitness costs associated with the over-production of detoxification enzymes have been previously described in various insect species (ffrench-Constant & Bass, 2017; Kliot & Ghanim, 2012). Direct measurement of energetic resources (e.g. lipids, glycogen and glucose) in *Cx. pipiens* mosquitoes over-expressing carboxylesterases suggested that resistant individuals carry up to 30% less energetic reserves than their susceptible counterparts (Rivero, Magaud, Nicot, & Vézilier, 2011). Such high metabolic cost may explain the favored selection of the shorter (less costly) haplotype B as observed in field populations and laboratory lines showing a high *CCE* amplification prevalence. Although further studies are required to quantify the relative importance of the different *CCE* genes included in this genomic amplification in insecticide resistance, the frequent over-expression of AAEL023844 in resistant populations, its inclusion in both amplification haplotypes and its ability to sequester and metabolize temephos (Grigoraki et al., 2016) support its central role in organophosphate resistance.

In addition to structural polymorphism, our study revealed extensive copy number variations between resistant individuals in both field populations and laboratory lines with *CCE* gene copies varying from 3 to ~80 as measured by our TaqMan assay. In addition, no increase gene copy number was observed before and after selection. This suggests that an important copy number polymorphism occurs in natural populations and that individuals carrying high gene copy number are not preferentially selected, probably as a consequence of a trade-off between insecticide survival and metabolic costs associated with the over-production of these enzymes. Such high copy number polymorphism also supports the occurrence of a single duplication event followed by multiple amplification events in *Ae. aegypti* as suggested in *Cx. pipiens* and *Ae. albopictus* (Grigoraki et al., 2017; Guillemaud et al., 1999; Qiao & Raymond, 1995).

Allelic variations of carboxylesterases have also been associated with organophosphate resistance in mosquitoes (Callaghan, Guillemaud, Makate, & Raymond, 1998; Mouchès et al.,

1987). Indeed, both allelic variations and genes duplications coexist in natural populations and can be captured by selection depending on fitness-to-environment relationships (Milesi, Lenormand, Lagneau, Weill, & Labbé, 2016). In *Ae. aegypti*, a few non-synonymous variations affecting the AAEL023844 gene were associated with temephos resistance in a Thai population (Poupardin et al., 2014). However, subsequent functional studies did not support the role of these variations in insecticide sequestration and metabolism (Grigoraki et al., 2016). More recently, we combined controlled crosses with pool-sequencing to segregate organophosphate resistance alleles in a multi-resistant population from French Guiana (Cattel et al., 2019). Such approach identified a strong selection signature associated with organophosphate at this *CCE* locus. Interestingly, several non-synonymous variations affecting *CCE* genes were positively associated with insecticide survival while no CNV were detected, suggesting that the selection of particular variants at this locus may also contribute to resistance. Further work is required to precise the interplay between *CCE* duplications, sequence polymorphism and their respective roles in insecticide resistance in *Ae. aegypti*.

4.3 | A novel TaqMan assay to track organophosphate resistance in *Ae. aegypti*

The present study confirmed the importance of a *CCE* amplifications in insecticide resistance in *Ae. aegypti*, confirming the usefulness of CNV markers for tracking in the field. Our cross-resistance data together with previous findings (Faucon et al., 2015, 2017; Grigoraki et al., 2016; Marcombe et al., 2019; Poupardin et al., 2014) support the routine use of this marker for the monitoring of resistance to various organophosphate insecticides in *Ae. aegypti*. In term of applicability, such CNV marker is highly superior to RNA markers classically used to detect *CCE* genes overexpression because i) genomic DNA can be extracted from dead specimens of any life stage stored at room temperature, ii) either pools or single individuals can be used if

allele frequency data are required and iii) CNV quantification by qPCR is fast, easy, affordable

and data are not affected by insect physiological state.

Genomic duplications can be detected by PCR in two different ways as illustrated in Weetman

et al., 2018. The first one consists of amplifying the junction between two copies by designing

specific primers toward both sides of the duplicated region. Such presence/absence assay is

cheap and low tech but i) does not quantify copy number, ii) requires the precise identification

of duplication breakpoints, which can be impaired by the high density of repeated elements in

flanking regions, and iii) may generate false negatives if duplication breakpoints vary in

position or sequence. The alternative approach adopted herein consists of comparing the copy

numbers between a target gene and a control gene only present as a single copy. Although this

approach is slightly more expensive and requires the use of a qPCR machine, time-to-result is

even shorter (no gel migration required) and results are not affected by structural

polymorphisms provided an appropriate target is defined. Though multiple *CCE* genes are

located within the amplified region, we selected AAEL023844 as target gene because of its

central position in the duplication, its inclusion in both structural haplotypes, its over-expression

in several resistant populations worldwide (Dusfour et al., 2015; Goindin et al., 2017;

Marcombe et al., 2019; Moyes et al., 2017) and its ability to sequester and metabolize temephos

(Grigoraki et al., 2016). By targeting a coding region showing no homology with other genomic

regions, we ensured a good assay specificity with limiting detrimental effects potentially caused

by polymorphisms variations. Though this approach was successfully used for CNV detection

with standard SybrGreen qPCR, it still required performing two distinct qPCR amplifications.

Time-to-results and specificity were then further improved by the development of a dual-color

TaqMan assay allowing the concomitant quantification of both target and control genes. This

assay still proved to be highly specific and allowed reducing time-to-results to ~2h for less than

1.5€ /sample including gDNA extraction, qPCR consumables/reagents, primers and Taqman

probes.

5 | CONCLUSION

While an increasing number of alternatives to chemical insecticides are being developed for

mosquito control (Achee et al., 2019) their optimization and deployment at a worldwide scale

will take at least a decade. Until then, preserving the efficacy of the few insecticides authorized

in public health by managing resistance is crucial to limit the impact of vector-borne diseases.

However, resistance management is often hampered by insufficient resistance monitoring

capacities, often leading to late or inappropriate implementations of management actions. In

this context, the deep comprehension of the genetic bases of resistance and the development of

molecular tools to track resistance alleles in the field still represent a significant capital gain for

public health. By combining experimental selection and deep sequencing, the present study

confirmed the key role of a genomic amplification of a carboxylesterase gene cluster in

organophosphate resistance in the mosquito *Ae. aegypti*. The spatial dynamics of this resistance

locus was investigated in SEA and a novel TaqMan assay was developed enabling its high-

throughput monitoring in field mosquito populations. The routine use of this assay in SEA, and

possibly in other tropical areas, should improve the monitoring of organophosphate resistance

in the arbovirus vector *Ae. aegypti*. From an evolutionary perspective, deciphering the

evolutionary history and the genetic events underlying the emergence and spread of this recent

adaptation undoubtedly deserves further attention.

ACKNOWLEDGEMENTS

732 The views expressed in this publication are those of the authors and do not necessarily reflect

733 the official policy or position of the Department of the Navy, Department of Defense, nor the

U.S. Government. This work was conducted in the framework of the U.S. Naval Medical Research Unit TWO projects BIO-LAO-2 (work unit number D1425) and ARBOVEC-PLUS (work unit number D1428), in support and funded by the Department of Defense Global Emerging Infections Surveillance Program and Military Infectious Disease Research Program. I (IWS and JCH) am a military Service member. This work was prepared as part of my official duties. Title 17, U.S.C., §105 provides that copyright protection under this title is not available for any work of the U.S. Government. Title 17, U.S.C., §101 defines a U.S. Government work as a work prepared by a military Service member or employee of the U.S. Government as part of that person's official duties. This publication was also supported by the project, Research Infrastructures for the control of vector-borne diseases (Infravec2), which has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 731060. Dr. Julien Cattel was supported by funding from the European Union's Horizon 2020 Research and Innovation Programme under ZIKAlliance Grant Agreement no. 734548. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The field study in Cambodia was supported by ECOMORE 2 project, coordinated by Institut Pasteur and financially supported by AFD (Agence Française pour le Développement). We thank Khaithong Lakeomany, Nothasin Phommavan, Somsanith Chonephetsarath, Somphat Nilaxay, and Phoutmany Thammavong for mosquito collections and rearing from Laos. Finally, we thank Dr. Pablo Tortosa for a critical reading of this manuscript.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ETHICAL APPROVAL

Blood feeding of adult mosquitoes was performed on mice. Mice were maintained in the animal house of the federative structure Environmental and Systems Biology (BEeSy) of Grenoble-Alpes University agreed by the French Ministry of animal welfare (agreement n° B 38 421 10 001) and used in accordance to European Union laws (directive 2010/63/UE). The use of animals for this study was approved by the ethic committee ComEth Grenoble-C2EA-12 mandated by the French Ministry of higher Education and Research (MENESR).

DATA AVAILABILITY STATEMENT

The sequence data from this study have been deposited to the European Nucleotide Archive (ENA; http://www.ebi.ac.uk/ena) under the accession numbers PRJEB37991 (RNA-seq data) and PRJEB37993 (whole genome pool-seq data).

REFERENCES

Achee, N. L., Grieco, J. P., Vatandoost, H., Seixas, G., Pinto, J., Ching-Ng, L., … Vontas, J. (2019). Alternative strategies for mosquito-borne arbovirus control. *PLoS Neglected Tropical Diseases*, *13*(1), 1–22. https://doi.org/10.1371/journal.pntd.0006822

Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count data. *Nature Precedings*, 1.

Assogba, B. S., Djogbénou, L. S., Milesi, P., Berthomieu, A., Perez, J., Ayala, D., … Weill, M. (2015). An ace-1 gene duplication resorbs the fitness cost associated with resistance in Anopheles gambiae, the main malaria mosquito. *Scientific Reports*, *5*(May), 19–21. https://doi.org/10.1038/srep14529

Bass, C., & Field, L. M. (2011). Gene amplification and insecticide resistance. *Pest Management Science*, *67*(8), 886–890. https://doi.org/10.1002/ps.2189

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., … Bolker, M. Ben. (2015). Package "lme4." *Convergence*, *12*(1), 2.

Benjamini, Y., & Hochberg, Y. (1995). Benjamini and Y FDR.pdf. *Journal of the Royal Statistical Society. Series B (Methodological)*, Vol. 57, pp. 289–300. https://doi.org/10.2307/2346101

Bisset, J. A., Marín, R., Rodríguez, M. M., Severson, D. W., Ricardo, Y., French, L., … Pérez, O. (2013). Insecticide Resistance in Two Aedes aegypti (Diptera: Culicidae) Strains From Costa Rica. *Journal of Medical Entomology*, *50*(2), 352–361. https://doi.org/10.1603/me12064

Boyer, S., Lopes, S., Prasetyo, D., Hustedt, J., Sarady, A. S., Doum, D., … Hii, J. (2018).
Resistance of Aedes aegypti (Diptera: Culicidae) Populations to Deltamethrin,
Permethrin, and Temephos in Cambodia. *Asia-Pacific Journal of Public Health*, *30*(2),
158–166. https://doi.org/10.1177/1010539517753876
Callaghan, A., Guillemaud, T., Makate, N., & Raymond, M. (1998). Polymorphisms and
fluctuations in copy number of amplified esterase genes in Culex pipiens mosquitoes.
Insect Molecular Biology, *7*(3), 295–300.
Campbell, C. D., & Eichler, E. E. (2013). Properties and rates of germline mutations in
humans. *Trends in Genetics*, *29*(10), 575–584.
Cattel, J., Faucon, F., Le Péron, B., Sherpa, S., Monchal, M., Grillet, L., … David, J. P.
(2019). Combining genetic crosses and pool targeted DNA-seq for untangling genomic
variations associated with resistance to multiple insecticides in the mosquito Aedes
aegypti. *Evolutionary Applications*, (April), 1–15. https://doi.org/10.1111/eva.12867
Chatonnet, A., Lenfant, N., Marchot, P., & Selkirk, M. E. (2017). Natural genomic
amplification of cholinesterase genes in animals. *Journal of Neurochemistry*, *142*, 73–81.
https://doi.org/10.1111/jnc.13990
Collins, F., Drumm, M. L., Cole, J. L., Lockwood, W. K., Woude, G. F. Vande, & Iannuzzi,
M. C. (1987). Construction of a general human chromosome jumping library, with
application to cystic fibrosis. *Science*, *235*, 1046–1050.
Cooper, G. M., Nickerson, D. A., & Eichler, E. E. (2007). Mutational and selective effects on
copy-number variants in the human genome. *Nature Genetics*, *39*(7s), S22.
Cuany, A., Handani, J., Bergé, J., Fournier, D., Raymond, M., Georghiou, G. P., & Pasteur,
N. (1993). Action of esterase b1 on chlorpyrifos in organophosphate-resistant culex
mosquitos. *Pesticide Biochemistry and Physiology*, Vol. 45, pp. 1–6.
https://doi.org/10.1006/pest.1993.1001
Dusfour, I., Vontas, J., David, J. P., Weetman, D., Fonseca, D. M., Corbel, V., … Chandre, F.
(2019). Management of insecticide resistance in the major Aedes vectors of arboviruses:
Advances and challenges. *PLoS Neglected Tropical Diseases*, Vol. 13, pp. 1–22.
https://doi.org/10.1371/journal.pntd.0007615
Dusfour, I., Zorrilla, P., Guidez, A., Issaly, J., Girod, R., Guillaumot, L., … Strode, C. (2015).
Deltamethrin Resistance Mechanisms in Aedes aegypti Populations from Three French
Overseas Territories Worldwide. *PLoS Neglected Tropical Diseases*, *9*(11), 1–17.
https://doi.org/10.1371/journal.pntd.0004226
Faucon, F., Dusfour, I., Gaude, T., Navratil, V., Boyer, F., Chandre, F., … David, J. (2015).
Unravelling genomic changes associated with insecticide resistance in the dengue
mosquito Aedes aegypti by deep targeted se. *Genome Research*, (August), 1347–1359.
https://doi.org/10.1101/gr.189225.115
Faucon, F., Gaude, T., Dusfour, I., Navratil, V., Corbel, V., Juntarajumnong, W., … David,
J.-P. (2017). In the hunt for genomic markers of metabolic resistance to pyrethroids in
the mosquito Aedes aegypti: An integrated next-generation sequencing approach. *PLOS*
Neglected Tropical Diseases, *11*(4), e0005526.
https://doi.org/10.1371/journal.pntd.0005526
ffrench-Constant, R. H., & Bass, C. (2017). Does resistance really carry a fitness cost?
Current Opinion in Insect Science, *21*, 39–46. https://doi.org/10.1016/j.cois.2017.04.011
Gambarra, W. P. T., Martins, W. F. S., Lucena Filho, M. L. de, Albuquerque, I. M. C. de,
Apolinário, O. K. dos S., & Beserra, E. B. (2013). Spatial distribution and esterase
activity in populations of Aedes (Stegomyia) aegypti (Linnaeus) (Diptera: Culicidae)
resistant to temephos. *Revista Da Sociedade Brasileira de Medicina Tropical*, *46*(2),
178–184.
Goindin, D., Delannay, C., Gelasse, A., Ramdini, C., Gaude, T., Faucon, F., … Fouque, F.

(2017). Levels of insecticide resistance to deltamethrin, malathion, and temephos, and associated mechanisms in Aedes aegypti mosquitoes from the Guadeloupe and Saint Martin islands (French West Indies). *Infectious Diseases of Poverty*, *6*(1), 38. https://doi.org/10.1186/s40249-017-0254-x

Grigoraki, L., Balabanidou, V., Meristoudis, C., Miridakis, A., Ranson, H., Swevers, L., & Vontas, J. (2016). Functional and immunohistochemical characterization of CCEae3a, a carboxylesterase associated with temephos resistance in the major arbovirus vectors Aedes aegypti and Ae. albopictus. *Insect Biochemistry and Molecular Biology*, *74*, 61–67. https://doi.org/10.1016/j.ibmb.2016.05.007

Grigoraki, L., Pipini, D., Labbé, P., Chaskopoulou, A., Weill, M., & Vontas, J. (2017). Carboxylesterase gene amplifications associated with insecticide resistance in Aedes albopictus: Geographical distribution and evolutionary origin. *PLoS Neglected Tropical Diseases*, *11*(4), 1–13. https://doi.org/10.1371/journal.pntd.0005533

Guillemaud, T., Raymond, M., Tsagkarakou, A., Bernard, C., Rochard, P., & Pasteur, N. (1999). Quantitative variation and selection of esterase gene amplification in Culex pipiens. *Heredity*, *83*(1), 87–99. https://doi.org/10.1038/sj.hdy.6885370

Harker, B. W., Behura, S. K., Debruyn, B. S., Lovin, D. D., Mori, A., Romero-Severson, J., & Severson, D. W. (2013). Stage-specific transcription during development of Aedes aegypti. *BMC Developmental Biology*, *13*(1). https://doi.org/10.1186/1471-213X-13-29

Hemingway, J., Hawkes, N. J., McCarroll, L., & Ranson, H. (2004). The molecular basis of insecticide resistance in mosquitoes. *Insect Biochemistry and Molecular Biology*, *34*(7), 653–665. https://doi.org/10.1016/j.ibmb.2004.03.018

Iskow, R. C., Gokcumen, O., & Lee, C. (2012). Exploring the role of copy number variants in human adaptation. *Trends in Genetics*, *28*(6), 245–257.

Kliot, A., & Ghanim, M. (2012). Fitness costs associated with insecticide resistance. *Pest Management Science*, *68*(11), 1431–1437. https://doi.org/10.1002/ps.3395

Kondrashov, F. A. (2012). Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1749), 5048–5057. https://doi.org/10.1098/rspb.2012.1108

Labbé, P., Berthomieu, A., Berticat, C., Alout, H., Raymond, M., Lenormand, T., & Weill, M. (2007). Independent duplications of the acetylcholinesterase gene conferring insecticide resistance in the mosquito Culex pipiens. *Molecular Biology and Evolution*, *24*(4), 1056–1067. https://doi.org/10.1093/molbev/msm025

Lucas, E. R., Miles, A., Harding, N. J., Clarkson, C. S., Lawniczak, M. K. N., Kwiatkowski, D. P., … Donnelly, M. J. (2019). Whole-genome sequencing reveals high complexity of copy number variation at insecticide resistance loci in malaria mosquitoes. *Genome Research*, *29*(8), 1250–1261. https://doi.org/10.1101/gr.245795.118

Marcombe, S., Fustec, B., Cattel, J., Chonephetsarath, S., Thammavong, P., Phommavanh, N., … Brey, P. T. (2019). Distribution of insecticide resistance and mechanisms involved in the arbovirus vector Aedes aegypti in Laos and implication for vector control. *PLoS Neglected Tropical Diseases*, *13*(12), e0007852. https://doi.org/10.1371/journal.pntd.0007852

Marcombe, S., Mathieu, R. B., Pocquet, N., Riaz, M. A., Poupardin, R., Sélior, S., … Chandre, F. (2012). Insecticide resistance in the dengue vector aedes aegypti from martinique: Distribution, mechanisms and relations with environmental factors. *PLoS ONE*, *7*(2). https://doi.org/10.1371/journal.pone.0030989

Marcombe, S., Poupardin, R., Darriet, F., Reynaud, S., Bonnet, J., Strode, C., … David, J.-P. (2009). Exploring the molecular basis of insecticide resistance in the dengue vector Aedes aegypti: a case study in Martinique Island (French West Indies). *BMC Genomics*, *10*(1), 494. https://doi.org/10.1186/1471-2164-10-494

Melo-Santos, M. A. V., Varjal-Melo, J. J. M., Araújo, A. P., Gomes, T. C. S., Paiva, M. H. S., Regis, L. N., … Ayres, C. F. J. (2010). Resistance to the organophosphate temephos: Mechanisms, evolution and reversion in an Aedes aegypti laboratory strain from Brazil. *Acta Tropica*, *113*(2), 180–189. https://doi.org/10.1016/j.actatropica.2009.10.015

Milesi, P., Lenormand, T., Lagneau, C., Weill, M., & Labbé, P. (2016). Relating fitness to long-term environmental variations in natura. *Molecular Ecology*, *25*(21), 5483–5499.

Mouchès, C., Magnin, M., Bergé, J. B., de Silvestri, M., Beyssat, V., Pasteur, N., & Georghiou, G. P. (1987). Overproduction of detoxifying esterases in organophosphate-resistant Culex mosquitoes and their presence in other insects. *Proceedings of the National Academy of Sciences of the United States of America*, *84*(8), 2113–2116. https://doi.org/10.1073/pnas.84.8.2113

Moyes, C. L., Vontas, J., Martins, A. J., Ng, L. C., Koou, S. Y., Dusfour, I., … Weetman, D. (2017). Contemporary status of insecticide resistance in the major Aedes vectors of arboviruses infecting humans. *PLoS Neglected Tropical Diseases*, *11*(7), 1–20. https://doi.org/10.1371/journal.pntd.0005625

Naqqash, M. N., Gökçe, A., Bakhsh, A., & Salim, M. (2016). Insecticide resistance and its molecular basis in urban insect pests. *Parasitology Research*, *115*(4), 1363–1373. https://doi.org/10.1007/s00436-015-4898-9

Paeporn, P., Komalamisra, N., Deesin, V., Rongsriyam, Y., Eshita, Y., Thongrungkiat, S., … Medical, O. (2013). Temephos Resistance in Two Forms of Aedes Aegypti and Its Significance for the Resistance. *Southeast Asian Journal of Tropical Medicine and Public Health.*, *34*(4).

Paiva, M. H. S., Lovin, D. D., Mori, A., Melo-Santos, M. A. V., Severson, D. W., & Ayres, C. F. J. (2016). Identification of a major Quantitative Trait Locus determining resistance to the organophosphate temephos in the dengue vector mosquito Aedes aegypti. *Genomics*, *107*(1), 40–48. https://doi.org/10.1016/j.ygeno.2015.11.004

Pethuan, S., Jirakanjanakit, N., Saengtharatip, S., Chareonviriyaphap, T., Kaewpa, D., & Rongnoparut, P. (2007). Biochemical studies of insecticide resistance in Aedes (Stegomyia) aegypti and Aedes (Stegomyia) albopictus (Diptera: Culicidae) in Thailand. *Tropical Biomedicine*, *24*(1), 7–15.

Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Research*, *29*(9), e45. https://doi.org/10.1093/nar/29.9.e45

Poupardin, R., Srisukontarat, W., Yunta, C., & Ranson, H. (2014). Identification of Carboxylesterase Genes Implicated in Temephos Resistance in the Dengue Vector Aedes aegypti. *PLoS Neglected Tropical Diseases*, *8*(3). https://doi.org/10.1371/journal.pntd.0002743

Qiao, C. L., & Raymond, M. (1995). The same esterase B1 haplotype is amplified in insecticide-resistant mosquitoes of the culex pipiens complex from the americas and china. *Heredity*, *74*(4), 339–345. https://doi.org/10.1038/hdy.1995.51

R Core Team. (2013). *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2013*. ISBN 3-900051-07-0.

Ranson, H., Burhani, J., Lumjuan, N., & Black IV, W. C. (2008). Insecticide resistance in dengue vectors. *TropIKA*, 1–12. Retrieved from http://journal.tropika.net

Raymond, M., Berticat, C., Weill, M., Pasteur, N., & Chevillon, C. (2001). Insecticide resistance in the mosquito *Culex pipiens*: what have we learned about adaptation? *Genetica*, *112–113*, 287–296. https://doi.org/10.1023/A:1013300108134

Raymond, M., Chevillon, C., Guillemaud, T., Lenormand, T., & Pasteur, N. (1998). An overview of the evolution of overproduced esterases in the mosquito Culex pipiens. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *353*(1376), 1707–1711. https://doi.org/10.1098/rstb.1998.0322

Rivero, A., Magaud, A., Nicot, A., & Vézilier, J. (2011). Energetic Cost of Insecticide
Resistance in Culex pipiens Mosquitoes. *Journal of Medical Entomology*, *48*(3), 694–
700. https://doi.org/10.1603/me10121
Rozen, S., & Skaletsky, H. (2000). Primer3 on the WWW for general users and for biologist
programmers. *Methods in Molecular Biology (Clifton, N.J.)*, *132*(August), 365–386.
https://doi.org/10.1385/1-59259-192-2:365
Saelim, V., Brogdon, W. G., Rojanapremsuk, J., Suvannadabba, S., Pandii, W., Jones, J. W.,
& Sithiprasasna, R. (2005). Bottle and biochemical assays on temephos resistance in
Aedes aegypti in Thailand. *Southeast Asian Journal of Tropical Medicine and Public
Health*, *36*(2), 417–425.
Schmidt, J. M., Good, R. T., Appleton, B., Sherrard, J., Raymant, G. C., Bogwitz, M. R., …
Robin, C. (2010). Copy number variation and transposable elements feature in recent,
ongoing adaptation at the Cyp6g1 locus. *PLoS Genetics*, *6*(6), 1–11.
https://doi.org/10.1371/journal.pgen.1000998
Vontas, J., Kioulos, E., Pavlidi, N., Morou, E., della Torre, A., & Ranson, H. (2012).
Insecticide resistance in the major dengue vectors Aedes albopictus and Aedes aegypti.
Pesticide Biochemistry and Physiology, *104*(2), 126–131.
https://doi.org/10.1016/j.pestbp.2012.05.008
Weetman, D., Djogbenou, L. S., & Lucas, E. (2018). Copy number variation (CNV) and
insecticide resistance in mosquitoes: evolving knowledge or an evolving problem?
Current Opinion in Insect Science, *27*, 82–88. https://doi.org/10.1016/j.cois.2018.04.005
Weill, M., Berthomieu, A., Berticat, C., Lutfalla, G., Nègre, V., Pasteur, N., … Raymond, M.
(2004). Insecticide resistance: a silent base. *Current Biology*, *14*(14), 552–553.
Zhang, J. (2003). Evolution by gene duplication : an update. *Trends in Ecology and Evolution*,
18(6), 292–298. https://doi.org/10.1016/S0169-5347(03)00033-8
968