FasR regulates fatty acid biosynthesis and is essential for virulence of

Mycobacterium tuberculosis

Sonia Mondino[a,*], Cristina Lourdes Vázquez[b], Matías Cabruja[a,*], Claudia Sala[c,*],

Amaury Cazenave-Gassiot[d], Federico Carlos Blanco[b], Markus R. Wenk[d], Fabiana

Bigi[b], Stewart T. Cole[c,*], Hugo Gramajo[a] and Gabriela Gago[a#]

[a] Laboratory of Physiology and Genetics of Actinomycetes, Instituto de Biología
Molecular y Celular de Rosario (IBR-CONICET), Facultad de Ciencias Bioquímicas
y Farmacéuticas, Universidad Nacional de Rosario, Argentina
[b] Instituto de Biotecnología-IABIMO, UEDD CONICET-Instituto Nacional de
Tecnología Agropecuaria (INTA), Argentina.
[c] Global Health Institute, Ecole Polytechnique Fédérale de Lausanne, Lausanne,
Switzerland.
[d] Singapore Lipidomics Incubator (SLING), Life Sciences Institute, and Department
of Biochemistry, Yong Loo Lin School of Medicine, National University of
Singapore, Singapore.
17

Running Head: FasR is essential for *M. tuberculosis* virulence

Address correspondence to Gabriela Gago, gago@ibr-conicet.gov.ar

[*] Sonia Mondino and Stewart T. Cole present address: Institut Pasteur, Paris
(France)

[*] Matías Cabruja present address: Stanford University (United States)

[*] Claudia Sala present address: Fondazione Toscana Life Sciences, Siena (Italy)

Sonia Mondino, Cristina L. Vázquez and Matías Cabruja contributed equally to this
work. Author order was determined on the basis of contributions.

The abstract has 247 words and the text has 5215 words (including Materials and
Methods).

Abstract

Mycobacterium tuberculosis, the etiologic agent of human tuberculosis, is the world's leading cause of death from an infectious disease. One of the main features of this pathogen is the complex and dynamic lipid composition of the cell envelope, which adapts to the variable host environment and defines the fate of infection by actively interacting with and modulating immune responses. However, while much has been learned about the enzymes of the numerous lipid pathways, little knowledge is available regarding the proteins and metabolic signals regulating lipid metabolism during *M. tuberculosis* infection. Previous studies identified FasR as an essential transcriptional activator of *de novo* fatty acid biosynthesis in *Mycobacterium smegmatis*. In this work, we constructed and characterized a FasR-deficient mutant in *M. tuberculosis* and demonstrated that FasR positively regulates *fas* and *acpS* expression. Lipidomic analysis of the wild type and mutant strains revealed complete rearrangement of most lipid components of the cell envelope, with phospholipids, mycolic acids, sulfolipids and phthiocerol dimycocerosates relative abundance severely altered. As a consequence, replication of the mutant strain was impaired in macrophages leading to reduced virulence in a mouse model of infection. Moreover, we show that the *fasR* mutant resides in acidified cellular compartments, suggesting that the lipid perturbation caused by the mutation prevented *M. tuberculosis* inhibition of phagolysosome maturation. This study identified FasR as a novel factor involved in regulation of

mycobacterial virulence and provides evidence for the essential role that modulation of lipid homeostasis plays in the outcome of *M. tuberculosis* infection.

Importance

Tuberculosis is the leading cause of death from a single infectious disease, resulting in approximately 1.5 million deaths per year worldwide. The composition and complexity of the cell envelope is the most distinctive feature of *Mycobacterium tuberculosis*, the etiologic agent of human tuberculosis. As the mycobacterial cell envelope is the interface with the host, the modulation of its structure and consequent responsiveness to immune factors might be important in determining the state of an infection. In this study, a *M. tuberculosis* mutant strain in FasR was constructed and analyzed. Inactivation of *fasR* in *M. tuberculosis* was not lethal for *in vitro* growth but led to rearrangement of the lipid composition of the cell envelope. Consequently, the mutant strain was less virulent in a mouse model of infection and showed impaired replication in macrophages, thus implying that lipid homeostasis plays an important role during *M. tuberculosis* infection.

Introduction

Tuberculosis (TB) is one of the top ten causes of death and the leading cause from a single infectious agent, exceeding both malaria and HIV. In 2018, there were an estimated 10 million new cases of active TB and 1.3 million patients died from active infection (1). The emergence of multidrug-resistant TB (MDR-TB) presents

an increasingly difficult therapeutic challenge; consequently, MDR-TB is now the

main cause of death due to antimicrobial resistance (1) thus transforming this long

neglected disease into a global health priority.

The composition and complexity of the cell envelope is the most distinctive feature

of *Mycobacterium tuberculosis*, the etiologic agent of human TB. Biochemical and

microscopic data revealed that the mycobacterial envelope consists of three

discrete entities: a plasma membrane surrounded by a complex cell wall and an

outermost layer called the capsule in pathogenic species (2, 3). The *M.*

tuberculosis cell wall consists of peptidoglycan covalently linked to arabinogalactan

esterified by mycolic acids (MA), which form the inner leaflet of the outer

membrane bilayer (mycomembrane). The outer leaflet of this membrane is

composed of free, non-covalently bound lipids and glycolipids like trehalose

monomycolates (TMM), trehalose dimycolates (TDM), glycerol monomycolates,

glucose monomycolates, phthiocerol dimycocerosates (PDIM), poly acylated

trehaloses (PAT), sulfolipids (SL), phosphatidylinositol mannosides (PIM), phenolic

glycolipids (PGL) and mannose-capped lipoarabinomannans (Man-Lam) (4, 5).

During the course of infection, intracellular pathogenic bacteria use different

strategies to survive, multiply and persist inside the host (6). *M. tuberculosis*

encounters heterogeneous host environments, including high oxygen tension in the

lung alveolus, reactive oxygen and nitrogen species in activated macrophages, as

well as nutrient depletion and hypoxia in tuberculous granulomas. How *M.*

tuberculosis survives these diverse environments is not fully understood. *M.*

tuberculosis has evolved a wide array of specific lipids and related metabolic

pathways that actively interact with the immune system, and multiple reports from

human and animal studies suggest an intimate link between *M. tuberculosis* lipid

metabolism and outcome of infection (7–9). Of particular interest is the challenge

that *M. tuberculosis* must face in order to adjust its own lipid metabolism while the

macrophage is changing its lipid content (10). Little is known about the mechanism

underlying the reorganization of *M. tuberculosis* envelope during infection (11).

Understanding how this pathogen selects and organizes the envelope components

to resist killing, and how these components make the envelope essential to *M.*

tuberculosis survival, virulence and pathogenesis, is still a major challenge.

Therefore, characterization of the regulatory elements that control these processes

should help improve understanding of the mechanisms involved in mycobacterial

pathogenicity and could provide new potential targets for anti-TB chemotherapy.

Previous studies on the regulation of lipid biosynthesis in mycobacteria allowed us

to identify FasR as the transcriptional activator of fatty acid biosynthesis in *M.*

smegmatis (12). Deletion of *fasR* from *M. smegmatis* was possible only in the

presence of an extra copy of the gene, suggesting that *fasR* is essential for

viability. Notably, we found that the affinity of FasR for its target promoter is

modulated by long chain acyl-CoAs ($\geq C_{16}$), the end products of the FAS I pathway.

FasR, together with MabR, the transcriptional regulator of MA biosynthesis, are the

only transcription factors involved in fatty acid biosynthesis regulation known to be

essential for *M. smegmatis* survival, reflecting the relevance of lipid homeostasis

for mycobacterial viability (12, 13).

In this study, a *M. tuberculosis* knockout strain in *fasR* was constructed and the

mutant used to analyze the implications of lipid homeostasis in *M. tuberculosis*

virulence. Inactivation of *fasR* in *M. tuberculosis* was not lethal for *in vitro* growth

but led to complete rearrangement of lipid composition of the envelope.

Consequently, the mutant strain was less virulent in a mouse model and showed

impaired replication in macrophages, thus implying that lipid homeostasis plays an

important role during *M. tuberculosis* infection.

Results

Construction and characterization of *M. tuberculosis fasR* mutant strain

To investigate the function of FasR in *M. tuberculosis,* and considering that *fasR* is

essential for *M. smegmatis* viability, we first constructed an *M. tuberculosis*

conditional knockdown strain in *fasR* (Fig 1A-B). Plasmids pFR33, providing a

second copy of *fasR* under the control of the TET-PIP OFF system at the

chromosomal L5 phage attachment site *attB* (14) and pGA80, were simultaneously

transformed in *M. tuberculosis* (Table S1)*.* Disruption of *fasR* was then obtained by

two-step allelic exchange using pFR35, which deleted the gene at the native locus.

Homologous recombination was confirmed by colony PCR and Southern blot

analysis, and the resulting conditional knockdown strain was named *MtbΔfasR*-

c*fasR* (Fig 1D). In this strain, expression of *fasR* was downregulated by addition of

anhydrotetracycline (ATc) to the culture medium, thus leading to protein depletion.

However, we were not able to detect any growth defect after addition of ATc,

suggesting that *fasR* was not essential for *M. tuberculosis* survival. This result

prompted us to try to obtain a *fasR* knockout (KO) mutant strain by replacing the

fasR complementing plasmid with the empty vector pND255 (Fig 1C). Consistent

with the ability of the conditional mutant to grow in the non-permissive condition,

plenty of colonies were obtained in the presence of Hyg, and the replacement was

successfully confirmed by colony PCR (Fig S1A). We selected two Hyg-resistant

clones and confirmed the depletion of FasR by Western blot using total protein

extracts (Fig S1B). The genotype of this strain was further analyzed by Southern

Blot (Fig 1D), thus validating the genomic deletion of *fasR* and the generation of

the *fasR* KO mutant strain, named *MtbΔfasR*.

Once the non-essentiality of *fasR* for *M. tuberculosis* survival was confirmed, we

then hypothesized that deletion of this transcriptional regulator could be deleterious

for bacterial growth under specific conditions. In particular, we analyzed the growth

of *MtbΔfasR* mutant using different culture media. As shown in Fig 2, growth of

MtbΔfasR in complete 7H9 medium (7H9 + OADC) was similar to that of the wild

type and the complemented strain (*MtbΔfasR-cfasR* grown without ATc), although

the KO mutant reached stationary phase earlier and at lower OD_{600}. Interestingly, if

the medium was not supplemented with oleate and BSA (i.e. 7H9 without OADC),

growth defects appeared to be more pronounced, suggesting that the presence of

lipids in complete 7H9 medium could partially rescue the absence of FasR.

Moreover, *MtbΔfasR* mutant showed severe growth defects in Sauton minimal

medium, reaching a maximal OD_{600} of 0.3 (Fig 2). Altogether, these experiments

showed that FasR is not essential for *M. tuberculosis* survival, although it is

required for optimal bacterial growth, especially under lipid-limited conditions.

FasR is a transcriptional activator of the *fas-acpS* operon in *M. tuberculosis*

We have previously demonstrated that FasR is a transcriptional activator of *fas* in

M. smegmatis (12). Thus, to validate the role of FasR in fatty acid biosynthesis in

M. tuberculosis, we analyzed expression of the *fas-acpS* operon genes in

MtbΔfasR and in the complemented strain *MtbΔfasR-cfasR*. The relative amounts

of *fas* and *acpS* mRNAs were measured by quantitative RT-PCR after 8 days of

growth in complete 7H9 medium and compared to those of *Mtb*H37Rv wild type

strain. As shown in Fig 3, transcription of *fas* and *acpS* was reduced by 70 % and

40 % respectively when cells were deprived of FasR, confirming that FasR is a

transcriptional activator of the *fas-acpS* operon in *M. tuberculosis*.

fasR knockout strongly impacts composition of *M. tuberculosis* cell envelope

Previous characterization of *M. smegmatis fasR* (12) and *fas* (15) conditional

mutant strains revealed that reduced FAS I levels had a strong impact on fatty acid

and phospholipid biosynthesis. Interestingly, MA remained being synthesized in the

FAS I mutant for several hours after addition of ATc (non-permissive conditions),

although with different abundance of the various species (15). Considering the

relevance that lipids have in *M. tuberculosis* survival and virulence, the impact of

the absence of FasR on the global lipid profile of *M. tuberculosis* was further

studied. We performed comparative lipidomic analyses between *Mtb*H37Rv,

MtbΔfasR and the complemented strain. Briefly, we extracted cell-associated lipids

as previously described (15) and analyzed the lipid content by high-performance

liquid chromatography-mass spectrometry with data-dependent tandem MS

(HPLC-MS/MS).

As illustrated in Fig 4, lipid analysis in the negative mode indicated that the relative

abundance (expressed as percentage of lipid MS signals) of phospholipids (PL)

and MA in the wild type and complemented strains was approximately 75 and 25

%, respectively. These values changed to 50 % for both PL and MA in the *fasR*

mutant strain, reflecting the strong impact that lower levels of *de novo* fatty acid

biosynthesis had in PL and MA relative abundance. The relative composition of PL

also showed a biased effect towards phosphatidylinositol (PI) and cardiolipins (CL),

which were significantly reduced in the mutant strain (Figs 4 and S2). Furthermore,

there was a strong reduction in the content of sulfolipids (Figs 4 and S2) and the

acyl chain substitutions of this lipid were consistently shorter in the mutant strain

(Fig S3). A relevant change observed in the qualitative composition of the PL was

the significant increase in the double bonds of the acyl residues in the mutant

strain, with those of the CL, phosphatidylglycerol (PG) and PI being most

significant (Fig S4).

Additionally, results obtained in the positive mode analysis showed that *M.*

tuberculosis Mtb∆fasR mutant strain displayed a significant reduction in the relative

content of PDIM (Figs 5 and S2), with acyl substitutions that are shorter in the

mutant strain, compared with the wild type or the complemented strain (Fig S3).

Most of the changes observed in the deletion mutant were reversed in the

complemented strain *Mtb∆fasR*-c*fasR*, indicating that *fasR* is needed to maintain

the global lipid composition of *M. tuberculosis* cell envelope.

FasR is required for *M. tuberculosis* replication in macrophages

To investigate whether FasR is required for the intracellular survival of *M. tuberculosis* in macrophages, we infected THP-1-derived macrophages with exponentially growing cultures of *Mtb*H37Rv, *MtbΔfasR* and the complemented strain *MtbΔfasR*-c*fasR*. To quantify bacterial uptake and intracellular replication, we determined the numbers of intracellular viable bacteria at different time points post-infection. As shown in Fig 6, infection with similar numbers of bacteria of each strain resulted in recovery of similar numbers of CFU for all of the strains at 24 and 48 h post-infection. However, 120 and 144 h after the infection, intracellular bacterial replication of *MtbΔfasR* strain was significantly reduced compared to that of the wild type strain. The wild type phenotype was partially restored in the *MtbΔfasR*-c*fasR* complemented strain at day 5 (120 h) and completely restored at day 6 (144 h), indicating that the intracellular growth defect was due to the deletion of *fasR* and that this regulator is necessary for optimal intracellular multiplication of *M. tuberculosis* in macrophages.

FasR regulates phagosome maturation in macrophages

We have shown that the absence of FasR, and consequently reduced levels of *fas-acpS*, have a strong impact on the composition of *M. tuberculosis* cell envelope, and also on bacterial multiplication within macrophages. We hypothesized that the altered bacterial cell envelope of the *MtbΔfasR* mutant may affect *M. tuberculosis*-driven phagosome maturation during infection. To investigate this, we analyzed the acidification of *M. tuberculosis MtbΔfasR* phagosome during infection of human macrophages by staining with Lysotracker dye and measuring the phagosomal acquisition of LAMP-3 (late endosomal/lysosomal markers). Interestingly, while

*Mtb*H37Rv prevented phagosome acidification, *Mtb*Δ*fasR* mutant strain resided in

phagosomes that were strongly stained by Lysotracker dye and that acquired

LAMP-3, suggesting that this strain cannot avoid phagosomal maturation. The

observed mutant phenotype was rescued in the *Mtb*Δ*fasR*-c*fasR* complemented

strain, indicating that the increased phagosome acidification was due to the

absence of FasR (Fig 7).

Disruption of lipid homeostasis attenuates mycobacterial virulence in mice

To characterize the role of FasR in an experimental model of infection, three

groups of BALB/c mice (10 per group) were infected intratracheally with 10^2 CFU of

*Mtb*H37Rv, *Mtb*Δ*fasR* and *Mtb*Δ*fasR*-c*fasR* strains. Mice were killed 60 days post-

infection and bacterial loads in lungs were determined. CFU counts were

significantly reduced for the *Mtb*Δ*fasR* strain compared to *Mtb*H37Rv, while the

wild type phenotype was restored in the *Mtb*Δ*fasR*-c*fasR* strain (Fig 8), indicating

that FasR is essential for survival and multiplication of *M. tuberculosis* in mice.

Discussion

In a previous investigation we identified FasR, a TetR-like transcriptional regulator

that specifically binds to the *fas* promoter region to activate its expression, and

found that FasR was essential for *in vitro* growth of *M. smegmatis* (12). However,

its role in *M. tuberculosis* viability was controversial. Using Himar 1 transposon

mutagenesis it was initially proposed that FasR was essential for *M. tuberculosis*

survival (16, 17). However, more recently, De Jesus *et al* found that FasR was

dispensable for growth, by assessing the composition of a highly saturated library

of transposon mutants *of M. tuberculosis* by deep sequencing (18). Moreover, the

relevance of FasR for mycobacterial survival in intracellular environments or animal

models had not been investigated. In this work, by constructing a deletion *fasR*

mutant and a complemented strain, we unequivocally established that FasR is not

essential for *M. tuberculosis* viability, although it is required for optimal growth of *M.*

tuberculosis in vitro and for virulence in macrophages and mice model of infection.

We had shown that FasR binds to three inverted repeats present in the promoter

region of the *fas-acpS* operon thereby activating fatty acid biosynthesis in *M.*

smegmatis (12). Furthermore, we also demonstrated, *in vitro* and *in vivo,* that FasR

binding to its target promoter is regulated by long-chain acyl-CoAs, the main

products of FAS I (12). The *fas* and *acpS* genes form a bicistronic operon, coding

for the FAS I synthase and the 4-phosphopantetheinyl transferase, respectively;

the latter being an essential enzyme to produce functional acyl carrier protein (holo

ACP), central to all fatty-acid biosynthesis systems. Transcriptional studies carried

out in this work showed that expression of the *fas-acpS* operon genes in *M.*

tuberculosis Δ*fasR* was reduced by 70 % and 40 %, respectively (Fig 3). This

result confirmed that FasR is a transcriptional activator of the *fas-acpS* operon in

M. tuberculosis. The activator nature of FasR highlights a remarkable difference

with most transcriptional regulators of fatty acid biosynthesis, which are repressor

proteins (19). Moreover, FasR is a member of the TetR family of regulators, and

the vast majority of these proteins are transcriptional repressors, with very few

acting as activators (20).

Since fatty acids are major constituents not only of phospholipids but also of

several complex lipids, inactivation of FasR was expected to modify synthesis of

several components of the cell envelope essential for mycobacterial viability and/or

virulence. Indeed, lipidomic analysis of *MtbΔfasR* and comparison with wild type

and complemented strains proved this hypothesis. As expected, reduced levels of

fatty acid biosynthesis in *MtbΔfasR* had a strong impact on relative abundance of

phospholipids and their relative and qualitative composition (Figs 4 and S2).

Phosphatidylinositol (PI) and cardiolipins (CL) were significantly reduced in the

mutant strain, while the acyl chains present in and these phospholipids exhibited a

significant increase in their double bond content (Fig S4). Mycobacterial

phospholipids have been implicated in immunomodulation. For example, CL is

processed into lysocardiolipin by the lysosomal phospholipase A2 during infection,

probably generating active lipid derivatives that play important roles during

infection or persistence within the host (21). The *MtbΔfasR* mutant strain also

displayed a strong reduction in the content of sulfolipids (Figs 4 and S2) and PDIM

(Figs 5 and S2). Interestingly, the acyl chain substitutions of both lipids were

consistently shorter compared to the wild type *Mtb*H37Rv strain (Fig S3). On this

regard, a *M. tuberculosis* mutant strain deficient in sulfolipids and PDIM has been

shown to be attenuated in macrophages and in mice models of infection (8).

Moreover, during infection, genes involved in the biosynthesis of sulfolipids and

PDIM are upregulated (22, 23) and the mass of these lipids increases (24, 25).

Altogether, our results showed that *M. tuberculosis fasR* mutant exhibits an

extensive rearrangement of the lipid components of the cell envelope, that are

reversed in the complemented strain *MtbΔfasR-cfasR*, confirming that FasR is

needed to maintain the lipid homeostasis of the *M. tuberculosis* cell envelope. As

the mycobacterial cell envelope is the interface with the host, FasR modulation of

its structure and consequent responsiveness to immune factors might be important

in determining the state of an infection.

Consistent with this hypothesis, we have shown that FasR is necessary for optimal

intracellular multiplication of *M. tuberculosis* in macrophages, therefore confirming

the role of cell wall components in the ability of the pathogen to survive and

multiply intracellularly (26). Persistence of the tubercle bacillus inside cells relies

upon inhibition of phagosome maturation and acidification (27). In this context,

several cell wall lipids have been shown to modulate host endocytic pathways. This

is the case for Man-Lam, TDM, PIM and PDIM, which, besides their role in

phagocytosis, also mediate intracellular trafficking and vacuole maturation arrest

induced by *M. tuberculosis* (8). More recently, PDIM and SL were shown to be

involved in controlling autophagy-related pathways in human macrophages (28).

Considering the changes observed in the PDIM content of the *MtbΔfasR* mutant,

we analyzed the acidification of phagosome during infection of human

macrophages. Interestingly, while the wild type *Mtb*H37Rv and the complemented

strains prevented phagosome acidification, the *MtbΔfasR* mutant was impaired in

avoiding phagosomal maturation, clearly indicating that increased phagosome

acidification was due to absence of FasR (Fig 7). These results suggested that

although FasR is not essential for *in vitro* growth, regulation of lipid biosynthesis,

mediated by FasR, is critical for macrophage infection. Moreover, using a mouse

model of infection we demonstrated the essentiality of FasR for *M. tuberculosis*

virulence *in vivo* (Fig 8), most probably as a consequence of the loss of various

mycobacterial lipids that contribute to pathogenicity.

Co-evolution with the human immune system prompted *M. tuberculosis* to generate

exquisite mechanisms of immunomodulation and immune evasion, many of which

depend on its cell envelope. In fact, comparison of cell envelope components

between *M. tuberculosis* and ancestral species revealed a restructuring of the

mycomembrane glycolipid components, including mycolic acids and free lipids of

the outer leaflet (11). The observed phenotype of the FasR mutant is probably a

consequence of additive effects, as several components of the cell envelope were

altered. *M. tuberculosis* interferes with innate immune signalling pathways, for

example by masking pathogen-associated molecular patterns with PDIM (29) and

inhibiting toll-like receptor 2 with sulfoglycolipids (30). PDIM is also crucial for other

steps of infection: it contributes to cell envelope properties, such as the

permeability barrier (31), improves the invasiveness of *M. tuberculosis* and its

capacity to arrest phagosome maturation, possibly by inducing changes in the

plasma membranes of human macrophages during infection (32), and directly

protects against reactive nitrogen species (33). Therefore, reduced content of

these molecules (and/or changes in their chemical composition) may contribute to

sensitivity to the bactericidal activities of phagocytes.

Altogether, our data indicate an important role for FasR in *M. tuberculosis*

pathogenesis and encourage further investigations into the contribution of lipid

regulation to virulence. It is well known that *M. tuberculosis* is able to modulate the

expression of its cell wall components in response to environmental changes,

however, little is known about their reorganization during infection, and to a lesser

extent about the regulatory mechanisms involved in this process (11). The

dynamism of the bacterial cell envelope should be a tightly regulated phenomenon,

ensuring *M. tuberculosis* survival during infection. Thus, specific regulatory

components involved in this process, as we have demonstrated for FasR, would

constitute interesting targets for the development of new treatment strategies.

Along this line of thinking, we recently solved the crystal structures of *M.*

tuberculosis FasR in complex with DNA and the acyl effector ligands, which

provided insights into the molecular sensory and transmission mechanisms of this

regulatory protein (34). This knowledge could lead now to perform structure-guided

drug discovery strategies to identify anti-mycobacterial molecules with novel

mechanisms of action.

Materials and methods

Ethics statement

All animal experiments were performed in compliance with the regulations of the

Institutional Animal Care and Use Committee (CICUAE) of INTA with the

authorization Number 20/2016.

Bacterial strains, culture and transformation conditions

The *E. coli* strain DH5α (35) was used for routine subcloning and was transformed

according to Sambrook *et al.* (36). *M. tuberculosis* H37Rv (37), *MtbΔfasR* and

MtbΔfasR-c*fasR* were grown at 37 ºC in Middlebrook 7H9 broth supplemented with

0.2 % glycerol, 0.03 % Tyloxapol and 10 % OADC or Middlebrook 7H10 agar

supplemented with 0.2 % glycerol and 10 % OADC. Hygromycin (Hyg, 50 µg/ml),

kanamycin (Kan, 15 µg/ml) and streptomycin (Str, 12 µg/ml) were added when

needed.

Recombinant plasmids and strain genotypes are listed in electronic supplementary

material (Tables S1 and S2).

DNA manipulation, plasmid construction and mutant generation

Genomic DNA of *M. tuberculosis* was obtained as described previously (38). For

the construction of the *fasR* mutant allele, a fragment of 946 bp containing 58 bp of

the *fasR* 5´end plus 882 bp of the upstream region was PCR amplified from

genomic DNA using oligonucleotides 3208_1up and 3208_2down (Table S3). The

PCR product was cloned in the pCR BluntII TOPO vector (Invitrogen). The

resulting plasmid was digested first with *Eco*RV and after with *Xho*I/*Nde*I. The 940

bp fragment obtained was cloned in the suicide vector pJG1100 (39, 40) digested

with the same enzymes, generating plasmid pFR34. Additionally, a fragment of 883

bp containing 87 bp of the *fasR* 3´end plus 784 bp of the downstream region was

PCR amplified from genomic DNA using oligonucleotides 3208_3up and

3208_4down (Table S3). The PCR product was cloned in the pCR BluntII TOPO

vector (Invitrogen). The resulting plasmid was digested first with *Eco*RI and after

with *Nde*I/*Spe*I and the fragment (877 bp) was cloned in pFR34 vector digested

with the same enzymes, generating plasmid pFR35.

For the construction of a *fasR* merodiploid strain, the fragment resulting from the digestion of the plasmid pFR3 (12) with *Xba*I and *Eco*RI was cloned between the *Avr*II and *Mfe*I sites in plasmid pGA44 (41). The resulting plasmid, pFR33, is an integrative plasmid that harbours *fasR* expressed under the control of *Ptr* promoter and the TetR/Pip OFF system necessary for its regulation.

Deletion of *fasR* was accomplished by homologous recombination. After transformation of *M. tuberculosis* with plasmid pFR35, the first recombination event was selected on 7H10 medium containing Kan and Hyg and resistant colonies were screened by colony PCR. The merodiploid strain was generated by integration of pFR33 at the L5 *attB* site and transformants were plated on 7H10 with Hyg, Kan and Str. Deletion of the wild type gene by allelic replacement was accomplished by plating bacteria on 7H10 agar plates supplemented with Str and 2.5 % sucrose. The Hyg^s, Kan^s, Str^r, Sucrose^r colonies were tested by PCR and the *fasR* conditional knockdown strain was named *MtbΔfasR*-c*fasR*. Generation of a *fasR* knockout strain (*MtbΔfasR*) was obtained by swapping the pFR33 plasmid at the L5 *attB* site with the pND255 plasmid. Transformants were selected on 7H10 plates containing Hyg. Deletion of *fasR* was confirmed by Southern Blot and colony PCR.

Southern Blot

10 µg of genomic *M. tuberculosis* DNA was digested overnight with an excess of *Xho*I, and the fragments were separated by electrophoresis in a 0.7% agarose gel. Southern blotting was carried out using Hybond-N$^+$ membrane (Amersham) and DIG High Prime DNA Labeling and Detection Starter kit II (Roche), following the indications of the manufacturer. The probe consisted of a 594 bp fragment

428 amplified with oligonucleotides Southern_1fasRTB and Southern_2fasRTB (Table

429 S3), corresponding to the upstream genomic region of *fasR*.

Western blot

Equal amounts of protein extracts were loaded on 4-12 % NuPAGE gels

(Invitrogen) and transferred onto nitrocellulose membranes using a dry

electrophoresis transfer apparatus (Invitrogen). Membranes were incubated in

TBS-Tween blocking buffer (20 mM Tris pH 7.5, 150 mM NaCl, 0.1 % Tween 20)

with 5 % (w/v) skimmed milk powder for 2 h at 4 $^{\circ}$C prior to overnight incubation

with 1:1000 dilution of monoclonal anti-FasR primary antibody (Alere^{TM}).

Membranes were washed in TBS-Tween three times at room temperature, and

then incubated with 1:100000 secondary antibody Anti-Mouse Kappa conjugated to

Horseradish Peroxidase (Southern Biotech) for 2 h before washing again. Signals

were detected using Chemiluminescent Peroxidase Substrate (Sigma-Aldrich).

Monoclonal mouse anti-RpoB antibodies were purchased from NeoClone and used

1:20000.

RNA techniques

RNA was extracted from *M. tuberculosis* strains grown for 8 days in complete 7H9

media using Trizol-phenol based methods (41) and treated, when needed, with

RQ1 RNase-Free DNase (Promega). qRT-PCR was performed using second

strand cDNA as the template, generated with SuperScript III Reverse

Transcriptase (Invitrogen), random primers and a green fluorochrome as the

indicator dye (qPCR master mix, Biodynamics). The expression of *fas*, *fasR* and

acpS was quantified after normalization of RNA levels to the expression of the *sigA*

gene as previously described (15). qPCR data are presented as relative

expression values normalized to *sigA* and to the wild type strain H37Rv. qPCR

cycling conditions were as follows: 95 ºC for 2 min followed by 40 cycles of 95 ºC

for 15 s, 58 ºC for 15 s and 68 ºC for 20 s. The sequences of all the primers used

are listed in electronic supplementary material (Table S3).

Lipid analysis

Cell-associated lipid for lipidomic analyses were prepared as previously described

(15). Briefly, cell pellets of cultures grown at 37 ºC in Middlebrook 7H9 broth

(DIFCO) supplemented with 0.2 % glycerol, 0.03 % Tyloxapol and 10 % OADC

were washed with 50 mM ammonium acetate, pH 7.8 and transferred to glass

tubes containing 5 ml $CHCl_3:CH_3OH$ (2:1, v/v). Samples were incubated overnight

at 4 ºC with gentle agitation. After centrifugation, bacterial pellets were subjected to

an additional extraction using $CHCl_3:CH_3OH$ (1:2, v/v) for 2 h. Organic extracts

were pooled and dried under nitrogen at 4 ºC. Lipids were resuspended in 3 ml of

$CHCl_3$, washed with 3 ml of H_2O and the organic phase was transferred to pre-

weighted glass tubes, dried under nitrogen at 4 ºC and re-weighted on a

microbalance. Extracts were dissolved in $CHCl_3:CH_3OH$ (1:1, v/v) at 1 mg ml^{-1} and

centrifuged at 3000 g for 5 min. Lipids were analyzed in an Agilent 1200 series

HPLC system with a Reprospher (Dr Maish) 100 C8 column (1.8 μm, 50 mm

length, 2 mm ID). The flow rate was 0.3 ml min^{-1} in binary gradient mode with the

following elution program: the column was equilibrated with 100 % mobile phase A

[$CH_3OH:H_2O$ (99:1, v/v), containing 0.05 mM $AcNH_4$]. 2 μl of each sample were

injected, and the same elution conditions continued for 1 min, followed by a 12 min

gradient to 100 % mobile phase B [isopropanol:hexane:H_2O (79:20:1, v/v),

containing 0.05 mM $AcNH_4$], finally holding this condition for 1 min. The column

was equilibrated for 2 min with 100 % mobile phase A before injection of the next

sample. An Agilent 6500 series Q-TOF instrument with a Dual AJS ESI was used

for mass analysis. Ionization gas temperature was maintained at 200 ºC with a 14 l

min^{-1} drying gas flow, a 35 psig nebulizer pressure and 3500 Volts. Spectra were

collected in positive and negative mode from *m/z* 115 to 3000 at 4 spectra s^{-1}.

Automatic tandem-MS (MS-MS) was obtained in data-dependent acquisition mode

at 15 MS-MS per cycle. Continuous infusion calibrants included *m/z* 121.051 and

922.010 in positive-ion mode and *m/z* 119.035 and 955.972 in negative-ion mode.

CID-MS was carried out with an energy of 50 volts. For the comparative analysis,

the column is conditioned by four successive mock injections before randomized

QCs and mycobacterial samples are analyzed. Lipid features were annotated

based on structural MS-MS information. Lipid classes' retention times (RT) were

identified using characteristics fragmentation patterns (42). Individual molecules in

each class were then identified based on exact mass and RT.

Colony-forming units (CFU) analysis

M. tuberculosis H37Rv, *MtbΔfasR* and *MtbΔfasR*-c*fasR* strains were used to infect

THP-1 cells (MOI=5). The day before infection $4x10^5$ THP-1 cells were plated in a

24-well plate in RPMI Medium with 4.5 g/L glucose, 10 % heat inactivated fetal calf

serum (Internegocios) and 2 mM L-glutamine (full medium) plus 50 nM phorbol 12-

myristate 13-acetate (PMA) to allow differentiation of the cells to macrophages. On

the infection day, cells were washed and a suspension of the different strains of *M.*

tuberculosis in RPMI medium was added. After 2 h of uptake, cells were washed 3

times with PBS and fresh full RPMI medium plus gentamicin 10 µg mL^{-1} was

added. Cells were incubated at 37 °C in 5 % CO_2 atmosphere. Subsequently, at

the indicated time points, cells were lysed in sterile water and serially diluted in

PBS-Tween 80 0.05 %. Dilutions were plated on Middlebrook 7H10 complete agar

medium (Difco Laboratories). Colonies were counted after 3 to 4 weeks to

determine CFU.

Mouse infections

Bacillary suspensions of the different strains were adjusted to 100 viable cells in

100 ml phosphate buffer saline (PBS). Each animal was anesthetized and

intratracheally inoculated with *M. tuberculosis* strains (*Mtb*H37Rv, *MtbΔfasR* and

MtbΔfasR-c*fasR*). Infected mice were kept in cages fitted with microisolators

connected to negative pressure. Groups of 10 mice were infected with the different

M. tuberculosis strains. The inoculum doses were determined by enumerating CFU

recovered from the lungs of three mice per strain, 24 h post-infection. Seven mice

per group were killed at 60 days after infection and lungs removed and

homogenized. Four dilutions of each homogenate were spread onto triplicate

plates of Middlebrook 7H10 complete agar medium. The number of colonies was

counted after 3 to 4 weeks.

Infection of THP-1 cells and indirect immunofluorescence (IFI)

M. tuberculosis strains were covalently stained with FITC (Sigma-Aldrich). Briefly,

1×10^9 bacteria were washed twice with 0.1 M PBS and suspended in 1 ml of the

same solution. FITC was added to a final concentration of 5 µg/ml and incubated in

the dark for 1 h at 37 °C. Bacteria were washed gently with PBS until

unincorporated dye was eliminated, and then were used to infect THP-1 cells.

THP-1 cells were maintained in complete full RPMI medium. The day before

infection 2×10^5 THP-1 cells per well were plated in a 24-well plate with coverslips in

full RPMI medium plus 50 nM PMA to allow differentiation of the cells to

macrophages. On the day of infection, cells were washed and infected with the

different stained *M. tuberculosis* strains (*Mtb*H37Rv, *Mtb*Δ*fasR* and *Mtb*Δ*fasR*-

c*fasR*) prepared in RPMI medium, at a MOI of 5. After 2 h of uptake, cells were

washed and incubated with 10 μM Lysotracker Red DND-99 for 1 hour to allow the

staining of acidic compartments, fixed with 4 % paraformaldehyde solution in PBS

for 30 min and subjected to immunofluorescence analysis.

Cells were quenched by incubation with 50 mM NH_4Cl solution in PBS for 30 min.

and permeabilized with 0.05 % saponin/1 % BSA in PBS for 15 min, followed by

overnight incubation with goat anti-LAMP-3 (Santa Cruz Biotechnology, Santa

Cruz, CA) primary antibody diluted 1:50 in PBS. Secondary anti-goat antibody

conjugated to Cy3 (Jackson Immuno Research Labs, Inc.) was diluted 1:600 in

PBS. Cells were mounted with mounting medium (Dako, Denmark) and analysed

by confocal microscopy using an SP5 AOBS confocal microscope (Leica

Microsystems, Germany). Mycobacterial internalization was monitored by following

FITC fluorescence (green). Lysotracker Red DND-99 or LAMP-3 association with

mycobacterial phagosomes was analysed in at least 250 cells using Fiji software

(U.S. National Institutes of Health, Bethesda, MD) as described previously (43).

The experiments were performed in duplicates in three independent experiments.

The statistical analysis was performed using two-tailed Student's t-test.

Acknowledgements

This research was supported by Agencia Nacional de Promoción de la

Investigación, el Desarrollo Tecnológico y la Innovación (ANPCyT) grants 2011-

0245, 2015-0796 and 2018-02539 to GG and 2015-2022 to HG and by National

Institutes of Health grant 1R01AI095183-01 to HG. The funders had no role in

study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

References

1.   WHO. 2019. Global tuberculosis report.

2.   Chiaradia L, Lefebvre C, Parra J, Marcoux J, Burlet-Schiltz O, Etienne G,

Tropis M, Daffé M. 2017. Dissecting the mycobacterial cell envelope and

defining the composition of the native mycomembrane. Sci Rep 7:12807.

3.   Daffé M, Marrakchi H. 2019. Unraveling the Structure of the Mycobacterial

Envelope. Microbiol Spectr 7:GPP3-0027–2018.

4.   Brennan   PJ,   Nikaido   H.   1995.   The   envelope   of   mycobacteria.

AnnuRevBiochem 64:29–63.

5.   Daffe M, Reyrat JM. 2008. The Mycobacterial Cell Envelope. ASM Press.

6.   Russell DG, Barry CE, Flynn JL. 2010. Tuberculosis: What We Don't Know

Can, and Does, Hurt Us. Science 328:852–856.

7.   Neyrolles O, Guilhot C. 2011. Recent advances in deciphering the

contribution of Mycobacterium tuberculosis lipids to pathogenesis.

Tuberculosis 91:187–195.

8.   Passemar C, Arbues A, Malaga W, Mercier I, Moreau F, Lepourry L,

Neyrolles O, Guilhot C, Astarie-Dequeker C. 2014. Multiple deletions in the

polyketide synthase gene repertoire of mycobacterium tuberculosis reveal

functional overlap of cell envelope lipids in host-pathogen interactions. Cell

Microbiol 16:195–213.

9.   Vromman F, Subtil A. 2014. Exploitation of host lipids by bacteria. Curr Opin

Microbiol 17:38–45.

10.   Teng O, En Ang CK, Guan XL. 2017. Macrophage-bacteria interactions-A

lipid-centric relationship. Front Immunol 8.

11.   Dulberger C, Rubin E, Boutte CC. 2020. The mycobacterial cell envelope —

a moving target. Nat Rev Microbiol 18:47–59.

12.   Mondino S, Gago G, Gramajo H. 2013. Transcriptional regulation of fatty

acid biosynthesis in mycobacteria. Mol Microbiol 89:372–387.

13.   Salzman V, Mondino S, Sala C, Cole ST, Gago G, Gramajo H. 2010.

Transcriptional regulation of lipid homeostasis in mycobacteria. Mol Microbiol

78:64–77.

14.   Boldrin F, Casonato S, Dainese E, Sala C, Dhar N, Palù G, Riccardi G, Cole

ST, Manganelli R. 2010. Development of a repressible mycobacterial

promoter system based on two transcriptional repressors. Nucleic Acids Res

38:1–11.

15.   Cabruja M, Mondino S, Tsai YT, Lara J, Gramajo H, Gago G. 2017. A

conditional mutant of the fatty acid synthase unveils unexpected cross talks

in mycobacterial lipid metabolism. Open Biol 7.

16.   Sassetti CM, Boyd DH, Rubin EJ. 2003. Genes required for mycobacterial

growth defined by high density mutagenesis. MolMicrobiol 48:77–84.

17.   Griffin JE, Gawronski JD, DeJesus MA, Ioerger TR, Akerley BJ, Sassetti CM.

2011. High-resolution phenotypic profiling defines genes essential for

mycobacterial growth and cholesterol catabolism. PLoS Pathog 7:1–9.

18.   DeJesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, Rubin EJ,

Schnappinger D, Ehrt S, Fortune SM, Sassetti CM, Ioerger R. 2017.

Comprehensive Essentiality Analysis of the Mycobacterium tuberculosis

Genome via sturating transposon mutagenesis. MBio 8:e02133–1.

19.   Zhu K, Zhang YM, Rock CO. 2009. Transcriptional regulation of membrane

lipid homeostasis in Escherichia coli. JBiolChem 284:34880–34888.

20.   Cuthbertson L, Nodwell JR. 2013. The TetR Family of Regulators. Microbiol

Mol Biol Rev 77:440–475.

21.   Fischer K, Chatterjee D, Torrelles J, Brennan PJ, Kaufmann SHE, Schaible

UE. 2001. Mycobacterial Lysocardiolipin Is Exported from Phagosomes

upon Cleavage of Cardiolipin by a Macrophage-Derived Lysosomal

Phospholipase A 2 . J Immunol 167:2187–2192.

22.   Graham JE, Clark-Curtiss JE. 1999. Identification of Mycobacterium

tuberculosis RNAs synthesized in response to phagocytosis by human

macrophages by selective capture of transcribed sequences (SCOTS).

Proc Natl Acad Sci 96:11554–11559.

23.   Rodríguez JE, Ramírez AS, Salas LP, Helguera-Repetto C, Gonzalez-y-
Merchand J, Soto CY, Hernández-Pando R. 2013. Transcription of Genes
Involved in Sulfolipid and Polyacyltrehalose Biosynthesis of Mycobacterium
tuberculosis in Experimental Latent Tuberculosis Infection. PLoS One 8.

24.   Griffin JE, Pandey AK, Gilmore SA, Mizrahi V, Mckinney JD, Bertozzi CR,
Sassetti CM. 2012. Cholesterol catabolism by Mycobacterium tuberculosis
requires transcriptional and metabolic adaptations. Chem Biol 19:218–227.

25.   Yang X, Nesbitt NM, Dubnau E, Smith I, Sampson NS. 2009. Cholesterol
Metabolism Increases the Metabolic Pool of Propionate in M. tuberculosis.
Biochemistry 48:3819–3821.

26.   Garcia-Vilanova A, Chan J, Torrelles JB. 2019. Underestimated Manipulative
Roles of Mycobacterium tuberculosis Cell Envelope Glycolipids During
Infection. Front Immunol 10.

27.   Rohde K, Yates RM, Purdy GE, Russell DG. 2007. Mycobacterium
tuberculosis and the environment within the phagosome. Immunol Rev
219:37–54.

28.   Bah A, Sanicas M, Nigou J, Guilhot C, Astarie-Dequeker C, Vergne I. 2020.
The Lipid Virulence Factors of Mycobacterium tuberculosis Exert
Multilayered Control over Autophagy-Related Pathways in Infected Human
Macrophages. Cells 9:666.

29.   Cambier CJ., Takaki KK., Larson RP., Hernandez RE., Tobin DM., Urdahl
KB., Cosma CL., Ramakrishnan L. 2014. Mycobacteria manipulate
macrophage recruitment through coordinated use of membrane lipids.
Nature 505:218–222.

30.   Blanc L, Gilleron M, Prandi J, Song O, Jang M, Gicquel B, Drocourt D,

Neyrolles O, Brodin P, Tiraby G, Vercellone A, Nigou J. 2017.

Mycobacterium tuberculosis inhibits human innate immune responses via the

production of TLR2 antagonist glycolipids. Proc Natl Acad Sci 114:11205–

11210.

31.   Camacho LR, Constant P, Raynaud C, Lane M, Triccas JA, Gicquel B, Daffe

M, Guilhot C. 2001. Analysis of the Phthiocerol Dimycocerosate Locus of

Mycobacterium tuberculosis. J Biol Chem 276:19845–19854.

32.   Astarie-Dequeker C, Le Guyader L, Malaga W, Seaphanh FK, Chalut C,

Lopez A, Guilhot C. 2009. Phthiocerol dimycocerosates of M. tuberculosis

participate in macrophage invasion by inducing changes in the organization

of plasma membrane lipids. PLoS Pathog 5:e1000289.

33.   Rousseau C, Winter N, Pivert E, Bordat Y, Neyrolles O, Avé P, Huerre M,

Gicquel B, Jackson M. 2004. Production of phthiocerol dimycocerosates

protects Mycobacterium tuberculosis from the cidal activity of reactive

nitrogen intermediates produced by macrophages and modulates the early

immune response to infection. Cell Microbiol 6:277–287.

34.   Lara J, Diacovich L, Trajtenberg F, Larrieux N, Malchiodi EL, Fernandez M,

Gago G, Gramajo H, Buschiazzo A. 2020. Mycobacterium tuberculosis FasR

senses long fatty acyl-CoA through a tunnel , inducing DNA-dissociation via

a transmission spine. Nat Comm, in press.

35.   Hanahan D. 1983. Studies on transformation of Escherichia coli with

plasmids. JMolBiol 166:557–580.

36.   Sambrook J. EFFTM. 1989. Molecular cloning: a laboratory manual, 2nd ed.

665   Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

666 37. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon S V,

667   Eiglmeier K, Gas S, Barry III CE, Tekaia F, Badcock K, Basham D, Brown D,

668   Chillingworth T, Connor R, Davies R, Devlin K, Feltwell T, Gentles S, Hamlin

669   N, Holroyd S, Hornsby T, Jagels K, Krogh A, McLean J, Moule S, Murphy L,

670   Oliver K, Osborne J, Quail MA, Rajandream MA, Rogers J, Rutter S, Seeger

671   K, Skelton J, Squares R, Squares S, Sulston JE, Taylor K, Whitehead S,

672   Barrell BG. 1998. Deciphering the biology of Mycobacterium tuberculosis

673   from the complete genome sequence. Nature 393:537–544.

674 38. Connell ND. 1994. Mycobacterium: isolation, maintenance, transformation,

675   and mutant selection. Methods Cell Biol 45:107–125.

676 39. Gomez JE, Bishai WR. 2000. whmD is an essential mycobacterial gene

677   required for proper septation and cell division. Proc Natl Acad Sci U S A

678   97:8554–8559.

679 40. Muñoz-Elías EJ, McKinney JD. 2006. Carbon metabolism of intracellular

680   bacteria. Cell Microbiol 8:10–22.

681 41. Kolly GS, Boldrin F, Sala C, Dhar N, Hartkoorn RC, Ventura M, Serafini A,

682   Mckinney JD, Manganelli R, Cole ST. 2014. Assessing the essentiality of the

683   decaprenyl-phospho-d-arabinofuranose  pathway  in  Mycobacterium

684   tuberculosis using conditional mutants. Mol Microbiol 92:194–211.

685 42. Layre E, Sweet L, Hong S, Madigan CA, Desjardins D, Young D, Cheng T,

686   Annand JW, Kim K, Shamputa I, McConnell MJ, Debono CA, Behar SM,

687   Minnaard AJ, Murray M, Barry III CE, Matsunaga I, Moody DB. 2011. A

688   comparative  lipidomics  platform  for  chemotaxonomic  analysis  of

Mycobacterium tuberculosis. Chem Biol 23:1537–1549.

43.   Vázquez CL, Bianco MV, Blanco FC, Forrellad MA, Gutierrez MG, Bigi F.

2017. Mycobacterium bovis Requires P27 (LprG) To Arrest Phagosome

Maturation and Replicate within Bovine Macrophages. Infect Immun 85:1–11.

Figure legends

Fig 1. Schematic representation of the genetic organization of the

chromosomal region of *fasR*.

(**A**) Wild type H37Rv strain (*Mtb*H37Rv), (**B**) *fasR* conditional knockdown strain

(*Mtb∆fasR*-c*fasR*) and (**C**) *fasR* KO strain (*Mtb∆fasR*). (**D**) Southern Blot analysis

of the different strains: chromosomal DNA was digested with *Xho*I and probed for

hybridization with the probe indicated with a black bar, corresponding to the 5´ end

of the *Rv3207* gene.

Fig 2. Growth of *M. tuberculosis* strains.

Growth curves of the wild type strain (*Mtb*H37Rv), the *fasR Mtb∆fasR*-c*fasR* strain

grown without ATc (complemented strain) and *fasR* KO strain (*Mtb∆fasR*) were

obtained in 3 different media (7H9 complete, 7H9 without OADC and Sauton).

Growth was followed by measuring $OD_{600\ nm}$. Results are representative of three

independent experiments.

Fig 3. Relative expression levels of *fasR, fas* and *acpS* mRNAs measured by quantitative RT-PCR.

Values were normalized to *sigA* and to the wild type strain (*Mtb*H37Rv). Samples for RNA extraction were collected at OD_{600nm} ~ 0.3.

Fig 4. Lipid profile in negative ion mode

LC-MS analyses (negative mode) of the phospholipids and mycolic acids present in the wild type strain (*Mtb*H37Rv), the *fasR* complemented strain (*Mtb*Δ*fasR*-c*fasR*) and *fasR* KO strain (*Mtb*Δ*fasR*). The values indicated represent the relative abundance of the MS signals corresponding to the different lipid classes found in the samples. Results are the means of three independent experiments. CL, cardiolipin; PI, phosphatidylinositol; PIM, phosphatidylinositol mannosides; PG, phosphatidylglycerol; PE, phosphatidylethanolamine; SL, sulfolipids; MA, mycolic acids.

Fig 5. Lipid profile in positive ion mode

LC-MS analyses (positive mode) of lipids present in the wild type strain (*Mtb*H37Rv), the *fasR* complemented strain (*Mtb*Δ*fasR*-c*fasR*) and *fasR* KO strain (*Mtb*Δ*fasR*). The values indicated represent the relative abundance of the MS signals corresponding to the different lipid classes found in the samples. Results are the means of three independent experiments. TAG, triacylglycerol; DAG, diacylglycerol; MMDAG, monomycolyl DAG; PDIM, phthiocerol dimycocerosates; GroMM, glycerol monomycolate; TMM, trehalose monomycolate.

Fig 6. FasR is required for the intracellular survival of *M. tuberculosis*.

THP-1-derived macrophages were infected with wild type *M. tuberculosis* H37Rv (*Mtb*H37Rv), *fasR* mutant (*Mtb*Δ*fasR*) and the complemented *fasR* strain (*Mtb*Δ*fasR*-c*fasR*) at a MOI =5. At different time post-infection (24, 48, 120, 144 hours) cells were washed and intracellular viable bacteria were counted on agar plates. Bars represent the average of three independent experiments performed in triplicate wells with standard deviation.

Fig 7. FasR regulates phagosome maturation in macrophages.

Confocal microscopy analysis of THP-1 macrophages infected with wild type *M. tuberculosis* H37Rv (*Mtb*H37Rv), *fasR* mutant (*Mtb*Δ*fasR*) and the complemented *fasR* strain (*Mtb*Δ*fasR*-c*fasR*). A) Representative images of THP-1 macrophages infected with the different strains labelled with FITC (green fluorescence) using a MOI=5. After 2 h of uptake, cells were washed and incubated with Lysotracker Red DND-99 for 1 hour to allow the staining of acidic compartments, fixed and subjected to immunofluorescence. The infected and fixed cells were used to detect LAMP-3 by indirect immunofluorescence using a specific antibody against LAMP-3. Scale bars: 10 μm. Merge indicates the location of the different strains in the phagolysosome. B, C) Fluorescence intensity quantification of Lysotracker and LAMP-3 associated with the phagosome of *M. tuberculosis Mtb*H37Rv (circles), *Mtb*Δ*fasR* (squares) and the complemented *fasR* strain *Mtb*Δ*fasR*-c*fasR* (triangles). Data were expressed in arbitrary units (a.u.). Fiji software was used for quantification. Data represent means ± SEM of three independent experiments.

Asterisks indicate significance: *, P ≤ 0.05; **, P ≤ 0.01; and ***, P ≤ 0.001. Data

were analyzed using a two-tailed Student *t* test.

Fig 8. FasR is essential for the multiplication of mycobacteria in mouse

lungs.

Three groups of BALB/c mice (10 per group) were infected intratracheally with 10^2

CFU of *Mtb*H37Rv, *MtbΔfasR* and *MtbΔfasR-cfasR* strains and 60 days post-

infection bacterial loads in lungs were determined by plating serial dilutions on

Middlebrook 7H10 complete agar medium. The graph shows the mean values and

error bar of CFU. Data were analyzed using one-way ANOVA analysis and

Bonferroni's post-test. (**, P <0.01; ***, P ≤ 0.001).

Supplemental material

Table S1. Plasmids used in this study

Table S2. Strains used in this study

Table S3. Oligonucleotides used in this study

Figure S1. Analysis of the knockout fasR mutant (A) Colony PCR of M.

tuberculosis transformants grown on Str, carrying the complementing fasR gene

(clones 1 and 2) and of transformants grown on Hyg, carrying the empty pND255

vector (clones 3 and 4). Primers used for the amplification of fasR are listed in

Table S3. The expected size of the PCR products is 687 bp for the fasR wild type

allele and 150 bp for the mutant fasR allele. Two fragments were amplified from the

Str-resistant clones, corresponding to the complementing fasR gene from the

integrated plasmid and the deleted fasR gene at the fasR locus; on the other hand,

only one fragment was amplified from the Hyg resistant clones (the partially

deleted fasR gene) (B) Western Blot analysis of crude protein extracts prepared

from cultures of wild type H37Rv strain (MtbH37Rv), fasR conditional knockdown

strain (MtbΔfasR-cfasR) and fasR KO strain (MtbΔfasR clones 3 and 4) grown at

OD600nm ~ 0,3. Detection was performed using monoclonal anti-FasRMT

antibodies (lower panel) and anti-RpoB as loading control (upper panel).

Figure S2. Relative abundance of lipid species. The tables show the relative

abundance as the total signal percentage of the different lipid classes. CL,

cardiolipin; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PI,

phosphatidylinositol; PIMs, phosphatidylinositol mannosides; SL, sulfoglycolipids;

MA, mycolic acids; DAG, diacylglycerol; TAGs, triacylglycerol; MMDAG,

monomycolyl-DAG; GroMM, glycerol monomycolate; PDIM, phthiocerol

dimycocerosates; TMM, trehalose monomycolate. Results are the mean of three

independent experiments ± SD (n = 3). p values were calculated with unpaired t

test with Benjamini-Hochberg's correction between MtbΔfasR and MtbH37Rv.

Figure S3. Detailed analysis of lipid species. (A) The tables show the weighted

average of total number of carbons atoms per molecule within the population of

mycolic acids (MA), phospholipids (PE, phosphatidylethanolamine; CL, cardiolipin;

PG, phosphatidylglycerol; PI, phosphatidylinositol); sulfoglycolipids (SL); DAG,

diacylglycerol; GroMM, glycerol monomycolate; TAGs, triacylglycerol; TMM

trehalose monomycolate and phthiocerol dimycocerosates (PDIM A). The

histograms show the distribution of the number of carbon atoms of SL (B) and

PDIM (C) present in the wild type strain (MtbH37Rv), the fasR complemented

strain (MtbΔfasR-cfasR) and fasR mutant strain (MtbΔfasR). Results are the mean

of three independent experiments ± SD (n = 3). p values were calculated with

unpaired t test with Benjamini-Hochberg's correction between MtbΔfasR and

MtbH37Rv.

Figure S4. Saturation level of the fatty acid substituents of the phospholipids.

Histograms represent the levels of unsaturation of the fatty acids constituents of

cardiolipin (CL), phosphatidylglycerol (PG) and phosphatidylinositol (PI). Results

are the means of three independent experiments.

Fig 1. **Schematic representation of the genetic organization of the chromosomal region of *fasR*.**
(**A**) Wild type H37Rv strain (*Mtb*H37Rv), (**B**) *fasR* conditional knockdown strain (*MtbΔfasR*-c*fasR*) and (**C**) *fasR* KO strain (*MtbΔfasR*). (**D**) Southern Blot analysis of the different strains: chromosomal DNA was digested with *Xho*I and probed for hybridization with the probe indicated with a black bar, corresponding to the 5´end of the *Rv3207* gene.

Fig 2. Growth of *M. tuberculosis* strains.
Growth curves of the wild type strain (*Mtb*H37Rv), the *fasR Mtb∆fasR-cfasR* strain grown without ATc (complemented strain) and *fasR* KO strain (*Mtb∆fasR*) were obtained in 3 different media (7H9 complete, 7H9 without OADC and Sauton). Growth was followed by measuring $OD_{600\ nm}$. Results are representative of three independent experiments.

Fig 3. Relative expression levels of *fasR*, *fas* and *acpS* mRNAs measured by quantitative RT-PCR.
Values were normalized to *sigA* and to the wild type strain (*Mtb*H37Rv). Samples for RNA extraction were collected at $OD_{600nm} \sim 0{,}3$.

Mycolic acids		Phospholipids	
*Mtb*H37Rv	25%	*Mtb*H37Rv	75%
*Mtb*Δ*fasR*	50%	*Mtb*Δ*fasR*	50%
*Mtb*Δ*fasR-cfasR*	22%	*Mtb*Δ*fasR-cfasR*	78%

Fig 4. Lipid profile in negative ion mode
 LC-MS analyses (negative mode) of the phospholipids and mycolic acids present in the wild type strain (*Mtb*H37Rv), the *fasR* complemented strain (*Mtb*Δ*fasR*-c*fasR*) and *fasR* KO strain (*Mtb*Δ*fasR*). The values indicated represent the relative abundance of the MS signals corresponding to the different lipid classes found in the samples. Results are the means of three independent experiments. CL, cardiolipin; PI, phosphatidylinositol; PIM, phosphatidylinositol mannosides; PG, phosphatidylglycerol; PE, phosphatidylethanolamine; SL, sulfolipids; MA, mycolic acids.

Fig 5. Lipid profile in positive ion mode
 LC-MS analyses (positive mode) of lipids present in the wild type strain (*Mtb*H37Rv), the *fasR* complemented strain (*MtbΔfasR-cfasR*) and *fasR* KO strain (*MtbΔfasR*). The values indicated represent the relative abundance of the MS signals corresponding to the different lipid classes found in the samples. Results are the means of three independent experiments. TAG, triacylglycerol; DAG, diacylglycerol; MMDAG, monomycolyl DAG; PDIM, phthiocerol dimycocerosates; GroMM, glycerol monomycolate; TMM, trehalose monomycolate.

Fig 6. FasR is required for the intracellular survival of *M. tuberculosis*.
THP-1-derived macrophages were infected with wild type *M. tuberculosis* H37Rv (*Mtb*H37Rv), *fasR* mutant (*MtbΔfasR*) and the complemented *fasR* strain (*MtbΔfasR*-c*fasR*) at a MOI =5. At different time post-infection (24, 48, 120, 144 hours) cells were washed and intracellular viable bacteria were counted on agar plates. Bars represent the average of three independent experiments performed in triplicate wells with standard deviation.

A

	Lysotracker	LAMP-3	Merge
*Mtb*H37Rv			
MtbΔfasR			
MtbΔfasR-cfasR			

B

Lysotracker associated to Mtb phagosome (a.u.)

*** **

MtbH37Rv   *MtbΔfasR*   *MtbΔfasR-cfasR*
M. tuberculosis

C

LAMP-3 associated to Mtb phagosome (a.u.)

*** *

MtbH37Rv   *MtbΔfasR*   *MtbΔfasR-cfasR*
M. tuberculosis

Fig 7. FasR regulates phagosome maturation in macrophages.

Confocal microscopy analysis of THP-1 macrophages infected with wild type *M. tuberculosis* H37Rv (*Mtb*H37Rv), *fasR* mutant (*MtbΔfasR*) and the complemented *fasR* strain (*MtbΔfasR-cfasR*). A) Representative images of THP-1 macrophages infected with the different strains labelled with FITC (green fluorescence) using a MOI=5. After 2 h of uptake, cells were washed and incubated with Lysotracker Red DND-99 for 1 hour to allow the staining of acidic compartments, fixed and subjected to immunofluorescence. The infected and fixed cells were used to detect LAMP-3 by indirect immunofluorescence using a specific antibody against LAMP-3. Scale bars: 10 µm. Merge indicates the location of the different strains in the phagolysosome. B, C) Fluorescence intensity quantification of Lysotracker and LAMP-3 associated with the phagosome of *M. tuberculosis Mtb*H37Rv (circles), *MtbΔfasR* (squares) and the complemented *fasR* strain *MtbΔfasR-cfasR* (triangles). The data were expressed in arbitrary units (a.u.). Fiji software was used for quantification. Data represent the means ±the SEM of three independent experiments. The asterisks indicate significance: *, $P \leq 0.05$; **, $P \leq 0.01$; and ***, $P \leq 0.001$. The data were analyzed using a two-tailed Student *t* test.

Fig 8. FasR is essential for the multiplication of mycobacteria in mouse lungs.
Three groups of BALB/c mice (10 per group) were infected intratracheally with 10^2 CFU of *Mtb*H37Rv, *MtbΔfasR* and *MtbΔfasR*-c*fasR* strains and 60 days post-infection bacterial loads in lungs were determined by plating serial dilutions on Middlebrook 7H10 complete agar medium. The graph shows the mean values and error bar of CFU. Data were analyzed using one-way ANOVA analysis and Bonferroni's post-test. (**, $P < 0.01$; ***, $P \leq 0.001$).

Table S1. Plasmids used in this study

Name	Description and features	References
pFR3	pET28a(+) with *fasR* His tag fusion gene, under the control of T7 promoter, Kan^R	(1)
pJG1100	Suicide vector for mutant construction, Hyg^R, Kan^R, *sacB*	(2)
pFR35	Suicide vector for *MtbΔfasR* mutant construction derived from pJG1100, Hyg^R, Kan^R, *sacB*	This study
pGA44	Integrative vector at L5 *attB* site, carrying the TetR/Pip OFF expression system and Ptr promoter, Str^R/Spect^R The *int* gene was removed in order to lock this plasmid at the integration site. Integration requires plasmid pGA80	(3)
pGA80	pMV261-derived vector, carrying the L5 *int* gene for expression in *trans*, lacking the mycobacterial origin of replication (Δ*oriM*), Kan^R	(3)
pFR33	pGA44-derived vector carrying *fasR* under the control of the Ptr promoter, Str^R/Spect^R	This study
pND255	L5-integrative vector for mutant construction, Hyg^R	(3)

Hyg^R, hygromicin resistance; Kan^R, kanamycin resistance; Str^R/Spect^R, streptomycin/spectinomycin resistance

Table S2. Strains used in this study

Name	Description or Genotype	References
DH5α	*E. coli K12 F- ΔlacU169 (ϕ80lacZΔM15) endA1 recA1 hsdR17 deoR supE44 thi-1λ- gyrA96 relA1*	(4)
*Mtb*H37Rv	*M. tuberculosis* wild type strain	(5)
MtbΔfasR	*Mtb*H37Rv background. Deletion of *fasR*	This study
MtbΔfasR-cfasR	*fasR* knockout complemented with *fasR* expressed *in trans*	This study

Table S3. Oligonucleotides used in this study

Name	Sequence (5'-3')
3208_1up	CTCGAGCCGGAATTCGTAGCC
3208_2down	CATATGGAGCGCTGCCGGACGATCTG
3208_3up	CATATGAAGCCGATTTCCAAGTCCGAC

3208_4down	ACTAGTTGCCACCGTGGTGACCGTCTC
Southern_1fasRTB	GAACACTTTGACGGTGCCTTG
Southern_2fasRTB	GGAATGACATTACTACCTGCG
L-*sigA*	AAACAGATCGGCAAGGTAGC
R-*sigA*	CTGGATCAGGTCGAGAAACG
L-*fasR*	TGAGCACGACTACCGACAAC
R-*fasR*	CTCGAAGATCAGCCGGTAAC
L-*fas*	GACATCCTGACCCGACTGAC
R-*fas*	TGAACTTCGTCGAGAGCTTG
fasRF	ATGAGCGATCTCGCCAAGACAG
fasRR	CTACGAGCGGGTAAGCGGGACG

References

1. Mondino S, Gago G, Gramajo H. 2013. Transcriptional regulation of fatty acid biosynthesis in mycobacteria. Mol Microbiol 89:372–387.

2. Gomez JE, Bishai WR. 2000. whmD is an essential mycobacterial gene required for proper septation and cell division. Proc Natl Acad Sci U S A 97:8554–8559.

3. Kolly GS, Boldrin F, Sala C, Dhar N, Hartkoorn RC, Ventura M, Serafini A, Mckinney JD, Manganelli R, Cole ST. 2014. Assessing the essentiality of the decaprenyl-phospho-d-arabinofuranose pathway in Mycobacterium tuberculosis using conditional mutants. Mol Microbiol 92:194–211.

4. Hanahan D. 1983. Studies on transformation of Escherichia coli with plasmids. JMolBiol 166:557–580.

5. Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon S V, Eiglmeier K, Gas S, Barry III CE, Tekaia F, Badcock K, Basham D, Brown D, Chillingworth T, Connor R, Davies R, Devlin K, Feltwell T, Gentles S, Hamlin N, Holroyd S, Hornsby T, Jagels K, Krogh A, McLean J, Moule S, Murphy L, Oliver K, Osborne J, Quail MA, Rajandream MA, Rogers J, Rutter S, Seeger K, Skelton J, Squares R, Squares S, Sulston JE, Taylor K, Whitehead S, Barrell BG. 1998. Deciphering the biology of Mycobacterium tuberculosis from the complete genome sequence. Nature 393:537–544.

A

Wildtype *fasR* →
Deleted *fasR* →

Deleted *fasR* →

3 4

B

RpoB

FasR

H37Rv (wild type *fasR*) expected size: 687 bp
pFR35 (mutant *fasR* allele) expected size: 150 bp

Figure S1. Analysis of the knockout *fasR* mutant (A) Colony PCR of *M. tuberculosis* transformants grown on Str, carrying the complementing *fasR* gene (clones 1 and 2) and of transformants grown on Hyg, carrying the empty pND255 vector (clones 3 and 4). Primers used for the amplification of *fasR* are listed in Table S3. The expected size of the PCR products is 687 bp for the *fasR* wild type allele and 150 bp for the mutant *fasR* allele. Two fragments were amplified from the Str-resistant clones, corresponding to the complementing *fasR* gene from the integrated plasmid and the deleted *fasR* gene at the *fasR* locus; on the other hand, only one fragment was amplified from the Hyg resistant clones (the partially deleted *fasR* gene) **(B)** Western Blot analysis of crude protein extracts prepared from cultures of wild type H37Rv strain (*Mtb*H37Rv), *fasR* conditional knockdown strain (*MtbΔfasR*-c*fasR*) and *fasR* KO strain (*MtbΔfasR* clones 3 and 4) grown at OD$_{600nm}$ ~ 0,3. Detection was performed using monoclonal anti-FasR$_{MT}$ antibodies (lower panel) and anti-RpoB as loading control (upper panel).

Lipid profile in negative ion mode: signal %			
	*MtbH*37Rv	*MtbΔfasR*	*MtbΔfasR-cfasR*	*p-value**
CL	36.37 ± 2.43	20.87 ± 3.34	31.52 ± 0.78	0.0234
PE	1.54 ± 1.43	2.31 ± 1.25	3.92 ± 1.23	0.624
PG	0.74 ± 0.11	0.82 ± 0.07	0.86 ± 0.25	0.4725
PI	16.71 ± 2.26	7.49 ± 0.09	16.45 ± 3.35	0.0234
PIMs	8.97 ± 2.43	6.01 ± 1.06	9.35 ± 2.74	0.1851429
MA alpha	7.54 ± 1.43	18.44 ± 0.61	6.7 ± 2.2	0.018
MA keto	1.54 ± 0.22	2.98 ± 0.42	1.5 ± 0.62	0.0234
MA methoxy	9.66 ± 2.3	16.78 ± 0.22	13 ± 8.58	0.03825
SL	0.89 ± 0.71	0.04 ± 0.03	0.16 ± 0.04	0.0234

* T-test *MtbΔfasR vs MtbH*37Rv. Benjamini-Hochberg adjusted *p-value*.

Lipid profile in positive ion mode: signal %			
	*MtbH*37Rv	*MtbΔfasR*	*MtbΔfasR-cfasR*	*p-value**
DAG	5.91 ± 1.37	1.84 ± 1.28	3.29 ± 0.19	0.0538
TAGs	67.15 ± 2.5	76.39 ± 0.61	72.78 ± 4.64	0.0246
MMDAG	2.43 ± 0.19	3.54 ± 0.25	2.57 ± 0.12	0.0246
GroMM	0.27 ± 0.13	0.03 ± 0.01	0.14 ± 0.047	0.0898
PDIM	23.1 ± 1.56	16.94 ± 0.24	20.22 ± 4.39	0.0246
TMM	0.02 ± 0.004	0.005 ± 0.001	0.1 ± 0.01	0.0246

* T-test *MtbΔfasR vs MtbH*37Rv. Benjamini-Hochberg adjusted *p-value*.

Figure S2. Relative abundance of lipid species. The tables show the relative abundance as the total signal percentage of the different lipid classes. CL, cardiolipin; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PI, phosphatidylinositol; PIMs, phosphatidylinositol mannosides; SL, sulfoglycolipids; MA, mycolic acids; DAG, diacylglycerol; TAGs, triacylglycerol; MMDAG, monomycolyl-DAG; GroMM, glycerol monomycolate; PDIM, phthiocerol dimycocerosates; TMM, trehalose monomycolate. Results are the mean of three independent experiments ± SD (n = 3). p values were calculated with unpaired *t* test with Benjamini-Hochberg's correction between *MtbΔfasR* and *Mtb*H37Rv.

A)

Lipid profile in negative ion mode: C – chain length			
	*MtbH*37Rv	*MtbΔfasR*	*MtbΔfasR-cfasR*	*p-value**
CL	68.71 ±0.02	68.65 ±0.14	68.95 ±0.05	0.017
PE	34.4 ±0.28	34.32 ±0.08	34.33 ±0.21	0.909
PG	34.2 ±0.11	34.17 ±0.17	34.3 ±0.03	0.219
PI	34.93 ±0.04	34.96 ±0.12	34.99 ±0.04	0.711
MA alpha	78.64 ±0.004	78.37 ±0.04	78.52 ±0.09	0.078
MA keto	84.64 ±0.03	83.65 ±0.28	84.3 ±0.08	0.016
MA methoxy	85.34 ±0.34	84.77 ±0.02	85.12 ±0.05	0.002
SL	151.04 ±0.27	144.01 ±1.61	148.84 ±0.72	0.013
Lipid profile in positive ion mode: C – chain length			
	*MtbH*37Rv	*MtbΔfasR*	*MtbΔfasR-cfasR*	*p-value**
DAG	35.49 ±0.09	36.82 ±0.96	35.28 ±0.06	0.031
GroMM	47.61 ±0.15	46.67 ±0.9	46.49 ±0.93	0.018
TAGs	55.87 ±0.21	58.65 ±0.43	56.34 ±0.33	0.003
TMM alpha	82.16 ±0.13	81.22 ±0.13	80.89 ±0.4	0.237
TMM keto	87.07 ±0.13	86.07 ±0.65	86.69 ±0.15	0.106
TMM methoxy	87.51 ±0.12	86.39 ±0.31	87.32 ±0.47	0.052
PDIM A	93.19 ±0.26	91.17 ±0.21	92.65 ±0.03	0.0005

* T-test *MtbΔfasR vs MtbH*37Rv. Benjamini-Hochberg adjusted *p-value*.

B)

C)

Figure S3. Detailed analysis of lipid species. (A) The tables show the weighted average of total number of carbons atoms per molecule within the population of mycolic acids (MA), phospholipids (PE, phosphatidylethanolamine; CL, cardiolipin; PG, phosphatidylglycerol; PI, phosphatidylinositol); sulfoglycolipids (SL); DAG, diacylglycerol; GroMM, glycerol monomycolate; TAGs, triacylglycerol; TMM trehalose monomycolate and phthiocerol dimycocerosates (PDIM A). The histograms show the distribution of the number of carbon atoms of SL **(B)** and PDIM **(C)** present in the wild type strain (*Mtb*H37Rv), the *fasR* complemented strain (*MtbΔfasR-cfasR*) and *fasR* mutant strain (*MtbΔfasR*). Results are the mean of three independent experiments ± SD (n = 3). p values were calculated with unpaired *t* test with Benjamini-Hochberg's correction between *MtbΔfasR* and *Mtb*H37Rv.

Figure S4. Saturation level of the fatty acid substituents of the phospholipids. Histograms represent the levels of unsaturation of the fatty acids constituents of cardiolipin (CL), phosphatidylglycerol (PG) and phosphatidylinositol (PI). Results are the means of three independent experiments.