Structural basis of client specificity in mitochondrial membrane-protein chaperones

Iva Sučec[1], Yong Wang[2,✉], Ons Dakhlaoui[1], Katharina Weinhäupl[1,5], Tobias Jores[3,6], Doriane Costa[1], Audrey Hessel[1,7], Martha Brennich[4,8], Doron Rapaport[3], Kresten Lindorff-Larsen[2], Beate Bersch[1,✉], and Paul Schanda[1,✉]

[1]Univ. Grenoble Alpes, CEA, CNRS, Institut de Biologie Structurale (IBS), 71, Avenue des Martyrs, F-38044 Grenoble, France
[2]Structural Biology and NMR Laboratory, the Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen, 2200 Copenhagen, Denmark
[3]Interfaculty Institute of Biochemistry, University of Tübingen, 72076 Tübingen, Germany
[4]European Molecular Biology Laboratory, 38042 Grenoble, France
[5]Present address: Universidade do Porto, I3S - Instituto de Investigacao e Inovacao em Saude, Porto, Portugal
[6]Present address: Department of Genome Sciences, University of Washington, Seattle, WA, USA
[7]Present address: Institute of Pharmacology and Structural Biology, 31077 Toulouse, France
[8]Present address: Bruker Optics, 76275 Ettlingen, Germany

Chaperones are essential for assisting protein folding, and for transferring poorly soluble proteins to their functional locations within cells. Hydrophobic interactions drive promiscuous chaperone–client binding, but our understanding of how additional interactions enable client specificity is sparse. Here we decipher what determines binding of two chaperones (TIM8·13, TIM9·10) to different integral membrane proteins, the all-transmembrane mitochondrial carrier Ggc1, and Tim23 which has an additional disordered hydrophilic domain. Combining NMR, SAXS and molecular dynamics simulations, we determine the structures of Tim23/TIM8·13 and Tim23/TIM9·10 complexes. TIM8·13 uses transient salt bridges to interact with the hydrophilic part of its client, but its interactions to the transmembrane part are weaker than in TIM9·10. Consequently, TIM9·10 outcompetes TIM8·13 in binding hydrophobic clients, while TIM8·13 is tuned to few clients with both hydrophilic and hydrophobic parts. Our study exemplifies how chaperones fine-tune the balance of promiscuity *vs.* specificity.

protein import | NMR spectroscopy | small-angle X-ray scattering | mitochondrial biogenesis | molecular dynamics

Correspondence: *yong.wang@bio.ku.dk* *beate.bersch@ibs.fr* *paul.schanda@ist.ac.at*

Introduction

Cellular survival and function fundamentally rely on an intact proteome. Proteins within cells need to be correctly folded to their functional conformation, and be present at the cellular location where they function. Chaperones play a central role in maintaining this cellular protein homeostasis (1), by either helping other proteins to reach their functional three-dimensional structure after synthesis, by transporting them across the cytosol or organelles, or by sustaining their native fold along their lifetime. More than 20,000 different proteins are required to fulfill the functions of human cells, and it is believed that the majority rely on chaperones to reach and maintain their native fold (2). Given the diversity of the client proteins, many chaperones promiscuously interact with tens of different 'client' proteins that may differ widely in size, structure and physico-chemical properties. However, the need for efficient binding and refolding of their clients also calls for some degree of specificity. Chaperones operate at this delicate balance of promiscuity and specificity to their clients. The interactions that determine the chaperone–client specificity are only partly understood.

Hydrophobic interactions play a crucial role for chaperone interactions as most chaperones bind to hydrophobic patches on their clients and shield them from aggregation. Electrostatic charges also play a role in some chaperone complexes (3). The interaction motifs recognized by different chaperones differ by their physico-chemical properties (4). For example, for interacting with the Hsp70 chaperone family, Ile, Phe, Leu and Val residues are particularly important (5, 6); the SecB chaperone recognizes 9-residue long stretches enriched in aromatic and basic residues (7); the chaperone Spy uses longer-range charge interactions for the formation of an initial encounter complex, followed by more tight binding mediated by hydrophobic interactions, (8) whereby structurally frustrated sites on the client protein are particularly prone to binding (9).

Our understanding of the underlying principles of chaperone-client interactions is hampered by the lack of atomic-level views onto the structure and dynamics of these complexes. Their inherently dynamic and often transient nature represents a significant experimental challenge towards structural characterization. Only a very limited number of chaperone complex structures have been reported (reviewed in (10)). The modes of interactions that they revealed range from rather well-defined binding poses of client polypeptides in the chaperone's binding pockets, reminiscent of complexes formed by globular proteins, to highly flexible ensembles of at least partly disordered conformations ('fuzzy complexes'). In the latter, a multitude of local chaperone-client interactions may result in a high overall affinity despite the low affinity and short life time of each individual inter-molecular contact.

Multiple molecular chaperones are present in the cell with mutually overlapping functions and 'clientomes' (2, 11, 12). It is poorly understood, however, whether a given client protein adopts a different conformation (or ensemble of conformations) when it is bound to different chaperones, and if different clients, when bound to a given chaperone, all show similar conformational properties. α-Synuclein appears to have similar interaction patterns with six different chaperones (13); outer-membrane proteins (OmpA, OmpX, FhuA)

have similar properties – essentially fully unfolded – when bound to SurA and Skp chaperones (14, 15), at least as judged by their NMR fingerprint spectra. Phosphatase A displays an extended dynamic conformation, but well-defined binding poses of its interacting parts, when bound to trigger factor (16), Hsp40 (17) or SecB (18). Thus, while these reports suggest that a given protein adopts similar properties on different chaperones, the scarcity of data and absence of a direct comparison of complex structures, leaves open which interactions may confer specificity.

A pair of 'holdase' chaperone complexes of the mitochondrial inter-membrane space (IMS), TIM8·13 and TIM9·10, are structurally highly similar, but have different substrate binding preferences. These chaperones transport precursors of membrane proteins (henceforth denoted as 'precursors') to the membrane-insertase machineries in the inner membrane (TIM22) and outer mitochondrial membranes (SAM) (19)). The TIM chaperones form hetero-hexameric structures of ca. 70 kDa, composed of an alternating arrangement of Tim9 and Tim10 or Tim8 and Tim13. TIM9·10 is essential to cellular viability (20–22); even single point mutations in Tim9 or Tim10 that keep the chaperone structure intact but affect precursor protein binding can impair yeast growth and cause lethality (23). Although TIM8·13 is not essential in yeast (24), yeast cells depleted of Tim8 and Tim13 show conditional lethality (25). Additionally, mutations in the human Tim8a protein have been identified as the cause of a neurodegenerative disorder known as Mohr-Tranebjærg (MTS) or Deafness-Dystonia-Optic Neuropathy (DDON) syndrome (26, 27).

In vivo experiments, predominantly in yeast, have identified mitochondrial membrane proteins whose biogenesis depends on small TIM chaperones. TIM9·10 is believed to interact with all members of the mitochondrial carrier (SLC25) family, which comprises more than 50 members in humans, such as the ADP/ATP carrier (Aac in yeast); TIM9·10 furthermore transports the central components of the TIM22 and TIM23 insertion machineries (Tim23, Tim17, Tim22) as well as outer-membrane β-barrel proteins (28). TIM8·13 has a narrower clientome, and was shown to bind the precursors of the inner-membrane proteins Tim23 (25, 29, 30) and Ca^{2+}-binding aspartate-glutamate carriers (31), as well as the outer-membrane β-barrel proteins VDAC/Porin, Tom40 (32) and Tob55/Sam50 (33). There is evidence that TIM8·13 does not bind the inner-membrane proteins ADP/ATP carrier (Aac) and Tim17 (25). The inner-membrane proteins that have been reported to interact with TIM8·13 have a hydrophilic domain in addition to trans-membrane domains (fig. S1) but this does not hold true for the outer-membrane β-barrels. Thus, the mechanisms by which TIM8·13 binds its clients remain unclear.

Recently, we obtained the first structure of a complex of a small TIM chaperone, TIM9·10, with the mitochondrial GDP/GTP carrier (Ggc1) (23). The structure, composed of two chaperone complexes holding one precursor protein, revealed a highly dynamic ensemble of Ggc1 conformers that form multiple short-lived and rapidly inter-converting (< 1

ms) interactions with a hydrophobic binding cleft of the chaperone (fig. S2). The TIM9·10-Ggc1 complex can be described as a "fuzzy complex", in which the high overall affinity is driven by a multitude of individually weak interactions with the hydrophobic trans-membrane (TM) parts of its clients.

To understand what confers specificity in the mitochondrial IMS chaperone system, we studied chaperone complexes of TIM9·10 and TIM8·13 with two precursor proteins, the GDP/GTP carrier (Ggc1) and the insertase component Tim23. In their native state, Ggc1 comprises six TM helices without soluble domains, and Tim23 four TM helices and a ca. 100-residue-long soluble inter-membrane space domain (Fig. 1A). By solving the complex structures of the two chaperone complexes holding Tim23, we reveal that the differential specificity of the two chaperones is based on an interplay of hydrophobic and hydrophilic interactions, which leads to different conformational properties of the precursor protein bound to these chaperones.

Results

TIM8·13 and TIM9·10 interact differently with membrane precursor proteins. We have developed an experimental protocol (23) to prepare complexes of the inherently insoluble membrane-protein precursors and chaperones (Fig. 1B,C). Briefly, the approach involves the recombinant production of His-tagged precursor protein, binding it to a His-affinity column in denaturing conditions, followed by removal of the denaturant and simultaneous addition of a chaperone. The chaperone-precursor complex is then eluted for further biochemical, biophysical and structural investigations.

The measurement of dissociation constants of chaperones and membrane precursor proteins, using methods such as isothermal titration calorimetry or surface plasmon resonance, is not possible, because the complexes cannot be formed in solution (e.g. flash-dilution methods, which work for other chaperones (14), failed; data not shown). Thus, to characterize the relative affinities of the precursor proteins to the two chaperones, we performed different types of competition experiments. In a first experiment, precursor protein was bound to the affinity resin, and both chaperones were simultaneously added, before washing excess chaperone, and eluting the chaperone-precursor complexes (Fig. 1C). NMR spectroscopy shows that the two chaperones do not form mixed hetero-hexameric complexes, implying that TIM9·10 and TIM8·13 stay intact in such competition experiments (fig. S3). In a second class of experiments, we prepared one type of complex (e.g. TIM9·10-Tim23) and added the other chaperone (e.g. TIM8·13) in its apo state, allowing the precursor protein to be transferred. These experiments also demonstrate that membrane precursor proteins can be transferred between these two chaperones, on the time scale we investigated (minutes to hours). We used SDS-PAGE analyses and electrospray ionization mass spectrometry (ESI-MS) to systematically quantify the amount of obtained complexes (Fig. 1D and fig. S4). Consistently, we find that Ggc1 has

Fig. 1. Biochemical characterization of TIM chaperone - membrane protein complexes. (A) Native topology of the two precursor proteins used in this study. (B) Schematic view of the pull-down experiment used to prepare chaperone-precursor complexes. (C) Formation of Tim23-chaperone complexes, monitored by SDS-PAGE. Either TIM8·13, TIM9·10 or a 1:1 mixture was added to NiNTA-bound Tim23. The lanes correspond to flow-through after applying chaperone (FT), additional wash (W), and imidazole elution (E). Protein bands corresponding to Tim10 and Tim8 overlap. (D) Relative amounts of complexes with Ggc1 and Tim23, obtained from three different experiments: (i) a pull-down assay where both chaperones were applied to bound precursor protein (black), (ii) preparation of a TIM9·10-precursor protein complex and addition of TIM8·13, and SDS-PAGE and mass spectrometry analysis after 1 and 3 hours (red), (iii) preparation of TIM8·13-Tim23 followed by TIM9·10 addition and SDS-PAGE (blue) as in (ii). The protein amounts were determined from LC/ESI-TOF-MS (fig. S4); error estimates from ≥ two experiments. (E) Sequence alignment of the small Tims, numbered from the N- and C-terminal conserved Cys residues ("0"). Red: conserved hydrophobic residues; blue: hydrophilic Tim8 residues, K30 and S36L. See fig. S5 for comprehensive alignment. (F) Location of the residues in the hydrophobic cleft. (G) Comparison of Kyte-Doolittle hydrophobicity of the residues in the binding cleft of wild-type (WT) native Tim proteins and Tim8$_{K30F,S36L}$. (H) Pull-down experiment of Ggc1 with TIM9·10, TIM8·13 and TIM8·13(Tim8$_{K30F,S36L}$). Lane descriptions are as in (C); additionally, the fraction obtained after final wash with urea and imidazole, to control the Ggc1 initially loaded onto the column, is shown (control, C). (I) Amount of complex obtained from pull-down experiments of WT and mutant chaperones; the same amount of Ggc1 was applied in all three experiments, and the total amount of eluted complex was determined spectroscopically.

a strong preference for TIM9·10 (ca. 5- to 10-fold), while Tim23 shows a slight preference for TIM8·13 (ca. 1.5-fold). Taken together, we established that the two chaperones bind with different affinities to two inner-membrane precursor proteins, whereby TIM8·13 is barely able to hold Ggc1, in contrast to TIM9·10, while it can hold Tim23 slightly better than TIM9·10.

The small TIM chaperones use a conserved hydrophobic cleft for membrane precursor protein binding. To understand the different binding properties, we performed

a sequence alignment of the small TIMs, which reveals a well conserved set of hydrophobic residues that point towards the binding cleft formed between the inner (N-terminal) and outer tentacles (23) (Fig. 1E,F). The overall hydrophobicity of these residues is lower in Tim8 and Tim13 than in Tim9 and Tim10 (Fig. 1G). In particular, Tim8 has a charged residue in position -14 (Lys30). (The sequences are numbered starting with negative numbering at the twin CX$_3$C motif towards the N-terminus, and positive numbering from the last Cys to the C terminus and the hydrophobic motif residues are at positions -20, -14, -11, -8, -7, +5, +11 and +15.) This

positive charge at position -14, either Lys or Arg, is conserved among eukaryotes (fig. S5). In yeast, position -8 of the hydrophobic motif is polar (Ser36), although this position is not strictly conserved. Overall, these residues make the hydrophobic binding cleft of Tim8 less hydrophobic than in the other small Tims.

We speculated that the less hydrophobic nature of TIM8·13's binding cleft reduces its affinity to TM parts of membrane precursor proteins. To test this hypothesis, we generated a mutant TIM8·13 with increased hydrophobicity (Tim8$_{K30F,S36L}$; Fig. 1G). This more hydrophobic TIM8·13(Tim8$_{K30F,S36L}$) chaperone allowed us to obtain significantly larger amounts of complex with Ggc1 than native TIM8·13, under otherwise identical conditions (Fig. 1H,I). This observation establishes the importance of the hydrophobic cleft for binding hydrophobic TM parts of precursor proteins. Equivalent experiments with the full-length Tim23, shown in fig. S6, reveal that the additional hydrophobic residues in the binding cleft of Tim8$_{K30F,S36L}$ do not improve its capacity to bind Tim23. This observation suggests that the binding mechanisms in place for binding these two different membrane-protein clients differ.

To better understand the client-binding properties of the two chaperones, we turned to structural studies. Solution-NMR spectra of apo TIM8·13 (Fig. 2A) and residue-wise resonance assignments allowed identifying the residues forming secondary structure and estimating their local flexibility. In agreement with the crystal structure, the core of rather rigid tentacles comprises the top part of the chaperone between the CX$_3$C motifs and ca. 15-25 residues before and after these motifs. About 10-20 residues on each the N- and C-termini are flexible (fig. S7). To probe the binding of a trans-membrane segment of a membrane precursor protein, we performed NMR-detected titration experiments of TIM8·13 with a cyclic peptide corresponding to the two C-terminal strands of the β-barrel voltage-dependent anion channel (VDAC$_{257-279}$) that has a propensity to form a β-turn (35). Addition of this cyclic VDAC$_{257-279}$ induces chemical-shift perturbations (Fig. 2B), that are primarily located in the hydrophobic cleft formed between the inner and the outer rings of helices (Fig. 2C,D). This binding site matches very closely the site on TIM9·10 to which VDAC$_{257-279}$ binds (23) (Fig. 2C). Interestingly, the VDAC$_{257-279}$-induced chemical-shift perturbation (CSP) effects in TIM8·13 are overall only about half of the magnitude of CSPs found in TIM9·10, pointing to a higher population of TIM9·10-VDAC$_{257-279}$ complex compared to TIM8·13-VDAC$_{257-279}$ at comparable conditions (Fig. 2C). This finding suggests a lower affinity of TIM8·13 to VDAC$_{257-279}$, as expected from its lower hydrophobicity.

Photo-induced cross-linking experiments of a Bpa-modified VDAC$_{257-279}$ peptide to TIM8·13 show that only the cyclic peptide forms cross-linking adducts while the linear, mostly disordered (35) form does not (Fig. 2F). The same behavior was also found for TIM9·10 (23) and yeast cytosolic chaperones Ssa1, Ydj1, Djp1, and Hsp104 (36). A rationale for this finding is the fact that in a β-turn the side chains of consecutive residues point to the two opposing faces thus creating one hydrophobic and one more hydrophilic face (Fig. 2E). In contrast, due to its disorder, the linear VDAC$_{257-279}$ peptide does not have a stable hydrophobic face, reducing its affinity to the hydrophobic binding cleft on the chaperone. In line with these findings, NMR titration data with the linear peptide shows small CSPs that are spread across the protein, thus pointing to unspecific interaction (fig. S8).

Collectively, the experiments with the client fragment VDAC$_{257-279}$ provide a first evidence that both chaperones use the same conserved binding cleft to interact with hydrophobic membrane precursor protein sequences, and that TIM9·10 interacts more efficiently with TM parts, and thus with Ggc1 and the VDAC fragment. We propose that the more hydrophobic nature of the binding cleft in TIM90·10 allows it to interact more strongly with TM parts of its clients. In light of this observation, how does TIM8·13 achieve a binding affinity to Tim23 which is slightly higher than the one of TIM9·10 (Fig. 1D)?

Hydrophilic fragments interact differently with TIM8·13 and TIM9·10. Tim23 has a hydrophilic N-terminal segment in addition to four TM helices (Fig. 3A), and we investigated whether this part interacts with the chaperones. NMR spectra of the soluble Tim23$_{IMS}$ fragment (residues 1-98) in isolation show the hallmark features of a highly flexible intrinsically disordered protein with low spectral dispersion of ^{1}H-^{15}N NMR signals (Fig. 3B,C, orange spectrum), as previously reported (37). Upon addition of TIM9·10, the Tim23$_{IMS}$ ^{1}H-^{15}N spectrum (Fig. 3B, left) shows only small changes: all cross-peaks are still detectable, and small chemical-shift perturbations (CSPs) are only observed for a few residues at the N-terminus, which has higher hydrophobicity (Fig. 3D). This finding suggests only very weak, possibly non-specific interactions between the very N-terminus of Tim23$_{IMS}$ and TIM9·10. In line with this finding, the interaction is not detectable by isothermal titration calorimetry (ITC) measurements (Fig. 3H).

The interaction of the hydrophilic Tim23$_{IMS}$ fragment with TIM8·13 is significantly stronger, with pronounced binding effects detected by ITC, and a dissociation constant of K$_d$ = 66 $\pm$ 8 μM (Fig. 3H, right panel; see Table S1). The ^{1}H-^{15}N NMR spectrum of Tim23$_{IMS}$ in the presence of TIM8·13 shows strongly reduced peak intensities for the majority of the residues (Fig. 3C, left). Such a peak broadening is expected when a highly flexible polypeptide binds to a relatively large object such as TIM8·13, thereby inducing faster nuclear spin relaxation and thus broader signals of lower magnitude. Analysis of the peak-intensity reduction reveals two regions of Tim23 that are particularly involved in the binding: (i) the N-terminal hydrophobic residues, which are also involved in interacting with TIM9·10, and (ii) a long sequence stretch comprising residues from ca. 30 to 80 (Fig. 3E). We also investigated whether TIM8·13 may interact with soluble proteins from the inter-membrane space, and performed ITC experiments with the globular protein cytochrome c. No interaction could be detected (fig. S9), suggesting that the TIM8·13-Tim23$_{IMS}$ interaction may be re-

Fig. 2. Solution-NMR and binding of a VDAC fragment to TIM8·13. (A) ^{1}H-^{15}N NMR spectrum of TIM8·13 at 35 °C. **(B)** Chemical-shift perturbation (CSP) in TIM8·13 upon addition of 5 molar equivalents of cyclic VDAC$_{257-279}$. **(C)** CSP effects of VDAC$_{257-279}$ binding. The data for TIM9·10 are from ref. (23). **(D)** Plot of CSP data on the TIM8·13 structure. **(E)** Schematic structure of the two last strands of VDAC, as found in the NMR structure (34) of the full β-barrel, showing that the hydrophobic and hydrophilic side chains cluster on the two opposite faces of the β-turn. **(F)** Photo-induced cross-linking of the linear (left) and cyclic (right) VDAC$_{257-279}$ peptides to TIM8·13. While hardly any adducts are observed for the linear one, the cyclic peptide forms cross-linking photo-adducts (PA), including of higher molecular weight, resulting from multiple cross-links, as reported earlier (23, 35, 36).

lated to the unfolded, flexible character of the latter.

To characterize the conformation of full-length (FL) Tim23 bound to TIM8·13 and TIM9·10, we prepared Tim23$_{FL}$-labeled Tim23–chaperone complexes using the method outlined in Fig. 1A. Very similar to the experiments with the Tim23$_{IMS}$ fragment, the signals corresponding to the N-terminal half of Tim23 are still intense in the Tim23-TIM9·10 complex (Fig. 3B,F). The small observed CSPs are localized primarily at the ten N-terminal residues. In contrast, when Tim23$_{FL}$ is bound to TIM8·13, the signals corresponding to its N-terminal half are severely reduced in intensity, revealing tight contact of the flexible N-terminal half of Tim23 to TIM8·13 (Fig. 3C,G).

In neither of the two Tim23$_{FL}$ complexes any additional signals, that may correspond to Tim23's TM helices, are visible. We ascribe this lack of detectable signals of residues in the TM part to extensive line broadening. The origin of this line broadening may be ascribed to the large size of the complex and likely to additional millisecond (ms) time scale dynamics of Tim23's TM parts in the hydrophobic binding cleft of

the chaperones. Such millisecond motions have been found in the TIM9·10–Ggc1 complex (23).

TIM8·13 uses an additional hydrophilic face for protein binding. We probed the binding sites that the chaperones use to interact with Tim23$_{IMS}$ or Tim23$_{FL}$ using NMR spectroscopy on samples in which only the chaperone was isotope-labeled. Interestingly, the CSPs in the two chaperones upon addition of Tim23$_{IMS}$ reveal distinct binding patterns (Fig. 4A): in TIM8·13, the largest effects involve residues in the hydrophilic top part of the chaperone, between the CX$_3$C motifs, as well as a few residues toward the C-terminal outer ring of helices; in contrast, the corresponding top part of TIM9·10 does not show any significant effects, but CSPs are observed at residues in the hydrophobic binding cleft, and in particular the N-terminal helix (Fig. 4B). This data, together with the Tim23$_{IMS}$-detected data in Fig. 3 establish that TIM8·13 uses its hydrophilic top part to bind Tim23's N-terminal half, while only a short stretch of hydrophobic residues at the very N-terminus of Tim23 inter-

Fig. 3. Tim23 has markedly different properties when binding to TIM8·13 and to TIM9·10. (A) Hydrophobicity of Tim23 (Kyte-Doolittle). **(B)** NMR spectra of the ^{15}N-labeled soluble Tim23$_{IMS}$ fragment in the presence of TIM9·10 (left, black), and of full-length Tim23 bound to TIM9·10 (right, black) are compared to the Tim23$_{IMS}$ fragment in isolation (orange), under identical buffer conditions and NMR parameters. **(C)** As in (B) but with TIM8·13 instead of TIM9·10. **(D)** Chemical-shift perturbation (CSP) of residues in Tim23$_{IMS}$ upon addition of 1 (light orange) or 5 (dark orange) molar equivalents of TIM9·10. **(E)** Intensity ratio of residues in Tim23$_{IMS}$ in the presence of 4 molar equivalents of TIM8·13 compared to Tim23$_{IMS}$ alone. **(F)** CSP of the detectable residues in full-length Tim23 attached to TIM9·10 (brown), compared to the soluble Tim23$_{IMS}$ fragment. **(G)** Intensity ratio of detectable residues in Tim23$_{FL}$ attached to TIM8·13. Note that the ratio was not corrected for differences in sample concentration, and the scale cannot be compared to the one in panel (E). **(H)** Calorimetric titrations for the interaction of TIM9·10 or TIM8·13 (54 µM in the calorimetric cell) with Tim23$_{IMS}$ (1.15 mM in the injecting syringe). Thermograms are displayed in the upper plots, and binding isotherms (ligand-normalized heat effects per injection as a function of the molar ratio, [Tim23$_{IMS}$]/[chaperone]) are displayed in the lower plots. Control experiments, injecting into a buffer, are shown in blue.

Fig. 4. Tim23$_{IMS}$ and full-length Tim23 differ in their interactions with TIM9·10 and TIM8·13 chaperones. **(A)** Chemical-shift perturbations observed upon addition of the Tim23$_{IMS}$ fragment to TIM9·10 (top) and TIM8·13 (bottom). The chaperone:Tim23$_{IMS}$ ratios were 1:1 (TIM8·13) and 1:3 (TIM9·10). Mapping of Tim23$_{IMS}$-induced CSPs on TIM9·10 **(B)** and TIM8·13 **(C)**, showing that while the top part of TIM9·10 does not show any significant CSPs, the corresponding part is the main interacting region of TIM8·13. CSP in complexes of TIM9·10 **(D)** and TIM8·13 **(E)** bound to full-length Tim23. Tim23$_{TM}$-induced CSP mapped on TIM9·10 **(F)**, showing similar binding as the full-length Tim23.

acts with the hydrophobic cleft of TIM9·10, which is also the binding site of TM parts (Figs. 1 and 2).

Chaperone-labeled complexes with Tim23$_{FL}$ confirm these findings, and point to the additional effects induced by the bound TM part: in TIM9·10-Tim23$_{FL}$, large CSP effects are located primarily in the binding cleft, in line with the view that the top part of TIM9·10 is not involved in binding Tim23. In contrast, Tim23$_{FL}$-induced CSPs are found across the whole TIM8·13, including the hydrophilic top and the hydrophobic cleft (Fig. 4C and fig. S10).

We furthermore prepared complexes of a truncated Tim23 fragment (Tim23$_{TM}$, residues 92-222), which allows to detect selectively the interaction of the trans-membrane part with the chaperones. The TIM9·10-Tim23$_{TM}$ complex features the largest CSPs in the hydrophobic binding cleft, qualitatively similar to the binding site detected with Tim23$_{FL}$ (Fig.

4F and fig. S11). The complex of Tim23$_{TM}$ with TIM8·13 appears to be much less stable than TIM9·10-Tim23$_{TM}$: in the pull-down experiment, only a very small amount of complex could be obtained, and the complex rapidly precipitated (not shown). This observation reflects that the hydrophobic cleft of TIM8·13 is less capable of holding a hydrophobic polypeptide than the one of TIM9·10.

Collectively, NMR, ITC and mutagenesis have revealed that the hydrophobic cleft of both TIM8·13 and TIM9·10 are essential to hold the hydrophobic parts of the clients, and that TIM8·13, but not TIM9·10, additionally interacts with the hydrophilic part of Tim23 to increase its affinity. This interaction, which is mediated by the hydrophilic top part of TIM8·13, reduces the conformational flexibility of Tim23's N-terminal half. This hydrophilically driven interaction supports previous observation during the protein import (25),

where TIM8·13 interacting with hydrophobic membrane precursor without long hydrophilic stretches (fig. S1) was detected only when they were fused to hydrophilic Tim23$_{IMS}$. We have also investigated whether a given full-length Tim23 chain may interact simultaneously with TIM9·10 and TIM8·13, using hydrophobic and hydrophilic interactions, respectively, thus forming a ternary complex. However, samples containing all three components do not contain detectable amounts of such complexes, suggesting that the affinity to simultaneously bind two chaperones is too low (fig. S12).

Structural ensembles of chaperone-Tim23 complexes.
We integrated the NMR data with further biophysical, structural and numerical techniques to obtain a full structural and dynamical description of the complexes. We first investigated the complex stoichiometry using size-exclusion chromatography coupled to multi-angle light scattering (SEC-MALS), NMR-detected diffusion-coefficient measurements, and analytical ultra-centrifugation. These methods, which provide estimates of molecular mass (and shape) from orthogonal physical properties (gel filtration and light scattering; translational diffusion), reveal properties best compatible with a 1:1 (chaperone:precursor) stoichiometry (Fig. 5A and fig. S13). Small-angle X-ray scattering (SAXS) data of both TIM9·10-Tim23 and TIM8·13-Tim23 also point to a molecular weight corresponding to a 1:1 complex (SAXS; Fig. 5B). This stoichiometry contrasts the 2:1 (chaperone:precursor) stoichiometry for TIM9·10 holding the 35 kDa-large carrier Ggc1 (23) (fig. S2).

Importantly, SAXS provides significantly more information, namely the overall shape of the ensemble of conformations present in solution. Given the flexibility of the complex, this SAXS data is best analyzed by considering explicitly a dynamic ensemble. We used molecular dynamics (MD) simulations to account for the breadth of possible conformations that, collectively, result in the observed scattering. To effectively sample the conformational space of the chaperone-Tim23 complex, we constructed two distinct structural models in which the N-terminal half of Tim23 is either modeled as a floppy unstructured tail or bound to the hydrophilic upper part of chaperone, denoted as 'N-tail unbound' and 'N-tail bound' conformations, respectively. In both models, the hydrophobic C-terminal transmembrane domain of Tim23 is bound to the hydrophobic cleft of the chaperone, as identified by NMR (Fig. 4 and fig. S10C,D) (23). Initiating from both conformations, explicit-solvent atomistic MD simulations (∼ 4.25 μs in total) were performed to collect the structures for the 'N-tail unbound' and 'N-tail bound' ensembles. In the case of TIM8·13, the 'N-tail bound' ensemble recapitulates the experimentally observed pattern better than the 'N-tail unbound' ensemble (Fig. 3 C and E). We then constructed a mixed ensemble consisting of a mixture of 'N-tail bound', and 'N-tail unbound' states. We used this pool of conformations for further ensemble refinement, with the relative populations of these two ensembles of states are free parameter, using the Bayesian Maximum Entropy (BME) method guided by experimental SAXS data (23, 38, 39). We found

that the experimental SAXS data of TIM8·13–Tim23 are very well reproduced when the mixed ensemble has >85 % of the 'N-tail bound' state (Fig. 5C, D). In contrast, the experimental data of TIM9·10–Tim23 are only well reproduced when the TIM9·10–Tim23 ensemble comprises predominantly the 'N-tail unbound' state. These refined ensembles guided by experimental SAXS data are in excellent agreement with the NMR data, which showed that (i) in the TIM9·10-Tim23 complex, the N-terminal part of Tim23 is predominantly free and flexible, and Tim23 makes contacts only to the hydrophobic cleft of the chaperone, while (ii) in TIM8·13-Tim23, the Tim23$_{Nter}$ part is largely bound to the upper part of the chaperone (Figs. 3 and 4).

The amount of 'N-tail bound' relative to 'N-tail unbound' states is expected to depend on the affinity of the N-tail of Tim23 to the chaperone. Indeed, the ITC-derived TIM8·13-Tim23$_{IMS}$ affinity (K_d=66 μM; Fig. 3H) predicts that the population of N-tail-bound states is of the order of 75 - 98 % (see Methods for details), in excellent agreement with the MD/SAXS derived value (> 85 %). This good match of data from the Tim23$_{IMS}$ fragment and Tim23$_{FL}$ suggests that the binding of Tim23's hydrophilic N-tail does not strongly depend on the presence of the TM parts. The low affinity of the N-tail to TIM9·10, reflected by the inability to detect TIM9·10-Tim23$_{IMS}$ binding by ITC, is mirrored by the small population of the 'N-tail bound' states in the full-length complex.

To identify the molecular mechanisms underlying the observed differences in N-tail binding, we studied the interactions formed between Tim23 and the chaperones along the MD simulation. An interesting pattern emerges from analysis of the electrostatic interactions. The top part of TIM8·13 has predominantly polar and negatively charged residues, which are in transient contact with the positive charges of Tim23 N-tail, within a dynamic ensemble of conformations (Fig. 6). For example, three key aspartate or glutamate residues in TIM8·13 appear to be involved in binding of lysine or arginine residues of Tim23$_{IMS}$. In Tim9, a lysine (K51) is present in the top part, and contributes a positive charge (the equivalent position in TIM8·13 is a non-charged, polar residue). We hypothesized that the less complementary electrostatic properties of TIM9·10's top part and Tim23's N-tail, as compared to TIM8·13, may diminish the affinity of the N-tail to TIM9·10. We attempted to investigate the importance of these charged residues experimentally, and prepared single- and double-mutants that alter the pattern of charged residues. In TIM8·13, we introduced lysine or arginine instead of negatively charged residues, expecting to reduce thereby the affinity to Tim23$_{IMS}$; conversely, in TIM9·10 we introduced negative charges to promote the Tim23$_{IMS}$ interaction. However, ITC experiments show that most of these mutants do not significantly differ in their binding affinity to Tim23$_{IMS}$ (fig. S9 and table S1). In one of the TIM8·13 mutants, the binding affinity even increases, despite the additional positive charge in the chaperone. These findings suggest that due to the disordered nature of Tim23's N-tail, its binding with the chaperones might not be dominated by a few strong interactions

Sučec *et al.* | Client specificity of mitochondrial TIM chaperones

Fig. 5. Architecture of the TIM8·13 and TIM9·10 holdases in complex with full-length Tim23. (A) (Left) Apparent molecular weights of apo and holo chaperone complexes from SEC-MALS, and AUC (red) circles. (Right) Translational diffusion coefficients of TIM9·10 (apo) and TIM9·10-Tim23$_{FL}$ from NMR DOSY measurements. Two independent samples were used for the complex, in which either the chaperone or the precursor protein was labeled, as indicated. See also fig. S13. **(B)** Small-angle X-ray scattering curves (top) and Kratky plot representations thereof for the two chaperone-precursor complexes. The lines are SAXS curves calculated from structural ensembles obtained over 4.25 μs long MD trajectories, in which the N-terminal half of Tim23 was either in a conformation bound to the top part of the chaperone (red) or in a loose unbound conformation (blue), or from an ensemble in which these two classes of states were present with optimized weights. **(C)** Goodness of fit of the back-calculated SAXS curves to the experimental SAXS data as a function of the relative weights of the two classes of conformations (bound/unbound). **(D)** Snapshots of conformations in which Tim23$_{N-tail}$ is either bound or unbound, and the best-fit relative weights of the two classes of states as derived from SAXS/MD. More SAXS/MD data and ensemble views are provided in fig. S14, and in Supplementary Movies 1 and 2.

but instead be contributed by a complex interaction network with many weak and widely distributed interactions which are tolerant to introduction of the single point mutants as we explored. The MD ensemble due to its limited time scale and force field imperfections (40) may only be able to identify a rather small number of key interaction sites. Interestingly, for the binding of the TM part of precursors, we were able to identify several single-point mutations in the TIM9·10 hydrophobic motif that abrogate the binding, with strong phenotype (23).

Discussion

Transfer chaperones (holdases) need to fulfill two contradicting requirements, holding their clients very tightly to avoid their premature release and aggregation, while at the same time allowing release at the downstream factor. This apparent contradiction is solved by a subtle balance of multiple individually weak interactions, and a resulting dynamic complex, wherein the precursor protein samples a wide range of differ-

ent conformations. This ensemble of conformations results in a high overall affinity, yet a downstream foldase/insertase can detach the precursor protein from the chaperone without significant energy barrier (41). Balancing the interaction strengths is, thus, crucial to chaperone function. Herein, we have revealed a fine-tuning of chaperone-client specificity that involves hydrophobic interactions with the chaperone's binding cleft and additional hydrophilic interactions, mostly mediated by charged residues, with the chaperone's top part. Lower hydrophobicity within the binding cleft of TIM8·13 compared to TIM9·10 arises by overall less hydrophobic residues and a positively charged residue (Lys/Arg) that is highly conserved in Tim8. As a consequence, TIM8·13 is less able to hold the TM parts of its clients than TIM9·10 by ca. one order of magnitude. As we showed, replacement of two charged/polar side chains in TIM8·13's cleft brings TIM8·13 to a similar level as TIM9·10 for holding an all-transmembrane client.

For binding of its native client Tim23, TIM8·13 uses addi-

Fig. 6. Tentative identification of electrostatic interactions from the MD ensemble. (A) The charged residue pairs forming salt bridges are connected by grey semi-transparent lines whose thickness linearly scales with the frequency of the corresponding salt bridge observed in MD simulations. Although more diverse salt bridges were observed in TIM9·10–Tim23 (10 in TIM9·10–Tim23 and 7 in TIM8·13–Tim23), these salt bridges were on average less stable than the ones in TIM8·13–Tim23, likely resulting in overall weaker interactions. **(B)** Snapshots of top views of the two chaperones along MD simulations of their holo forms in complex with TIM23. The top views of the chaperones in the apo forms are shown in Fig. S5E and F. Residues are color-coded according to the scheme reflected below the figure. **(C)** Ensemble view of the N-tail-bound state of TIM8·13–Tim23. The red surface represents the negatively charged E59 of Tim13 and E50 and D54 of Tim8. Blue stick-and-ball represents the side chain of positively charged residues (K8, K25, K27, K32, R57 and K66) of Tim23, which is shown as an ensemble of 25 structures. Ensemble view of the N-tail-bound state of TIM9·10–Tim23 is shown in fig. S14.

tional hydrophilic interactions to its client's IMS segment, which is ineffective in the TIM9·10-Tim23 interaction. The additional interaction effectively compensates for the lower affinity of TIM8·13 to the client's TM part. In the case of Tim23, this additional interaction involves a sequence stretch of at least 35-40 residues (Fig. 3). Remarkably, TIM8·13 has also been shown to be involved in the transport of a Ca²⁺-regulated mitochondrial carrier, the Asp/Glu carrier (31), which has an additional soluble calmodulin-like

domain. Whether this soluble domain is folded or disordered while the TM domain is attached to the hydrophobic chaperone cleft remains to be investigated. It is tempting to speculate that interactions between TIM8·13's top part and the calmodulin-like hydrophilic part of these carriers are important for this binding, similarly as for the case of Tim23 (fig. S1). Interestingly, membrane precursor proteins that have been shown *not* to interact with TIM8·13, such as mitochondrial carriers (Ggc, Aac) and Tim17, lack extended hydrophilic stretches, underlining the importance of those parts in binding (fig. S1). From the sequences of known clients and known 'non-clients' of TIM8·13, we propose that a minimum sequence length of about 20-25 residues is required for binding.

The nature of these additional hydrophilic interactions appears to involve primarily charged residues which form a complex and wide-connected interaction network that could be hard to suppress by mutating individual sites.

This study provides a rationale why mitochondria contain two very similar IMS chaperone complexes, the essential TIM9·10 and the non-essential TIM8·13 complex. The observation that this dual system is conserved even in humans suggests that the presence of the TIM8·13 system is not just the result of gene duplication, which appears rather often in yeast. The current results propose that for some substrates (like Tim23, or Asp-Glu carrier; see fig. S1), TIM8·13 can contribute stabilizing interactions with the hydrophilic soluble parts. Our competition experiments have also revealed that mitochondrial membrane precursor proteins may be transferred from one TIM chaperone to the other, opening the possibility that these two chaperones truly cooperate in precursor protein transfer to downstream insertases.

Taken together, our study reveals how a subtle balance of hydrophobic and hydrophilic interactions is used to tune promiscuity versus specificity in molecular chaperones. We propose that a similar balance of interactions determines the clientome of the cellular chaperones.

Methods

Plasmids. Genes coding for *Saccharomyces cerevisiae* Tim8 and Tim13 were cloned in the co-expression plasmid pET-Duet1. The expressed protein sequences were MSSLSTSD LASLDDTSKKEIATFLEGENSKQKVQMSIHQFTNICFK KCVESVNDSNLSSQEEQCLSNCVNRFLDTNIRIVNGL QNTR (Tim8) and MGSSHHHHHHSQDPSQDPENLYFQ GGLSSIFGGGAPSQQKEAATTAKTTPNPIAKELKNQIA QELAVANATELVNKISENCFEKCLTSPYATRNDACID QCLAKYMRSWNVISKAYISRIQNASASGEI (Tim13). A tobacco etch virus (TEV) cleavage site on Tim13 allows generating the final construct starting with GGLSS (the native Tim13 sequence starts with MGLSS). The same approach was used for preparing TIM9·10, including co-expression of the two proteins, with a cleavable His₆ tag on one of the proteins (Tim10), as described elsewhere (23). The gene coding for full-length *S. cerevisiae* Tim23 (C98S, C209S, C213A) with a C-terminal His₆-tag was cloned in the expression plasmid pET21b(+). The plasmid for expression

of the intrinsically disordered N-terminal domain of *S. cerevisiae* Tim23$_{IMS}$ (residues 1 to 98) with a N-terminal glutathion S-transferase (GST) tag is described in ref. (37), and the pET10N plasmid encoding Tim23$_{TM}$ in ref. (42). The *S. cerevisiae* Ggc1(C222S) construct was designed with a C-terminal 6xHis-tag in pET21a expression plasmid, reported earlier (23).

Protein expression and purification. We found that TIM9·10 and TIM8·13 chaperone complexes can be obtained either by over-expression in SHuffle T7 or BL21(DE3) *Escherichia coli* cells. Expression in the former results in soluble protein with correctly formed disulfide bonds, while the latter requires refolding from inclusion bodies. The proteins obtained with either method have indistinguishable properties (SEC, NMR). For TIM9·10, SHuffle expression results in better yield, while we obtain higher TIM8·13 yields with refolding from BL21(DE3). Accordingly, TIM9·10 and un-labelled TIM8·13 were over-expressed in the SHuffle T7 *Escherichia coli* cells and purified as described previously (23). Over-expression of the isotope labelled TIM8·13 chaperone complex from the BL21(DE3) *E. coli* cells was induced with 1 mM IPTG, and the cells were incubated for 4 hours at 37ºC. Cell pellets were sonicated and the inclusion-body fraction was resuspended sequentially, first in buffer A (50 mM Tris(tris(hydroxymethyl)aminomethane), 150 mM NaCl, pH 7.4) supplemented with 1% LDAO and 1% Triton100, then in buffer A supplemented with 1 M NaCl and 1M urea and lastly in buffer B (50 mM Tris, 250 mM NaCl, pH 8.5). The last pellet fraction was solubilized in buffer B supplemented with 50 mM DTT and 3 M guanidine-HCl at 4 ºC over night. The TIM8·13 complex was refolded by rapid dilution in buffer B containing 5 mM glutathione (GSH) and 0.5 mM glutathione disulfide (GSSG). The complex was purified on a NiNTA affinity column and the affinity tag was removed with TEV protease and an additional NiNTA purification step. Full-length precursor proteins, Tim23 and Ggc1, were expressed as inclusion bodies from BL21(DE3) cells, at 37 °C during 1.5 and 3 hours, respectively, after adding 1 mM IPTG. Precursor proteins were solubilized in buffer A supplemented with 4 M guanidine-HCl for Tim23 and 6 M guanidine-HCl for Ggc1, at 4 ºC over night. Precursor proteins were purified by affinity chromatography, in the same denaturating conditions used for solubilization. Imidazole was removed from the precursor protein sample with dialysis in buffer A supplemented with 4 M guanidine-HCl.

GST-tagged Tim23$_{IMS}$ was expressed in the soluble protein fraction from BL21(DE3)Ril+ cells during 4 h at 25 °C, after adding 1 mM IPTG. After sonication of the cell pellets, the soluble protein fraction was incubated with glutathion-agarose resin for 2 hours at 4 ºC. After washing the unspecifically bound proteins with 10 CV of Buffer A, the GST-tag was cleaved from the Tim23$_{IMS}$ by incubating the resin with 1 mg of TEV protease per 50 mg of precursor protein, at 4 ºC over night. Cleaved Tim23$_{IMS}$ and the protease were collected in the flow-through and an additional NiNTA purification step was applied to remove the TEV protease from the protein sample. Soluble Tim23$_{IMS}$ was subjected to gel fil-

tration on a Superdex 75 10/300 column and stored in buffer A.

Tim23$_{TM}$, comprising residues 92-222 (42), was produced in *E. coli* BL21(DE3)Ril+ during 3 hours at 37ºC and purified in denaturing conditions as described for the full-length Tim23. Chaperones proteins used for detection by NMR experiments were expressed in D_2O M9 minimal medium and labeled either with 1 g/L $^{15}NH_4Cl$ and 2 g/L D-[^{2}H,^{13}C] glucose or specifically labeled on isoleucine, alanine, leucine, valine side chains using the QLAM-A$^{\beta}$I$^{\delta1}$L^{proR}V^{proR} kit from NMR-Bio (www.nmr-bio.com) according to the manufacturer's instructions. The protein not detected by NMR in complex samples (i.e. the precursor proteins in complexes directed towards chaperone-detection, or the chaperone in preprotein-detected experiments) were unlabeled and produced in LB medium. Chaperone-bound Tim23$_{FL}$ was deuterated (produced in D_2O M9 mdeium), while the Tim23$_{IMS}$ fragment was prepared in H_2O M9 medium.

The fragments of human VDAC1 peptide (cyclic or linear VDAC$_{257-279}$) were prepared by solid-phase synthesis as described elsewhere (35), lyophilized, and resolubilized first in DMSO and then step-wise diluted into buffer, as described elsewhere (23). The peptide used for photo-induced cross-linking differed from the one used for NMR by the substitution of L263 by a Bpa side chain, as used earlier (23, 35).

Preparation of chaperone-precursor protein complexes. Purified precursor protein, i.e., either full-length Tim23, the trans-membrane Tim23$_{TM}$ fragment or Ggc1, was bound to NiNTA resin in 4 M guanidine-HCl. The column was washed with five column volumes (CV) of buffer A supplemented with 4 M guanidine-HCl, and with 5 CV of buffer A. A twofold excess of the chaperone complex was passed through the column twice. The column was washed with 10 CV of buffer A and the precursor-chaperone complex was eluted in 5 CV of buffer A supplemented with 300 mM imidazole. The precursor-chaperone complex was immediately subjected to dialysis against buffer A prior to concentrating on Amicon 30 kDa MWCO centrifugal filters (1000 g). Immediate removal of imidazole was particularly important for the preparation of the less stable Tim23$_{FL}$-TIM9·10 complex. Complexes of Tim23$_{IMS}$ with TIM8·13 or TIM9·10 were prepared by mixing two purified protein samples, and dialysis against buffer A. Formation of the precursor-chaperone complex was verified by size-exclusion chromatography (SEC) on a Superdex 200 column. The resulting complex was further characterized by SEC coupled to multi-angle light scattering (MALS). TIM8·13 and TIM8·13-Tim23 were furthermore analyzed by analytical ultra-centrifugation (AUC). Both experiments were performed at 10°C in Buffer A. The amount of eluted complex was estimated from the protein concentration, measured absorbance of the sample at 280 nm and the sum of the molecular weights and extinction coefficients of the chaperone and the precursor protein.

Competition assays. The first competition assay was performed by adding an equimolar mixture of TIM8·13 and TIM9·10 chaperones to the NiNTA bound precursor protein,

Tim23$_{FL}$ or Ggc1. After washing the column, precursor-chaperone complex was eluted in Buffer A supplemented with 300 mM imidazole. In the time dependent competition assay, the complex of a precursor protein and one of the chaperones (TIM8·13 or TIM9·10) was prepared, and then an equimolar amount of the other chaperone was added (time point 0). The reaction mixture was incubated at 30°C. After 0.5, 1 and 3 hours, an aliquot of the reaction mixture was taken and (newly formed) precursor-chaperone complex was isolated on a NiNTA affinity column. The difference in the amount of specific chaperone, TIM8·13 or TIM9·10, bound to precursor protein was analysed by SDS-PAGE and liquid chromatography coupled with mass-spectrometry (LC ESI-TOF MS, 6210, Agilent Technologies, at the MS platform, IBS Grenoble). Samples for analysis by MS were heat shocked for 15' at 90°C, resulting in the dissociation and precipitation of the precursor protein, while the apo-chaperones were recovered in the supernatant after cooling the sample and centrifugation for 10' at 39k g. As a reference, samples of precursor proteins, Tim23$_{FL}$ and Ggc1, bound to individual chaperone, TIM8·13 or TIM9·10, were prepared and analysed in parallel. To be noted, preparation of the TIM8·13-Ggc1 complex, in quantity sufficient for the analysis, was unsuccessful. To calculate the difference in the amount of specific chaperone bound to precursor protein, normalized areas under the chromatography peaks corresponding to each Tim monomer were used.

Isothermal Titration Calorimetry experiments. Calorimetric binding experiments of Tim23$_{IMS}$ and TIM chaperones were performed using a MicroCal ITC200 instrument (GE Healthcare). Sixteen successive 2.5 µl aliquots of 1.15 mM Tim23$_{IMS}$ were injected into a sample cell containing 55 µM TIM9·10 or TIM8·13. All ITC data were acquired in Buffer A at 20°C. Control experiments included titrating Tim23$_{IMS}$ into the Buffer A. The enthalpy accompanying each injection was calculated by integrating the resultant exotherm, which corresponds to the released heat as a function of ligand concentration added at each titration point. ITC data were analysed via the MicroCal Origin software using a single site binding model and nonlinear least squares fit of thermodynamic binding parameters (ΔH, K, and n). An identical procedure was performed for TIM8·13-cytochrome c ITC experiments. We also performed ITC experiments with the VDAC peptides; no effects could be detected, in line with a millimolar affinity, as already reported for the TIM9·10–cyclic-VDAC$_{257-279}$ peptide (23).

Cross-linking of VDAC$_{257-279}$. *In vitro* cross-linking of VDAC$_{257-279}$ used precisely the protocol described in ref. (23) for TIM9·10. Briefly, 5 µM TIM8·13 was mixed with VDAC$_{257-279}$ at 0, 25 or 50 µM, incubated 10 min on ice, and UV-illuminated (30 min, 4 °C).
We detected several cross-linking adducts of the β-hairpin peptide and the TIM components. Such multi-band behavior is similar to the pattern of cross-linking products of this peptide with either TOM subunits or cytosolic chaperones (35, 36). We suppose that this variability can result from a variable number of peptides bound to one molecule of protein and from different cross-linking sites on the protein, which in turn can cause different migration behavior in the SDS-PAGE.

SEC-MALS experiments. SEC-MALS experiments were performed at the Biophysical platform (AUC-PAOL) in Grenoble. The experimental setup comprised a HPLC (Schimadzu, Kyoto, Japan) consisting of a degasser DGU-20AD, a LC-20AD pump, a autosampler SIL20-ACHT, a column oven XL-Therm (WynSep, Sainte Foy d'Aigrefeuille, France), a communication interface CBM-20A, a UV-Vis detector SPD-M20A, a static light scattering detector miniDawn Treos (Wyatt, Santa-Barbara, USA), a dynamic light scattering detector DynaPro NANOSTAR, a refractive index detector Optilab rEX. The samples were stored at 4 °C, and a volume of 20, 40, 50 or 90 µl was injected on a Superdex 200, equilibrated at 4 °C; the buffer was 50 mM Tris, 150 mM NaCl filtred at 0.1 µm, at a flow rate of 0.5 ml/min. Bovine serum albumine was used for calibration. Two independent sets of experiments conducted with two different batches of protein samples were highly similar.

Analytical ultra-centrifugation. AUC experiments of TIM8·13 and TIM8·13-Tim23 were performed at 50000 rpm and 10 °C, on an analytical ultracentrifuge XLI, with a rotor Anti-60 and anti-50 (Beckman Coulter, Palo Alto, USA) and double-sector cells of optical path length 12 and 3 mm equipped of Sapphire windows (Nanolytics, Potsdam, DE). Acquisitions were made using absorbance at 250 and 280 nm wave length and interference optics. The reference is the buffer 50 mM Tris, 150 mM NaCl. The data were processed by Redate software v 1.0.1. The c(s) and Non Interacting Species (NIS) analysis was done with the SEDFIT software, version 15.01b and Gussi 1.2.0, and the Multiwavelenght analysis (MWA) with SEDPHAT software version 12.1b.

NMR spectroscopy. All NMR experiments were performed on Bruker Avance-III spectrometers operating at 600, 700, 850 or 950 MHz ^{1}H Larmor frequency. The samples were in the NMR buffer (50 mM NaCl, 50 mM Tris, pH 7.4) with 10% (v/v) D$_2$O, unless stated differently. All multidimensional NMR data were analyzed with CCPN (version 2) (43). DOSY data were analyzed with in-house written python scripts. For calculating chemical-shift perturbation data, the contribution of each different nuclei was weighted by the gyromagnetic ratios of the respective nucleus: e.g. the combined ^{1}H-^{15}N CSP was calculated as $\sqrt[2]{[\mathrm{CSP}^2_{1H} + \mathrm{CSP}^2_{15N} \cdot (\gamma_{15N}/\gamma_{1H})]}$, where the γ are the gyromagnetic ratios.

TIM8·13 and Tim23$_{IMS}$ resonance assignments. For the resonance assignment of TIM8·13, the following experiments were performed : 2D ^{15}N-^{1}H- BEST-TROSY HSQC, 3D BEST-TROSY HNCO, 3D BEST-TROSY HNcaCO, 3D BEST-TROSY HNCA, 3D BEST-TROSY HNcoCA, 3D BEST-TROSY HNcocaCB and 3D BEST-TROSY HNcaCB (44, 45) and a 3D ^{15}N-NOESY HSQC. The experiments

were performed with a 0.236 mM [^{2}H,^{15}N,^{13}C]-labeled TIM8·13, at 308 K and 333K. The NMR resonance assignment of TIM9·10 was reported earlier (23). We collected BEST-TROSY HNCA, HNCO and HNcoCA experiments to assign Tim23$_{IMS}$, aided by the previously reported assignment (37).

VDAC titration experiments. Cyclic hVDAC1$_{257-279}$ peptide was synthesized and lyophilized as described elsewhere (35). The peptide was dissolved in DMSO, and the DMSO concentration was reduced to 10% by step-wise addition of NMR buffer (1:1 in each step). Chaperone, TIM9·10 or TIM8·13, in buffer A was added to yield a final DMSO concentration of 6% and a chaperone concentration of 0.15 mM (TIM9·10) or 0.1 mM (TIM8·13). Combined ^{15}N-^{1}H chemical shift-perturbation (CSP) was calculated from the chemical shifts obtained from the ^{15}N-^{1}H HSQC spectra of the complex samples with molar ratio of 1:4 for TIM9·10:VDAC, and ratio of 1:5 for TIM8·13:VDAC, in comparison to the chemical shifts from the apo-chaperone spectrum. The NMR experiments were performed at 308K.

Tim23$_{IMS}$ titration experiments. For each titration point individual samples were prepared by mixing two soluble protein samples, and monitored using ^{15}N-^{1}H- BEST-TROSY HSQC experiments at 283K (for Tim23 observed experiment) or at 308K (for chaperone observed experiments). Titration samples with 100 μM [^{15}N]-labeled Tim23$_{IMS}$ with the molar ratios for Tim23$_{IMS}$:TIM8·13 from 1:0 to 1:4, and the molar ratios for Tim23$_{IMS}$:TIM9·10 from 1:0 to 1:5, were used. For the chaperone observed experiments, used samples contained 200 μM [^{2}H,^{13}C,^{15}N]-labeled TIM8·13 with molar ratios of Tim23$_{IMS}$ 1:0 and 1:1, and 350 μM [^{2}H,^{13}C,^{15}N]-labeled TIM9·10 with 1:0 and 1:3 molar ratios of Tim23$_{IMS}$.

NMR experiments with the Tim23$_{FL}$. Complexes of the chaperones with the full-length Tim23 were prepared as indicated above (Preparation of chaperone-preursor protein complexes). Peak positions (chemical shifts) of the amide backbone sites of TIM8·13, apo- and in the complex with Tim23$_{FL}$, were obtained from the ^{1}H-^{15}N HSQC experiments at 308K, with 120 μM [^{13}CH$_3$-ILV]-TIM8·13-Tim23$_{FL}$ sample. Similarly, to calculate combined ^{15}N-^{1}H and ^{13}C-^{1}H CSPs, chemical shifts of the amide backbone and ILVA-^{13}CH$_3$ groups of TIM9·10, apo- and in the complex with Tim23$_{FL}$, were obtained from the ^{1}H-^{15}N HSQC and ^{1}H-^{13}C HMQC experiments at 288K. Sample of the [^{13}CH$_3$-ILVA]-TIM9·10 with the Tim23$_{FL}$ was at 140 μM concentration. For the CSP calculations with the complexes of [^{2}H-^{15}N]-labelled Tim23$_{FL}$ and the chaperones (190 μM complex with TIM8·13, and 61 μM complex with TIM9·10), chemical shifts from ^{1}H-^{15}N HSQC experiments at 288K were used in comparison to the chemical shifts of the apo-Tim23$_{IMS}$.

Diffusion ordered spectroscopy. Diffusion-ordered NMR spectroscopy (DOSY) experiments were performed at 288K and 600 MHz ^{1}H Larmor frequency. Diffusion constants were derived from a series of one-dimensional ^{1}H spectra

either over the methyls region (methyl-selective DOSY experiments, for ^{13}CH$_3$-ILVA-labeled apo- and Tim23$_{FL}$ bound TIM9·10) or over the amides region (for [^{15}N]Tim23$_{FL}$-TIM9·10). Diffusion coefficients were obtained from fitting integrated 1D intensities as a function of the gradient strength at constant diffusion delay.

Small-angle X-ray scattering data collection and analysis. SAXS data were collected at ESRF BM29 beam line (46) with a Pilatus 1M detector (Dectris) at a distance of 2.872 m from the 1.8 mm diameter flow-through capillary. Data on TIM8·13 were collected in a batch mode. The X-ray energy was 12.5 keV and the accessible q-range 0.032 nm^{-1} to 4.9 nm^{-1}. The incoming flux at the sample position was in the order of 1012 photons/s in 700x700 mm^2. All images were automatically azimuthally averaged with pyFAI (47). SAXS data of pure TIM8·13 was collected at 1, 2.5 and 5 mg/mL using the BioSAXS sample changer (48). Ten frames of one second were collected for each concentration. Exposures with radiation damage were discarded, the remaining frames averaged and the background was subtracted by an online processing pipeline (49). Data from the three concentrations were merged following standard procedures to create an idealized scattering curve, using Primus from the ATSAS package (50). The pair distribution function p(r) was calculated using GNOM (51).

Online purification of the TIM8·13–Tim23$_{FL}$ and TIM9·10–Tim23$_{FL}$ complexes using gelfiltration column (HiLoad 16/600 Superdex S200 PG) was performed with a high pressure liquid chromatography (HPLC) system (Shimadzu, France), as described in reference (52). The HPLC system was directly coupled to the flow-through capillary of SAXS exposure unit. The flow rate for all online experiments was 0.2 mL/min. Data collection was performed continuously throughout the chromatography run at a frame rate of 1 Hz. All SAXS data have been deposited on SASBDB.

Molecular dynamics simulations and fitting of SAXS data. The inital model of Tim23 was built using I-TASSER (Iterative Threading ASSEmbly Refinement) (53) and QUARK webservers (54) which predicted a long unstructured N-terminal tail and four/five helical structures in the transmembrane domain. The structure of TIM9·10 hexamer built in our previous work (23) was used as the initial model of TIM9·10 chaperone, and as the template to build the model of TIM8·13 chaperone based on the sequence of yeast Tim8 and Tim13 (UniProt IDs: P57744 and P53299) by homology modeling with MODELLER (55). (Note that in the crystal structure of TIM8·13 (PDB-ID 3CJH), more than 75 residues are missing in each Tim8-Tim13 pair, thus requiring model building.) The disulfide bonds related to the twin CX$_3$C motif were also kept in these models. The structures of the TIM8·13 hexamer and Tim23 were subsequently used to build the full structure of the TIM8·13-Tim23 complex by manually wrapping the helical structures of the transmembrane domain of Tim23 around the hydrophobic cleft of TIM8·13 which has been identified by NMR, and leaving the unstructured N-terminal of Tim23 as a floppy tail. The com-

plex structure was further optimized by energy minimization and relaxation in 100 ns MD simulations using the simulation protocol as described in the following section. This model was used to generate the so-called 'N-tail unbound' ensemble in which the N-terminal half of Tim23 is free in solution. Based on the 'N-tail unbound' model of the TIM8·13-Tim23 complex, we further constructed the 'N-tail bound' model in which the N-terminal half of Tim23 is in contact with the upper part of TIM8·13. This was achieved by adding a restraint term in the force field using PLUMED plugin ([56]), $V_{\text{restraints}}$, which is a half-harmonic potential of the form of $k(R - R_0)^2$ when R is larger than R_0, and zero when R less than R_0. Here R is the distance between the center of mass of Tim23Nter and the top part of the chaperone. We used $R_0 =$ 1 nm and k = 400 kJ·mol⁻¹. The 'N-tail bound' and unbound models for the TIM9·10-Tim23 complex were constructed by replacing TIM8·13 with TIM9·10 based on the corresponding TIM8·13-Tim23 models.

The TIM8·13-Tim23 complex in the 'N-tail unbound' conformation was placed into a periodic cubic box with sides of 17.5 nm solvated with TIP3P water molecules containing Na⁺ and Cl⁻ ions at 0.10 M, resulting in ∼700,000 atoms in total. To reduce the computational cost, the complex in 'N-tail bound' conformation was placed in a smaller cubic box with sides of 12.9 nm, resulting in ∼300,000 atoms in total. The systems of the TIM9·10-Tim23 complex have similar size as the TIM8·13-Tim23 systems in the corresponding states. The apo TIM8·13 chaperone was placed into a periodic cubic box with sides of 12.0 nm, containing ∼230,000 atoms.

The Amber ff99SB-disp force field ([57]) was used for all simulations. The temperature and pressure were kept constant at 300 K using the v-rescale thermostat and at 1.0 bar using the Parrinello-Rahman barostat with a 2 ps time coupling constant, respectively. Neighbor searching was performed every 10 steps. The PME algorithm was used for electrostatic interactions. A single cut-off of 1.0 nm was used for both the PME algorithm and Van der Waals interactions. A reciprocal grid of 96 x 96 x 96 cells was used with 4th order B-spline interpolation. The hydrogen mass repartitioning technique ([58]) was employed with a single LINCS iteration (expansion order 6) ([59]), allowing simulations to be performed with an integration time step of 4 fs. MD simulations was performed using Gromacs 2018 or 2019 ([60]).

A total of 4.25 μs trajectories were collected to sample the conformational space of the chaperone-Tim23 complexes in both 'N-tail bound' and 'N-tail unbound' states. Four μs trajectories were also collected to sample the ensemble of apo TIM8·13 chaperone. These sampled conformations were used for further ensemble refinement using the Bayasian Maximum Entropy (BME) method ([38], [39]) guided by experimental SAXS data as decribed in our previous work ([23]). The distribution of both states in principle could be identified from the force field but needs substantial sampling. Therefore instead of estimating the prior by large-scaled MD simulations, we assigned equal weight (50%) for both states by inputting equal number of conformations (5000) into the mixed

ensemble so without bias to either state. By tuning the regularization parameter in the BME reweighting algorithm, we adjusted the conformational weights in variant degrees to improve the fitting with experimental SAXS data.

The hydrogen bond and salt bridge formation between the N-terminal tail of Tim23 (residue 1-100) and the top surface of the chaperones was analyzed by GetContacts scripts (https://getcontacts.github.io/), and visulized using Flareplot (https://gpcrviz.github.io/flareplot/). Protein structures were visualized with PyMOL ([61]) and VMD ([62]).

Calculations of affinities and populations.

Estimation of the population of 'N-tail bound' states from ITC-derived K_d. We attempted to link the ITC-derived dissociation constant of the Tim23$_{\text{IMS}}$ fragment to the populations of bound and unbound states in the Tim23$_{\text{FL}}$-chaperone complexes, using a rationale akin to the one outlined earlier for binding of disordered proteins to two sub-sites ([63]). Briefly, we treat the N-terminal tail of Tim23 as a ligand, and the remaining bound complex as the target protein, then the relationship between the population of the bound state (P$_{\text{bound}}$) and the binding affinity (K$_d$) can be written as P$_{\text{bound}}$/(1-P$_{\text{bound}}$) = C$_{\text{eff}}$/K$_d$ where C$_{\text{eff}}$ is the effective concentration of the disordered N-tail which was estimated to be between 0.2-3 mM from the MD simulations, resulting in the estimation of P$_{\text{bound}}$ to be between 75% and 98%.

Estimation of the K_d ratio from competition assays. Determining dissociation constants of TIM chaperones to its insoluble client proteins is hampered by the impossibility to form the complexes by solution methods such as titration, as it requires the pull-down method outlined in Fig. [1]A. Nonetheless, the amount of TIM8·13-Tim23 and TIM9·10-Tim23 complexes obtained in the competition assays (Fig. [1]) can provide an estimate of the relative affinities. The dissociation constants can be written from the concentrations as follows:

$$K_d^{\text{TIM8·13-Tim23}} = \frac{[\text{TIM8·13}] \times [\text{Tim23}]}{[\text{TIM8·13-Tim23}]}$$

$$K_d^{\text{TIM9·10-Tim23}} = \frac{[\text{TIM9·10}] \times [\text{Tim23}]}{[\text{TIM9·10-Tim23}]}$$

where [TIM8·13-Tim23] denotes the concentration of the formed chaperone-precursor complex, and [TIM8·13] and [Tim23] are the concentrations of free chaperone and precursor protein in solution. The latter is negligible, as no free precursor protein is eluted from the column (some aggregated precursor protein was removed from the equilibrium). Both chaperones have been applied at the same concentration c_0 = [TIM8·13]+[TIM8·13-Tim23] = [TIM9·10]+[TIM9·10-Tim23] to the resin-bound precursor protein that was present at a concentration b_0 = [Tim23]+[TIM9·10-Tim23]+[TIM8·13-Tim23]. Using the ratio of formed complex obtained in the competition assay,

$r = [\text{TIM9·10-Tim23}]/[\text{TIM8·13-Tim23}]$ leads to

$$\frac{K_d^{\text{TIM9·10-Tim23}}}{K_d^{\text{TIM8·13-Tim23}}} = \frac{c_0 \times (1+r) - r \times b_0}{[c_0 \times (1+r) - b_0] \times r}$$

The experimental protocol does not allow to determine with precision the concentrations of precursor protein (b_0) and each chaperone (c_0), as the former is bound to a resin. As the chaperone was added in excess, and some of the precursor protein precipitated on the column, we can safely assume $c_0 \leq b_0$. With an experimentally found ratio of formed complexes of $r = 5$ and assuming that c_0/b_0 assumes the values of 1-5, the K_d ratio falls in the range of 1:25 to 1:6, i.e. ca. one order of magnitude.

Acknowledgements

We thank Nils Wiedemann and Caroline Lindau for many insightful discussions and exploratory in vivo experiments. We thank Aline Le Roy and Christine Ebel (SEC-MALS; AUC), Luca Signor (mass spectrometry) and Caroline Mas (ITC) for excellent support. We thank Hubert Kalbacher (Univ. Tübingen) for synthesis of the VDAC peptides, and Karine Giandoreggio-Barranco for preparing Tim23 samples and Undina Guillerm for support with protein production. We are grateful to Bernhard Brutscher, Alicia Vallet and Adrien Favier for excellent NMR platform operation and management. We are grateful to Markus Zweckstetter and Nikolaus Pfanner for providing the Tim23$_{\text{IMS}}$ and the Tim23$_{\text{TM}}$ plasmids, respectively. This study was supported by the European Research Council (StG-2012-311318-ProtDyn2Function), and the Agence Nationale de la Recherche (ANR-18-CE92-0032). This work used the platforms of the Grenoble Instruct-ERIC center (ISBG; UMS 3518 CNRS-CEA-UGA-EMBL) within the Grenoble Partnership for Structural Biology (PSB), supported by FRISBI (ANR-10-INBS-05-02) and GRAL, financed within the University Grenoble Alpes graduate school (Ecoles Universitaires de Recherche) CBH-EUR-GS (ANR-17-EURE-0003). Y. W. and K.-L. were supported by the BRAINSTRUC structural biology initiative from the Lundbeck Foundation. We acknowledge access to computational resources from the Danish National Supercomputer for Life Sciences (Computerome) and the ROBUST Resource for Biomolecular Simulations (supported by the Novo Nordisk Foundation). This work was furthermore supported by the Deutsche Forschungsgemeinschaft (RA 1028/8-1,2 to D. R.)

Competing Interests

The authors declare that they have no competing financial interests.

Author contributions

K. W. and P. S. initiated this project. I. S., A. H., O. D., D. C., B. B. and K. W. prepared protein samples. I.S. performed the pull-down and competition assays. I.S., K. W., B. B., O. D. and P. S. performed and analyzed NMR data. K. W. and M. B. performed and analyzed SAXS experiments. Y. W. and K. L.-L. performed and analyzed MD simulations. T. J. and D. R. performed/analyzed cross-linking experiments. All authors contributed ideas and discussion. I.S., Y. W. and P.S. made the figures. P. S., I. S. and Y. W. wrote the manuscript with input from all authors.

Data availability

The chemical shift assignments of TIM8·13 have been deposited in the BioMagResBank (www.bmrb.wisc.edu) under accession number 50213. All MD models and SAXS data have been deposited in the SASBDB (www.sasbdb.org) under accession numbers SASDH89 (TIM8·13-Tim23), SASDJP4 (TIM9·10-Tim23), SASDJQ4 (TIM8·13) and SASDEF2 (TIM9·10 (23)). All chemical-shift perturbation data have been deposited on Mendeley data (http://dx.doi.org/10.17632/8cr8rvtddm.1).

Supplementary materials

Fig. S1: Kyte-Doolittle hydrophobicity of membrane precursor protein.
Fig. S2: Ensemble representation of the structure of TIM9·10 (blue/green) holding full-length Ggc.
Fig. S3: NMR spectra showing that TIM8·13 and TIM9·10 do not form mixed complexes.
Fig. S4: LC/ESI–TOF MS analysis used to quantify amounts of chaperone complexes.
Fig. S5: Sequence alignment of the small Tims across eukaryotes.
Fig. S6: Pull-down experiment of Tim23$_{\text{FL}}$ with wild-type TIM9·10, wild-type TIM8·13 and TIM8·13$_{\text{K30F,S36L}}$.
Fig. S7: Local order parameters in TIM8·13 and TIM9·10 from TALOS.
Fig. S8: Chemical-shift perturbations on TIM8·13 upon addition of the linear VDAC$_{257-279}$ peptide.
Fig. S9: ITC curves of interactions of wild-type and mutant chaperones.
Fig. S10: NMR data showing the Tim23$_{\text{FL}}$ interactions with TIM9·10 and TIM8·13.
Fig. S11: NMR data showing the interaction of Tim23$_{\text{TM}}$ with TIM9·10.
Fig. S12: SEC-MALS data of samples containing TIM9·10, Tim23$_{\text{FL}}$ and TIM8·13.
Fig. S13: Experimental characterization of the size of TIM chaperones and their precursor protein complexes.
Fig. S14: SAXS data of chaperone complexes and views of the electrostatics of TIM chaperones.
Table S1: Thermodynamic parameters of TIM8·13 - Tim23$_{\text{IMS}}$ interaction from ITC.

References

1. Powers, E. T., Morimoto, R. I., Dillin, A., Kelly, J. W. & Balch, W. E. Biological and Chemical Approaches to Diseases of Proteostasis Deficiency. *Ann. Rev. Biochem.* **78**, 959–991 (2009).

2. Kim, Y. E., Hipp, M. S., Bracher, A., Hayer-Hartl, M. & Hartl, U. F. Molecular Chaperone Functions in Protein Folding and Proteostasis. *Ann. Rev. Biochem.* **82**, 323–355 (2013).

3. Kim, D. H. *et al.* Cytosolic targeting factor AKR2A captures chloroplast outer membrane-localized client proteins at the ribosome during translation. *Nat. Commun.* **6**, 1–13 (2015).

4. Bose, D. & Chakrabarti, A. Substrate specificity in the context of molecular chaperones. *IUBMB Life* **69**, 647–659 (2017).

5. Rüdiger, S., Germeroth, L., Schneider-Mergener, J. & Bukau, B. Substrate specificity of the DnaK chaperone determined by screening cellulose-bound peptide libraries. *EMBO J.* (1997).

6. Blond-Elguindi, S. *et al.* Affinity panning of a library of peptides displayed on bacteriophages reveals the binding specificity of BiP. *Cell* **75**, 717–728 (1993).

7. Knoblauch, N. T. M. *et al.* Substrate Specificity of the SecB Chaperone. *J. Biol. Chem.* **274**, 34219–34225 (1999).

8. Koldewey, P., Stull, F., Horowitz, S., Martin, R. & Bardwell, J. C. Forces Driving Chaperone Action. *Cell* **166**, 369–379 (2016).

9. He, L., Sharpe, T., Mazur, A. & Hiller, S. A molecular mechanism of chaperone-client recognition. *Sci. Adv.* **2**, e1601625–e1601625 (2016).

10. Hiller, S. & Burmann, B. M. Chaperone–client complexes: A dynamic liaison. *J. Magn. Reson.* **289**, 142–155 (2018).

11. Genevaux, P. *et al.* In vivo analysis of the overlapping functions of DnaK and trigger factor. *EMBO Rep.* **5**, 195–200 (2004).

12. Bukau, B., Deuerling, E., Pfund, C. & Craig, E. A. Getting Newly Synthesized Proteins into Shape. *Cell* **101**, 119–122 (2000).

13. Burmann, B. M. *et al.* Regulation of α-synuclein by chaperones in mammalian cells. *Nature* **577**, 127–132 (2020).

14. Burmann, B. M., Wang, C. & Hiller, S. Conformation and dynamics of the periplasmic membrane-protein–chaperone complexes OmpX–Skp and tOmpA–Skp. *Nat. Struct. Mol. Biol.* **20**, 1265–1272 (2013).

15. Thoma, J., Burmann, B. M., Hiller, S. & Müller, D. J. Impact of holdase chaperones Skp and SurA on the folding of β-barrel outer-membrane proteins. *Nat. Struct. Mol. Biol.* **22**, 795–802 (2015).

16. Saio, T., Guan, X., Rossi, P., Economou, A. & Kalodimos, C. G. Structural Basis for Protein Antiaggregation Activity of the Trigger Factor Chaperone. *Science* **344**, 1250494–1250494 (2014).

17. Jiang, Y., Rossi, P. & Kalodimos, C. G. Structural basis for client recognition and activity of Hsp40 chaperones. *Science* **1319**, 1313–1319 (2019).

18. Huang, C., Rossi, P., Saio, T. & Kalodimos, C. G. Structural basis for the antifolding activity of a molecular chaperone. *Nature* **537**, 202–206 (2016).

19. Wiedemann, N. & Pfanner, N. Mitochondrial Machineries for Protein Import and Assembly. *Annu. Rev. Biochem.* **86**, 685–714 (2017).

20. Sirrenberg, C. *et al.* Carrier protein import into mitochondria mediated by the intermembrane proteins Tim10/Mrs11 and Tim12/Mrs5. *Nature* **391**, 912–915 (1998).

21. Koehler, C. M. *et al.* Import of Mitochondrial Carriers Mediated by Essential Proteins of the Intermembrane Space. *Science* **279**, 369–373 (1998).

22. Lionaki, E. *et al.* The essential function of Tim12 in vivo is ensured by the assembly interactions of its C-terminal domain. *J. Biol. Chem.* **283**, 15747–53 (2008).

23. Weinhäupl, K. *et al.* Structural Basis of Membrane Protein Chaperoning through the Mitochondrial Intermembrane Space. *Cell* **175**, 1365–1379.e25 (2018).

24. Koehler, C. M. *et al.* Human deafness dystonia syndrome is a mitochondrial disease. *Proc. Natl. Acad. Sci.* **96**, 2141–2146 (1999).

25. Paschen, S. A. *et al.* The role of the TIM8-13 complex in the import of Tim23 into mitochondria. *EMBO J.* **19**, 6392–6400 (2000).

26. Roesch, K., Curran, S. P., Tranebjaerg, L. & Koehler, C. M. Human deafness dystonia syndrome is caused by a defect in assembly of the DDP1/TIMM8a-TIMM13 complex. *Hum. Mol. Genet.* **11**, 477–486 (2002).

27. Kang, Y. *et al.* Function of hTim8a in complex IV assembly in neuronal cells provides insight into pathomechanism underlying Mohr-Tranebjærg syndrome. *Elife* **8**, 1–32 (2019).

28. Morgenstern, M. *et al.* Definition of a high-confidence mitochondrial proteome at quantitative scale. *Cell Rep.* **19**, 2836–2852 (2017).

29. Leuenberger, D., Bally, N. A., Schatz, G. & Koehler, C. M. Different import pathways through the mitochondrial intermembrane space for inner membrane proteins. *EMBO J.* **18**, 4816–4822 (1999).

30. Davis, A. J., Sepuri, N. B., Holder, J., Johnson, A. E. & Jensen, R. E. Two intermembrane space TIM complexes interact with different domains of Tim23p during its import into mitochondria. *J. Cell Biol.* **150**, 1271–1282 (2000).

31. Roesch, K., Hynds, P. J., Varga, R., Tranebjaerg, L. & Koehler, C. M. The calcium-binding aspartate/glutamate carriers, citrin and aralar1, are new substrates for the DDP1/TIMM8a–TIMM13 complex. *Hum. Mol. Genet.* **13**, 2101–2111 (2004).

32. Hoppins, S. C. & Nargang, F. E. The Tim8-Tim13 Complex of Neurospora crassa Functions in the Assembly of Proteins into Both Mitochondrial Membranes. *J. Biol. Chem.* **279**, 12396–12405 (2004).

33. Habib, S. J., Waizenegger, T., Lech, M., Neupert, W. & Rapaport, D. Assembly of the TOB complex of mitochondria. *J. Biol. Chem.* **280**, 6434–6440 (2005).

34. Hiller, S. *et al.* Solution structure of the integral human membrane protein VDAC-1 in detergent micelles. *Science* **321**, 1206–1210 (2008).

35. Jores, T. *et al.* Characterization of the targeting signal in mitochondrial β-barrel proteins. *Nat. Commun.* **7**, 12036 (2016).

36. Jores, T. *et al.* Cytosolic Hsp70 and Hsp40 chaperones enable the biogenesis of mitochondrial β-barrel proteins. *J. Cell Biol.* **217**, jcb.201712029 (2018).

37. de la Cruz, L., Bajaj, R., Becker, S. & Zweckstetter, M. The intermembrane space domain of Tim23 is intrinsically disordered with a distinct binding region for presequences. *Protein Sci.* **19**, 2045–2054 (2010).

38. Orioli, S., Larsen, A. H., Bottaro, S. & Lindorff-Larsen, K. How to learn from inconsistencies: Integrating molecular simulations with experimental data. *Prog. Mol. Biol. Transl. Sci.* **1**, 123–176 (2020).

39. Bottaro, S., Bengtsen, T. & Lindorff-Larsen, K. Integrating molecular simulation and experimental data: a bayesian/maximum entropy reweighting approach. In *Structural Bioinformatics*, 219–240 (Springer, 2020).

40. Ahmed, M. C., Papaleo, E. & Lindorff-Larsen, K. How well do force fields capture the strength of salt bridges in proteins? *PeerJ* **6**, e4967 (2018).

41. Hiller, S. Chaperone-Bound Clients: The Importance of Being Dynamic. *Trends Biochem. Sci.* **44**, 517–527 (2019).

42. Truscott, K. N. *et al.* A presequence- and voltage-sensitive channel of the mitochondrial preprotein translocase formed by Tim23. *Nat. Struct. Biol.* **8**, 1074–1082 (2001).

43. Vranken, W. F. *et al.* The CCPN data model for NMR spectroscopy: development of a software pipeline. *Proteins* **59**, 687–96 (2005).

44. Schanda, P., Van Melckebeke, H. & Brutscher, B. Speeding up three-dimensional protein NMR experiments to a few minutes. *J. Am. Chem. Soc.* **128**, 9042–9043 (2006).

45. Favier, A. & Brutscher, B. Recovering lost magnetization: polarization enhancement in biomolecular NMR. *J. Biomol. NMR* **49**, 9–15 (2011).

46. Pernot, P. *et al.* Upgraded ESRF BM29 beamline for SAXS on macromolecules in solution. *J. Synchrotron Radiat.* **20**, 660–664 (2013).

47. Ashiotis, G. *et al.* The fast azimuthal integration Python library: PyFAI. *J. Appl. Crystallogr.* **48**, 510–519 (2015).

48. Round, A. *et al.* BioSAXS Sample Changer: A robotic sample changer for rapid and reliable high-throughput X-ray solution scattering experiments. *Acta Crystallogr. Sect. D Biol. Crystallogr.* **71**, 67–75 (2015).

49. Brennich, M. E. *et al.* Online data analysis at the ESRF bioSAXS beamline, BM29. *J. Appl. Crystallogr.* **49**, 203–212 (2016).

50. Petoukhov, M. V. *et al.* New developments in the ATSAS program package for small-angle scattering data analysis. *J. Appl. Crystallogr.* **45**, 342–350 (2012).

51. Svergun, D. I. Determination of the regularization parameter in indirect-transform methods using perceptual criteria. *J. Appl. Crystallogr.* **25**, 495–503 (1992).

52. Brennich, M. E., Round, A. R. & Hutin, S. Online Size-exclusion and Ion-exchange Chromatography on a SAXS Beamline. *J. Vis. Exp.* e54861 (2017).

53. Yang, J. *et al.* The I-TASSER Suite: protein structure and function prediction. *Nat. Methods* **12**, 7–8 (2015).

54. Xu, D. & Zhang, Y. Ab initio protein structure assembly using continuous structure fragments and optimized knowledge-based force field. *Proteins* **80**, 1715–1735 (2012).

55. Webb, B. & Sali, A. Comparative protein structure modeling using modeller. *Curr. Protoc. bioinfo.* **47**, 5–6 (2014).

56. Bonomi, M. *et al.* Promoting transparency and reproducibility in enhanced molecular simulations. *Nat. Methods* **16**, 670–673 (2019).

57. Robustelli, P., Piana, S. & Shaw, D. E. Developing a molecular dynamics force field for both folded and disordered protein states. *Proc. Natl. Acad. Sci. USA* **115**, E4758–E4766 (2018).

58. Hopkins, C. W., Le Grand, S., Walker, R. C. & Roitberg, A. E. Long-time-step molecular dynamics through hydrogen mass repartitioning. *J. Chem. Theory Comput.* **11**, 1864–1874 (2015).

59. Feenstra, K. A., Hess, B. & Berendsen, H. J. Improving efficiency of large time-scale molecular dynamics simulations of hydrogen-rich systems. *J. Comput. Chem.* **20**, 786–798 (1999).

60. Abraham, M. J. *et al.* Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **1**, 19–25 (2015).

61. Schrödinger, LLC. The PyMOL molecular graphics system, version 2.3 (2020).

62. Humphrey, W., Dalke, A. & Schulten, K. VMD – Visual Molecular Dynamics. *J. Mol. Graph.* **14**, 33–38 (1996).

63. Zhou, H. X., Pang, X. & Lu, C. Rate constants and mechanisms of intrinsically disordered proteins binding to structured targets. *Phys. Chem. Chem. Phys.* **14**, 10466–10476 (2012).

64. Kyte, J. & Doolittle, R. F. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**, 105–132 (1982).

65. Shen, Y. & Bax, A. Protein backbone and sidechain torsion angles predicted from NMR chemical shifts using artificial neural networks. *J. Biomol. NMR* **56**, 227–241 (2013).

Protein	χ^2 / DoF	N	K [M^{-1}]	- ΔH [cal/mol]	ΔS [cal/mol/deg]	K$_d$ [µM]
TIM8·13$_{WT}$	592.9	2.02 ± 0.09	(1.38 ± 0.14) x 10^4	3606 ± 241	6.64	66 *
TIM8·13$_{E50K}$	789.5	2.34 ± 0.06	(2.27 ± 0.27)×10^4	2285 ± 104	12.1	44
TIM8·13$_{K47D}$	3965	2.19 ± 0.17	(1.49 ± 0.37)×10^4	3435 ± 455	7.37	67
TIM8·13$_{E59K,E50K}$	1195	2.50 ± 0.05	(3.44 ± 0.44)×10^4	2120 ± 74	13.5	42 *
TIM8·13$_{E59K,D54T}$	4112	2.05 ± 0.06	(1.93 ± 0.53)×10^5	1327 ± 53	19.7	10 *
TIM8·13$_{D54R}$	1484	1.83 ± 0.12	(1.27 ± 0.16)×10^4	4964 ± 469	1.83	79

Table S1. Thermodynamic parameters of TIM8·13 - Tim23$_{IMS}$ interaction from isothermal titration calorimetry measurements for wild-type and mutant TIM8·13 variants. The mutants have selected charge inversions in the top region of the chaperone. Using the 'One Set of Sites' model, the fitted parameters were: N (number of sites), K (binding constant in M^{-1}) and ΔH (heat change in cal/mol). Entropy change ΔS (in cal/mole/deg) was calculated from ΔH and K after the fitting, and dissociation constant K$_d$ (in µM) was calculated as the reciprocal of the binding constant. The goodness of the fit is reported as the sum of the squares of the deviations of the theoretical curves from the experimental points (χ^2) divided by the degrees of freedom (DoF, the total number of experimental points minus the number of adjustable parameters).

Fig. S1. Kyte-Doolittle hydrophobicity of membrane precursor proteins known not to bind to TIM8·13 (Tim17, Ggc1, Aac2) and proteins known to bind (Glu-Asp carrier Agc1, Tim23 and, putatively, Sal1 - the human form of Sal1, Citrin, is known to depend on the human homolog of TIM8·13). The Kyte-Doolittle hydrophobicity (64) has been determined with the web server of ExPASy, using a window size of 13 and the standard linear weight variation model. Trans-membrane helices (but not other secondary structure elements) are indicated above each plot, as determined from either UniProt or a modelling with SwissModeller (for Sal1, using the structure of Aac2 as a template).

Tim9
Tim10
Ggc1

hydrophobic residues in
the binding cleft of TIM9·10

Fig. S2. Ensemble representation of the structure of TIM9·10 (blue/green) holding full-length Ggc1 (orange), as reported in ref. (23). In contrast to the complexes presented in this study, the complex has a 2:1 stoichiometry, and essentially the entire precursor protein is located to the hydrophobic binding cleft of the chaperone.

(A)

TIM9•10
TIM9•10+TIM8•13, o/n

Fig. S3. TIM8·13 and TIM9·10 do not form mixed complexes. NMR spectra of a $^2H,^{15}N,^{13}C$ labelled TIM9·10 sample (blue) and a mixture of this sample with unlabelled (NMR-invisible) TIM8·13 after overnight incubation. If TIM9·10 formed mixed oligomers with TIM8·13, the environment around each of the Tim9 and Tim10 subunits, as they would be surrounded by Tim8 or Tim13 subunits, would be different. Thus, the spectrum of TIM9·10 would feature additional peaks corresponding to those alternate environments. The spectrum after over-night incubation does not show any additional peaks and features, only the one set of peaks corresponding to the hexameric TIM9·10. Therefore, this data demonstrates that TIM9·10 does not form mixed hetero-oligomers with TIM8·13.

Fig. S4. Liquid chromatography coupled with electrospray ionization, time-of-flight mass spectrometry (LC/ESI–TOF MS) analysis used to read out the difference in the amount of specific chaperone, TIM8·13 and TIM9·10, bound to precursor protein.

(A) Mass chromatograms of the control sample, complex of TIM9·10 chaperone bound to Ggc1 (left), and the competition reaction three hours after adding TIM8·13 to the pre-formed Ggc1–TIM9·10 complex (right). Deconvoluted mass values are indicated next to the corresponding peaks. Mass obtained for the unspecific peak, indicated with asterisk, was 10 226.53 (left) and 10 226.53 amu (right). **(B)** Same as in panel A, with the Tim23$_{FL}$ as a substrate precursor protein. Mass of the unspecific peak in the left chromatogram was 10 198.69 amu.

(C) Mass chromatogram of the TIM8·13 chaperone bound to Tim23$_{FL}$ (left) as a control, and of the competition reaction three hours upon adding TIM9·10 to the pre-formed Tim23$_{FL}$–TIM8·13 complex (right). In the left chromatogram, obtained mass from the unspecific peaks (impurities) was 10 207.94 (left asterisk) and 11 208.27 amu (right asterisk).

(D) Deconvoluted mass spectra for each of the chromatography peaks from the chromatogram shown in panel C, right.

Reaction mixtures were heat-shocked before the analysis, resulting in the presence of the chaperone only in the analysed sample (see methods).

Additional experiments were performed with a mixture of TIM8·13 and TIM9·10 of known quantities, in order to correlate observed integrals of these chromatograms to molar ratios between the two chaperones (not shown).

Fig. S5. Sequence alignment of the small Tims across eukaryotes, including *Saccharomyces cerevisiae* (Sc), *Schizosaccharomyces pombe* (Sp), *Neurospora crassa* (Nc), *Arabidopsis thaliana* (At), *Drosophila melanogaster* (Dm), *Caenorhabditis elegans* (Ce), *Danio rerio* (Dr), *Rattus norvegicus* (Rn), *Mus musculus* (Mm) and *Homo sapiens* (Hs). The positions of the hydrophobic motif are highlighted in red, and was identified by analyzing the mean hydrophobicity across the alignment, as reporter earlier (23). The respective numbering of the hydrophobic position is provided below the sequence alignment. The strictly conserved cysteines in dark grey, and the top part of the chaperones in light grey. Positions discussed in the main text are numbered above the alignment, whereby the numbering refers to *Sc* Tim8.

Fig. S6. Results of a pull-down binding experiment of Tim23$_{FL}$ with wild-type TIM9·10, wild-type TIM8·13 and TIM8·13$_{K30F,S36L}$. **(A)** SDS-PAGE of the different fractions of the pull-down experiments. Lane descriptions are as in Figure 1H. **(B)** SDS-PAGE of the soluble (S) and insoluble (P) fraction of the Tim23$_{FL}$ complex with TIM9·10$_{WT}$, TIM8·13$_{WT}$ or TIM8·13$_{K30F,S36L}$ showing that under the same conditions, Tim23$_{FL}$–TIM8·13$_{K30F,S36L}$ complex is less stable. Immediately after the elution of the complex during the pull-down experiment, dialysis was performed. Dialysed sample was centrifuged in order to obtain the remaining complex in the soluble fraction and an insoluble Tim23$_{FL}$ in the pellet. **(C)** Quantification of the amount of complex obtained after imidazole-elution. The three pull-down experiments (TIM9·10, TIM8·13, TIM8·13$_{K30F,S36L}$) were done in parallel, binding each time the same amount of Tim23$_{FL}$ to the column, and performing the same wash, elution and end-wash (denaturing conditions) steps. The experiment was repeated twice, with Tim23$_{FL}$ from two different production batches (shown in red and grey, respectively). As the protein was from different purifications, the amount of Tim23$_{FL}$ bound onto the column was not exactly identical; for this reason, the amount obtained in each experiment (of the order of 3 mg complex) was normalized to the amount of TIM9·10-Tim23$_{FL}$ complex.

Fig. S7. Local order parameters in TIM8·13 and TIM9·10, derived from assigned backbone chemical shifts and the TALOS-N (65) software. This data shows that the core of the chaperones is rather rigid, and the tentacles become increasingly flexible towards the N- and C-termini.

TIM8·13-VDAC$_{linear}$

HN $\Delta\delta$ [ppm]
>0.08
>0.06
>0.04
<0.04

Fig. S8. ^{1}H-^{15}N chemical-shift perturbations on TIM8·13 upon addition of the linear VDAC$_{257-279}$ peptide, mapped onto the structure of TIM8·13. In contrast to the cyclic VDAC$_{257-279}$ peptide, we observe widely spread CSPs with the linear peptide, indicating unspecific binding across the chaperone.

Fig. S9. Isothermal titration calorimetry (ITC) experiments with TIM8·13 and TIM9·10 chaperones. Thermograms (thermal power as a function of time) are displayed in the upper plots, and binding isotherms (ligand-normalized heat effects per injection as a function of the molar ratio, [Tim23$_{IMS}$]/[chaperone]) are displayed in the lower plots. **(A)** Calorimetric titration of wild-type TIM8·13 and TIM9·10 with cytochrome c. No binding of chaperones with folded, soluble cytochrome c was detected, supporting that interaction of the chaperone with client proteins requires either a hydrophobic region, or an unfolded stretch of hydrophilic residues. **(B)** ITC of TIM9·10 and TIM8·13 with Tim23$_{IMS}$. **(C)** Calorimetric titration of TIM8·13 mutants with Tim23$_{IMS}$. Variants were obtained by charge inversion of negatively charged residues in the top region of the chaperone. The highly conserved Lys47 was exchanged for a negatively charged residue to investigate its relevance for binding of Tim23. All variants show similar isothermal binding curves, despite altered charges. This observation points to more complex binding interactions, which were not significantly perturbed in the chosen mutants. **(D)** Calorimetric titration of TIM9·10 variants with Tim23$_{IMS}$. Variants were obtained by inverting the charge of selected positively charged residues in the top region of the chaperone. No binding was detected between Tim23$_{IMS}$ and any of the TIM9·10 variants.

Fig. S10. Full-length Tim23 interactions with TIM9·10 and TIM8·13 chaperones.
(A) Chemical-shift perturbations observed upon addition of the Tim23$_{FL}$ to TIM9·10 (top) and TIM8·13 (bottom). Comparison of CSPs on TIM8·13 induced by Tim23$_{FL}$ and VDAC$_{257-279}$ (Fig. 2C), shows that for binding of full-length Tim23, besides conserved hydrophobic binding site, TIM8·13 employs its hydrophilic top region where 3 fold increase in CSPs can be seen (mapped on the structure in Fig. 4C). Asterisk marks residues 16Val and 26Leu which showed biggest methyl CSP upon adding Tim23$_{FL}$. **(B)** Methyl spectra overlay of the ILVA-labeled apo-TIM9·10 chaperone sample (blue) and ILVA-TIM9·10 in complex with the full-length Tim23 (red). In the zoomed-in regions, showing resides 22Ala and 28Leu, difference in the peak intensity is shown with the 1D traces (hydrogen spectra) at the corresponding carbon frequency of the selected peak. The biggest methyl CSP for the residue 16Val is indicated with an red arrow. **(C)** Methyl-detected CSP of the ILVA-labeled TIM9·10 upon addition of the full-length Tim23 (top). **(D)** Mapping of Tim23$_{FL}$-induced methyl CSPs and biggest peak-broadening on TIM9·10, showing absence of interaction with the top part of TIM9·10.

Fig. S11. Interaction of trans-membrane Tim23 with the TIM9·10 chaperone.
(A) ^{1}H-^{15}N chemical-shift perturbations observed upon adding the trans-membrane construct Tim23$_{TM}$ to TIM9·10. **(B)** Methyl-detected chemical-shift perturbations of the ILVA-labeled TIM9·10 upon addition of the trans-membrane Tim23. **(C)** Mapping of Tim23$_{TM}$-induced methyl CSPs on TIM9·10, showing binding in the hydrophobic cleft, similarly to the binding detected with Tim23$_{FL}$.

Fig. S12. Investigating the presence of ternary complexes in a sample containing TIM9·10, Tim23$_{FL}$ and TIM8·13, using multi-angle light scattering coupled to size-exclusion chromatography. TIM9·10-Tim23$_{FL}$ complex was prepared using the pull-down method of Figure 1B, and TIM8·13 was added. If TIM8·13 binds to the soluble N-terminal half of Tim23 with sufficient affinity, a ternary complex may be detected.
(A) Elution profiles of TIM9·10–Tim23$_{FL}$ complex (blue) and TIM9·10–Tim23$_{FL}$ after adding slight excess of TIM8·13 (pink). The elution profiles of the two samples are similar, with no additional peak appearing at higher molecular weights, suggesting that there is no ternary complex present. Protein fraction eluting at 10 mL, in the column void volume, corresponds to the aggregates present in both samples. **(B)** Multi-angle light scattering analysis of two main peaks of the size-exclusion chromatography of the TIM9·10–Tim23$_{FL}$ after adding slight excess of TIM8·13. A protein fraction eluting at 20 mL, corresponds for 36% of the injected mass and it corresponds to a co-elution of TIM9·10 (Mw$_{theor}$ = 60.8 kDa) and TIM 8·13 (Mw$_{theor}$= 62.4 kDa), while the fraction at 18.2 mL, corresponds for 49% of the injected mass and it corresponds to a co-elution of the complex of TIM9·10–Tim 23$_{FL}$ (Mw$_{theor}$= 85 kDa) and the complex of TIM8·13–Tim 23$_{FL}$ (Mw$_{theor}$= 86.6 kDa).

Fig. S13. Experimental characterization of the size of TIM chaperones and their precursor protein complexes.

(A) Size-exclusion-chromatography coupled to multi-angle light scattering (SEC-MALS) of TIM9·10–Tim23 and TIM8·13–Tim23. For TIM9·10 the dominant contribution (> 85 %) corresponds to 56 kDa, or ca. 5.5 subunits; for TIM9·10–Tim23, the major contribution (71 %) corresponds to a mass of 77.5 kDa, in reasonable agreement with the expected 23 kDa increase from Tim23-binding, with a clearly visible 'shoulder' at the position of the apo-TIM9·10. For TIM8·13, a peak corresponding to ca. 86 % of the signal is at a mass of 57.9 kDa; in TIM8·13–Tim23, the peak is at 74.2 kDa, in reasonable agreement with the expected 23 kDa increase. A control experiment with BSA yielded an observed mass of 62 kDa (theoretical: 66 kDa).

(B) Analytical ultra-centrifugation of TIM8·13 and TIM8·13-Tim23 complexes. For the TIM8·13 we observe a main contribution, $98 \pm 1\%$ of the total signal, at 2.89 ± 0.01 S (s20w = 3.97 ± 0.01S). The Non Interacting Species analysis gives Mw = 57 ± 5 kDa. For the TIM8·13-Tim23 sample we observe a main contribution ($67 \pm 2\%$ of the total signal) at 3.76S (s20w = 5.17S), at 0.7 mg/mL sample concentration, slightly shifted to s20w = 5.67S, at 2 mg/mL. The Non Interacting Species analysis results in a Mw = 70 ± 5 kDa, similar to SEC-MALS. A further contribution is detected at 3 ± 0.05S (s20w = 4.1 ± 0.1S), for 33 ± 5 % of the total signal. The Non Interacting Species analysis gives Mw = 50 ± 3 kDa, could be imprecise. This contribution superimposes to the main one of TIM8·13 alone.

(C) NMR diffusion-ordered spectroscopy curves (DOSY), obtained as integrals of one-dimensional ^{1}H spectra over either methyls (first three) or amides (fourth panel) as a function of the gradient strength. The four panels correspond to three different samples, ^{13}CH$_3$-ILVA-labeled apo-TIM9·10 (first panel), TIM9·10-ILVA methyl labeled TIM9·10-Tim23$_{FL}$ complex (second panel) and Tim23$_{FL}$-^{15}N labeled TIM9·10-Tim23$_{FL}$ complex (last two panels). The first two used a ^{13}C-filtered experiment, the third one uses a ^{1}H methyl-frequency selective scheme and the fourth uses a ^{15}N-filtered version. The latter thus detects amide protons; as these are water exchangeable, in particular in disordered proteins, the observed diffusion coefficient is likely over-estimated, as it contains contributions from the diffusion coefficient of water; thus, this value is to be seen as an upper limit. The diffusion coefficient for a spherical particle scales with the cubic root of the molecular weight, which would lead to an expected change in the diffusion coefficient of $(71.1/94.1)^{1/3}$, i.e. ca. 10 % lower in the complex compared to the apo state; additional domain motions, such as the flexibility of the Tim23 tail, decreases the diffusion coefficient, i.e. would lead to a larger relative change. This expected 10-15 % effect on diffusion coefficients is in good agreement with the experimental effect (11-20 %). A TIM9·10:Tim23 stoichiometry, as found for TIM9·10–Ggc1 would lead to an expected DOSY effect of 32 % (considering only mass), thus in worse agreement with the data.

Fig. S14. (A) P(r) distributions of TIM9·10 in the apo form and in complex with Tim23 or Ggc1. **(B)** P(r) distributions of TIM8·13 in the apo form and in complex with Tim23. **(C)** SAXS curves of TIM8·13 ensemble. **(D)** Kratky plot of SAXS curves of TIM8·13 ensemble. **(E)** Top view of a snapshot of MD simulations of apo TIM8·13. **(F)** Top view of a snapshot of MD simulations of apo TIM9·10. **(G)** Ensemble view of the N-tail-bound state of TIM9·10–Tim23.