Influenza-induced activation of recruited alveolar macrophages during the early

inflammatory phase drives lung injury and lethality

Clarissa M Koch[1], Kishore R Anekalla[1], Yuan-Shih Hu[1], Jennifer M. Davis[1], Mark Ciesielski[1],

Gaurav Gadhvi[2], Shang-Yang Chen[2], Margaret Turner[1], Yuan Cheng[1], Bria M Coates[1,3,4], Hiam

Abdala-Valencia[1], Paul A Reyfman[1], Alexander V Misharin[1], GR Scott Budinger[1], Deborah R

Winter[2,±], Karen M Ridge[1,5,±]

[1]Pulmonary and Critical Care Division, [2]Rheumatology Division, Department of Medicine,

[3]Department of Pediatrics, Northwestern University Feinberg School of Medicine, Chicago, IL USA,

[4]Ann & Robert H. Lurie Children's Hospital of Chicago, Chicago IL, [5]Department of Cell and

Developmental Biology, Northwestern University Feinberg School of Medicine, Chicago, IL.

15
16 ± Contributed equally to this paper
17

18 Corresponding author:
19
Karen M. Ridge, Ph.D.
Northwestern University
Feinberg School of Medicine
Division of Pulmonary and Critical Care Medicine
Simpson Querrey Biomedical Research Center
Department of Cell and Developmental Biology
303 E. Superior, 5-520
Chicago, IL 60611
Kridge@northwestern.edu
312-503-1648 – office

30
31

Abstract

Severe respiratory virus infections initiate a robust host immune response that contributes to disease severity. Immunomodulatory strategies that limit virus-initiated inflammation are of critical importance. In this study, we compared the host response to influenza A virus (IAV) infection in susceptible animals (wild-type, WT) to resilient mice (*Vimentin*[-/-] mice). We identified distinct gene expression patterns in recruited monocyte-derived alveolar macrophages (MoAMs) associated with three phases (Infiltrating, Early Inflammatory, Late Inflammatory) that evolve in sequence over the course of IAV infection. We report a core set of pro-inflammatory genes involved in the WT MoAM Early Inflammatory response that is suppressed in *Vim*[-/-] MoAMs. Moreover, we identify CEBPB, Jun-AP1, and IRF transcriptions factors as regulators of this attenuated inflammatory response. We performed causal experiments using bone-marrow chimeras to credential that *Vim*[-/-] MoAMs with suppressed pro-inflammatory genes confer protection from influenza-induced mortality in WT susceptible mice. Taken together, these data support the notion that vimentin plays a causal role in determining the pro-inflammatory function of recruited MoAMs and drives IAV-induced lung injury.

Introduction

Severe viral pneumonia due to influenza A virus (IAV) or severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is characterized by the release of proinflammatory mediators that damage the alveolar epithelial and capillary barrier to cause acute respiratory distress syndrome (ARDS)[1, 2]. This points to the importance of a balanced and regulated immune response, where infection can be controlled while keeping tissue damage to a minimum. This hypothesis is clinically supported by a recent analysis of patients enrolled in the ARDSnet, in which patients with a hyperinflammatory endotype of ARDS, characterized in part by increased levels of circulating cytokines and chemokines, had worse clinical outcomes [3].

Shortly after IAV or SARS-CoV-2 infection, tissue-resident alveolar macrophages (TRAMs) are depleted, and monocytes are released from the bone marrow and recruited into the infected, injured lung [1, 4-6]. Upon entering the lung, these recruited monocytes, referred to as monocyte-derived alveolar macrophages (MoAMs), drive the immune response in humans and mice. MoAMs activate inflammatory signaling pathways in response to pathogen- or damage-associated molecular patterns (PAMPs or DAMPs) released from infected and dying cells [1, 3, 4, 7]. Preventing the recruitment of MoAMs into the lung by deleting CCR2 or its ligand reduces the severity of the dysregulated immune response (e.g., no cytokine storm) without affecting viral clearance [4]. These data suggest that recruited MoAMs mediate the excessive pro-inflammatory response associated with tissue damage. Indeed, elevated levels of IL-1β and IL-18, cytokines produced by the NLRP3 inflammasome, in the alveolar exudate of IAV-infected patients serve as poor prognostic markers of viral pneumonia outcomes [8-10]. Inhibiting the NLRP3 inflammasome early in IAV infection impairs viral clearance and increases morbidity and mortality in mice [11]. In contrast, inhibiting the NLRP3 inflammasome post-viral clearance reduces inflammation-induced tissue damage and promotes lung repair [11, 12]. These studies highlight the importance of understanding that recruited alveolar macrophages may play different roles in the virus-infected lung depending on the

stage of infection (e.g., pre- versus post-viral clearance). As a result, therapies that target recruited

MoAMs may be of interest [4, 13, 14]; however, no study has yet investigated the time-dependent

transcriptional signature and function of MoAMs in response to IAV infection.

Vimentin, a type III intermediate filament, is ubiquitously expressed in macrophages and plays a

role in several critical immune response processes. Vimentin expression is increased in COVID-

19 patients' lungs compared with healthy, control and COPD patients [15]. In patients with

ulcerative colitis and Crohn's disease, vimentin expression was also significantly increased in

inflamed tissues [16]. Consistent with these observations, vimentin-null ($Vim^{-/-}$) mice develop

significantly less gut inflammation compared to wild-type mice with acute colitis induced by dextran

sodium sulfate [17]. We have shown that $Vim^{-/-}$ mice are protected from LPS-induced lung injury

and death by suppressing pro-inflammatory cytokines [18]. Vimentin is a known ligand for some

PRRs, including nucleotide-binding oligomerization domain-containing protein 2 (NOD2) and NLR

Family Pyrin Domain Containing 3 (NLRP3). We, and others, reported that vimentin is required to

activate the NLRP3 inflammasome [18, 19] and subsequent NF-κB activation [20], which is

consistent with the observation that pro-inflammatory gene expression is altered in microglia lacking

vimentin in a murine model of Alzheimer's disease [21]. Interestingly, bulk RNA-seq transcriptomic

analysis showed significant upregulation of Vim, which was associated with decreased viral host

resistance in cultured embryonic stem cells and embryos [22]. Collectively, these studies support

the notion that vimentin may play a role in the immune response to IAV infection.

The temporal transcriptional response of recruited MoAMs that drives the immune response in the

IAV-infected host has not been previously reported. For the first time, our data identifies the distinct

gene expression patterns in recruited MoAMs associated with three phases (Infiltrating, Early

Inflammatory, Late Inflammatory) that evolve in sequence following infection with influenza A virus.

Additionally, we report that wild-type mice mount a sustained pro-inflammatory response

contributing to increased lung injury and death. In contrast, $Vim^{-/-}$ mice are remarkably resilient to

IAV infection. Whether vimentin plays a causal role in determining the pro-inflammatory function of

recruited MoAMs in the IAV-infected lung has not been previously investigated. If vimentin indeed

affects MoAMs' pro-inflammatory function, is it a cell-autonomous phenomenon? Using isochronic

adoptive MoAM transfer experiments and molecular profiling in mice, we sought to determine

whether the loss of vimentin-related protection from influenza-induced lung injury is intrinsic to Vim^{-}

$^{/-}$MoAMs. Our data supports a paradigm in which recruited MoAMs upregulate vimentin expression,

activate maladaptive responses, and consequently exhibit a persistent pro-inflammatory response,

promoting virus-induced lung injury.

Results

116
Vimentin null mice are protected from influenza-induced lung injury and mortality

Our group has recently reported that persistent inflammation in patients with severe SARS-CoV-2

pneumonia is sustained via stable circuits between recruited alveolar macrophages and T cells[1].

Using single-cell RNA-seq, distinct clusters of monocytes-derived alveolar macrophages (MoAMs)

were identified from the bronchoalveolar lavage fluid (BALF) collected from patients with severe

SARS-CoV-2 pneumonia (**Figure 1a, Supplemental Figure 1a**) [1]. MoAM1 cluster was

characterized by high expression of pro-inflammatory cytokine genes, including *IL1B, CXCL2, and*

CCL3, and low expression of reparative genes *MRC1* and *C1QA* (**Figure 1c, d)** [23-25]. In contrast,

cluster MoAM3 exhibited the highest expression of *MRC1, C1QA, and MERTK. MERTK* encodes

a receptor necessary for phagocytosis of apoptotic cells – the uptake of apoptotic cells reduces the

expression of pro-inflammatory cytokines and chemokines from macrophages. This process has

been mechanistically linked to transcriptional responses leading to a change in macrophage

phenotype from pro-inflammatory to reparative [26, 27] (**Figure 1c-d).** By querying this single-cell

RNA-seq dataset, we found that *VIM* was among the genes differentially expressed between

MoAM1 and MoAM3: with the highest expression in the pro-inflammatory MoAM1 population and

decreasing in the reparative MoAM3 population (**Figure 1b-c).** These data from patients with severe

SARS-CoV-2 pneumonia show that higher *VIM* expression is associated with a pro-inflammatory

MoAM phenotype in viral pneumonia, while lower levels of *VIM* expression characterize MoAMs

with a reparative phenotype.   Importantly, these data suggest that there may be a causal

relationship between *VIM* expression and MoAM phenotype.

Next, we used a clinically relevant, causal murine model of viral pneumonia to determine whether

the host response to influenza A virus (IAV) infection is altered in $Vim^{-/-}$ mice compared to WT mice.

WT and $Vim^{-/-}$ mice were infected with a lethal dose of IAV (Strain A/WSN/1933 (H1N1), 200 pfu)

and assessed for morbidity and mortality. WT mice had a 100% mortality rate, whereas >80% of

the $Vim^{-/-}$ mice were alive 10 days post-infection (p.i.) (**Figure 1e**). By 14 days p.i., 50% of the

surviving $Vim^{-/-}$ mice had recovered from mild lethargy, coat ruffling, febrile shaking, and eye-

watering. $Vim^{-/-}$ mice exhibited a lower wet-to-dry lung weight ratio and decreased total protein

concentration in BALF than WT mice infected with IAV, indicating less damage to the alveolar-

capillary barrier (**Figure 1f-g**). Histopathological examination of IAV-infected $Vim^{-/-}$ mice lungs also

demonstrated less severe alveolar damage, characterized by reduced alveolar edema and less

lung tissue damage than in IAV-infected WT mice (**Figure 1h**). Vimentin-deficiency did not interfere

with IAV replication or clearance, as evidenced by comparable viral titers (**Figure 1i**) and similar

levels of the H1N1 viral RNA molecules, such as matrix protein 2, nucleoprotein, and

neuraminidase, which were detected in recruited MoAMs from WT and $Vim^{-/-}$ mice by RNA-seq

(**Supplemental Figure 1b**). These results show that $Vim^{-/-}$ mice are surprisingly resilient to IAV

infection with improved survival rates and decreased lung injury, independent of viral load.

The influx of immune cells into the airspace during IAV infection and the resulting production of pro-

inflammatory cytokines is a major driver of lung injury. We performed a systematic flow cytometric

analysis of whole-lung lysates to characterize the immune cell recruitment into the lungs

(**Supplemental figure 1c**). The number of tissue-resident alveolar macrophages (TRAMs) rapidly

decreased after IAV infection at a similar rate in both genotypes, with a more than 80% reduction

by 6 days p.i. (**Figure 1j**). MoAMs are typically not present in the alveolar compartment in normal,

healthy lung tissue at steady-state (e.g., 0 days p.i.). During IAV infection, monocytes are recruited

and rapidly differentiated into MoAMs in the lung parenchyma [5, 28]. WT and $Vim^{-/-}$ mice infected

with IAV recruited comparably increasing number of MoAMs into the lung from 2 days p.i. onward

(**Figure 1j**). Similarly, neutrophil recruitment peaked on day 3 p.i. in both genotypes (**Supplemental**

Figure 1d). Despite similar immune cell recruitment, $Vim^{-/-}$ mice had decreased levels of pro-

inflammatory cytokines and chemokines, such as IL-1β, TNFα, RANTES, MIP-2, MCP-1, and Type

I interferons IFNα/β, compared to WT mice following IAV infection (**Figure 1k**). In summary, wild-

type mice, but not $Vim^{-/-}$ mice, exhibited a pro-inflammatory phenotype in response to IAV infection.

As there was no difference in MoAM recruitment between genotypes, these data suggest that

vimentin plays an essential role in determining the pro-inflammatory function of MoAMs.

Three distinct temporal patterns characterize the transcriptional response of MoAMs to

influenza A virus

To investigate the response of WT MoAMs to IAV infection, we performed RNA-seq on flow-sorted

MoAMs to assess global gene expression profiles on 2, 3, 4, and 6 days p.i. (**Supplemental Figure**

2a, 2b). We identified 2623 differentially expressed genes over this time course and, using k-means

clustering, defined three clusters that correspond temporally to three phases of the IAV response:

Infiltrating (2 days p.i.), Early Inflammatory (3-4 days p.i.), and Late Inflammatory (6 days p.i.)

(**Figure 2a**).

The first cluster is characterized by genes with the highest expression in the **Infiltrating Phase** at

2 days p.i. (**Figure 2a-b, Supplemental Figure 2c**). Gene Ontology (GO) analysis finds this cluster

is enriched for genes associated with *cell migration* and *cell adhesion*, such as *Cx3cr1, Rock1,*

Tln1, and Cxcl14 (**Figure 2c-d**), likely involved in transendothelial migration of leukocytes[29-32].

These results are consistent with the requirement for MoAMs to leave the vasculature and migrate

across the endothelium to reach the alveolar space soon after IAV infection*.* We also observed

genes in this cluster associated with the recruitment of lymphocytes, such as *Cd86,* MHC Class II

gene *H2-Aa,* and *Ccl8* (**Figure 2c-d, Supplemental Figure 2d**). Furthermore, our results suggest

that MoAMs may be proliferating during this phase due to the enrichment of GO processes, such

as *DNA replication initiation*, and the expression of cell cycle genes, such as *Mcm5* (**Figures 2c-f**).

Taken together, these data show that the primary role of MoAMs in the Infiltrating Phase of the IAV

response is to enter the lung parenchyma and promote the recruitment of immune cells.

The second cluster is characterized by genes that peak in expression during the **Early**

Inflammatory Phase at 3 days p.i. (**Figure 2a-b, Supplemental Figure 2c**). This cluster is

enriched for genes associated with *inflammatory response* and *regulation of cytokine secretion*

(**Figure 2c)** – such as *Tlr2*, *Il1b,* and *Il1r2* (**Figure 2d)**. These gene expression changes correspond

with the secretion of inflammatory cytokines, such as IL-1β and TNFα, beginning on Day 3 of IAV

response (**Figure 1k)** and binding of viral PAMPs through pattern recognition receptors (PRRs),

including TLR2 [28]. The expression of genes associated with *chemotaxis,* such as *Ccl3, Ccl2,* and

Cxcl2, also peak during the Early Inflammatory phase (**Figure 2c-d**) and corresponds with the

secretion of their respective proteins MCP-1, MIP1-α and MIP-2 measured in the BALF (**Figure**

1k). These data collectively suggest that WT MoAMs adopt a pro-inflammatory state during the

Early Inflammatory Phase of the IAV response.

The third cluster is characterized by genes that display a continuous increase in expression until

the **Late Inflammatory Phase** at 6 days p.i. (**Figure 2a-b, Supplemental Figure 2c**). This cluster

is enriched for genes associated with the GO processes of *cell differentiation* and *adaptive immune*

response – such as *Irf8, Mafb, Ifng, Ctsh, Ccr1* and*, Arg2* (**Figure 2d, Supplemental Figure 2d).**

The increasing expression of *Irf8* and *Mafb* further supports the notion that recruited monocytes

differentiate into monocyte-derived alveolar macrophages [33, 34]. Interestingly*,* the gene encoding

vimentin increased in expression over time, suggesting that vimentin may play a role throughout

the IAV response (**Figure 2d**). Although GO analysis implicated metabolic processes in this cluster

(**Figure 2c**), and macrophage activation has been associated with cellular respiration changes[35-

37], we did not observe preferential expression of genes in the glycolysis or oxidative

phosphorylation pathways at any time point (**Figure 2e-f**).

Since interferon pathways are an essential element of the viral immune response, we investigated

the temporal expression of Type I and Type II interferon genes. We found Type I IFNα response

genes peaked in expression on day 2, which coincided with their enrichment in the Infiltrating cluster

(**Figure 2e-f)**. This is also consistent with Type I IFNs' role in viral clearance.   In contrast, Type II

IFNy response genes did not reach their highest expression level until 6 days p.i. and were enriched

in the Late Inflammatory cluster (**Figure 2d-f)**.   Type II IFNy has been shown to stimulate

macrophages and monocytes to produce more chemoattractants and other inflammatory

molecules, resulting in a cycle of persistent inflammation [1, 38]. These data support Type I

interferon's role as an early effector of the immune response, while Type II interferon is involved in

the later immune response.

While GO analysis can be informative, many terms, such as *inflammatory response* and *innate*

immune response, lack cell-type specificity. *In vivo* macrophages are characterized by considerable

diversity and plasticity and respond to environmental cues by acquiring distinct functional

phenotypes. *In vitro* macrophages can be polarized toward a pro-inflammatory (M1) phenotype or

a reparative (M2) phenotype through the administration of LPS or IL-4, respectively [25, 39, 40]. To

further define the generic *inflammatory response* in recruited MoAMs, we generated annotations

for M(LPS) and M(IL-4) responses by integrating gene sets from several published studies on LPS

(M(LPS)), and IL-4 (M(IL-4)) stimulated macrophages *in vitro* (see Methods)[25, 39-41]. We found that

M(IL-4) genes were enriched in the Infiltrating cluster, and their expression peaked on day 2 p.i.

Notably, recruited WT MoAMs quickly downregulated M(IL-4) genes on days 3, 4, and 6 p.i. (**Figure**

2e-f). Furthermore, we found that the M(LPS) genes peaked on day 3 p.i. in WT MoAMs **(Figure 2e**)

and were enriched in the Early Inflammatory cluster (**Figure 2f**). M(LPS) genes were also enriched

in the Late Inflammatory cluster (days 4 and 6 p.i.) but to a lesser degree. (**Figure 2f**). Taken

together, our results support a critical switch in WT MoAM function as they transition from their initial

transcriptional state in the Infiltrating phase to a pro-inflammatory phenotype during the Early and

Late Inflammatory phases.

***Vim*^{-/-} MoAMs modulate expression of Infiltrating and Early Inflammatory, but not Late**

Inflammatory phase genes.

To gain a better understanding of the conferred protection from IAV infection in *Vim*^{-/-} mice (**Figure**

1), we performed RNA-seq on flow-sorted MoAMs from *Vim*^{-/-} mice at 2, 3, 4, and 6 days p.i.

(**Supplemental Figures 2a, 2b, 3a**). We compared global gene expression between genotypes

each day p.i., identifying down- and upregulated genes in *Vim*^{-/-} compared to WT MoAMs. The

greatest differential gene expression occurred on day 3 p.i. (**Figure 3a**). To characterize processes

with differential expression between genotypes, we ran GO analysis on up- and downregulated

genes on each day p.i. (**Figure 3b**). Top GO processes that enrich on at least two days p.i. uncover

a pattern of downregulated expression of genes involved in the inflammatory response and stress

response and increased expression of genes involved in cell cycle and extracellular matrix

organization in *Vim*^{-/-} MoAMs throughout IAV infection (**Figure 3b)**. Furthermore, using GSEA we

found that gene sets associated with inflammation – including M(LPS) and Type I Interferon – were

significantly decreased in expression in *Vim*^{-/-} MoAMs on every day p.i. (**Supplemental Figure 3b)**,

In contrast, *Cell Cycle* and M$_{(IL-4)}$ associated genes had increased expression in *Vim*^{-/-} MoAMs

compared to WT MoAMs (**Supplemental Figure 3b)**.

Using the prior clustering of 2623 genes across the three phases of the IAV response (**Figure 2a**),

we compared gene expression between MoAMs in WT and *Vim*^{-/-} mice during each phase (**Figure**

3c-e, Supplemental 3f-h). In *Vim*^{-/-} MoAMs, genes in the Infiltrating cluster were increased in

expression while genes in the Early Inflammatory cluster tended to be decreased (**Figure 3c,d,f,**

Supplemental Figure 3g,h). In contrast, genes in the Late Inflammatory cluster were less changed

between genotypes (**Figure 3e, Supplemental Figure 3g,h**). In line with our previous findings,

day 3 p.i. represented the most significant difference between genotypes in any cluster (**Figure 3c-**

f). Indeed, day 2 and day 3 *Vim*^{-/-} MoAMs were globally more similar to each other than their WT

counterparts, suggesting a greater transcriptional shift occurs between day 2 to 3 p.i. in WT

compared to *Vim*^{-/-} MoAMs (**Supplemental Figure 3f).** These transcriptional differences between

genotypes were mirrored in the data for processes associated with each phase (**Figure 2f**). M$_{(IL-4)}$

and Cell Cycle genes, associated with the Infiltrating phase, exhibited a higher and sustained level

of expression in *Vim*^{-/-} MoAMs compared to WT (**Figure 3c**). These results were confirmed using

GSEA (**Supplemental Figure 3b**). Type I IFN response genes were differentially expressed

between genotypes, but were downregulated over time in both genotypes. In contrast, M$_{(LPS)}$ genes

associated with the Early Inflammatory phase were downregulated on 2 and 3 days p.i. in *Vim-/-*

compared to WT MoAMs (**Figure 3d**), suggesting an early attenuation. Type II IFN response gene

expression did not significantly differ between genotypes (**Figure 3e**). We next asked whether any

of the individual gene sets correlated with each other. We observed that *M$_{(IL-4)}$* and Cell cycle gene

expression are highly correlated in both genotypes, suggesting they may be part of a more

extensive network of co-regulated processes (**Supplemental Figure 3c, d**). None of the gene sets

significantly differed in expression in steady-state tissue-resident alveolar macrophages,

suggesting that vimentin-deficiency has a larger effect during the response to IAV-infection than in

steady-state (**Supplemental Figure 3e**).

These results support an altered function of $Vim^{-/-}$ MoAMs in response to IAV that favors the

maintenance of Infiltrating phase genes and the suppression of Early Inflammatory phase genes to

confer protection from influenza-induced lung injury. Of note, these data demonstrate that the

greatest difference between genotypes occurs on day 3 p.i. and this represents the key timepoint

for the protective phenotype in $Vim^{-/-}$ mice.

$Vim^{-/-}$ MoAMs fail to activate core pro-inflammatory genes in the early response to IAV

Thus far, our results suggest that the maintenance of Infiltrating phase gene expression and

suppression of pro-inflammatory genes in $Vim^{-/-}$ MoAMs, particularly in the Early Inflammatory

phase, is responsible for the resilient phenotype observed in $Vim^{-/-}$ mice infected with IAV (see

Figure 1). To identify macrophage-specific genes involved in the Early Inflammatory Phase, we

focused on the $M_{(LPS)}$ response associated with the increased expression on day 3 p.i. in WT mice.

In MoAMs from $Vim^{-/-}$ mice, fewer $M_{(LPS)}$ genes exhibited peak expression on day 3 p.i. (**Figure 4a**).

Moreover, $M_{(LPS)}$ genes were enriched among those genes exhibiting the greatest expression

decrease in $Vim^{-/-}$ MoAMs compared with WT during the course of IAV response (**Supplemental**

Figure 3b). Of the $M_{(LPS)}$ genes most decreased on at least one day, we find that the majority failed

to intensify expression on day 3 p.i. in $Vim^{-/-}$ MoAM compared with WT MoAM (**Figure 4b-c,**

Supplemental Figure 4a). This core set of pro-inflammatory genes includes interferon-stimulated

genes (ISGs), such as *Ifit1, Irgm1, and Rsad2*, genes associated with chemokine pathways,

including *Ccrl2* and *Cxcl2,* and genes associated with pathogen sensing, *Fcgr1, and Tlr2* (**Figure**

4b-c). Complementary to the decrease in $M_{(LPS)}$ gene expression, $M_{(IL-4)}$ genes, such as *Cd36*,

Cd11c, and *Cd83*, previously associated with the Infiltrating phase, exhibited higher expression on

all days post IAV infection in $Vim^{-/-}$ MoAMs compared with WT MoAMs (**Supplemental Figures 3b,**

c, 4b). Collectively, these data suggest $Vim^{-/-}$ MoAMs fail to activate core pro-inflammatory genes

associated with the Early Inflammatory phase in response to IAV.

Prior work has demonstrated that macrophage activation is largely defined by the activity of cell-

type-specific transcription factors (TF) that poise regulatory elements for activation by stimulus-

specific TFs [42-44]. To better characterize the impact on gene regulation observed in $Vim^{-/-}$

MoAMs, we searched for TF binding sites enriched in the promoters of vimentin-dependent genes

associated with the Early Inflammatory phase. First, we classified the 440 Early Inflammatory

genes into those "most changed" and "least changed" in $Vim^{-/-}$ MoAMs based on their fold-change

compared with WT MoAMs (**Figure 4d-e, Supplemental Figure 4c**). Next, we performed motif-

finding on each gene set using a known sequence motifs database (see Methods). We found that

TF families previously implicated in an inflammatory response, such as *Jun-AP1* and *Rela*, were

significantly enriched among the most changed but not the least changed genes (**Figure 4f**). Some

Irf family motifs were differentially enriched between the two gene sets, while others were shared

(**Figure 4f**). As supported by induction of gene expression in IAV and the decrease observed in

$Vim^{-/-}$, those associated with ISGs (*Irf1/7/9*) were more likely to be affected by vimentin-deficiency

(**Figure 4f**). At the same time, *Irf8*, which is more generally involved in macrophage function,

exhibited similar enrichment in both groups (**Supplemental Figure 4e**). Furthermore, the CEBP

motif was associated with the most changed genes in $Vim^{-/-}$ MoAMs, and *Cebpb* was expressed

highly at both baseline and throughout IAV infection (**Figure 4f**). This observation is consistent with

Cebpb, a known macrophage TF that can also act in combination with AP1 factors (**Figure 4f),**

specifying poised regulatory elements in macrophages that activate genes on stimulation[45, 46].

On the other hand, ETS was the main TF family with motifs enriched in the least changed but not

the most changed genes (**Supplemental Figure 4d**). This family marginally includes PU.1, a well-

known myeloid lineage factor, but PU.1 is unlikely to have a specific role in IAV response

(**Supplemental Figure 4e**). Instead, the expression of *Elf1* and *Elf4* suggests that they may have

an interferon-independent antiviral role in influenza as previously reported [47], that is not impacted

by vimentin-deficiency.   In summary, vimentin-deficiency impedes the stimulus-dependent

regulation of pro-inflammatory genes in MoAMs during the Early Inflammatory phase of IAV

infection.   Although this effect leads to subtle changes in overall gene expression, it appears

sufficient to alter the MoAM function.

Vimentin-deficient MoAMs are sufficient to confer protection from IAV-induced mortality

We examined whether the loss of vimentin *in vitro* paralleled our *in vivo* observations of suppressed

pro-inflammatory gene expression (**Figure 3,4**). We polarized bone marrow-derived macrophages

(BMDMs) from WT and $Vim^{-/-}$ mice toward a pro-inflammatory (M1/M$_{(LPS)}$) phenotype or a reparative

(M2/ M$_{(IL-4)}$) phenotype through the administration of LPS or IL-4, respectively (**Figure 5a**). In line

with our *in vivo* observations, WT M1/M$_{(LPS)}$ macrophages had increased expression of pro-

inflammatory genes, such as *Tnf* and *Il1b*. The loss of vimentin suppressed the expression of M1

pro-inflammatory genes (**Figure 5b**) and increased the expression of M2/ M$_{(IL-4)}$ reparative genes

(*Mrc1, Arg1)* compared to WT macrophages (**Figure 5b**). Collectively, these data suggest that

vimentin plays a critical role in defining the macrophage pro-inflammatory phenotype.

To determine if vimentin plays a causal role in defining the pro-inflammatory function of recruited

MoAMs we used isochronic adoptive MoAM transfer experiments and molecular profiling in IAV-

infected mice. Bone marrow derived cells from WT or $Vim^{-/-}$ donor mice (CD45.2) were transferred

into irradiated WT recipient mice (CD45.1), allowing for the specific reconstitution of MoAMs with a

WT or $Vim^{-/-}$ genotype (**Figure 5c**). All mice showed reconstitution of donor hematopoietic cells in

peripheral blood above 85% (**Supplementary Figure 5a**). Following confirmed engraftment and

recovery, mice were infected with a lethal dose of IAV (**Figure 5c**). We found that IAV-infected WT

mice who received $Vim^{-/-}$ donor bone marrow exhibited significantly lower total protein concentration

in the BALF and lower wet-to-dry lung weight ratio than mice that received WT donor bone marrow,

indicating less severe alveolar damage (**Figure 5d-g**). Mice showed no difference in IAV replication

or clearance regardless of donor genotype (**Figure 5f**). Histopathological examination of IAV-

infected WT mice who received *Vim*$^{-/-}$ donor bone marrow demonstrated less severe alveolar

damage than those who received WT bone marrow (**Figure 5g**). As in *Vim*$^{-/-}$ mice (**Figure 1**), WT

mice with *Vim*$^{-/-}$ MoAMs exhibited lower levels of pro-inflammatory cytokines and chemokines

(**Figure 5h**). Taken together, these data support the notion that vimentin plays a causal role in

determining the pro-inflammatory function of recruited MoAM and drives IAV-induced lung injury.

Discussion

Monocytes are recruited to the lung during injury, where they differentiate into monocyte-derived alveolar macrophages (MoAMs) in response to cues provided by the injured microenvironment. We used a clinically relevant murine model of severe viral pneumonia to identify monocyte-derived alveolar macrophage-specific gene expression patterns associated with three distinct phases of IAV infection: Infiltrating, Early Inflammatory, and Late Inflammatory. WT MoAMs began expressing pro-inflammatory genes and secreting tissue-damaging pro-inflammatory cytokines in the Early Inflammatory phase, resulting in increased morbidity and mortality in influenza-infected wild-type mice. Conversely, upon entering the IAV-infected alveolar microenvironment, $Vim^{-/-}$ MoAMs failed to upregulate pro-inflammatory genes and suppressed the secretion of pro-inflammatory cytokines conferring protection to the host. By leveraging the $Vim^{-/-}$ mouse's protective phenotype, we investigated the transcriptional response following IAV infection and identified a core set of pro-inflammatory genes that drive lung injury and mortality. Although prior studies have examined macrophage gene expression at steady-state or single time points following IAV infection [48, 49], by analyzing gene expression over a time course post-IAV infection, our results identified critical temporal dynamics that shape the function of recruited monocyte-derived alveolar macrophages in response to respiratory infections.

During IAV-infection, infected cells, and cells dying from necroptosis or pyroptosis, release PAMPs and DAMPs, which activate signaling pathways to recruit additional effector cells, including MoAMs. We reason that as monocytes extravasate and infiltrate the lung parenchyma, the differentiation of WT MoAMs toward a pro-inflammatory phenotype involves activation of specific transcriptional programs in response to the IAV-infected lung microenvironment. Understanding how MoAMs communicate with resident cell populations to drive persistent inflammation (e.g., WT MoAMs) or promote reparative processes (e.g., $Vim^{-/-}$ MoAMs) represents an active area of investigation. During homeostasis, alveolar epithelial cells interact with tissue-resident macrophages via

receptor/ligand pairs CD200/CD200R, signal regulatory protein-α (SIRPα)/CD47, and CSF2R/GM-

CSF and immune/epithelial E-cadherin interactions [50, 51]. It follows that similar interactions occur

between alveolar epithelial cells and recruited MoAMs in the infected lung. These interactions, or

the disruption of these interactions, may activate specific transcriptional programs contributing to

the pro-inflammatory phenotype we observed in WT MoAMs. Investigating receptor/ligand

interactions and the secretion of molecules is limited by their short half-life and the requirement for

mass spectroscopy for their detection. Importantly, little information is known about specific

receptor/ligand interactions in recruited MoAMs during lung injury. We demonstrated that WT and

Vim^-/- MoAMs encountered similar environmental signals associated with IAV infection (e.g.,

equivalent viral load, similar immune cell recruitment – see Figure 1). Notably, *Vim^-/-* MoAMs

suppress pro-inflammatory genes and cytokines and maintain an increased expression of

reparative $M_{(IL-4)}$ genes throughout IAV infection. We hypothesize that the protective phenotype

displayed by recruited *Vim^-/-* MoAMs may be the result of disrupted or altered receptor/ligand

interactions in the IAV-infected lung microenvironment. It will be important to investigate how the

temporal transcriptional changes observed in our study impact the ability of recruited MoAMs to

interact with and respond to other cell types in the lung microenvironment.

In vitro studies have established vimentin's role in essential cellular processes, such as cell motility,

cell signaling, and protein complex scaffolding [52-55]. Retinoic acid-inducible gene I, a cytosolic

PRR that detects viral RNA and mediates type I interferon expression, leads to increased vimentin

expression[56]. Our data support this previous observation, as type, I interferon genes peaked on

day 2 p.i., preceding the increase in vimentin gene expression in WT MoAMs. We confirmed

previous reports that *vimentin-deficient* mice fail to secrete mature IL-1b as assembly and activation

of the NLRP3 inflammasome requires vimentin [18, 19]. Complementary to our findings, vimentin's

interaction with NOD2, a cytoplasmic PRR, activated NF-κB and induced a pro-inflammatory

immune response in an inflammatory bowel disease model [20]. Of note, the suppressed pro-

inflammatory response observed in *Vim^-/-* MoAMs does not appear to result from differing

macrophage maturation. In fact, genes involved in the transition from monocyte to macrophage-

like state, such as *Mafb*, were not differentially expressed between genotypes. Collectively, this

suggests that vimentin mediates the innate immune response(s) by regulating the expression of

pro-inflammatory genes.

Our findings demonstrate that depleting vimentin in MoAMs modulates the pro-inflammatory

response and protects the host, providing a rationale that vimentin could serve as a potential

therapeutic target during viral infection. Our group and others have shown that vimentin can be

therapeutically targeted in non-small cell lung cancer and other diseases such as idiopathic

pulmonary fibrosis [57, 58]. Molecules such as withaferin A, FiVe1, and simvastatin were used to

perturb the expression and/or assembly of vimentin filaments and led to the downregulation of

vimentin was accompanied by decreased cancer progression, decreased fibrosis and intermediate

filament disorganization, and cell death [58-61]. Collectively, these and our studies serve as a proof-

of-principle to support the development of pharmacological agents to target vimentin selectively.

These tools may be clinically beneficial in modulating the hyperinflammatory microenvironment of

the virus-infected lung.

Our study has several limitations. We used a well-established bone-marrow chimera mouse model

to demonstrate that *Vim^-/-* MoAMs failed to upregulate pro-inflammatory genes and secrete pro-

inflammatory cytokines, which was sufficient to confer protection from IAV-induced lung injury and

mortality. Additionally, we showed that *Vim^-/-* BMDMs expressed higher levels of M2/M$_{(IL-4)}$ markers

and lower levels of M1/M$_{(LPS)}$ markers, which appears to reflect the macrophage plasticity we

observed in our murine model of viral pneumonia. We caution the reader that while the framework

of M1/M2 polarization has provided a useful system to study macrophages in vitro, it is unlikely that

this occurs in tissue. This was shown by Xue *et al*., who found that the M1/M2 paradigm failed to

describe the transcriptome of human monocyte-derived and alveolar macrophages stimulated with

LPS/IFN-γ or IL-4/IL-13 in the presence of factors that recapitulate different tissue or disease

microenvironments [62]. Future studies could be improved by using a lineage mark to track the

recruitment and phenotype changes associated with recruited MoAMs *in vivo*. By generating a

Vim$^{fl/fl}$Cx3cr1^{CreERT2}zsGreen transgenic mouse, we could selectively delete vimentin and track the

recruitment of *GFP$^+$Vim$^{-/-}$* alveolar macrophages during IAV infection. Regrettably, the vimentin-

floxed mouse is not yet available. Second, our RNA-seq data identified distinct transcriptional

phases of the IAV response in MoAMs; however, it is unclear whether different MoAM

subpopulations enter the lung at different time points from a single compartment. For example,

MoAMs may differentiate directly from circulating monocytes that extravasate from the pulmonary

capillaries into the alveolar space. Alternatively, MoAMs may stem from interstitial macrophages

already residing in the lung at steady-state. Using a *Vim$^{fl/fl}$Cx3cr1^{CreERT2}zsGreen* mouse, pulsed

lineage tracing would address these important questions.

In summary, our results define the temporal, transcriptional dynamics in recruited alveolar

macrophages that shape the response to respiratory infection and demonstrate that vimentin plays

an essential role in modifying macrophage phenotype and function. Moreover, we defined a core

set of pro-inflammatory genes whose upregulation during the early response to IAV drives lung

injury and mortality in wild-type, but not *Vim$^{-/-}$* mice. We show that vimentin-deficient recruited

alveolar macrophages exhibit a dampened inflammatory response in a cell-autonomous manner,

sufficient to confer protection from IAV-induced mortality. Our data demonstrate that vimentin plays

a causal role in shaping the pro-inflammatory response in recruited MoAMs, promoting virus-

induced lung injury following IAV infection.

Materials & Methods

In silico single-cell RNA-seq data analysis. We queried a single-cell RNA-seq transcriptomic data set derived from bronchoalveolar lavage samples from 10 patients with severe SARS-CoV-2 viral pneumonia[1]. We downloaded normalized counts tables and visualized the pre-determined cell population clusters using Scanpy [63]. 40,000 single cells representing the four monocyte-derived alveolar macrophages (MoAMs) populations were visualized as a UMAP. Mean expression data of individual genes per cell population cluster were visualized as dot plots in Scanpy.

Mice. All experiments were performed using wild-type 129S or vimentin-null ($Vim^{-/-}$) 129S/V mice, age 8-12 weeks. Wild-type mice were purchased from Taconic, $Vim^{-/-}$ mice were a gift from Albee Messing (University of Wisconsin, Madison, WI). All animals were kept under pathogen-free conditions, and all experiments were approved in accordance with Northwestern University's Institutional Animal Care and Use Committee (IACUC) guidelines.

Bone marrow chimeras. 8-12 week old recipient mice (Ptprc^a/Pep-Boy, CD45.1) were lethally irradiated with a single dose of 1,000 cGy γ-radiation using a Cs-137–based Gammacell-40 irradiator (Nordion). Within 24 hours, fresh bone marrow cells (2×10^6) from either WT donor mice or $Vim^{-/-}$ donor mice (CD45.2) were injected via retro-orbital injection. Mice were housed together in individually ventilated cages and were given *ad libitum* access to food and water supplemented with Trimetoprim/Sulfamethoxazole (TMP/SMX 40/8 mg formula, Hi-Tech Pharmacal) for 4 weeks and then switched to standard housing regimen. Reconstituted mice were used for influenza infection experiments ~4 weeks after transplantation. Engraftment was assessed after two weeks, using flow cytometry to measure the percentage of CD45.1+ and CD45.2+ cells in peripheral blood collected from the facial vein. All mice used for experiments showed reconstitution of the donor (CD45.2) circulating leukocytes in peripheral blood above 85%.

Bone marrow derived macrophage (BMDM) polarization and RT-PCR. BMDMs were isolated from wild-type 129S or vimentin-null ($Vim^{-/-}$) 129S/V mice, age 8-12 weeks, and treated with 100

498 ng/mL LPS or 20 ng/mL IL-4 to induce $M_{(LPS)}$ and $M_{(IL-4)}$ polarization, respectively. Cells were lysed

499 after 24 hours and RNA was extracted using RNeasy Mini kit (Qiagen) according to manufacturer's

500 protocol. RNA concentration was assessed using a NanoDrop platform (Thermo Fisher). RT-PCR

501 was run to assess gene expression changes (BioRad CFX).

502 *Virus*. H1N1 Influenza virus strain A/WSN/1933 (WSN) was propagated in fertile chicken eggs

503 (Sunnyside Hatchery) as previously described[4, 5]. Viral titers were determined by measuring

504 plaque-forming units (pfu) in Madin-Darby canine kidney epithelial cells (ATCC).

505 **Plaque Assay.** Confluent monolayers of Madin-Darby canine kidney epithelial cells were infected

506 with stock WSN virus in 1% BSA DMEM for 2h at $37°C$. Plates were washed with PBS and an

507 overlay of 50% 2x replacement media (2x DMEM, 0.12M $NaHCO_3$, 2% penicillin-streptomycin, and

508 1% HEPES), 50% Avicel (2.35%), and N-acetyl trypsin ($1.5\mu g/mL$) remained on the cells for 72h at

509 $37°C$. Plates were washed with PBS and fixed with 0.2% PFA before staining the monolayers with

510 naphthalene blue-black stain.

511 **In vivo influenza virus infection.** 8-12 week old WT and $Vim^{-/-}$ 129S/V male mice were

512 anesthetized using isoflurane and intratracheally (i.t.) infected with 200 pfu WSN. Mice were

513 monitored daily, and weight was recorded every 24 hours.

514 *Histology*. Mice were euthanized, after which a 20-gauge angiocath was sutured into the trachea.

515 Heart and lungs were removed *en bloc*, and lungs were inflated with 0.8mL of 4% paraformaldehyde

516 (PFA) at a pressure not exceeding 16 cm H_2O. Tissue was fixed and embedded in paraffin. 5 μm

517 sections were stained with H&E by the Mouse Histology Phenotyping Laboratory (Northwestern

518 University, Chicago, IL). Tissue sections were visualized using TissueGnostics, a Tissue/Cell High

519 Throughput Imaging Analysis Systems (Vienna, Austria) and captured using TissueFAXS software

520 (TissueGnostics, Los Angeles, CA) at the Northwestern University Cell Imaging Facility

521 (Northwestern University, Chicago, IL).

Bronchoalveolar Lavage. A 20-gauge angiocath was ligated into the trachea, and 1mL of sterile

PBS was instilled into the lungs, then removed through the angiocath. This process was repeated

three times. Bronchoalveolar lavage fluid (BALF) was centrifuged at 500 x *g* for 10 min Protein

levels in the supernatant were measured by Bradford assay (Bio-Rad), and cytokine levels were

measured using a custom multiplex ELISA (ThermoFisher Scientific).

Wet-to-dry weight ratio. Mice were euthanized, and lungs were surgically removed *en bloc*. The

left lung was ligated, excised, and weighed in a tared container. The left lung was dried at 70°C in

a Speed-Vac SC100 evaporator (Thermo Scientific, Waltham, MA) until a constant weight was

obtained, and the wet-to-dry weight ratio was calculated.

Flow cytometry analysis of lung cell populations. Mice were euthanized, and lungs were

perfused with 10mL HBSS with Ca^{2+} and Mg^{2+}. Lung loves were removed and inflated with enzyme

solution (5mL of 0.2 mg/mL DNase I and 2 mg/mL collagenase D in HBSS with Ca^{2+} and Mg^{2+})

using a 30-gauge needle. According to the manufacturer's instructions, tissue was minced and

processed in gentleMACS dissociator tubes (Miltenyi Biotec). Processed lungs were passed

through a 40μm cell strained, and RBCs were lysed using BD PharmLyse (BD Biosciences, San

Jose, CA). The remaining cells were counted using a Countess cell counter (Invitrogen, Grand

Island, NY). CD45 microbeads were added, and cells were eluted according to the Miltenyi

manufacturer's instructions. Cells were stained with viability dye Aqua (Invitrogen), and non-specific

antibody (Ab) binding was inhibited by adding Fc Block (553142, clone 2.4G2; BD Pharmingen).

Blocked cells were stained with a mixture of fluorochrome-conjugated Abs (see **Online**

Supplemental Table 1 for list of Abs). Data were acquired on a BD LSR II flow cytometer using BD

FACSDiva software (BD Biosciences), and data analyses were performed using FlowJo software

(Tree Star, Ashland, OR). Cell populations were identified using a sequential gating strategy

(**Supplemental Figure 1b)**. Cell number was normalized to body weight.

Macrophage harvesting and flow sort. Mice were euthanized and perfused with 5 mL sterile PBS by cardiac perfusion through the right ventricle. A 20-gauge angiocath was ligated into the trachea, and the lung and trachea were removed *en bloc*. Lungs were instilled with 1 mL dispase (Corning) through the angiocath, and a second suture was tied around the trachea to avoid leakage. Lungs were placed in 50 mL tubes (Falcon) containing 750 μL cold dispase and rocked for 45 min at RT. Trachea and angiocath were removed, and lungs were minced using curved-edge scissors. Digested tissue was poured into 50 mL tubes containing 7 mL DMEM (Corning) and 14 μL DNase I (50 mg/mL, Sigma). Tubes were rocked for 10 minutes at RT. Using serological pipettes, tissue was gently further disrupted and subsequently filtered through 70 μm cell strainers and 40 μm strainers (Corning). Tubes were rinsed with DMEM + 5% FBS and filtered through the same strainers. Cell suspensions were centrifuged for 10 min at 1300 rpm, 4°C and pellets were resuspended in 350 μL MACS buffer (Milteny). 100 μL CD11c magnetic microbeads (Milteny) were added, and cells were eluted according to the Miltenyi manufacturer's instructions. Cells were stained for CD11b (Biolegend), Siglec F (BD), Ly6C (eBioscience), CD45 (eBioscience), MHCII (Biolegend), CD64 (Biolegend), Ly6G (BD), and NK1.1(BD). Viability dye (eFluor506, eBioscience) was used to sort living cells only (**Online Supplemental Table 1**). Using sequential gating (**Supplemental Figure 2b),** recruited alveolar macrophages (MoAMs) and tissue-resident alveolar macrophages (TRAMs) were flow-sorted using a BD FACSaria III 5-laser cell sorter and 100 μm nozzle (BD Biosciences). Of note, MoAMs were collected in the same gate as interstitial macrophages (Ly6C-, CD64+); however, by 2 days p.i., interstitial macrophages were no longer present in this gate (see **Online Supplemental Figure 2**). Flow-sorted cells were collected into 1.5 mL Eppendorf tubes containing 200 μL MACS buffer and centrifuged at 5,000 rpm for 5 min at RT. Pellets were resuspended and lysed in RLT lysis buffer (Qiagen) and stored at -80°C until all time points were collected.

RNA sequencing (RNA-seq). Samples collected from infected mice that at flow sort appeared

uninfected (had no infiltrating MoAM) were excluded from RNA isolation. RNA was isolated in all

samples at the same time to minimize the batch effect. All libraries for a given cell type were

prepared as a single batch for sequencing. Total RNA was isolated using RNeasy Plus Mini kits

(Qiagen). RNA quality (RIN) and quantity were assessed using a TapeStation 4200 (Agilent

Technologies). RIN scores ranged from 7.2 to 9.4, with an average of 8.6. Using poly(A) mRNA

magnetic enrichment kits from NEBNext (New England Biolabs), mRNA was isolated. cDNA

libraries were prepared using NEBNext Ultra DNA library kits (New England BioLabs) for Illumina.

Libraries were sequenced using high-output 75-cycle single-end kits on an Illumina NextSeq 500

sequencer to an average depth of 4.1 million reads per sample. Reads were demultiplexed

(bcl2fastq), trimmed, and aligned to reference genome mm10 using TopHat2[64]. Aligned reads

were mapped to genes using HTSeq with an Ensembl annotation (version 78)[65].

RNA-seq analysis. For k-means clustering, we used gene counts normalized to FPKM (Fragments

per Kilobase of transcript per Million mapped reads). Low counts were removed, with a cut-off set

as row sum >27 (# of samples used for analysis), yielding a gene counts table of 9,169 expressed

genes. One-way ANOVA with Benjamini-Hochberg (BH) FDR correction was used to define

differentially expressed genes over time. Genes with an FDR <0.05 were considered differentially

expressed, resulting in 2,623 DEGs. Clustering was performed on the 2,623 DEGs using k-means

clustering (GENE-E, https://software.broadinstitute.org/GENE-E/index.html).

For pairwise comparison, low counts were trimmed if CPM <2 in all replicates in the smallest group

of the comparison. EdgeR was used to define pairwise differentially expressed genes between WT

and *Vim*[-/-] MoAMs[66, 67]. Principal Component analysis (PCA) plots were generated using *pca*

function in R. Pearson correlation matrix, and heatmaps were generated using GENE-E. Gene

ontology (GO) analysis was run using Gorilla[68, 69]. Expressed genes, filtered for low counts

(9,169) were used as background. Gene set enrichment analysis (GSEA) was run using the

weighted pre-ranked setting based on the $\log_2$FC reported by edgeR [70, 71]. Gene sets associated

with specific processes were downloaded from KEGG (Cell Cycle) or Hallmark pathways

(Glycolysis, Oxidative Phosphorylation, IFN alpha (Type I interferon), IFNγ (Type II Interferon)

except for the custom gene sets described below.   The 85 overlapping genes between the IFNα

and IFNγ Hallmark pathways were excluded from the IFNγ gene set. Overlaps between gene lists

were visualized using Venny (https://bioinfogp.cnb.csic.es/tools/venny/index.html).

Custom gene set generation. Custom gene sets for LPS/IFNγ and IL-4 stimulated were generated

by querying primary published data available in relevant experimental models. The final curated

gene sets, $M_{(LPS/IFN\gamma)}$ and $M_{(IL-4)}$, were derived from four primary publications [25, 39-41]. All four

publications used mouse bone-marrow-derived macrophages (BMDM) and stimulated them *in vitro*

with either LPS+IFNγor IL-4. Methodologies used included LC-ESI-MS/MS mass spectrometry,

microarray, and RNA-seq. A summary of the experimental approach, methods, and results for all

publications can be found in **Online Supplemental Table 2**. We included all genes that were

upregulated in at least one data set and could be converted into a unique Ensembl ID. Three genes

were found in both $M_{(LPS/IFN\gamma)}$ and $M_{(IL-4)}$ and were removed to avoid any overlap between gene sets.

These trimming steps resulted in two annotated gene sets of 110 LPS/IFNγ stimulated genes and

101 IL-4 stimulated genes (**Online Supplemental Table 2**).

Motif finding. We generated two gene sets of the 186 most and 254 least changed genes in Vim^{-}

$^{/-}$ MoAM by splitting the 440 genes of the Early Inflammatory cluster (Figure 2) based on whether

their $\log_2(Vim^{-/-}/WT)$ expression on day 3 p.i. was above or below the mean of -0.23. (**Figure 6d,e)**.

To identify motifs that were enriched in either of these gene sets, we ran findmotifs.pl from the

HOMER package with default parameters and the mouse mm10 genome as background [45]. We

identified significantly enriched motifs as those with p-value < 0.05 in the known motif results.

Viral reads. To identify viral reads within libraries, unmapped reads were aligned to the H1N1

reference genome. Reads for the H1N1 matrix protein 2 (M2), nucleoprotein (NP), and

neuraminidase (NA) were normalized by sample to the number of uniquely aligned (mm10) reads.

Statistics*.* Analyses for cytokine levels, BALF protein, viral titer, and wet-to-dry ratio were

performed using GraphPad Prism (GraphPad Software). P values were calculated using a two-

tailed unpaired t-test assuming equal variance.

Acknowledgments

Histology services were provided by the Northwestern University Mouse Histology and Phenotyping

Laboratory, which is supported by NCI P30-CA060553 awarded to the Robert H Lurie

Comprehensive Cancer Center. Flow Cytometry was supported by the Northwestern University

RHLCCC Flow Cytometry Facility and a Cancer Center Support Grant (NCI CA060553). This

research was supported in part through the computational resources and staff contributions

provided by the Genomics Compute Cluster which is jointly supported by the Feinberg School of

Medicine, the Center for Genetic Medicine, and Feinberg's Department of Biochemistry and

Molecular Genetics, the Office of the Provost, the Office for Research, and Northwestern

Information Technology. The Genomics Compute Cluster is part of Quest, Northwestern

University's high performance computing facility, with the purpose to advance research in

genomics. Graphic design was created with BioRender.com. Kishore R. Anekalla was supported

by NIH T32 HL076139. Shang-Yang Chen was supported by a pre-doctoral AHA award

(19PRE34380200). Alexander V. Misharin was supported by N.I.H. grants U19AI135964,

P01AG049665, R01HL153312, and NUCATS COVID-19 Rapid Response Grant. Karen M. Ridge

was supported by N.I.H. grants P01AG049665 and P01GM096971. Bria M. Coates was supported

by funds provided by the Manne Research Institute COVID-19 Springboard Exploratory Research

Award and N.I.H. grant K08HL143127. Deborah R. Winter `was supported by the American Lung

Association, Arthritis National Research Foundation, American Thoracic Society, Scleroderma

Foundation, American Federation for Aging Research, American Heart Association

18CDA34110224, and NIH/NIAID U19AI135964. G.R.S.B. was supported by NIH grants

U19AI135964, P01AG049665, P01AG04966506S1, R01HL147575 and Veterans Affairs grant

I01CX001777.

Author contributions: C.M.K., K.R.A., Y.H., B.M.C., A.V.M., D.R.W., K.M.R., contributed to the

acquisition, analysis, and interpretation of the data and drafting the manuscript.

M.C., G.G., S.C., M.T., Y.C., J.M.D., contributed to the acquisition and interpretation of data.

C.M.K., K.R.A., Y.H., B.M.C., H.A.V., P.A.R., A.V.M., G.R.S.B., D.R.W., K.M.R. contributed to the

conception and design of the study and interpretation of the data. All authors critically revised and

approved the final version of the manuscript.

655
656

References

1. Grant, R.A., et al., *Circuits between infected macrophages and T cells in SARS-CoV-2 pneumonia.* 2021.
2. Matthay, M.A., et al., *Acute respiratory distress syndrome.* 2019. **5**(1): p. 18.
3. Calfee, C.S., et al., *Subphenotypes in acute respiratory distress syndrome: latent class analysis of data from two randomised controlled trials.* Lancet Respir Med, 2014. **2**(8): p. 611-20.
4. Coates, B.M., et al., *Inflammatory Monocytes Drive Influenza A Virus-Mediated Lung Injury in Juvenile Mice.* J Immunol, 2018. **200**(7): p. 2391-2404.
5. Misharin, A.V., et al., *Monocyte-derived alveolar macrophages drive lung fibrosis and persist in the lung over the life span.* Journal of Experimental Medicine, 2017. **214**(8): p. 2387-2404.
6. Budinger, G.R.S., et al., *Distinctive features of severe SARS-CoV-2 pneumonia.* 2021. **131**(14).
7. Yona, S., et al., *Fate Mapping Reveals Origins and Dynamics of Monocytes and Tissue Macrophages under Homeostasis.* Immunity, 2013. **38**(1): p. 79-91.
8. Wang, Z., et al., *Early hypercytokinemia is associated with interferon-induced transmembrane protein-3 dysfunction and predictive of fatal H7N9 infection.* 2014. **111**(2): p. 769-74.
9. Yu, L., et al., *Clinical, virological, and histopathological manifestations of fatal human infections by avian influenza A(H7N9) virus.* 2013. **57**(10): p. 1449-57.
10. Guo, J., et al., *The Serum Profile of Hypercytokinemia Factors Identified in H7N9-Infected Patients can Predict Fatal Outcomes.* 2015. **5**: p. 10942.
11. Tate, M.D., et al., *Reassessing the role of the NLRP3 inflammasome during pathogenic influenza A virus infection via temporal inhibition.* Sci Rep, 2016. **6**: p. 27912.
12. Coates, B.M., et al., *Inhibition of the NOD-Like Receptor Protein 3 Inflammasome Is Protective in Juvenile Influenza A Virus Infection.* Front Immunol, 2017. **8**: p. 782.
13. Dawson, T.C., et al., *Contrasting effects of CCR5 and CCR2 deficiency in the pulmonary inflammatory response to influenza A virus.* Am J Pathol, 2000. **156**(6): p. 1951-9.
14. Lin, K.L., et al., *CCR2-antagonist prophylaxis reduces pulmonary immune pathology and markedly improves survival during influenza infection.* J Immunol, 2011. **186**(1): p. 508-15.
15. Gaurav, R., et al., *IL-33 Depletion in COVID-19 Lungs.* 2021. **160**(5): p. 1656-1659.
16. Nimmo, E.R., et al., *TLE1 modifies the effects of NOD2 in the pathogenesis of Crohn's disease.* 2011. **141**(3): p. 972-981.e1-2.
17. Mor-Vaknin, N., et al., *Murine colitis is mediated by vimentin.* Sci Rep, 2013. **3**: p. 1045.
18. dos Santos, G., et al., *Vimentin regulates activation of the NLRP3 inflammasome.* Nat Commun, 2015. **6**: p. 6574.
19. Lang, T., et al., *Macrophage migration inhibitory factor is required for NLRP3 inflammasome activation.* Nat Commun, 2018. **9**(1): p. 2223.
20. Stevens, C., et al., *The intermediate filament protein, vimentin, is a regulator of NOD2 activity.* Gut, 2013. **62**(5): p. 695-707.
21. Kamphuis, W., et al., *GFAP and vimentin deficiency alters gene expression in astrocytes and microglia in wild-type mice and changes the transcriptional response of reactive glia in mouse model for Alzheimer's disease.* Glia, 2015. **63**(6): p. 1036-56.
22. Abdulhasan, M., et al., *Stress Decreases Host Viral Resistance and Increases Covid Susceptibility in Embryonic Stem Cells.* 2021. **17**(6): p. 2164-2177.
23. Bucher, H., et al., *Neutralization of both IL-1alpha/IL-1beta plays a major role in suppressing combined cigarette smoke/virus-induced pulmonary inflammation in mice.* Pulmonary Pharmacology & Therapeutics, 2017. **44**: p. 96-105.

707   24.   Liu, M., et al., *C1q/TNF-related protein-9 promotes macrophage polarization and improves*
708   *cardiac dysfunction after myocardial infarction.* 2019. **234**(10): p. 18731-18747.
709   25.   Gensel, J.C., et al., *Predictive screening of M1 and M2 macrophages reveals the*
710   *immunomodulatory effectiveness of post spinal cord injury azithromycin treatment.* Sci
711   Rep, 2017. **7**: p. 40144.
712   26.   Cai, B., et al., *MerTK signaling in macrophages promotes the synthesis of inflammation*
713   *resolution mediators by suppressing CaMKII activity.* Sci Signal, 2018. **11**(549).
714   27.   Xu-Vanpala, S., et al., *Functional heterogeneity of alveolar macrophage population based*
715   *on expression of CXCL2.* Sci Immunol, 2020. **5**(50).
716   28.   Herold, S., et al., *Influenza virus-induced lung injury: pathogenesis and implications for*
717   *treatment.* Eur Respir J, 2015. **45**(5): p. 1463-78.
718   29.   Imai, T., et al., *Identification and molecular characterization of fractalkine receptor*
719   *CX3CR1, which mediates both leukocyte migration and adhesion.* Cell, 1997. **91**(4): p.
720   521-30.
721   30.   Worthylake, R.A. and K. Burridge, *RhoA and ROCK promote migration by limiting*
722   *membrane protrusions.* J Biol Chem, 2003. **278**(15): p. 13578-84.
723   31.   Lammermann, T., et al., *Rapid leukocyte migration by integrin-independent flowing and*
724   *squeezing.* Nature, 2008. **453**(7191): p. 51-5.
725   32.   Starnes, T., et al., *The chemokine CXCL14 (BRAK) stimulates activated NK cell migration:*
726   *implications for the downregulation of CXCL14 in malignancy.* Exp Hematol, 2006. **34**(8):
727   p. 1101-5.
728   33.   Watanabe, S., et al., *The role of macrophages in the resolution of inflammation.* 2019.
729   **129**(7): p. 2619-2628.
730   34.   Kratofil, R.M., P. Kubes, and J.F. Deniset, *Monocyte Conversion During Inflammation and*
731   *Injury.* Arterioscler Thromb Vasc Biol, 2017. **37**(1): p. 35-42.
732   35.   Sanchez, K.K., et al., *Cooperative Metabolic Adaptations in the Host Can Favor*
733   *Asymptomatic Infection and Select for Attenuated Virulence in an Enteric Pathogen.* 2018.
734   **175**(1): p. 146-158.e15.
735   36.   Smallwood, H.S., et al., *Targeting Metabolic Reprogramming by Influenza Infection for*
736   *Therapeutic Intervention.* Cell Reports, 2017. **19**(8): p. 1640-1653.
737   37.   Singer, B.D. and N.S. Chandel, *Immunometabolism of pro-repair cells.* 2019. **129**(7): p.
738   2597-2607.
739   38.   Ivashkiv, L.B., *IFNy: signalling, epigenetics and roles in immunity, metabolism, disease*
740   *and cancer immunotherapy.* Nat Rev Immunol, 2018. **18**(9): p. 545-558.
741   39.   Jablonski, K.A., et al., *Novel Markers to Delineate Murine M1 and M2 Macrophages.* PLoS
742   One, 2015. **10**(12): p. e0145342.
743   40.   Hans, C.P., et al., *Transcriptomics Analysis Reveals New Insights into the Roles of Notch1*
744   *Signaling on Macrophage Polarization.* 2019. **9**(1): p. 7999.
745   41.   Becker, L., et al., *Unique proteomic signatures distinguish macrophages and dendritic*
746   *cells.* PLoS ONE [Electronic Resource], 2012. **7**(3): p. e33297.
747   42.   Gosselin, D., et al., *Environment drives selection and function of enhancers controlling*
748   *tissue-specific macrophage identities.* Cell, 2014. **159**(6): p. 1327-40.
749   43.   Ostuni, R., et al., *Latent enhancers activated by stimulation in differentiated cells.* Cell,
750   2013. **152**(1-2): p. 157-71.
751   44.   Lavin, Y., et al., *Tissue-resident macrophage enhancer landscapes are shaped by the*
752   *local microenvironment.* Cell, 2014. **159**(6): p. 1312-26.
753   45.   Heinz, S., et al., *Simple combinations of lineage-determining transcription factors prime*
754   *cis-regulatory elements required for macrophage and B cell identities.* 2010. **38**(4): p. 576-
755   89.
756   46.   Heinz, S., et al., *Effect of natural genetic variation on enhancer selection and function.*
757   Nature, 2013. **503**(7477): p. 487-92.

47. Seifert, L.L., et al., *The ETS transcription factor ELF1 regulates a broadly antiviral program distinct from the type I interferon response.* PLoS Pathog, 2019. **15**(11): p. e1007634.

48. Zhang, J., et al., *Two waves of pro-inflammatory factors are released during the influenza A virus (IAV)-driven pulmonary immunopathogenesis.* 2020. **16**(2): p. e1008334.

49. Tate, M.D., et al., *Critical role of airway macrophages in modulating disease severity during influenza virus infection of mice.* Journal of Virology, 2010. **84**(15): p. 7569-80.

50. Hussell, T. and T.J. Bell, *Alveolar macrophages: plasticity in a tissue-specific context.* 2014. **14**(2): p. 81-93.

51. Bissonnette, E.Y., et al., *Cross-Talk Between Alveolar Macrophages and Lung Epithelial Cells is Essential to Maintain Lung Homeostasis.* 2020. **11**: p. 583042.

52. Li, Q.F., et al., *Critical role of vimentin phosphorylation at Ser-56 by p21-activated kinase in vimentin cytoskeleton signaling.* J Biol Chem, 2006. **281**(45): p. 34716-24.

53. Ivaska, J., et al., *Novel functions of vimentin in cell adhesion, migration, and signaling.* Exp Cell Res, 2007. **313**(10): p. 2050-62.

54. Mendez, M.G., S. Kojima, and R.D. Goldman, *Vimentin induces changes in cell shape, motility, and adhesion during the epithelial to mesenchymal transition.* FASEB J, 2010. **24**(6): p. 1838-51.

55. Yoon, M., et al., *Motile properties of vimentin intermediate filament networks in living cells.* J Cell Biol, 1998. **143**(1): p. 147-57.

56. de Rivero Vaccari, J.P., et al., *Astrogliosis involves activation of retinoic acid-inducible gene-like signaling in the innate immune response after spinal cord injury.* Glia, 2012. **60**(3): p. 414-21.

57. Richardson, A.M., et al., *Vimentin Is Required for Lung Adenocarcinoma Metastasis via Heterotypic Tumor Cell-Cancer-Associated Fibroblast Interactions during Collective Invasion.* 2018. **24**(2): p. 420-432.

58. Berr, A.L., et al., *Vimentin is Required for Tumor Progression and Metastasis in a Mouse Model of Non-Small Cell Lung Cancer.* bioRxiv, 2020: p. 2020.06.04.130963.

59. Bargagna-Mohan, P., et al., *Withaferin A effectively targets soluble vimentin in the glaucoma filtration surgical model of fibrosis.* 2013. **8**(5): p. e63881.

60. Bollong, M.J., et al., *A vimentin binding small molecule leads to mitotic disruption in mesenchymal cancers.* 2017. **114**(46): p. E9903-E9912.

61. Trogden, K.P., et al., *An image-based small-molecule screen identifies vimentin as a pharmacologically relevant target of simvastatin in cancer cells.* 2018. **32**(5): p. 2841-2854.

62. Xue, J., et al., *Transcriptome-based network analysis reveals a spectrum model of human macrophage activation.* 2014. **40**(2): p. 274-88.

63. Wolf, F.A., P. Angerer, and F.J. Theis, *SCANPY: large-scale single-cell gene expression data analysis.* Genome Biol, 2018. **19**(1): p. 15.

64. Kim, D., et al., *TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions.* 2013. **14**(4): p. R36.

65. Anders, S., P.T. Pyl, and W. Huber, *HTSeq--a Python framework to work with high-throughput sequencing data.* Bioinformatics, 2015. **31**(2): p. 166-9.

66. Robinson, M.D., D.J. McCarthy, and G.K. Smyth, *edgeR: a Bioconductor package for differential expression analysis of digital gene expression data.* Bioinformatics, 2010. **26**(1): p. 139-40.

67. McCarthy, D.J., Y. Chen, and G.K. Smyth, *Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation.* Nucleic Acids Res, 2012. **40**(10): p. 4288-97.

68. Eden, E., et al., *Discovering motifs in ranked lists of DNA sequences.* PLoS Comput Biol, 2007. **3**(3): p. e39.

69. Eden, E., et al., *GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists.* BMC Bioinformatics, 2009. **10**: p. 48.

70.   Mootha, V.K., et al., *PGC-1alpha-responsive genes involved in oxidative phosphorylation*
are coordinately downregulated in human diabetes. 2003. **34**(3): p. 267-73.
71.   Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*
interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43): p.
15545-50.

814
815

Figure Legends

Figure 1. Vimentin-null mice are protected from influenza-induced lung injury and mortality.
a) Uniform Manifold Approximation and Projection (UMAP) of 40,248 cells annotated as monocyte-
derived alveolar macrophages (MoAMs) obtained from 10 patients with severe COVID-19 disease
within 48 hours after intubation. **b)** Fraction of cells expressing *VIM* and mean expression in each
MoAM cluster. **c)** Visualization of *IL1-B, CCL3, CXCL2, C1QA*, and *MERTK* expression in MoAM
cells using density projection plots. Scale bar represents expression averaged within hexagonal
areas on UMAP. **d)** Fraction of cells expressing the representative genes and their mean expression
in each MoAM cluster. **e)** Survival of wild-type (WT) and $Vim^{-/-}$ mice intratracheally infected with a
lethal dose of influenza A virus (IAV) (200pfu) was measured and compared. Survival curves were
analyzed using log-rank test, p-value =0.001. Data shown are n=12 per group. **f)** Lung wet-to-dry
ratio measured at day 3 and 5 p.i. (n = 8-11/timepoint) **g)** Bronchoalveolar lavage fluid (BALF)
protein levels measured by Bradford assay, n=5/timepoint. In **f** and **g**, differences between
genotypes on each day were assessed by t-test. p-values were 0.0015 and <0.0001, for 3 and 5
days p.i., respectively. **h)** Representative images of murine lung architecture at day 7 p.i. assessed
by H&E staining, scale bar 100 um. **i**) Viral titer assessed by plaque assay. The difference between
genotypes was statistically not significant. n=5-9/timepoint. **j)** Flow cytometric analysis of tissue-
resident (TRAMs) and infiltrating monocyte-derived alveolar macrophages (MoAMs) out of the total
lung CD45+ cell population normalized to body weight during the IAV response, n=4/timepoint. **k)**
Levels of inflammatory cytokines in BALF were assessed using a custom multiplex ELISA. n=3-
4/timepoint. Bar graphs represent mean, error bars represent SD, differences between genotypes
on each day were assessed by t-test. * p<0.05

Supplemental Figure 1. **a**) UMAP of 77,146 cells annotated by cell type obtained from 10 patients
with severe COVID-19 disease within 48 hours after intubation. **b)** Quantification of H1N1 viral RNA
molecules detected among RNA isolated from MoAMs, normalized to the number of reads uniquely
aligned to the mouse genome (mm10) for each sample. Bar graphs represent mean, error bars
represent SD. **c)** Gating strategy used for flow cytometry analysis of neutrophils, TRAMs, interstitial
macrophages (IM), and MoAMs. **d)** Flow cytometric analysis demonstrated no differences in the
number of neutrophils out of the total lung CD45+ cell population normalized to body weight
between $Vim^{-/-}$ and WT mice during the IAV response.

Figure 2. Wild-type monocyte-derived alveolar macrophages exhibit gene expression
patterns associated with three phases of influenza A viral infection. a) Heatmap of normalized
expression of 2,623 differentially expressed genes (ANOVA p-value <0.05). Z-score normalization
for each row was applied to gene expression (FPKM) data. K-means clustering (k=3) identified
three clusters: Infiltrating (1131 genes), Early Inflammatory (440 genes), Late Inflammatory (1052
genes). **b)** Mean z-scores calculated across genes for each cluster **c)** Gene Ontology (GO) term
enrichment for each phase. **d)** Bar graphs of gene expression (FPKM) for representative genes in
each phase. Error bars represent SD. (ANOVA p-value <0.05). **e)** Bubble plot representing the
proportion of genes (%) within each process that had an expression above the row mean. Color
gradient represents the mean z-score of expression across genes. **f)** Enrichment of gene sets in
each cluster as defined in A. Asterisk represents significantly enriched overlaps (hypergeometric p-
value <0.0018, FWER = 0.05). The dotted line indicates an expected enrichment score of 1. n=3-4
per group. Ox Phos = Oxidative phosphorylation.

Supplemental Figure 2. a) Schematic of experimental design and workflow. WT or $Vim^{-/-}$ mice
were infected with a lethal dose of influenza A virus (200 pfu). MoAMs were sorted by flow cytometry
(FACS) from lungs on 2, 3, 4, and 6 days p.i. Total RNA was isolated, mRNA was enriched, and
bulk RNA-seq was performed. Principal component analysis (PCA) was used to visualize replicates.

b) Sequential gating strategy used for FACS collection of TRAMs and MoAMs **c)** Violin plots of log$_2$(FPKM+1) counts data for all genes in each cluster. Solid lines represent median, dotted lines represent quartiles. **d)** Bar graphs of gene expression (FPKM) in MoAMs for representative genes in each phase. Error bars represent SD. (ANOVA p-value <0.05).

Figure 3. *Vim$^{-/-}$* MoAMs modulate expression of Infiltrating and Early Inflammatory, but not Late Inflammatory phase genes. a) Volcano plots of differentially expressed genes (edgeR p-value<0.05, and edgeR reported log$_2$FC>|1.0|) between *Vim$^{-/-}$* and WT MoAMs on each day post-infection. The number of genes downregulated (blue) and upregulated (red) in *Vim$^{-/-}$* compared to WT MoAMs are shown. **b)** GO term enrichment across days p.i. in the downregulated (blue) and upregulated (red) genes in *Vim$^{-/-}$* compared to WT MoAMs. Scale represents -log$_{10}$ p-value. **c-e)** Line graphs on the left represent mean adapted z-scores, calculated across genes for each cluster in WT MoAMs and *Vim$^{-/-}$* MoAMs. Shading represents WT MoAMs inter-quartile range (IQR). Dot plots to the right represent % of genes at each timepoint expressed above their WT mean. Each dot represents a replicate. Lines represent the median **f)** % genes in each cluster that exhibit decreased or increased expression in *Vim$^{-/-}$* MoAMs compared to WT MoAMs (n=3-4 per group). Decreased genes defined as log$_2$FC < -0.58 (FC of 1.5); increased genes defined as log$_2$FC>0.58.

Supplemental Figure 3. a) Principal component analysis (PCA) of *Vim$^{-/-}$* MoAMs replicates. **b)** Gene Set Enrichment Analysis (GSEA) enrichment plots. GSEA was performed on gene lists ranked by edgeR reported fold-change between *Vim$^{-/-}$* and WT MoAMs at each time point. Normalized enrichment scores (NES) are shown. Asterisks indicate statistically significant adjusted p-values <0.05. **c)** Scatter plot of z-score normalized gene expression across genes in M$_{(IL-4)}$ vs Cell Cycle. Pearson correlation coefficients (r) across WT or *Vim$^{-/-}$* MoAM replicates, and associated p-values are shown. **d)** Heatmap matrix of Pearson correlation between processes, asterisk indicates p-value <0.05. **e)** Violin plots representing the log$_2$FC between *Vim$^{-/-}$* and WT in tissue-resident alveolar macrophages (TRAMs) at steady-state (0 dpi) for each gene by biological cluster and biological process. Solid lines represent median, dotted lines represent quartiles **f)** Heatmap showing pairwise Pearson correlation of mean gene expression across replicates between each time point in WT and *Vim$^{-/-}$* MoAMs. Pearson correlation coefficient values are shown within each square. Scale bar represents Pearson correlation coefficient. **g)** Heatmap of normalized expression of 2,623 genes as defined in Figure 2a. *Vim$^{-/-}$* data was normalized to the WT mean and standard deviation (adapted z-score). **h)** Violin plots representing the log$_2$FC between *Vim$^{-/-}$* and WT for each gene by cluster in MoAMs. Solid lines represent median, dotted lines represent quartiles. Asterisks indicate adjusted p-value <0.05

Figure 4. Protective phenotype of *Vim$^{-/-}$*mice is associated with the failure of *Vim$^{-/-}$*MoAMs to activate core pro-inflammatory genes in the early response to IAV. a) Percent (%) M$_{(LPS)}$ genes peaking in expression on each day post-infection in WT and *Vim$^{-/-}$* MoAMs **b)** Bar graphs of gene expression (FPKM) for representative M$_{(LPS)}$ genes present in the leading edge of at least 1 day from GSEA. Error bars represent SD. **c)** Heatmap of normalized expression of 84 M$_{(LPS)}$ genes present in the GSEA leading edge on at least 1 day (indicated by a color-coded bar on the left). Adapted Z-score normalization using WT mean and standard deviation for each row were applied to gene expression (FPKM) data. **d)** Histogram representing the distribution of log$_2$FC between *Vim$^{-/-}$* and WT MoAMs on 3 dpi of the Early Inflammatory phase (440 genes). The dashed line represents the mean of the distribution. The number of genes less than (186) and greater than (254) the mean are shown. Genes with log$_2$FC between *Vim$^{-/-}$* and WT MoAMs less than the mean represent those genes that are most changed between *Vim$^{-/-}$* and WT MoAMs. **e)** Mean adapted z-scores were calculated across genes for most (186) and least (254) changed genes on 3 dpi in the Early

Inflammatory Phase in WT and *Vim*$^{-/-}$ MoAMs. **f)** HOMER known motif analysis using genomic
background of most changed genes on 3 dpi in the Early Inflammatory Phase (186 genes). Motif
logos are shown below motif names. -log$_{10}$pvalues of enrichment in most or least changed gene set
is shown. Log$_2$(FPKM+1) average gene expression of transcription factor family members are
shown for WT TRAMs and WT IMs at steady-state (0 dpi) and 2-6 dpi for WT MoAMs. log$_2$FC
between *Vim*$^{-/-}$ and WT MoAM gene expression of transcription factor family members on each dpi.
(n=3-4)
925
Supplemental Figure 4. a) Mean adapted z-scores for M$_{(LPS)}$ and M$_{(IL-4)}$ genes present in the
leading edge of at least 1 day from GSEA in WT and *Vim*$^{-/-}$ MoAMs. **b)** Heatmap of normalized
expression of M$_{(IL-4)}$ genes present in the leading edge of at least 1 day from GSEA. Adapted Z-
score normalization for each row was applied to gene expression (FPKM) data. Bar graphs of gene
expression (FPKM) for representative M$_{(IL-4)}$ genes present in the leading edge of at least 1 day
from GSEA. Error bars represent SD. **c)** Heatmap of normalized expression of 186 most changed,
and 254 least changed genes on 3dpi of the Early Inflammatory Phase. Adapted Z-score
normalization for each row was applied to gene expression (FPKM) data. **d)** HOMER known motif
analysis using genomic background of least changed genes on 3 dpi in the Early Inflammatory
Phase (254 genes). Motif logos are shown below motif names. Log$_2$(FPKM+1) average gene
expression of transcription factor family members is shown for WT TRAMs and WT IMs at steady-
state (0 dpi) and 2-6 dpi for WT MoAMs. log$_2$FC between *Vim*$^{-/-}$ and WT MoAM gene expression
of transcription factor family members on each dpi. (n=3-4). **e)** Motif logos are shown below motif
names. -log$_{10}$pvalues of enrichment in most or least changed gene set is shown.
940
Figure 5. Vimentin-deficiency in the MoAM compartment is sufficient to protect from
influenza A virus-induced lung damage and mortality. a) Schematic of experimental design.
BMDM were isolated from either WT or Vim-/- mice. BMDM were treated with LPS or IL-4 to polarize
macrophages towards an M$_{(LPS)}$ or M$_{(IL-4)}$ state, respectively. Gene expression was assessed by
qPCR **b)** Expression of genes characterizing a pro-inflammatory and reparative macrophage state
in WT and *Vim*$^{-/-}$ macrophages. n = 2, in duplicate. Bar graphs represent mean relative fold change,
error bars represent SEM. Differences between genotypes were assessed by t-test. **c)** Schematic
of the experimental design. Recipient mice (CD45.1) were irradiated and reconstituted with bone
marrow cells from WT or *Vim*$^{-/-}$ donor mice (CD45.2). After a 30-day recovery period, chimeric mice
were intratracheally (i.t.) infected with a lethal dose of IAV at 5 days p.i. and subjected to
analyses. **d)** Bronchoalveolar lavage fluid (BALF) protein levels measured by Bradford assay, n=3-
4 **e)** Lung wet-to-dry ratio measured at day 5 p.i. n=7-10. **f)** Viral titer assessed by plaque assay.
n=11-14. **g)** Representative images of murine lung architecture at day 5 p.i. assessed by H&E
staining, scale bar 100 um. **h)** Levels of inflammatory cytokines in BALF assessed using a custom
multiplex ELISA. n=4-9. Bar graphs represent mean, error bars represent SEM. Differences
between genotypes were assessed by t-test. * p<0.05.
957
958   **Supplemental Figure 5.** Representative flow cytometry dot plots demonstrate successful
959   engraftment of WT and *Vim*$^{-/-}$ CD45.2 donor cells into CD45.1 recipient mice.
960

Figure 1

Supplemental Figure 1

a Cluster

b
Matrix Protein 2 (M2) H1N1

Nucleoprotein (NP) H1N1

Neuraminidase (NA) H1N1

☐ WT
▨ Vim-/-

c

d Neutrophils

☐ WT
▨ Vim⁻/⁻

Figure 2

Supplemental Figure 2

Figure 3

Supplemental Figure 3

Figure 4

Supplemental Figure 4

Figure 5

Supplemental Figure 5