The epidemic consequences of virulence and free-living survival relationships in models of emerging outbreaks

Lourdes M. Gomez[1][*], Victor A, Meszaros[1][*], Wendy C. Turner[2], and C. Brandon Ogbunugafor[1][#]

[1]Department of Ecology and Evolutionary Biology, Brown University, Providence RI 02912

[2]Department of Biological Sciences, University at Albany – State University of New York, Albany, New York 12222, USA

[#]Corresponding Author: brandon_ogbunu@brown.edu

[*]These authors contributed equally to this manuscript

ABSTRACT

The relationship between parasite virulence and transmission is a pillar of evolutionary theory that has specific implications for public health. Part of this canon involves the idea that virulence and free-living survival (a key component of transmission) may have different relationships in different host-parasite systems. Most examinations of the evolution of virulence-transmission relationships—theoretical or empirical in nature—tend to focus on the evolution of virulence, with transmission a secondary consideration. In virus evolution, recent studies have examined how the evolution of free-living survival can influence other traits. However, few studies have examined the epidemic-scale consequences of evolution across the different virulence-survival relationships, and specifically, of increased free-living survival. In this study, we utilize a mathematical model motivated by aspects of SARS-CoV-2 natural history to investigate how evolutionary changes in survival traits, and their subsequent relationship to virulence traits, may influence several aspects of disease dynamics. Specifically, we examine how each of two different hypothesized relationships between virulence and survival, one deemed "*the Curse of the Pharaoh hypothesis"* and the other more classically referred to as the "tradeoff hypothesis." Across virulence-survival frameworks, we found that small changes (5% above and below the nominal value) in survival can have a meaningful effect on certain outbreak features, including the R_0, and the size of the infectious peak in the population. These results highlight the importance of properly understanding the relationship between virulence and parasite survival, as they offer considerably different predictions for how the evolution of free-living survival will influence the trajectory of an emerging epidemic.

INTRODUCTION

The interactions between the life history of a pathogen and the environment in which it is embedded drive the evolution of virulence. These interactions thus dictate both the experience of disease at the individual host level, and the shape of disease dynamics in host populations [1,2]. The nature of the interaction between virulence and transmission has been the object of both theoretical and empirical examination [2-8]. Free-living survival, here defined as the ability of a pathogen to persist outside of its host, is one of many transmission life history traits associated with virulence. The relationship between the two varies between host-pathogen types and different environments [4,8-10].

Two hypotheses serve as canon in the evolution of virulence, theorizing on its relationship to transmission traits. The *Curse of the Pharaoh* hypothesis--named after a tale about a mythical curse that torments individuals who dig up tombs of Egyptian pharaohs [11] suggests that if a parasite has high free-living survival, then it is far less dependent on its host for transmission, and consequently, will have no evolutionary incentive to decrease virulence [2,4,12]. Spreading through a host population with high mortality can be supported by persisting in the environment until the arrival of new hosts or the recovery of the host population from survivors. That is, any presumptive selection on beneficence would be relaxed: parasites can detrimentally affect the health of hosts at no cost to transmission, because most of their life cycle is spent outside of a host. Previous studies support a positive correlation between free-living survival and mortality per infection (a common proxy for virulence) [13].

Alternatively, the "tradeoff" hypothesis suggests that virulence and transmission are inversely correlated, and there is some intermediate level of parasite virulence [3,6,14] that is optimal for a given setting. In this scenario, too high of virulence kills host and parasite, and too low of virulence leads to a failure to transmit. Applying this hypothesis specifically to free-living survival would suggest that selection for increased free-living survival should come at the expense of virulence (producing a pathogen that is less harmful to the host). Mechanistically, as a consequence of increased adaptation to a non-host environment, a virus may be less fit to replicate inside a host [9,15]. For example, a more robust viral capsid may help to survive harsh environmental conditions, but may make it more difficult to package RNA/DNA [15]. More generally, the tradeoff hypothesis can be framed in the context of a life history tradeoff: investment in certain parts of the life cycle often comes at the expense of others [2,16]. For

balance with the colorful name of the other hypothesis, here we call the tradeoff hypothesis the

"*Goldilocks hypothesis*" for the remainder of this study.

That support for each hypothesis exists suggests that different virulence-survival relationships

apply across varying host-pathogen and environmental contexts. Understanding which

hypothesis applies in specific settings provides a rough guideline for how the evolution of free-

living survival may influence the evolution of virulence in real world epidemics. By extension,

this may allow one to predict how the relationship between evolution of survival and evolution of

virulence will influence the intensity of outbreaks at the population level.

In this study, we compare the *Curse of the Pharaoh* (CP) and *Goldilocks hypothesis* (tradeoff)

(GH) with respect to how the evolution of free-living survival would influence the dynamics of an

epidemic resembling COVID-19. In order to measure how pathogen survival influences disease

dynamics, we included an "environmental compartment" in our model, which represents

"contaminated" environments that act as a reservoir for persisting pathogens, causing disease

spread when they come in contact with susceptible individuals (infection via "fomites") [17,18].

We also include an asymptomatic compartment—representing individuals who are infectious but

not displaying symptoms. In the COVID-19 pandemic, asymptomatic transmission is a

contentious topic, but several studies have suggested that it plays a role in the spread of

disease [19-21]. Though environmental transmission of SARS-CoV-2 has not been studied

rigorously, it is plausible that asymptomatic individuals may spread disease through frequent

contact with the environment, thus increasing the proportion of virus that is free-living [22].

We find that the identity of the virulence-free-living survival relationship has distinct implications

for how an epidemic will unfold. Some, but not all, features of an outbreak are dramatically

influenced by the nature of the underlying virulence-survival relationship. This indicates that

signatures for adaptation in a pathogen population will manifest more conspicuously in certain

characteristics of an outbreak. We reflect on these findings in light of their theoretical

implications on the evolution and ecology of infectious disease, and for their potential utility in

the public health interventions.

METHODS

Model motivation and application

The mathematical model explored in this study is adapted from a recent one developed to

investigate environmental transmission of SARS-CoV-2 during the early-stage outbreak

dynamics of COVID-19, with parameter values based on fits to actual country outbreak data

[23]. In this study, we utilize this model to examine questions about the evolution of free-living

survival. While the phenomenon we examine is a very relevant one that manifests in the actual,

biological world, we want to emphasize that none of the methods or results in this study are

intended to apply to the active COVID-19 pandemic. This study is an attempt at responsible

theoretical biology, with data-informed models and inferences that are germane to the natural

world. However, we do not support the extrapolation of these findings to any particular aspect of

COVID-19, nor should they inform a policy or intervention.

Model Description

The model is implemented via a set of ordinary differential equations, defined by Equations 1.1

– 1.6. It implements viral free-living survival via the "Waterborne Abiotic or other Indirect

Transmission (WAIT)" modelling framework, coupling individuals and the pathogen within the

environment [24,25].

Within the model a β_w term allows for individuals to become infected via viral pathogen

deposited in the environment, and terms $\square_A$ and $\square_I$ allow asymptomatic and symptomatic

individuals to deposit pathogen into the environment, respectively. Adapted from the more

traditional SEIR (susceptible-exposed-infected-recovered), the SEAIR-W (susceptible-exposed-

asymptomatic-infected-recovered-WAIT) model interrogates the consequences of the two

hypotheses outlined above, while representing the dynamics of a very relevant disease system

(SARS-CoV-2) that includes an asymptomatic infectious population.

$$\frac{dS}{dt} = \mu(N-S) - (\frac{\beta_A A + \beta_I I}{N} + \beta_W W)S \qquad (1.1)$$

$$\frac{dE}{dt} = (\frac{\beta_A A + \beta_I I}{N} + \beta_W W)S - (\epsilon + \mu)E \qquad (1.2)$$

$$\frac{dA}{dt} = \epsilon E - (\omega + \mu)A \qquad (1.3)$$

$$\frac{dI}{dt} = (1-p)\omega A - (v + \mu_I)I \qquad (1.4)$$

$$\frac{dR}{dt} = p\omega A + v I - \mu R \qquad (1.5)$$

$$\frac{dW}{dt} = (\frac{\sigma_A A + \sigma_I I}{N})(1-W) - kW \qquad (1.6)$$

Figure 1 depicts the compartmental diagram for the model. The direction of the arrows outline

the flow of the individuals and the pathogen through the system. Note that individuals can move

from the asymptomatically infected compartment to the recovered compartment, what we call a

"mild track". The dashed arrows represent the WAIT coupling to the environment. The model is

inspired by a model developed to interrogate environmental transmission of SARS-CoV-2 [23].

Simulations of outbreaks

The system was numerically integrated using the "odeint" solver in the Scipy 1.4 - the Python

scientific computation suite [26]. The simulations track the populations for each of the

compartments listed in Figure 1. Each model run occurred over 250 days, which amounts to

over 8 months of epidemic or 5x the peak of the outbreak. This length of time is consistent with

the antecedent SARS-CoV-2 model (23), long enough for the dynamics of the system to

manifest. Note, however, that for this study, we are especially interested in the early window of

an outbreak, the first 30 days. We focus on this window because this is the area of an outbreak

that best captures real-world dynamics, as 30 days is often before populations have been able

to adjust their individual behaviors. In that way, the first month is a measure of the true physics

of an epidemic. The code constructed for the analysis in this study is publicly available on

github: https://github.com/OgPlexus/Pharaohlocks

Figure 1. Compartmental diagram of the SEAIR-W model: Here we present the compartmental diagram for the model system highlighted in this study. The model is similar to a mathematical model used to interrogate environmental transmission of SARS-CoV-2 (see [23]).

Population definitions & parameter values

Table 1 outlines the definitions of each population, and provides the initial population values used for all simulations conducted in this study. The nominal parameter values used are defined in Table 2. The initial values are drawn from the aforementioned COVID-19 outbreak study, derived from empirical findings and country-level outbreak data [23].

Symbols	Value	Units	Definitions
S_0	57.05×10^6	people	Susceptible individuals
E_0	66.50	people	Exposed individuals
A_0	13.30	people	Asymptomatic individuals
I_0	13.30	people	Symptomatic individuals
Rec_0	0	people	Recovered individuals
W_0	1%	unitless	% of viral pathogen in environment

Table 1. Definitions and initial values for the populations represented by each compartment. The values here are the averages of the model values across all the countries developed in a prior COVID-19 study [23].

Symbols	Values	Units	Definitions
μ	1/(80.3 * 365)	1/day	Natural death rate (Reciprocal of the upper bound of average human lifespan)
μ_I	0.00159	1/day	Infected death rate (Natural death rate + disease induced death rate)
$\eta = (?{-}\varepsilon^{-1})$	5.5	days	SARS-CoV-2 incubation period
$1/?$	$\eta - \varepsilon^{-1}$	days	Rate of transfer from asymptomatic to symptomatic
v	0.0305	1/day	Recovery rate (Average of 3 to 6 weeks)
p	95.6%	percent	Percent that move along the "mild" recovery track
k	0.649	1/day	Waning virus rate in environment (using average of all material values, wood, steal, cardboard, plastic)
β_a	0.550	1/day	(Contact rate of people with people) x (transmission probability of people to people by A-person)
β_I	0.491	1/day	(Contact rate of people with people) x (transmission probability of people to people by I-person)
β_w	0.031	1/day	(Contact rate of person with environment) x (transmission probability of environment to people)
σ_a	3.404	1/day	(Contact rate of person with environment) x (probability of shedding by A-people to environment)
σ_I	13.492	1/day	(Contact rate of person with environment) x (probability of shedding by I-people to environment)
$1/\varepsilon$	2.478	days	Average number of days before infectious

173 **Table 2.** Definitions for the nominal parameter values used in this study. Parameter values were
174 developed from empirical findings and country-level data, as discussed in another study [23].

Virulence definition

In this study, we define virulence as a set of parameters that uniformly increase the rate or probability of causing symptomatic disease, or the severity of those symptoms (including death). Our definition is more comprehensive than many other models of parasite virulence (e.g., [4,13]), which tend to focus on a single aspect of the natural history of disease associated with harm to a host (e.g., the fitness consequences of an infection on the host population, or the case fatality rate). Instead of having to justify a definition built around a single term (e.g. the term associated with fatality), we took a collective approach to defining virulence through all terms that foment the viral-induced onset of symptomatic disease, and death. This more flexible definition allows for the reality of pleiotropic effects in viral pathogens, where adaptations can have multiple effects on the natural history of disease [2,27].

This iteration of virulence negates the potential for overly-weighting only one, or a small set, of parameters under a large umbrella of virulence. The collection of parameters that we use to define virulence are as follows: the infected population death rate ($\square_I$), the incubation period of SARS-CoV-2 ($\square$), the rate of transfer from asymptomatic to symptomatic ($1/\square$), the infected population recovery rate (v), the percent of individuals that move from the asymptomatic to the recovered compartment without showing symptoms (the "mild" recovery track, p), the contact rate of people with people *X* transmission probability of people to people by an asymptomatic individual (β_A), the contact rate of people with people *X* transmission probability of people to people by asymptomatically infected person (β_I), the contact rate of people with environment *X* probability of shedding by asymptomatic individual to the environmental ($\square_A$), the contact rate of people with the environment *X* probability of symptomatically infected individuals shedding into the environment ($\square_I$), and the average number of days before infection ($1/\varepsilon$).

Table 3 outlines the direction in which each of the virulence-associated parameters are modulated as virulence decreases or increases. An up arrow (↑) indicates the parameter increases (by an equivalent percent) when the percent virulence is changed. A down arrow (↓) indicates the parameter decreases (by an equivalent percent), when the percent change in virulence is applied. Changes in virulence are then defined, in this study, as an equivalent uniform (percent) change in each of the parameters listed above. For the purposes of our study, we modify virulence by changing all parameters associated with virulence by 5%. It would also be easy to disambiguate virulence into changes in individual subcomponents, however, that is not the focus on the current study.

Symbols	Definition	Virulence Increased	Virulence Decreased
μ_I	Infected death rate (Natural death rate + disease induced death rate)	↑	↓
$\eta = (\square - \varepsilon^{-1})$	SARS-CoV-2 incubation period	↓	↑
$1/\square$	Rate of transfer from asymptomatic to symptomatic	↑	↓
v	Recovery rate (Average of 3 to 6 weeks)	↓	↑
p	Percent that move along the "mild" recovery track	↓	↑
β_A	(Contact rate of people with people) x (transmission probability of people to people by A-person)	↑	↓
β_I	(Contact rate of people with people) x (transmission probability of people to people by I-person)	↑	↓
σ_A	(Contact rate of person with environment) x (probability of shedding by A-people to environment)	↑	↓
σ_I	(Contact rate of person with environment) x (probability of shedding by I-people to environment)	↑	↓
$1/\varepsilon$	Average number of days before infectious	↓	↑

221 **Table 3. Virulence parameters.** This is a list of uniformly modulated parameters, and the
222 direction in which they change, when virulence is increased or decreased. When virulence
223 changes, an up arrow (↑) indicates the parameter is increased (by an equivalent percent), and a
224 down arrow (↓) indicates the parameter is decreased (by an equivalent percent).
225

Survival definition

Survival is defined as the set of parameters that when uniformly modulated, increase the pathogen's probability of surviving the outside environment and successfully infected a new host. In our model this includes both the waning virus rate in the environment (k) and the contact rate of an individual with environment **X** transmission probability of environment to people (β_w). Table 4 outlines the direction (increasing or decreasing) in which these parameters are modulated when survival is decreased or increased. Within both models a (percent) change in survival is defined as an equivalent uniform (percent) change in the survival parameters.

Symbols	Definition	Survival Increased	Survival Decreased
k	Waning virus rate in environment (using average of all material values, wood, steal, cardboard, plastic)	↓	↑
β_w	(Contact rate of person with environment) x (transmission probability of environment to people)	↑	↓

Table 4. Survival parameters. This is a list of parameters that are uniformly modulated, and the direction in which they change, when survival is increased or decreased. An up arrow (↑) indicates the parameter is increased by some percent, when the equivalent (percent) change in survival is applied. A down arrow (↓) indicates the parameter is decreased by some percent, when the equivalent (percent) change in survival is applied.

Throughout this study, the impact of changes in virulence and survival are assessed with respect to the following four epidemic metrics: the number of infected individuals (asymptomatic and symptomatic) at the maximum (when the outbreak is at its most severe), the rate at which the peak infected population is reached (t_{max}^{-1}), the total infected population after 30 days, and the basic reproductive ratio (R_0).

Basic reproductive ratio

259 **Basic reproductive ratio**

261
$$R_0 = \frac{R_p + \sqrt{R_p^2 + 4R_e^2}}{2} \qquad (2.1)$$

264
$$R_p = \frac{\epsilon(\beta_A(\mu_I + v) + \beta_I(1-p)\omega)}{(\mu+\epsilon)(\mu+\omega)(\mu_S+v)} \qquad (2.2)$$

267
$$R_e^2 = \frac{\epsilon\beta_W(\sigma_A(\mu_I+v) + \sigma_I(1-p)\omega)}{k(\mu+\epsilon)(\mu+\omega)(\mu_S+v)} \qquad (2.3)$$

Equations 2.1 - 2.3 give the analytic expression of the basic reproductive ratio (R_0) for the model

used in this study. This expression for R_0 can be deconstructed into two components. Equation

2.2 only contains parameters associated with person to person transmission (R_p), while

equation 2.3 solely contains parameters associated with transmission from the environment

(R_e).

Applying the parameters values in Table 2, the numerical value of the basic reproductive ratio is

given as: $R_0 \sim 2.82$.

RESULTS

Model sensitivity analysis.

Figure 2 depicts a tornado plot that communicates the sensitivity of the model to permutations in

parameters. The model is most sensitive to parameters that are considered virulence-

associated (Table 3), and is relatively less

sensitive to survival- associated parameters

(Table 4).

Figure 2. Tornado plot showing the sensitivity of R_0 to individual parameter changes.
Filled bars indicate the value of R_0 when the associated parameter is increased by 5.0% from its nominal value. White bars indicate the value of R_0 when the associated parameter is decreased by 5.0%. Blue coloring with checkered patterning indicates a parameter associated with survival, and orange coloring with lined patterning indicates a parameter associated with virulence.

The R_0 of the model is most sensitive to the parameters ▯, β_A, and v. The sensitivity of R_0 to

changes in ▯ reflects the importance of the rate of conversion into the symptomatic state on

model dynamics. In addition, the β_A has a very important influence on the model, consistent with

other findings for COVID-19 that have emphasized the importance of asymptomatic

transmission in disease spread [19-21].

Illustrative dynamics of model system

Applying the parameter values in Table 2, Figure 3a demonstrates the base dynamics of the

model playing out over the first 100 days while Figure 3b shows the dynamics within the

environment over the course of 250 days. In these dynamics, the population begins to be fixed

for susceptible hosts. The disease dynamics manifest in the shapes of the curves corresponding

to the exposed, asymptomatic, and symptomatic individuals. Note the long tail of the curve

corresponding to contamination by the environment. That is, the environment remains infectious

even after the infected populations have declined in number. The length and shape of this tail

that is influenced by the free-living survival of the virus.

A   **B**

327 **Figure 3. Sample dynamics for the model system.** (**A**) Gives the dynamics for all host
328 compartments within the model **(B)** corresponds to the fraction of environmental reservoirs in a
329 setting that are contaminated with infectious virus.
330

The epidemic consequences of varying virulence and survival.

In the next analysis, we examine the epidemic consequences of varying traits associated with

survival and virulence. In Figure 4, we observe how dynamics of the outbreak are influenced by

different combinations of traits altering virulence (see Table 3 for a list of virulence-associated

parameters) and survival (see Table 4 for a list of survival-associated parameters), changed by

+/- 5% (10% overall). In Figure 4D, we demonstrate how changes in virulence and free-living

survival traits influence R_0, with variation in virulence-related traits having the largest effect on

R_0. Of note is how the range in R_0 values varies widely across virulence-survival values, from

nearly 2.0 to 3.7 (Figure 4D).

 A

B

C   **D**

366
367
Figure 4. The impact of varying virus virulence and survival assessed on four key
epidemic metrics. Here we present heatmaps expressing the change in (A) the number of
infected individuals (asymptomatic and symptomatic) at the epidemic peak, (B) the rate at which
the epidemic peak is reached, t_{max}^{-1}, (C) the total infected population after 30 days, and (D) the
basic reproductive ratio (R_0) when virulence and survival are modulated by +/- 5% within the
model.
Curse of the Pharaoh **vs. the** *Goldilocks hypothesis***: Implications at their extremes**

Having observed how outbreak dynamics are influenced by variation in traits that alter virulence-

survival phenotypes, we then examined how each outbreak metric is influenced by extreme (+/-

5%) values of these trait combinations. We compared how the evolution of traits that enhance

survival in turn affect outbreak dynamics, according to the two hypotheses considered, and their

relationships between virulence and survival (Figure 5).

● The *Curse of the Pharaoh* hypothesis would manifest as a positive relationship between

virulence and survival. High values for survival would be associated with high values for

virulence [4,13]. Because the correlations we observe are often not exactly linear, we

utilize quadrants to express a trend, allowing for some variance around the expected

"line". In Figure 5, 5he *Curse of the Pharaoh* hypothesis can be represented by

combinations of virulence and survival residing in quadrants I and III.

● The *Goldilocks hypothesis* (tradeoff) would manifest as a negative relationship between

virulence and survival. In this scenario, high values for survival would be associated with

low values for virulence, and low peak total infected population [2,9,15]. Pathogens with

a life history that follow the tradeoff hypothesis exhibit virulence-survival associations

that would likely appear in quadrants II and IV in Figure 5.

If host-pathogen evolution proceeds according to the *Curse of the Pharaoh* hypothesis, all

outbreak metrics would show an increase in severity as both survival and virulence increase.

Across the range of variation in virulence and survival traits considered (5% above and below

the nominal value), peak number of infected individuals increases by ~35%, the rate at which

the peak is reached increases by ~16%, the total number of infected individuals after 30 days

increases by ~98%, and the R_0 increases by ~94% (Figure 6 and Table 5).

Under the *Goldilocks hypothesis*, outbreak severity generally decreases as survival increases.

Across the measured range of variation of virulence-survival traits, the peak number of infected

individuals decreases by ~23%, the rate at which the epidemic peak is reached decreases by

0.15%, the total number of infected individuals decreases by 3%, and the R_0 decreases by

~84% (Figure 6 and Table 6). Across all metrics considered, the effects of increased viral

survival on outbreak dynamics is more extreme under the *Curse of the Pharaoh* than the

Goldilocks hypothesis (Figure 6).

In Figure 7, we observe the disease dynamics at survival extremes, and dynamics

corresponding to the fraction of the environment that is contaminated with infectious virus.

Consistent with the data represented in Figure 6, we observe that minimum and maximum

simulations differ more substantially for extreme survival scenarios in the *Curse of the Pharaoh*

scenario than for the *Goldilocks hypothesis*. The simulation dynamics should be analyzed with

respect to the dynamics of hosts, and of the environmental reservoirs. The feature of the

different outbreaks that differs most ostensibly between the parameters is the time needed to

reach the peak number of infected individuals. In the *Curse of the Pharaoh* simulations, one can

observe the low virulence, low survival scenario (Figure 5A and 5B) take longer to reach the

peak number of infected individuals. Most notably, however, the low virulence, low survival

setting has far smaller peak of environmental contamination, and shorter tail relative to its high

virulence, high survival counterpart (Figure 5D). Similarly, intriguing findings exist in the

comparison between the simulation sets corresponding to extremes in the *Goldilocks*

hypothesis setting (Figure 5E-H). Especially notable is the difference in the length of the tail of

the environment contamination for the high virulence, low survival combination (Figure 5F) vs.

the low virulence, low survival combination variant (Figure 5H). The explanation is that, in this

model, high virulence influences (among many other things) the rate at which virus is shed into

the environment from either the asymptomatic (σ_A) or symptomatic (σ_I) host. We observe how

the high virulence, low survival simulation (Figure 5E) features a symptomatic peak that is larger

in size, and prolonged relative to the low virulence counterpart (Figure 5G). This relatively large

symptomatic population is shedding infectious virus into the environment for a longer period of

time, contributing to the long tail of contaminated environments observed in Figure 5F.

437
438

Figure 5. The expected effect of increasing survival on virulence for the two hypotheses considered. Here we present a schematic of how the different hypotheses for the relationship between virulence and survival manifest on a map with a structure similar to the heat maps shown in Figure 4. The directions of the arrows depict how increasing survival would affect virulence under the two hypotheses: the blue arrow indicates the flow of an increasing "*Curse of the Pharaoh*" dynamic, while the direction of the orange arrow indicates an increasing "*Goldilocks hypothesis*" dynamic.

455 *Curse of the Pharaoh*

Epidemic Metric	min virulence, min survival	max virulence, max survival	% difference between min survival and max survival
Peak total infected (people)	5.68×10^{06}	7.64×10^{06}	+34.51%
t_{max}^{-1} (days^{-1})	1.99×10^{-02}	2.23×10^{-02}	+12.06%
Total after 30 days (people)	8.18×10^{07}	1.62×10^{08}	+98.04%
Basic reproductive ratio (R_0)	1.95	3.78	+93.84%

456 **Table 5. Comparing epidemic metrics under low survival/low virulence versus high
457 survival/high virulence scenarios (as in the *Curse of the Pharaoh* hypothesis).** For each
458 metric analysed, these are the heatmap values for the bottom left [at "coordinates" (Vir , Sur)
459 → (-5% , -5%)] and top right [at "coordinates" (Vir , Sur) → (+5% , +5%)] corners.
475 *Goldilocks hypothesis*

Epidemic Metric	max virulence, min survival	min virulence , max survival	% difference between min survival and max survival
Peak total infected (people)	7.39×10^{06}	5.99×10^{06}	-23.47%
t_{max}^{-1} (days^{-1})	2.10×10^{-02}	2.23×10^{-02}	-0.15%
Total infected after 30 days (people)	1.16×10^{08}	1.13×10^{08}	-2.68%
Basic reproductive ratio (R_0)	3.67	1.99	-84.39%

476 **Table 6. Comparing epidemic metrics under low survival/low virulence versus high**
477 **survival/high virulence scenarios (as in the *Goldilocks hypothesis*).** For each metric
478 analysed, these are the global heatmap values for the top left [at "coordinates" (Vir , Sur) → (
479 +5% , -5%)] and bottom right corners [at "coordinates" (Vir , Sur) → (-5% , +5%)]

491
492

Figure 6. The percent change in SEAIR-W outbreak metrics as survival increases from -5% to +5% comparing virulence-survival hypotheses. For each metric analysed we present the percent difference between the minimum and maximum survival values given the two hypotheses tested, *Curse of the Pharaoh* (comparing low virulence/low survival to high virulence/high survival) and the *Goldilocks hypothesis* (comparing high virulence/low survival to high survival/low virulence). The bars here correspond to the values (percents) in the third columns of Table 5 and Table 6, which denote the differences between the minimum and maximum values.

521   **A**   **B**

524   **C**   **D**

Figure 7. Virus outbreak dynamics for the extreme values of virulence and free-living survival considered for the two hypotheses. These plots are similar to the illustrative dynamics in Figure 1. Here we observe the dynamics of disease corresponding to the extreme values presented in Tables 5 and 6. Figures 7A, C, E, and G depict disease dynamics, and 7B, D, F and H the dynamics of contaminated environments. Figures 7A - D correspond to the parameter values considered for the *Curse of the Pharaoh*, while Figures 7E - H correspond to the *Goldilocks hypothesis.*

DISCUSSION

The virulence-survival relationship drives the consequences of virus evolution during an outbreak. In this study, we examined how different virulence-survival relationships may dictate different features of outbreaks as pathogens evolve (according to the *Curse of the Pharaoh* or the *Goldilocks hypothesis*). When the parameter space for virulence and survival is mapped, we find that certain outbreak metrics are more sensitive to change in free-living survival traits than others, and that the nature of this sensitivity differs depending on whether evolution of heightened survival proceeds according to *Curse of the Pharaoh* (CP) or *Goldilocks hypothesis* (GH).

In the *Curse of the Pharaoh* setting, as free-living survival varies between 5% below and above the nominal value, we observed a dramatic change in the total number of infected individuals in the first 30 days (98% increase from minimum survival to maximum survival), and the R_0 nearly doubles (94% increase). These two traits are, of course, connected: the theoretical construction of the R_0 metric specifically applies to settings where a pathogen is spreading in a population of susceptible hosts [28,29]. Strikingly, evolution of survival in the *Goldilocks hypothesis* setting has different outbreak implications: while the R_0 difference between minimum and maximum survival is significant (~84% decrease), the total number of infected individuals only changes by roughly 3%. The difference between the *Curse of the Pharaoh* and the *Goldilocks hypothesis* resides entirely in how virulence maps onto survival in the two models: at extreme survival values in the *Goldilocks hypothesis*, the hypothetical virus has a reduction across virulence parameters. This has a dramatic impact on the R_0. Notably, this large difference between the R_0 at high and low survival values does not translate to a difference in the total number of infected individuals in the first 30 days of an infection (the early outbreak window). That is, in a *Goldilocks hypothesis* scenario, a highly virulent and less virulent virus population can have similar signatures on a population with respect to the number of infected individuals in the first month. That is, simply measuring the number of infected individuals in the first month of an outbreak is not sufficient to reveal how a pathogen is evolving.

Across frameworks, we observe that relatively small changes in virulence and survival can have large impacts on outbreak dynamics. That the entirety of the results in this study took place within a relatively small range of parameter values (+/- 5%) is significant. This suggests that it

doesn't take large phenotypic changes in free-living survival in order to have a notable imprint
on outbreak dynamics.

Practical implications for the understanding of outbreaks caused by emerging viruses.
In summary, our results suggest that carefully constructed, mechanistically sound models of
epidemics are important, both for capturing the dynamics of an outbreak, and for understanding
how evolution of survival and virulence influence disease dynamics. For example, our results
highlight that questions surrounding the evolution of viruses like SARS-CoV-2 during an
epidemic require very specific types of evidence. Detecting the consequences of virus evolution
would depend on which feature of an outbreak an epidemiologist is measuring: from our
analysis, the R_0 is most significantly impacted by changes in virulence and survival. In addition,
the total number of infected individuals in the early window, and the size of the infected "peak"
would each be impacted most readily by changes in virulence-survival traits. The rate at which
the epidemic peak was reached, on the other hand, showed relatively little change as survival
increased, or between the two hypotheses.

In the context of COVID-19, speculation has surfaced about the potential for local adaptation in
certain virus populations that is associated with an attenuation in disease severity (or decreased
virulence, in our parlance). As of June 1, 2020, evidence for these types of adaptations is
tenuous, and any grand conclusions or predictions are premature. We propose that the process
of predicting the evolutionary consequences of viral adaptation should encompass several
discrete steps. Firstly, we should determine whether molecular signatures exist for adaptive
evolution. Adaptive evolution would manifest in observable differences in genotype and
phenotype, perhaps manifesting in the natural history of disease. Secondly, we should aim to
attain knowledge of the underlying mechanistic relationship between survival and virulence. This
knowledge is not necessarily easy to attain (it requires extensive laboratory studies) but would
allow us predictive power: we may be able to extrapolate how changes in some traits influence
others. Finally, with this information, mechanistically-informed models can be constructed that
may give useful insight into the future trajectory of an outbreak.

While the stochastic nature of epidemics renders them challenging to predict [30], we suggest
that canons such as life history theory and the evolution of virulence provide useful lenses that
can aid in making modest predictions. More generally, we propose that in an age of great
accumulation of genomic and phenotypic data in many pathogen-host systems, that we
continue to responsibly apply or modify existing theory, in order to collate said data into an

organized picture—and potentially paradigm—for how different key components of pathogen
biology are related to one another, and influence the shape of viral outbreaks of various kinds.
REFERENCES

1. Ewald, P. W. Host-parasite relations, vectors, and the evolution of disease severity.
Annu. Rev. Ecol. Syst. **14**, 465–485 (1983).
618

619   2. Goldhill, D. H. & Turner, P. E. The evolution of life history trade-offs in viruses. *Curr.*
620   *Opin. Virol.* **8**, 79–84 (2014).
621

622   3. Anderson, R. M. & May, R. M. Trends and Perspectives: Coevolution of hosts and
623   parasites. *Parasitology* **85**, 411–426 (1982).
624

625   4. Bonhoeffer, S., Lenski, R. E. & Ebert, D. The curse of the pharaoh: The evolution of
626   virulence in pathogens with long living propagules. *Proc. R. Soc. B Biol. Sci.* **263**, 715–
627   721 (1996).
628

629   5. Gandon, S. The Curse of the Pharaoh Hypothesis. *Proc. R. Soc. B Biol. Sci.* **265**, 1545–
630   1552 (1998).
631

632   6. Alizon, S., Hurford, A., Mideo, N. & Van Baalen, M. Virulence evolution and the trade-off
633   hypothesis: History, current state of affairs and the future. *J. Evol. Biol.* **22**, 245–259
634   (2009).
635

636   7. Cressler, C. E., McLeod, D. V., Rozins, C., Van Den Hoogen, J. & Day, T. The adaptive
637   evolution of virulence: A review of theoretical predictions and empirical tests.
638   *Parasitology* **143**, 915–930 (2016).
639

640   8. Rafaluk-Mohr, C. The relationship between parasite virulence and environmental
641   persistence: A meta-analysis. *Parasitology* **146**, 897–902 (2019).
642

643   9. Ogbunugafor, C.B. *et al.* Evolution of increased survival in RNA viruses specialized on
644   cancer-derived cells. *Am. Nat.* **181**, 585–595 (2013).
645

646   10. Rafaluk, C., Jansen, G., Schulenburg, H. & Joop, G. When experimental selection for
647   virulence leads to loss of virulence. *Trends Parasitol.* **31**, 426–434 (2015).
648
649   11. Corelli, M. The curse of Osiris. London Daily News, Apri 5, p. 1 (1923).
650
651   12. Roche, B., Drake, J. M. & Rohani, P. The curse of the Pharaoh revisited: Evolutionary bi-
652   stability in environmentally transmitted pathogens. *Ecol. Lett.* **14**, 569–575 (2011).
653
654   13. Walther, B. A. & Ewald, P. W. Pathogen survival in the external environment and the
655   evolution of virulence. *Biol. Rev. Camb. Philos. Soc.* **79**, 849–869 (2004).
656
657   14. Frank, S. A. Models of parasite virulence. *Q. Rev. Biol.* **71**, 37–78 (1996).
658
659   15. De Paepe, M. & Taddei, F. Viruses' life history: Towards a mechanistic basis of a trade-
660   off between survival and reproduction among phages. *PLoS Biol.* **4**, 1248–1256 (2006).
661
662   16. Stearns, S. C. Trade-Offs in Life-History Evolution. *Funct. Ecol.* **3**, 259–268 (1989).
663
664   17. Barker, J., Stevens, D. & Bloomfield, S. F. Spread and prevention of some common viral
665   infections in community facilities and domestic homes. *J. Appl. Microbiol.* **91**, 7–21
666   (2001).
667
668   18. Boone, S. A. & Gerba, C. P. Significance of fomites in the spread of respiratory and
669   enteric viral disease. *Appl. Environ. Microbiol.* **73**, 1687–1696 (2007).
670
671   19. Bai, Y. *et al.* Presumed Asymptomatic Carrier Transmission of COVID-19. *JAMA* **323**,
672   1406–1407 (2020).
673
674   20. Park, S. W., Cornforth, D. M., Dushoff, J. & Weitz, J. S. The time scale of asymptomatic
675   transmission affects estimates of epidemic potential in the COVID-19 outbreak. *medRxiv*
676   2020.03.09.20033514 (2020) doi:10.1101/2020.03.09.20033514.
677
678   21. Tao, Y. *et al.* High incidence of asymptomatic SARS-CoV-2 infection, Chongqing, China.
679   *medRxiv* (2020).

22. Santarpia, A. J. L. *et al.* Transmission Potential of SARS-CoV-2 in Viral Shedding
    Observed at the University of Nebraska Medical Center. 1–12 (2020).

23. Ogbunugafor, C. Brandon, et al. "The intensity of COVID-19 outbreaks is modulated by
    SARS-CoV-2 free-living survival and environmental transmission." *medRxiv* (2020).

24. Miller-Dickson, M. D., Meszaros, V. A., Almagro-Moreno, S. & Brandon Ogbunugafor, C.
    Hepatitis C virus modelled as an indirectly transmitted infection highlights the centrality
    of injection drug equipment in disease dynamics. J. R. Soc. Interface 16, (2019).

25. Meszaros, V. A., Miller-Dickson, M. D., Junior, F. B. A., Almagro-Moreno, S. & Brandon
    Ogbunugafor, C. Direct transmission via households informs models of disease and
    intervention dynamics in cholera. *PLoS One* **15**, 1–14 (2020).

26. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python.
    Nat. Methods **17**, 261–272 (2020).

27. Duffy, S., Turner, P. E., & Burch, C. L. Pleiotropic costs of niche expansion in the RNA
    bacteriophage Φ6. *Genetics* **172.2** 751-757 (2006).

28. Jones, J. H. Notes on R0. *California: Department of Anthropological Sciences* **323**, 1–19
    (2007).

29. Diekmann, O., Heesterbeek, J. A. P., & Roberts, M. G. The construction of next-
    generation matrices for compartmental epidemic models. J*. R. Soc. Interface.* **7**, 873–
    885 (2010).

30. Scarpino, S.V., Petri, G. On the predictability of infectious disease outbreaks. *Nature*
    Communications **10**, 898 (2019).