Endochondral bone in an Early Devonian 'placoderm' from Mongolia

Martin D. Brazeau[1,2], Sam Giles [2,3,4], Richard P. Dearden[1,5], Anna Jerve[1,6], Y.A.

Ariunchimeg[7], E. Zorig[8], Robert Sansom[9], Thomas Guillerme[10], Marco Castiello[1]

[1] *Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst*

Rd, Ascot, SL5 7PY, UK;

[2] *Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7*

5BD, UK;

[3] *School of Geography, Earth and Environmental Sciences, University of Birmingham,*

Birmingham, UK;

[4] *Department of Earth Sciences, University of Oxford, South Parks Road, Oxford, OX1 3AN,*

UK;

[5] *CR2P Centre de Recherche en Paléontologie – Paris, Muséum national d'Histoire*

naturelle, Sorbonne Universités, CNRS, CP 38, 57 Rue Cuvier, 75231, Paris, Cedex 05,

France

[6] *Department of Organismal Biology, Subdepartment of Evolution and Development,*

Uppsala University, Norbyvägen 18A, 752 36 Uppsala, Sweden;

[7] *Natural History Museum, P.O. Box 46/52, Ulaanbaatar 1420, Mongolia*

[8] *Institute of Paleontology, Mongolian Academy of Science, P.O. Box 46/650, S. Danzan*

Street 3/1, Chingeltei District. Ulaanbaatar 15160, Mongolia;

[9] *School of Earth and Environmental Sciences, University of Manchester, Manchester M13*

9PT, UK;

[10] *Department of Animal and Plant Sciences, The University of Sheffield, Sheffield S10 2TN,*

UK;

Endochondral bone is the main internal skeletal tissue of nearly all osteichthyans[1,2] —

the group comprising more than 60,000 living species of bony fishes and tetrapods.

Chondrichthyans (sharks and their kin) are the living sister group of osteichthyans and

have cartilaginous endoskeletons, long considered the ancestral condition for all jawed

vertebrates (gnathostomes)[3,4]. The absence of bone in modern jawless fishes and the

absence of endochondral ossification in early fossil gnathostomes appears to lend

support to this conclusion. Here we report the discovery of extensive endochondral bone

in a new genus of 'placoderm'-like fish from the Early Devonian (Pragian) of western

Mongolia described using x-ray computed microtomography (XR-µCT). The fossil

consists of a partial skull roof and braincase with anatomical details providing strong

evidence of placement in the gnathostome stem group. However, its endochondral space

is filled with an extensive network of fine trabeculae resembling the endochondral bone

of osteichthyans. Phylogenetic analyses place this new taxon as a proximate sister group

of the gnathostome crown. These results provide direct support for theories of

generalised bone loss in chondrichthyans[5,6]. Furthermore, they revive theories of a

phylogenetically deeper origin of endochondral bone and its absence in

chondrichthyans as a secondary condition[7,8].

Systematic palaeontology

45   Placodermi M'Coy 1848

46   *Minjinia turgenensis* gen. et sp. nov.

Etymology. Generic name honours the memory of Chuluun Minjin for his extensive

contributions to the Palaeozoic stratigraphy of Mongolia, his enthusiastic support of this

work, and introducing us to the Yamaat River locality. Specific name recognises the

provenance of the fossil from the Turgen region, Uvs aimag of western Mongolia.

Holotype. Institute of Paleontology, Mongolian Academy of Sciences MPC-FH100/9.1, a

partial braincase and skull roof.

Type locality. Turgen Strictly Protected Area, Uvs province, western Mongolia; near the top

of the stratigraphic sequence that occurs between the Tsagaan-Salaat and Yamaat Rivers.

Formation and age. Upper part of Tsagaansalaat Formation, Pragian (Early Devonian)[9,10].

Diagnosis. 'placoderm'-grade stem gnathostome with endochondral bone, deep epaxial
muscle cavities flanking a slender occipital ridge, and the following possible autapomorphies:
dermal bones covered in sparsely placed tubercles, penultimate spino-occipital nerve canal
substantially larger in diameter than others.

Description

MPC-FH100/9.1 consists of a partial braincase and skull roof (Fig. 1). The skull roof is
ornamented with sparsely distributed stellate tubercles resembling those of the Siberian
'placoderm' *Dolganosteus*[11]. Towards the midline of the skull roof, the tubercles are larger
and more pointed, and are more broadly separated from each other by unornamented fields.
The specimen shows signs of extensive post-mortem transport, with angles of the braincase
worn off and much of the skull roof and some of the braincase preserved as a mould.
Individual skull roof ossifications cannot be identified, although this may be due to the
dominantly mouldic preservation. There appears to have been a prominent nuchal plate
eminence comparable to acanthothoracids[12-14].

Endoskeletal tissue. The braincase of MPC-FH100/9.1 is well ossified, comprising an
external bony sheath filled with an extensive matrix of spongy tissue (Fig. 2a-b; Extended
Data Fig. 1; Supplementary Videos 1 & 2). The trabecles forming this tissue are irregular and
branching, less than 1mm thick and often curved, and resemble most closely the
endochondral tissue of osteichthyans (Fig. 2c-d). As such, we interpret this as endochondral
bone. Notably, this is found in all preserved regions of the braincase, in contrast to the
isolated trabeculae identified as endochondral bone in *Boreaspis*[15] and *Bothriolepis*[16]. The
margins of the braincase, the endocranial walls, and the boundaries of nerve and blood
canals, are formed from a thicker tissue which we interpret as perichondral bone. This
suggests that the endoskeleton of *Minjinia* comprises osteichthyan-like endochondral bone,
with an ossified perichondrium. To address the possible alternative explanation that it is an
aberrant instance of calcified cartilage, we compared the structure of this tissue with rarely-
preserved mineralized cartilage in the stem chondrichthyan *Diplacanthus crassismus*
(National Museums of Scotland specimen NMS 1891.92.334; Fig. 2e-f) observed using
synchrotron tomography. The cancellae within the endochondral tissue of *Minjinia* are
irregular, with a diameter of approximately 1-2mm. This tissue is distinctly unlike the

calcified cartilage of *Diplacanthus* in appearance, which consists of a densely packed matrix

of irregularly stacked chondrons between 20-60 μm in diameter.

Neurocranium. The braincase is preserved from the level of the right posterior orbital wall

to the posterior end of the occipital ridge. Occipital glenoid condyles are not preserved, but

broad, flat parachordal plates are present, separated by a midline groove that accommodated a

relatively narrow notochordal tunnel. A transverse fissure spans the basicranial surface at

about mid-length of the preserved portion. It clearly demarcates the anterior margin of the

parachordal plates and may correspond to the ventral cranial fissure of crown-group

gnathostomes. However, unlike in crown gnathostomes, it is traversed by a substantial

anterior extension of the cranial notochord. The courses of the lateral dorsal aortae are

marked by a pair of sulci on the lateral margins of the parachordal plates. A narrow, shallow

sulcus for the efferent hyoid artery is present on the preserved right side of the specimen,

immediately behind the level of the orbit (Fig. 1a).

The lateral surface of the braincase is preserved on the right side as a mouldic

impression in the matrix (Fig. 1). A sharply demarcated hyoid fossa is present on the lateral

wall of the otic region (Fig. 1). Posterior to this, a stout but pronounced vagal process with a

pair of rounded eminences likely corresponds to the branchial arch articulations. There is no

evidence for a pair of anterior and posterior divisions to the vagal process, which are

typically seen in other 'placoderms'. A well-developed 'placoderm'-like craniospinal

process is absent; its homologous position is instead covered in perichondral bone and

marked by a low ridge (Fig. 1).

In posterior view, a tall, narrow median otic ridge is evident and resembles the

morphology of *Romundina*[17] and *Arabosteus*[18]. Similar to these taxa, the median otic ridge is

flanked by two large occipital fossae for the epaxial musculature. The notochordal tunnel is

approximately the same size as or smaller than the foramen magnum, as in 'placoderms' and

in contrast with crown-group gnathostomes. A metotic fissure is absent.

Endocast. A partial cranial endocast is preserved, consisting of the hindbrain cavity, partial

midbrain cavity, labyrinth cavities, and posteromedial corner of the orbital region. The two

primary trunk canals of the trigeminal nerve (N.V$_1$ and N.V$_{2,3}$) are preserved (Fig. 1;

Extended Data Fig. 2). The acoustic (N.VIII) and facial nerve (N.VII) canals share a common

trunk canal behind the trigeminal nerves, as in many other 'placoderms' [17,19-21]. The

supraopthalmic branch opens into the rear wall of the orbit and part of its supraorbital course

is preserved (Extended Data Figs. 2, 3). A slender branch extends below the labyrinth and

divides into palatine and hyomandibular branches (Extended Data Figs. 2, 3). As in other

'placoderm'-grade taxa, the vagus nerve (N. X) trunk canal is very large in diameter and exits

from immediately behind the labyrinth cavity (Fig. 1; Extended Data Fig. 2). The spino-

occipital region resembles other 'placoderms' in being extended. At least four spino-occipital

nerve canals are present in a linear series, and the penultimate canal is largest in diameter

(Fig. 1; Extended Data Fig. 2). Intercalating these is a network of occipital artery canals

branching from the dorsal aortae.

The skeletal labyrinth is not complete on either side of the specimen, but can mostly

be reconstructed according to the assumption of bilateral symmetry. The most significant

feature is that the labyrinth and endolymphatic cavity are joined to the main endocavity

chamber (Fig. 1). This is a striking contrast to other 'placoderms' and closely resembles

crown-group gnathostomes[22]. The endolymphatic canals are elongate and tubular, extending

posterolaterally to reach the skull roof, though external openings cannot be clearly identified.

The anterior semi-circular canal follows the saccular cavity closely as in petalichthyids[23](Fig.

1; Extended Data Fig. 2). However, the horizontal and posterior canals appear to extend well

away from the saccular chamber (Fig. 1, Extended Data Fig. 2). The dorsal junctions of the

anterior and posterior canals are joined in a crus commune, as in *Romundina*[17] and

Jagorina[19]. A sinus superior is absent.

Phylogenetic analyses

We conducted phylogenetic analyses under four different protocols: equal weights

parsimony, implied weights parsimony, an unpartitioned Bayesian analysis, and a Bayesian

analysis with characters partitioned by fit determined under implied weights parsimony[24] (see

Extended Data Figs. 4-7). All phylogenetic analyses consistently place *Minjinia* as a stem-

group gnathostome, proximate to the gnathostome crown (Fig. 3, Extended Data Figs 4,5).

Equal weights parsimony recovers *Minjinia* in a position crownward of arthrodires but

outside of a grade consisting of *Entelognathus*, *Ramirosuarezia*, and *Janusiscus*. Under

implied weights, these three taxa move onto the osteichthyan stem and *Minjinia* is placed as

the immediate sister taxon of the gnathostome crown. Under Bayesian analyses, arthrodires

are resolved as more crownward than *Minjinia*. However, the latter analyses fail to recover

arthrodires as a clade and the node uniting them with the crown to the exclusion of *Minjinia*

is extremely weakly supported (posterior probability: 0.52-0.55). Under parsimony, the

crownward position of *Minjinia* is unambiguously supported by the skeletal labyrinth and

endolymphatic duct being confluent with the main cranial cavity[22] (Fig. 3). In common with

arthrodires and the gnathostome crown, *Minjinia* possesses a division of the facial nerve deep

to the transverse otic process (Fig. 1; Extended Data Fig. 2). However, *Minjinia* is excluded

from the gnathostome crown group due to the absence of a metotic fissure and a posterior

dorsal fontanelle, and presence of broad, flat parachordal plates expanded behind the saccular

cavity (Fig. 3, Supplementary Information).

We undertook ancestral states reconstructions to assess the evolutionary history of

endochondral bone (Fig. 3, Extended Data Figs. 6 & 7). Interestingly, parsimony analysis

fails to recover secondary homology of this trait between *Minjinia* and osteichthyans. The

crownward placement of *Minjinia* is, in fact, based on independent evidence relating to

anatomical features of the braincase and endocast. However, the strict precision of parsimony

reconstructions makes it insensitive to underlying uncertainty. To explore this, we used

likelihood reconstructions and compared the ancestral state reconstructions under equal rates

(ER) and all rates different (ARD) variants of the Mkv model on branch-length-rescaled

parsimony trees and Bayesian trees. On the parsimony trees, both models show substantial

non-zero probabilities (0.23 for ER; 0.39 for ARD; Extended Data Table 1) for the presence

of endochondral bone in the common node of *Minjinia* and Osteichthyes (Extended Data Fig.

6) in the parsimony trees. The ARD model shows the best likelihood score and a better AIC

fit for endochondral bone (Extended Data Table 1), favouring repeated losses of this tissue

over multiple gains (see Discussion). The values are substantially lower in the Bayesian trees

(Extended Data Fig. 7, Extended Data Table 1), but this results from the relative positions of

Minjinia and arthordires, which is not well supported in those trees.

Discussion

Minjinia presents an unusual discovery of extensive endochondral bone in a 'placoderm'-

grade fish, with repercussions for the phylogenetic origin of this tissue and the problem of

early gnathostome relationships more generally. The vertebrate skeleton is split into two

systems: the exoskeleton (external achondral dermal bones) and endoskeleton (internal

chondral bones) [1]. Dermal bone evolved at least 450 million years ago in jawless stem

gnathostomes, but the endoskeleton in these taxa is not endochondrally ossified (but see

below). More crownward stem gnathostomes (osteostracans and 'placoderms') surround their

cartilaginous endoskeleton in a sheath of perichondral bone. Extant chondrichthyans lack

both dermal and perichondral bone, possessing a cartilaginous endoskeleton enveloped by

prismatic calcified cartilage. Endochondral bone, in which the cartilaginous endoskeletal

precursor is invaded by and eventually replaced by bone, is widely considered an

osteichthyan apomorphy based on clear prior polarity[3,7,25,26]. However, recent work has cast

doubt on this assertion. The recognition that dermal bone is secondarily lost in

chondrichthyans[27,28] is consonant with prior knowledge of the loss of perichondral bone in

this same lineage [29]. Taken together, this has revived uncertainty about the true phylogenetic

timing of the origin of endochondral ossification[8].

Minjinia does not represent the first report of endochondral bone outside of

Osteichthyes. However, it is by far the most extensive and unequivocal example, and raises

explicit questions in light of the proximity of *Minjinia* to the gnathostome crown. Isolated

examples of trabecular bone, typically restricted to a small region of the neurocranium, have

historically been reported in boreaspid osteostracans[15,30], buchanosteid arthrodires[31] and

petalichthyids[32]. However, these reports have all been dismissed as misidentifications[26],

possibly representing the retreat of perichondral bone deposited during cartilage growth[33].

Most recently, trabeculae in supposed endoskeletal bones of *Bothriolepis* have been termed

endochondral bone[16], although the small scale of these is in line with 'superficial'

perichondral trabeculae seen elsewhere. In line with ref. 26, we found no evidence of

endochondral bone in material of *Buchanosteus* held in the Natural History Museum,

London, or indeed in any other 'placoderms' we have examined. The *Epipetalichthys*

holotype (Museum für Naturkunde, Berlin specimen MB.f.132.1-3) shows an apparently

spongiose infilling in the anterior region of the braincase, but the identity of this structure, or

even whether it is biological, cannot be determined.

Does endochondral bone have a deep origin within the gnathostome stem group? This

would imply repeated losses of this tissue. We do find some statistical support for this

hypothesis (Fig. 3, Extended Data Figs. 6, 7; Extended Data Table 1), and the model is well

justified on prior phylogenetic and biological grounds. Endochondral bone has long been

known to be inconsistently developed across 'primitive' bony fishes: incomplete or entirely

absent ossification of the endoskeleton is known in both Palaeozoic actinopterygians[34] and

sarcopterygians[35], as well as more recent taxa[36]. The frequent absence of endochondral bone

is considered secondary, and other controlling factors such as body size, maturity, mechanical

stress, and buoyancy can determine its degree of development[1]. Our findings are also in

agreement with studies establishing a genetic basis for secondary loss of all bone types within

chondrichthyans[5,37,38], with the failure to produce endochondral bone likely representing

arrested development of chondrocytes as opposed to a primary lack of ability[6].

Another confounding factor in this question is the problem of 'placoderm'

relationships. Although currently resolved as a deeply pectinate grade along the gnathostome

stem, the backbone of this arrangement has poor statistical support and there is a lack of

consistency in the arrangement of plesia. *Minjinia* itself highlights this uncertainty, given its

highly unexpected character combinations. Notwithstanding its endochondral bone and

crown-gnathostome-like inner ear structure, it strongly resembles 'acanthothoracids'—the

'placoderms' widely considered among the most removed from the gnathostome crown (i.e.

most 'primitive'). This apparent character conflict could perhaps be more easily reconciled

with a more coherent (though not necessarily monophyletic) 'placoderm' assemblage.

Indeed, the highly pectinate structure of the 'placoderm' grade seems symptomatic of an

overemphasis on characters and taxa resembling the crown group, thereby undersampling

characters that could stabilise a clear picture of 'placoderm' interrelationships.

Minjinia reveals new data on 'placoderm' endoskeleton and tissue diversity from

Mongolia—an otherwise extremely poorly known biogeographic realm for early

gnathostomes. The phylogenetic placement of this 'acanthothoracid'-like taxon crownward of

all non-maxillate 'placoderms', in conjunction with possession of extensive endochondral

bone, highlights the importance of material from traditionally undersampled geographic

areas. The presence of endochondral bone renews the hypothesis that this tissue is

evolutionarily ancient and was lost secondarily in chondrichthyans[7,8]. This view is overall

consistent with evidence of generalised bone loss in chondrichthyans, potentially as a result

of the suppression of bone-generating molecular genetic pathways[6,38]. Continued work in

Mongolia and re-evaluation of phylogenetic datasets will be necessary to address this, with

the results likely to lead to substantial re-evaluation of gnathostome phylogeny.

Acknowledgements. M. Bolortsetseg generously assisted MDB with contacts and field

experience in Mongolia. Fieldwork was supported by National Geographic Society grants

CRE 8769-10 and GEFNE35-12 to MDB. ALJ's field contributions were supported by funds

from the Anna Maria Lundin's stipend from Smålands Nation, Uppsala University. RS's field

contributions were supported by a Royal Society Research Grant and the University of

Manchester. The majority of this work was supported by the European Research Council

(ERC) under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC

Grant Agreement number 311092 to MDB. RPD was also supported by the Île-de-France

DIM (domaine d'intérêt majeur) matériaux anciens et patrimoniaux grant PHARE. Stig

Walsh is thanked for access and loan of specimen at the National Museums of Scotland.

Synchrotron tomography was performed at the ESRF (application LS 2451) with the
assistance of Paul Tafforeau. SG was supported by a Royal Society Dorothy Hodgkin
Research Fellowship. Matt Friedman is thanked for undertaking the X-ray computed
microtomography analysis. TNT was made available with the support of the Willi Hennig
Society.

Author Contributions: MDB conceived and designed the study. MDB, AJ, YAA, and EZ
participated in all field seasons. RPD and AJ undertook preliminary CT scanning and
segmentation that revealed the fossil was a 'placoderm' and had endochondral bone. RS
discovered the first vertebrate remains in the first field season at Yamaat Gol in 2010. SG did
most of the segmentation of *Minjinia* with input from MDB. AJ performed segmentation of
Diplacanthus tissue. MC provided input on occipital comparative morphology of
'placoderms'. RPD provided data and comparative analyses and data for endoskeletal tissue.
YAA provided background on the geology, palaeontology, and stratigraphy of the type
location; EZ and YAA organized field logistics and permitting. MDB, SG, MC, RPD, and AJ
undertook the anatomical interpretation and prepared the figures. MDB and SG conducted the
phylogenetic analyses. RS conducted the parsimony branch support analyses. TG wrote the
script for generating MrBayes partitions from TNT's character fits table and conducted the
likelihood and model-fitting analyses. The manuscript was written by MDB, RPD, and SG.

Author Information:

Correspondence and requests for materials should be addressed to
m.brazeau@imperial.ac.uk.

Methods

X-ray computed microtomography. We scanned MPC-FH100/9.1 using the Nikon XT
225s at the Museum of Paleontology, University of Michigan with the following parameters:
200kV, 140μA, over 3123 projections and a voxel size of 32.92μm. We conducted
segmentation using Mimics 19.0 (http://biomedical.materialise.com/mimics; Materialise,
Leuven, Belgium) and we imaged models for publication using Blender
(https://www.blender.org).

Synchrotron light propagation phase contrast tomography. We imaged *Diplacanthus*

crassismus specimen NMS 1891.92.334 on Beamline 19 of the European Synchrotron

Radiation Facility, using propagation phase-contrast synchrotron microtomography. We

performed a spot scan with an energy of 116keV, achieving a voxel size of 0.55 μm. We

processed the resulting tomograms using VG StudioMax 2.2 (Volume Graphics, Germany),

and prepared images in Blender.

Phylogenetic analysis. We conducted a parsimony analysis using TNT 1.5 [39] and Bayesian

analysis using MrBayes v 3.2.7[40]. The dataset consisted of 95 taxa and 284 discrete

characters based on a pre-existing dataset[41]. We employed Osteostraci and Galeaspida as

composite outgroups. We conducted parsimony analysis using both equal weights and

implied weights methods. Global settings were 1000 search replicates and a hold of up to 1

million trees. Equal weights parsimony analyses were conducted using the ratchet with

default settings. Implied weights parsimony used a concavity parameter of 3 and the search

was without the ratchet. Command lists are included in Supplementary Information. We

conducted Bayesian analysis using both a partitioned and unpartitioned dataset. We used the

Mkv model [42] and gamma rate distribution. We ran the analyses for 5 million generations

with a relative burn-in fraction of 0.25. Runs were checked for convergence using Tracer[43].

We partitioned the dataset using a newly proposed method[24] that partitions the data according

to homoplasy levels. Using the results of implied weights parsimony conducted in TNT, we

created a text table of character fit values. We wrote an R[44] script to generate a list of

partition commands for MrBayes. To reduce the number of partitions with small numbers of

characters, we concatenated the partitions by rounding the fitness scores to 2 significant

figures, yielding 10 individual partitions.

We assessed parsimony ancestral states visually using Mesquite[45]. Likelihood and

Bayesian ancestral states were estimated in R using the castor package[46]. Prior to calculating

likelihood ancestral states on parsimony trees, we scaled branch lengths using PAUP*[47] and

calculated the likelihood scores for all of the trees under the Mkv model. The trees were then

exported with branch lengths. To account for overall uncertainty in tree estimates, we

estimated ancestral states on 100 trees randomly selected from the fundamental set of most

parsimonious trees and two times 50 trees selected from the 75% last trees of each posterior

tree distribution from the Bayesian analysis. We then run an ancestral states estimation Mk

model (using the castor R package) using both the Equal Rates (ER) and All Rates Different

(ARD) models. This resulted in 400 ancestral states estimations. For each estimation we

extracted the overlap log likelihood, the AIC (counting one parameter for the ER model and

two for the ARD model) and the scaled log likelihood (probability) for the presence and

absence of the endochondral bone character (character 4) for the last common node of

Minjinia and crown-group gnathostomes. We present the median value of these distributions

of the estimations overall log likelihoods, AICs and presence or absence of endochondral

bone in Extended Data Table 1.

Data availability

The holotype specimen of *Minjinia turgenensis* will be permanently deposited in the

collections of the Institute of Paleontology, Mongolian Academy of Sciences. Original

tomograms are available at (doi:10.6084/m9.figshare.12301229) and rendered models are

available at (doi:10.6084/m9.figshare.12301223). The phylogenetic character list and dataset

are available as Supplementary Information S1 and S2. The LifeScience Identifier for

Minjinia turgenensis is urn:lsid:zoobank.org:act:82A1CEEC-B990-47FF-927A-

D2F0B59AEA87

344
345   **Code availability**

346   R code for generating partitions based on character fits and code for likelihood ancestral

347   states reconstructions and plots are available in the Supplementary Information.

Fig. 1 | MPC-FH100/9.1 a 'placoderm' skull roof and braincase from the Early Devonian of Mongolia. a, Ventral view. **b**, Dorsal view. **c**, Left lateral view. **d**, Posterior view. **e**, Braincase endocavity in dorsal view. Taupe: endoskeleton; grey: mould; pink: endocavity; blue: exoskeleton. a.scc., anterior semicircular canal; cav.end., endolymphatic cavity; crsp.ri, craniospinal ridge; d.end., endolymphatic duct; e.hy.a., sulcus for the efferent hyoid artery; f.m.ep., epaxial muscle fossa; fo.mag., foramen magnum; h.scc., horizontal semicicular canal; l.d.ao, sulcus for the lateral dorsal aorta; N.V, trigeminal nerve canal; N.VII, facial nerve canal; N.VIII, acoustic nerve canal; N.X, vagus nerve canal; nch., notochordal canal; occ.ri, occipital ridge; orb., orbit; p.scc, posterior semicircular canal; pr.pv., paravagal process; sac., sacculus; soc., spino-occipital nerve canals. Scale bar, 20 mm.

Fig. 2 | Endoskeletal mineralisation in fossil gnathostomes. a, Transverse tomographic slice through MPC-FH100/9.1. **b**, Three-dimensional rendering of trabecular bone structure. **c**, Transverse tomographic section through the braincase of the osteichthyan *Ligulalepis*. **d**, Three-dimensional rendering of the trabecular bone in *Ligulalepis* (**c** and **d** use data from[41]). **e**, Synchrotron tomography image of the calcified cartilage of the stem-group chondrichthyan *Diplacanthus crassisimus* specimen NMS 1891.92.334. **f**, Semi-transparent three-dimensional structure of calcified cartilage of NMS 1891.92.334. Scale bars, **a** and **b,** 10 mm; c and d, 1 mm ; **c** and **h**, 150 µm.

Fig. 3 | Summary phylogenetic relations of early gnathostomes showing distribution of endochondral bone and exoskeletal armour. Squares at nodes indicate parsimony reconstruction for endochondral bone. Pie charts at nodes show likelihood reconstructions for the same character under the all-rates-different model (see Extended Data Figs 6 & 7 for competing reconstructions). Grey box indicates uncertainty. Loss of endochondral bone maps closely with generalised loss of bone in chondrichthyans where exoskeletal armour and perichondral bone are also absent.

Extended Data Fig. 1 | Tomograms of endoskeletal ossification in *Minjinia*. Top row:

semi-coronal sections through braincase. Double-headed arrows indicate anterior-posterior

(a-p) dorsal-ventral (d-v) axes. Bottom row: semi-transverse sections through posterior part

of endocranium. Voids of black space represent mouldic preservation. Scale bars, 10 mm and

apply across each row of panels.

Extended Data Fig. 2 | Braincase endocavity of *Minjinia*. a, Semi-transparent rendering of

skull roof and braincase (grey and blue) showing extent of endocavity (pink). **b,** Ventral

view. **c,** Dorsal view. a.scc., anterior semicircular canal; cav.end., endolymphatic cavity;

d.end., endolymphatic duct; h.scc., horizontal semicicular canal; l.d.ao., sulcus for the lateral

dorsal aorta; N.V, trigeminal nerve canal; N.VIIhm, hyomandibular branch of facial nerve

canal; N.VIIpal, palatine branch of facial nerve canal; N.VIII, acoustic nerve canal; N.X,

vagus nerve canal, N.X$_a$, anterior branch of vagus nerve canal; N.Xp, posterior branch of

vagus nerve canal; occ.a, occipital artery canals; p.scc, posterior semicircular canal; sac.,

sacculus; soc., spino-occipital nerve canals; sup.opth, canal for supra-ophtalmic nerve. Scale

bars, 20 mm (upper scale bar associates with **a**, lower scale bar associates with **b** and **c**).

397
Extended Data Fig. 3 | Right orbital wall and innervation pattern of *Minjinia*. a, orbit in anterolateral view showing disposition of nerve openings (pink infill). **b**, endocast in the same perspective showing the relationship between never canals and endocast.

Extended Data Fig. 4 | Results of phylogenetic parsimony analysis. Dataset consists of 95

taxa and 284 characters. Both trees are strict consensus topologies. Equal weights parsimony

analysis using the ratchet resulted in 240 trees with a length of 832 steps. Implied weights

parsimony analysis using random addition sequence + branch-swapping resulted in two

optimal trees with score 85.23240. Double-digit figures above internal branches are bootstrap

values of 50% and over; single-digit figures below branches are Bremer decay index values.

Blue shading: osteichthyan total group (dark blue: crown group); orange shading:

chondrichthyan total group (dark orange: crown group).

Extended Data Fig. 5 | Results of Bayesian phylogenetic analysis using both partitioned
and unpartitioned data. Majority-rules consensus trees with posterior probabilities shown
along branches. Blue shading: osteichthyan total group (dark blue: crown group); orange
shading: chondrichthyan total group (dark orange: crown group).

**Endochondral bone estimation
(parsimony – ARD)**

**Endochondral bone estimation
(parsimony – ER)**

Extended Data Fig. 6 | Likelihood ancestral state mapping of endochondral bone on equal weights parsimony results. ARD, all rates different model; ER, equal rates model.

Extended Data Fig. 7 | Likelihood ancestral state mapping of endochondral bone on unpartitioned Bayesian analysis results. ARD, all rates different model; ER, equal rates model.

425 **Extended Data Table 1 | Tree distribution (100) ancestral states estimation results.** ER =

426 Equal rates model; ARD = All Rates Different model. The columns AIC and log.lik represent

427 the median AIC and log.lik across the 100 parsimony and bayesian trees (for both models).

428 The columns Absent and Present represent the median scaled likelihood for the endochondral

429 bone state.

Node	Tree	Model	log.lik	AIC	Absent	Present
Minjinia:crown gnathostomes	Parsimony	ER	-27.60	57.20	0.94	0.06
		ARD	-25.47	54.93	0.61	0.39
	Bayesian	ER	-29.94	61.89	0.98	0.02
		ARD	-27.69	59.38	0.82	0.18

430
431
432 **References**
433

1.   Hall, B. K. *Bones and Cartilage*. (Academic Press, 2005).
2.   Janvier, P. *Early Vertebrates*. (Oxford University Press, 1996).
3.   Friedman, M. & Brazeau, M. A Reappraisal of the Origin and Basal Radiation of the
Osteichthyes. *J Vertebr Paleontol* **30,** 36–56 (2010).
4.   Brazeau, M. D. & Friedman, M. The characters of Palaeozoic jawed vertebrates. *Zool*
J Linn Soc **170,** 779–821 (2014).
5.   Eames, B. F. *et al.* Skeletogenesis in the swell shark Cephaloscyllium ventriosum.
Journal of Anatomy **210,** 542–554 (2007).
6.   Marconi, A., Hancock-Ronemus, A. & Gillis, J. A. Adult chondrogenesis and
spontaneous cartilage repair in the skate, *Leucoraja erinacea*. *eLife Sciences* **9,** 2813
(2020).
7.   Maisey, J. G. Heads and tails: a chordate phylogeny. *Cladistics* **2,** 201–256 (1986).
8.   Zhu, M. Bone gain and loss: insights from genomes and fossils. *Natl Sci Rev* **1,** 490–
492 (2014).
9.   Alekseeva, R. E., Mendbayar, B. & Erlanger, O. A. *Brachiopods and Biostratigraphy*
of the Lower Devonian of Mongolia. **16,** (Nauka, 1981).
10.   Alekseeva, R. E. *Devonian Biostratigraphy of Mongolia*. (Nauka, 1993).
11.   Mark-Kurik, E. in *Morphology, Phylogeny and Paleobiogeography of Fossil Fishes*
(eds. Elliott, D. K., Maisey, J., Yu, X. & Miao, D.) 101–106 (2010).
12.   Stensiö, E. A. Contributions to the knowledge of the vertebrate fauna of the Silurian
and Devonian of western Podolia. II. Notes on two arthrodires from the Downtonian of
Podolia. **35A,** 1–83 (1944).
13.   Ørvig, T. Description, with special reference to the dermal skeleton, of a new radotinid
arthrodire from the Gedinnian of Arctic Canada. *Colloque international C.N.R.S. no.*
218 43–71 (1975).
14.   Vaškaninová, V. & Ahlberg, P. E. Unique diversity of acanthothoracid placoderms
(basal jawed vertebrates) in the Early Devonian of the Prague Basin, Czech Republic:
A new look at Radotina and Holopetalichthys. *PLoS ONE* **12,** e0174794 (2017).

15.   Wängsjö, G. The Downtonian and Devonian vertebrates of Spitsbergen. IX. *Norsk*
Polarinstitutt Skrifter **97,** 1–611 (1952).

16.   Charest, F., Johanson, Z. & Cloutier, R. Loss in the making: absence of pelvic fins and
presence of paedomorphic pelvic girdles in a Late Devonian antiarch placoderm
(jawed stem-gnathostome). *Biology Letters* **14,** 20180199 (2018).

17.   Dupret, V., Sanchez, S., Goujet, D. & Ahlberg, P. E. The internal cranial anatomy of
Romundina stellina Ørvig, 1975 (Vertebrata, Placodermi, Acanthothoraci) and the
origin of jawed vertebrates—Anatomical atlas of a primitive gnathostome. *PLoS ONE*
12, e0171241 (2017).

18.   Olive, S., Goujet, D., Lelièvre, H. & Janjou, D. A new Placoderm fish
(Acanthothoraci) from the Early Devonian Jauf Formation (Saudi Arabia).
Geodiversitas **33,** 393–409 (2011).

19.   Stensiö, E. A. La cavité labyrinthique, l'ossification sclérotique et l'orbite de *Jagorina*.
Colloques internationaux du Centre national de la Recherche scientifique **21,** 9–43
(1950).

20.   Stensiö, E. A. Anatomical studies on the arthrodiran head. *K Sven Vetenskapsakad*
Handl **9,** 1–419 (1963).

21.   Goujet, D. *Les poissons placodermes du Spitsberg*. (Cahiers de Paléontologie, Editions
du CNRS, 1984).

22.   Davis, S. P., Finarelli, J. A. & Coates, M. I. *Acanthodes* and shark-like conditions in
the last common ancestor of modern gnathostomes. *Nature* **486,** 247–250 (2012).

23.   Castiello, M. & Brazeau, M. D. Neurocranial anatomy of the petalichthyid placoderm
Shearsbyaspis oepiki Young revealed by X☐ray computed microtomography.
Palaeontology **21,** 754 (2018).

24.   Rosa, B. B., Melo, G. A. R. & Barbeitos, M. S. Homoplasy-Based Partitioning
Outperforms Alternatives in Bayesian Analysis of Discrete Morphological Data.
Systematic Biology **54,** 373 (2019).

25.   Rosen, D. E., Forey, P. L., Gardiner, B. G. & Patterson, C. Lungfishes, tetrapods,
paleontology, and plesiomorphy. *Bull Am Mus Nat Hist* **167,** 159–276 (1981).

26.   Gardiner, B. G. The relationships of the palaeoniscid fishes, a review based on new
specimens of *Mimia* and *Moythomasia* from Upper Devonian of Western Australia.
Bull Br Mus nat Hist (Geol) **37,** 173–428 (1984).

27.   Zhu, M. *et al.* A Silurian placoderm with osteichthyan-like marginal jaw bones. *Nature*
502, 188–193 (2013).

28.   Giles, S., Friedman, M. & Brazeau, M. D. Osteichthyan-like cranial conditions in an
Early Devonian stem gnathostome. *Nature* **520,** 82–85 (2015).

29.   Miles, R. S. in *Interrelationships of Fishes* (eds. Greenwood, P. H., Miles, R. S. &
Patterson, C.) 63–103 (Academic Press London, 1973).

30.   Janvier, P. *Les céphalaspides du Spitsberg*. (Éditions du Centre National de la
Recherche Scientifique, 1985).

31.   Young, G. C. New information on the structure and relationships of *Buchanosteus*
(Placodermi: Euarthrodira) from the Early Devonian of New South Wales. *Zool J Linn*
Soc **66,** 309–352 (1979).

32.   Stensiö, E. A. On the head of the macropetalichthyids. *Field Museum of Natural*
History Publication Geological Series **4,** 87–197 (1925).

33.   Ørvig, T. Notes on some Paleozoic lower vertebrates from Spitsbergen and North
America. *Norsk Geologisk Tidsskrift* **37,** 285–353 (1957).

34.   Schaeffer, B. The braincase of the holostean fish *Macrepistius*, with comments on
neurocranial ossification in the Actinopterygii. *American Museum Novitates* **2459,** 1–
34 (1971).

35.   Cloutier, R. in *Devonian Fishes and Plants* (eds. Schultze, H.-P. & Cloutier, R.) 227–
513   247 (Verlag Dr. Friedrich Pfeil, 1996).
514   36.   Grande, L. & Bemis, W. E. Osteology and Phylogenetic Relationships of Fossil and
515   Recent Paddlefishes (Polyodontidae) with Comments on the Interrelationships of
516   Acipenseriformes. *Memoir (Society of Vertebrate Paleontology)* **11 (Supplement to**
517   **Number 1),** 1–121 (1991).
518   37.   Venkatesh, B. *et al.* Elephant shark genome provides unique insights into gnathostome
519   evolution. *Nature* **505,** 174–179 (2014).
520   38.   Ryll, B., Sanchez, S., Haitina, T., Tafforeau, P. & Ahlberg, P. E. The genome of
521   *Callorhinchus* and the fossil record: a new perspective on SCPP gene evolution in
522   gnathostomes. *Evolution & Development* **16,** 123–124 (2014).
523   39.   Goloboff, P. A. & Catalano, S. A. TNT version 1.5, including a full implementation of
524   phylogenetic morphometrics. *Cladistics* **32,** 221–238 (2016).
525   40.   Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under
526   mixed models. *Bioinformatics* **19,** 1572–1574 (2003).
527   41.   Clement, A. M. *et al.* Neurocranial anatomy of an enigmatic Early Devonian fish sheds
528   light on early osteichthyan evolution. *eLife Sciences* **7,** e34349 (2018).
529   42.   Lewis, P. O. A likelihood approach to estimating phylogeny from discrete
530   morphological character data. *Systematic Biology* **50,** 913 (2001).
531   43.   Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. Posterior
532   Summarization in Bayesian Phylogenetics Using Tracer 1.7. *Systematic Biology* **67,**
533   901–904 (2018).
534   44.   R Core Team. *R: A language and environment for statistical computing.* (R
535   Foundation for Statistical Computing, 2019).
536   45.   Maddison, D. R. & Maddison, W. P. *Mesquite.* (2019).
537   46.   Louca, S. & Doebeli, M. Efficient comparative phylogenetics on large trees.
538   *Bioinformatics* **34,** 1053–1055 (2017).
539   47.   Swofford, D. L. *PAUP*.* 4.0a166, (2019).
540
541