Endochondral bone in an Early Devonian 'placoderm' from Mongolia

Martin D. Brazeau[1,2]*, Sam Giles [2,3,4], Richard P. Dearden[1,5], Anna Jerve[1,6], Y.A.

Ariunchimeg[7], E. Zorig[8], Robert Sansom[9], Thomas Guillerme[10], Marco Castiello[1]

[1] *Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst*

Rd, Ascot, SL5 7PY, UK;

[2] *Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7*

5BD, UK;

[3] *School of Geography, Earth and Environmental Sciences, University of Birmingham,*

Birmingham, UK;

[4] *Department of Earth Sciences, University of Oxford, South Parks Road, Oxford, OX1 3AN,*

UK;

[5] *CR2P Centre de Recherche en Paléontologie – Paris, Muséum national d'Histoire*

naturelle, Sorbonne Universités, CNRS, CP 38, 57 Rue Cuvier, 75231, Paris, Cedex 05,

France

[6] *Department of Organismal Biology, Subdepartment of Evolution and Development,*

Uppsala University, Norbyvägen 18A, 752 36 Uppsala, Sweden;

[7] *Natural History Museum, P.O. Box 46/52, Ulaanbaatar 1420, Mongolia*

[8] *Institute of Paleontology, Mongolian Academy of Science, P.O. Box 46/650, S. Danzan*

Street 3/1, Chingeltei District. Ulaanbaatar 15160, Mongolia;

[9] *School of Earth and Environmental Sciences, University of Manchester, Manchester M13*

9PT, UK;

[10] *Department of Animal and Plant Sciences, The University of Sheffield, Sheffield S10 2TN,*

UK;

* Author for correspondence.

Endochondral bone is the main internal skeletal tissue of nearly all osteichthyans—the

group comprising more than 60,000 living species of bony fishes and tetrapods.

Chondrichthyans (sharks and their kin) are the living sister group of osteichthyans and

have cartilaginous endoskeletons, long considered the ancestral condition for all jawed

vertebrates (gnathostomes). The absence of bone in modern jawless fishes and the

absence of endochondral ossification in early fossil gnathostomes appears to lend

support to this conclusion. Here we report the discovery of extensive endochondral bone

in _Minjinia turgenensis_, a new genus and species of 'placoderm'-like fish from the Early

Devonian (Pragian) of western Mongolia described using x-ray computed

microtomography (XR-μCT). The fossil consists of a partial skull roof and braincase

with anatomical details providing strong evidence of placement in the gnathostome stem

group. However, its endochondral space is filled with an extensive network of fine

trabeculae resembling the endochondral bone of osteichthyans. Phylogenetic analyses

place this new taxon as a proximate sister group of the gnathostome crown. These

results provide direct support for theories of generalised bone loss in chondrichthyans.

Furthermore, they revive theories of a phylogenetically deeper origin of endochondral

bone and its absence in chondrichthyans as a secondary condition.

The vertebrate skeleton comprises two main systems: the exoskeleton (external achondral

dermal bones) and endoskeleton (internal chondral bones and cartilages, as well as some

intramembranous bones)[1]. An ossified exoskeleton evolved at least 450 million years ago in

jawless stem gnathostomes[2,3], but the endoskeleton in these taxa is not endochondrally

ossified. Endochondral bone, in which the cartilaginous endoskeletal precursor is invaded by

and eventually replaced by bone, is widely considered an osteichthyan apomorphy[3-7] and

such a reliable identifying character gives the group its name. Extant chondrichthyans lack

dermal bone and possess a mainly cartilaginous endoskeleton enveloped by a structurally

diverse range of tessellate calcified cartilage[8]. Outgroups of the gnathostome crown also lack

endochondral ossification. Galeaspids surround their cartilaginous skeleton in globular

calcified cartilage{NianZhong:2005tj}, while osteostracan and 'placoderm' endoskeletons

were sheathed in perichondral bone[3]. Consequently, the last common ancestor of jawed

vertebrates was long thought to have been perichondrally ossified, but lacking endochondral

ossification[3].

In this paper, we describe a new genus and species of 'placoderm' from the Early

Devonian of western Mongolia. Although Mongolia is known for some of the geologically

oldest putative gnathostome fossils (isolated chondrichthyan-like scales [9-12]), it remains a

poorly sampled region of the world with respect to early vertebrates. 'Placoderms' were until

now known from only a single fragmentary occurrence[13] in the early Middle Devonian

(Eifelian). Our new data highlight the importance of Mongolia as a key region for studies of

early gnathostome evolution. We describe a braincase and partial skull roof representing the

first substantial body fossil of an early gnathostome from Mongolia and displaying an

unexpected occurrence of endochondral bone analysed using XR-μCT. We conducted

phylogenetic analyses to reconstruct the evolutionary relationships of this new taxon. To

explore the evolutionary history of endochondral bone in light of this new discovery, we used

parsimony and maximum likelihood ancestral states reconstruction. Finally, we discuss these

results in the context of earlier statements about endochondral bone in non-osteichthyans,

new developments in understanding the complexity and diversity of chondrichthyan

endoskeletal tissues, and current uncertainties about early gnathostome phylogenetic

relationships.

Systematic palaeontology

78   Gnathostomata Gegenbaur, 1874[14]

79   *Minjinia turgenensis* gen. et sp. nov.

Etymology. Generic name honours the memory of Chuluun Minjin for his extensive

contributions to the Palaeozoic stratigraphy of Mongolia, his enthusiastic support of this

work, and introducing us to the Yamaat River locality. Specific name recognises the

provenance of the fossil from the Turgen region, Uvs aimag of western Mongolia.

Holotype. Institute of Paleontology, Mongolian Academy of Sciences MPC-FH100/9.1, a

partial braincase and skull roof.

Type locality. Turgen Strictly Protected Area, Uvs province, western Mongolia; near the top

of the stratigraphic sequence that occurs between the Tsagaan-Salaat and Yamaat Rivers.

Formation and age. Upper part of Tsagaansalaat Formation, Pragian (Early Devonian) [15,16].

Diagnosis. 'Placoderm'-grade stem gnathostome with endochondral bone, deep epaxial

muscle cavities flanking a slender occipital ridge, and the following possible autapomorphies:

dermal bones covered in sparsely placed tubercles, penultimate spino-occipital nerve canal

substantially larger in diameter than others.

Description

MPC-FH100/9.1 consists of a partial braincase and skull roof (Fig. 1). The skull roof is

ornamented with sparsely distributed finely ridged tubercles resembling those of the Siberian

'placoderm' *Dolganosteus*[17]; the tubercles become more broadly separated towards the

midline of the skull. They are distinct from those of *Dolganosteus* in that towards the midline

of the skull roof, the tubercles are larger and more pointed. The specimen shows signs of

extensive post-mortem transport, with angles of the braincase worn off and much of the skull

roof and some of the braincase preserved as a mould. Individual skull roof ossifications

cannot be identified, although this may be due to the dominantly mouldic preservation. There

appears to have been a prominent nuchal plate eminence comparable to certain

acanthothoracids such as *Romundina*[18] and *Arabosteus*[19].

Endoskeletal tissue. The braincase of MPC-FH100/9.1 is well ossified, comprising an

external bony sheath filled with an extensive matrix of spongy tissue (Fig. 2a-b; Extended

Data Fig. 1; Supplementary Video 1). The trabecles forming this tissue are irregular and

branching, less than 1 mm thick and often curved, and resemble most closely the

endochondral tissue of osteichthyans (Fig. 2c-d; Supplementary Video 2). As such, we

interpret this as endochondral bone. Notably, this is found in all preserved regions of the

braincase, in contrast to the isolated trabeculae previously identified as endochondral bone in

Boreaspis[20] and *Bothriolepis*[21]. The margins of the braincase, the endocranial walls, and the

boundaries of nerve and blood canals, are formed from a thicker tissue which we interpret as

perichondral bone. This suggests that the endoskeleton of *Minjinia* comprises osteichthyan-

like endochondral bone, with an ossified perichondrium. To address the possible alternative

explanation that it is an aberrant instance of calcified cartilage, we compared the structure of

this tissue with rarely-preserved mineralized cartilage in the stem chondrichthyan

Diplacanthus crassismus (National Museums of Scotland specimen NMS 1891.92.334; Fig.

2e-f) observed using synchrotron tomography. The cancellae within the endochondral tissue

of *Minjinia* are irregular, with a diameter of approximately 1-2 mm. This tissue is distinctly

unlike the calcified cartilage of *Diplacanthus* in appearance, which consists of a densely

packed matrix of irregularly stacked chondrons between 20-60 μm in diameter.

Neurocranium. The braincase is preserved from the level of the right posterior orbital wall

to the posterior end of the occipital ridge. Occipital glenoid condyles are not preserved, but

much of the rest of the broad, flat parachordal region is present, separated by a midline

groove that accommodated a relatively narrow notochordal tunnel. An asymmetric transverse

fissure spans the basicranial surface at about mid-length of the preserved portion. It appears

to demarcate the anterior margin of the parachordal plates and may correspond to the ventral

cranial fissure of crown-group gnathostomes. However, unlike in crown gnathostomes, it is

traversed by a substantial anterior extension of the cranial notochord. The courses of the

lateral dorsal aortae are marked by a pair of sulci on the lateral margins of the parachordal

plates, though only a short part of the canal is preserved on the right side of the specimen. A

narrow, shallow sulcus for the efferent hyoid artery is present on the preserved right side of

the specimen, immediately behind the level of the orbit (Fig. 1a).

The lateral surface of the braincase is preserved on the right side as a mouldic

impression in the matrix (Fig. 1). A sharply demarcated hyoid fossa is present on the lateral

wall of the otic region (Fig. 1). Posterior to this, a stout but pronounced vagal process with a

pair of rounded eminences likely corresponds to the branchial arch articulations. There is no

evidence for a pair of anterior and posterior divisions to the vagal process, which are

typically seen in other 'placoderms'. A well-developed 'placoderm'-like craniospinal

process is absent; its homologous position is instead covered in perichondral bone and

marked by a low ridge (Fig. 1).

In posterior view, a tall, narrow median occipital ridge is evident and resembles the

morphology of *Romundina*[22] and *Arabosteus*[19]. Similar to these taxa, the median otic ridge is

flanked by two large occipital fossae for the epaxial musculature. The notochordal tunnel is

approximately the same size as or smaller than the foramen magnum, as in 'placoderms' and

in contrast with crown-group gnathostomes. A metotic fissure is absent.

Endocast. A partial cranial endocast is preserved, consisting of the hindbrain cavity, partial

midbrain cavity, labyrinth cavities, and posteromedial corner of the orbital region. The two

primary trunk canals of the trigeminal nerve (N.V_1 and N.$V_{2,3}$) are preserved (Fig. 3). The

acoustic (N.VIII) and facial nerve (N.VII) canals share a common trunk canal behind the

trigeminal nerves, as in many other 'placoderms' [22-25]. The facial nerve canal branches into

palatal and hyomandibular branches between the saccular chamber and rear orbit wall (Fig. 3;

Extended Data Fig. 2), indicating this division was internal (deep) to the otic process. The

supraophthalmic branch opens into the rear wall of the orbit and part of its supraorbital

course is preserved (Fig. 3; Extended Data Fig. 2). A slender branch extends below the

labyrinth and divides into palatine and hyomandibular branches (Fig 3; Extended Data Fig.

2). As in other 'placoderm'-grade taxa, the vagus nerve (N. X) trunk canal is very large in

diameter and exits from immediately behind the labyrinth cavity (Fig. 3). The spino-occipital

region resembles other 'placoderms' in being extended. At least four spino-occipital nerve

canals are present in a linear series, and the penultimate canal is largest in diameter (Fig. 3).

Intercalating these is a network of occipital artery canals branching from the dorsal aortae.

The skeletal labyrinth is not complete on either side of the specimen, but can mostly

be reconstructed according to the assumption of bilateral symmetry. The most significant

feature is that the labyrinth and endolymphatic cavity are joined to the main endocavity

chamber (Fig. 3). This is a striking contrast to other 'placoderms' and closely resembles

crown-group gnathostomes[26]. The endolymphatic canals are elongate and tubular, extending

posterolaterally to reach the skull roof, though external openings cannot be clearly identified.

The anterior semi-circular canal follows the saccular cavity closely as in petalichthyids[27](Fig.

3). However, the horizontal and posterior canals appear to extend well away from the

saccular chamber (Fig. 3). The dorsal junctions of the anterior and posterior canals are joined

in a crus commune, as in *Romundina*[22] and *Jagorina*[23]. A sinus superior is absent.

Phylogenetic analyses

We conducted phylogenetic analyses under four different protocols: equal weights

parsimony, implied weights parsimony, an unpartitioned Bayesian analysis, and a Bayesian

analysis with characters partitioned by fit determined under implied weights parsimony[28] (see

Extended Data Figs. 3-6). All phylogenetic analyses consistently place *Minjinia* as a stem-

group gnathostome, proximate to the gnathostome crown (Fig. 4, Extended Data Figs 3, 4).

Minjinia is recovered in a position crownward of arthrodires but outside of a grade consisting

of *Entelognathus*, *Ramirosuarezia*, and *Janusiscus*. Under implied weights parsimony, these

three taxa move onto the osteichthyan stem and *Minjinia* is placed as the immediate sister

taxon of the gnathostome crown. Under parsimony, the crownward position of *Minjinia* is

unambiguously supported by the skeletal labyrinth and endolymphatic duct being confluent

with the main cranial cavity[26] (Supplementary Information). In common with arthrodires and

the gnathostome crown[29], *Minjinia* possesses a division of the facial nerve(Fig. 3; Extended

Data Fig. 2) deep to the transverse otic process. However, *Minjinia* is excluded from the

gnathostome crown group due to the absences of a metotic fissure and a posterior dorsal

fontanelle, and presence of broad, flat parachordal plates expanded behind the saccular cavity

(Fig. 3, Supplementary Information).

We undertook ancestral states reconstructions to assess the evolutionary history of

endochondral bone (Fig. 4; Extended Data Figs. 5 & 6; Supplementary Information).

Interestingly, parsimony analysis fails to recover secondary homology of this trait between

Minjinia and osteichthyans. The crownward placement of *Minjinia* is, in fact, based on

independent evidence relating to anatomical features of the braincase and endocast. However,

the resolution becomes ambiguous if missing data in either *Entelognathus* or *Ramirosuarezia*

are resolved as having endochondral bone. The reconstruction becomes similarly ambiguous

if *Janusiscus* is moved a single branch (requiring only two additional steps) onto the

chondrichthyan stem. The strict precision of parsimony reconstructions makes it insensitive

to this underlying uncertainty. To explore this, we used likelihood reconstructions and

compared the ancestral state reconstructions under equal rates (ER) and all rates different

(ARD) variants of the Mkv model on branch-length-rescaled parsimony trees and Bayesian

trees. Both models show substantial non-zero marginal likelihoods if endochondral bone is

assumed present in the common node of *Minjinia* and Osteichthyes, with ARD strongly

favouring its presence (0.33 for ER; 0.81 for ARD; Fig. 3, Table 1, Extended Data Figs. 5, 6,

Supplementary Table 1). Under the ARD model, there is nearly equivocal support for

presence or absence of endochondral bone at the gnathostome crown node (Table 1). The

ARD model shows the best fit for endochondral bone (likelihood ratios 4.75 for parsimony [p

= 0.029] and 5.26 for Bayesian, [p = 0.022]) (Table 1, Supplementary Table 1), favouring

repeated losses of this tissue over multiple gains (see Discussion).

Discussion

Minjinia turgenensis presents an unexpected discovery of extensive endochondral bone in a

'placoderm'-grade fish, with repercussions for the phylogenetic origin of this tissue and the

problem of early gnathostome relationships more generally. The prevailing hypothesis has

been that endochondral bone is an osteichthyan apomorphy[3,7,29]. However, recent discoveries

have cast doubt on this assertion. The recognition that dermal bone is secondarily lost in

chondrichthyans[30,31] (Fig. 4) is consonant with prior knowledge of the loss of perichondral

bone in this same lineage[32]. Taken together, this has revived uncertainty about the true

phylogenetic timing of the origin of endochondral ossification[33]. *Minjinia* provides direct

corroboration for a more ancient origin.

Minjinia does not represent the first report of endochondral bone outside of

Osteichthyes. However, it is by far the most extensive and unequivocal example and raises

explicit questions in light of the proximity of *Minjinia* to the gnathostome crown (Fig. 4;

Extended Data Figs. 3, 4). Isolated examples of trabecular endoskeletal bone have historically

been reported in boreaspid osteostracans[20,34], a rhenanid[35], arthrodires[36], a ptyctodont[37], and a

petalichthyid[38,39]. However, these reports are nearly all unillustrated statements; they have all

been considered tenuous[3] or dismissed as misidentifications[5]. In line with these assessments,

we found no evidence of endochondral bone in material of *Buchanosteus* held in the Natural

History Museum, London, or indeed in any other 'placoderms' we have examined. The

Epipetalichthys holotype (Museum für Naturkunde, Berlin specimen MB.f.132.1-3) shows an

apparently spongiose infilling in the anterior region of the braincase, but the identity of this

structure, or even whether it is biological, cannot be determined. The *Epipetalichthys* tissue

figured by Stensiö[38] was very superficial, and possibly represents the retreat of perichondral

bone deposited during cartilage growth[39]. Most recently, trabeculae in supposed endoskeletal

pelvic bones of *Bothriolepis* have been termed endochondral bone[21], although the small scale

of these is in line with 'superficial' perichondral trabeculae seen elsewhere[38]. The reported

examples in boreaspid osteostracans have also been dismissed by later authors[3,5]. Although

they warrant further study, their tissue structures are unlikely to be homologous to

osteichthyans owing to their phylogenetic remoteness and nested position in the

Osteostraci[40].

Among chondrichthyans, endochondral bone has been mentioned in 'acanthodians'[3,41]

and superficial bone-like tissues have been reported in the skeletons of extant

chondrichthyans. We are unable to substantiate statements about acanthodians: no authors

have cited primary sources or specimens. One possible source is Watson's[42] description of

"massive ossification" of the endoskeleton of *Diplacanthus*. However, our synchrotron data

of this same specimen (Fig. 2) shows that this tissue is undoubtably calcified cartilage. Some

authors have speculated that the superficial mineralised tissue in the jaws of acanthodians or

chondrichthyans may have developed in an endochondral position[39]. Histological studies

show that endoskeletal mineralization in the jaws of acanthodians is globular calcification

and occasionally 'sub-tessellate'[8,43]. Recent comparative histology and development in extant

chondrichthyans has shown the presence of an extensive canalicular network in the tesserae[44]

and a trabecular tesseral network in some vertebral elements[45], both resembling bone.

Whether these represent homologues of osteichthyan examples remains open to debate;

future works could employ synchrotron microtomography of stem-chondrichthyan cartilages

to address these questions.

Does endochondral bone have a deep origin within the gnathostome stem group? This

would imply repeated losses of this tissue. We do find statistical support for this hypothesis

(Fig. 4, Table 1, Extended Data Figs. 5, 6, Supplementary Table 1), and the model is well

justified on prior phylogenetic and biological grounds. Endochondral bone has long been

known to be inconsistently developed across 'primitive' bony fishes: incomplete,

polymorphic, or entirely absent ossification of the endoskeleton is known in both Palaeozoic

actinopterygians[41,46,47] and sarcopterygians{Cloutier:wm}, as well as more recent taxa[48]. The

frequent absence of endochondral bone in osteichthyans is considered secondary, and other

controlling factors such as body size, maturity, mechanical stress, and buoyancy can

determine its degree of development[1]. Our findings are also in agreement with studies

establishing a genetic basis for secondary loss of all bone types within chondrichthyans[49-51],

with the failure to produce endochondral bone likely representing arrested development of

chondrocytes as opposed to a primary lack of ability[52].

Another confounding factor in this question is the problem of 'placoderm'

relationships. Although currently resolved in most analyses as a deeply pectinate grade along

the gnathostome stem (Fig. 4), the backbone of this arrangement has poor statistical support,

even in the present analysis (Extended Data Figs. 3). There is a lack of consistency in the

arrangement of plesia and Bayesian tip-dating methods have even recovered a monophyletic

Placodermi[53]. *Minjinia* itself highlights this uncertainty, given its highly unexpected character

combinations. Notwithstanding its endochondral bone and crown-gnathostome-like inner ear

structure, it resembles 'acanthothoracids'—the 'placoderms' widely considered among the

most removed from the gnathostome crown (i.e. most 'primitive'): it possesses deep epaxial

fossae either side of a prominent occipital ridge and a nuchal eminence otherwise seen only

in acanthothoracids such as *Romundina*[18] and *Arabosteus*[19]. This apparent character conflict

could perhaps be more easily reconciled with a more coherent (though not necessarily

monophyletic) 'placoderm' assemblage. Indeed, the highly pectinate structure of the

'placoderm' grade seems symptomatic of an overemphasis on characters and taxa resembling

the crown group, thereby undersampling characters that could stabilise a clear picture of

'placoderm' interrelationships.

Minjinia turgenensis reveals new data on 'placoderm' endoskeleton and tissue

diversity recorded from Mongolia—an otherwise extremely poorly known biogeographic

realm for early gnathostomes. The phylogenetic placement of this 'acanthothoracid'-like

taxon crownward of all non-maxillate 'placoderms', in conjunction with possession of

extensive endochondral bone, highlights the importance of material from traditionally

undersampled geographic areas. The presence of endochondral bone renews the hypothesis

that this tissue is evolutionarily ancient and was lost secondarily in chondrichthyans[6,33]. This

view is overall consistent with evidence of generalised bone loss in chondrichthyans,

potentially as a result of the suppression of bone-generating molecular genetic pathways[51,52].

Continued work in Mongolia and re-evaluation of phylogenetic datasets will be necessary to

address this, with the results likely to lead to substantial re-evaluation of gnathostome

phylogeny.

Methods

X-ray computed microtomography. We scanned MPC-FH100/9.1 using the Nikon XT

225s at the Museum of Paleontology, University of Michigan with the following parameters:

200kV, 140μA, over 3123 projections and a voxel size of 32.92μm. We conducted

segmentation using Mimics 19.0 (http://biomedical.materialise.com/mimics; Materialise,

Leuven, Belgium) and we imaged models for publication using Blender

(https://www.blender.org).

Synchrotron light propagation phase contrast tomography. We imaged *Diplacanthus*

crassismus specimen NMS 1891.92.334 on Beamline 19 of the European Synchrotron

Radiation Facility, using propagation phase-contrast synchrotron microtomography. We

performed a spot scan with an energy of 116keV, achieving a voxel size of 0.55 μm. We

processed the resulting tomograms using VG StudioMax 2.2 (Volume Graphics, Germany),

and prepared images in Blender.

Phylogenetic analysis. We conducted a parsimony analysis using TNT 1.5[54] and Bayesian

analysis using MrBayes v 3.2.7[55]. The dataset consisted of 95 taxa and 284 discrete

characters based on a pre-existing dataset[56]. We employed Osteostraci and Galeaspida as

composite outgroups. We conducted parsimony analysis using both equal weights and

implied weights methods. Global settings were 1000 search replicates and a hold of up to 1

million trees. Equal weights parsimony analyses were conducted using the ratchet with

default settings. Implied weights parsimony used a concavity parameter of 3 and the search

was without the ratchet. Command lists are included in Supplementary Information. We

conducted Bayesian analysis using both a partitioned and unpartitioned dataset. We used the

Mkv model[57] and gamma rate distribution. We ran the analyses for 5 million generations with

a relative burn-in fraction of 0.25. Runs were checked for convergence using Tracer[58]. We

partitioned the dataset using a newly proposed method[28] that partitions the data according to

homoplasy levels. Using the results of implied weights parsimony conducted in TNT, we

created a text table of character fit values. We wrote an R[59] script to generate a list of

partition commands for MrBayes.

We assessed parsimony ancestral states visually using Mesquite[60]. Likelihood and

Bayesian ancestral states were estimated in R using the castor package[61] version 1.5.7. Prior

to calculating likelihood ancestral states on parsimony trees, we scaled branch lengths using

PAUP*[62] and calculated the likelihood scores for all of the trees under the Mkv model with

gamma rate parameter. The trees were then exported with branch lengths. To account for

overall uncertainty in tree estimates, we estimated ancestral states on 100 trees randomly

selected from the fundamental set of most parsimonious trees and two times 50 trees selected

from the 75% last trees of each posterior tree distribution from the Bayesian analysis. We

then run an ancestral states estimation Mk model (using the castor R package) using both the

Equal Rates (ER) and All Rates Different (ARD) models. This resulted in 400 ancestral states

estimations. For each estimation we extracted the overlap log likelihood, the AIC (counting

one parameter for the ER model and two for the ARD model) and the scaled log likelihood

(probability) for the presence and absence of the endochondral bone character (character 4)

for the last common node of *Minjinia* and crown-group gnathostomes. We present the median

value of these distributions of the estimations overall log likelihoods, AICs and presence or

absence of endochondral bone in Table 1.

Data availability

The holotype specimen of *Minjinia turgenensis* will be permanently deposited in the

collections of the Institute of Paleontology, Mongolian Academy of Sciences. Original

tomograms are available at (doi:10.6084/m9.figshare.12301229) and rendered models are

available at (doi:10.6084/m9.figshare.12301223). The phylogenetic character list and dataset

are available as Supplementary Information S1 and S2. The LifeScience Identifier for

Minjinia turgenensis is urn:lsid:zoobank.org:act:82A1CEEC-B990-47FF-927A-

D2F0B59AEA87

Code availability

R code for generating partitions based on character fits and code for likelihood ancestral

states reconstructions and plots are available in the Supplementary Information.

References

1.   Hall, B. K. *Bones and Cartilage*. (Academic Press, 2005).

2.   Janvier, P. *Early Vertebrates*. (Oxford University Press, 1996).

3.   Donoghue, P. C. J., Sansom, I. & Downs, J. P. Early evolution of vertebrate skeletal

tissues and cellular interactions, and the canalization of skeletal development. *Journal

of Experimental Zoology* **306B,** 278–294 (2006).

4.   Rosen, D. E., Forey, P. L., Gardiner, B. G. & Patterson, C. Lungfishes, tetrapods,

paleontology, and plesiomorphy. *Bull Am Mus Nat Hist* **167,** 159–276 (1981).

5.   Gardiner, B. G. The relationships of the palaeoniscid fishes, a review based on new

specimens of *Mimia* and *Moythomasia* from Upper Devonian of Western Australia.

Bull Br Mus nat Hist (Geol) **37,** 173–428 (1984).

6.   Maisey, J. G. Heads and tails: a chordate phylogeny. *Cladistics* **2,** 201–256 (1986).

7.   Friedman, M. & Brazeau, M. A Reappraisal of the origin and basal radiation of the

Osteichthyes. *J Vertebr Paleontol* **30,** 36–56 (2010).

8.   Maisey, J. G., Denton, J. S. S., Burrow, C. & Pradel, A. Architectural and

ultrastructural features of tessellated calcified cartilage in modern and extinct

chondrichthyan fishes. *Journal of Fish Biology* **15,** 37–23 (2020).

9.   Karatajūtė-Talimaa, V. & Novitskaya, L. *Sodolepis*—a new representative of

Mongolepidida (Chondrichthyes?) from the lower Silurian of Mongolia.

Paleontologicheskii Zhurnal **5,** 96–103 (1997).

10.   Karatajūtė-Talimaa, V. & Novitskaya, L. *Teslepis*—a new representative of

mongolepid elasmobranchs from the Lower Silurian of Mongolia. *Paleontologicheskii*

Zhurnal **4,** 36–43 (1992).

11.   Karatajūtė-Talimaa, V., Novitskaya, L. & Rozman, K. S. *Mongolepis*—a new lower

Silurian genus of elasmobranchs from Mongolia. *Paleontologicheskii Zhurnal* **1,** 76–

86 (1990).

12.   Andreev, P. *et al.* The systematics of the Mongolepidida (Chondrichthyes) and the

Ordovician origins of the clade. *PeerJ* **4,** e1850–e1850 (2016).

13.   Vorobjeva, E. I. & Janvier, P. Observations to the note by E Vorobjeva: 'First

discovery of Palaeozoic fishes on the territory of the Mongolian Republic'.

Palaeontology and Biostratigraphy of Mongolia, Moscow, 3, 134-136 (in Russian).

Geobios **17,** 379–380 (1984).

14.   Gegenbaur, C. *Grundriss der vergleichenden Anatomie*. 1–680 (Wilhelm Engelmann,

400   1874).

401   15.   Alekseeva, R. E., Mendbayar, B. & Erlanger, O. A. *Brachiopods and Biostratigraphy*

402   *of the Lower Devonian of Mongolia*. **16,** (Nauka, 1981).

403   16.   Alekseeva, R. E. *Devonian Biostratigraphy of Mongolia*. (Nauka, 1993).

404   17.   Mark-Kurik, E. in *Morphology, Phylogeny and Paleobiogeography of Fossil Fishes*

405   (eds. Elliott, D. K., Maisey, J., Yu, X. & Miao, D.) 101–106 (2010).

406   18.   Ørvig, T. Description, with special reference to the dermal skeleton, of a new radotinid

407   arthrodire from the Gedinnian of Arctic Canada. *Colloque international C.N.R.S. no.*

408   *218* 43–71 (1975).

409   19.   Olive, S., Goujet, D., Lelièvre, H. & Janjou, D. A new Placoderm fish

410   (Acanthothoraci) from the Early Devonian Jauf Formation (Saudi Arabia).

411   *Geodiversitas* **33,** 393–409 (2011).

412   20.   Wängsjö, G. The Downtonian and Devonian vertebrates of Spitsbergen. IX. *Norsk*

413   *Polarinstitutt Skrifter* **97,** 1–611 (1952).

414   21.   Charest, F., Johanson, Z. & Cloutier, R. Loss in the making: absence of pelvic fins and

415   presence of paedomorphic pelvic girdles in a Late Devonian antiarch placoderm

416   (jawed stem-gnathostome). *Biology Letters* **14,** 20180199 (2018).

417   22.   Dupret, V., Sanchez, S., Goujet, D. & Ahlberg, P. E. The internal cranial anatomy of

418   *Romundina stellina* Ørvig, 1975 (Vertebrata, Placodermi, Acanthothoraci) and the

419   origin of jawed vertebrates—Anatomical atlas of a primitive gnathostome. *PLoS ONE*

420   **12,** e0171241 (2017).

421   23.   Stensiö, E. A. La cavité labyrinthique, l'ossification sclérotique et l'orbite de *Jagorina*.

422   *Colloques internationaux du Centre national de la Recherche scientifique* **21,** 9–43

423   (1950).

24.   Stensiö, E. A. Anatomical studies on the arthrodiran head. *K Sven Vetenskapsakad*

Handl **9,** 1–419 (1963).

25.   Goujet, D. *Les poissons placodermes du Spitsberg*. (Cahiers de Paléontologie, Editions

du CNRS, 1984).

26.   Davis, S. P., Finarelli, J. A. & Coates, M. I. *Acanthodes* and shark-like conditions in

the last common ancestor of modern gnathostomes. *Nature* **486,** 247–250 (2012).

27.   Castiello, M. & Brazeau, M. D. Neurocranial anatomy of the petalichthyid placoderm

Shearsbyaspis oepiki Young revealed by X ray computed microtomography.

Palaeontology **61,** 369–389 (2018).

28.   Rosa, B. B., Melo, G. A. R. & Barbeitos, M. S. Homoplasy-based partitioning

outperforms alternatives in Bayesian analysis of discrete morphological data.

Systematic Biology **54,** 373 (2019).

29.   Brazeau, M. D. & Friedman, M. The characters of Palaeozoic jawed vertebrates. *Zool*

J Linn Soc **170,** 779–821 (2014).

30.   Zhu, M. *et al.* A Silurian placoderm with osteichthyan-like marginal jaw bones. *Nature*

502, 188–193 (2013).

31.   Giles, S., Friedman, M. & Brazeau, M. D. Osteichthyan-like cranial conditions in an

Early Devonian stem gnathostome. *Nature* **520,** 82–85 (2015).

32.   Miles, R. S. in *Interrelationships of Fishes* (eds. Greenwood, P. H., Miles, R. S. &

Patterson, C.) 63–103 (Academic Press London, 1973).

33.   Zhu, M. Bone gain and loss: insights from genomes and fossils. *Natl Sci Rev* **1,** 490–

492 (2014).

34.   Janvier, P. *Les céphalaspides du Spitsberg*. (Éditions du Centre National de la

Recherche Scientifique, 1985).

35.   Ørvig, T. Histologic studies of placoderms and fossil elasmobranchs. *Arkiv för Zoologi*

2, 321–454 (1951).

36.   Young, G. C. New information on the structure and relationships of *Buchanosteus*

(Placodermi: Euarthrodira) from the Early Devonian of New South Wales. *Zool J Linn*

Soc **66,** 309–352 (1979).

37.   Miles, R. S. & Young, G. C. in *Problems in Vertebrate Evolution* (eds. Andrews, S.

M., Miles, R. S. & Walker, A. D.) 123–198 (Academic Press, 1977).

38.   Stensiö, E. A. On the head of the macropetalichthyids. *Field Museum of Natural*

History Publication Geological Series **4,** 87–197 (1925).

39.   Ørvig, T. Notes on some Paleozoic lower vertebrates from Spitsbergen and North

America. *Norsk Geologisk Tidsskrift* **37,** 285–353 (1957).

40.   Sansom, R. Phylogeny, classification and character polarity of the Osteostraci

(Vertebrata). *Journal of Systematic Palaeontology* **7,** 95–115 (2009).

41.   Schaeffer, B. The braincase of the holostean fish *Macrepistius*, with comments on

neurocranial ossification in the Actinopterygii. *American Museum Novitates* **2459,** 1–

34 (1971).

42.   Watson, D. M. S. The acanthodian fishes. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences **228,** 49–146 (1937).

43.   Burrow, C. J., Davidson, R. G., Blaauwen, den, J. L. & Newman, M. J. Revision of

Climatius reticulatus Agassiz, 1844 (Acanthodii, Climatiidae), from the Lower

Devonian of Scotland, based on new histological and morphological data. *J Vertebr*

Paleontol **35,** e913421 (2015).

44.   Dean, M. N., Socha, J. J., Hall, B. K. & Summers, A. P. Canaliculi in the tessellated

skeleton of cartilaginous fishes. *Journal of Applied Ichthyology* **26,** 263–267 (2010).

45.   Atake, O. J., Cooper, D. M. L. & Eames, B. F. Bone-like features in skate suggest a

473   novel elasmobranch synapomorphy and deep homology of trabecular mineralization

474   patterns. *Acta Biomaterialia* **84,** 424–436 (2019).

475   46.   Pearson, D. M. & Westoll, T. S. The Devonian actinopterygian *Cheirolepis* Agassiz.

476   *Transactions of the Royal Society of Edinburgh: Earth Sciences* **70,** 337–399 (1979).

477   47.   Giles, S. *et al.* Endoskeletal structure in *Cheirolepis* (Osteichthyes, Actinopterygii), An

478   early ray   finned fish. *Palaeontology* **58,** 849–870 (2015).

479   48.   Grande, L. & Bemis, W. E. Osteology and Phylogenetic Relationships of Fossil and

480   Recent Paddlefishes (Polyodontidae) with Comments on the Interrelationships of

481   Acipenseriformes. *Memoir (Society of Vertebrate Paleontology)* **11 (Supplement to**

482   **Number 1),** 1–121 (1991).

483   49.   Eames, B. F. *et al.* Skeletogenesis in the swell shark *Cephaloscyllium ventriosum*.

484   *Journal of Anatomy* **210,** 542–554 (2007).

485   50.   Venkatesh, B. *et al.* Elephant shark genome provides unique insights into gnathostome

486   evolution. *Nature* **505,** 174–179 (2014).

487   51.   Ryll, B., Sanchez, S., Haitina, T., Tafforeau, P. & Ahlberg, P. E. The genome of

488   *Callorhinchus* and the fossil record: a new perspective on SCPP gene evolution in

489   gnathostomes. *Evolution & Development* **16,** 123–124 (2014).

490   52.   Marconi, A., Hancock-Ronemus, A. & Gillis, J. A. Adult chondrogenesis and

491   spontaneous cartilage repair in the skate, *Leucoraja erinacea. eLife Sciences* **9,** 2813

492   (2020).

493   53.   King, B., Qiao, T., Lee, M. S. Y., Zhu, M. & Long, J. A. Bayesian morphological

494   clock methods resurrect placoderm monophyly and reveal rapid early evolution in

495   jawed vertebrates. *Systematic Biology* **66,** 499–516 (2016).

496   54.   Goloboff, P. A. & Catalano, S. A. TNT version 1.5, including a full implementation of

497   phylogenetic morphometrics. *Cladistics* **32,** 221–238 (2016).

498   55.   Ronquist, F. & Huelsenbeck, J. P. MrBayes 3: Bayesian phylogenetic inference under

499   mixed models. *Bioinformatics* **19,** 1572–1574 (2003).

500   56.   Clement, A. M. *et al.* Neurocranial anatomy of an enigmatic Early Devonian fish sheds

501   light on early osteichthyan evolution. *eLife Sciences* **7,** e34349 (2018).

502   57.   Lewis, P. O. A likelihood approach to estimating phylogeny from discrete

503   morphological character data. *Systematic Biology* **50,** 913 (2001).

504   58.   Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. Posterior

505   summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67,**

506   901–904 (2018).

507   59.   R Core Team. *R: A language and environment for statistical computing.* (R

508   Foundation for Statistical Computing, 2019).

509   60.   Maddison, D. R. & Maddison, W. P. *Mesquite.* (2019).

510   61.   Louca, S. & Doebeli, M. Efficient comparative phylogenetics on large trees.

511   *Bioinformatics* **34,** 1053–1055 (2017).

62.   Swofford, D. L. *PAUP*.* 4.0a166, (2019).

Acknowledgements. M. Bolortsetseg generously assisted MDB with contacts and field

experience in Mongolia. Fieldwork was supported by National Geographic Society grants

CRE 8769-10 and GEFNE35-12 to MDB. AJ's field contributions were supported by funds

from the Anna Maria Lundin's stipend from Smålands Nation, Uppsala University. RS's field

contributions were supported by a Royal Society Research Grant and the University of

Manchester. The majority of this work was supported by the European Research Council

(ERC) under the European Union's Seventh Framework Programme (FP/2007-2013)/ERC

521 Grant Agreement number 311092 to MDB. RPD was also supported by the Île-de-France

522 DIM (domaine d'intérêt majeur) matériaux anciens et patrimoniaux grant PHARE. Stig

523 Walsh is thanked for access and loan of specimen at the National Museums of Scotland.

524 Synchrotron tomography was performed at the ESRF (application LS 2451) with the

525 assistance of Paul Tafforeau. SG was supported by a Royal Society Dorothy Hodgkin

526 Research Fellowship. Matt Friedman is thanked for undertaking the X-ray computed

527 microtomography analysis. This study includes data produced in the CTEES facility at

528 University of Michigan, supported by the Department of Earth & Environmental Sciences

529 and College of Literature, Science, and the Arts. TNT was made available with the support of

530 the Willi Hennig Society.

532 **Author Contributions:** MDB conceived and designed the study. MDB, AJ, YAA, and EZ

533 participated in all field seasons. RPD and AJ undertook preliminary CT scanning and

534 segmentation that revealed the fossil was a 'placoderm' and had endochondral bone. RS

535 discovered the first vertebrate remains in the first field season at Yamaat Gol in 2010. SG

536 undertook the segmentation of *Minjinia* with input from MDB. AJ performed segmentation

537 of *Diplacanthus* tissue. MC provided input on occipital comparative morphology of

538 'placoderms'. RPD provided data and comparative analyses and data for endoskeletal tissue.

539 YAA provided background on the geology, palaeontology, and stratigraphy of the type

540 location; EZ and YAA organized field logistics and permitting. MDB, SG, MC, RPD, and AJ

541 undertook the anatomical interpretation and prepared the figures. MDB and SG conducted the

542 phylogenetic analyses. RS conducted the parsimony branch support analyses. TG wrote the

543 script for generating MrBayes partitions from TNT's character fits table and conducted the

544 likelihood and model-fitting analyses. The manuscript was written by MDB, RPD, and SG.

Competing interests statement:

The authors declare no competing interests.

Fig. 1 | MPC-FH100/9.1 a 'placoderm' skull roof and braincase from the Early

Devonian of Mongolia. a, Ventral view. **b**, Dorsal view. **c**, Left lateral view. **d**, Posterior

view. **e**, Braincase endocavity in dorsal view. Taupe: endoskeleton; grey: mould; blue:

exoskeleton. crsp.ri, craniospinal ridge; e.hy.a, sulcus for the efferent hyoid artery; f.m.ep,

epaxial muscle fossa; fo.mag., foramen magnum; hy.fo, hyodean fossa; l.d.ao, sulcus for the

lateral dorsal aorta; N.V, trigeminal nerve canal; N.VII, facial nerve canal; N.VIII, acoustic

nerve canal; nch, notochordal canal; occ.ri, occipital ridge; orb, orbit; pr.pv, paravagal

process. Scale bar, 10 mm.

Fig. 2 | Endoskeletal mineralisation in fossil gnathostomes. a, Transverse tomographic

slice through MPC-FH100/9.1. **b**, Three-dimensional rendering of trabecular bone structure.

c, Transverse tomographic section through the braincase of the osteichthyan *Ligulalepis*. **d**,

Three-dimensional rendering of the trabecular bone in *Ligulalepis* (**c** and **d** use data from[56]).

e, Synchrotron tomography image of the calcified cartilage of the certatohyal of the stem-

group chondrichthyan *Diplacanthus crassisimus* specimen NMS 1891.92.334. **f**, Semi-

transparent three-dimensional structure of calcified cartilage of NMS 1891.92.334. Scale

bars, **a** and **b,** 10 mm; **c** and **d**, 1 mm; **e** and **f**, 150 µm.

Fig. 3 | Braincase endocavity of *Minjinia*. a, Semi-transparent rendering of skull roof and

braincase (grey and blue) showing extent of endocavity (pink). **b,** Ventral view. **c,** Dorsal

view. a.scc, anterior semicircular canal; cav.end, endolymphatic cavity; d.end, endolymphatic

duct; h.scc, horizontal semicicular canal; N.V, trigeminal nerve canal; N.VIIhm,

hyomandibular branch of facial nerve canal; N.VIIpal, palatine branch of facial nerve canal;

N.VIII, acoustic nerve canal; N.X, vagus nerve canal, $N.X_a$, anterior branch of vagus nerve

canal; $N.X_p$, posterior branch of vagus nerve canal; occ.a, occipital artery canals; p.scc,

posterior semicircular canal; sac, sacculus; soc, spino-occipital nerve canals; sup.opth, canal

for supra-ophtalmic nerve. Scale bars, 10 mm (upper scale bar associates with **a**, lower scale

bar associates with **b** and **c**).

Fig. 4 | Strinct consensus tree from parsimony analysis of early gnathostomes showing

distribution of endochondral bone and exoskeletal armour. Squares at nodes indicate

parsimony reconstruction for endochondral bone. Pie charts at nodes show likelihood

reconstructions for the same character under the all-rates-different model (see Extended Data

Figs 6 & 7 for competing reconstructions). Grey box indicates uncertainty. Loss of

endochondral bone maps closely with generalised loss of bone in chondrichthyans where

exoskeletal armour and perichondral bone are also absent.

Table 1 | Tree distribution (n=100) ancestral states estimation results. ER = Equal rates

model; ARD = All Rates Different model. The columns AIC and log.like represent the

median AIC and log.lik across the 100 parsimony and Bayesian trees (for both models). The

like.ratio column is the likelihood ratio for the models compared on these trees. The columns

Absent and Present represent the median scaled likelihood for the endochondral bone state.

trees	model	log.like.	like.ratio	AIC	node	Absent	Present
Parsimony	ER	-28.91	4.74	59.82	Minjinia:Gnathostomes	0.67	0.33
(equal weights)	ARD	-26.54		57.09		0.19	0.81
	ER				Crown Gnathostomes	0.91	0.09
	ARD					0.46	0.54
Bayesian	ER	-29.66	5.26	61.32	Minjinia:Gnathostomes	0.73	0.27
(unpartitioned)	ARD	-27.03		58.06		0.17	0.83
	ER				Crown Gnathostomes	0.79	0.21
	ARD					0.22	0.78

Extended Data Fig. 1 | Tomograms of endoskeletal ossification in *Minjinia*. Top row:

semi-coronal sections through braincase. Double-headed arrows indicate anterior-posterior

(a-p) dorsal-ventral (d-v) axes. Bottom row: semi-transverse sections through posterior part

of endocranium. Voids of black space represent mouldic preservation. Scale bars, 10 mm and

apply across each row of panels.

Extended Data Fig. 2 | Right orbital wall and innervation pattern of *Minjinia*. a, orbit in

anterolateral view showing disposition of nerve openings (pink infill). **b**, endocast in the

same perspective showing the relationship between nerve canals and endocast. a.scc., anterior

semicircular canal; N.V$_{2,3}$ trunk of the trigeminal nerve canal for branches 2 and 3; N.VIIhm,

hyomandibular branch of facial nerve canal; N.VIIpal, palatine branch of facial nerve canal;

sac., sacculus; sup.opth, canal for supra-ophtalmic nerve. Scale bars, 20 mm (upper scale bar

associates with **a**, lower scale bar associates with **b** and **c**).

Extended Data Fig. 3 | Results of phylogenetic parsimony analysis. Dataset consists of 95

taxa and 284 characters. Both trees are strict consensus topologies. Equal weights parsimony

analysis using the ratchet resulted in 240 trees with a length of 831 steps. Implied weights

parsimony analysis using random addition sequence + branch-swapping resulted in 8 optimal

trees with score 85.20513. Double-digit figures above internal branches are bootstrap values

of 50% and over; single-digit figures below branches are Bremer decay index values. Blue

shading: osteichthyan total group (dark blue: crown group); orange shading: chondrichthyan

total group (dark orange: crown group).

Extended Data Fig. 4 | Results of Bayesian phylogenetic analysis using both partitioned

and unpartitioned data. Majority-rules consensus trees with posterior probabilities shown

along branches. Blue shading: osteichthyan total group (dark blue: crown group); orange

shading: chondrichthyan total group (dark orange: crown group).

Extended Data Fig. 5 | Likelihood ancestral state mapping of endochondral bone on

equal weights parsimony results. a, ARD, all rates different model; **b**, ER, equal rates

model.

Extended Data Fig. 6 | Likelihood ancestral state mapping of endochondral bone on

unpartitioned Bayesian analysis results. a, ARD, all rates different model; **b**, ER, equal

rates model.

Endochondral bone
Exoskeleton

Absent — Tessellated
Present — Armoured

Missing / polymorphism

Galeaspida
Osteostraci
Gemuendina
Jagorina
Parayunnanolepis — Antiarchi
Yunnanolepis
Bothriolepis
Pterichthyodes
Brindabellaspis
Lunaspis
Macropetalichthys
Austroptyctodus
Campbellodus
Rhamphodopsis
Eurycaraspis
Romundina
Kujdanowiaspis — Arthrodira
Cowralepis
Dicksonosteus
Buchanosteus
Coccosteus
Incisoscutum
Minjinia
Entelognathus
Ramirosuarezia
Janusiscus
Dialipina
"Ligulalepis"
Guiyu
Sparalepis
Psarolepis
Achoania
Meemannia
Cheirolepis
Howqualepis
Mimipiscis
Raynerius
Moythomasia — Osteichthyes
Kentuckia
Lawrenciella
Kansasiella
Ptyctolepis
Onychodus
Qingmenodus
Diplocercides
Miguashaia
Gavinia
Styloichthys
Eusthenopteron
Gogonasus
Powichthys
Glyptolepis
Porolepis
Youngolepis
Diabolepis
Dipterus
Uranolophus
Culmacanthus
Tetanopsyrus
Diplacanthus
Gladiobranchus
Ischnacanthus
Poracanthodes
Cassidiceps
Euthacanthus
Mesacanthus
Promesacanthus
Cheiracanthus
Acanthodes
Homalacanthus
Latviacanthus
Brochoadmones
Kathemacanthus
Brachyacanthus
Climatius
Parexus — Chondrichthyes
Ptomacanthus
Vernicomacanthus
Lupopsyrus
Gyracanthides
Obtusacanthus
Pucapampella
Doliodus
Cladodoides
Orthacanthus
Tamiobatis
Akmonistion
Cladoselache
Cobelodus
Tristychius
Hamiltonichthys
Onychoselache
Helodus
Chondrenchelys
Debeerius

"Placodermi"

Crown-group Gnathostomata

Parsimony: equal weights

- Galeaspida
- Osteostraci
- *Gemuendina* — 70, 3
- *Jagorina*
- *Parayunnanolepis* — 73
- *Yunnanolepis*
- *Bothriolepis* — 83, 3
- *Pterichthyodes*
- *Brindabellaspis*
- *Lunaspis*
- *Macropetalichthys*
- *Austroptyctodus* — 62, 3
- *Campbellodus*
- *Rhamphodopsis*
- *Eurycaraspis*
- *Romundina*
- *Kujdanowiaspis* — 2
- *Cowralepis*
- *Dicksonosteus*
- *Buchanosteus* — 57
- *Coccosteus* — 79, 3
- *Incisoscutum*
- *Minjinia*
- *Entelognathus*
- *Ramirosuarezia*
- *Janu. siscus*
- *Dialipina* — 56, 2
- *"Ligulalepis"*
- *Guiyu*
- *Sparalepis*
- *Psarolepis*
- *Achoania*
- *Meemannia* — 66, 3
- *Cheirolepis*
- *Howqualepis*
- *Mimipiscis* — 82, 2, 58
- *Raynerius*
- *Moythomasia*
- *Kentuckia* — 66, 67
- *Lawrenciella*
- *Kansasiella*
- *Ptyctolepis*
- *Onychodus* — 56, 2
- *Qingmenodus*
- *Diplocercides* — 76
- *Miguashaia*
- *Gavinia*
- *Styloichthys*
- *Eusthenopteron* — 96, 6
- *Gogonasus*
- *Powichthys* — 54, 2, 74, 3
- *Glyptolepis*
- *Porolepis*
- *Youngolepis*
- *Diabolepis* — 84, 4, 94
- *Dipterus* — 5
- *Uranolophus*
- *Culmacanthus*
- *Tetanopsyrus*
- *Diplacanthus* — 56, 2
- *Gladiobranchus*
- *Ischnacanthus* — 68
- *Poracanthodes*
- *Cassidiceps*
- *Euthacanthus*
- *Mesacanthus*
- *Promesacanthus*
- *Cheiracanthus* — 74, 65
- *Acanthodes*
- *Homalacanthus*
- *Latviacanthus*
- *Brochoadmones*
- *Kathemacanthus*
- *Brachyacanthus*
- *Climatius*
- *Parexus* — 2
- *Ptomacanthus*
- *Vernicomacanthus*
- *Lupopsyrus*
- *Gyracanthides* — 2
- *Obtusacanthus*
- *Pucapampella*
- *Doliodus*
- *Cladodoides*
- *Orthacanthus*
- *Tamiobatis* — 2
- *Akmonistion*
- *Cladoselache*
- *Cobelodus*
- *Tristychius*
- *Hamiltonichthys* — 61
- *Onychoselache*
- *Helodus*
- *Chondrenchelys* — 83, 2
- *Debeerius*

Parsimony: implied weights

- Galeaspida
- Osteostraci
- *Parayunnanolepis*
- *Yunnanolepis*
- *Bothriolepis*
- *Pterichthyodes*
- *Brindabellaspis*
- *Romundina*
- *Eurycaraspis*
- *Lunaspis*
- *Macropetalichthys*
- *Austroptyctodus*
- *Campbellodus*
- *Rhamphodopsis*
- *Gemuendina*
- *Jagorina*
- *Cowralepis*
- *Kujdanowiaspis*
- *Dicksonosteus*
- *Buchanosteus*
- *Coccosteus*
- *Incisoscutum*
- *Minjinia*
- *Diplacanthus*
- *Janusiscus*
- *Ramirosuarezia*
- *Tetanopsyrus*
- *Entelognathus*
- *Dialipina*
- *"Ligulalepis"*
- *Meemannia*
- *Cheirolepis*
- *Kentuckia*
- *Lawrenciella*
- *Kansasiella*
- *Howqualepis*
- *Moythomasia*
- *Mimipiscis*
- *Raynerius*
- *Guiyu*
- *Sparalepis*
- *Psarolepis*
- *Achoania*
- *Ptyctolepis*
- *Onychodus*
- *Qingmenodus*
- *Gavinia*
- *Diplocercides*
- *Miguashaia*
- *Styloichthys*
- *Eusthenopteron*
- *Gogonasus*
- *Powichthys*
- *Glyptolepis*
- *Porolepis*
- *Youngolepis*
- *Diabolepis*
- *Dipterus*
- *Uranolophus*
- *Cassidiceps*
- *Mesacanthus*
- *Promesacanthus*
- *Cheiracanthus*
- *Acanthodes*
- *Homalacanthus*
- *Euthacanthus*
- *Ischnacanthus*
- *Poracanthodes*
- *Latviacanthus*
- *Gladiobranchus*
- *Culmacanthus*
- *Kathemacanthus*
- *Brochoadmones*
- *Ptomacanthus*
- *Climatius*
- *Vernicomacanthus*
- *Brachyacanthus*
- *Parexus*
- *Gyracanthides*
- *Lupopsyrus*
- *Obtusacanthus*
- *Pucapampella*
- *Doliodus*
- *Cladodoides*
- *Tristychius*
- *Hamiltonichthys*
- *Onychoselache*
- *Cladoselache*
- *Akmonistion*
- *Cobelodus*
- *Orthacanthus*
- *Tamiobatis*
- *Helodus*
- *Chondrenchelys*
- *Debeerius*

Bayesian: unpartitioned

Bayesian: partitioned by character fits