1   <u>Title:</u> Extracellular DNA traps in a ctenophore demonstrate conserved immune cell behaviors in
2   a non-bilaterian
3

4   <u>Authors and Affiliations:</u> Lauren E. Vandepas[a,b,c,e*†], Caroline Stefani[c*], Phillip P. Domeier[c], Nikki
5   Traylor-Knowles[d], Frederick W. Goetz[b], William E. Browne[e], Adam Lacy-Hulbert[c]
6
7   [a]NRC Research Associateship Program; [b]Northwest Fisheries Science Center, National
8   Oceanographic and Atmospheric Administration, Seattle, WA 98112; [c]Benaroya Research
9   Institute at Virginia Mason, Seattle, WA 98101; [d]University of Miami Rosenstiel School of Marine
10 and Atmospheric Sciences, Miami, FL 33149; [e]University of Miami Department of Biology, Coral
11 Gables, FL 33146; [†]Corresponding author; [*]Equal contribution

17
18   **Abstract**
19

The formation of extracellular DNA traps (ETosis) is a first response mechanism by specific
immune cells following exposure to microbes [1,2]. Initially characterized in vertebrate neutrophils,
cells capable of ETosis have been discovered recently in diverse non-vertebrate taxa [3] [4-6]. To
assess the conservation of ETosis between evolutionarily distant non-vertebrate phyla, we
observed and quantified ETosis using the model ctenophore *Mnemiopsis leidyi* and the oyster
Crassostrea gigas. Here we report that ctenophores – thought to have diverged very early from
the metazoan stem lineage [7-10]– possess immune cell types capable of phagocytosis and ETosis.
We demonstrate that both *Mnemiopsis* and *Crassostrea* immune cells undergo ETosis after
exposure to diverse microbes and chemical agents that stimulate ion flux. Our results support
ETosis as an evolutionarily ancient metazoan defense against pathogens.

Main

Response to invading microbes is a primary physiological task for all metazoans. In vertebrates, specialized immune cell types such as macrophages and neutrophils detect the presence of microbes. These immune cells perform specific behaviors like phagocytosis or secretion of antimicrobial compounds to sequester and eliminate microbial invaders [11,12]. The release of extracellular DNA traps (ETs) is a relatively recently described immune cell behavior – a morphologically and molecularly distinct cell death process called ETosis – during which immune cells cast filamentous nets composed of nuclear chromatin material or mitochondrial DNA into the surrounding extracellular space, trapping and killing invading microbes [13,14]. Initially believed to be a behavior exclusive to vertebrate neutrophils, recent studies have highlighted ETosis as an anti-microbial response in non-vertebrate taxa [4,5,15-18]. The independent evolution of ETs has also been proposed in taxa as divergent as social amoebas [19] and plants [20].

In mammalian cells, ETosis and other immune responses are often initiated by pattern recognition receptors (PRRs) that bind to molecular motifs on microbes commonly referred to as pathogen-associated molecular patterns (PAMPs) [21]. PRR protein sequences and domain architectures can vary between taxa, as they evolve to detect host specific pathogens [22-24]. For example, mussels have a radiation of TLRs that may recognize a suite of specific pathogens [25], while *Drosophila* uses its Toll receptors to identify the protein Spätzle [26]. Despite these broad patterns of clade-specific immune receptor diversity, the signaling cascades and effectors downstream of PRRs have been shown to be well conserved [30], including intracellular signaling pathways involving calcium, MAP kinases, and reactive oxygen species (ROS) [27-29].

In vertebrate leukocytes, exposure to cytokines, microbes, PAMPs, or pharmacological agents activate specific cellular pathways involved in ET formation, such as a response to ion imbalance (calcium, potassium) or reactive oxygen species (ROS) burst [31,32]. Although the detailed mechanisms of these pathways remain unknown in non-vertebrates [5,6,14,31,33], genes involved in vertebrate immune cell signaling cascades have been identified in diverse metazoans, including non-bilaterians [34-38].

Non-vertebrate ETosis has been best studied in molluscan model systems, particularly in bivalves like the model oyster *Crassostrea gigas* [4-6,16,39,40]. However, attempts to stimulate production of ETs in *Crassostrea* via exposure to some microbial signatures as well as ETosis-stimulating drugs has yielded variable results [6]. For example, robust ETotic response in oyster hemocytes was observed following challenge with *Vibrio* [4], however induction of ETosis with other microbes or PAMPs has been either modest or not observed [6]. Without a consensus assessment of ETosis stimulation, the conservation of ET induction in non-vertebrates remains unclear, and it is unknown whether ETosis represents a fundamental metazoan immune response [4-6].

Ctenophores, also known as "comb jellies", are a clade of gelatinous planktonic marine animals that diverged early from the animal stem lineage and represent one of the most ancient extant metazoan phyla [8-10]. Ctenophores have two distinct germ layers (ectoderm and endomesoderm) separated by a jelly-like layer of collagenous mesoglea and lack a circulatory system. Understanding functional attributes of ctenophore physiological systems has provided fundamental insights into the conservation of animal cell types and signaling pathways, as well as revealing mechanisms for the emergence of evolutionary novelties [36,41-43]. Currently, the ctenophore immune system remains almost entirely undescribed [36,44]. Specific immune cell types have not been explicitly identified in ctenophores, though they possess motile amoebocyte-like cells that are abundant in the mesoglea and are capable of phagocytosis [36,45]. Whether ctenophores have additional specialized cellular immune mechanisms has not been reported.

Here we demonstrate that the model ctenophore *Mnemiopsis leidyi* possesses immune cell types capable of ETosis in response to diverse stimuli known to activate conserved signaling cascades in vertebrate immune cells. To rapidly and accurately quantify ETosis we developed a novel, automated imaging pipeline to identify ETs. We applied this imaging pipeline to

82   comparatively assess ET production in *Mnemiopsis* and the bivalve mollusk *Crassostrea gigas,*
83   a model for invertebrate ETosis. Our data from both non-vertebrate species suggest that cells
84   undergoing ETosis following a range of stimuli represent evolutionarily ancient anti-microbial
85   defense mechanisms in metazoans.
86
87   ## Results
88
Isolated ctenophore cells display immune behaviors
To investigate whether the model ctenophore *Mnemiopsis leidyi* possesses cell types with
specialized immune functions, we mechanically disassociated whole *Mnemiopsis* and examined
isolated cells [46,47]. We observed morphologically distinct motile cell types displaying prominent
intracellular granules and/or vesicles, including amoebocyte-like cells with relatively short
pseudopodia and stellate cells with long pseudopodia (**Fig. 1A-C, Supp. Video 1**). Time-lapse
microscopy shows that some stellate cells have a dynamic morphology and can rapidly vary the
length and number of pseudopodia (**Supp. Video 2**). These motile cells are highly active in
primary cell cultures and their morphologies and scavenging behaviors are reminiscent of immune
cell types, such as macrophage-like cells, that have been described in diverse metazoans [11].
A major role for macrophage-like cells is host defense against microbes. We therefore
assessed whether the motile cells present in *Mnemiopsis* exhibited fundamental immune cell
behaviors in response to bacteria. *Mnemiopsis* cells were incubated with killed fluorescently
labeled *Escherichia coli* and imaged by combined DIC and fluorescence microscopy. We found
Mnemiopsis cells exhibiting diverse morphologies internalized fluorescent *E. coli* [36]; these
included motile stellate and amoeboid cells (**Fig. 1D-F**). Furthermore, timelapse DIC microscopy
revealed that amoeboid cells actively maneuvered their pseudopodia to engulf nearby bacteria,
supporting a potential role in sensing and removing microbes (**Supp. Video 3**).
Following exposure to *E. coli* a conspicuous subset of stellate cells changed their
morphology dramatically by retracting their pseudopodial processes and undergoing nuclear
rotation, followed by a rapid expulsion of cellular material (**Fig. 1G; Supp. Video 4; Supp. Video
5**). This behavior is remarkably similar to vertebrate monocyte behavior preceding extracellular
DNA trap (ET) formation [31,48]. This led us to speculate that some ctenophore immune cells were
producing ETs in response to the presence of microbes.
113
Ctenophore immune cells produce extracellular traps composed of chromatin
In vertebrates, ETs are released in response to recognition of microbial components, and
protect against infection by immobilizing and killing invading microbes[1]. To test whether incubation
with microbes promoted extrusion of DNA from *Mnemiopsis* cells we stained isolated cells with
Hoechst to label DNA. In isolated cells incubated in seawater media, we observed intact nuclei
with concentrated Hoechst labeling (**Fig. 2A**). After treatment with heat-killed fluorescent *E. coli*
we observed networks of Hoechst-stained DNA filaments cast in large areas around some
individual *Mnemiopsis* cell bodies. Three-dimensional rendering of confocal z-stacks showed that
Mnemiopsis cells extruding DNA had bacteria entangled in the DNA networks, consistent with
ETs (**Fig. 2B; Supp. Video 6;** [49]).
In vertebrate immune cells, ETs can originate either from the cell nucleus or from
mitochondria [50,51]. To assess the organellar origin of the extruded DNA, we stained *Mnemiopsis*
cells with an antibody that recognizes an array of nuclear histone proteins (H1, H2A, H2B, H3,
H4). Nuclei of non-ETotic *Mnemiopsis* cells were labeled with this histone antibody, which showed
stereotypical patterns of concentrated DNA and histone labeling, confirming the specificity of this
antibody in *Mnemiopsis* cells (**Fig. 2C**) [52,53]. Notably, in cells that have undergone ETosis, the
extracellular ET DNA also stained with this pan-histone antibody, demonstrating that the extruded
DNA filaments are composed of chromatin (**Fig. 2C, arrows**). Together, these data identify that

Mnemiopsis cells produce *bona fide* extracellular DNA traps and further concluded that these ETs
are derived from the nucleus [14].
134
Ctenophore ETosis is an immune response to microbial exposure
To assess whether the formation of ETs in *Mnemiopsis* is a response to microbial
challenge, we measured ET production after incubation of *Mnemiopsis* cells with *E. coli* and two
additional microbial particles: heat-killed gram-positive *S. aureus*, and the yeast cell wall extract
zymosan. We found that all three microbial particles stimulated ET formation in *Mnemiopsis* cells
when compared with untreated cells incubated with media alone (**Fig. 2D**). Numbers of ETotic
cells were manually quantified based on release of Hoechst-positive material by fluorescent
microscopy (**Fig. 2E**). Exposure to *S. aureus* and zymosan promoted considerably higher rates
of ET formation (mean increase of 98.2% and 117.7% over control respectively) than *E. coli*
(43.7%) (**Fig. 2E**). These data confirm that production of ETs represents a *Mnemiopsis* immune
cell-specific response to the presence of microbes.
146
Ctenophore ETs form filamentous networks that capture microbes
Scanning electron microscopy (SEM) is commonly used to confirm the structure of ETs,
and is informative in distinguishing ETosis from DNA release secondary to other forms of cell
death such as necrosis or apoptosis [54]. To examine the structure of the extruded DNA in
Mnemiopsis cell cultures, we performed SEM on isolated cells that had been incubated with *S.*
aureus or with seawater media alone. Images of untreated *Mnemiopsis* cells confirm the presence
of diverse morphological cell types observed by DIC, including stellate cells (**Fig. 3A; Supp. Fig.**
1). Following incubation with *S. aureus*, we observed long filamentous networks surrounding
some *Mnemiopsis* cells (**Fig. 3B-C; Supp. Fig. 1**), which are similar to ETs produced by
vertebrate and crustacean immune cells [1,5,55,56]. We also observed a close association between
individual *S. aureus* and the filamentous nets produced by *Mnemiopsis* cells, providing further
evidence that these structures are able to capture microbes, a critical function of ETs (**Fig. 3B-C**
insets) [57].
160
PMA induces ETosis in *Mnemiopsis*
We hypothesized that ETosis in ctenophores may be initiated by engaging evolutionarily
conserved signaling pathways known to precede ETosis in other metazoans. For example,
phorbol 12-myristate 13-acetate (PMA) is commonly used to induce ET production in vertebrate
immune cells through activation of NADPH-mediated ROS [31]. This pathway has been proposed
to induce ET formation in immune cells in annelids and crustaceans [5,15], suggesting that it may
be evolutionarily ancient and well conserved in metazoans. We treated *Mnemiopsis* cells with
PMA for four hours, stained DNA with Hoechst, and visualized cells by fluorescence microscopy.
We observed the extrusion of DNA after PMA treatment, which was similar ot that seen after
micobial challenge (**Fig. 2D; Fig. 4A**).
Importantly, exposure to PMA can also trigger non-ETotic cell death pathways [58], raising
the possibility that the extracellular DNA we observed was a result of cell rupture and release of
DNA secondary to cell death. To distinguish between these possible scenarios, we used a
combination of two DNA dyes: Hoechst, which is a membrane-permeable dye that labels DNA in
both live cells and dead cells, and SytoxGreen, a non-permeable cell stain that selectively labels
DNA in cells with compromised cell membranes, as observed in cells undergoing cell death (**Fig.**
4A). Both DNA dyes label ETs [5,59], and ETotic cells have both diffuse gradients of Hoechst and
Sytox staining (**Fig. 4A-B**). We then developed a semi-automated imaging analysis approach that
allowed us to measure Hoescht and SytoxGreen labeling in individual cell nuclei and accurately
quantify both ETosis and cell death simultaneously (**Fig. 4B; Supp. Fig. 2**). ET production has
been successfully identified and quantified in neutrophils using a combination of machine learning
and high-throughput imaging [60,61], we deployed similar techniques to assess ETosis in ctenophore

cells. An approach similar to those implemented in flow cytometry analyses was used in which
we plotted the intensity of SytoxGreen fluorescence against Hoechst signal for each individual
cell (**Fig. 4B-C**). This allowed for the identification of three distinct populations in PMA-treated
Mnemiopsis cells: Hoechst^{high}/ SytoxGreen^{low} live cells that show no DNA release and intact cell
membranes; Hoechst^{low}/ SytoxGreen^{low} cells which display diffuse extracellular Hoechst and
SytoxGreen fluorescence, representing ETotic cells [5,59]; and SytoxGreen^{high} cells with a range of
Hoescht staining, which are dead or dying non-ETotic cells.
To confirm that the image analysis pipeline accurately distinguishes non-ETotic cell death
from ETosis, we induced death in *Mnemiopsis* cells by exposure to a moderate osmotic shock
(dilute saline solution, 2X PBS). We measured a significant increase in non-ETotic dead cells
compared with normal seawater media controls (**Fig. 4C**). Notably, dying cells were clearly
separated from cells undergoing ETosis. Using this pipeline, we then quantified ETosis and cell
death in cells treated with PMA. Confirming our initial observations by microscopy, PMA promoted
significant induction of ETosis, with a mean increase of 57.2% over cells incubated with seawater
media alone (**Fig. 4C-D**). Importantly, non-ETotic cell death did not significantly increase following
PMA incubation.
We sought to establish that the automated image analysis pipeline could detect ETosis
induced by microbial exposure. Because we observed clear ETs with SEM following *S. aureus*
challenge, we then applied this automated imaging approach to this microbial stimulus. Samples
that are incubated with microbes contain more background fluorescence due to Hoescht and
Sytox labeling of *S. aureus* nuclear material; however, we observed a clear increase in ETosis
following *S. aureus* challenge using automated imaging analysis, as well as an increase in non-
ETotic cell death (**Fig. 4C and 4D**). Together, these data confirm that PMA induces ETosis in
Mnemiopsis. Further, the novel semi-automated image analysis pipeline presented here can
accurately detect and quantify invertebrate ETosis and distinguish ETosis from total cell death.
208
ETosis in *Mnemiopsis* is initiated by agonists that induce ion flux
In vertebrates, ET formation can be triggered by membrane ion flux triggered by exposure
to bacterial toxins that act as ionophores [31,53,62]. Because stimulation of ETosis via ion flux has
limited reports in non-vertebrates [6,32], we assessed whether well-characterized bacterially-derived
ionophores nigericin (potassium ionophore) and calcimycin (A23187; calcium ionophore) could
also elicit ETosis in *Mnemiopsis* cells. We incubated *Mnemiopsis* cells with the K$^+$ ionophore
nigericin and observed a mean increase of 367% over untreated cells (**Fig. 5A and 5B; Supp.**
Video 6; Supp. Fig. 3). Incubation with the Ca^{2+} ionophore calcimycin resulted in a mean increase
of 225% over untreated control cells (**Fig. 5A and 5B**). Furthermore, although incubation with
either ionophore also significantly increased *Mnemiopsis* cell death, consistent with their effects
in vertebrate cells [63-65], our image analysis pipeline was able to clearly distinguish ETosis from
other forms of cell death. These data therefore support an evolutionarily conserved pathway for
ET formation in response to immune stimulation or cellular damage.
222
Microbes, PMA, and calcium ionophore stimulates ETosis in *Crassostrea gigas* hemocytes
Our observations that ET formation in *Mnemiopsis* can be triggered by similar signaling
pathways to those involved in mouse and human immune cell ETosis supports the hypothesis
that ETosis represents an evolutionarily ancient response to infection or cell damage activated by
intracellular signaling pathways that are widely conserved throughout animals [66-68]. However,
stimulation of ETosis has not been widely assessed outside of vertebrates [6,32] or compared
between phyla. To compare stimulation of *Mnemiopsis* cells with another non-vertebrate
metazoan that is an established model for analysis of ETosis, we extended our analysis to the
bivalve mollusc, *Crassostrea gigas*, a marine non-vertebrate bilaterian with a circulatory system
possessing specialized immune cells (hemocytes). Some hemocyte cell types have been shown
to display immune behaviors such as phagocytosis [69]. We incubated isolated hemocytes with

fluorescent *E. coli* for 4 hours to assess whether *C. gigas* cells were responding to the presence
of bacteria by phagocytosing the *E. coli* (**Fig. 6A-B)** or undergoing ETosis (**Fig. 6B,** inset). When
observing hemocytes by confocal microscopy, we found that some hemocytes are highly
phagocytic, with individual cells ingesting multiple bacteria (**Fig. 6B)**. Fluorescent imaging also
identified networks of extracellular DNA surrounding some hemocytes after exposure to microbial
particles, similar to previous reports of bivalve ET formation [4] (**Fig. 6B, insets**).
While ETs have been reported in multiple bivalve species, successful induction of ETosis
in hemocytes following incubation with either microbial stimuli or pharmaceuticals has varied [4-6].
To directly compare ET production in oyster hemocytes with observations in *Mnemiopsis*, we
assessed ETosis in *C. gigas* hemocytes using the same panel of microbes. We first confirmed
that *Crassostrea* immune cells undergo ETosis in response to microbial stimulation using the
automated imaging analysis pipeline (**Fig. 6C-D**). Hemocytes were incubated with heat-killed *E.*
coli, S. aureus, or zymosan, and then assessed using the automated image analysis pipeline for
quantification of ETosis. We found that ETosis increased after exposure to all three microbes,
when compared to hemocytes incubated *in vitro* with seawater media alone (**Fig 6C-D)**.
Crassostrea hemocytes undergo ETosis in response to bacterial challenge with both gram-
negative *E. coli* and gram-positive *S. aureus* with mean increases of 52.3% and 41.5% over
untreated hemocytes, respectively (**Fig. 6D)**. The fungal PAMP zymosan also elicited a significant
ETotic response in oyster hemocytes, with a mean increase of 112% over seawater media
controls. Interestingly, *S. aureus* stimulation did not stimulate significant increases in non-ETotic
cell death in hemocytes (**Fig. 6D)**, in contrast to what we observed in *Mnemiopsis* cells (**Fig. 2D-**
E). Hemocyte total non-ETotic cell death increased significantly with either *E. coli* or zymosan
exposure.
We also measured ETosis in *Crassostrea* hemocytes after four-hour incubations with
PMA, nigericin, calcimycin, or seawater media controls (**Fig. 6E-F**). Exposure to PMA induced
significant ET production in isolated *Crassostrea* hemocytes, with a relative mean increase of
95.7% (**Fig. 6F**). A large proportion of *Crassostrea* hemocytes also produced ETs after exposure
to calcimycin, with a mean increase of 209% over controls (**Fig. 6F**). These results are similar to
the robust response reported in prior studies of bivalve hemocytes to Ca^{2+} ionophore stimulation
[6]. In contrast to *Mnemiopsis* cells, incubation of hemocytes with the K^+ ionophore nigericin showed
no significant induction of ETosis (compare **Fig. 2D-E and Fig. 6F**). Notably, non-ETotic cell death
does not increase in *C. gigas* hemocytes following drug incubation. Unexpectedly, both PMA and
calcimycin treatment resulted in a relatively minor, but significant, decrease in non-ETotic cell
death.
268
269 **<u>Discussion</u>**
270
Identifying the core molecular mechanisms of immune cells and their evolutionary origins
is critical for understanding the evolution of metazoan immune cell function and behavior. Here
we report that ctenophores, which diverged very early from the metazoan stem lineage, have
immune cells capable of ETosis, a cellular response to microbes first described in mammalinan
neutrophils. We also show that cells isolated from the model ctenophore *Mnemiopsis leidyi*
undergo ETosis following exposure to diverse microbial components, and stimulation of
intracellular signaling cascades known to precede ETosis in vertebrates. For example, we find
that incubation with the potassium ionophore nigericin stimulates ETosis in *Mnemiopsis* cells, as
has been reported for mammalian neutrophils, suggesting that K^+ efflux may be an evolutionarily
conserved trigger for ETosis. We conclude from our results that ETosis represents an ancient
mechanism of cellular immune defense in metazoans (**Fig. 7**).
Since the first description of neutrophil extracellular traps [11], multiple pathways involved in
the generation of mammalian ETs have been identified [31]. However, the intracellular actors
necessary for ET formation in non-vertebrate taxa are less well understood. For example, gene

homologs for proteins known to be essential for ETosis in mammalian neutrophils, such as
peptidylarginine deiminase 4 (PAD4), neutrophil elastase, myeloperoxidase (MPO) and pannexin-
1 are absent outside of vertebrates. There is strong evidence for the presence of multiple ETosis
pathways in non-vertebrates that can be activated via stimulation with microbial signatures,
parasites, PMA, A23187, and UV light [4-6,15-17]. Potassium ion efflux, for example, is classically
associated with cellular damage, including during bacterial infection [70]. Calcium flux has been
implicated in ET formation in vertebrates either as a component of immune receptor-mediated
signaling or as an activator of downstream effectors that drive DNA condensation preceding
ETosis [71]. In addition to intracellular ion flux, components of vertebrate ETosis pathways, such as
ERK, p38, or Akt, are foundational components of metazoan intracellular signaling [66-68].
Our ability to make further direct comparisons of the molecular mechanisms of ETosis
between vertebrates and *Mnemiopsis* is currently hampered by a lack of tractable forward genetic
approaches and is therefore largely restricted to pharmacological perturbation. However, our data
supports considerable apparent conservation of ET formation; firstly, bacterial and fungal
components (*E. coli*, intact *S. aureus* and fungal particles) significantly stimulated ETosis in
Mnemiopsis and *Crassostrea* hemocytes in our study. Our results are in agreement with work in
other non-vertebrates and supports the concept that ET formation is a fundamental component
of metazoan anti-microbial defense [4-6,15-17]. We also demonstrate that ETosis in *Mnemiopsis* and
Crassostrea is stimulated by both PMA, which in mammalian cells stimulates protein kinase C
and Ca^{2+} flux, and by the Ca^{2+} ionophore A23187. These findings align with observations in
vertebrate neutrophils [31] supporting an evolutionarily conserved role for Ca^{2+} transients in ETosis.
Finally, the K^+ ionophore nigericin promoted robust ETosis in *Mnemiopsis*, indicating additional
conservation of K^+ sensing and signaling pathways. Interestingly, *Crassostrea* hemocytes did not
produce ETs in response to nigericin, raising the possibility that the nigericin-stimulated pathway
is lost in this mollusc species. Signaling pathway inhibitors optimized for use in marine
invertebrate cells will be useful for future studies of molecular actors involved in immune
responses.
In contrast to vertebrate immune cell nuclear morphologies that correspond with discrete
immune cell functions (e.g. lobed nuclei in neutrophils and eosinophils; [72]), no obvious differences
in nuclear morphologies have been reported in either bivalve hemocytes or ctenophore immune
cells. We did observe that *Mnemiopsis* immune cells that underwent ETosis *in vitro* after exposure
to *E. coli* did not appear to have phagocytosed bacteria in significant amounts (**Supp. Video 2**).
Intriguingly, we also observed other motile cells that had phagocytosed large amounts of bacteria
without undergoing ETosis (**Supp. Fig. 2, Supp. Video 4; Supp. Video 5**). Our functional
characterization of ETosis-competent *Mnemiopsis* cells suggests that future studies should
assess whether ETosis-competent and highly phagocytic non-ETotic cells represent discrete
immune cell types in ctenophores.
Ctenophores have not had specific immune cell antimicrobial behaviors described beyond
phagocytosis [36,45]. We demonstrate that *Mnemiopsis leidyi* possesses cells functionally competent
for ETosis in response to a range of microbial challenges. It remains unclear whether non-
vertebrate immune cells are capable of vital (mitochondrial) ETosis, which involves production of
ETs derived from mitochondrial DNA while maintaining cell viability [18]. Our data demonstrates that
both *Mnemiopsis leidyi* and *Crassostrea gigas* immune cells can be stimulated to produce ETs
with distinct pharmaceutical agents that, in vertebrate immune cells, differentially induce specific
intracellular signaling pathways to activate ET production. The production of extracellular traps in
Mnemiopsis suggests this specific immune cell antimicrobial defense behavior was likely present
early in metazoan evolution, prior to the diversification of extant metazoan lineages.
332
333 **Figure Legends**
334

Figure 1. Stellate and amoebocyte-like *Mnemiopsis* cells display immune behaviors. (A) DIC image of a highly granular amoebocyte-like cell. (B) DIC image of a granular stellate cell displaying multiple processes. (C) DIC image of a stellate cell with pseudopodia. (D) Merged brightfield and fluorescent image of a live *Mnemiopsis* motile, stellate cell that has phagocytosed fluorescent *E. coli* (green). Nuclei are labeled with Hoechst (blue). Lysosomes are labeled with Lysotracker-redDND99 (red). Scale bar is 10 µm. (E)-(F) Combined DIC and fluorescent images of live granular cells with pseudopodia that have phagocytosed fluorescent *E. coli*. (G) Still images from Movie S2 showing a motile stellate cell retracting its processes, undergoing nuclear rotation (spinning), and extruding its nuclear contents after exposure to *E. coli*.

Figure 2. Ctenophore immune cells undergo ETosis when exposed to microbes. (A)-(C) Confocal images of *Mnemiopsis* cell nuclei stained with Hoechst. (A) Left – Nuclei of unstimulated *Mnemiopsis* cells. Right – A cell exposed to fluorescent TxRed-*E. coli* has undergone ETosis; a large web-like pattern of chromatin has been extruded from the cell. Individual *E. coli* can be seen enmeshed by chromatin filaments (white arrowheads). (B) Still images from a 3D projection of *Mnemiopsis* ET. (C) Confocal image of *Mnemiopsis* extracellular traps composed of DNA and histones. Histone 11-4 antibody (green) and Hoechst (white) staining are visible in intact and ETosed *Mnemiopsis* cells treated with the potassium ionophore nigericin. White arrowhead marks extracellular DNA+histone chromatin nets. (D) Fluorescent microscope (Cytation) images of Hoechst-labeled *Mnemiopsis* cells. Representative images of intact nuclei are outlined in white dotted lines, while boundaries of ETs are marked with blue dotted lines. (E) Incubation with *S. aureus*, *E. coli*, and zymosan significantly induced ETosis in *Mnemiopsis* cells. Scale bar is 10 µm in all images.

Figure 3. SEM of *Mnemiopsis* ETs. Scanning electron microscopy (SEM) images of isolated *Mnemiopsis* cells. (A) Images of untreated *Mnemiopsis* stellate cells showing multiple pseudopodial processes. (B)-(C) *Mnemiopsis* cells incubated with *S. aureus* produce extracellular traps composed of long filamentous networks surrounding the cells. Individual *S. aureus* bacterium can be seen closely associated with the filaments. Insets show high magnification of *S. aureus* closely associated with the *Mnemiopsis* ETs. (*S. aureus* are pseudocolored in high magnification for clarity).

Figure 4. PMA stimulation in *Mnemiopsis* evaluated by an automated imaging analysis pipeline. All samples were incubated with their respective stimuli or seawater media alone for 4 hours and imaged. (A) Representative images of untreated controls (top) and PMA-incubated (bottom) *Mnemiopsis* cells labeled with Hoechst and Sytox Green. White dotted lines denote rough boundaries of intact nuclei. Blue dotted lines outline ETs. (B) Representative images of nuclei from SytoxGreen[high] cells which are dead or dying non-ETotic cells (top), Hoechst[low]/ SytoxGreen[low] ETotic cells showing diffuse Hoechst and SytoxGreen fluorescence (middle), Hoechst[high]/ SytoxGreen[low] live cells that show intact cell membranes. (C) Representative FlowJo plot showing distribution of fluorescent signals from *Mnemiopsis* cells labeled with Hoechst and SytoxGreen following respective treatments. (D) ETosis is significantly stimulated in *Mnemiopsis* cells are incubated with PMA gram-positive *S. aureus in vitro*. Total cell death increases significantly after incubation with dilute saline compared to seawater control, while detection of ETosis events is unchanged.

Figure 5. ETosis in *Mnemiopsis* is initiated by agonists that induce ion flux. (A) Representative Hoechst-labeled nuclei of untreated controls (left) and cells incubated with either K^+ ionophore nigericin or Ca^{2+} ionophore A23187. White dotted lines denote rough boundaries of intact nuclei. Blue dotted lines outline ETs. (B) Representative FlowJo plot showing distribution of fluorescent signals from cells labeled with Hoechst and SytoxGreen labeled following respective

386 treatments. (C) ETosis is significantly stimulated in *Mnemiopsis* cells incubated with K^+ ionophore
387 nigericin and Ca^{2+} ionophore A23187. Total cell death also increases significantly after incubation
388 with both ionophores.
389
Figure 6. Diverse stimuli induce ETosis in Crassostrea gigas hemocytes. (A) Confocal image
showing *C. gigas* hemocytes that have phagocytosed fluorescent *E. coli*. (B) Confocal image with
C. gigas hemocytes where some have phagocytosed fluorescent bacteria (arrow) and one cell
has produced an ET (arrowhead). Insets show *E. coli* and diffuse DNA staining indicative of
ETosis of a hemocyte. (C) (top) Representative images of hemocyte nuclei stained with Hoechst
that were either incubated with media alone or exposed to fungal or bacterial PAMPs. White
dotted lines denote rough boundaries of intact nuclei. Blue dotted lines outline ETs. (bottom)
Representative FlowJo plot showing distribution of fluorescent signals from *C. gigas* hemocytes
labeled with Hoechst and SytoxGreen following respective treatments. (D) ETosis is significantly
stimulated in *Crassostrea* hemocytes exposed to zymosan, gram-negative *E. coli*, and gram-
positive *S. aureus* compared to seawater media controls after 4 hours of incubation. Non-ETotic
cell death decreases slightly following zymosan incubation, increases slightly with *E. coli*
treatment. No significant change in non-ETotic cell death is detected with *S. aureus* treatment.
(E) (top) Representative images of hemocyte nuclei stained with Hoechst that were either
incubated with media alone or with PMA, K^+ ionophore nigericin, or Ca^{2+} ionophore A23187. White
dotted lines denote rough boundaries of intact nuclei. Blue dotted lines outline ETs. (bottom)
Representative FlowJo plot showing distribution of fluorescent signals from *C. gigas* hemocytes
labeled with Hoechst and SytoxGreen following respective treatments. (F) PMA and A23187, but
not nigericin, significantly stimulate ETosis in hemocytes. Observations of non-ETotic cell death
in hemocytes treated with PMA or A23187 decrease.
Scale bar is 10 μm in all images.
411
Figure 7. ETosis is an ancient metazoan immune response. Summary of ETosis phenomena
across Metazoa. The presence of cells competent for ETosis in diverse non-bilaterian,
protostome, and deuterostome taxa support the production of extracellular DNA traps as an
ancient animal immune defense mechanism. Taxa abbreviations: Ch – Choanoflagellata, Ct –
Ctenophora, Cn – Cnidaria, Tu – Tunicata, Ve – Vertebrata, An – Annelida, Mo – Mollusca, Cr –
Crustacea, He – Hexapoda. Red – this study. Grey – other studies. * – Conflicting reports of
ETosis stimulation.
419
420 ## **Methods**
421
Animal maintenance
Laboratory cultures of *Mnemiopsis leidyi* were maintained as previously described (Presnell, 2021
#35). Adult *Mnemiopsis* cells were isolated from whole animals following established protocols
[36,46,47]. *Crassostrea gigas* were maintained under flowing seawater at approximately 13°C and
hemolymph was extracted from the adductor muscle with a syringe. Isolated cells from both taxa
were maintained *in vitro* under sterile conditions in 0.22μm filtered seawater (FSW) + 1%
penicillin/streptomycin.
429
Scanning electron microscopy (SEM)
Mnemiopsis cells were seeded onto coverslips treated with poly-L-lysine. Cells were incubated
with or without heat-killed *E. coli* for 4 hours to induce ET formation. Cells were fixed in ½ strength
Karnovsky's fixative (2.5% glutaraldehyde, 2% paraformaldehyde in 0.1M sodium cacodylate
buffer, pH 7.3) overnight at 4°C. Samples were then rinsed with 0.1M cacodylate buffer and
treated with 1% osmium tetroxide for 1 hour. The samples were then dehydrated through a graded
series of alcohols and critical point dried (Autosamdri, Tousimis Corp, Rockville, MD). Samples

were mounted on stubs, sputter coated with gold/palladium (Denton Desk IV, Denton Vacuum,
Moorestown, NJ) and imaged on a JSM 6610 LV scanning electron microscope at 5kV (JEOL,
Tokyo, Japan).
440
Stimulation of ETosis and imaging
For stimulation and quantification of ETosis, cells were isolated from 24 individual animals, plated
in 96 well plates, and exposed to pHrodo-*E. coli*, *Staphylococcus aureus*, zymosan particles
(Sigma Aldrich), 25 uM nigericin (Thermo Fisher Scientific), 1 mg/mL PMA (Sigma Aldrich), or 4
uM A23187 (Sigma). Each experimental condition for each animal was performed in triplicate.
Live cell staining was performed following (2) and live imaging was performed using a JuLI Stage
(NanoEntek).
For immunofluorescence, ctenophore cells were prepared as previously described [46,47,73] and
labeled with mouse anti-histone H11-4 (EMD Millipore) and anti-mouse Alexa Fluor 488 (Thermo
Fisher Scientific). The H11-4 histone antibody was selected because it recognizes histones H1,
H2A/B, H3, and H4 proteins across diverse species. Cells were imaged using ×60 objective and
×100 oil objective, on a Nikon Ti (Eclipse) inverted microscope with Ultraview Spinning Disc (CSU-
X1) confocal scanner (Perkin Elmer). Images were captured with an Orca-ER Camera using
Volocity (Quorum technologies). Post-acquisition analysis such as contrast adjustment,
deconvolution through iterative restoration and colocalization were performed using Volocity
software.
For quantification of ETosis and total cell death, cells were treated with Hoechst 33342 (Sigma
Aldrich) and SytoxGreen (Invitrogen) for 20 min, before imaging at 20X on an automated imaging
plate reader, Cytation 3 (Biotek, software Gen5 v4.2).
460
Automated image-based profiling
We analyzed over 28,000 total images using CellProfiler (v4.1.3). Image quality was assessed by
calculating a focus score using two classes Otsu thresholding method, weighted variance on
20x20 pixel measurements. We calculated and applied an illumination correction for each
fluorescent channel (SytoxGreen and Hoechst) using a background illumination function of 50
pixels block size, without smoothing. Each corrected image was then segmented using a global
robust background method (0.05-50), with a smoothing scale of 1.3488 and a correction factor of
0.89. Clumped objects were identified and split by shape. For each segmented object we
measured the number and intensity of pixels in each fluorescent channel.
Each image and each segmented object, along with Metadata, were exported as csv files by
experiment. R (v4.0.5) software with tidyverse (v1.3.1), dplyr (v1.0.7) and readr (v1.4.0) packages
were then used to transform the datasets. Data from images and objects were merged, and
measurements from individual images with a Focus Score <0.2 were removed from further
analysis. This allowed us to identify and select only images that were in focus. Surface area,
Hoechst intensity and SytoxGreen intensity per object (nucleus) and per individual animal were
then imported into FlowJo (v10.8.0), and percentages of cells per delineated population
(dead/dying cell, live cell, and ETotic cell) were calculated. Dying and ETotic cells were gated as
indicated on the figures. Finally, percentages per individual animal surveyed were combined and
tested for statistical significance using GraphPad Prism (v9.2.0). All statistical tests were
performed using two-tailed unpaired student t-test *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$.
481

Acknowledgements
The authors are grateful to Anna Bruchez and Rachel Prins for imaging support. This research
was supported by the National Oceanographic and Atmospheric Administration, a National
Research Council Postdoctoral Fellowship to LEV, and the National Science Foundation under
Grant No. 2013692. CS and ALH were supported by National Institutes of Health Grants
R33AI119341 and R01GM102482. The authors thank Steve Ziegler for technical and

experimental support. The authors gratefully acknowledge the Fred Hutchinson Cancer Center
Electron Microscopy Shared Resource (EMSR). The EMSR is supported in part by the Cancer
Center Support Grant P30 CA015704-40.
491
Author contributions
Conceptualization, C.S. and L.E.V.; Methodology, L.E.V. and C.S.; Formal Analysis, C.S. and
L.E.V.; Investigation, L.E.V. and C.S.; Visualization, C.S., L.E.V., and W.E.B.; Writing – Original
Draft, L.E.V. and C.S.; Writing – Review & Editing, C.S., A.L.H., W.E.B., F.W.G., and L.E.V.;
Funding Acquisition, A.L.H., W.E.B., N.T.K., F.W.G., and L.E.V.; Resources, A.L.H., W.E.B., and
F.W.G.; Supervision, A.L.H., W.E.B., and F.W.G.
498
499 Competing interests
500 The authors declare no competing interests.
501
502 **Supplemental Material**
503
504 **Supp. Figure 1:** Widefield views of SEM images of unstimulated (left panel) and microbe-
505 challenged (right panel) *Mnemiopsis* cells *in vitro*. Unstimulated cells from whole *Mnemiopsis*
506 show diverse sizes and morphologies indicating the presence of multiple cell types. Microbe-
507 exposed *Mnemiopsis* cells were incubated with *S. aureus* for 3 hours.
508
509 **Supp. Figure 2:** Workflow schematic of the automatic imaging analysis pipeline.
510
511
512 **Supp. Video 1:** Video of DIC timelapse images of isolated *Mnemiopsis* cells *in vitro*. An
513 amoebocyte-like granular cell (arrowhead) and highly motile stellate cell (arrow) are visible.
514
515 **Supp. Video 2**: Video of DIC timelapse images of an isolated *Mnemiopsis* stellate cell *in vitro*.
516 The cell initially has two large processes, then absorbs them and subsequently produces several
517 additional processes. The cell then begins to crawl out of view.
518
519 **Supp. Video 3:** Merged brightfield and fluorescent video of a live *Mnemiopsis* motile, stellate cell
520 that is phagocytosing fluorescent *E. coli* (green). *Mnemiopsis* cells were labeled with Lysotracker
521 Red. Scale bar is 10 μm.
522
523 **Supp. Video 4**: Merged brightfield and fluorescent video of a live *Mnemiopsis* motile, stellate cell
524 undergoing ETosis after *in vitro* exposure to pHrodo-*E. coli*. The cell moves into view, retracts its
525 processes, spins, and exudes its nuclear material. DNA (Hoechst, blue), pHrodo-*E. coli* (red)
526
527 **Supp. Video 5:** 3D reconstruction of confocal stack of *Mnemiopsis* extracellular DNA traps. DNA
528 (Hoechst, white) is extracellular, in filamentous "nets"; lysosomal marker (Lysotracker, green)
529 denotes cellular debris; bacteria are ensnared in the "nets" (pHrodo-*E. coli*, red)
530
531 **Supp. Video 6:** Merged brightfield and fluorescent video of a live *Mnemiopsis* motile, stellate cell
532 that is phagocytosing fluorescent *E. coli* (red). Scale bar is 10 μm.
533
534
535 **REFERENCES:**
536

1   Brinkmann, V. *et al.* Neutrophil extracellular traps kill bacteria. *Science* **303**, 1532-1535
(2004). https://doi.org:10.1126/science.1092385

2 Wartha, F., Beiter, K., Normark, S. & Henriques-Normark, B. Neutrophil extracellular traps:
casting the NET over pathogenesis. *Curr Opin Microbiol* **10**, 52-56 (2007).
https://doi.org:10.1016/j.mib.2006.12.005
3 Han, Y., Zhang, Q., Chen, L., Yang, D. & Zhao, J. Mitochondria are essential for
antibacterial extracellular trap formation mediated by zymosan in hemocytes of Ruditapes
philippinarum. *Dev Comp Immunol* **121**, 104094 (2021).
https://doi.org:10.1016/j.dci.2021.104094
4 Poirier, A. C. *et al.* Antimicrobial histones and DNA traps in invertebrate immunity:
evidences in Crassostrea gigas. *J Biol Chem* **289**, 24821-24831 (2014).
https://doi.org:10.1074/jbc.M114.576546
5 Robb, C. T., Dyrynda, E. A., Gray, R. D., Rossi, A. G. & Smith, V. J. Invertebrate
extracellular phagocyte traps show that chromatin is an ancient defence weapon. *Nat
Commun* **5**, 4627 (2014). https://doi.org:10.1038/ncomms5627
6 Romero, A., Novoa, B. & Figueras, A. Extracellular traps (ETosis) can be activated through
NADPH-dependent and -independent mechanisms in bivalve mollusks. *Dev Comp
Immunol* **106**, 103585 (2020). https://doi.org:10.1016/j.dci.2019.103585
7 Dunn, C. W. *et al.* Broad phylogenomic sampling improves resolution of the animal tree of
life. *Nature* **452**, 745-749 (2008). https://doi.org:10.1038/nature06614
8 Li, Y., Shen, X. X., Evans, B., Dunn, C. W. & Rokas, A. Rooting the Animal Tree of Life.
Mol Biol Evol **38**, 4322-4333 (2021). https://doi.org:10.1093/molbev/msab170
9 Redmond, A. K. & McLysaght, A. Evidence for sponges as sister to all other animals from
partitioned phylogenomics with mixture models and recoding. *Nat Commun* **12**, 1783
(2021). https://doi.org:10.1038/s41467-021-22074-7
10 Schultz, D. T. *et al.* Ancient gene linkages support ctenophores as sister to other animals.
Nature **618**, 110-117 (2023). https://doi.org:10.1038/s41586-023-05936-6
11 Buchmann, K. Evolution of Innate Immunity: Clues from Invertebrates via Fish to
Mammals. *Front Immunol* **5**, 459 (2014). https://doi.org:10.3389/fimmu.2014.00459
12 Mandujano-Tinoco, E. A., Sultan, E., Ottolenghi, A., Gershoni-Yahalom, O. & Rosental, B.
Evolution of Cellular Immunity Effector Cells; Perspective on Cytotoxic and Phagocytic
Cellular Lineages. *Cells* **10** (2021). https://doi.org:10.3390/cells10081853
13 Fuchs, T. A. *et al.* Novel cell death program leads to neutrophil extracellular traps. *J Cell
Biol* **176**, 231-241 (2007). https://doi.org:10.1083/jcb.200606027
14 Sollberger, G., Tilley, D. O. & Zychlinsky, A. Neutrophil Extracellular Traps: The Biology
of Chromatin Externalization. *Dev Cell* **44**, 542-553 (2018).
https://doi.org:10.1016/j.devcel.2018.01.019
15 Homa, J., Ortmann, W. & Kolaczkowska, E. Conservative Mechanisms of Extracellular
Trap Formation by Annelida Eisenia andrei: Serine Protease Activity Requirement. *PLoS
One* **11**, e0159031 (2016). https://doi.org:10.1371/journal.pone.0159031
16 Lange, M. K. *et al.* Gastropod-derived haemocyte extracellular traps entrap
metastrongyloid larval stages of Angiostrongylus vasorum, Aelurostrongylus abstrusus
and Troglostrongylus brevior. *Parasit Vectors* **10**, 50 (2017).
https://doi.org:10.1186/s13071-016-1961-z
17 Nascimento, M. T. C. *et al.* DNA extracellular traps are part of the immune repertoire of
Periplaneta americana. *Dev Comp Immunol* **84**, 62-70 (2018).
https://doi.org:10.1016/j.dci.2018.01.012
18 Ramos-Martinez, E. *et al.* Multiple Origins of Extracellular DNA Traps. *Front Immunol* **12**,
621311 (2021). https://doi.org:10.3389/fimmu.2021.621311
19 Zhang, X., Zhuchenko, O., Kuspa, A. & Soldati, T. Social amoebae trap and kill bacteria
by casting DNA nets. *Nat Commun* **7**, 10938 (2016).
https://doi.org:10.1038/ncomms10938

20 Wen, F., White, G. J., VanEtten, H. D., Xiong, Z. & Hawes, M. C. Extracellular DNA is
required for root tip resistance to fungal infection. *Plant Physiol* **151**, 820-829 (2009).
https://doi.org:10.1104/pp.109.142067
21 Ward, J. R. *et al.* A central role for monocytes in Toll-like receptor-mediated activation of
the vasculature. *Immunology* **128**, 58-68 (2009). https://doi.org:10.1111/j.1365-
2567.2009.03071.x
22 Zhang, Q., Zmasek, C. M. & Godzik, A. Domain architecture evolution of pattern-
recognition receptors. *Immunogenetics* **62**, 263-272 (2010).
https://doi.org:10.1007/s00251-010-0428-1
23 Snoeck, S. *et al.* Evolutionary gain and loss of a plant pattern-recognition receptor for
HAMP recognition. *Elife* **11** (2022). https://doi.org:10.7554/eLife.81050
24 Tassia, M. G., Whelan, N. V. & Halanych, K. M. Toll-like receptor pathway evolution in
deuterostomes. *Proc Natl Acad Sci U S A* **114**, 7055-7060 (2017).
https://doi.org:10.1073/pnas.1617722114
25 Saco, A., Novoa, B., Greco, S., Gerdol, M. & Figueras, A. Bivalves Present the Largest
and Most Diversified Repertoire of Toll-Like Receptors in the Animal Kingdom, Suggesting
Broad-Spectrum Pathogen Recognition in Marine Waters. *Mol Biol Evol* **40** (2023).
https://doi.org:10.1093/molbev/msad133
26 Parthier, C. *et al.* Structure of the Toll-Spatzle complex, a molecular hub in Drosophila
development and innate immunity. *Proc Natl Acad Sci U S A* **111**, 6281-6286 (2014).
https://doi.org:10.1073/pnas.1320678111
27 Dong, C., Davis, R. J. & Flavell, R. A. MAP kinases in the immune response. *Annu Rev*
Immunol **20**, 55-72 (2002). https://doi.org:10.1146/annurev.immunol.20.091301.131133
28 Gostner, J. M., Becker, K., Fuchs, D. & Sucher, R. Redox regulation of the immune
response. *Redox Rep* **18**, 88-94 (2013).
https://doi.org:10.1179/1351000213Y.0000000044
29 Vig, M. & Kinet, J. P. Calcium signaling in immune cells. *Nat Immunol* **10**, 21-27 (2009).
https://doi.org:10.1038/ni.f.220
30 Akira, S., Uematsu, S. & Takeuchi, O. Pathogen recognition and innate immunity. *Cell*
124, 783-801 (2006). https://doi.org:10.1016/j.cell.2006.02.015
31 Kenny, E. F. *et al.* Diverse stimuli engage different neutrophil extracellular trap pathways.
Elife **6** (2017). https://doi.org:10.7554/eLife.24437
32 Auguste, M., Balbi, T., Ciacci, C. & Canesi, L. Conservation of Cell Communication
Systems in Invertebrate Host-Defence Mechanisms: Possible Role in Immunity and
Disease. *Biology (Basel)* **9** (2020). https://doi.org:10.3390/biology9080234
33 Douda, D. N., Khan, M. A., Grasemann, H. & Palaniyar, N. SK3 channel and mitochondrial
ROS mediate NADPH oxidase-independent NETosis induced by calcium influx. *Proc Natl*
Acad Sci U S A **112**, 2817-2822 (2015). https://doi.org:10.1073/pnas.1414055112
34 Ryan, J. F. *et al.* The genome of the ctenophore Mnemiopsis leidyi and its implications for
cell type evolution. *Science* **342**, 1242592 (2013).
https://doi.org:10.1126/science.1242592
35 Moroz, L. L. *et al.* The ctenophore genome and the evolutionary origins of neural systems.
Nature **510**, 109-114 (2014). https://doi.org:10.1038/nature13400
36 Traylor-Knowles, N., Vandepas, L. E. & Browne, W. E. Still Enigmatic: Innate Immunity in
the Ctenophore Mnemiopsis leidyi. *Integr Comp Biol* **59**, 811-818 (2019).
https://doi.org:10.1093/icb/icz116
37 Srivastava, M. *et al.* The Amphimedon queenslandica genome and the evolution of animal
complexity. *Nature* **466**, 720-726 (2010). https://doi.org:10.1038/nature09201
38 Putnam, N. H. *et al.* Sea anemone genome reveals ancestral eumetazoan gene repertoire
and genomic organization. *Science* **317**, 86-94 (2007).
https://doi.org:10.1126/science.1139158

Bachere, E. *et al.* The new insights into the oyster antimicrobial defense: Cellular, molecular and genetic view. *Fish Shellfish Immunol* **46**, 50-64 (2015). https://doi.org:10.1016/j.fsi.2015.02.040

Jiang, S. *et al.* Functional characterisation of phagocytes in the Pacific oyster Crassostrea gigas. *PeerJ* **4**, e2590 (2016). https://doi.org:10.7717/peerj.2590

Babonis, L. S., Ryan, J. F., Enjolras, C. & Martindale, M. Q. Genomic analysis of the tryptome reveals molecular mechanisms of gland cell evolution. *Evodevo* **10**, 23 (2019). https://doi.org:10.1186/s13227-019-0138-1

Jokura, K. *et al.* CTENO64 Is Required for Coordinated Paddling of Ciliary Comb Plate in Ctenophores. *Curr Biol* **29**, 3510-3516 e3514 (2019). https://doi.org:10.1016/j.cub.2019.08.059

Presnell, J. S. & Browne, W. E. Kruppel-like factor gene function in the ctenophore Mnemiopsis leidyi assessed by CRISPR/Cas9-mediated genome editing. *Development* **148** (2021). https://doi.org:10.1242/dev.199771

Bolte, S. *et al.* Specific immune priming in the invasive ctenophore Mnemiopsis leidyi. *Biol Lett* **9**, 20130864 (2013). https://doi.org:10.1098/rsbl.2013.0864

Franc, J. M. La matrice mésogléenne des Cténaires. *Bulletin de la Société zoologique de France* **113**, 347-447 (1988).

Dieter, A. C., Vandepas, L. E. & Browne, W. E. Isolation and Maintenance of In Vitro Cell Cultures from the Ctenophore Mnemiopsis leidyi. *Methods Mol Biol* **2450**, 347-358 (2022). https://doi.org:10.1007/978-1-0716-2172-1_18

Vandepas, L. E., Warren, K. J., Amemiya, C. T. & Browne, W. E. Establishing and maintaining primary cell cultures derived from the ctenophore Mnemiopsis leidyi. *J Exp Biol* **220**, 1197-1201 (2017). https://doi.org:10.1242/jeb.152371

Boe, D. M., Curtis, B. J., Chen, M. M., Ippolito, J. A. & Kovacs, E. J. Extracellular traps and macrophages: new roles for the versatile phagocyte. *J Leukoc Biol* **97**, 1023-1035 (2015). https://doi.org:10.1189/jlb.4RI1014-521R

Apel, F. *et al.* The cytosolic DNA sensor cGAS recognizes neutrophil extracellular traps. *Sci Signal* **14** (2021). https://doi.org:10.1126/scisignal.aax7942

Yousefi, S. *et al.* Catapult-like release of mitochondrial DNA by eosinophils contributes to antibacterial defense. *Nat Med* **14**, 949-953 (2008). https://doi.org:10.1038/nm.1855

Lood, C. *et al.* Neutrophil extracellular traps enriched in oxidized mitochondrial DNA are interferogenic and contribute to lupus-like disease. *Nat Med* **22**, 146-153 (2016). https://doi.org:10.1038/nm.4027

Nagase, H., Ueki, S. & Fujieda, S. The roles of IL-5 and anti-IL-5 treatment in eosinophilic diseases: Asthma, eosinophilic granulomatosis with polyangiitis, and eosinophilic chronic rhinosinusitis. *Allergol Int* **69**, 178-186 (2020). https://doi.org:10.1016/j.alit.2020.02.002

Fukuchi, M. *et al.* How to detect eosinophil ETosis (EETosis) and extracellular traps. *Allergol Int* **70**, 19-29 (2021). https://doi.org:10.1016/j.alit.2020.10.002

de Buhr, N. & von Kockritz-Blickwede, M. How Neutrophil Extracellular Traps Become Visible. *J Immunol Res* **2016**, 4604713 (2016). https://doi.org:10.1155/2016/4604713

Barnes, B. J. *et al.* Targeting potential drivers of COVID-19: Neutrophil extracellular traps. *J Exp Med* **217** (2020). https://doi.org:10.1084/jem.20200652

Zhao, M. L., Chi, H. & Sun, L. Neutrophil Extracellular Traps of Cynoglossus semilaevis: Production Characteristics and Antibacterial Effect. *Front Immunol* **8**, 290 (2017). https://doi.org:10.3389/fimmu.2017.00290

Brinkmann, V. & Zychlinsky, A. Neutrophil extracellular traps: is immunity the second function of chromatin? *J Cell Biol* **198**, 773-783 (2012). https://doi.org:10.1083/jcb.201203170

58   Seo, K. S. *et al.* PMA synergistically enhances apicularen A-induced cytotoxicity by
disrupting microtubule networks in HeLa cells. *BMC Cancer* **14**, 36 (2014).
https://doi.org:10.1186/1471-2407-14-36

59   Riegger, J. *et al.* Histopathological evaluation of thrombus in patients presenting with stent
thrombosis. A multicenter European study: a report of the prevention of late stent
thrombosis by an interdisciplinary global European effort consortium. *Eur Heart J* **37**,
1538-1549 (2016). https://doi.org:10.1093/eurheartj/ehv419

60   Elsherif, L. *et al.* Machine Learning to Quantitate Neutrophil NETosis. *Sci Rep* **9**, 16891
(2019). https://doi.org:10.1038/s41598-019-53202-5

61   Gupta, S., Chan, D. W., Zaal, K. J. & Kaplan, M. J. A High-Throughput Real-Time Imaging
Technique To Quantify NETosis and Distinguish Mechanisms of Cell Death in Human
Neutrophils. *J Immunol* **200**, 869-879 (2018). https://doi.org:10.4049/jimmunol.1700905

62   Remijsen, Q. *et al.* Dying for a cause: NETosis, mechanisms behind an antimicrobial cell
death modality. *Cell Death Differ* **18**, 581-588 (2011). https://doi.org:10.1038/cdd.2011.1

63   Ojcius, D. M., Zychlinsky, A., Zheng, L. M. & Young, J. D. Ionophore-induced apoptosis:
role of DNA fragmentation and calcium fluxes. *Exp Cell Res* **197**, 43-49 (1991).
https://doi.org:10.1016/0014-4827(91)90477-c

64   Yu, S. P. Regulation and critical role of potassium homeostasis in apoptosis. *Prog
Neurobiol* **70**, 363-386 (2003). https://doi.org:10.1016/s0301-0082(03)00090-x

65   Zhu, S. *et al.* Nigericin treatment activates endoplasmic reticulum apoptosis pathway in
goldfish kidney leukocytes. *Fish Shellfish Immunol* **134**, 108616 (2023).
https://doi.org:10.1016/j.fsi.2023.108616

66   Cai, X. Unicellular Ca2+ signaling 'toolkit' at the origin of metazoa. *Mol Biol Evol* **25**, 1357-
1361 (2008). https://doi.org:10.1093/molbev/msn077

67   Gandara, A. C. P., Torres, A., Bahia, A. C., Oliveira, P. L. & Schama, R. Evolutionary
origin and function of NOX4-art, an arthropod specific NADPH oxidase. *BMC Evol Biol* **17**,
92 (2017). https://doi.org:10.1186/s12862-017-0940-0

68   Tompkins-Macdonald, G. J. *et al.* Expression of a poriferan potassium channel: insights
into the evolution of ion channels in metazoans. *J Exp Biol* **212**, 761-767 (2009).
https://doi.org:10.1242/jeb.026971

69   P.M., H. The inter-relationships of bivalve haemocytes. *Fish & Shellfish Immunology* **9**,
pp.367-385. (1999).

70   Do, E. A. & Gries, C. M. Beyond Homeostasis: Potassium and Pathogenesis during
Bacterial Infections. *Infect Immun* **89**, e0076620 (2021). https://doi.org:10.1128/IAI.00766-
20

71   Khan, M. A. & Palaniyar, N. Transcriptional firing helps to drive NETosis. *Sci Rep* **7**, 41749
(2017). https://doi.org:10.1038/srep41749

72   Skinner, B. M. & Johnson, E. E. Nuclear morphologies: their diversity and functional
relevance. *Chromosoma* **126**, 195-212 (2017). https://doi.org:10.1007/s00412-016-0614-
5

73   Acharya, M. *et al.* alphav Integrins combine with LC3 and atg5 to regulate Toll-like receptor
signalling in B cells. *Nat Commun* **7**, 10917 (2016). https://doi.org:10.1038/ncomms10917
731

Figures

Figure 1

Figure 3

737
738

741
742

Figure 7