Organization of cortico-hippocampal networks in the human brain

Short Title: Organization of cortico-hippocampal networks

Alexander J. Barnett*[1], Walter Reilly[1], Halle R. Dimsdale-Zucker[2], Eda Mizrak[1,3], Zachariah Reagh[1,4,5], Charan Ranganath[1]

1. University of California, Davis, Center for Neuroscience, 1544 Newton Ct, Davis, CA 95618, United States
2. Columbia University, Department of Psychology, 312 Schermerhorn Hall 1190 Amsterdam Avenue New York, NY 10027
3. University of Zurich, Department of Psychology, Binzmühlestrasse 14, 8050 Zürich, Switzerland
4. University of California, Davis, Department of Neurology, 3160 Folsom Blvd., Suite 2100, Sacramento, CA 95816
5. Washington University in St. Louis, Department of Psychological and Brain Sciences, 1 Brookings Dr. St. Louis, MO 63130

*Corresponding Author
Alexander J. Barnett
1544 Newton Ct, Davis, CA 95618, United States
Center for Neuroscience, Room 209
Email: ajbarnett@ucdavis.edu

Alexander J. Barnett: https://orcid.org/0000-0001-5891-7880
Walter Reilly: https://orcid.org/0000-0003-2303-3094
Halle Dimsdale-Zucker: https://orcid.org/0000-0002-8611-111X
Eda Mizrak: https://orcid.org/0000-0002-1765-7047
Zachariah Reagh: https://orcid.org/0000-0002-4763-4268
Charan Ranganath: https://orcid.org/0000-0001-5835-6091

Author Contributions
AJB, WR, HDZ, and CR designed research; AJB, HDZ, EM, and ZR performed research; EM contributed new reagents or analytic tools; AJB and WR analyzed data. AJB and CR wrote the paper

Abstract

Episodic memory depends on interactions between the hippocampus and the interconnected regions comprising default mode network (DMN). Here, using data-driven analyses of resting-state fMRI data, we identified the networks that interact with the hippocampus—the DMN and a "Medial Temporal Network" (MTN) that included regions in the medial temporal lobe and retrosplenial cortex. We observed that the MTN plays a critical role in connecting the visual network to the DMN and hippocampus. The DMN could be further divided into three subnetworks: a "Posterior-Medial" Subnetwork comprised of posterior cingulate, and lateral parietal cortices, an "Anterior-Temporal" Subnetwork comprised of regions in the temporopolar, and dorsomedial prefrontal cortex, and a "Medial-Prefrontal" Subnetwork comprised of regions primarily in the medial prefrontal cortex. These networks vary in their functional connectivity along the hippocampal long-axis and represent different kinds of information during memory-guided decision-making. Finally, a Neurosynth meta-analysis of fMRI studies suggests new hypotheses regarding the functions of the MTN and DMN subnetworks, providing a framework to guide future research on the neural architecture of episodic memory.

Episodic memory allows us to relive past events that happened at a particular place and time (1).

Most research investigating the neurobiology of episodic memory retrieval has focused on the

hippocampus and medial temporal lobe (MTL) cortex (perirhinal (PRC), parahippocampal

(PHC), and entorhinal cortex) highlighting the interactions among these regions (2–4). It is

generally assumed, however, that the hippocampus supports memory by integrating information

represented across distributed areas in the neocortex (5). Consistent with this idea, fMRI studies

of memory retrieval have shown activity and connectivity in a cortico-hippocampal network

composed of medial parietal, lateral parietal, lateral temporal, and medial prefrontal neocortical

regions along with the MTL (6–10), and lesion studies have shown that damage within this

network can cause amnesia (11).

This distributed set of cortical regions that are recruited during episodic retrieval overlaps

extensively with regions that comprise the default-mode network (DMN). The DMN is a large-

scale network that is typically identified in studies that use intrinsic functional connectivity

analysis of functional magnetic resonance imaging (fMRI) data (12), and some studies suggest

that the DMN can be differentiated into different subnetworks (13–15). One approach to partition

the DMN has been to view DMN subnetworks as an extension of the connectivity differences

within the MTL (16–18). Guided by neuroanatomical studies of MTL connectivity in rodents and

nonhuman primates (19–22), studies of intrinsic functional connectivity in fMRI data (23–26)

have differentiated between cortical regions that preferentially affiliate with the PRC and anterior

hippocampus and cortical regions that preferentially affiliate with the PHC and posterior

hippocampus (27, 28). For example, Libby et al. (24) demonstrated that the PHC has relatively

higher functional connectivity with a network of Posterior Medial (PM) cortical regions,

including the posterior cingulate and retrosplenial cortex and posterior hippocampus, whereas the

PRC has relatively higher connectivity with a network of Anterior Temporal (AT) cortical

regions, including the orbitofrontal and temporopolar cortex and anterior hippocampus. Drawing

on these findings, Ranganath and Ritchey (29) reviewed evidence converging on the idea that

PM and AT networks support different aspects of memory-guided behavior. This "PM/AT

framework" has provided a valuable framework for interpreting memory phenomena in imaging

(9, 30–32), stimulation (33, 34), and disease (35–37). However, recent work has come to

question the homogeneity of the PM network (38), with several studies highlighting a

dissociation between PHC, retrosplenial and precuneal cortex from posterior cingulate cortex

(PCC) (39, 40) under conditions that seem to tax perceptual relative to abstract event processing,

or scene relative to face processing.

A second approach to segmenting the DMN has used data-driven methods and these

studies have also found partially conflicting network delineations (15, 41, 42). These studies

have generally partitioned the DMN into three subnetworks: an MTL subnetwork, a midline

subnetwork of posterior cingulate and medial prefrontal cortex, and a third network composed of

dorsal medial prefrontal, lateral temporal, and ventrolateral prefrontal cortex, but have focussed

largely on cortical regions. Critically, it is unclear how these data-driven, cortical networks

pertain to memory functioning and previous frameworks that have strongly characterized

hippocampal connections.

Here, we seek to unify these two approaches. The MTL-centric approach has rarely

considered the interplay of regions outside of the MTL, and a recent review has suggested that

considerable heterogeneity may exist within the previously described PM network (38),

prompting a need for re-examination of these network properties. Conversely, the data-driven

approach has rarely described the hippocampal interactions with these networks, focusing mainly

on cortico-cortical interactions limiting the applicability to understanding episodic memory. The

goal of this paper was to comprehensively characterize cortico-hippocampal networks that

contribute to episodic memory. Using a whole-brain, data-driven approach to examine functional

connectivity in resting-state fMRI, we sought to 1. identify and partition the DMN into

subnetworks and examine whether these subnetworks converge with the PM/AT framework, 2.

examine the connectivity of the hippocampus to the identified networks, and 3. determine

whether these cortico-hippocampal networks play different roles in episodic memory.

Results

Data-Driven Partition of Neocortical Networks

Our first objective was to use a data-driven approach to identify large-scale resting-state

networks. Using 25 minutes of resting-state fMRI data acquired from 40 participants, we

partitioned the brain into canonical resting networks using state-of-the-art techniques (43). We

extracted the average confound-corrected timeseries from each region in a recently published

atlas of the human neocortex (44) and computed Fisher z-transformed Pearson's correlations

between the timeseries of each cortical region and every other cortical region. We then created a

group-averaged functional connectivity matrix to be used for community detection. The Louvain

community detection algorithm (45) was run for 1000 iterations, tuning the resolution (how large

or small communities might be) across a range of resolution parameters to identify a partition

solution that showed both high network modularity (i.e. higher within community connectivity

than would be expected by chance) and high network grouping consistency (quantified using the

z-Rand index (43, 46)). The solution was also subject to a qualitative criterion that the partition

should separate the primary sensory networks (43). The resolution parameter of gamma = 2.005

satisfied our criteria and produced a network partition (Figure 1) that corresponds closely with

previously reported partitions in other recent studies (43, 47, 48). This partition identified the

DMN, but, interestingly, some areas (such as the PHC, retrosplenial cortex, and PRC) that are

often identified as part of the DMN were grouped within a separate network that included MTL

and parietal regions. This set of regions overlaps with what has been previously described as the

MTL network (13), or the contextual association network (49), and coactivates particularly

during recall of places (50, 51). Here we will describe this network as the medial temporal

network (MTN).

<u>Figure 1. Louvain community detection identifies large-scale resting-state networks including</u>

<u>the default-mode and medial temporal networks that are thought to interface with the</u>

<u>hippocampus.</u> A. Inflated cortical surface, colored according to community membership. B.

Connectivity matrix reordered by community to demonstrate the community structure of the

group-averaged brain. Colors along the axis demonstrate which rows/columns belong to a given

community. Color bar represents Fisher Z-transformed correlation values. C. Force-directed

graph of the group-averaged networks color coded by community membership of the selected

partition created using the ForceAtlas2 algorithm (52). Attn, attention; DMN, Default mode

network; Front Par, Frontoparietal; MTN, Medial temporal network; Som-Mot, Somatomotor.

We next interrogated the functional connectivity between the networks defined above and the

hippocampus. We segmented the hippocampus in each participant using FreeSurfer v6.0

(http://surfer.nmr.mgh.harvard.edu/, (53)), and then manually segmented it further into anterior

and posterior regions, as these subdivisions of the hippocampus are known to have somewhat

different functional properties (16, 20). The posterior hippocampus was defined as all the

hippocampus posterior to the last slice of the uncal notch (16). We calculated the connectivity of

the hippocampus to every cortical region in the atlas and averaged together functional

connectivity weights of cortical regions within the same network based on network affiliations.

This was done for each participant. One-sample t-tests revealed that two networks were

functionally connected to every hippocampal ROI in our sample—the DMN and the MTN

(DMN range: $t(38) = 10.7–12.9$, *all P < .001*; MTN range: $t(38) = 4.2–9.3$, *all P < .01*). The

language network and somatomotor network both showed mixed connectivity with the

hippocampus, showing significant anterior, but not posterior hippocampal connectivity (see

Supplemental Table 1). The robust connectivity of the MTN and DMN to the hippocampus

corroborates previous investigations of hippocampal connectivity (54).

<u>Network analysis support an interfacing role of the MTN</u>

A recent review discussing the heterogeneity within the PM network suggested that

parahippocampal, and lateral parietal cortex interface with lower level sensory cortex in service

of higher-order feature encoding prior to binding in the hippocampus (29, 38). They also note

that these regions also interface with other DMN regions such as the mPFC and PCC (regions

found in the DMN, here) that track long-timescale event structure and provide conceptual

knowledge pertaining to event schemas (38, 55). Thus, the MTN may serve as a bridge between

external, low level perception in the visual network and the hippocampus and DMN. By plotting

the structure of the network interactions in Figure 2A (left) we do indeed see that the MTN

shows functional connectivity with both the DMN, hippocampus, and the visual network.

Treating the brain as an abstract graph (56), we sought to test the hypothesis that the MTN

mediates network communication between the visual network and the DMN and hippocampus.

To do so, we used graph theory to calculate the shortest path length between the visual network

and the DMN and hippocampus. Path length refers to the shortest number of links that needs to

be traversed to link one node in a network to a target node. If two regions are directly connected

by a strong link, they will have a short path length. If two regions are not directly connected by a

positive link but can be connected by multiple intermediate nodes that are positively connected,

then they would have a longer path length. To examine the influence of the MTN on

internetwork communication between the DMN and visual network, we compared the path

length between the visual network and DMN when the MTN was removed relative to the path

length when we removed all other networks. If the MTN mediates the connectivity between the

visual network and DMN and hippocampus, then removal of the MTN should disproportionately

increase the average path length between visual network and DMN over and above any path

length increases caused by removal of all other networks. This was done at the individual-subject

level to determine the significance of the effect and is similar to methods used by (57).

We found that across subjects, removal of the MTN led to a disproportionately large

increase in DMN–visual network path length compared to removal of any other network (all

$t(39) > 6.7$, p < .001; Figure 2B), indicating that the MTN plays a critical role in cross-network

communication between the visual network and DMN. Importantly, this effect cannot be

explained simply by the number of nodes removed from the whole network, as the MTN has

fewer nodes (30 nodes) than the somatomotor (34 nodes), salience (35 nodes), frontoparietal (44

nodes) and auditory (33 nodes) networks. To further confirm this, we randomly removed 30

nodes from each subject's network (excluding DMN, and visual nodes) and recalculated the path

length values. This was performed 1000 times and we compared the DMN-visual network path

length of the true removal of the MTN nodes compared to the distribution of DMN-visual

network path length following random node removals. We observed that the MTN removal still

resulted in greater average path length between the DMN and visual network relative to removal

of random nodes, $t(39) = 9.4$, $p < .001$. Another possible explanation is that due to the high level

of connectivity between the DMN and MTN, removal of the MTN will non-specifically lead to a

disproportionate increase in path length of the DMN to all other networks. However, the path

length effects caused by removal of the MTN were specific to the DMN – visual network path

lengths (S1 Figure).

Figure 2. Removal of the MTN disproportionately decreases the network connectivity of the

DMN and visual network. (A) Force-directed graph of the group-averaged networks color coded

by community membership of the selected partition created using the ForceAtlas2 algorithm (52)

with the MTN present (left) and with the MTN excluded (right). (B) Boxplots displaying the

path length between the DMN and visual network or the path length between the hippocampi and

visual network following removal of each network. *** p < .001.

As mentioned above, it is well-known that MTL regions such as the perirhinal and

parahippocampal cortex provide the hippocampus with higher-order representations of objects

and scenes (4, 58). Therefore, we repeated the previous analysis, this time examining the effects

of MTN removal on the pathlength between the hippocampus and visual network. We observed

that removal of the MTN lead to greater path length between the hippocampus and visual

network compared to removal of any other networks (all $t(39) > 6$, p < .001; Figure 2B). Again,

MTN removal led to disproportionately higher path length than removal of random nodes ($t(39)$

= 3.15, $p = .003$). This effect was specifically between the hippocampus and visual network path

length, rather than leading to disproportionately higher path length between the hippocampus and

any other network, S2 Figure.

<u>Partitioning the DMN</u>

We next sought to identify whether the DMN could be broken up into subnetworks using similar

methods that we used in our whole-brain network partition. We selected the DMN regions

identified in our whole-brain partition and performed Louvain community detection on the

functional connections between the regions in the network. This was done for 1000 iterations

across a range of resolution parameters, and we again selected the partition yielding the highest

modularity and stability. The solution with the highest modularity-weighted stability produced a

partition of the DMN with three communities which we refer to as the PM subnetwork, the AT

subnetwork and the medial prefrontal (MP) subnetwork. The PM subnetwork and AT

subnetwork were named as such based on their resemblance to previous spatial maps of these

networks (24). The PM subnetwork included posterior cingulate cortex, posterior angular gyrus,

and dorsal prefrontal cortex. The AT subnetwork included temporopolar cortex, lateral

orbitofrontal cortex, temporoparietal junction and dorsomedial prefrontal cortex. The MP

subnetwork included medial prefrontal regions and the entorhinal cortex (Figure 3).

Figure 3. The default-mode network can be partitioned into three interconnect subnetworks. A. Functional connectivity matrix organized by subnetwork. Color bar represents Fisher Z-transformed correlation values. B. Inflated cortical surface, colored according to community membership of subnetworks. AT, anterior temporal; MP, medial prefrontal; PM, posterior medial.

To visualize the topology of connections between these DMN sub-networks and the other large-scale networks defined earlier, we constructed a force-directed graph in abstract graph space

(Figure 4). The ForceAtlas2 algorithm was used to construct the layout by modeling the graph as

a physical system in which nodes repel each other and the functional connectivity between the

nodes act as springs (52). Regions within the PM subnetwork tightly clustered and several nodes

showed prominent cross-network connectivity with the MTN and frontoparietal network. The

AT subnetwork showed cross-network interactions with the language and frontoparietal network

and the MP subnetwork was less clustered with fewer out of DMN connections.

Figure 4. <u>The subnetworks of the DMN interact with language, frontoparietal and medial</u>

<u>temporal networks.</u> A. Force-directed graph of the group-averaged networks with the DMN

relabelled according to subnetwork membership. B. Boxplots showing participation coefficient

for DMN nodes to the rest of the brain (Whole Brain) and participation coefficient for DMN

nodes to subnetworks of the DMN (Within DMN). AT, anterior temporal; Attn, attention; Front

Par, Frontoparietal; MP, medial prefrontal; MTN, medial temporal network; PM, posterior

medial; Som-Mot, Somatomotor. *indicates significant difference at p < .05.

To quantify the diversity of connections in the DMN subnetworks to the rest of the brain, we

calculated participation coefficients—a graph theory metric that describes the extent to which a

node's functional connections are spread across different communities. Nodes with a high

participation coefficient are thought to be critical for communication between communities, and

disruption of such nodes can lead to altered network function and cognitive impairment (59–62).

Of the subnetworks, the AT subnetwork had the greatest diversity of connections to nodes

outside of the DMN (AT > PM: $t(43) = 3$, $P =.004$; AT > MP: $t(32) = 2.7$, $P =.01$). As shown in

Figure 4B, the PM subnetwork had several nodes with relatively high participation. These nodes

connect to MTN, and the frontoparietal network, but the majority of the PM subnetwork showed

less out of network connectivity. The MP subnetwork, on the other hand, had relatively low

participation outside of the DMN.

266   We next addressed the extent to which nodes in the DMN showed diverse connectivity

with nodes in the other DMN subnetworks. Interestingly, regions in the MP subnetwork

exhibited the highest participation coefficients across the DMN subnetworks (MP > PM: $t(28) =$

2.7, $P = .01$; MP > AT: $t(35) = 3.7$, $P < .0007$). These findings substantiate what can be seen in

Figure 4A—that nodes in the MP subnetwork are positioned to integrate information across

DMN subnetworks, whereas nodes in the AT subnetwork and PM subnetwork may interface

with networks outside of the DMN.

Cortico-hippocampal network connectivity

Having identified cortical networks and subnetworks that connect to the hippocampus (cortico-

hippocampal networks), we then examined whether these networks showed a preference in

connectivity along the hippocampal long-axis. Previous research has shown differential

functional connectivity along the long-axis axis of the hippocampus (24, 28, 63, 64), with the

anterior hippocampus demonstrating greater connectivity to orbitofrontal cortex, and temporal

pole and the posterior hippocampus demonstrating greater connectivity to retrosplenial cortex

and precuneus. We therefore hypothesized that the anterior hippocampus should have

preferential connectivity to the AT subnetwork, whereas the posterior hippocampus would have

preferred connectivity with the PM subnetwork. These predictions were partially confirmed by

our findings. The anterior hippocampus showed stronger connectivity to the AT and MP

subnetworks $(t(37) = 8.2, P < .00001; t(37) = 6.5, P < .00001)$ and the posterior hippocampus

showed stronger connectivity to the MTN than the anterior hippocampus $(t(37) = 2.6, P = .01)$,

but there was no significant difference in connectivity between the hippocampal regions and PM

subnetwork $(t(37) = .41, P = .68)$ (Figure 5A). To determine whether these differences were

evenly spread across the networks or driven by particular regions within the networks, we

contrasted anterior and posterior hippocampal connectivity at a region-to-region level. As

expected from the network-level analysis, we observed significantly greater anterior

hippocampal connectivity to medial prefrontal, anterior temporal, and also anterior medial

temporal cortex, whereas the posterior hippocampus had significantly greater connectivity to the

parietal occipital sulcus, precuneus, and dorsal posterior cingulate cortex. Thus, the anterior-

posterior differences are fairly consistent in the AT and MP subnetworks, but within the MTN

the overall anterior-posterior differences are driven by relatively higher posterior hippocampal

connectivity in the medial parietal cortex (Figure 5B).

Based on neuropsychological evidence demonstrating distinct functional roles of the left

and right hippocampus (65–67), we next investigated whether the left and right hippocampus

differed in their connectivity to the cortical networks. We observed significant hemispheric

laterality effects with the right hippocampus having greater connectivity to the MTN than the left

($t(37) = 3.8$, $P = .0001$) and the left hippocampus having greater connectivity to AT subnetwork

($t(37) = 2.6$, $P = .01$). There was no significant laterality effect in the PM ($t(37) = .29$, $P = .77$)

and MP subnetworks ($t(37) = .98$, $P = .33$). This connectivity difference might help to explain

differentiated function of the left and right hippocampus.

A.

B.

Figure 5. Regions within the MTN and DMN subnetworks exhibit differential functional

connectivity with the anterior and posterior hippocampus. A: Bar graphs of functional

connectivity from each of the cortical networks to the anterior and posterior hippocampus. Error

bars show 95% confidence interval. B: T-contrast of anterior hippocampal connectivity versus

posterior hippocampal connectivity, with warm colors showing regions that had stronger anterior

hippocampal connectivity and cool colors showing regions that had stronger posterior

hippocampal connectivity. Cortico-hippocampal network boundaries are outlined on the surface

of the brain. These results indicate that the temporopolar and medial prefrontal cortex show

preferential connectivity with the anterior hippocampus, whereas posterior medial parietal cortex

shows preferential connectivity with the posterior hippocampus. Ant, anterior; AT, anterior

temporal; HC, hippocampus; MP, medial prefrontal; MTN, medial temporal network; PM,

posterior medial; Post, posterior. $* P < .05$, $*** P < .001$.

<u>Regions within the same community represent similar kinds of information during a memory</u>

<u>task</u>

We next sought to determine whether network membership was related to functional relevance

of these subnetworks using an independent dataset in which participants performed a memory

retrieval task (described in Mizrak et al. (68) and in *Methods* section) by examining the

representational profile similarity between regions within the same network or in different

networks (17, 69). Multivoxel pattern similarity was calculated between trials which created a

trial-by-trial representational similarity matrix for every ROI in our cortico-hippocampal

networks. Here, we assume that trial-by-trial fluctuations in pattern similarity for a given ROI

are driven by trial-by-trial fluctuations in features for which the ROI is sensitive. These

fluctuations across trials represent the ROI's *representational profile*. We sought to determine

whether regions within the same network had greater similarity in their representational

profiles—and thus represent similar features—compared to regions outside their network.

Here, we calculated the similarity of representational profiles between ROIs by

correlating each ROI's representational similarity matrix to every other ROI. We then averaged

the representational profile similarities for pairs of ROIs within the same network to create a

mean within-network similarity value, for each subject (we removed the similarity of each ROI

with itself to avoid inflating the within network similarity). To create a mean between-network

similarity value, for each subject, we averaged together the representational profile similarity

values for pairings of ROIs that were in different networks. We contrasted the mean within-

network similarity with the mean between-network similarity using a repeated-measures

ANOVA and observed that regions within the same network had higher profile similarity

compared to regions in different networks, $t(21) = 18.5$, $P < .001$ (Figure 6). This effect persists

even when accounting for the similarity in trial-by-trial mean BOLD activity ($t(21) = 12.1$, p <

.001), suggesting it is not simply a result of univariate activity similarity.

Figure 6. Regions show higher representational profile similarity with community members

compared to regions in other communities. Left: Individual subject representational profile

similarity of regions within the same network or between network in grey lines with the group

average in red, with 95% confidence intervals represented by error bars. Right: Boxplot of

average region-to-region representational profile similarity between each cortical-hippocampal

network (labelled along the x-axis) to itself and every other cortico-hippocampal network (each

represented by a colored bar) across the sample. Individual participants represented as dots. AT,

anterior temporal; MP, medial prefrontal; MTN, medial temporal network; PM, posterior medial.

Another possible explanation of these findings is that regions in the same network tend to be

closer in spatial proximity and there is some spatial non-independence of adjacent ROIs of these

representational effects. To account for this spatial proximity, we created an ROI-by-ROI

distance matrix by calculating the Euclidean distance between the center of gravity of all pairs of

ROIs. We then regressed out the influence of spatial proximity from the representational profile

matrix using this distance matrix and repeated the analysis. After statistically removing the

influence of spatial proximity, we still observed that ROIs within the same network had greater

representational profile similarity compared to pairs of ROIs in different networks, $t(21) = 9.3$, p

$< .001$.

<u>Neurosynth Decoding</u>

Having found evidence that regions within the same cortico-hippocampal networks carry similar

information, we sought to identify the cognitive terms typically associated with activity in these

networks. To better understand the role these networks may play in cognition, we used

Neurosynth (70) to determine what cognitive terms are typically associated with activation in

these subnetworks. The meta-analytic activation map of every term in the Neurosynth database

was correlated with the binarized volumetric masks of each cortico-hippocampal network. By

examining the top cognitive terms associated with each network we observed that the term

autobiographical was the sole common term in the top 20 across all networks. Activation in

MTN and PMN is disproportionately associated with studies involving episodic memory, but this

was not the case for the ATN and MPN in which the term episodic memory did not appear in the

top 20 search terms. The MTN also showed strong overlap with meta-analytic maps for

preference for navigation *(r = .15)*, scenes *(r = .12)*, and the PM subnetwork showed strong

overlap with self-referential *(r = .13)*, and theory of mind terms *(r = .08)*. The top terms

associated with AT subnetwork related to theory of mind *(r = .22)*, intentions *(r = .17)*, mental

states *(r = .22)*, and social cognition *(r = .19)*, while the top terms associated with MP

subnetwork were value *(r = .17)*, fear *(r = .16)*, emotion *(r = .16)* , and terms related to emotional

valence *(r = .14)*. Weightings for each network are available in Supplementary Table 2.

<u>Figure 7. Terms associated with activation of each cortico-hippocampal network.</u> Word clouds

are colored according to the network they describe which are displayed on the inflated brain. The

size of the terms in the word cloud relates to how strongly the meta-analytic activity of that term

correlates with the spatial extent of the network. Red, AT subnetwork; Blue, PM subnetwork;

Yellow, MP subnetwork; Green, Medial temporal network.

Discussion

We have taken a data-driven approach using resting-state fMRI to identify and characterize

cortico-hippocampal network connectivity, finding a set of subnetworks that interact with the

hippocampus. Our analyses showed that the MTN, a cluster of regions in the medial temporal

and dorsomedial parietal lobes, could be differentiated from the DMN, which included medial

and lateral parietal, temporal, and frontal regions. This MTN mediates connectivity between the

visual network and the DMN and between the visual network and hippocampus. We also found

that the DMN could be meaningfully subdivided into three subnetworks: the PM subnetwork,

encompassing the posterior cingulate, lateral parietal, and dorsal lateral prefrontal cortex, the AT

subnetwork, encompassing the temporopolar, lateral orbitofrontal, and dorsal medial prefrontal

cortex, and the MP subnetwork, encompassing the ventral medial prefrontal, and entorhinal

cortex. These cortico-hippocampal subnetworks vary in connectivity strength along the long-axis

of the hippocampus, and meaningfully segregate based on feature representation as measured by

multivoxel activity patterns in an independent dataset.

Prior work on the neurobiology of memory has drawn on the idea that the hippocampal

formation primarily interacts with the PHC and PRC, such that these areas collectively comprise

a "MTL memory system" (2) that is functionally distinct from surrounding cortical areas. More

recently, we and others proposed that PHC and PRC are embedded in larger scale cortico-

hippocampal networks as evidenced by divergent anatomical pathways found in rodents and non-

human primates (19–22) and by differing functional connectivity of the PHC and PRC in humans

(24, 25, 71). In this "PM/AT" framework, the PHC is a core region in the PM network, which

also includes retrosplenial, posterior cingulate and lateral parietal cortex, whereas the PRC is a

core region in the AT network, which also includes temporopolar, and orbitofrontal cortex (29).

As shown in Figure 8, the present results were not fully consistent with either the original PMAT

framework. We found that the PRC, PHC, retrosplenial cortex, inferior lateral parietal, and

dorsal medial parietal formed the MTN, and this network could be distinguished from the DMN.

This network corresponds closely to the "parieto-medial temporal pathway" described by Kravitz

et al. (72). This network division converges with previous data-driven community detection

studies (15, 42, 47, 73) showing that the MTN can be differentiated from the DMN and support

recent suggestions that heterogeneity exists within the PM system (38, 50). Critically, both the

MTN and DMN showed strong functional connectivity with the hippocampus.

Figure 8. Left. Hypothesized PM/AT divisions with PM nodes as blue and AT nodes as red (29,

74). Right. Data-driven network assignments.

Although the present work shows that the MTN could be differentiated from the DMN,

they do not imply that regions within the MTN are functionally or anatomically identical in

terms of connectivity. Prior anatomical studies in animal models (75, 76) and fMRI studies in

humans have shown that the PRC and PHC have different functional connectivity patterns (24–

26, 77). In fMRI studies, these differences are most apparent when the functional connectivity

maps of the PRC and PHC are directly contrasted to identify regions that show

disproportionately high connectivity to the PRC or the PHC. When not contrasted, the two

regions frequently show strong functional connectivity to each other and considerable overlap in

the spatial extent of their functional connectivity (24, 25, 77). For example, Wang et al (25) and

Zhuo et al. (77) showed that the PHC and posterior PRC show high connectivity with each other,

and with medial parietal, medial prefrontal, lateral parietal and temporopolar regions. Therefore,

the evidence presented here should not be taken to reflect the presence or absence of functional

distinctions between PHC and PRC (as prior work has clearly shown that certain regions connect

more to PHC or PRC), but rather they demonstrate that when taking into account the topology of

functional connections across the entire neocortex, the MTN can be segregated from the DMN.

This evidence suggests that heterogeneity exists within the previously reported PM network.

Treating the brain as an abstract graph, we observed that the MTN serves as a critical

bridge between the visual network and the DMN and hippocampus, such that removal of the

MTN results in considerable loss of cross-network connectivity between the visual network and

DMN/hippocampus. The medial temporal cortex is thought to provide high-level feature

representation of the external world to the hippocampus which then binds those representations

in context (4, 22, 29, 58). Damage to the medial temporal cortex is often associated with varying

level of circumscribed memory impairment. Our results here suggest that loss of MTN

connections would result in disconnection between the external visual world and the

hippocampus and DMN, which could result in memory impairment. Using a novel analytic

method, a recent study mapped the functional connectivity of regions in the brain where lesion

damage resulted in amnesia and found a common functional network to all the lesions (11). This

lesion functional connectivity map encompassed the DMN, MTN and encroached into visual

network regions as described in this study, underscoring the importance of these networks in

memory processing. Another study of amnesia following traumatic brain injury observed

reduced functional connectivity between the PHC and PCC was related to memory impairment

and normalized when memory functioning improved (78).

The distinction between the MTN and the DMN is consistent with recent studies

suggesting heterogeneity within the PM network (14, 38, 50, 79). Based on its network

properties, the MTN sits in a reasonable position to represent higher-order features of the visual

world to the DMN which is spatially distal from networks involved in primary sensation. Recent

theories of event cognition have suggested a separation between content (higher order

representations of current environment) and structure (abstracted knowledge pertaining the

typical way in which sequences of events unfold) (80). One possible instantiation of this

separation is that the MTN represents content of events whereas the DMN represents abstracted

structure. Indeed, a recent study has shown that when orienting to retrieval of perceptual features

during autobiographical recall, there is increased activity in the MTL, retrosplenial, and dorsal

precuneus (regions in the MTN) compared to when recall is oriented to conceptual or thematic

elements that preferentially resulted in activity of posterior cingulate, medial prefrontal, temporal

polar and lateral parietal cortex (regions in the DMN) (39). Another study found a similar

separation between the MTN and DMN by having participants encode a set of movie clips that

were considered typical (events that one would be familiar with through day to day life or

through movies/stories) or atypical (very unusual or dissimilar to anything you would encounter

in day to day life or in movie/stories) (40). Regions in the DMN showed increased activity

during encoding of typical compared to atypical events and the reverse was true for regions in

the MTN. During encoding of typical events, activity in the DMN was interpreted as being

related to retrieval of relevant prior event knowledge to assist in comprehension and integration

of incoming information. Conversely, during encoding of atypical events, the activity in the

MTN was interpreted to reflect the need to encode the relations between items, contexts and

actions in the absence of a strong semantic framework. While these studies present strong

evidence to suggest a distinction between the MTN and DMN, it is important to note that in

many circumstances involving memory recall, these networks will be working in tandem to

support retrieval of both content and structure with some degree of bias that may favor retrieval

of item-level, scene-level or event-level elements. For example, one might expect that, even

within the MTN, a contrast that focusses on the distinction between different types of content,

such as scenes versus objects, may split regions within the MTN that have a bias towards those

content types (35). These findings demonstrate the necessity of our intrinsic connectivity

analyses to clarify that these differences in patterns of functional activity are likely due to the

presence of distinct functional networks.

The second major finding in the present study is that the DMN could be differentiated

into three subnetworks: the PM, AT, and MP subnetworks. The PM subnetwork consisted of

posterior cingulate, angular gyrus, and dorsal prefrontal cortex, whereas the AT subnetwork

consisted on temporopolar, lateral temporal, orbitofrontal and dorsal medial prefrontal cortex.

The collection of regions in the PM and AT subnetwork are in line with the PM/AT framework

(24, 25), with the notable exception that the parahippocampal, retrosplenial cortex were not in

the PM subnetwork, and the perirhinal cortex was not in the AT subnetwork. The PM and AT

subnetworks also resemble DMN subnetworks identified by previous data-driven methods (15,

41, 42), confirming that the PM and AT distinctions theorized do indeed persist, but they form

distinct communities from their MTN counterparts when examining their connectivity

architecture.

The MP subnetwork consists of medial prefrontal cortex, along with the entorhinal cortex

and was not explicitly predicted by prior theories. The medial prefrontal cortex has long been

distinguished from the orbitofrontal cortex based on anatomical connectivity (21, 22, 81), and is

often grouped together with the posterior cingulate cortex as one of the "core hubs" of the DMN

(15, 41, 42). The entorhinal cortex certainly has dense interconnections with the medial

prefrontal cortex (81), but also is strongly interconnected with the hippocampus and adjacent

medial temporal cortex (19, 22, 82). Interestingly, the entorhinal cortex often is part of the DMN

even when the MTL cortex is distinguished from the rest of the DMN ((73, 83), but see (15, 47)).

Furthermore, network analysis on histologically-defined axonal connections between cortical

regions in rodents, showed that the lateral entorhinal cortex was part of a community with the

prefrontal cortex, consistent with our partition, though the medial entorhinal cortex was placed in

a community with the hippocampus, retrosplenial cortex and subiculum (84). A distinction in the

functional connectivity of entorhinal cortex subregions has been shown in high-resolution human

resting state data (23, 37), but the spatial resolution of the current study is unable to resolve these

subregions and future work will be important to further characterize its connectivity in humans.

We examined the topology of how these subnetworks connect to each other and the rest

of the brain finding that the MP subnetwork had the highest within-DMN participation

coefficient, with nodes connecting broadly to AT and PM regions. Conversely, the MP

subnetwork had a low out of DMN participation coefficient, especially relative to the AT

subnetwork. Our results provide converging evidence for a recent study published by Gordon et

al (85) that also found that a set of lateral networks resembling the AT subnetwork here, showed

the highest participation coefficient out of network. Their AT equivalent showed

intercommunication with language and frontoparietal networks, as was also seen here. The

participation coefficient findings suggest that the AT subnetwork and several high participation

PM nodes are positioned to mediate information transfer in and out of the DMN, whereas the MP

subnetwork may integrate and coordinate information within the DMN. A recent review of the

human lesion literature has reported that robust memory impairment can be observed following

medial prefrontal cortex damage and suggested that this may be due to the fact that the medial

prefrontal cortex has a role in initiating and coordinating cognitive processes for retrieval and

episodic simulation (86). Given the network findings here, and implications from the lesion

literature, the MP subnetwork is well positioned for higher-order integration and coordination of

memory-guided activity.

Differentiation of the three DMN subnetworks was reinforced by evidence from an

independent dataset in which we found that regions grouped in the same network represented

similar kinds of information, as compared to regions in different networks. These findings

replicate and extend results from previous studies that identified different functional differences

between different cortico-hippocampal networks (17, 69). For instance, Ritchey et al. (17) used

community detection to differentiate networks of regions hypothesized to show different patterns

of MTL connectivity. Ritchey et al. (17) interrogated these networks in an independent study of

episodic memory retrieval, as did Inhoff et al. (69) during a concept learning task. The functional

distinction of these networks was supported by findings showing that regions grouped in the

same network had greater representational profile similarity (17) and activation profile similarity

than regions grouped in different networks during learning and memory tasks (17, 69). In these

past studies, the regions used in network construction were those that showed a significant

difference in functional connectivity between the PRC and PHC, being motivated by anatomical

evidence suggesting parallel pathways in the MTL (19). Whereas the seed selection that defined

these networks was based specifically on regions that had different functional connectivity

between PHC and PRC, in the current study, we did not impose any pre-existing assumptions,

and identified networks holistically in the context of the entire brain. We saw reliably higher

representational profile similarity for regions in the same network—a pattern consistently found

in every subject, highlighting the functional significance of these cortico-hippocampal networks

and validating our delineation of these cortico-hippocampal networks.

We also found that the MTN and DMN subnetworks show variation in their connectivity

across the anterior and posterior hippocampus. The hippocampus is known to vary along its long

axis in terms of intrinsic connectivity, extrinsic connectivity, receptor distribution, and gene

expression (19, 20, 87–90). In humans, resting state functional connectivity has been the primary

tool for examining differences in connectivity along the hippocampal longitudinal axis, though

distinctions between the anterior and posterior hippocampus have been observed using DTI (28).

The anterior hippocampus has been shown to have greater connectivity to inferior temporal,

temporopolar, orbitofrontal, medial prefrontal and PRC, whereas the posterior hippocampus has

been shown to have greater connectivity to PHC, retrosplenial cortex, inferior parietal cortex,

precuneus, the anterior thalamus (24, 26–28, 63). Here, we replicated and extended these

findings to characterize hippocampal connectivity to the networks formed by these cortical

regions. We found that the anterior hippocampus has relatively greater connectivity to the MP

and AT subnetworks, whereas the posterior hippocampus has relatively greater connectivity to

the MTN. The preferential connectivity to the anterior hippocampus was relatively consistent

across all regions in the MP and AT subnetworks, but there was heterogeneity in the MTN's

preference for the anterior and posterior hippocampus with the anterior hippocampus having

stronger connectivity to anterior MTL cortex and the posterior hippocampus having relatively

stronger connectivity to dorsal medial parietal cortex. Theories regarding the functional

difference of the anterior and posterior hippocampus are often informed by preferential long-axis

connectivity to the rest of the brain (16, 91), but they do not consider how the rest of the brain

connects to form networks. The results presented here suggest the possibility that functional

differences between the anterior and posterior hippocampus may reflect differences in their

connections to the subnetworks of the DMN.

To better understand the functional specializations of these networks, we used meta-

analytic maps from the Neurosynth database to identify the cognitive constructs that have been

reliably associated with regions in the DMN subnetworks. Activity within the MTN was

associated with a set of terms heavily rooted in episodic memory and navigation. Activation in

the MP subnetwork, in contrast, was highly associated with terms pertaining to emotion, and

value. As noted above, the MTN (in particular the dorsal precuneus and dorsal posterior

cingulate) had greater connectivity to the posterior relative to the anterior hippocampus, whereas

the MP subnetwork had greater connectivity with the anterior compared to posterior

hippocampus. These connectivity patterns and neural decoding results converge with theories

linking the posterior hippocampus to cognitive and navigational processes and the anterior

hippocampus in motivation and emotional behavior (92). We also found that activation in the AT

subnetwork was associated with terms that relate to social cognition, theory of mind, and beliefs,

in line with contemporary theories (93) and lesion evidence (94) of this network. In the PM

subnetwork, Neurosynth decoding revealed cognitive terms related to episodic memory and the

self. This is echoed by recent work, taking a highly subject-specific approach, that found a

distinction within the DMN between regions involved in episodic versus theory of mind-like

processes (95). In this study DiNicola et al (95) identified two distinct subnetworks of the DMN

on a subject-by-subject, voxel-wise basis. One subnetwork resembled the AT subnetwork and the

other resembled the PM subnetwork described here. The same subjects also performed both

episodic projection tasks and theory of mind tasks and it was observed that the voxels classified

as being in the AT-like network showed greater activity for the theory of mind tasks, whereas the

voxels classified as being in the PM-like network showed greater activity for the episodic

projection tasks. Interestingly, the Neurosynth analysis showed that the term "autobiographical"

was strongly associated with all four cortico-hippocampal networks. This broad association may

be due to the fact that autobiographical memory tasks usually involve retrieval of episodically

rich events that dynamically unfold involving social interactions, and relevance to the self (96).

Although the Neurosynth decoding analysis is exploratory, and by definition, post-hoc,

they can guide the generation of hypotheses to be tested in future studies. Further studies, similar

to the work of DiNicola et al (95), should examine whether MTN and DMN subnetworks

segregate based on spatial, valuation, and social processes, within the same group of individuals.

For example, during encoding and recall of complex events, we might expect regions within the

MTN to represent high-level object and context features, regions within the AT subnetwork to

represent interpretations of theory of mind and social interactions, and regions within the MP

subnetwork to represent the emotional valence or perceived value of the event. Indeed, a recent

preprint demonstrated a dissociation between the MTN and MP subnetwork during event

simulation, with increased activity in the MTN occurring when subjects were instructed to

simulate past or future events using cues that were selected to elicit high levels of vividness,

whereas increased MP activity was observed when cues elicited simulations with high levels of

valence (97). Further, based on the functional connectivity of the MP subnetwork, we

hypothesize that it would be well suited to serve a coordinating role in network rearrangement to

accomplish task goals and demands. Again, a recent study by Nawa and Ando (98) found that

when elaborating on autobiographical memories, the vmPFC drove hippocampal activity and this

effect was augmented for memories that were of greater emotional valence. Further, damage to

the medial prefrontal cortex is associated with memory impairments characterized by difficulties

in elaborating on memories unfold over time and events (86). We hypothesize that these

impairments may be associated with dysfunction of network dynamics due to a loss of the

coordination normally provided by the MP subnetwork.

Finally, we note that recent studies have demonstrated the value in taking subject-specific

approaches to delineating functional networks in a voxel-wise fashion (14, 47). This work has

shown that there are individual differences in the location of boundaries between functional

networks, though it is notable that the majority of the cortex is labelled consistently in the

majority of participants (73). A key goal of the present study was to provide a clear framework to

guide analyses in future task-fMRI studies, in which it might not be feasible to obtain

independent, subject-specific parcellations. We have listed each region and its community label

in Supplementary Table 3, and the HCP-MMP atlas is available online

(https://balsa.wustl.edu/study/show/RVVG, (44)). Task fMRI studies that use group-level

analyses can use the group-level characterization of cortico-hippocampal networks reported here

in order to rigorously test hypotheses about the functions of these networks and how they may be

implicated in memory disorders, as resting connectivity has shown to predict spread of pathology

in Alzheimer's (99), and DMN subnetwork perturbation has been demonstrated in MTL amnesia

(100).

<u>Conclusions</u>

The hippocampus affiliates with a broad set of regions that enable episodic retrieval. Here, we

have shown that the MTN and three subnetworks of the DMN can be differentiated on the basis

of their whole-brain functional connectivity. The three DMN subnetworks vary in connectivity

along the hippocampal long axis, have distinct representational roles during a memory task, and

topologically are connected by a set of hubs in the medial prefrontal subnetwork. This

subnetwork organization offers a novel framework to investigate event cognition and memory

retrieval.

Methods

<u>Participants</u>

Forty-five healthy, young adult participants were recruited from the University of California,

Davis and surrounding area ($N_{Females}$ = 26, mean age = 25.6 years [SD = 4.2 years]). All

participants were right-handed and neurologically healthy. The study was approved by the

Institutional Review Board of the University of California at Davis and all participants provided

informed consent prior to participation. Participants were compensated $20/hour for their time.

This sample size is comparable to the cohort sample sizes from the seminal Power et al. (101)

study investigating functional brain organization. There is no specific effect size that can be

taken to approximate power for identifying network communities, but previous studies have

shown high reliability across subjects in network identification when using greater than 10

minutes of resting-state data (73, 102).

<u>MRI Acquisition</u>

MRI scanning was performed using a 3T Siemens Skyra scanner system with a 32-channel head

coil. A T1-weighted structural images was acquired using a magnetization prepared rapid

acquisition gradient-echo pulse sequence (TR = 2100ms; TE = 2.98ms; field of view = 256 mm^2

; flip angle = 7°; image matrix = 256 × 256, 208 axial slices with 1.0 mm^3 thickness; GRAPPA

acceleration factor 2 with 24 reference lines). Functional images were acquired using a gradient

EPI sequence (TR = 1220 ms; TE = 24 ms; field of view = 192 mm^2 ; image matrix = 64 × 64;

flip angle = 67°; bandwidth = 2442 Hx/pixel; multiband factor = 2; 38 interleaved axial slices,

voxel size = 3 mm^3 isotropic). Five runs of five minutes in duration were acquired for rest scans

for a total of 25 minutes of resting state fMRI data per subject. Participants were instructed to lay

as still as possible and try not to fall asleep.

<u>Anatomical Preprocessing</u>

T1-weighted anatomical scans were preprocessed using FreeSurfer which included intensity

normalization, removal of non-brain tissue, transformation to Talaraich space, and segmentation

of gray matter, white matter, and CSF. Surfaces were calculated for the white matter-gray matter,

and gray matter-pial interface. Surface-based registration to the HCP-MMP1.0 atlas (44) was

performed and subject-specific cortical regions were calculated according to atlas boundaries.

Surface-based cortical regions were converted to volumetric regions of interest and transformed

into functional native space. The hippocampus was segmented in FreeSurfer in an automated

fashion. Manual adjustments were done to correct misclassified voxels and the hippocampus was

divided into anterior and posterior segments based off of the disappearance of the uncal notch

(16), with the posterior hippocampus designated as all of the hippocampus posterior to the

disappearance of the uncal notch on a coronal slice.

<u>Functional MRI Preprocessing</u>

Functional preprocessing was performed using *fMRIPrep* 1.4.1 ((103, 104),

RRID:SCR_016216), which is based on *Nipype* 1.2.0 ((105, 106), RRID:SCR_002502). For each

of the 5 BOLD runs found per subject (across all tasks and sessions), the following preprocessing

was performed. First, a reference volume and its skull-stripped version were generated using a

custom methodology of *fMRIPrep*. A deformation field to correct for susceptibility distortions

was estimated based on two echo-planar imaging (EPI) references with opposing phase-encoding

directions, using 3dQwarp from Cox and Hyde (107) (AFNI 20160207). Based on the estimated

susceptibility distortion, an unwarped BOLD reference was calculated for a more accurate co-

registration with the anatomical reference. The BOLD reference was then co-registered to the

T1w reference using flirt (FSL 5.0.9, (108)) with the boundary-based registration (109) cost-

function. Co-registration was configured with nine degrees of freedom to account for distortions

remaining in the BOLD reference. Head-motion parameters with respect to the BOLD reference

(transformation matrices, and six corresponding rotation and translation parameters) are

estimated before any spatiotemporal filtering using mcflirt (FSL 5.0.9, (110)). The BOLD time-

series were resampled onto their original, native space by applying a single, composite transform

to correct for head-motion and susceptibility distortions. These resampled BOLD time-series will

be referred to as *preprocessed* BOLD. Several confounding time-series were calculated based on

the *preprocessed BOLD* as part of *fMRIPrep*, however, only the head-motion estimates

calculated in the correction step and frame displacement (FD) were used in subsequent analyses.

All resamplings can be performed with *a single interpolation step* by composing all the pertinent

transformations (i.e. head-motion transform matrices, susceptibility distortion correction when

available, and co-registrations to anatomical space). Gridded (volumetric) resamplings were

performed using antsApplyTransforms (ANTs), configured with Lanczos interpolation to

minimize the smoothing effects of other kernels (111). Many internal operations

of *fMRIPrep* use *Nilearn* 0.5.2 ((112), RRID:SCR_001362), mostly within the functional

processing workflow. For more details of the pipeline, see

https://fmriprep.readthedocs.io/en/stable/workflows.html.

The preprocessed BOLD timeseries, anatomical images, and native space Glasser parcels

were imported into CONN Toolbox version 18b (www.nitrc.org/projects/conn,

RRID:SCR_009550). Based on Ciric et al.'s (113) assessment of 14 common denoising

protocols, we selected the '9p' protocol because it was shown to facilitate the highest functional

network identifiability (see (113) Figure 5b). BOLD timeseries were demeaned, linear and

quadratic trends were removed, and bandpass filtered between .008 and .09 hz. The 9p protocol

includes as confound regressors six motion parameters, WM, CSF, and global signal regression.

All regressors were band-pass filtered to maintain the same frequency range as the data. Visual

inspection of the frequency distributions of functional connectivity values as well as quality

control plots of the correlation between mean FD and functional connectivity values was

performed to identify any aberrant runs or subjects. Five outlier subjects were identified which

each corresponded to previously identified subjects having a mean FD value of greater than

.15mm. These subjects were excluded from further analyses. Functional connectivity matrices

were created for each subject by computing Pearson's correlations between all possible pairs of

each regions' confound-corrected timeseries. Finally, each correlation value was Fisher z-

transformed with the inverse hyperbolic tangent function.

<u>Community Detection</u>

Cortical functional connectivity (FC) matrices from all subjects were thresholded to exclude

negative connections that may be introduced by global signal regression (114, 115) and then

averaged together. Using the resulting group averaged connectivity matrix, community detection

was performed using the Louvain method (45) via the brain connectivity toolbox

(https://sites.google.com/site/bctnet/) which iteratively performs a greedy optimization of

modularity by randomly selecting nodes and merging them into the community that maximally

increases modularity, until no more gains in modularity are possible. Modularity (Q) describes

how well a network can be divided into communities that have higher within-community

connectivity than would be expected by chance and can range from -1 to 1, with negative values

indicating fewer intracommunity connections than would be expected by chance and positive

values indicating higher intracommunity connections than would be expected by chance (56).

We used a method of the Louvain community detection algorithm adapted to accommodate

weighted graphs (116) which accepts the weighted, group average functional connectivity

matrix. This algorithm can be tuned using a resolution parameter, gamma, that biases the

algorithm towards producing few, large networks (low gamma) or towards many, smaller

networks (high gamma). To determine the resolution parameter in a principled manner, we

adopted the approach used by Ji et al. (43). The criteria included: i) separation of the primary

sensory-motor networks (visual, auditory, somatomotor) from all other networks, ii) high

stability across nearby parameters (similar network partitions across neighboring parameter

settings), and iii) optimized modularity.

754   Given that the Louvain algorithm detection is influenced by a random starting point, for

every tested resolution parameter, we ran the algorithm 1000 times. To determine the optimal

partitioning solution at each resolution, gamma, we examined how consistent a given partitioning

solution was to every other solution produced over the 1000 iterations at the same resolution.

Consistency was calculated using the z-Rand score (46) (http://commdetect.weebly.com/). For

each partition solution, we weighted the mean z-Rand score (consistency) of the partition with

the modularity value of that partition to select a partition solution that was both highly consistent

at a given resolution and had high modularity. These methods are in keeping with Ji et al. (43).

Using the default parameter, gamma = 1, we could not satisfy criteria i) as the somatomotor and

auditory network were grouped together (also seen in Ji et al. (43)). Thus, we ran a parameter

sweep from gamma 1 – 2.8 by increments of 0.005 until we identified a resolution parameter at

which the somatomotor and auditory network separated consistently and produced the highest

modularity-weighted z-Rand score.

Path Length Analysis

Using the brain connectivity toolbox (https://sites.google.com/site/bctnet/), weighted path length

was calculated between all nodes in the brain for each subject, after excluding negative

connectivity weights. Path length is the fewest number of links that need to be traversed to

connect two nodes. To calculate the weighted path length, the functional connectivity weights

were inverted creating a distance matrix in which nodes with high functional connectivity had

low distance. Then, the shortest path between each pair of nodes was calculated providing a

value that represented the minimal the total weighted distance that needed to be traversed to

connect two nodes.

We hypothesized that removal of the MTN would reduce the ability of the visual network

and DMN/hippocampus to connect with each other. To test this hypothesis, all the nodes from

the MTN were removed from each subject's network and the average weighted path length was

calculated between visual network and DMN and between visual network and hippocampus.

Because removal of nodes can only lead to an increase in path length (shorter paths can be

removed, but will never be added following removal of nodes), we compared the weighted path

length following removal of the MTN to the weighted path length following removal of every

other network, using paired t-tests. As a secondary control, we also compared the weighted path

length following removal of the MTN to the distribution of weighted path lengths following

removal of 30 random nodes (other than the DMN, visual network and hippocampus) over 1000

iterations. For this, we calculated the z-score of the weighted path length following removal of

the MTN compared to the distribution of weighted path lengths following removal of 30 nodes

over 1000 iterations. This was done for each subject such that each subject had a corresponding

z-score. A one-sample t-test was then performed on the z-score values to examine whether they

were significantly greater than 0.

Community Partitioning of the Default Mode Network

From the identified whole brain partitioning, we selected the default mode network based on

visual similarity to previously identified DMN solutions (42, 43, 117). We repeated the

community detection procedure described above on the identified DMN (1000 iterations of

Louvain community detection at each resolution parameter) and calculated modularity-weighted

z-rand scores across a range of resolution parameters, 0.75 – 1.1 at increments of 0.05. We

selected local modularity-weighted z-rand peaks as solutions for further exploration.

Connection Diversity in the Default Mode Network

The participation coefficient, which quantifies the degree to which a node is connected to a

diverse set of communities, was calculated using the tools from the brain connectivity toolbox

(https://sites.google.com/site/bctnet/). Since the participation coefficient calculation uses node

strength (the sum of the connectivity weights across the network) in the denominator of its

calculation, nodes with unusually low strength can produce unstably large participation

coefficient values. We, thus, excluded regions with node strength in the bottom 25% of the

network hubs. Participation coefficient calculations were performed across a range of network

densities from 5-20% at 1% intervals. These methods for calculating participation coefficient are

in line with best practices in the literature (57, 59, 62, 118, 119). Whole network participation

coefficient was calculated for each DMN region, using the whole brain connectivity matrix and

community labels generated from the whole-brain Louvain community detection. Within DMN

participation coefficient was also calculated for each DMN node, using only the connections

within the DMN and using the DMN subnetwork labels. Participation coefficient values were

compared between DMN subnetworks using a within-subjects mixed model using mixedlm from

statsmodels in Python 3 (120).

Hippocampal-cortical network connectivity

To identify hippocampal-cortical networks we calculated functional connectivity between each

hippocampal ROI and every cortical ROI in the brain. To identify average connectivity of the

hippocampus to a given network within-subject, functional connectivity weights were averaged

together across cortical nodes, based on community affiliation. A one-sample t-test was

performed to determine what networks were, on average, significantly connected to the

hippocampus. Significance was set at $p<.05$, Bonferroni corrected. We examined the

connectivity of the resulting significant networks and subnetworks to the hippocampus as a

function of the long-axis (anterior versus posterior) and hemisphere (left versus right) in a

within-subjects model as described above.

ROI level functional connectivity difference between the anterior and posterior

hippocampus were performed in CONN toolbox version 18b. Using the cortical ROIs that were a

part of large-scale networks connected to the hippocampus, a within-subjects analysis was

performed which contrasted functional connectivity in the left and right anterior hippocampus

against the left and right posterior hippocampus. Significance was set at $p<.05$, *FWE* corrected.

<u>Representational Similarity Analysis</u>

Using an independent dataset collected during a memory-based decision-making task (68), we

examined the representational profile of each ROI in the DMN subnetworks (See SI Appendix

for a detailed description of the task). If the hippocampal-cortical networks identified using

resting-state functional connectivity are functionally meaningful, then regions within those

networks should have similar representational profiles.

In brief, 22 participants in this dataset viewed a set of 8 grocery items one at a time, in a

pre-scanning session, and learned through trial and error whether each grocery item was

desirable to a hypothetical customer within a store context or not. In each trial, a grocery item

appeared in the context of a grocery store. There were four different grocery stores and for half

of the items the particular grocery store modulated their desirability and for the other half, the

grocery store context was irrelevant to their desirability. For example, the apple at grocery store

'A' may be desirable, but the apple at grocery store 'B' may be undesirable which makes apple's

desirability context-dependent. Alternatively, desirability of a carton of milk may be the same

regardless or grocery store which makes it context-invariant. Overall preference averaged to 50%

desirable for both context dependent and invariant. Further, of the four contexts, 50% shared the

same preference rules. Following successful learning, participants were put in the scanner and

shown the context and a food item, one at a time and asked to remember whether the food item

was desirable to the hypothetical customer at the store context based on their learning phase.

Feedback was not provided during scanning. In the original study, Mizrak et al. (68) examined

representational similarity between memory-based decisions depending on a) shared features

between grocery items such as being modulated by the context or not, b) shared features between

store contexts such as having similar desirability for the same grocery items or not.

Here, using the RSA toolbox (https://www.mrc-cbu.cam.ac.uk/methods-and-

[resources/toolboxes/license/](https://www.mrc-cbu.cam.ac.uk/methods-and-resources/toolboxes/license/), (121)), pattern similarity was calculated between memory based

decision making trials for each ROI using Pearson's correlation, excluding a) trials that occurred

during the same scanning run, and b) incorrect trials. Thus, fluctuations in pattern similarity

would be driven by variations in shared features. Correlations were then calculated between

ROIs to calculate the similarity of their representational profiles. We then examined the

representational similarity within cortico-hippocampal subnetworks versus between using a

paired t-test to examine whether these networks identified using resting-state functional

connectivity did indeed create functionally relevant communities.

To account for univariate effects on the multivariate analysis, we further calculated the

mean BOLD activity within an ROI for each trial and correlated the trial-wise mean BOLD

activity for each pair of ROIs to create an ROI-by-ROI activation similarity matrix. Using the

activation similarity matrix, we regressed out the influence of univariate activity from the RDM

similarity matrix and re-ran the statistical analysis contrasting within- vs. between-network RDM

similarity.

To account for the effects of spatial proximity between ROIs, we calculated the

Euclidean distance between the centroids of each pair of ROIs to create an ROI-by-ROI distance

matrix. Using the distance matrix, we regressed out the influence of spatial proximity from the

RDM similarity matrix and re-ran the statistical analysis contrasting within- vs. between-network

RDM similarity.

Cognitive characterization of hippocampal-cortical networks

We decoded cortico-hippocampal networks to determine what terms are most frequently

descriptive of the spatial distribution of the networks using the repository data version 0.7

(https://github.com/neurosynth/neurosynth-data) of Neurosynth

(https://github.com/neurosynth/neurosynth) (70). The meta-analytic activation of every term in

the Neurosynth database was correlated with the binarized volumetric masks of the hippocampal-

cortical networks. For each hippocampal-cortical network we examined the top terms whose

meta-analytic activation correlated most strongly with the spatial layout of the hippocampal-

cortical network. We removed terms that related to anatomy (e.g. ventromedial prefrontal

cortex), network (e.g. default mode), technique (e.g. independent component) and report terms

associated with cognitive functioning (e.g. navigation). We also removed terms that were

redundant with other terms (e.g. "autobiographic" and "autobiographical memory"). When

removing redundant terms, the term that was more specific was retained (e.g. "autobiographical

memory" retained in favor of "memory"), otherwise the term with the higher correlation to the

network was retained. Word cloud visualizations were made using wordcloud 1.7.0

(https://pypi.org/project/wordcloud/).

References

1.   E. Tulving, Memory and consciousness. *Can. Psychol.* **26**, 1–12 (1985).

2.   L. R. Squire, S. Zola-Morgan, The medial temporal lobe memory system. *Science* **253**,

1380–6 (1991).

3.   H. Eichenbaum, A. P. Yonelinas, C. Ranganath, The medial temporal lobe and recognition

memory. *Annu. Rev. Neurosci.* **30**, 123–52 (2007).

4.   L. Davachi, Item, context and relational episodic encoding in humans. *Curr. Opin.*

Neurobiol. **16**, 693–700 (2006).

5.   T. J. Teyler, P. DiScenna, The Hippocampal Memory Indexing Theory. *Behav. Neurosci.*

100, 147–154 (1986).

6.   M. D. Rugg, K. L. Vilberg, Brain networks underlying episodic memory retrieval. *Curr.*

Opin. Neurobiol. **23**, 255–260 (2013).

7.   C. McCormick, M. Moscovitch, A. B. Protzner, C. G. Huber, M. P. McAndrews,

Hippocampal-neocortical networks differ during encoding and retrieval of relational

memory: Functional and effective connectivity analyses. *Neuropsychologia* **48**, 3272–

3281 (2010).

8.   C. Sestieri, M. Corbetta, G. L. Romani, G. L. Shulman, Episodic memory retrieval,

parietal cortex, and the default mode network: Functional and topographic analyses. *J.*

Neurosci. **31**, 4407–4420 (2011).

9.   R. A. Cooper, M. Ritchey, Cortico-hippocampal network connections support the

multidimensional quality of episodic memory. *Elife* **8**, 1–22 (2019).

10.   B. Bellana, Z. X. Liu, N. B. Diamond, C. L. Grady, M. Moscovitch, Similarities and

917   differences in the default mode network across rest, retrieval, and future imagining. *Hum.*

918   *Brain Mapp.* **38**, 1155–1171 (2017).

919   11.   M. A. Ferguson, *et al.*, A human memory circuit derived from brain lesions causing

920   amnesia. *Nat. Commun.* **10**, 1–9 (2019).

921   12.   M. D. Greicius, B. Krasnow, A. L. Reiss, V. Menon, Functional connectivity in the resting

922   brain: A network analysis of the default mode hypothesis. *Proc. Natl. Acad. Sci. U. S. A.*

923   **100**, 253–258 (2003).

924   13.   J. R. Andrews-Hanna, J. Smallwood, R. N. Spreng, The default network and self-

925   generated thought: Component processes, dynamic control, and clinical relevance. *Ann. N.*

926   *Y. Acad. Sci.* **1316** (2014).

927   14.   R. M. Braga, K. R. A. Van Dijk, J. R. Polimeni, M. C. Eldaief, R. L. Buckner, Parallel

928   distributed networks resolved at high resolution reveal close juxtaposition of distinct

929   regions. *J. Neurophysiol.* **121**, 1513–1534 (2019).

930   15.   T. J. Akiki, C. G. Abdallah, Determining the Hierarchical Architecture of the Human

931   Brain Using Subject-Level Clustering of Functional Networks. *Sci. Rep.* **9**, 19290 (2019).

932   16.   J. Poppenk, H. R. Evensmoen, M. Moscovitch, L. Nadel, Long-axis specialization of the

933   human hippocampus. *Trends Cogn. Sci.* **17**, 230–40 (2013).

934   17.   M. Ritchey, A. P. Yonelinas, C. Ranganath, Functional connectivity relationships predict

935   similarities in task activation and pattern information during associative memory

936   encoding. *J. Cogn. Neurosci.* **26**, 1085–99 (2014).

937   18.   M. J. Sekeres, G. Winocur, M. Moscovitch, The hippocampus and related neocortical

938   structures in memory transformation. *Neurosci. Lett.* **680**, 39–53 (2018).

19.   R. D. Burwell, The parahippocampal region: corticocortical connectivity. *Ann. N. Y. Acad.*

Sci. **911**, 25–42 (2000).

20.   B. A. Strange, M. P. Witter, E. S. Lein, E. I. Moser, Functional organization of the

hippocampal longitudinal axis. *Nat. Rev. Neurosci.* **15**, 655–669 (2014).

21.   H. Kondo, K. S. Saleem, J. L. Price, Differential connections of the perirhinal and

parahippocampal cortex with the orbital and medial prefrontal networks in macaque

monkeys. *J. Comp. Neurol.* **493**, 479–509 (2005).

22.   J. P. Aggleton, Multiple anatomical systems embedded within the primate medial

temporal lobe: Implications for hippocampal function. *Neurosci. Biobehav. Rev.* **36**,

1579–1596 (2012).

23.   A. Maass, D. Berron, L. A. Libby, C. Ranganath, E. Düzel, Functional subregions of the

human entorhinal cortex. *Elife* **4**, 1–20 (2015).

24.   L. A. Libby, A. D. Ekstrom, J. D. Ragland, C. Ranganath, Differential Connectivity of

Perirhinal and Parahippocampal Cortices within Human Hippocampal Subregions

Revealed by High-Resolution Functional Imaging. *J. Neurosci.* **32**, 6550–6560 (2012).

25.   S. Wang, M. Ritchey, L. a Libby, C. Ranganath, Functional connectivity based

parcellation of the human medial temporal lobe. *Neurobiol. Learn. Mem.* **134**, 123–134

(2016).

26.   I. Kahn, J. R. Andrews-Hanna, J. L. Vincent, A. Z. Snyder, R. L. Buckner, Distinct

cortical anatomy linked to subregions of the medial temporal lobe revealed by intrinsic

functional connectivity. *J. Neurophysiol.* **100**, 129–39 (2008).

27.   J. Poppenk, M. Moscovitch, A hippocampal marker of recollection memory ability among

healthy young adults: contributions of posterior and anterior segments. *Neuron* **72**, 931–7

(2011).

28.   A. Adnan, *et al.*, Distinct hippocampal functional networks revealed by tractography-

964   based parcellation. *Brain Struct. Funct.* **221**, 2999–3012 (2016).

965   29.   C. Ranganath, M. Ritchey, Two cortical systems for memory-guided behaviour. *Nat. Rev.*

966   *Neurosci.* **13**, 713–726 (2012).

967   30.   L. A. Libby, Z. M. Reagh, N. R. Bouffard, J. D. Ragland, C. Ranganath, The

968   Hippocampus Generalizes across Memories that Share Item and Context Information. *J.*

969   *Cogn. Neurosci.* **31**, 24–35 (2019).

970   31.   M. Ritchey, M. E. Montchal, A. P. Yonelinas, C. Ranganath, Delay-dependent

971   contributions of medial temporal lobe regions to episodic memory retrieval. *Elife* **2015**, 1–

972   19 (2015).

973   32.   L. T. Hsieh, M. J. Gruber, L. J. Jenkins, C. Ranganath, Hippocampal Activity Patterns

974   Carry Information about Objects in Temporal Context. *Neuron* **81**, 1165–1178 (2014).

975   33.   S. Kim, *et al.*, Selective and coherent activity increases due to stimulation indicate

976   functional distinctions between episodic memory networks. *Sci. Adv.* **4**, 1–10 (2018).

977   34.   K. N. Warren, M. S. Hermiller, A. S. Nilakantan, J. L. Voss, Stimulating the Hippocampal

978   posteriormedial network enhances task-dependent connectivity and memory. *Elife* **8**, 1–21

979   (2019).

980   35.   A. Maass, *et al.*, Alzheimer's pathology targets distinct memory networks in the ageing

981   brain. *Brain* **142**, 2492–2509 (2019).

982   36.   D. Berron, D. van Westen, R. Ossenkoppele, O. Strandberg, O. Hansson, Medial temporal

983   lobe connectivity and its associations with cognition in early Alzheimer's disease. *Brain*

984   **143**, 1233–1248 (2020).

37.   J. N. Adams, A. Maass, T. M. Harrison, S. L. Baker, W. J. Jagust, Cortical tau deposition

follows patterns of entorhinal functional connectivity in aging. *Elife* **8**, 1–22 (2019).

38.   M. Ritchey, R. A. Cooper, Deconstructing the Posterior Medial Episodic Network. *Trends*

Cogn. Sci., 1–15 (2020).

39.   L. Gurguryan, S. Sheldon, Retrieval orientation alters neural activity during

autobiographical memory recollection. *Neuroimage* **199**, 534–544 (2019).

40.   K. Bonasia, *et al.*, Prior knowledge modulates the neural substrates of encoding and

retrieving naturalistic events at short and long delays. *Neurobiol. Learn. Mem.* **153**, 26–39

(2018).

41.   J. R. Andrews-Hanna, J. S. Reidler, J. Sepulcre, R. Poulin, R. L. Buckner, Functional-

Anatomic Fractionation of the Brain's Default Network. *Neuron* **65**, 550–562 (2010).

42.   B. T. T. Yeo, *et al.*, The organization of the human cerebral cortex estimated by intrinsic

functional connectivity. *J. Neurophysiol.* **106**, 1125–1165 (2011).

43.   J. L. Ji, *et al.*, Mapping the human brain's cortical-subcortical functional network

organization. *Neuroimage* **185**, 35–57 (2019).

44.   M. F. Glasser, *et al.*, A multi-modal parcellation of human cerebral cortex. *Nature* **108**,

125–38 (2016).

45.   V. D. Blondel, J. L. Guillaume, R. Lambiotte, E. Lefebvre, Fast unfolding of communities

in large networks. *J. Stat. Mech. Theory Exp.* **2008** (2008).

46.   A. L. Traud, E. D. Kelsic, P. J. Mucha, M. A. Porter, Comparing community structure to

characteristics in online collegiate social networks. *SIAM Rev.* **53**, 526–543 (2011).

47.   E. M. Gordon, *et al.*, Precision Functional Mapping of Individual Human Brains. *Neuron*

95, 791-807.e7 (2017).

48.   M. W. Cole, D. S. Bassett, J. D. Power, T. S. Braver, S. E. Petersen, Intrinsic and task-

evoked network architectures of the human brain. *Neuron* **83**, 238–251 (2014).

49.   A. W. Gilmore, S. M. Nelson, K. B. McDermott, The Contextual Association Network

Activates More for Remembered than for Imagined Events. *Cereb. Cortex* **26**, 611–617

(2016).

50.   E. H. Silson, A. Steel, A. Kidder, A. W. Gilmore, C. I. Baker, Distinct subdivisions of

human medial parietal cortex support recollection of people and places. *Elife* **8**, 554915

(2019).

51.   M. Peer, R. Salomon, I. Goldberg, O. Blanke, S. Arzy, Brain system for mental orientation

in space, time, and person. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11072–11077 (2015).

52.   M. Jacomy, T. Venturini, S. Heymann, M. Bastian, ForceAtlas2, a continuous graph

layout algorithm for handy network visualization designed for the Gephi software. *PLoS*

One **9**, 1–12 (2014).

53.   B. Fischl, *et al.*, Whole brain segmentation: automated labeling of neuroanatomical

structures in the human brain. *Neuron* **33**, 341–55 (2002).

54.   J. L. Vincent, *et al.*, Coherent Spontaneous Activity Identifies a Hippocampal-Parietal

Memory Network. *J. Neurophysiol.* **96**, 3517–3531 (2006).

55.   C. Baldassano, *et al.*, Discovering Event Structure in Continuous Narrative Perception and

Memory. *Neuron* **95**, 709-721.e5 (2017).

56.   M. Rubinov, O. Sporns, Complex network measures of brain connectivity: Uses and

interpretations. *Neuroimage* **52**, 1059–1069 (2010).

57.   E. M. Gordon, *et al.*, Three Distinct Sets of Connector Hubs Integrate Human Brain

Function. *Cell Rep.* **24**, 1687-1695.e4 (2018).

58.   R. A. Diana, A. P. Yonelinas, C. Ranganath, Imaging recollection and familiarity in the

medial temporal lobe: a three-component model. *Trends Cogn. Sci.* **11**, 379–386 (2007).

59.   C. J. Lynch, *et al.*, Precision Inhibitory Stimulation of Individual-Specific Cortical Hubs

Disrupts Information Processing in Humans. *Cereb. Cortex* **29**, 3912–3921 (2019).

60.   D. E. Warren, *et al.*, Network measures predict neuropsychological outcome after brain

injury. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14247–14252 (2014).

61.   C. Gratton, E. M. Nomura, F. Pérez, M. D'Esposito, Focal brain lesions to critical

locations cause widespread disruption of the modular organization of the brain. *J. Cogn.*

Neurosci. **24**, 1275–1285 (2012).

62.   M. A. Bertolero, B. T. T. Yeo, M. D'Esposito, The diverse club. *Nat. Commun.* **8**, 1–10

(2017).

63.   L. E. Frank, C. R. Bowman, D. Zeithamova, Differential Functional Connectivity along

the Long Axis of the Hippocampus Aligns with Differential Role in Memory Specificity

and Generalization. *J. Cogn. Neurosci.* **31**, 1958–1975 (2019).

64.   I. Przeździk, M. Faber, G. Fernández, C. F. Beckmann, K. V. Haak, The functional

organisation of the hippocampus along its long axis is gradual and predicts recollection.

Cortex **119**, 324–335 (2019).

65.   M. P. McAndrews, M. Cohn, Neuropsychology in Temporal Lobe Epilepsy: Influences

from Cognitive Neuroscience and Functional Neuroimaging. *Epilepsy Res. Treat.* **2012**,

1–13 (2012).

66.   J. Miller, *et al.*, Lateralized hippocampal oscillations underlie distinct aspects of human

spatial memory and navigation. *Nat. Commun.* **9** (2018).

67.   K. Iglói, C. F. Doeller, A. Berthoz, L. Rondi-Reig, N. Burgess, Lateralized human

hippocampal activity predicts navigation based on sequence or place memory. *Proc. Natl.*

Acad. Sci. U. S. A. **107**, 14466–14471 (2010).

68.   E. Mizrak, N. R. Bouffard, L. A. Libby, E. Boorman, C. Ranganath, Representation of

1057   Task Structure in Human Hippocampus. *bioRxiv*, 794305 (2019).

1058   69.   M. Inhoff, L. Libby, T. Noguchi, B. Love, C. Ranganath, Dynamic integration of

1059   conceptual information during learning. *PLoS One* **13**, 1–26 (2018).

1060   70.   T. Yarkoni, R. A. Poldrack, T. E. Nichols, D. C. Van Essen, T. D. Wager, Large-scale

1061   automated synthesis of human functional neuroimaging data. **8** (2011).

1062   71.   I. Kahn, D. Shohamy, Intrinsic connectivity between the hippocampus, nucleus

1063   accumbens, and ventral tegmental area in humans. *Hippocampus* **23**, 187–192 (2013).

1064   72.   D. J. Kravitz, K. S. Saleem, C. I. Baker, M. Mishkin, A new neural framework for

1065   visuospatial processing. *Nat. Rev. Neurosci.* **12**, 217–230 (2011).

1066   73.   E. M. Gordon, T. O. Laumann, B. Adeyemo, S. E. Petersen, Individual Variability of the

1067   System-Level Organization of the Human Brain. *Cereb. Cortex* **27**, 386–399 (2017).

1068   74.   Z. M. Reagh, C. Ranganath, What does the functional organization of cortico-

1069   hippocampal networks tell us about the functional organization of memory? *Neurosci.*

1070   *Lett.* **680**, 69–76 (2018).

1071   75.   W. L. Suzuki, D. G. Amaral, Perirhinal and parahippocampal cortices of the macaque

1072   monkey: Cortical afferents. *J. Comp. Neurol.* **350**, 497–533 (1994).

1073   76.   P. Lavenex, W. A. Suzuki, D. G. Amaral, Perirhinal and parahippocampal cortices of the

1074   macaque monkey: Projections to the neocortex. *J. Comp. Neurol.* **447**, 394–420 (2002).

1075   77.   J. Zhuo, *et al.*, Connectivity profiles reveal a transition subarea in the parahippocampal

1076   region that integrates the anterior temporal–posterior medial systems. *J. Neurosci.* **36**,

1077   2782–2795 (2016).

1078   78.   S. De Simoni, *et al.*, Disconnection between the default mode network and medial

1079   temporal lobes in post-traumatic amnesia Keywords: post-traumatic amnesia; traumatic

1080   brain injury; functional connectivity; default mode network; memory Abbreviations:

1081   DMN = default mode network; M. *Brain* **139**, 3137–3150 (2016).

1082   79.   E. R. Chrastil, S. M. Tobyne, R. K. Nauer, A. E. Chang, C. E. Stern, Converging meta-

1083   analytic and connectomic evidence for functional subregions within the human

1084   retrosplenial region. *Behav. Neurosci.* **132**, 339–355 (2018).

1085   80.   S. M. Frankland, J. D. Greene, Concepts and Compositionality: In Search of the Brain's

1086   Language of Thought. *Annu. Rev. Psychol.* **71**, 273–303 (2020).

1087   81.   S. T. Carmichael, J. L. Price, Sensory and premotor connections of the orbital and medial

1088   prefrontal cortex of macaque monkeys. *J. Comp. Neurol.* **363**, 642–664 (1995).

1089   82.   W. A. Suzuki, D. G. Amaral, Topographic organization of the reciprocal connections

1090   between the monkey entorhinal cortex and the perirhinal and parahippocampal cortices. *J.*

1091   *Neurosci.* **14**, 1856–1877 (1994).

1092   83.   S. Marek, *et al.*, Identifying reproducible individual differences in childhood functional

1093   brain networks: An ABCD study. *Dev. Cogn. Neurosci.* **40** (2019).

1094   84.   M. Bota, O. Sporns, L. W. Swanson, Architecture of the cerebral cortical association

1095   connectome underlying cognition. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E2093–E2101

1096   (2015).

1097   85.   E. M. Gordon, *et al.*, Default-mode network streams for coupling to language and control

1098   systems. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 17308–17319 (2020).

1099   86.   C. McCormick, E. Ciaramelli, F. De Luca, E. A. Maguire, Comparing and contrasting the

1100   cognitive effects of hippocampal and ventromedial prefrontal cortex damage: A review of

human lesion studies. *Neuroscience* (2017)

https:/doi.org/10.1016/j.neuroscience.2017.07.066.

87. H. Kondo, P. Lavenex, D. G. Amaral, Intrinsic connections of the macaque monkey

hippocampal formation: II. CA3 connections. *J. Comp. Neurol.* **377**, NA-NA (2009).

88. I. K. Brunec, *et al.*, Multiple Scales of Representation along the Hippocampal

Anteroposterior Axis in Humans. *Curr. Biol.* **28**, 2129-2135.e6 (2018).

89. J. W. Vogel, *et al.*, A molecular gradient along the longitudinal axis of the human

hippocampus informs large-scale behavioral systems. *Nat. Commun.* **11**, 1–17 (2020).

90. A. J. Barnett, V. Man, M. P. McAndrews, Parcellation of the Hippocampus Using Resting

Functional Connectivity in Temporal Lobe Epilepsy. *Front. Neurol.* **10**, 1–12 (2019).

91. C. L. Grady, Meta-analytic and functional connectivity evidence from functional magnetic

resonance imaging for an anterior to posterior gradient of function along the hippocampal

axis. *Hippocampus*, 1–16 (2019).

92. M. S. Fanselow, H. W. Dong, Are the Dorsal and Ventral Hippocampus Functionally

Distinct Structures? *Neuron* **65**, 7–19 (2010).

93. I. R. Olson, D. McCoy, E. Klobusicky, L. A. Ross, Social cognition and the anterior

temporal lobes: A review and theoretical framework. *Soc. Cogn. Affect. Neurosci.* **8**, 123–

133 (2013).

94. M. Cohn, M. St-Laurent, A. Barnett, M. P. McAndrews, Social inference deficits in

temporal lobe epilepsy and lobectomy: Risk factors and neural substrates. *Soc. Cogn.*

Affect. Neurosci. **10**, 636–644 (2015).

95. L. M. DiNicola, R. M. Braga, R. L. Buckner, Parallel distributed networks dissociate

episodic and social functions within the individual. *J. Neurophysiol.* **123**, 1144–1179

(2020).

96.   J. R. Andrews-Hanna, R. Saxe, T. Yarkoni, Contributions of episodic retrieval and

1126   mentalizing to autobiographical thought: Evidence from functional neuroimaging, resting-

1127   state connectivity, and fMRI meta-analyses. *Neuroimage* **91**, 324–335 (2014).

1128   97.   S. Lee, T. Parthasarathi, J. W. Kable, The dorsal and ventral default mode networks are

1129   dissociably modulated by the valence and vividness of imagined events. *bioRxiv*,

1130   2020.05.21.109728 (2020).

1131   98.   N. E. Nawa, H. Ando, Effective connectivity within the ventromedial prefrontal cortex-

1132   hippocampus-amygdala network during the elaboration of emotional autobiographical

1133   memories. *Neuroimage* **189**, 316–328 (2019).

1134   99.   N. Franzmeier, *et al.*, Functional brain architecture is associated with the rate of tau

1135   accumulation in Alzheimer's disease. *Nat. Commun.* **11**, 1–17 (2020).

1136   100.   S. M. Hayes, D. H. Salat, M. Verfaellie, Default network connectivity in medial temporal

1137   lobe amnesia. *J. Neurosci.* **32**, 14622–14630 (2012).

1138   101.   J. D. Power, *et al.*, Functional Network Organization of the Human Brain. *Neuron* **72**,

1139   665–678 (2011).

1140   102.   E. M. Gordon, *et al.*, Individual-specific features of brain systems identified with resting

1141   state functional correlations. *Neuroimage* **146**, 918–939 (2017).

1142   103.   O. Esteban, *et al.*, fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat.*

1143   *Methods* **16**, 111–116 (2019).

1144   104.   O. Esteban, *et al.*, fMRIPrep: a robust preprocessing pipeline for functional MRI (2019)

1145   https:/doi.org/10.5281/ZENODO.2859286 (March 31, 2020).

105.   K. Gorgolewski, *et al.*, Nipype: A Flexible, Lightweight and Extensible Neuroimaging

Data Processing Framework in Python. *Front. Neuroinform.* **5** (2011).

106.   K. J. Gorgolewski, *et al.*, nipy/nipype: 1.2.0 (2019)

https:/doi.org/10.5281/ZENODO.2685428 (March 31, 2020).

107.   R. W. Cox, J. S. Hyde, Software tools for analysis and visualization of fMRI data. *NMR*

Biomed. **10**, 171–8.

108.   M. Jenkinson, S. Smith, A global optimisation method for robust affine registration of

brain images. *Med. Image Anal.* **5**, 143–156 (2001).

109.   D. N. Greve, B. Fischl, Accurate and robust brain image alignment using boundary-based

registration. *Neuroimage* **48**, 63–72 (2009).

110.   M. Jenkinson, P. Bannister, M. Brady, S. Smith, Improved optimization for the robust and

accurate linear registration and motion correction of brain images. *Neuroimage* **17**, 825–

841 (2002).

111.   C. Lanczos, Evaluation of noisy data. *J. Soc. Ind. Appl. Math. Ser. B Numer. Anal.* **1**, 76–

85 (1964).

112.   A. Abraham, *et al.*, Machine learning for neuroimaging with scikit-learn. *Front.*

Neuroinform. **8** (2014).

113.   R. Ciric, *et al.*, Benchmarking of participant-level confound regression strategies for the

control of motion artifact in studies of functional connectivity. *Neuroimage* **154**, 174–187

(2017).

114.   A. Weissenbacher, *et al.*, Correlations and anticorrelations in resting-state functional

connectivity MRI: A quantitative comparison of preprocessing strategies. *Neuroimage* **47**,

1408–1416 (2009).

115.   J. D. Power, *et al.*, Methods to detect, characterize, and remove motion artifact in resting

state fMRI. *Neuroimage* **84**, 320–341 (2014).

116. M. Rubinov, O. Sporns, Weight-conserving characterization of complex functional brain

networks. *Neuroimage* **56**, 2068–2079 (2011).

117. C. Gratton, *et al.*, Functional Brain Networks Are Dominated by Stable Group and

Individual Factors, Not Cognitive or Daily Variation. *Neuron* **98**, 439-452.e5 (2018).

118. B. A. Seitzman, *et al.*, A set of functionally-defined brain regions with improved

representation of the subcortex and cerebellum. *Neuroimage* **206**, 116290 (2020).

119. M. A. Bertolero, B. T. T. Yeo, D. S. Bassett, M. D'Esposito, A mechanistic model of

connector hubs, modularity and cognition. *Nat. Hum. Behav.* **2**, 765–777 (2018).

120. S. Seabold, J. Perktold, Statsmodels: econometric and statistical modeling with python.

Proc. 9th Python Sci. Conf. , June 28 - July 3, Austin, Texas, 57–61 (2010).

121. H. Nili, *et al.*, A Toolbox for Representational Similarity Analysis. *PLoS Comput. Biol.*

10, e1003553 (2014).

1183